1 **Replication protein-A, RPA, plays a pivotal role in the maintenance of**
2 **recombination checkpoint in yeast meiosis**
3

4 **Arivarasan Sampathkumar, Masaru Ito, Yurika Fujita, and Akira Shinohara***
5
6 Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan
7

8 Running title: Essential role of RPA in recombination checkpoint
9
10 Keywords: meiosis; homologous recombination; RPA; recombination checkpoint;
11 Rfa1, auxin degron, synapsis checkpoint
12
13
14

15 *Corresponding author:
16 Akira Shinohara; ashino@protein.osaka-u.ac.jp
17
Lead contact:
Akira Shinohara
Institute for Protein Research, Osaka University,
3-2 Yamadaoka, Suita, Osaka 565-0871 JAPAN
Phone: 81-6-6879-8624
FAX: 81-6-6879-8626
E-mail: ashino@protein.osaka-u.ac.jp
ORCID: 0000-0003-4207-8247
26

Summary

DNA double-strand breaks (DSBs) activate DNA damage responses (DDR) in both mitotic and meiotic cells. Meiotic DSBs induce homologous recombination monitored by a meiotic DDR called the recombination checkpoint for the pachytene exit in meiotic prophase I. In this study, we showed the essential role of a single-stranded DNA (ssDNA) binding protein, Replication protein-A (RPA), in the maintenance of the recombination checkpoint during *S. cerevisiae* meiosis. The depletion of an RPA subunit, Rfa1, in a recombination-defective *dmc1* mutant, fully alleviates the pachytene arrest with the persistent unrepaired DSBs. RPA depletion downregulates a meiosis-specific CHK2 homolog, Mek1, which in turn activates Ndt80 transcriptional activator for pachytene exit. These support the idea that RPA is a sensor of ssDNAs for the activation of meiotic DDR. Rfa1 depletion also accelerates the prophase I delay induced by the *zip1* mutant defective in both chromosome synapsis and the recombination, suggesting that the accumulation of ssDNAs rather than defective synapsis triggers prophase I relay in the *zip1* mutant.

Introduction

A eukaryotic single-stranded (ss)DNA binding protein, Replication protein-A (RPA), plays an essential role in DNA replication, DNA repair, and homologous recombination [1-3]. RPA consists of three subunits; RPA70(RPA1), RPA32(RPA2), and RPA14(RPA3), which are highly conserved from yeast to humans. In *S. cerevisiae*, the knockout of three genes (*RFA1*, *-2* and *-3*) encoding the RPA subunits leads to lethality, consistent with its essential role in DNA replication [4].

DNA double-strand breaks (DSBs) initiate homologous recombination. RPA binds to ssDNA formed by the 5'-3' resection of the DSB ends to prevent the formation of the secondary structure of ssDNAs and to protect ssDNAs from degradation by various nucleases [5-7]. The assembly of Rad51 filaments on RPA-coated ssDNA is assisted by Rad51 mediators such as Rad52, Rad55-57, and Psy3-Csm2-Shu1-Shu2 in the budding yeast [8-10], and the Rad51 filament promotes homology search and strand exchange with a homologous double-stranded (ds)DNA [11,12]. The invasion of the ssDNA into the homologous dsDNA results in the formation of a displacement loop (D-loop). The 3'-OH end of the invading strand becomes a substrate for DNA synthesis. When a newly synthesized DNA strand is dissociated from the D-loop, the displaced ssDNA of the DSB end can capture the ssDNA of the other DSB end, which leads to a

gapped DNA for further second DNA synthesis. On the other hand, the stable
association of the synthesized ssDNA with the D-loop results in the extension of
the D-loop. The annealing of the displaced ssDNA with the ssDNA of the other
DSB end results in the recombination intermediate with a Holliday structure. In
these steps, RPA promotes DNA synthesis by DNA polymerases and the
annealing between ssDNAs by Rad52 protein [13,14].

DNA damage activates the DNA damage response (DDR). DSB ends
stimulate a DDR kinase, ATM/Tel1 (Ataxia telangiectasia mutated) while ssDSBs
do an ATM homolog, ATR/Mec1 (Ataxia telangiectasia mutated related), both of
which belong to the PI3 kinase family [15,16]. The activation of ATR is mediated by
the interaction with an ATR partner, ATRIP/Ddc2, which directly binds to RPA [17-
19]. In mammals, ATR activation is also mediated by the second RPA-binding
protein, ETAA1 [20,21]. Thus, RPA is considered to work as an essential DNA
damage sensor to detect ssDNAs, as well as a DNA damage processing protein.
The Mec1/ATR activates downstream DDR responses such as DNA repair, cell-
cycle arrest/delay, and/or apoptosis through a mediator kinase such as
CHK2/Rad53.

Meiosis rewires DNA transactions such as homologous recombination/DSB
repair as well as DDR response to guarantee the faithful segregation of two
paternal homologous chromosomes at meiosis I for the reduction of chromosome
numbers in gametes. In budding yeast, meiotic recombination is initiated by the
DSB formation at recombination hotspots by the TopoVI-like protein, Spo11,
which is followed by the formation of ssDNAs [22,23]. The ssDNA is used by a
meiosis-specific Rad51 homolog, Dmc1 [24], for D-loop formation, which is aided
by Rad51 [25] and its mediator, Mei5-Sae3 complex [26-28]. Further processing
generates a recombination intermediate with double-Holliday junctions (dHJs),
which are specifically resolved into crossovers between homologs. Importantly,
the interhomolog bias for the partner choice in the recombination is mediated by
meiosis-specific chromosome-axis proteins, Hop1-Red1-Mek1/Mre4, via an
unknown mechanism [29-31]. Mek1/Mre4 (hereafter Mek1) kinase is a meiosis-
specific homolog of Rad53/CHK2 kinase, which suppresses Rad51-mediated
recombination in meiosis [32,33]. This suppression occurs through the Mek1-
mediated phosphorylation of two proteins, Rad54 and Hed1 [34,35]. Rad54 belongs
to the Swi2/Snf2 family protein essential for Rad51-mediated strand invasion [36,37].
Mek1-mediated phosphorylation of Rad54 suppresses its interaction with Rad51
[35]. Hed1, which is expressed only in meiosis, inhibits Rad51 by binding to the

Rad51-ssDNA complex [38]. The inhibitory activity of Hed1 against Rad51 and its
protein stability are augmented by the Mek1 phosphorylation [34].
Meiotic events such as recombination and chromosome synapsis are strictly
coupled with each other and also with meiosis progression (Figure 0) [39]. Similar
to mitosis, DSBs in yeast meiosis activate Tel1(ATM) and Mec1(ATR). Different
from mitotic DNA damage checkpoint, downstream events of Mec1 or Tel1
activation such as the Rad53 activation are largely suppressed in meiotic cells
with poorly-defined mechanisms [40,41]. On the other hand, Mec1 and/or Tel1
phosphorylates a meiosis-specific chromosomal axis protein, Hop1 [42]. Mec1
activation is assisted by DDR clamp, Ddc1-Rad17-Mec1 (RAD9-RAD1-HUS1, 9-
1-1 complex), and clamp loader, Rad24(RAD17) [42]. Together with chromosomal-
bound Red1, phosphorylated Hop1 activates Mek1 kinase, whose
autophosphorylation provides positive feedback on its activity [43]. Mek1-mediated
phosphorylation promotes the inactivation of the Ndt80 transcription factor [44],
which is essential for the exit from the mid-pachytene stage, where chromosome
synapsis and the formation of the recombination intermediates with dHJs are
completed [43,45]. By activating Ndt80 (inactivating Mek1), meiotic cells exit the
pachytene stage by expressing downstream targets such as Cdc5 Polo-like
kinase (PLK) and a cyclin, Clb1 [46] for the resolution of dHJs and the disassembly
of the synaptonemal complex (SC), tripartite chromosome structure specific to
meiosis.
It is widely believed that meiotic cells surveil not only homologous
recombination but also chromosome synapsis (Figure 1). Like a recombination-
defective *dmc1* mutant, the budding yeast *zip1* mutant, which is defective in the
assembly of the central region of the SCs, thus synapsis, shows a delay in
meiosis progression [47]. In nematodes and mice, synapsis-defective mutants also
delay the meiotic prophase I progression and/or induce apoptosis [48,49]. However,
in meiosis, chromosome synapsis is tightly coupled with meiotic recombination:
e.g. a recombination-deficient mutant display also abnormal synapsis [50,51]; even
the *zip1* mutant completely defective in SC formation impairs the processing of
meiotic recombination intermediates. These complicate the presence of the
synapsis checkpoint in meiotic cells. Moreover, genes involved in DDR response
also play a role in meiotic recombination, which also provides a caveat in the
interpretation of the meiotic phenotypes of DDR mutants [52,53].
Although the role of RPA in DDR response is widely accepted, the
involvement *in vivo* remains unaddressed. A previous report showed that Rfa1

depletion in G2/M-arrested mitotic cells in budding yeast does not affect
checkpoint activation to a single DSB [5]. Moreover, the depletion of Rpa1 in mouse
spermatocytes eliminated RPA foci on meiotic chromosomes but is proficient in
ATR recruitment to the chromosomes [54]. These studies cast doubt on the role of
RPA in ATR-dependent checkpoint activation. In this study, we define the role of
RPA in the meiotic recombination checkpoint by using a conditional-depletion
mutant of Rfa1, RPA70 subunit of yeast RPA. The depletion of Rfa1 in the *dmc1*-
induced pachytene arrest facilitates the pachytene exits, thus entry into meiosis I
and II. Importantly, the RPA depletion accumulates meiotic DSBs without any
repair, which causes abnormal segregation of chromosomes in the division. This
confirms the essential role of RPA in the maintenance of the meiotic
recombination checkpoint. Moreover, we also found that Rfa1 depletion also
alleviates delayed meiosis progression in the *zip1* mutant, which is deficient in
SC formation and recombination. This suggests that the *zip1* mutation mainly
triggers an RPA-dependent response of the prophase delay, thus the
recombination checkpoint rather than the synapsis checkpoint.
151
Results
Depletion of Rfa1 alleviates *dmc1*-induced arrest of meiosis
The formation of ssDNA triggers cell cycle delay or arrest in both mitotic and
meiotic cells. To know the role of a major eukaryotic ssDNA-binding protein, RPA,
in the DNA damage-induced cell cycle arrest in budding yeast meiosis, we used
the *dmc1* mutant, which is unable to repair meiotic DSBs and as a result, an arrest
at mi-pachytene stage during meiotic prophase I [24]. This response to unrepaired
DSBs is referred to as the recombination checkpoint. Since all subunits of RPA,
Rfa1, -2, and -3, are essential for cell viability, we constructed a conditional
depletion mutant of the *RFA1* gene encoding the largest RPA subunit in the
budding yeast by the auxin-induced degron (AID) system. The AID tag was added
to the C-terminal of the Rfa1 protein. To know the role of RPA in the recombination
checkpoint in meiosis, *RFA1-AID* was introduced in the *dmc1* deletion mutant;
dmc1 RFA1-AID. Like the *dmc1* mutant, without auxin addition, *dmc1 RFA1-AID*
shows an arrest in meiotic prophase I without division (e.g. Figure 2B), indicating
that the Rfa1 tagging did not affect the recombination checkpoint in the absence
of Dmc1.
RPA is also involved in DNA replication and S-phase checkpoint [1,2], which
indirectly affects the G2 event. To avoid the effect in pre-meiotic DNA replication,

we depleted Rfa1 protein by adding auxin to a meiotic culture of *dmc1 RFA1-AID*
cells at 4 h after the induction of meiosis in the sporulation medium (SPM), after
pre-meiotic DNA replication, which occurs at 1.5-3 h. At 5 h, after the 1-hour
addition of auxin, the protein level of Rfa1-AID was decreased to ~15% level
relative to the original level at 4 h (Figure 2A), indicating the efficient depletion of
Rfa1 under the condition. We also checked the focus formation of Rfa2, the
second subunit of RPA, on meiotic chromosome spreads. Rfa2 shows punctate
staining on the spreads with other recombination proteins such as Rad51 [55],
which corresponds with ongoing recombination (Figure 2B). In the *dmc1* mutant,
Rfa2 focus-staining appears at 4 h, reaches a plateau at 5 h, and persists after
that, indicating little turnover of Rfa2 due to stalled recombination in the mutant
(Figure 2C, Supplemental Figure S1A). When auxin was added at 4 h, Rfa2 foci
disappeared within one hour (Figure 2A), and Rfa2 foci were rarely observed
during further incubation by 10 h (Figure 2B, C). This suggests that chromosome-
bound RPA is highly dynamic with an exchange of chromosome-bound ones with
that in a pool of nucleoplasm. We concluded that Rfa1 depletion at 4 h in meiosis
eliminates most chromosome-bound RPA complexes by 5 h.
188
Checkpoint alleviation in Rfa1 depletion in *dmc1* meiosis
DAPI staining of DNAs confirmed a single DAPI body in a *dmc1 RFA1-AID* cell
even at 24 h, thus an arrest in mid-pachynema, as shown previously [24]. Rfa1
depletion at 4 h induced fragmentation of DAPI bodies at late time points. In some
cells, 2-4 irregular-sized DAPI bodies with or without smaller DAPI bodies were
detected in a single cell (Figure 2D, top). The cells with abnormal DAPI bodies
appeared at 6 h and gradually accumulated by 24 h (Figure 2D, bottom). This
suggests the presence of nuclear division, thus the onset of meiosis I and/or II in
the *dmc1* cells upon Rfa1 depletion.
We also checked the effect of Rfa1 depletion on meiotic DNA replication.
The FACS analysis showed that the addition of auxin at 4 h did not affect the
transition from 2C to 4C (Figure 2E).
To evaluate the meiosis progression precisely, we examined the status of
spindle morphology in the *dmc1 RFA1-AID* cells in the absence or presence of
auxin (Figure 2F-H). Tubulin staining revealed a single focus associated with
single or multiple short projections in meiotic prophase I of the cell in the absence
of auxin (Figure 2G). The *dmc1 RFA1-AID* mutant maintained this arrangement
of spindles in the absence of auxin by 24 h, consistent with an arrest of the *dmc1*

mutant in prophase I (Figure 2H, left). On the other hand, the Rfa1 depletion
induced the elongation of spindles after one hour of addition (at 5 h). At 6 h, ~50%
dmc1 RFA1-AID cells contained single short and long spindles corresponding
with metaphase I and anaphase I, respectively (Figure 2H). At 8 h, in addition to
the metaphase I and anaphase I spindle, ~70% of cells contained metaphase II
or anaphase II spindles (Figure 2G). At 10 h, most of the cells showed diffuse
staining of microtubules in multiple DAPI bodies in a cell. These show that Rfa1
depletion almost alleviates meiotic prophase I arrest induced by the *dmc1*
mutation.
Cdc5/PLK is expressed after the pachytene exit [46]. Like *dmc1* cells, without
auxin addition, the *dmc1 RFA1-AID* mutant cells did not express Cdc5 (Figure
2A). On the other hand, the addition of auxin at 4 h induced the expression of
Cdc5 from 5 h. This confirms the exit of the pachytene stage of *dmc1* cells in the
absence of RPA.
221
No DSB repair in *dmc1* meiosis with Rfa1 depletion
One possibility to suppress the *dmc1* arrest is to channel unrepaired DSBs into
the repair pathway: e.g. the *rad51* mutation partially suppresses the *dmc1* arrest
[56,57]. On the other hand, the acute depletion of RPA did not affect the status of
meiotic DSBs. The CHEF gel electrophoresis confirmed the accumulation of
fragmented chromosomes in the *dmc1* mutant irrespective of the presence or
absence of RPA. In the *dmc1 RFA1-AID* mutant without auxin treatment, smear
bands of full-length chromosomes appeared at 3 h and most of the chromosomes
became fragmented with a size of less than 700 kbp by 5 h. After 6 h, fragmented
chromosomes persisted by 24 h (Figure 3A). The depletion of Rfa1 from 4 h did
not affect the persistence of fragmented chromosomes. This suggests that
meiotic DSBs remain unrepaired under the depletion of RPA in the *dmc1* mutant.
We also confirmed the DSB repair by immuno-staining analysis of Rad51
on meiotic chromosomes (Figure 3B, C). Like Rfa2 foci, in the *dmc1 RFA1-AID*
mutant in the absence of auxin, Rad51 foci appeared at 4 h, reached a plateau
at 5 h, and persisted at late times as shown previously [58]. The addition of auxin
at 4 h rapidly induced the disappearance of Rad51 foci at 5 h. The reduced Rad51
foci are seen at late time points (Figure 3C and Supplemental Figure 1B). This
suggests that RPA is critical for the maintenance of Rad51 foci on the
chromosomes. The absence of Rad51 foci in the Rfa1 depletion supports the
inability of *dmc1 RFA1-AID* cells without RPA to repair meiotic DSBs. Thus, RPA

might play an essential role in the post-synaptic stage of Rad51 assembly.

245   The Hop1/Mek1-Ndt80 axis was down-regulated after RFA1 depletion

Un-repaired DSBs in the *dmc1* mutant activate the recombination checkpoint,

which down-regulates a meiosis-specific transcription activator, Ndt80 for the

pachytene exit (Figure 1). DSB-activated Mec1/ATR kinase phosphorylates a

HORMAD protein, Hop1, which stimulates a meiosis-specific kinase, Mek1. Mek1

binds and phosphorylates Ndt80 to inactivate it. For pachytene exit, further

phosphorylation of Nd80 by Ime2 kinase promotes the activation of Ndt80 [59]. In

dmc1 RFA1-AID cells, the Ndt80 appeared at 4 h with multiple bands and

accumulated more during further incubation in SPM without a change of band-

shift status (Figure 2A). At 6 h, after two hours of addition of auxin, a slower-

migrating Ndt80 band appeared and gradually decreased in intensity. Importantly,

as shown above, under the depletion condition, Cdc5/PLK, which is a major target

of activated Ndt80 and is necessary for pachytene exit, appeared at 6 h (Figure

2A). This is consistent with meiosis progression induced by Rfa1-depletion as

shown above.

We also checked the behavior of the upstream regulators of Ndt80, Hop1,

and Mek1 by western blotting. In the *dmc1* mutant, Hop1 appeared as a non-

phosphorylated form at 2 h and phosphorylated Hop1 with slower migrated bands

by Mec1(ATR) or Tel1(ATM) persisted from 4 h (Figure 2A) as shown previously

[42]. After Rfa1 depletion, major bands of phosphorylated Hop1 almost disappeared

from 6 h with a similar steady-state level of non-phosphorylated Hop1 during the

time course. This suggests that Hop1 phosphorylation is actively de-

phosphorylated during pachytene exit. On chromosome spreads, Hop1 shows

beads-in-line staining for the localization of chromosome axes (Figure 3D) [60,61].

In the *dmc1* mutant without Rfa1 depletion, Hop1 localization remains unchanged

during meiosis (Figure 3D). The addition of auxin at 4 h induced the

disappearance of Hop1-positive spreads from 5 h. At 6 h, ~35% of cells are

positive for Hop1 while ~95% are positive in a control (Figure 3D, E). At 8 h, most

of the Hop1 signals on the chromosomes disappeared.

One of the Mek1 kinase targets is histone H3T11 phosphorylation [62]. Under

the *dmc1* arrest condition, H3T11 phosphorylation accumulated (Figure 2A). On

the other hand, the Rfa1 depletion rapidly induced the disappearance of the

signal (Figure 2A), suggesting the inactivation of Mek1 under Rfa1 depletion.

These results clearly showed that Rfa1, thus RPA, is critical for the maintenance

of the recombination checkpoint through the Hop1/Mek1-Ndt80 axis.

Bypass of *dmc1* arrest by Rfa1 depletion depends on Ndt80

We confirmed whether the pachytene exit of the *dmc1* cells by Rfa1 depletion

depends on the *NDT80* (Figure 4). The addition of auxin to *dmc1 ndt80 RFA1-*

AID cells efficiently reduced the amount of Rfa1 protein (Figure 4A) and Rfa2 foci

from meiotic chromosomes (Figure 4B). However, different from the *dmc1 RFA1-*

AID mutant, the addition of auxin to *dmc1 ndt80 RFA1-AID* mutant did not induce

the elongation of spindles (Figure 4C) or Hop1 disassembly from the

chromosomes (Figure 4D). Thus, the pachytene exit of *dmc1* cells by Rfa1

depletion still depends on Ndt80. Interestingly, the amount of the phosphorylated

Hop1 is reduced slightly compared to that in the control (Figure 4A). This is not

drastic compared to the presence of Ndt80, supporting an auto-feedback

mechanism between Mec1 activation and Ndt80 activation [63]. Moreover, the

Hop1 dephosphorylation is not a prerequisite for Hop1 dissociation from meiotic

chromosomes.

Checkpoint alleviation in Rfa1 depletion in *zip1* meiosis

Meiotic yeast cells monitor not only ongoing meiotic recombination but also

chromosome synapsis [49]. However, the presence of the so-called "synapsis

checkpoint" is under debate [63]. The mutant in the *ZIP1* gene, which encodes the

transverse filament of the central region of the SC [47], is deficient in chromosome

synapsis with a delay of the entry into meiosis I and is used to analyze the

synaptic checkpoint. We examined the effect of Rfa1 depletion on the *zip1-*

induced delay of meiosis I (Figure 5). As reported previously [51], in the SK1 strain

background, *zip1 RFA1-AID* cells without auxin addition started meiosis I at 8 h

while wild-type cells entered at 5 h (and finished meiosis II by 10 h) (Figure 5B).

When auxin was added at 4 h, which induced rapid depletion of Rfa1 and the

dissociation of Rfa2 from meiotic chromosomes (Figure 5A, C, I), metaphase I

spindles appeared at 5 h, and at 6 h, ~35% of the cells contained metaphase-I

and anaphase-I spindles (Figure 5D, E). At 8 h, ~30% of the cells contained

anaphase-II spindle. However, compared to the control, the entry into meiosis I

and II is not efficient.

The *zip1 RFA1-AID* cells without auxin expressed Cdc5 from 8 h (Figure

5A), consistent with delayed progression of prophase I in the mutant. The addition

of auxin at 4 h accelerated the Cdc5 expression at 5 h. The *zip1* mutant

accumulated T11 phosphorylation of histone H3 signal from 4 h to 8 h (Figure 5A).
The Rfa1 depletion accelerated the disappearance of this phosphorylation from
5 h. The accelerated disappearance of the phosphor-Hop1 protein is also seen
by the depletion. These showed the Rfa1-dependent inactivation of the
Hop1/Mek1-Ndt80 signaling pathway induced by the *zip1* mutation. Thus, the
delay induced by the *zip1* mutation largely depends on the Rfa1 function.

In addition to SC formation, Zip1 plays a critical role in meiotic recombination
as a pro-CO factor. Indeed, the *zip1* mutant shows delayed DSB repair. The
CHEF analysis showed the presence of broken chromosomes in the *zip1* mutant
from 3 h to 8 h (Figure 5H). Importantly, full-length chromosomes were
reassembled at 10 h without Rfa1 depletion, indicating that most of the DSBs in
the mutant are repaired with the delay. On the other hand, Rfa1 depletion from 4
h induced fragmentation of the chromosomes from 5 h. Even after 24 h, little
recovery of full-length chromosomes was seen under the condition. This suggests
that Rfa1 is essential for meiotic DSB repair in the *zip1* mutant.

Rfa1 depletion does not alleviate delay induced by *rad50S* meiosis
The *rad50S* mutant, which is defective in the end-processing of meiotic DSBs,
also shows delayed meiosis I progression [64]. This delay in the *rad50S* mutant is
mediated by Tel1(ATM) and Rad50-Mre11-Xrs2, which recognize DSB ends [65].
The pathway is distinct from the Mec1(ATR)-mediated pathway shown above. We
also examined the effect of Rfa1 depletion on the *rad50S*-induced delay (Figure
6). As reported, without auxin, *rad50S RFA1-AID* cells with auxin addition delay
the onset of meiosis I as shown by whole-cell staining of tubulin (Figure 6B).
Rather, the addition of auxin delayed the onset of meiosis I induced by the *rad50S*
mutation. We cannot detect Rfa2 foci in the r*ad50S* mutant even in the presence
of Rfa1 (Figure 6C), which accumulates DSBs in late meiosis I as shown by the
CHEF analysis (Figure 6D). The Rfa1 depletion did not affect the kinetics of the
accumulation of chromosome fragments (Figure 6D). Moreover, Hop1
phosphorylation is not observed in the mutant (Figure 6A). This is consistent with
little formation of ssDNAs without DSB processing in the mutant [64], thus little
Mec1/ATR activation. This is not surprising since the *rad50S* mutant does not
form any ssDNAs, which is a substrate for RPA. The accumulation of
chromosomal Hop1 in the mutant is not affected by the Rfa1 depletion (Figure
6E).

Discussion

RPA plays multiple roles in DNA replication, DNA repair, and recombination through its ability to bind to ssDNAs with a high affinity. RPA is also important for DDR as a sensor of DNA damage, thus ssDNAs. Although the role of RPA in Mec1/ATR activation through the binding of Mec1/ATR-interaction protein, Ddc1(ATRIP), to RPA has been documented well [17,18,66], the *in vivo* evidence to support the role of RPA as the DDR sensor is still missing. In this paper, we showed the pivotal role of RPA in the maintenance of DDR in meiosis, called recombination checkpoint.

The role of RPA in recombination checkpoint

Although the role of RPA on ssDNA for the recruitment of Mec1-Ddc2(ATR-ATRIP) for Mec1/ATR-activation in the DDR response is well-established [17,18,66], There is no provided *in vivo* evidence that RPA acts as an ssDNA sensor. Previously, in the budding yeast, two *RFA1*-degron mutants, *rfa1-td* and *td-RFA1*, were constructed and characterized to study the role of Rfa1 in DDR activation. Indeed, the *rfa1-td*, as well as the *rft1-t11 (rfa1-K45E)* mutant, reduce the recruitments of Ddc2 to a DSB site [66]. On the other hand, it is shown that, by using the *td-RFA1*, G2/M-arrested haploid cells depleted Rfa1 still activate Rad53 upon a single DSB [5]. The latter study concludes that a low level of RPA is sufficient for the activation of Mec1-Ddc2. More recently, an RPA-independent pathway for ATR activation, which requires APE1, was identified in *Xenopus* extracts [67]. We evaluated the role of RPA in DDR activation by studying the effect of Rfa1 depletion in the meiotic recombination checkpoint which shares the Mec1-dependent DDR pathway with a canonical DNA damage checkpoint in yeast mitotic cells. By using the auxin-degron, we depleted Rfa1 in the *dmc1* mutant which arrests in meiotic prophase I with hyper-resected ssDNAs in ~150 DSBs [24]. As expected, the Rfa1-depletion induced the pachytene exit by downregulating Mec1-Mek1 activity, which in turn activates the Ndt80-dependent gene expression of key regulators for the pachytene exit such as Cdc5 [43]. Importantly, the Rfa1 depletion did not affect the accumulation of ssDNAs in the *dmc1* cells with persistent unrepaired DSBs. This is particularly important since mutations in DDR genes such as *RAD17*, *-24*, *DDC1*, and *MEC3* even in the *MEC1* gene bypass the *dmc1*-induced arrest by activating the other DSB repair pathway such as ectopic recombination [52,53,68]. The results described here provide strong *in vivo* evidence that RPA is a sensor for ssDNAs, which creates a scaffold for the

assembly of checkpoint effectors assembly such as Mec1-Ddc1(ATR-ATRIP). In
meiosis, rather than activating Rad53 kinase, a meiosis-specific Mek1 is activated
through the phosphorylation of Hop1 by Mec1 kinase[42]. The Rfa1-depletion
induces rapid dephosphorylation of Hop1 and the dissociation from meiotic
chromosomes without affecting the protein level, suggesting the presence of
active de-phosphorylation, probably by Mec1-bound phosphatase, PP4 [69,70].
393

The chromosome synapsis checkpoint in yeast meiosis
In addition to the recombination checkpoint, meiotic cells show a response to
abnormal chromosome synapsis called synapsis checkpoint. This idea was
proposed that mutants defective in chromosome synapsis show a delay in the
progression of prophase I and apoptosis in yeasts and mice, respectively.
However, the presence of the synapsis checkpoint is still under debate [48,49], since
in most organisms meiotic recombination and chromosome synapsis are tightly
coupled. The budding yeast *zip1* mutant, which is completely defective in SC
formation, thus, chromosome synapsis, delays or arrest in meiotic prophase I [47,51].
This arrest is alleviated by the mutant of genes encoding a component of
checkpoint such as *PCH2* [71]. Pch2 modulates Hop1 conformation and regulates
the association and dissociation of Hop1 on meiotic chromosomes [56,72,73]. On the
other hand, the *zip1* mutant also shows defective DSB repair and/or the crossing-
over [50,51], which suggests delayed progression of meiosis I in the *zip1* mutant is
caused by the impaired recombination. In this study, by using the *RFA1-AID*, we
could test whether the delay in *zip1* mutant depends on RPA or not and found
that the Rfa1-depletion eliminates the delay induced by the *zip1* mutation. This
strongly suggests that the recombination checkpoint delays the meiosis I
progression in the *zip1* mutant as seen in the *dmc1* mutant. Moreover, consistent
with this, the other yeast mutants, which are defective in the synapsis with normal
recombination such as *ecm11* and *gmc2*, do not show a drastic delay in the
meiosis I progression [74,75]. Thus, at least in budding yeast which couples the
recombination and chromosome synapsis, rather than the status of chromosome
synapsis, a recombination process, the presence of ssDNAs, can be monitored
by a meiosis-specific surveillance mechanism to guarantee the completion of
meiotic recombination directly and chromosome synapsis indirectly.
420

Materials and Methods
Strains and plasmids

All strains described here are derivatives of *S. cerevisiae* SK1 diploids,
NKY832/833 (*MATα/MATa, lys2/'', ura3/'', leu2::hisG/'', trp1::hisG/''*). The
genotypes of each strain used in this study are described in Supplementary Table
S1.
427

Strain Construction

429   The *RFA1-AID* gene was constructed by one-step DNA integration using a
430   template plasmid, which was provided by Dr. Neil Hunter, with primers shown in
431   Table S2. DNA changes were confirmed by DNA sequencing.
432

Anti-serum and antibodies

Anti-myc (Santa Cruz), anti-tubulin (MCA77G, Bio-Rad/Serotec, Ltd), anti-Cdc5
(Santa Cruz), and histone H3 phospho-T11 (Abcam) were used for western
blotting. Rabbit anti-Rad51 [25] and guinea pig anti-Rad51 [76], rabbit anti-Hop1 and
rabbit anti-Rfa2 were used for staining. Rabbit anti-Ndt80 was a gift by M. Lichten.
The secondary antibodies for immuno-staining were Alexa Fluor 488 (Goat) and
594 (Goat) IgG used at a 1/2000 dilution (Thermo Fisher Scientific). For western
blotting, alkaline phosphatase-conjugated secondary antibodies (Promega) were
used.
442

Meiotic Time course

Saccharomyces cerevisiae SK1 strains were patched onto YPG plates (2%
bacteriological peptone, 1% yeast extract, 3% glycerol, 2% bacteriological agar)
and incubated at 30 °C for 12 h. Cells were inoculated onto YPD plates (2%
bacteriological peptone, 1% yeast extract, 2% glucose, 2% bacteriological agar)
and grown for two days to obtain isolated colonies. A single colony was inoculated
into 3 ml of YPD liquid medium and grown to saturation at 30 °C overnight. To
synchronize cell cultures, the overnight cultures were transferred to pre-warmed
SPS medium (1% potassium acetate, 1% bacteriological peptone, 0.5% yeast
extract, 0.17% yeast nitrogen base with ammonium sulfate and without amino
acids, 0.5% ammonium sulfate, 0.05 M potassium biphthalate) and grown for 16–
17 h. Meiosis was induced by transferring the SPS-cultured cells to pre-warmed
SPM (1% potassium acetate, 0.02% raffinose). The SPM-cultured cells were
harvested at various times after transfer. Auxin was added at a final concentration
of 4 mM.
458

Cytology

Chromosome spreads were immunostained as described previously [53,76]. Spheroplasts were burst in the presence of 1% paraformaldehyde and 0.1% lipsol. Stained samples were stained with primary and secondary antibodies observed using a fluorescence microscope (BX51; Olympus/Evident) with a 100× objective (NA1.3). Images were captured by CCD camera (CoolSNAP; Roper), and then processed using IP lab and/or iVision (Bio Vision Technologies) and Photoshop (Adobe) software. For focus counting, ~30 nuclei were counted at each time point.

Whole-cell staining was carried out as follows. Yeast cells are fixed with formaldehyde (3.7%) and placed on a glass slide coated with poly-lysine. The cells were treated with cold ethanol and acetone and stained with primary and secondary antibodies.

Western blotting

Western blotting was performed as described previously [27,77], using cell lysates extracted by the TCA method. After being harvested and washed twice with 20% TCA, cells were roughly disrupted with zirconia beads by the Multi-beads shocker (Yasui Kikai). The protein precipitate recovered by centrifugation was suspended in SDS-PAGE sample buffer adjusted to pH 8.8 and then incubated at 95 °C for 10 min. After electrophoresis, the proteins were transferred onto a nylon membrane (Immobilon-P, Millipore) and incubated with primary antibodies in a blocking buffer (1× PBS, 0.5% BSA) and then with alkaline phosphatase-conjugated secondary antibody (Promega). The color reaction was developed with NBT/BCIP solution (Nacalai Tesque).

CHEF analysis Pulsed-field gel electrophoresis

For pulsed-field gel electrophoresis (PFGE), chromosomal DNA was prepared in agarose plugs as described in [78] and run at 14 °C in a CHEF DR-III apparatus (BioRad) using the field 6V/cm at a 120° angle. Switching times followed a ramp from 15.1 to 25.1 seconds. Duration of electrophoresis was 41 h for chromosome III.

FACS analysis

Cells in 1 ml of SPM culture were collected by centrifugation. The pellet was then resuspended in 1 ml of 70% ethanol. Fixed cells were recollected and washed by suspending 1 ml of 50 mM sodium citrate buffer. The collected cells were

resuspended in 0.5 ml of sodium citrate buffer and sonicated for the 20s at room
temperature. Sonicated cells were incubated with 12.5 μl of 10 mg/mL RNAseA
for 1 h at 50 °C and in 25 μl of 15mg/ml proteinase K for 1h at 50 °C. 0.5ml of 16
μlg/mL PI solution was added and incubated for 30 min at room temperature in
the dark. The samples were then analyzed using SA-3800 Spectral analyzer
(Sony). Floreada.io was used for gating and plotting graphs.
501
502 **Statistics**
503 Means $\pm$ S.D values are shown. Datasets were compared using the Mann-
504 Whitney U-test. χ^2-test was used for proportion. Multiple test correction was done
505 with Bonferroni's correction. *, **, and *** show *P*-values of <0.05, <0.01 and
506 <0.001, respectively.
507

Acknowledgments
We are grateful to Drs. Neil Hunter for providing the yeast AID system and
Michael Lichten for anti-Ndt80. A.Sa. is a JICA fellow. Y.F. and M.I. were
supported by the Institute for Protein Research. This work was supported by a
grant from Institute for Fermentation, Osaka (IFO) and from Uehara Memorial
Foundation to Life Sciences; to A.Sh.

Competing interest
The authors declare no competing financial interest.
517

518 **Author contributions**
519 A.Sh. conceived and designed the experiments. A.Sa. carried out experiments
520 with help from Y.F. and M.I.. A.Sa, Y.F., M.I., and A.Sh. analyzed the data. A.Sh.
521 wrote the manuscript with inputs from help from others.
522
523

References

1   Bhat, K. P. & Cortez, D. RPA and RAD51: fork reversal, fork protection, and genome stability. *Nat Struct Mol Biol* **25**, 446-453 (2018). https://doi.org:10.1038/s41594-018-0075-z

2   Caldwell, C. C. & Spies, M. Dynamic elements of replication protein A at the crossroads of DNA replication, recombination, and repair. *Crit Rev Biochem Mol Biol* **55**, 482-507 (2020). https://doi.org:10.1080/10409238.2020.1813070

3   Wold, M. S. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Rev Biochem* **66**, 61-92 (1997). https://doi.org:10.1146/annurev.biochem.66.1.61

4   Brill, S. J. & Stillman, B. Replication factor-A from Saccharomyces cerevisiae is encoded by three essential genes coordinately expressed at S phase. *Genes Dev* **5**, 1589-1600 (1991). https://doi.org:10.1101/gad.5.9.1589

5   Chen, H., Lisby, M. & Symington, L. S. RPA coordinates DNA end resection and prevents formation of DNA hairpins. *Mol Cell* **50**, 589-600 (2013). https://doi.org:10.1016/j.molcel.2013.04.032

6   Deng, S. K., Chen, H. & Symington, L. S. Replication protein A prevents promiscuous annealing between short sequence homologies: Implications for genome integrity. *Bioessays* **37**, 305-313 (2015). https://doi.org:10.1002/bies.201400161

7   Ruff, P., Donnianni, R. A., Glancy, E., Oh, J. & Symington, L. S. RPA Stabilization of Single-Stranded DNA Is Critical for Break-Induced Replication. *Cell Rep* **17**, 3359-3368 (2016). https://doi.org:10.1016/j.celrep.2016.12.003

8   Sasanuma, H. *et al.* A new protein complex promoting the assembly of Rad51 filaments. *Nat Commun* **4**, 1676 (2013). https://doi.org:10.1038/ncomms2678

9   Shinohara, A. & Ogawa, T. Stimulation by Rad52 of yeast Rad51-mediated recombination. *Nature* **391**, 404-407 (1998). https://doi.org:10.1038/34943

10   Sung, P. Yeast Rad55 and Rad57 proteins form a heterodimer that functions with replication protein A to promote DNA strand exchange by Rad51 recombinase. *Genes Dev* **11**, 1111-1121 (1997).

https://doi.org:10.1101/gad.11.9.1111

11 Sung, P. Catalysis of ATP-dependent homologous DNA pairing and strand
exchange by yeast RAD51 protein. *Science* **265**, 1241-1243 (1994).
https://doi.org:10.1126/science.8066464

12 Sung, P. & Robberson, D. L. DNA strand exchange mediated by a RAD51-
ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell*
82, 453-461 (1995). https://doi.org:10.1016/0092-8674(95)90434-4

13 Shinohara, A., Shinohara, M., Ohta, T., Matsuda, S. & Ogawa, T. Rad52
forms ring structures and co-operates with RPA in single-strand DNA
annealing. *Genes Cells* **3**, 145-156 (1998). https://doi.org:10.1046/j.1365-
2443.1998.00176.x

14 Sugiyama, T., New, J. H. & Kowalczykowski, S. C. DNA annealing by
RAD52 protein is stimulated by specific interaction with the complex of
replication protein A and single-stranded DNA. *Proceedings of the National*
Academy of Sciences of the United States of America **95**, 6049-6054
(1998). https://doi.org:10.1073/pnas.95.11.6049

15 Friedel, A. M., Pike, B. L. & Gasser, S. M. ATR/Mec1: coordinating fork
stability and repair. *Curr Opin Cell Biol* **21**, 237-244 (2009).
https://doi.org:10.1016/j.ceb.2009.01.017

16 Williams, R. M., Yates, L. A. & Zhang, X. Structures and regulations of ATM
and ATR, master kinases in genome integrity. *Curr Opin Struct Biol* **61**, 98-
105 (2020). https://doi.org:10.1016/j.sbi.2019.12.010

17 Biswas, H., Goto, G., Wang, W., Sung, P. & Sugimoto, K. Ddc2ATRIP
promotes Mec1ATR activation at RPA-ssDNA tracts. *PLoS Genet* **15**,
e1008294 (2019). https://doi.org:10.1371/journal.pgen.1008294

18 Deshpande, I. *et al.* Structural Basis of Mec1-Ddc2-RPA Assembly and
Activation on Single-Stranded DNA at Sites of Damage. *Mol Cell* **68**, 431-
445 e435 (2017). https://doi.org:10.1016/j.molcel.2017.09.019

19 Yates, L. A., Tannous, E. A., Morgan, R. M., Burgers, P. M. & Zhang, X. A
DNA damage-induced phosphorylation circuit enhances Mec1(ATR)
Ddc2(ATRIP) recruitment to Replication Protein A. *Proceedings of the*
National Academy of Sciences of the United States of America **120**,
e2300150120 (2023). https://doi.org:10.1073/pnas.2300150120

20 Bass, T. E. *et al.* ETAA1 acts at stalled replication forks to maintain genome
integrity.   *Nat   Cell   Biol*   **18**,   1185-1195   (2016).
https://doi.org:10.1038/ncb3415

21 Haahr, P. *et al.* Activation of the ATR kinase by the RPA-binding protein
ETAA1. *Nat Cell Biol* **18**, 1196-1207 (2016).
https://doi.org:10.1038/ncb3422
22 Bergerat, A. *et al.* An atypical topoisomerase II from Archaea with
implications for meiotic recombination. *Nature* **386**, 414-417 (1997).
https://doi.org:10.1038/386414a0
23 Keeney, S., Giroux, C. N. & Kleckner, N. Meiosis-specific DNA double-
strand breaks are catalyzed by Spo11, a member of a widely conserved
protein family. *Cell* **88**, 375-384 (1997).
24 Bishop, D. K., Park, D., Xu, L. & Kleckner, N. DMC1: a meiosis-specific
yeast homolog of E. coli recA required for recombination, synaptonemal
complex formation, and cell cycle progression. *Cell* **69**, 439-456 (1992).
25 Shinohara, A., Ogawa, H. & Ogawa, T. Rad51 protein involved in repair
and recombination in S. cerevisiae is a RecA-like protein. *Cell* **69**, 457-470
(1992).
26 Cloud, V., Chan, Y. L., Grubb, J., Budke, B. & Bishop, D. K. Rad51 is an
accessory factor for Dmc1-mediated joint molecule formation during
meiosis. *Science* **337**, 1222-1225 (2012).
https://doi.org:10.1126/science.1219379
27 Hayase, A. *et al.* A protein complex containing Mei5 and Sae3 promotes
the assembly of the meiosis-specific RecA homolog Dmc1. *Cell* **119**, 927-
940 (2004).
28 Tsubouchi, H. & Roeder, G. S. The budding yeast mei5 and sae3 proteins
act together with dmc1 during meiotic recombination. *Genetics* **168**, 1219-
1230 (2004). https://doi.org:10.1534/genetics.103.025700
29 Hong, S. *et al.* The logic and mechanism of homologous recombination
partner choice. *Mol Cell* **51**, 440-453 (2013).
https://doi.org:10.1016/j.molcel.2013.08.008
30 Kim, K. P. *et al.* Sister cohesion and structural axis components mediate
homolog bias of meiotic recombination. *Cell* **143**, 924-937 (2010).
https://doi.org:10.1016/j.cell.2010.11.015
31 Niu, H. *et al.* Partner choice during meiosis is regulated by Hop1-promoted
dimerization of Mek1. *Molecular biology of the cell* **16**, 5804-5818 (2005).
https://doi.org:10.1091/mbc.e05-05-0465
32 Leem, S. H. & Ogawa, H. The MRE4 gene encodes a novel protein kinase
homologue required for meiotic recombination in Saccharomyces

632   cerevisiae. *Nucleic Acids Res* **20**, 449-457 (1992).
633   https://doi.org:10.1093/nar/20.3.449

634   33   Rockmill, B. & Roeder, G. S. A meiosis-specific protein kinase homolog
635   required for chromosome synapsis and recombination. *Genes Dev* **5**,
636   2392-2404 (1991). https://doi.org:10.1101/gad.5.12b.2392

637   34   Callender, T. L. *et al.* Mek1 Down Regulates Rad51 Activity during Yeast
638   Meiosis by Phosphorylation of Hed1. *PLoS Genet* **12**, e1006226 (2016).
639   https://doi.org:10.1371/journal.pgen.1006226

640   35   Niu, H. *et al.* Regulation of meiotic recombination via Mek1-mediated
641   Rad54 phosphorylation. *Mol Cell* **36**, 393-404 (2009).
642   https://doi.org:10.1016/j.molcel.2009.09.029

643   36   Jiang, H. *et al.* Direct association between the yeast Rad51 and Rad54
644   recombination proteins. *J Biol Chem* **271**, 33181-33186 (1996).
645   https://doi.org:10.1074/jbc.271.52.33181

646   37   Petukhova, G., Stratton, S. & Sung, P. Catalysis of homologous DNA
647   pairing by yeast Rad51 and Rad54 proteins. *Nature* **393**, 91-94 (1998).
648   https://doi.org:10.1038/30037

649   38   Tsubouchi, H. & Roeder, G. S. Budding yeast Hed1 down-regulates the
650   mitotic recombination machinery when meiotic recombination is impaired.
651   *Genes Dev* **20**, 1766-1775 (2006). https://doi.org:10.1101/gad.1422506

652   39   Borner, G. V., Hochwagen, A. & MacQueen, A. J. Meiosis in budding yeast.
653   *Genetics* (2023). https://doi.org:10.1093/genetics/iyad125

654   40   Cartagena-Lirola, H., Guerini, I., Manfrini, N., Lucchini, G. & Longhese, M.
655   P. Role of the Saccharomyces cerevisiae Rad53 checkpoint kinase in
656   signaling double-strand breaks during the meiotic cell cycle. *Molecular and*
657   *cellular biology* **28**, 4480-4493 (2008).
658   https://doi.org:10.1128/MCB.00375-08

659   41   Usui, T. & Shinohara, A. Rad9, a 53BP1 Ortholog of Budding Yeast, Is
660   Insensitive to Spo11-Induced Double-Strand Breaks During Meiosis. *Front*
661   *Cell Dev Biol* **9**, 635383 (2021). https://doi.org:10.3389/fcell.2021.635383

662   42   Carballo, J. A., Johnson, A. L., Sedgwick, S. G. & Cha, R. S.
663   Phosphorylation of the axial element protein Hop1 by Mec1/Tel1 ensures
664   meiotic interhomolog recombination. *Cell* **132**, 758-770 (2008).
665   https://doi.org:10.1016/j.cell.2008.01.035

666   43   Hollingsworth, N. M. Mek1/Mre4 is a master regulator of meiotic
667   recombination in budding yeast. *Microb Cell* **3**, 129-131 (2016).

https://doi.org:10.15698/mic2016.03.487

44   Xu, L., Ajimura, M., Padmore, R., Klein, C. & Kleckner, N. NDT80, a

meiosis-specific gene required for exit from pachytene in Saccharomyces

cerevisiae. *Molecular and cellular biology* **15**, 6572-6581 (1995).

https://doi.org:10.1128/MCB.15.12.6572

45   Hochwagen, A. Meiosis: making a synaptonemal complex just got easier.

Curr Biol **19**, R849-851 (2009). https://doi.org:10.1016/j.cub.2009.07.001

46   Sourirajan, A. & Lichten, M. Polo-like kinase Cdc5 drives exit from

pachytene during budding yeast meiosis. *Genes Dev* **22**, 2627-2632

(2008). https://doi.org:10.1101/gad.1711408

47   Sym, M., Engebrecht, J. A. & Roeder, G. S. ZIP1 is a synaptonemal

complex protein required for meiotic chromosome synapsis. *Cell* **72**, 365-

378 (1993).

48   Huang, Y. & Roig, I. Genetic control of meiosis surveillance mechanisms

in mammals. *Front Cell Dev Biol* **11**, 1127440 (2023).

https://doi.org:10.3389/fcell.2023.1127440

49   MacQueen, A. J. & Hochwagen, A. Checkpoint mechanisms: the puppet

masters of meiotic prophase. *Trends Cell Biol* **21**, 393-400 (2011).

https://doi.org:10.1016/j.tcb.2011.03.004

50   Borner, G. V., Kleckner, N. & Hunter, N. Crossover/noncrossover

differentiation, synaptonemal complex formation, and regulatory

surveillance at the leptotene/zygotene transition of meiosis. *Cell* **117**, 29-

45 (2004).

51   Storlazzi, A., Xu, L., Schwacha, A. & Kleckner, N. Synaptonemal complex

(SC) component Zip1 plays a role in meiotic recombination independent

of SC polymerization along the chromosomes. *Proceedings of the National*

Academy of Sciences of the United States of America **93**, 9043-9048

(1996). https://doi.org:10.1073/pnas.93.17.9043

52   Grushcow, J. M. *et al.* Saccharomyces cerevisiae checkpoint genes MEC1,

RAD17 and RAD24 are required for normal meiotic recombination partner

choice. *Genetics* **153**, 607-620 (1999).

https://doi.org:10.1093/genetics/153.2.607

53   Shinohara, M., Sakai, K., Ogawa, T. & Shinohara, A. The mitotic DNA

damage checkpoint proteins Rad17 and Rad24 are required for repair of

double-strand breaks during meiosis in yeast. *Genetics* **164**, 855-865

(2003). https://doi.org:10.1093/genetics/164.3.855

54 Shi, B. *et al.* Dual functions for the ssDNA-binding protein RPA in meiotic
recombination. *PLoS Genet* **15**, e1007952 (2019).
https://doi.org:10.1371/journal.pgen.1007952

55 Gasior, S. L., Wong, A. K., Kora, Y., Shinohara, A. & Bishop, D. K. Rad52
associates with RPA and functions with rad55 and rad57 to assemble
meiotic recombination complexes. *Genes Dev* **12**, 2208-2221 (1998).
https://doi.org:10.1101/gad.12.14.2208

56 Raina, V. B. & Vader, G. Homeostatic Control of Meiotic Prophase
Checkpoint Function by Pch2 and Hop1. *Curr Biol* **30**, 4413-4424 e4415
(2020). https://doi.org:10.1016/j.cub.2020.08.064

57 Shinohara, A., Gasior, S., Ogawa, T., Kleckner, N. & Bishop, D. K.
Saccharomyces cerevisiae recA homologues RAD51 and DMC1 have
both distinct and overlapping roles in meiotic recombination. *Genes Cells*
2, 615-629 (1997). https://doi.org:10.1046/j.1365-2443.1997.1480347.x

58 Bishop, D. K. RecA homologs Dmc1 and Rad51 interact to form multiple
nuclear complexes prior to meiotic chromosome synapsis. *Cell* **79**, 1081-
1092 (1994).

59 Chen, X. *et al.* Mek1 coordinates meiotic progression with DNA break
repair by directly phosphorylating and inhibiting the yeast pachytene exit
regulator Ndt80. *PLoS Genet* **14**, e1007832 (2018).
https://doi.org:10.1371/journal.pgen.1007832

60 Hollingsworth, N. M., Goetsch, L. & Byers, B. The HOP1 gene encodes a
meiosis-specific component of yeast chromosomes. *Cell* **61**, 73-84 (1990).
https://doi.org:10.1016/0092-8674(90)90216-2

61 Smith, A. V. & Roeder, G. S. The yeast Red1 protein localizes to the cores
of meiotic chromosomes. *J Cell Biol* **136**, 957-967 (1997).
https://doi.org:10.1083/jcb.136.5.957

62 Kniewel, R. *et al.* Histone H3 Threonine 11 Phosphorylation Is Catalyzed
Directly by the Meiosis-Specific Kinase Mek1 and Provides a Molecular
Readout of Mek1 Activity in Vivo. *Genetics* **207**, 1313-1333 (2017).
https://doi.org:10.1534/genetics.117.300359

63 Hollingsworth, N. M. & Gaglione, R. The meiotic-specific Mek1 kinase in
budding yeast regulates interhomolog recombination and coordinates
meiotic progression with double-strand break repair. *Curr Genet* **65**, 631-
641 (2019). https://doi.org:10.1007/s00294-019-00937-3

64 Alani, E., Padmore, R. & Kleckner, N. Analysis of wild-type and rad50

mutants of yeast suggests an intimate relationship between meiotic
chromosome synapsis and recombination. *Cell* **61**, 419-436 (1990).
https://doi.org:10.1016/0092-8674(90)90524-i

65　Usui, T., Ogawa, H. & Petrini, J. H. A DNA damage response pathway
controlled by Tel1 and the Mre11 complex. *Mol Cell* **7**, 1255-1266 (2001).
https://doi.org:10.1016/s1097-2765(01)00270-2

66　Zou, L. & Elledge, S. J. Sensing DNA damage through ATRIP recognition
of RPA-ssDNA complexes. *Science* **300**, 1542-1548 (2003).
https://doi.org:10.1126/science.1083430

67　Lin, Y. *et al.* APE1 recruits ATRIP to ssDNA in an RPA-dependent and -
independent manner to promote the ATR DNA damage response. *Elife* **12**
(2023). https://doi.org:10.7554/eLife.82324

68　Shinohara, M., Bishop, D. K. & Shinohara, A. Distinct Functions in
Regulation of Meiotic Crossovers for DNA Damage Response Clamp
Loader Rad24(Rad17) and Mec1(ATR) Kinase. *Genetics* **213**, 1255-1269
(2019). https://doi.org:10.1534/genetics.119.302427

69　Hustedt, N. *et al.* Yeast PP4 interacts with ATR homolog Ddc2-Mec1 and
regulates checkpoint signaling. *Mol Cell* **57**, 273-289 (2015).
https://doi.org:10.1016/j.molcel.2014.11.016

70　Li, K., Yoshimura, K. & Shinohara, M. Meiotic DNA double-strand break-
independent role of protein phosphatase 4 in Hop1 assembly to promote
meiotic chromosome axis formation in budding yeast. *Genes Cells* (2023).
https://doi.org:10.1111/gtc.13049

71　San-Segundo, P. A. & Roeder, G. S. Pch2 links chromatin silencing to
meiotic checkpoint control. *Cell* **97**, 313-324 (1999).
https://doi.org:10.1016/s0092-8674(00)80741-2

72　Borner, G. V., Barot, A. & Kleckner, N. Yeast Pch2 promotes domainal axis
organization, timely recombination progression, and arrest of defective
recombinosomes during meiosis. *Proceedings of the National Academy of*
Sciences of the United States of America **105**, 3327-3332 (2008).
https://doi.org:10.1073/pnas.0711864105

73　Herruzo, E. *et al.* The Pch2 AAA+ ATPase promotes phosphorylation of
the Hop1 meiotic checkpoint adaptor in response to synaptonemal
complex defects. *Nucleic Acids Res* **44**, 7722-7741 (2016).
https://doi.org:10.1093/nar/gkw506

74　Humphryes, N. *et al.* The Ecm11-Gmc2 complex promotes synaptonemal

complex formation through assembly of transverse filaments in budding
yeast. *PLoS Genet* **9**, e1003194 (2013).
https://doi.org:10.1371/journal.pgen.1003194

75   Lee, M. S. *et al.* The synaptonemal complex central region modulates
crossover pathways and feedback control of meiotic double-strand break
formation. *Nucleic Acids Res* **49**, 7537-7553 (2021).
https://doi.org:10.1093/nar/gkab566

76   Shinohara, M., Gasior, S. L., Bishop, D. K. & Shinohara, A. Tid1/Rdh54
promotes colocalization of rad51 and dmc1 during meiotic recombination.
Proceedings of the National Academy of Sciences of the United States of
America **97**, 10814-10819 (2000).
https://doi.org:10.1073/pnas.97.20.10814

77   Shinohara, M., Oh, S. D., Hunter, N. & Shinohara, A. Crossover assurance
and crossover interference are distinctly regulated by the ZMM proteins
during yeast meiosis. *Nat Genet* **40**, 299-309 (2008).
https://doi.org:10.1038/ng.83

78   Bani Ismail, M., Shinohara, M. & Shinohara, A. Dot1-dependent histone
H3K79 methylation promotes the formation of meiotic double-strand
breaks in the absence of histone H3K4 methylation in budding yeast. *PLoS*
One **9**, e96648 (2014). https://doi.org:10.1371/journal.pone.0096648

796
797

 Figure legends

Figure 1. A pathway of recombination checkpoint in meiosis.

Figure 2. Rfa1 depletion alleviates *dmc1*-induced pachytene arrest

(A) Expression of various proteins in meiosis. Lysates obtained from the *dmc1*
RFA1-AID (SAY63/64) cells at various time points during meiosis with or
without the addition of auxin at 4 h were analyzed by western blotting using
anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-histone H3-phospho-T11,
anti-Ndt80, anti-Cdc5 or anti-tubulin (lower) antibodies.

(B) Rfa2 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with
or without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and
DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.

(C) Kinetics of Rfa2 assembly/disassembly. The number of Rfa2-positive cells
(with more than 5 foci) was counted at each time point. At each time point,
more than 100 cells were counted.

(D) Meiotic cell cycle progression. The entry into meiosis I and II in *dmc1 RFA1-*
AID (SAY63/64) cells were analyzed by DAPI staining. The number of DAPI
bodies in a cell was counted. A cell with 2, 3, and 4, and 3 and 4 DAPI bodies
was defined as a cell that passed through meiosis I and meiosis II,
respectively. The graph shows the percentages of cells that completed MI or
MII at the indicated time points. More than 200 cells were counted at each
time point. The representative results (*n*=3) are shown. *dmc1 RFA1-AID*
without auxin, blue closed circles; *dmc1 RFA1-AID* with auxin addition at 4 h,
red closed circles.

(E) FACS profile of the *dmc1 RFA1-AID* mutant without or with auxin addition at
4 h. Fixed cells at each time point were stained with PI and analyzed by FACS
cell sorter.

(F) Tubulin staining in the *dmc1 RFA1-AID* mutant. Whole cells were fixed and
stained with anti-tubulin (green) and DAPI (blue). Representative images for
each class are shown. Bar = 2 μm.

(G) Whole cells were fixed and stained with anti-tubulin (green) and DAPI (blue).
Representative images at 8 h of *dmc1 RFA1-AID* (SAY63/64) cells with or
without the addition of auxin at 4 h are shown. Bar = 2 μm.

(H) Classification of tubulin staining at each time point of meiosis in *dmc1 RFA1-*

AID (SAY63/64) cells with (right) or without (left) the addition of auxin at 4 h.

Dot, short line, and long line tubulin-staining with single DAPI mass were

defined as prophase I (blue), metaphase I (red), anaphase I (orange),

metaphase II (blue), anaphase II (green) and post-meiosis II (pale green).

Short and long tubulin-staining were defined as metaphase II and anaphase

II, respectively. At each time point, more than 100 cells were counted. The

representative results are shown (*n*=3).

Figure 3. Rfa1 depletion does not affect DSB turnovers in *dmc1* mutant

(A) CHEF analysis of meiotic DSB repair. Chromosomal DNAs from *dmc1 RFA1-*

AID (SAY63/64) cells with or without the addition of auxin at 4 h were studied

by CHEF electrophoresis. Auxin was added at 4 h in the incubation with SPM.

(B) Rad51 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with

or without the addition of auxin at 4 h were stained with anti-Rad51 (red), and

DAPI (blue). Representative images at each time point under the two

conditions are shown. Bar = 2 μm.

(C) Kinetics of Rad51 assembly/disassembly. The number of Rad51-positive cells

(with more than 5 foci) was counted at each time point. At each time point,

more than 100 cells were counted.

(D) Hop1 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with

or without the addition of auxin at 4 h were stained with anti-Hop1 (red), and

DAPI (blue). Representative images at each time point under the two

conditions are shown. Bar = 2 μm.

(E) Kinetics of Hop1 assembly/disassembly. The number of Rad51-positive cells

(with more than 5 foci) was counted at each time point. At each time point,

more than 100 cells were counted.

Figure 4. *NDT80* is necessary for pachytene exit by Rfa1 depletion

(A) Expression of various protein in meiosis. Lysates obtained from the *ndt80*

dmc1 RFA1-AID (SAY80/81) cells at various time points during meiosis with

or without the addition of auxin at 4 h were analyzed by western blotting using

anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5 or anti-tubulin (lower)

antibodies.

(B) Rfa2 staining. Nuclear spreads from *ndt80 dmc1 RFA1-AID* (SAY80/81) cells

with or without the addition of auxin at 4 h were stained with anti-Rfa2 (red),

and DAPI (blue). Representative images at each time point under the two

conditions are shown. Bar = 2 μm.

(C) Classification of tubulin staining at each time point of meiosis in *ndt80 dmc1*
RFA1-AID (SAY80/81) cells with (right) or without (left) the addition of auxin at
4 h. Dot, short line, and long line tubulin-staining with single DAPI mass were
defined as prophase I (blue), metaphase I (red), anaphase I (orange),
metaphase II (blue), anaphase II (green) and post-meiosis II (pale green).
Short and long tubulin-staining were defined as metaphase II and anaphase
II, respectively. At each time point, more than 100 cells were counted. The
representative results are shown (*n*=2).

(D) Hop1 staining. Nuclear spreads from *ndt80 dmc1 RFA1-AID* (SAY SAY80/81)
cells with or without the addition of auxin at 4 h were stained with anti-Hop1
(red), and DAPI (blue). Representative images at each time point under the
two conditions are shown. Bar = 2 μm.

Figure 5. Rfa1 depletion alleviates *zip1*-induced delay in meiosis I

(A) Expression of various protein in meiosis. Lysates obtained from the *zip1*
RFA1-AID (SAY72/73) cells at various time points during meiosis with or
without the addition of auxin at 4 h were analyzed by western blotting using
anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5, anti-histone H3-
phospho-T11 or anti-tubulin (lower) antibodies.

(B) Meiotic cell cycle progression. The entry into meiosis I and II in *zip1 RFA1-*
AID (SAY72/73) cells were analyzed by DAPI staining as shown in Figure 1D.
More than 200 cells were counted at each time point. The representative
results (*n*=3) are shown. *dmc1 RFA1-AID* without auxin, blue closed circles;
dmc1 RFA1-AID with auxin addition at 4 h, red closed circles.

(C) Rfa2 staining. Nuclear spreads from *zip1 RFA1-AID* (SAY72/73) cells with or
without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and DAPI
(blue). Representative images at each time point under the two conditions are
shown. Bar = 2 μm.

(D) Tubulin staining in the *zip1 RFA1-AID* mutant. Whole cells were fixed and
stained with anti-tubulin (green) and DAPI (blue). Representative images for
each class are shown. Bar = 2 μm.

(E) Classification of tubulin staining at each time point of meiosis in *zip1 RFA1-*
AID (SAY72/73) cells with (right) or without (left) the addition of auxin at 4 h.
The quantification was performed as shown in Figure 1G. At each time point,
more than 100 cells were counted. The representative results are shown (*n*=2).

(F) Hop1 staining. Nuclear spreads from *zip1 RFA1-AID* (SAY72/73) cells with or
without the addition of auxin at 4 h were stained with anti-Hop1 (red), and
DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.
(G) CHEF analysis of meiotic DSB repair. Chromosomal DNAs from *zip1 RFA1-*
AID (SAY72/73) cells with or without the addition of auxin at 4 h were studied
by CHEF electrophoresis. Auxin was added at 4 h in the incubation with SPM.
(H) Kinetics of assembly/disassembly of Rfa2, Rad51, and Hop1. The number of
focus-positive cells (with more than 5 foci) was counted at each time point. At
each time point, more than 100 cells were counted.
916

Figure 6. Rfa1 depletion does not suppress *rad50S*-induced delay in
meiosis I
(A) Expression of various proteins in meiosis. Lysates obtained from the *rad50S*
RFA1-AID (SAY68/69) cells at various time points during meiosis with or
without the addition of auxin at 4 h were analyzed by western blotting using
anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5 or anti-tubulin (lower)
antibodies.
(B) Classification of tubulin staining at each time point of meiosis in *rad50S RFA1-*
AID (SAY68/69) cells with (right) or without (left) the addition of auxin at 4 h.
The quantification was performed as shown in Figure 1G. At each time point,
more than 100 cells were counted. The representative results are shown (*n*=1).
(C) CHEF analysis of meiotic DSB repair. Chromosomal DNAs from *zip1 RFA1-*
AID (SAY68/69) cells with or without the addition of auxin at 4 h were studied
by CHEF electrophoresis. Auxin was added at 4 h in the incubation with SPM.
(D) Rfa2 staining. Nuclear spreads from *rad50S RFA1-AID* (SAY68/69) cells with
or without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and
DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.
(E) Hop1 staining. Nuclear spreads from *rad50S RFA1-AID* (SAY68/69) cells with
or without the addition of auxin at 4 h were stained with anti-Hop1 (red), and
DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.
939

Supplemental Figure S1.
(A) The number of Rfa2 foci per spread in *dmc1 RFA1-AID* (SAY72/73) cells with

(red) or without (blue) the addition of auxin at 4 h was counted at different time

points. The mean and SD are shown in the plot.

(B) The number of Rad51 foci per spread *dmc1 RFA1-AID* (SAY72/73) cells with

(red) or without (blue) the addition of auxin at 4 h was counted at different time

points. The mean and SD are shown in the plot.

(C) The number of Rfa2 foci per spread in *zip1 RFA1-AID* (SAY72/73) cells with

(red) or without (blue) the addition of auxin at 4 h was counted at different time

points. The mean and SD are shown in the plot.

(D) The number of Rad51 foci per spread *zip1 RFA1-AID* (SAY72/73) cells with

(red) or without (blue) the addition of auxin at 4 h was counted at different time

points. The mean and SD are shown in the plot.

Supplemental Figure S 2.

CHEF analysis of meiotic DSB repair. Chromosomal DNAs from Wild type

(MSY831/833) and the *dmc1* mutant cells were studied by CHEF electrophoresis.

Auxin was added at 4 h in the incubation with SPM.

 Supplementary Information

960 **Table S1. Strain List**

Strain number	Genotypes
MSY831/833	*MATa/ α, ho::LYS2/", lys2/", ura3/", leu2::hisG/", trp1::hisG/"*
NKY1551	*MATa/ α, ho::LYS2/", lys2/", ura3/", leu2::hisG/", his4X-LEU2(BamHI)-URA3/his4B-LEU2(MluI), arg4-bgl/arg4-nsp*
SAY15/16	MSY831/833 with *Rfa1-AID::HYG/", p-CUP1-Os-Tir1::URA3*
SAY63/64	SAY15/16 with *dmc1::URA3/"*
SAY68/69	SAY15/16 with *rad50S::LEU2/"*
SAY72/73	SAY15/16 with *zip1::LEU2/"*
SAY80/81	SAY15/16 with *ndt80::LEU2/"*

962 **Table S2. Oligonucleotide list**

963 **Table S2. Primer List**

Primer name	Sequence
Rfa1-AID-Hygro-F	5'-TTG AAT TAC AGG GCT GAA GCC GAC TAT CTT GCC GAT GAG TTA TCC AAG GCT TTG TTA GCT CGT ACG CTG CAG GTC GAC
Rfa1-AID-Hygro-R	5'-TTT TTT TTT TAC ATT TCT CAT ATG TTA CAT AGA TTA AAT AGT ACT TGA TTA TTT GAT ACA ATC GAT GAA TTC GAG CTC G
RFA1-F	5'-GTG AGA AGT GCG ACA CCA ATA
RFA1-R	5'-GTC GAC TAT TTG GAG AAG GAA G

Figure 1. Arivarasan S. et al.

Figure 2. Arivarasan S. et al.

Figure 3. Arivarasan S. et al.

Figure 4. Arivarasan S. et al.

Figure 5. Arivarasan S. et al.

Figure 6. Arivarasan S. et al.

Supplementary Figure S1. Arivarasan S. et al.

Supplementary Figure S2. Arivarasan S. et al.