1   **Replication protein-A, RPA, plays a pivotal role in the maintenance of**
2   **recombination checkpoint in yeast meiosis**
3

4   **Arivarasan Sampathkumar, Zhong Chen, Yuting Tang, Yurika Fujita,**
5   **Masaru Ito, and Akira Shinohara***
6
7   Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan
8

9   Running title: Essential role of RPA in recombination checkpoint
10
11 Keywords: meiosis; homologous recombination; RPA; recombination checkpoint;
12 Rfa1, auxin degron, synapsis checkpoint
13
14
15

16 *Corresponding author:
17 Akira Shinohara; ashino@protein.osaka-u.ac.jp
18
Lead contact:
Akira Shinohara
Institute for Protein Research, Osaka University,
3-2 Yamadaoka, Suita, Osaka 565-0871 JAPAN
Phone: 81-6-6879-8624
FAX: 81-6-6879-8626
E-mail: ashino@protein.osaka-u.ac.jp
ORCID: 0000-0003-4207-8247
27

Summary

DNA double-strand breaks (DSBs) activate DNA damage responses (DDR) in

both mitotic and meiotic cells. A single-stranded DNA (ssDNA) binding protein,

Replication protein-A (RPA) binds to the ssDNA formed at DSBs to activate

ATR/Mec1 kinase for the response. Meiotic DSBs induce homologous

recombination monitored by a meiotic DDR called the recombination checkpoint

that blocks the pachytene exit in meiotic prophase I. In this study, we showed the

essential role of RPA in the maintenance of the recombination checkpoint during

S. cerevisiae meiosis. The depletion of an RPA subunit, Rfa1, in a recombination-

defective *dmc1* mutant, fully alleviates the pachytene arrest with the persistent

unrepaired DSBs. RPA depletion decreases the activity of a meiosis-specific

CHK2 homolog, Mek1 kinase, which in turn activates the Ndt80 transcriptional

regulator for pachytene exit. These support the idea that RPA is a sensor of

ssDNAs for the activation of meiotic DDR. Rfa1 depletion also accelerates the

prophase I delay in the *zip1* mutant defective in both chromosome synapsis and

the recombination, suggesting that the accumulation of ssDNAs rather than

defective synapsis triggers prophase I delay in the *zip1* mutant.

Introduction

A eukaryotic single-stranded (ss)DNA binding protein, Replication protein-A

(RPA), plays an essential role in DNA replication, DNA repair, and homologous

recombination [1-3]. RPA consists of three subunits; RPA70(RPA1), RPA32(RPA2),

and RPA14(RPA3), which are highly conserved from yeast to humans. In *S.*

cerevisiae, the knockout of either one of three genes (*RFA1*, *-2* and *-3*) encoding

the RPA subunits leads to lethality, consistent with its essential role in DNA

replication [4]. RPA also plays a role in DNA damage response (DDR) [3,5].

In homologous recombination. RPA binds to ssDNAs to prevent the

formation of the secondary structure of ssDNAs and to protect ssDNAs from

degradation by various nucleases [6]. The assembly of Rad51 filaments on RPA-

coated ssDNA is assisted by Rad51 mediators such as Rad52 in the budding

yeast [7-9]. RPA assists Rad51-mediated homology search and strand exchange

with a homologous double-stranded (ds)DNA [10]. The invasion of the ssDNA into

the homologous dsDNA results in the formation of a displacement loop (D-loop).

The 3'-OH end of the invading strand provides a substrate for DNA synthesis,

which is followed by further processing and engaging with the ssDNA in the other

DSB end to form a joint molecule. The joint molecule is finally matured into a

recombination product. RPA also works in recombination-associated DNA
synthesis as well as the annealing in the second-end capture.

DNA double-strand break (DSB) activates the DDR. DSB ends stimulate a
DDR kinase, ATM/Tel1 (Ataxia telangiectasia mutated) while ssDNAs do an ATM
homolog, ATR/Mec1 (Ataxia telangiectasia mutated related), both of which
belong to the PI3 kinase family [11,12]. The activation of ATR is mediated by the
interaction with an ATR partner, ATRIP/Ddc2, which directly interacts with RPA
bound to ssDNAs [13-15]. In mammals, ATR activation is also mediated by the
second RPA-binding protein, ETAA1 [16,17]. Thus, RPA is considered to work as an
essential DNA damage sensor to ssDNAs as well as a DNA damage processing
protein.

Meiosis rewires DNA transactions such as homologous recombination/DSB
repair as well as DDR to guarantee the faithful segregation of two paternal
homologous chromosomes at meiosis I for the reduction of chromosome
numbers in gametes. In budding yeast, meiotic recombination is initiated by the
DSB formation at recombination hotspots by the TopoVI-like protein, Spo11,
which is followed by the production of ssDNAs [18,19]. The ssDNA is used by a
meiosis-specific Rad51 homolog, Dmc1 [20], for D-loop formation, which is aided
by Rad51 [21] and its mediator, Mei5-Sae3 complex [22,23]. Further processing
generates a recombination intermediate with double-Holliday junctions (dHJs),
which are specifically resolved into crossovers between homologs. Importantly,
the partner choice in the recombination is mediated by meiosis-specific
chromosome-axis proteins (Hop1, Red1, and Mek1/Mre4), promoting
interhomolog bias via an unknown mechanism [24-26]. Mek1/Mre4 (hereafter Mek1),
a meiosis-specific homolog of Rad53/CHK2 kinase [27,28], suppresses Rad51-
mediated recombination in meiosis [25,26]. This suppression occurs through the
Mek1-mediated phosphorylation of two proteins, Rad54 and Hed1 [29,30]. Rad54
belongs to the Swi2/Snf2 family protein essential for Rad51-mediated strand
invasion [31]. Mek1-mediated phosphorylation of Rad54 abolishes its interaction
with Rad51 [30]. Hed1, which is expressed only in meiosis, inhibits Rad51 by
binding to the Rad51-ssDNA complex [32]. The inhibitory activity of Hed1 against
Rad51 and the stability of Hed1 are augmented by the Mek1-mediated
phosphorylation [29].

Meiotic events such as recombination and chromosome synapsis are strictly
coupled with each other (Fig. 1) [33]. Similar to mitosis, DSBs in yeast meiosis
activate Tel1(ATM) and Mec1(ATR). Different from mitotic DNA damage

checkpoint, downstream events of Mec1 or Tel1 activation such as the Rad53
activation are largely suppressed in meiotic cells with poorly-defined mechanisms
[34,35]. On the other hand, Mec1 and/or Tel1 phosphorylates Hop1 on chromosome
axis [36]. Together with chromosomal-bound Red1, phosphorylated Hop1 activates
Mek1 kinase, whose autophosphorylation provides positive feedback on its
activity [37]. Mek1-mediated phosphorylation promotes the inactivation of the Ndt80
transcription factor [38] essential for the exit from the pachytene stage, where
chromosome synapsis and the formation of the dHJs are completed [37,39]. By
activating Ndt80 (through inactivating Mek1), meiotic cells exit the pachytene
stage by expressing downstream targets such as Cdc5 Polo-like kinase (PLK)
and a cyclin, Clb1 [40] for the resolution of dHJs and the disassembly of the
synaptonemal complex (SC), a meiosis-specific tripartite chromosome structure,
an indicative of chromosome synapsis.

It is widely accepted that meiotic cells surveil not only homologous
recombination but also chromosome synapsis, referred to as the "recombination"
and "synapsis" checkpoints (Fig. 1). Like a recombination-defective *dmc1* mutant,
the budding yeast *zip1* mutant, which is defective in the assembly of the central
region of the SCs, thus synapsis shows a delay in meiosis progression [41]. In
nematodes and mice, synapsis-defective mutants also delay the meiotic
prophase I progression and/or induce apoptosis [42,43]. In yeast, chromosome
synapsis is tightly coupled with meiotic recombination: e.g. a recombination-
deficient mutant also displays abnormal synapsis [44,45]; the *zip1* mutant completely
defective in SC formation also impairs the processing of meiotic recombination
intermediates. This complicates the presence of the synapsis checkpoint.
Moreover, DDR proteins also play a role in meiotic recombination, which also
provides a caveat in the interpretation of the meiotic phenotypes of DDR mutants
[46-48].

The role of RPA in DDR is widely accepted. In a pioneer work [5], the depletion
of RPA70 in human cell lines reduced the ATRIP/Ddc2 binding to the DNA
damage as well as the activation of downstream events. In the same study, it was
also shown that the depletion of Rfa1 in mitotic yeast cells by a temperature-
induced degron reduced the binding of Ddc2 to a single DSB. On the other hand,
the other report showed that Rfa1 depletion in G2/M-arrested mitotic cells in
budding yeast by the degron does not affect checkpoint activation to a single DSB
[6]. A point mutant in the *RFA1* gene encoding RPA70 subunit of yeast RPA called
rfa1-t11 with K45E substitution in the N-terminal protein-protein interaction

domain is partially defective in the DDR in the response to telomere dysfunction
[49]. Rfa1-t11 showed normal binding to the DSB, but the mutant decreased
recruitment of Ddc2 to the DSB [5]. In meiosis, the *rfa1-t11* with reduced Ddd2-
focus formation in nuclei weakly suppresses the *dmc1*-induced delay in meiosis
progression [50] while RPA containing Rfa1-t11 showed normal localization on
meiotic chromosomes [51]. The *rfa1-t11* mutant alters recombination kinetics [51],
which may indirectly affect the checkpoint. Furthermore, the conditional depletion
of Rpa1 in mouse spermatocytes eliminated RPA foci on meiotic chromosomes
but is proficient in ATR recruitment to the chromosomes [52]. Therefore, we still
need careful evaluation of the role of RPA in the DDR in meiotic cells.
In this study, we revisited the role of RPA in the meiotic recombination
checkpoint by using a conditional-depletion mutant of Rfa1. The depletion of Rfa1
in the *dmc1*-induced pachytene arrest facilitates the pachytene exit, thus entry
into meiosis I and II. Importantly, the RPA depletion did not change the
accumulation unrepaired meiotic DSBs in the *dmc1*, which causes abnormal
segregation of chromosomes in the division. This confirms the essential role of
RPA in the maintenance of the meiotic recombination checkpoint. Moreover, we
found that Rfa1 depletion also alleviates delayed meiosis progression in the *zip1*
mutant. This suggests that the *zip1* mutation mainly triggers an RPA-dependent
response of the prophase I delay, thus the recombination checkpoint rather than
the synapsis checkpoint.
157
Results
Depletion of Rfa1 alleviates *dmc1*-induced arrest of meiosis
Three RPA subunits, Rfa1, -2, and -3, are essential for the viability of budding
yeast cells [4]. To know the role of RPA in the DDR in yeast meiosis, we constructed
a conditional depletion mutant of the *RFA1* gene encoding the largest RPA
subunit, Rfa1, by the auxin-induced degron (AID) system [53]. The AID tag was
added to the C-terminal of the Rfa1 protein. The addition of auxin as well as
copper, which induced Tir1 F-box protein, in the medium largely inhibited the
mitotic growth of *RFA1-AID* cells (Supplemental Fig. 1a), consistent with the
previous study [4]. *RFA1-AID* diploid cells were induced for the entry of meiosis by
incubating in the sporulation medium (SPM). When auxin was added at 4 h
incubation with SPM after pre-meiotic DNA replication, which induced the
disappearance of Rfa1 from 5 h (Supplemental Fig. 1b), the cells formed few
spores after 24 h while more than 90% of the control wild-type cells sporulated

(Supplemental Fig. 1c). Most of the Rfa1-depleted cells showed abnormal
nuclear morphology (Supplemental Fig. 1c). Indeed, rarely formed tetrad of the
cells are all dead (0 viable spores in 100 tetrads) while without the auxin treatment,
the viability of spores from *RFA1-AID* cells was 91.8% (among 100 tetrads). The
pulse-field gel electrophoresis revealed the accumulation of meiotic DSBs on
chromosomes (Supplemental Fig. 1d), confirming a defect in DSB repair [51].
These show that the AID-tag did not interfere with Rfa1 function in both miotic
and meiotic cells and that Rfa1 is essential for both mitosis and meiosis.
To know the role of RPA in the recombination checkpoint in meiosis, we
used the *dmc1* mutant, which is unable to repair meiotic DSBs and, as a result,
shows an arrest at mid-pachytene stage during meiotic prophase I [20]. This
response to unrepaired DSBs is referred to as the recombination checkpoint.
RFA1-AID was introduced in the *dmc1* deletion mutant; *dmc1 RFA1-AID*. Like the
dmc1 mutant, without auxin addition, *dmc1 RFA1-AID* shows an arrest in meiotic
prophase I without division (e.g. Fig. 2d), indicating that the Rfa1 tagging did not
affect the recombination checkpoint in the absence of Dmc1. RPA is also involved
in DNA replication and S-phase checkpoint [1,2], which indirectly affects the G2
event. To avoid the effect in pre-meiotic DNA replication, we depleted Rfa1 protein
by adding auxin to a meiotic culture of *dmc1 RFA1-AID* cells at 4 h after the
induction of meiosis in the sporulation medium (SPM), thus, after pre-meiotic DNA
replication, which occurs at 1.5-3 h. At 5 h, after the 1-hour addition of auxin, the
protein level of Rfa1-AID was decreased to ~15% level relative to the original
level at 4 h (Fig. 2a, Supplementary Figs. S1e and S2), indicating the efficient
depletion of Rfa1 under the condition. We also checked the focus formation of
Rfa2, the second subunit of RPA, on meiotic chromosome spreads. Rfa2 shows
punctate staining on the spreads with other recombination proteins such as
Rad51 [54], which corresponds with ongoing recombination (Fig. 2b). In the *dmc1*
mutant, Rfa2 focus-staining appears at 2.5 h, reaches a plateau at 5 h, and
persists after that, indicating little turnover of Rfa2 due to stalled recombination in
the mutant (Fig. 2c, Supplementary Fig. S3a). When auxin was added at 4 h,
Rfa2 foci disappeared within one hour, and Rfa2 foci were rarely observed during
further incubation by 10 h (Fig. 2b, c). This suggests that RPA is highly dynamic
and easily dissociated from the chromosome. We concluded that Rfa1 depletion
at 4 h in meiosis eliminates most chromosome-bound RPA complexes by 5 h.
206
Checkpoint alleviation in Rfa1 depletion in *dmc1* meiosis

DAPI staining of DNAs confirmed a single DAPI body in a *dmc1 RFA1-AID* cell
even at 24 h, thus an arrest in prophase I as shown previously [20]. Rfa1 depletion
at 4 h induced fragmentation of DAPI bodies at late time points. In some cells, 2-
4 irregular-sized DAPI bodies with smaller DAPI bodies were detected in a single
cell (Fig. 2d). The cells with abnormal DAPI bodies appeared at 6 h and gradually
accumulated by 24 h (Fig. 2e). This suggests the presence of nuclear division,
thus the onset of meiosis I and/or II in the *dmc1* cells upon Rfa1 depletion. We
also checked the effect of Rfa1 depletion on meiotic DNA replication. The FACS
analysis showed that the addition of auxin at 4 h did not affect the transition from
2C to 4C (Supplementary Fig. S1f).
To evaluate the meiosis progression precisely, we examined the status of
spindle morphology in the *dmc1 RFA1-AID* cells in the absence or presence of
auxin (Fig. 2f, g). Tubulin staining revealed a single focus associated with single
or multiple short projections in meiotic prophase I of the cell in the absence of
auxin (Fig. 2f). The *dmc1 RFA1-AID* mutant maintained this arrangement of
spindles in the absence of auxin by 24 h, consistent with an arrest of the *dmc1*
mutant in prophase I (Fig. 2f). On the other hand, the Rfa1 depletion induced the
elongation of spindles after one hour of addition (at 5 h). At 6 h, ~50% *dmc1*
RFA1-AID cells contained single short and long spindles corresponding with
metaphase I and anaphase I, respectively (Fig. 2e). At 8 h, in addition to the
metaphase I and anaphase I spindle, ~70% of cells contained metaphase II or
anaphase II spindles (Fig. 2f, g). At 10 h, most of the cells showed diffuse staining
of microtubules in multiple DAPI bodies in a cell. These show that Rfa1 depletion
almost alleviates meiotic prophase I arrest induced by the *dmc1* mutation.
Cdc5/PLK is expressed after the pachytene exit [40,55]. Like *dmc1* cells,
without auxin addition, the *dmc1 RFA1-AID* mutant cells did not express Cdc5
(Fig. 2a). On the other hand, the addition of auxin at 4 h induced the expression
of Cdc5 from 5 h. This confirms the exit of the pachytene stage of *dmc1* cells in
the absence of RPA.
237
No DSB repair in *dmc1* meiosis with Rfa1 depletion
One possibility to suppress the *dmc1* arrest is to channel unrepaired DSBs into
the repair pathway. On the other hand, the acute depletion of RPA did not affect
the status of meiotic DSBs under *dmc1* arrest. The CHEF (Counter-clamped
homogeneous electric field) gel electrophoresis confirmed the accumulation of
fragmented chromosomes in the *dmc1* mutant irrespective of the presence or

absence of RPA. In the *dmc1 RFA1-AID* mutant without auxin treatment, smear
bands of full-length chromosomes appeared at 3 h and most of the chromosomes
became fragmented with a size of less than 700 kbp by 5 h. After 6 h, fragmented
chromosomes persisted by 24 h (Fig. 3a, Supplementary Fig. S4 for wild-type
control). The depletion of Rfa1 from 4 h did not affect the persistence of
fragmented chromosomes. This suggests that meiotic DSBs remain unrepaired
under the depletion of RPA in the *dmc1* mutant.
We also indirectly checked the DSB status by analyzing Rad51 foci on
meiotic chromosomes (Fig. 3b, c). Like Rfa2 foci, in the *dmc1 RFA1-AID* mutant
in the absence of auxin, Rad51 foci appeared at 4 h, reached a plateau at 5 h,
and persisted at late times as shown previously [56]. The addition of auxin at 4 h
rapidly induced the disappearance of Rad51 foci at 5 h. The reduced Rad51 foci
are seen at late time points (Fig. 3c and Supplementary Fig. 3b). This suggests
that RPA is critical for the maintenance of Rad51 foci on the chromosomes. The
absence of Rad51 foci in the Rfa1 depletion supports the inability of *dmc1 RFA1-*
AID cells without RPA to repair meiotic DSBs by Rad51-dependent recombination.
260
The Hop1/Mek1-Ndt80 axis was down-regulated after RFA1 depletion
Un-repaired DSBs in the *dmc1* mutant activate the recombination checkpoint,
which down-regulates a meiosis-specific transcription activator, Ndt80 for the
pachytene exit (Fig. 1). DSB-activated Mec1/ATR kinase phosphorylates a
HORMAD protein, Hop1, which stimulates a meiosis-specific kinase, Mek1. Mek1
binds and phosphorylates Ndt80 to inactivate it. For pachytene exit, further
phosphorylation of Nd80 by Ime2 kinase promotes the activation of Ndt80 [38]. In
dmc1 RFA1-AID cells, the Ndt80 appeared at 4 h with multiple bands and
accumulated more during further incubation in SPM without a change of band-
shift status (Fig. 2a). At 6 h, after two hours of addition of auxin, a slower-
migrating Ndt80 band appeared and accumulated. Importantly, as shown above,
under the depletion condition, Cdc5/PLK, which is a major target of activated
Ndt80 and is necessary for pachytene exit, appeared at 6 h (Fig. 2a). This is
consistent with meiosis progression of the *dmc1* mutant cells induced by Rfa1-
depletion.
We also checked the behavior of the upstream regulators of Ndt80, and
Hop1 by western blotting. In the *dmc1* mutant, Hop1 appeared as a non-
phosphorylated form at 2 h and phosphorylated Hop1 with slower migrated bands
persisted from 4 h (Fig. 2a) as shown previously [36]. After Rfa1 depletion, major

bands of phosphorylated Hop1 almost disappeared from 6 h with a similar steady-
state level of non-phosphorylated Hop1 during the time course. This suggests
that Hop1 is de-phosphorylated during pachytene exit. On chromosome spreads,
Hop1 shows beads-in-line staining for the localization of chromosome axes (Fig.
3d) [57,58]. In the *dmc1* mutant without Rfa1 depletion, Hop1 localization remains
unchanged during meiosis (Fig. 3d). The addition of auxin at 4 h induced the
disappearance of Hop1-positive spreads from 5 h. At 6 h, ~35% of cells are
positive for Hop1 while ~95% are positive in a control (Fig. 3d, e). At 8 h, most of
the Hop1 signals on the chromosomes disappeared.

One of the Mek1 kinase targets is histone H3T11 phosphorylation [59]. Under
the *dmc1* arrest condition, H3T11 phosphorylation accumulated (Fig. 2a). On the
other hand, the Rfa1 depletion rapidly induced the disappearance of the signal
(Fig. 2a), suggesting the downregulation of Mek1 activity under Rfa1 depletion.
These results clearly showed that Rfa1, thus RPA, is critical for the maintenance
of the recombination checkpoint through the Hop1/Mek1-Ndt80 axis.

The *dmc1*-induced arrest in meiosis is also partially suppressed by the
rad51 [60]. A recent study showed that the suppression of *dmc1*-arrest by the *rad51*
mutation is mediated by partial elongation of SCs with concomitant loss of Hop1
[61]. Rad51 loss from the chromosomes in the *dmc1* mutant upon Rfa1-depletion
such as 5-h time point by 4-h addition of auxin seems to mimic a condition seen
in the *rad51 dmc1* double mutant. To address whether Rad51-focus loss in the
dmc1 mutant by Rfa1 depletion induced SC formation, we checked SC formation
by staining Zip1, a component of SC central region [41]. The Zip1 staining is
classified into three classes; dots, short lines, and long lines. As shown previously
[20], the *dmc1* mutant is defective in SC elongation (Supplemental Fig. 5a, b). Rfa1
depletion promotes the SC disassembly by the pachytene exit. At 5 h when most
of Rad51 foci disappeared, we did not see elongation of Zip1 lines relative to
control with Rad51 foci and rather observed increased Zip1 polycomplex, which
is indicative of defective SC assembly (Supplemental Fig. 5c). Moreover, Hop1
localization one-hour after Rfa1 depletion is similar to that without the depletion
(Fig. 3d, e). These suggests that the suppression of arrest by Rfa1 depletion is
unlikely to come from altered Hop1 functions as seen in the *rad51 dmc1* double
mutant [61].

Bypass of *dmc1* arrest by Rfa1 depletion depends on Ndt80
We tested whether the pachytene exit of the *dmc1* cells by Rfa1 depletion

depends on the *NDT80* (Fig. 4). The addition of auxin to *dmc1 ndt80 RFA1-AID*
cells efficiently reduced the amount of Rfa1 protein (Fig. 4a, Supplementary Fig.
S6) and Rfa2 foci from meiotic chromosomes (Fig. 4b). However, different from
the *dmc1 RFA1-AID* mutant, the addition of auxin to *dmc1 ndt80 RFA1-AID*
mutant did not induce the elongation of spindles (Fig. 4c) or Hop1 disassembly
from the chromosomes (Fig. 4d). Thus, the pachytene exit of *dmc1* cells by Rfa1
depletion still depends on Ndt80. Interestingly, the amount of the phosphorylated
Hop1 is reduced slightly compared to that in the control (Fig. 4a). This is less
drastic compared to the presence of Ndt80, supporting an auto-feedback
mechanism between Mek1 activation and Ndt80 activation [62].

Checkpoint alleviation in Rfa1 depletion in *zip1* meiosis
Meiotic yeast cells monitor not only ongoing meiotic recombination but also
chromosome synapsis [43]. However, the presence of the so-called "synapsis
checkpoint" in yeast meiosis is under debate [62]. The mutant in the *ZIP1* gene is
deficient in chromosome synapsis with a delay of the entry into meiosis I and is
used to analyze the synaptic checkpoint [41]. We examined the effect of Rfa1
depletion on the *zip1*-induced delay of meiosis I (Fig. 5). As reported previously
[45], in the SK1 strain background, *zip1 RFA1-AID* cells without auxin addition
started meiosis I at 8 h while wild-type cells entered at 5 h (and finished meiosis
II by 10 h) (Fig. 5b). When auxin was added at 4 h, which induced rapid depletion
of Rfa1 and the dissociation of Rfa2 from meiotic chromosomes (Fig. 5a, c, h,
Supplementary Fig. S7), metaphase I spindles appeared at 5 h, and at 6 h, ~35%
of the cells contained metaphase-I and anaphase-I spindles (Fig. 5d, e). At 8 h,
~30% of the cells contained anaphase-II spindle. Importantly, DAPI staining
showed the entry of the *zip1* cells into meiosis I upon Rfa1 depletion is similar to
that in wild-type cells (Fig. 5b). The spore viability of *zip1 RFA1-AID* cells treated
with auxin was 0% while cells without auxin was 35.5% (among 100 tetrads).
The *zip1 RFA1-AID* cells without auxin expressed Cdc5 from 8 h (Fig. 5a),
consistent with delayed progression of prophase I in the mutant. The addition of
auxin at 4 h accelerated the Cdc5 expression from 5 h. The *zip1* mutant
accumulated H3T11 phosphorylation of histone H3 signal from 4 h to 8 h (Fig.
5a). The Rfa1 depletion accelerated the disappearance of this phosphorylation
from 5 h. The accelerated disappearance of the phospho-Hop1 protein is also
seen by the depletion (Fig. 5a, f). These showed the Rfa1-dependent inactivation
of the Hop1/Mek1-Ndt80 signaling pathway induced by the *zip1* mutation. Thus,

the delay induced by the *zip1* mutation largely depends on the Rfa1.

In addition to SC formation, Zip1 plays a critical role in meiotic recombination

as a pro-crossover factor. Indeed, the *zip1* mutant shows delayed DSB repair.

The CHEF analysis showed the presence of broken chromosomes in the *zip1*

mutant from 3 h to 7 h (Fig. 5g). Importantly, full-length chromosomes were

reassembled at 10 h without Rfa1 depletion, indicating that most of the DSBs in

the *zip1* mutant were repaired with the delay. On the other hand, Rfa1 depletion

from 4 h induced fragmentation of the chromosomes from 5 h. Even after 24 h,

little recovery of full-length chromosomes was seen under the condition. This

suggests that Rfa1 is essential for meiotic DSB repair in the *zip1* mutant.

Rfa1 depletion does not alleviate delay induced by *rad50S* meiosis

The *rad50S* mutant, which is defective in the end-processing of meiotic DSBs,

also shows delayed meiosis I progression [63]. This delay in the *rad50S* mutant is

mediated by Tel1(ATM) and Rad50-Mre11-Xrs2, which recognize DSB ends [64].

The pathway is distinct from the Mec1(ATR)-mediated pathway shown above. We

also examined the effect of Rfa1 depletion on the *rad50S*-induced delay (Fig. 6a,

Supplementary Fig. S8). As reported, without auxin, *rad50S RFA1-AID* cells delay

the onset of meiosis I as shown by whole-cell staining of tubulin (Fig. 6b). Rather,

the addition of auxin to the *rad50S RFA1-AID* cells delayed the onset of meiosis

I more than without the auxin. We cannot detect Rfa2 foci in the r*ad50S* mutant

even in the presence of Rfa1 (Fig. 6c), confirming that the *rad50S* accumulates

unresected DSBs [63]. Moreover, Hop1 phosphorylation is not observed in the

mutant (Fig. 6a). This is consistent with little formation of ssDNAs without DSB

processing in the mutant [63], thus little Mec1/ATR activation. This is expected since

the *rad50S* mutant does not form any ssDNAs, which is a substrate for RPA. The

accumulation of chromosomal Hop1 in the mutant is not affected by the Rfa1

depletion (Fig. 6d).

Discussion

RPA plays multiple roles in DNA replication, DNA repair, and recombination

through its ability to bind to ssDNAs with a high affinity. RPA is also important for

DDR as a sensor of DNA damage, thus ssDNAs. The role of RPA in Mec1(ATR)

activation through the binding of Mec1(ATR)-interaction protein, Ddc2(ATRIP), to

RPA has been documented well [5,13,14]. In this paper, we showed the pivotal role

of RPA in the maintenance of DDR in meiosis, called recombination checkpoint.

The role of RPA in recombination checkpoint

Previously, in the budding yeast, two *RFA1*-degron mutants, *rfa1-td* and *td-RFA1*,
were constructed and characterized to study the role of Rfa1 in DDR activation.
Indeed, the *rfa1-td*, as well as the *rft1-t11 (rfa1-K45E)* mutant, reduce the
recruitments of Ddc2 to a DSB site [5]. On the other hand, it is shown that, by using
the *td-RFA1*, G2/M-arrested haploid cells depleted for Rfa1 still activate Rad53
upon a single DSB [6]. The latter study concludes that a low level of RPA is
sufficient for the activation of Mec1-Ddc2. More recently, an RPA-independent
pathway for ATR activation, which requires APE1, was identified in *Xenopus* egg
extracts [65]. We revisited the role of RPA in DDR by studying the effect of Rfa1
depletion in DDR such as the meiotic recombination checkpoint. The
recombination checkpoint in meiotic cells responds to the ssDNA through RPA-
Mec1-Ddc2 axis as seen in a canonical DNA damage checkpoint in mitotic cells.
By using the auxin-degron, we depleted Rfa1 in the *dmc1* mutant which arrests
in meiotic prophase I with hyper-resected ssDNAs in ~150 DSBs [20]. The Rfa1-
depletion induced the pachytene exit by downregulating Mec1 and Mek1
activities, which in turn activates the Ndt80-dependent gene expression of key
regulators for the pachytene exit such as Cdc5 [40,55]. This is consistent with the
previous study which showed that the *rfa1-t11* with reduced Ddd2-focus formation
in nuclei weakly suppresses the *dmc1*-induced delay [50]. Importantly, the Rfa1
depletion did not affect the accumulation of ssDNAs in the *dmc1* cells with
persistent unrepaired DSBs. This is particularly important since mutations in DDR
genes such as *RAD17*, *-24*, *DDC1*, and *MEC3* even in the *MEC1* gene bypass
the *dmc1*-induced arrest by repairing DSBs by an alternative pathway such as
ectopic recombination [46,48,66]. The results described here supported the idea that
RPA is a sensor for ssDNAs, which creates a scaffold for the assembly of
checkpoint effectors such as Mec1-Ddc2(ATR-ATRIP). In meiosis, meiosis-
specific Mek1 is activated through the phosphorylation of Hop1 by Mec1 kinase
[36]. The Rfa1-depletion induces rapid dephosphorylation of Hop1 and the
dissociation from meiotic chromosomes without affecting the protein level,
suggesting the presence of an active de-phosphorylation pathway of Hop1.

The chromosome synapsis checkpoint in yeast meiosis

In addition to the recombination checkpoint, meiotic cells responds to abnormal
chromosome synapsis, which is called synapsis checkpoint. This idea was

proposed that mutants defective in chromosome synapsis show a delay in the
progression of prophase I and apoptosis in yeasts and mice or nematodes,
respectively. However, the presence of the synapsis checkpoint is still under
debate [42,43], since in most organisms meiotic recombination and chromosome
synapsis are tightly coupled. The budding yeast *zip1* mutant, which is completely
defective in SC formation, thus, chromosome synapsis, delays or arrest in meiotic
prophase I [41,45]. This arrest is alleviated by the mutant of genes encoding a
component of checkpoint such as *PCH2* [67]. Pch2 modulates Hop1 conformation
and regulates the association and dissociation of Hop1 on meiotic chromosomes
[61,68,69]. On the other hand, the *zip1* mutant also shows defective DSB repair
and/or the crossing-over [44,45], which suggests delayed progression of meiosis I in
the *zip1* mutant is caused by the impaired recombination. In this study, by using
the *RFA1-AID*, we could test whether the delay in *zip1* mutant depends on RPA
or not and found that the Rfa1-depletion eliminates the delay induced by the *zip1*
mutation. This strongly suggests that the recombination checkpoint delays the
meiosis I progression in the *zip1* mutant as seen in the *dmc1* mutant. Moreover,
consistent with this, the other yeast mutants, which are fully defective in the
synapsis with a weak recombination defect such as *ecm11* and *gmc2*, do not
show a drastic delay in the meiosis I progression [70,71]. Thus, at least in budding
yeast which couples the recombination and chromosome synapsis, rather than
the status of chromosome synapsis, a recombination process, the presence of
ssDNAs, can be monitored by a meiosis-specific surveillance mechanism to
guarantee the completion of meiotic recombination directly and chromosome
synapsis indirectly. Our idea is consistent with the idea proposed by recent
reports claiming the merging of the recombination and synapsis checkpoints in
meiosis [61,72].
450
Methods
Strains and plasmids
All strains described here are derivatives of *S. cerevisiae* SK1 diploids,
MSY832/833 (*MATα/MAT**a**, lys2/", ura3/", leu2::hisG/", trp1::hisG/"*). The
genotypes of each strain used in this study are described in Supplementary Table
S1
457
Strain Construction
The *RFA1-AID* gene was constructed by one-step DNA integration. The AID-

9Myc tag was added to the C-terminus of the *RFA1* open reading frame by the
transformation of yeast cells (MSY823) with a DNA fragment amplified by PCR
using p7AID-9myc (provided by Dr. Neil Hunter) as a template and a pair of
primers (Rfa1-AID-Hygoro-F and -R in Supplementary Table S2). The
transformants were selected against hygromycin. Genomic DNAs from the
candidate colonies were purified and the genomic DNAs were checked for the
correct tagging by PCR and DNA sequencing using the primer described in
Supplementary Table S2.
468
Anti-serum and antibodies
Anti-myc (9E10, sc-40, Santa Cruz), anti-tubulin (MCA77G, Bio-Rad/Serotec,
Ltd), anti-Cdc5 (sc-33625, Santa Cruz), and histone H3 phospho-T11 (ab5168,
Abcam) were used for western blotting. Rabbit anti-Rad51 [21] and guinea pig anti-
Rad51 [73], rabbit anti-Hop1 [74] and rabbit anti-Rfa2 [75] were home-made and used
for staining. Rabbit anti-Ndt80 was a gift by M. Lichten. The secondary antibodies
for immuno-staining were Alexa Fluor 488 (Goat, A-21206, Thermo Fisher
Scientific) and 594 (Goat, A-211012, Thermo Fisher Scientific) IgG used at a
1/2000 dilution. For western blotting, alkaline phosphatase-conjugated secondary
antibodies (Promega) were used.
479
Meiotic Time course
Saccharomyces cerevisiae SK1 strains were patched onto YPG plates (2%
bacteriological peptone, 1% yeast extract, 3% glycerol, 2% bacteriological agar)
and incubated at 30 °C for 12 h. Cells were inoculated onto YPD plates (2%
bacteriological peptone, 1% yeast extract, 2% glucose, 2% bacteriological agar)
and grown for two days to obtain isolated colonies. A single colony was inoculated
into 3 ml of YPD liquid medium and grown to saturation at 30 °C overnight. To
synchronize cell cultures, the overnight cultures were transferred to pre-warmed
SPS medium (1% potassium acetate, 1% bacteriological peptone, 0.5% yeast
extract, 0.17% yeast nitrogen base with ammonium sulfate and without amino
acids, 0.5% ammonium sulfate, 0.05 M potassium biphthalate) and grown for 16–
17 h. Meiosis was induced by transferring the SPS-cultured cells to pre-warmed
SPM (1% potassium acetate, 0.02% raffinose). The SPM-cultured cells were
harvested at various times after transfer. For Rfa1 depletion from 4 h, $CuSO_4$ was
added to 50 mM at 3.5 h. Auxin (3-indoleacetic acid) solved in dimethyl sulfoxide
(DMSO) was added at a final concentration of 2 mM at 4 h. In the control

experiment (without auxin), we added an equivalent DMSO solution alone.
497

Cytology

Chromosome spreads were immunostained as described previously [48,73].
Spheroplasts were burst in the presence of 1% paraformaldehyde and 0.1% lipsol.
The spreads were stained with primary and secondary antibodies observed using
a fluorescence microscope (BX51; Olympus/Evident) with a 100× objective
(NA1.3). Images were captured by CCD camera (CoolSNAP; Roper), and then
processed using IP lab and/or iVision (Bio Vision Technologies) and Photoshop
(version, 25.4.0; Adobe; https://www.adobe.com) software. For focus counting,
~30 nuclei were counted at each time point.

Whole-cell staining was carried out as follows. Yeast cells are fixed with
formaldehyde (a final concentration of 3.7%) and collected. The pellets were
suspended in the ZK buffer (25 mM Tris-HCl [pH 7.5], 80 mM KCl) and treated
with DTT (20 mM) and then with Zymolyase 100T (2.5 μg/mL). After washing with
PBS, the cells were placed on a glass slide coated with poly-lysine (1 mg/ml).
The cells were treated with cold ethanol (100% 6 min) and acetone (100% 0.5
min), and washed with PBS. After blocking with PBS with 1% BSA, cells were
stained with primary and secondary antibodies.
515

Western blotting

Western blotting was performed as described previously [22,76], using cell lysates
extracted by the TCA method. After being harvested and washed twice with 20%
TCA, cells were roughly disrupted with zirconia beads by the Multi-beads shocker
(Yasui Kikai). The protein precipitate recovered by centrifugation was suspended
in SDS-PAGE sample buffer adjusted to pH 8.8 and then incubated at 95 °C for
10 min. After electrophoresis, the proteins were transferred onto a nylon
membrane (Immobilon-P, Millipore) and incubated with primary antibodies in a
blocking buffer (1× PBS, 0.5% BSA) and then with alkaline phosphatase-
conjugated secondary antibody (Promega). The color reaction was developed
with NBT/BCIP solution (Nacalai Tesque).
527

CHEF analysis Pulsed-field gel electrophoresis

For pulsed-field gel electrophoresis (PFGE), chromosomal DNA was prepared in
1.3% of low-melting point agarose plugs as described in [74] and run at 14 °C with
1X TBE buffer in a CHEF DR-III apparatus (BioRad) using the field 6V/cm at a

120° angle for 48 h. Switching times followed a ramp from 15.1 to 25.1 seconds.
After the electrophoresis, the gel was stained with 0.5 µg/mL ethidium bromide
for 30 min and washed with distilled water. Images were taken using ImageQuant
LAS4000 BioImager (GE Healthcare).
536
FACS analysis
Cells in 1 ml of SPM culture were collected by centrifugation. The pellet was then
resuspended in 1 ml of 70% ethanol. Fixed cells were recollected and washed by
suspending 1 ml of 50 mM sodium citrate buffer. The collected cells were
resuspended in 0.5 ml of sodium citrate buffer and sonicated for the 20s at room
temperature. Sonicated cells were incubated with 12.5 µl of 10 mg/mL RNAseA
for 1 h at 50 °C and in 25 µl of 15mg/ml proteinase K for 1h at 50 °C. 0.5ml of 16
µlg/mL PI solution was added and incubated for 30 min at room temperature in
the dark. The samples were then analyzed using SA-3800 Spectral analyzer
(Sony). Floreada.io was used for gating and plotting graphs.
547
548   **Data availability**
549   Uncropped western blotting images are provided in Supplementary Figures. The
550   raw data used in plots and graphs are also shown in Source Data. The raw
551   microscopy images used for figures and quantification are available from
552   corresponding authors upon request.
553

Acknowledgments
We are grateful to Drs. Neil Hunter for providing the yeast AID system and
Michael Lichten for anti-Ndt80. A.Sa. is a JICA fellow. Y.F. and M.I. were
supported by the Institute for Protein Research. This work was supported by a
grant from Institute for Fermentation, Osaka (IFO) and from Uehara Memorial
Foundation to Life Sciences; to A.Sh.

560
561   **Competing interest**
562   The authors declare no competing financial interest.
563

564   **Author contributions**
565   A.Sh. conceived and designed the experiments. A.Sa., Z.C., and Y.T. carried out
566   experiments with help from Y.F. and M.I.. A.Sa, Z.C., Y.T., Y.F., M.I., and A.Sh.
567   analyzed the data. A.Sh. wrote the manuscript with inputs from help from others.

References

References

1   Bhat, K. P. & Cortez, D. RPA and RAD51: fork reversal, fork protection,
and genome stability. *Nat Struct Mol Biol* **25**, 446-453 (2018).
https://doi.org:10.1038/s41594-018-0075-z

2   Caldwell, C. C. & Spies, M. Dynamic elements of replication protein A at
the crossroads of DNA replication, recombination, and repair. *Crit Rev*
*Biochem   Mol   Biol   ***55***,   482-507   (2020).
https://doi.org:10.1080/10409238.2020.1813070

3   Wold, M. S. Replication protein A: a heterotrimeric, single-stranded DNA-
binding protein required for eukaryotic DNA metabolism. *Annu Rev*
Biochem   **66**,   61-92   (1997).
https://doi.org:10.1146/annurev.biochem.66.1.61

4   Brill, S. J. & Stillman, B. Replication factor-A from Saccharomyces
cerevisiae is encoded by three essential genes coordinately expressed at
S   phase.   *Genes   Dev*   **5**,   1589-1600   (1991).
https://doi.org:10.1101/gad.5.9.1589

5   Zou, L. & Elledge, S. J. Sensing DNA damage through ATRIP recognition
of   RPA-ssDNA   complexes.   *Science*   **300**,   1542-1548   (2003).
https://doi.org:10.1126/science.1083430

6   Chen, H., Lisby, M. & Symington, L. S. RPA coordinates DNA end resection
and prevents formation of DNA hairpins. *Mol Cell* **50**, 589-600 (2013).
https://doi.org:10.1016/j.molcel.2013.04.032

7   Sasanuma, H. *et al.* A new protein complex promoting the assembly of
Rad51   filaments.   *Nat   Commun*   **4**,   1676   (2013).
https://doi.org:10.1038/ncomms2678

8   Shinohara, A. & Ogawa, T. Stimulation by Rad52 of yeast Rad51-mediated
recombination. *Nature* **391**, 404-407 (1998). https://doi.org:10.1038/34943

9   Sung, P. Yeast Rad55 and Rad57 proteins form a heterodimer that
functions with replication protein A to promote DNA strand exchange by
Rad51   recombinase.   *Genes   Dev*   **11**,   1111-1121   (1997).
https://doi.org:10.1101/gad.11.9.1111

10   Sung, P. Catalysis of ATP-dependent homologous DNA pairing and strand
exchange by yeast RAD51 protein. *Science* **265**, 1241-1243 (1994).
https://doi.org:10.1126/science.8066464

11   Friedel, A. M., Pike, B. L. & Gasser, S. M. ATR/Mec1: coordinating fork

stability and repair. *Curr Opin Cell Biol* **21**, 237-244 (2009).
https://doi.org:10.1016/j.ceb.2009.01.017

12   Williams, R. M., Yates, L. A. & Zhang, X. Structures and regulations of ATM
and ATR, master kinases in genome integrity. *Curr Opin Struct Biol* **61**, 98-
105 (2020). https://doi.org:10.1016/j.sbi.2019.12.010

13   Biswas, H., Goto, G., Wang, W., Sung, P. & Sugimoto, K. Ddc2ATRIP
promotes Mec1ATR activation at RPA-ssDNA tracts. *PLoS Genet* **15**,
e1008294 (2019). https://doi.org:10.1371/journal.pgen.1008294

14   Deshpande, I. *et al.* Structural Basis of Mec1-Ddc2-RPA Assembly and
Activation on Single-Stranded DNA at Sites of Damage. *Mol Cell* **68**, 431-
445 e435 (2017). https://doi.org:10.1016/j.molcel.2017.09.019

15   Yates, L. A., Tannous, E. A., Morgan, R. M., Burgers, P. M. & Zhang, X. A
DNA damage-induced phosphorylation circuit enhances Mec1(ATR)
Ddc2(ATRIP) recruitment to Replication Protein A. *Proceedings of the*
National Academy of Sciences of the United States of America **120**,
e2300150120 (2023). https://doi.org:10.1073/pnas.2300150120

16   Bass, T. E. *et al.* ETAA1 acts at stalled replication forks to maintain genome
integrity. *Nat Cell Biol* **18**, 1185-1195 (2016).
https://doi.org:10.1038/ncb3415

17   Haahr, P. *et al.* Activation of the ATR kinase by the RPA-binding protein
ETAA1. *Nat Cell Biol* **18**, 1196-1207 (2016).
https://doi.org:10.1038/ncb3422

18   Bergerat, A. *et al.* An atypical topoisomerase II from Archaea with
implications for meiotic recombination. *Nature* **386**, 414-417 (1997).
https://doi.org:10.1038/386414a0

19   Keeney, S., Giroux, C. N. & Kleckner, N. Meiosis-specific DNA double-
strand breaks are catalyzed by Spo11, a member of a widely conserved
protein family. *Cell* **88**, 375-384 (1997).

20   Bishop, D. K., Park, D., Xu, L. & Kleckner, N. DMC1: a meiosis-specific
yeast homolog of E. coli recA required for recombination, synaptonemal
complex formation, and cell cycle progression. *Cell* **69**, 439-456 (1992).

21   Shinohara, A., Ogawa, H. & Ogawa, T. Rad51 protein involved in repair
and recombination in S. cerevisiae is a RecA-like protein. *Cell* **69**, 457-470
(1992).

22   Hayase, A. *et al.* A protein complex containing Mei5 and Sae3 promotes
the assembly of the meiosis-specific RecA homolog Dmc1. *Cell* **119**, 927-

940 (2004).

Tsubouchi, H. & Roeder, G. S. The budding yeast mei5 and sae3 proteins act together with dmc1 during meiotic recombination. *Genetics* **168**, 1219-1230 (2004). https://doi.org:10.1534/genetics.103.025700

Hong, S. *et al.* The logic and mechanism of homologous recombination partner choice. *Mol Cell* **51**, 440-453 (2013). https://doi.org:10.1016/j.molcel.2013.08.008

Kim, K. P. *et al.* Sister cohesion and structural axis components mediate homolog bias of meiotic recombination. *Cell* **143**, 924-937 (2010). https://doi.org:10.1016/j.cell.2010.11.015

Niu, H. *et al.* Partner choice during meiosis is regulated by Hop1-promoted dimerization of Mek1. *Molecular biology of the cell* **16**, 5804-5818 (2005). https://doi.org:10.1091/mbc.e05-05-0465

Leem, S. H. & Ogawa, H. The MRE4 gene encodes a novel protein kinase homologue required for meiotic recombination in Saccharomyces cerevisiae. *Nucleic Acids Res* **20**, 449-457 (1992). https://doi.org:10.1093/nar/20.3.449

Rockmill, B. & Roeder, G. S. A meiosis-specific protein kinase homolog required for chromosome synapsis and recombination. *Genes Dev* **5**, 2392-2404 (1991). https://doi.org:10.1101/gad.5.12b.2392

Callender, T. L. *et al.* Mek1 Down Regulates Rad51 Activity during Yeast Meiosis by Phosphorylation of Hed1. *PLoS Genet* **12**, e1006226 (2016). https://doi.org:10.1371/journal.pgen.1006226

Niu, H. *et al.* Regulation of meiotic recombination via Mek1-mediated Rad54 phosphorylation. *Mol Cell* **36**, 393-404 (2009). https://doi.org:10.1016/j.molcel.2009.09.029

Shinohara, M. *et al.* Characterization of the roles of the Saccharomyces cerevisiae RAD54 gene and a homologue of RAD54, RDH54/TID1, in mitosis and meiosis. *Genetics* **147**, 1545-1556 (1997). https://doi.org:10.1093/genetics/147.4.1545

Tsubouchi, H. & Roeder, G. S. Budding yeast Hed1 down-regulates the mitotic recombination machinery when meiotic recombination is impaired. *Genes Dev* **20**, 1766-1775 (2006). https://doi.org:10.1101/gad.1422506

Borner, G. V., Hochwagen, A. & MacQueen, A. J. Meiosis in budding yeast. *Genetics* (2023). https://doi.org:10.1093/genetics/iyad125

Cartagena-Lirola, H., Guerini, I., Manfrini, N., Lucchini, G. & Longhese, M.

P. Role of the Saccharomyces cerevisiae Rad53 checkpoint kinase in

signaling double-strand breaks during the meiotic cell cycle. *Molecular and*

cellular biology **28**, 4480-4493 (2008).

https://doi.org:10.1128/MCB.00375-08

35   Usui, T. & Shinohara, A. Rad9, a 53BP1 Ortholog of Budding Yeast, Is

Insensitive to Spo11-Induced Double-Strand Breaks During Meiosis. *Front*

Cell Dev Biol **9**, 635383 (2021). https://doi.org:10.3389/fcell.2021.635383

36   Carballo, J. A., Johnson, A. L., Sedgwick, S. G. & Cha, R. S.

Phosphorylation of the axial element protein Hop1 by Mec1/Tel1 ensures

meiotic interhomolog recombination. *Cell* **132**, 758-770 (2008).

https://doi.org:10.1016/j.cell.2008.01.035

37   Hollingsworth, N. M. Mek1/Mre4 is a master regulator of meiotic

recombination in budding yeast. *Microb Cell* **3**, 129-131 (2016).

https://doi.org:10.15698/mic2016.03.487

38   Chen, X. *et al.* Mek1 coordinates meiotic progression with DNA break

repair by directly phosphorylating and inhibiting the yeast pachytene exit

regulator Ndt80. *PLoS Genet* **14**, e1007832 (2018).

https://doi.org:10.1371/journal.pgen.1007832

39   Hochwagen, A. Meiosis: making a synaptonemal complex just got easier.

Curr Biol **19**, R849-851 (2009). https://doi.org:10.1016/j.cub.2009.07.001

40   Sourirajan, A. & Lichten, M. Polo-like kinase Cdc5 drives exit from

pachytene during budding yeast meiosis. *Genes Dev* **22**, 2627-2632

(2008). https://doi.org:10.1101/gad.1711408

41   Sym, M., Engebrecht, J. A. & Roeder, G. S. ZIP1 is a synaptonemal

complex protein required for meiotic chromosome synapsis. *Cell* **72**, 365-

378 (1993).

42   Huang, Y. & Roig, I. Genetic control of meiosis surveillance mechanisms

in mammals. *Front Cell Dev Biol* **11**, 1127440 (2023).

https://doi.org:10.3389/fcell.2023.1127440

43   MacQueen, A. J. & Hochwagen, A. Checkpoint mechanisms: the puppet

masters of meiotic prophase. *Trends Cell Biol* **21**, 393-400 (2011).

https://doi.org:10.1016/j.tcb.2011.03.004

44   Borner, G. V., Kleckner, N. & Hunter, N. Crossover/noncrossover

differentiation, synaptonemal complex formation, and regulatory

surveillance at the leptotene/zygotene transition of meiosis. *Cell* **117**, 29-

45 (2004).

45 Storlazzi, A., Xu, L., Schwacha, A. & Kleckner, N. Synaptonemal complex
(SC) component Zip1 plays a role in meiotic recombination independent
of SC polymerization along the chromosomes. *Proceedings of the National*
Academy of Sciences of the United States of America **93**, 9043-9048
(1996). https://doi.org:10.1073/pnas.93.17.9043

46 Grushcow, J. M. *et al.* Saccharomyces cerevisiae checkpoint genes MEC1,
RAD17 and RAD24 are required for normal meiotic recombination partner
choice. *Genetics* **153**, 607-620 (1999).
https://doi.org:10.1093/genetics/153.2.607

47 Shinohara, M., Hayashihara, K., Grubb, J. T., Bishop, D. K. & Shinohara,
A. DNA damage response clamp 9-1-1 promotes assembly of ZMM
proteins for formation of crossovers and synaptonemal complex. *Journal*
of cell science **128**, 1494-1506 (2015). https://doi.org:10.1242/jcs.161554

48 Shinohara, M., Sakai, K., Ogawa, T. & Shinohara, A. The mitotic DNA
damage checkpoint proteins Rad17 and Rad24 are required for repair of
double-strand breaks during meiosis in yeast. *Genetics* **164**, 855-865
(2003). https://doi.org:10.1093/genetics/164.3.855

49 Kim, H. S. & Brill, S. J. Rfc4 interacts with Rpa1 and is required for both
DNA replication and DNA damage checkpoints in Saccharomyces
cerevisiae. *Molecular and cellular biology* **21**, 3725-3737 (2001).
https://doi.org:10.1128/MCB.21.11.3725-3737.2001

50 Refolio, E., Cavero, S., Marcon, E., Freire, R. & San-Segundo, P. A. The
Ddc2/ATRIP checkpoint protein monitors meiotic recombination
intermediates. *Journal of cell science* **124**, 2488-2500 (2011).
https://doi.org:10.1242/jcs.081711

51 Joo, J. H. *et al.* RPA interacts with Rad52 to promote meiotic crossover
and noncrossover recombination. *Nucleic Acids Res* (2024).
https://doi.org:10.1093/nar/gkae083

52 Shi, B. *et al.* Dual functions for the ssDNA-binding protein RPA in meiotic
recombination. *PLoS Genet* **15**, e1007952 (2019).
https://doi.org:10.1371/journal.pgen.1007952

53 Nishimura, K., Fukagawa, T., Takisawa, H., Kakimoto, T. & Kanemaki, M.
An auxin-based degron system for the rapid depletion of proteins in
nonplant cells. *Nat Methods* **6**, 917-922 (2009).
https://doi.org:10.1038/nmeth.1401

54 Gasior, S. L., Wong, A. K., Kora, Y., Shinohara, A. & Bishop, D. K. Rad52

associates with RPA and functions with rad55 and rad57 to assemble
meiotic recombination complexes. *Genes Dev* **12**, 2208-2221 (1998).
https://doi.org:10.1101/gad.12.14.2208

55   Chu, S. *et al.* The transcriptional program of sporulation in budding yeast.
Science **282**, 699-705 (1998).
https://doi.org:10.1126/science.282.5389.699

56   Bishop, D. K. RecA homologs Dmc1 and Rad51 interact to form multiple
nuclear complexes prior to meiotic chromosome synapsis. *Cell* **79**, 1081-
1092 (1994).

57   Hollingsworth, N. M., Goetsch, L. & Byers, B. The HOP1 gene encodes a
meiosis-specific component of yeast chromosomes. *Cell* **61**, 73-84 (1990).
https://doi.org:10.1016/0092-8674(90)90216-2

58   Smith, A. V. & Roeder, G. S. The yeast Red1 protein localizes to the cores
of meiotic chromosomes. *J Cell Biol* **136**, 957-967 (1997).
https://doi.org:10.1083/jcb.136.5.957

59   Kniewel, R. *et al.* Histone H3 Threonine 11 Phosphorylation Is Catalyzed
Directly by the Meiosis-Specific Kinase Mek1 and Provides a Molecular
Readout of Mek1 Activity in Vivo. *Genetics* **207**, 1313-1333 (2017).
https://doi.org:10.1534/genetics.117.300359

60   Shinohara, A., Gasior, S., Ogawa, T., Kleckner, N. & Bishop, D. K.
Saccharomyces cerevisiae recA homologues RAD51 and DMC1 have
both distinct and overlapping roles in meiotic recombination. *Genes Cells*
2, 615-629 (1997). https://doi.org:10.1046/j.1365-2443.1997.1480347.x

61   Raina, V. B. & Vader, G. Homeostatic Control of Meiotic Prophase
Checkpoint Function by Pch2 and Hop1. *Curr Biol* **30**, 4413-4424 e4415
(2020). https://doi.org:10.1016/j.cub.2020.08.064

62   Hollingsworth, N. M. & Gaglione, R. The meiotic-specific Mek1 kinase in
budding yeast regulates interhomolog recombination and coordinates
meiotic progression with double-strand break repair. *Curr Genet* **65**, 631-
641 (2019). https://doi.org:10.1007/s00294-019-00937-3

63   Alani, E., Padmore, R. & Kleckner, N. Analysis of wild-type and rad50
mutants of yeast suggests an intimate relationship between meiotic
chromosome synapsis and recombination. *Cell* **61**, 419-436 (1990).
https://doi.org:10.1016/0092-8674(90)90524-i

64   Usui, T., Ogawa, H. & Petrini, J. H. A DNA damage response pathway
controlled by Tel1 and the Mre11 complex. *Mol Cell* **7**, 1255-1266 (2001).

784   https://doi.org:10.1016/s1097-2765(01)00270-2

785 65 Lin, Y. *et al.* APE1 recruits ATRIP to ssDNA in an RPA-dependent and -
786   independent manner to promote the ATR DNA damage response. *Elife* **12**
787   (2023). https://doi.org:10.7554/eLife.82324

788 66 Shinohara, M., Bishop, D. K. & Shinohara, A. Distinct Functions in
789   Regulation of Meiotic Crossovers for DNA Damage Response Clamp
790   Loader Rad24(Rad17) and Mec1(ATR) Kinase. *Genetics* **213**, 1255-1269
791   (2019). https://doi.org:10.1534/genetics.119.302427

792 67 San-Segundo, P. A. & Roeder, G. S. Pch2 links chromatin silencing to
793   meiotic checkpoint control. *Cell* **97**, 313-324 (1999).
794   https://doi.org:10.1016/s0092-8674(00)80741-2

795 68 Borner, G. V., Barot, A. & Kleckner, N. Yeast Pch2 promotes domainal axis
796   organization, timely recombination progression, and arrest of defective
797   recombinosomes during meiosis. *Proceedings of the National Academy of*
798   *Sciences of the United States of America* **105**, 3327-3332 (2008).
799   https://doi.org:10.1073/pnas.0711864105

800 69 Herruzo, E. *et al.* The Pch2 AAA+ ATPase promotes phosphorylation of
801   the Hop1 meiotic checkpoint adaptor in response to synaptonemal
802   complex defects. *Nucleic Acids Res* **44**, 7722-7741 (2016).
803   https://doi.org:10.1093/nar/gkw506

804 70 Humphryes, N. *et al.* The Ecm11-Gmc2 complex promotes synaptonemal
805   complex formation through assembly of transverse filaments in budding
806   yeast. *PLoS Genet* **9**, e1003194 (2013).
807   https://doi.org:10.1371/journal.pgen.1003194

808 71 Lee, M. S. *et al.* The synaptonemal complex central region modulates
809   crossover pathways and feedback control of meiotic double-strand break
810   formation. *Nucleic Acids Res* **49**, 7537-7553 (2021).
811   https://doi.org:10.1093/nar/gkab566

812 72 Gonzalez-Arranz, S. *et al.* Functional Impact of the H2A.Z Histone Variant
813   During Meiosis in Saccharomyces cerevisiae. *Genetics* **209**, 997-1015
814   (2018). https://doi.org:10.1534/genetics.118.301110

815 73 Shinohara, M., Gasior, S. L., Bishop, D. K. & Shinohara, A. Tid1/Rdh54
816   promotes colocalization of rad51 and dmc1 during meiotic recombination.
817   *Proceedings of the National Academy of Sciences of the United States of*
818   *America* **97**, 10814-10819 (2000).
819   https://doi.org:10.1073/pnas.97.20.10814

74   Bani Ismail, M., Shinohara, M. & Shinohara, A. Dot1-dependent histone
H3K79 methylation promotes the formation of meiotic double-strand
breaks in the absence of histone H3K4 methylation in budding yeast. *PLoS*
One **9**, e96648 (2014). https://doi.org:10.1371/journal.pone.0096648
75   Sasanuma, H., Furihata, Y., Shinohara, M. & Shinohara, A. Remodeling of
the Rad51 DNA strand-exchange protein by the Srs2 helicase. *Genetics*
194, 859-872 (2013). https://doi.org:10.1534/genetics.113.150615
76   Shinohara, M., Oh, S. D., Hunter, N. & Shinohara, A. Crossover assurance
and crossover interference are distinctly regulated by the ZMM proteins
during yeast meiosis. *Nat Genet* **40**, 299-309 (2008).
https://doi.org:10.1038/ng.83
831
832

Figure 1. A pathway of recombination checkpoint in meiosis.

Figure 2. Rfa1 depletion alleviates *dmc1*-induced pachytene arrest

(a) Expression of various proteins in meiosis. Lysates obtained from the *dmc1 RFA1-AID* (SAY63/64) cells at various time points during meiosis with the addition of auxin (2 mM) in DMSO or DMSO alone at 4 h were analyzed by western blotting using anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-histone H3-phospho-T11, anti-Ndt80, anti-Cdc5 or anti-tubulin (lower) antibodies.

(b) Rfa2 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with or without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and DAPI (blue). Representative images at each time point under the two conditions are shown. Bar = 2 μm.

(c) Kinetics of Rfa2 assembly/disassembly. The number of Rfa2-positive cells
(with more than 5 foci) was counted at each time point. At each time point,
more than 100 cells were counted.

(d) Meiosis progression. *dmc1 RFA1-AID* (SAY63/64) cells were analyzed by
DAPI staining. Representative image of cells treated with/without 4 h-auxin
addition at 10 h are shown. Bar = 2 μm.

(e) Meiosis progression. The entry into meiosis I and II in *dmc1 RFA1-AID*
(SAY63/64) cells were analyzed by DAPI staining. The number of DAPI
bodies in a cell was counted. A cell with 2, 3, and 4, and 3 and 4 DAPI bodies
was defined as a cell that passed through meiosis I and meiosis II,
respectively. The graph shows the percentages of cells that completed MI or
MII at the indicated time points. More than 200 cells were counted at each
time point. The representative results (*n*=3) are shown. *dmc1 RFA1-AID*
without auxin (with DMSO), blue closed circles; *dmc1 RFA1-AID* with auxin
addition at 4 h, red closed circles; Wild type (MSY823/833), gray circles.

(f) Whole cells were fixed and stained with anti-tubulin (green) and DAPI (blue).
Representative images at 8 h of *dmc1 RFA1-AID* (SAY63/64) cells with or
without the addition of auxin at 4 h are shown. Bar = 2 μm.

(g) Classification of tubulin staining at each time point of meiosis in *dmc1 RFA1-*
AID (SAY63/64) cells with (right) or without (left) the addition of auxin at 4 h.
Dot, short line, and long line tubulin-staining with single DAPI mass were
defined as prophase I (blue), metaphase I (red), anaphase I (orange),
metaphase II (green) and anaphase II (light green). At each time point, more
than 100 cells were counted. The representative results are shown (*n*=3).

Figure 3. Rfa1 depletion does not affect DSB turnover in *dmc1* mutant

(a) CHEF analysis of meiotic DSB repair. Chromosomal DNAs from *dmc1 RFA1-*
AID (SAY63/64) cells with the addition of auxin (2 mM) in DMSO or DMSO
alone at 4 h were studied by CHEF electrophoresis. The gel is stained with
EtBr. Auxin was added at 4 h in the incubation with SPM.

(b) Rad51 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with
or without the addition of auxin at 4 h were stained with anti-Rad51 (green),
and DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.

(c) Kinetics of Rad51 assembly/disassembly. The number of Rad51-positive cells
(with more than 5 foci) was counted at each time point. At each time point,
more than 100 cells were counted.

(d) Hop1 staining. Nuclear spreads from *dmc1 RFA1-AID* (SAY63/64) cells with
or without the addition of auxin at 4 h were stained with anti-Hop1 (green),
and DAPI (blue). Representative images at each time point under the two

conditions are shown. Bar = 2 μm.

(e) Kinetics of Hop1 assembly/disassembly. The number of Rad51-positive cells

(with more than 5 foci) was counted at each time point. At each time point,

more than 100 cells were counted.

Figure 4. *NDT80* is necessary for pachytene exit by Rfa1 depletion

(a) Expression of various protein in meiosis. Lysates obtained from the *ndt80 dmc1 RFA1-AID* (SAY80/81) cells at various time points during meiosis with the addition of auxin (2 mM) in DMSO or DMSO alone were analyzed by western blotting using anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5 or anti-tubulin (lower) antibodies.

(b) Rfa2 staining. Nuclear spreads from *ndt80 dmc1 RFA1-AID* (SAY80/81) cells with or without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and DAPI (blue). Representative images at each time point under the two conditions are shown. Bar = 2 μm.

(c) Classification of tubulin staining at each time point of meiosis in *ndt80 dmc1 RFA1-AID* (SAY80/81) cells with (right) or without (left) the addition of auxin at 4 h. Dot, short line, and long line tubulin-staining with single DAPI mass were defined as prophase I (blue), metaphase I (red), anaphase I (orange), metaphase II (blue), anaphase II (green) and post-meiosis II (pale green). Short and long tubulin-staining were defined as metaphase II and anaphase

II, respectively. At each time point, more than 100 cells were counted. The
representative results are shown (*n*=2).
(d) Hop1 staining. Nuclear spreads from *ndt80 dmc1 RFA1-AID* (SAY SAY80/81)
cells with or without the addition of auxin at 4 h were stained with anti-Hop1
(green), and DAPI (blue). Representative images at each time point under the
two conditions are shown. Bar = 2 μm.
918

Figure 5. Rfa1 depletion alleviates *zip1*-induced delay in meiosis I

(a) Expression of various proteins in meiosis. Lysates obtained from the *zip1 RFA1-AID* (SAY72/73) cells at various time points during meiosis with the addition of auxin (2 mM) in DMSO or DMSO alone were analyzed by western

blotting using anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5, anti-
histone H3-phospho-T11 or anti-tubulin (lower) antibodies.

(b) Meiotic cell cycle progression. The entry into meiosis I and II in *zip1 RFA1-*
AID (SAY72/73) cells were analyzed by DAPI staining as shown in Figure 2D.
More than 200 cells were counted at each time point. The representative
results (*n*=3) are shown. *dmc1 RFA1-AID* without auxin, blue closed circles;
dmc1 RFA1-AID with auxin addition at 4 h, red closed circles.

(c) Rfa2 staining. Nuclear spreads from *zip1 RFA1-AID* (SAY72/73) cells with or
without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and DAPI
(blue). Representative images at each time point under the two conditions are
shown. Bar = 2 μm.

(d) Tubulin staining in the *zip1 RFA1-AID* mutant. Whole cells were fixed and
stained with anti-tubulin (green) and DAPI (blue). Representative images for
each class are shown. Bar = 2 μm.

(e) Classification of tubulin staining at each time point of meiosis in *zip1 RFA1-*
AID (SAY72/73) cells with (right) or without (left) the addition of auxin at 4 h.
The quantification was performed as shown in Figure 1G. At each time point,
more than 100 cells were counted. The representative results are shown (*n*=2).

(f) Hop1 staining. Nuclear spreads from *zip1 RFA1-AID* (SAY72/73) cells with or
without the addition of auxin at 4 h were stained with anti-Hop1 (green), and
DAPI (blue). Representative images at each time point under the two
conditions are shown. Bar = 2 μm.

(g) CHEF analysis of meiotic DSB repair. Chromosomal DNAs from *zip1 RFA1-*
AID (SAY72/73) cells with or without the addition of auxin at 4 h were studied
by CHEF electrophoresis. Auxin was added at 4 h in the incubation with SPM.

(h) Kinetics of assembly/disassembly of Rfa2 (left), Rad51 (middle), and Hop1
(right). The number of focus-positive cells (with more than 5 foci) was counted
for Rfa2 and Rad51 at each time point. At each time point, more than 100
cells were counted.

Figure 6. Rfa1 depletion does not suppress *rad50S*-induced delay in meiosis I

(a) Expression of various proteins in meiosis. Lysates obtained from the *rad50S RFA1-AID* (SAY68/69) cells at various time points during meiosis with the addition of auxin (2 mM) in DMSO or DMSO alone were analyzed by western blotting using anti-myc (Rfa1-myc-AID, upper), anti-Hop1, anti-Cdc5 or anti-tubulin (lower) antibodies.

(b) Classification of tubulin staining at each time point of meiosis in *rad50S RFA1-AID* (SAY68/69) cells with (right) or without (left) the addition of auxin at 4 h. The quantification was performed as shown in Figure 1G. At each time point, more than 100 cells were counted. The representative results are shown (*n*=1).

(c) Rfa2 staining. Nuclear spreads from *rad50S RFA1-AID* (SAY68/69) cells with or without the addition of auxin at 4 h were stained with anti-Rfa2 (red), and DAPI (blue). Representative images at each time point under the two conditions are shown. Bar = 2 μm.

(d) Hop1 staining. Nuclear spreads from *rad50S RFA1-AID* (SAY68/69) cells with

971   or without the addition of auxin at 4 h were stained with anti-Hop1 (red), and

972   DAPI (blue). Representative images at each time point under the two

973   conditions are shown. Bar = 2 μm.