1 **Disruption of Synaptic Endoplasmic Reticulum Luminal Protein Containment in**
2 ***Drosophila Atlastin* Mutants**
3

Mónica C. Quiñones-Frías[1], Dina M. Ocken[1] and Avital Rodal[1,*]

[1]Department of Biology, Brandeis University, Waltham, MA

*Corresponding author:

Email: arodal@brandeis.edu

<u>**Abstract:**</u>

The endoplasmic reticulum (ER) extends throughout neurons and regulates many neuronal
functions, including neurite outgrowth, neurotransmission, and synaptic plasticity. Mutations in
proteins that control ER shape are linked to the neurodegenerative disorder Hereditary Spastic
Paraplegia (HSP), yet the ultrastructure and dynamics of neuronal ER remain largely
unexplored, especially at presynaptic terminals. Using super-resolution and live imaging in *D.*
melanogaster larval motor neurons, we investigated ER structure at presynaptic terminals of
wild-type animals and null mutants of the ER shaping protein and HSP-linked gene, Atlastin.
Previous studies using an ER luminal marker reported diffuse localization at *Atlastin* mutant
presynaptic terminals, which was attributed to ER fragmentation. However, using an ER
membrane marker, we discovered that *Atlastin* mutant ER forms robust networks with only mild
defects in structural dynamics, indicating the primary defect is functional rather than
architectural. We demonstrate that *Atlastin* mutants progressively displace overexpressed
luminal ER proteins to the cytosol during larval development, specifically at synapses, while
these proteins remain correctly localized in cell bodies, axons, and muscles. This synaptic-
specific displacement phenotype, previously unreported in non-neuronal cells, emphasizes the
importance of studying neurons to understand HSP pathogenesis.

<u>**Introduction:**</u>

The endoplasmic reticulum (ER) is a continuous organelle that extends to the periphery
of neurons and regulates many neuronal functions, including neurite outgrowth,
neurotransmission, and synaptic plasticity (de Juan-Sanz et al., 2017; Liu et al., 2019; Oliva et
al., 2020; Perez-Alvarez et al., 2020; Raffaello et al., 2016; Raiborg et al., 2015). Structurally,
this essential organelle consists of two domains: ER sheets (enriched in cell bodies) and ER
tubules (abundant in dendrites and axons) (Öztürk et al., 2020; Wu et al., 2017). Recent studies

have revealed even finer levels of organization, with increasing evidence to suggest further structural and functional differences in neuronal ER tubules. For instance, different regions of the neuron show distinct features, including varying densities of ER-organelle contacts (Wu et al., 2017), specialized ER-endosome contacts that regulate lysosome size in cell bodies (Özkan et al., 2021), and unique ER ladder structures in developing axons and dendrites (Benedetti et al., 2024; Zamponi et al., 2022). These regional specializations underscore the importance of examining ER structure and function across all neuronal regions. However, the cell biology and ultrastructure of the neuronal ER, particularly at presynaptic terminals, have been under-investigated. Conventional microscopy, used in most studies of neuronal ER structure, lacks the resolution needed to visualize individual ER tubules in small structures like presynaptic terminals (Chouhan et al., 2010; Oliva et al., 2020; Orso et al., 2009; Pérez-Moreno et al., 2023; Summerville et al., 2016; Yalçın et al., 2017). Other techniques such as electron microscopy (EM) have typically been limited to thin 2D sampling and cannot be used in live samples. These limitations make it difficult to distinguish the ER from synaptic vesicles at synapses and prevent the examination of live dynamics (Orso et al., 2009; Wu et al., 2017; Zhu et al., 2022).

We must overcome the technical challenges of studying neuronal ER as mutations in ER-shaping genes cause neurodegenerative disorders such as Hereditary Spastic Paraplegia (HSP) and Charcot Marie Tooth Disease, indicating that ER structure is uniquely essential for neuronal physiology (Boutry et al., 2019; Fowler et al., 2019; Öztürk et al., 2020). One essential ER tubule-shaping protein is Atlastin, a GTPase that regulates the homotypic fusion and tethering of ER tubules to form 3-way junctions (Lü et al., 2020). In humans, mutations in *Atlastin* underlie one of the most common forms of autosomal dominant HSP, highlighting the importance of examining its neuron-specific functions in regulating ER structure (Finsterer et al., 2012). In various experimental systems, *Atlastin* mutants exhibit numerous neuronal phenotypes, including defects in synaptic vesicle cycling and trafficking of neuronal cargoes (Gregorio et al., 2017), synapse development (Gregorio et al., 2017; Lee et al., 2009; Zhu et al., 2006), and axon regeneration (Rao et al., 2016). However, significant gaps remain in our understanding of how Atlastin-dependent changes in the structure and dynamics of neuronal ER lead to these functional defects. For example, using electron microscopy others found *Drosophila Atlastin* mutants exhibited shorter ER tubules in their motor neuron cell bodies(Orso et al., 2009) and diffuse localization of an ER lumen marker in their presynaptic terminals, attributed to extensive ER fragmentation (Summerville et al., 2016). However, this fragmentation has not been validated

by EM or by fluorescence using ER membrane markers. Further, in *C. elegans, Atlastin* mutant

ER networks in neuronal dendrites were not fragmented but were prone to retraction, leading to

disrupted microtubule structures and mitochondrial fission (Liu et al., 2019). Finally, a ladder-like

ER expansion was observed in spinal axons in a mouse model of Atlastin-dependent HSP (Zhu

et al., 2022). In contrast to these neuronal findings, knockdown of Atlastin in Hela or Cos-7 cells

leads to long and unbranched ER tubules (Hu et al., 2009, 2015; Niu et al., 2019). Thus, loss of

Atlastin causes distinct ER phenotypes depending on several factors, including cell type, which

region of the neuron was examined, and the reporter used to visualize the ER. To resolve the

functions of this vital protein, comprehensive imaging studies of ER organization in the different

regions of the neuron are needed, using complementary labeling strategies.

Another critical gap is our limited understanding of neuronal ER dynamics. High-

resolution imaging in non-neuronal cells has revealed transitions from ER tubules to ER sheets

(Lu et al., 2009; Sun et al., 2011), ER fragmentation (Espadas et al., 2019), tubule extension

and retraction (Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014), and alterations

in the number of 3-way junctions (Spits and Neefjes, 2021). These dynamics have been linked

to ER functions such as endosome or mitochondrial fission (Friedman et al., 2011; Hoyer et al.,

2018; Rowland et al., 2014) and calcium signaling (Crapart et al., 2023; Sun et al., 2011). In

neurons, ER rearrangements have been suggested to alter calcium compartmentalization

(Subramanian and Meyer, 1997), local protein synthesis (Deng et al., 2021; Schwarz and

Blower, 2016), organelle contacts (Lu et al., 2020) and transport within the ER (Scott et al.,

2021). However, the types and functions of neuronal ER dynamics, particularly in presynaptic

terminals, remain largely unknown due to the challenges of imaging ER networks over time in

small compartments. New insights into the mechanisms that regulate ER structure and ER

dynamics in neurons are crucial for understanding why and how mutations in ER shaping

proteins, such as Atlastin, predominantly affect the nervous system. Here, we addressed this

gap by examining the structure and dynamics of the ER at the *Drosophila* neuromuscular

junction (NMJ) of wild-type and *Atlastin* mutants using *in vivo* imaging and super-resolution

Airyscan microscopy.

Results

<u>High resolution imaging of the ER in *Drosophila* motor neuron cell bodies, axons, and presynaptic terminals.</u>

At presynaptic terminals of the *Drosophila* larval NMJ, the ER is highly sensitive to fixation, limiting imaging experiments to live samples, often acquired on upright microscopes using water dipping objectives (Chouhan et al., 2010; Oliva et al., 2020; O'Sullivan et al., 2012; Pérez-Moreno et al., 2023; Summerville et al., 2016; Yalçın et al., 2017). However, the axial resolution limit of typical water dipping objectives is >300 nm, which cannot distinguish the densely packed ER tubules at presynaptic terminals (Espadas et al., 2019; Wu et al., 2017). To overcome this limitation, we created a reusable microscopy imaging slide to capture live super-resolution images or movies of neuronal ER in *Drosophila* larval fillets through glass coverslips, utilizing high numerical aperture oil objectives and Airyscan microscopy (**Figure 1A**). Our system provides an axial resolution of ~170 nm after image processing (Del Signore et al., 2022). We imaged larval fillets using this imaging slide on both inverted and upright microscopes within 40-50 min from dissection, a time frame in which this preparation maintains muscle potential and the capacity for evoked neurotransmitter release, even once the axon is severed from the cell body (Feng et al., 2004).

To study the live dynamics of the ER, we used Vglut-Gal4 to express either the luminal ER marker BiP:sfGFP:HDEL (**Figure 1B-D**, **2C**) or the transmembrane ER marker tdTomato:Sec61β (**Figure 2B**) in *Drosophila* 3rd instar larval motor neurons. These markers were previously shown to co-localize at this synapse, indicating they are comparable reporters for visualizing synaptic ER (Summerville et al., 2016). Our imaging approach allowed more precise visualization of ER tubules and networks labeled by these reporters in cell bodies, axons, and presynaptic terminals compared to previously reported results (**Figure 1B-D, 2**). As previously observed using lower-resolution imaging (Summerville et al., 2016), we detected ER tubule structures in all neuronal regions examined, including ER tubules and ER sheets in cell bodies, and ER tubules in axons and presynaptic terminals. In axons, ER tubules were long, and we sometimes observed a ladder-like structure consisting of pairs of interconnected ER tubules joined by rungs or cross-bridges, similar to structures previously described in developing axons of mammalian cultured neurons and *Drosophila* dendrites (Benedetti et al., 2024; Zamponi et

al., 2022) (**Figure 1C**). At presynaptic terminals, ER tubules in some boutons appeared in a
highly interconnected basket-like network, while others were transversed by several parallel ER
tubules, similar to the ladder-like structures observed in axons (**Figure 1D**). Using this improved
imaging approach, we achieved high-resolution visualization of individual ER tubules within
networks of *Drosophila* presynaptic terminals, enabling detailed analysis of their structure and
dynamics over time.

We next examined ER dynamics in *en passant* and terminal boutons at the *Drosophila*
NMJ (**Figure 2A**). Terminal boutons have distinct cytoskeletal organization and signaling that
could regulate the dynamics of the ER differently (Deshpande et al., 2016; Fernandes et al.,
2023; Roos et al., 2000). To examine the dynamics of the ER at presynaptic terminals, we
collected Z stacks of single boutons expressing the ER membrane marker for 40 sec at 0.92
sec/stack (**Movie S1-S3**) or the luminal ER marker for 90 sec at 0.73 sec/stack (**Movie S4-S6**).
Both markers were distributed unevenly across the ER network (**Figure 2B-2C**). In particular,
the luminal ER marker exhibited numerous discrete round foci that moved within ER tubules with
fast dynamics (<1 sec) (**Figure 2C**). These foci were less frequently observed with the ER
membrane marker (**Figure 2B**, **bottom**), consistent with the model that proteins in the ER lumen
and ER membrane may differentially cluster and move within the ER (Holcman et al., 2018;
Obara et al., 2024). Conversely, static regions of high intensity (which we saw for both markers)
may reflect stacked ER tubules (Costantini et al., 2012).

We focused on the ER membrane marker to define dynamic events since its labeling
pattern had fewer confounding foci. Analysis revealed three distinct types of ER dynamics: (1)
ER budding events, in which a fragment of ER dissociates from the network; (2) Focus dynamics,
in which round foci move within the ER network; and (3) ER tubule dynamics, including overall
displacement of the tubule in the XY plane, as well as lengthening (extension) and shortening
(retraction) of the tubule (**Figure 2D**). We qualitatively categorized boutons as "static" if we
observed no change in ER network structure or "dynamic" if we observed at least one change.
On many occasions, we observed multiple types of ER dynamics within the same bouton. We
found that terminal boutons are highly dynamic (89%) while *en passant* boutons were
moderately dynamic (38%) (**Figure 2E**). This difference in dynamics suggests that the ER is
regulated differently in different regions of presynaptic terminals and could have specialized
functions.

Next, we characterized the frequency of the three main types of ER network dynamics at
terminal boutons, where we observed the most remodeling. Among terminal boutons, 93%
exhibited tubule movements, making this the most common type of ER remodeling (**Figure 2F**).
Less frequently, we observed that 28% of boutons had ER budding and 12% had moving foci.
We further classified tubule movements into four categories: displacement, extension, retraction,
and extension followed by retraction. The most common category was extension followed by
retraction, observed in 81% of terminal boutons (**Figure 2G**). Simple displacement occurred in
44% of boutons, 24% showed extension alone, and 18% showed retraction alone. These results
demonstrate our ability to capture diverse ER behaviors at high resolution in live *Drosophila*
motor neurons.

<u>Presynaptic terminals of *Atlastin* mutants form robust ER networks.</u>

To investigate the relationship between ER structure and function at synapses, we
examined mutants of Atlastin, a GTPase that regulates ER tubule fusion. *Drosophila* has a single
homolog while mammals have three Atlastin homologs, with Atlastin-1 enriched in the brain
(Rismanchi et al., 2008). The diverse neuronal phenotypes in *Atlastin* mutants, including defects
in synaptic growth, vesicle cycling, cargo trafficking, and axon regeneration, make it an excellent
model for understanding how ER architecture impacts cellular function (Audhya et al., 2007;
Gregorio et al., 2017; Lee et al., 2009; Rao et al., 2016; Zhu et al., 2006). We analyzed
transheterozygotes of an *Atlastin* null allele (*atl²*) and a deficiency removing the *Atlastin* locus to
avoid second-site effects (Lee et al., 2009). These *Atlastin* null mutants (hereafter referred to as
Atlastin mutants) were mainly pharate lethal and showed increased satellite boutons at larval
NMJs (**Figure 3A-B**), indicating defects in synaptic development as previously reported
(Gregorio et al., 2017; Lee et al., 2009).

Using confocal microscopy, Summerville et al. (2016) observed that the ER lumen marker
BiP:sfGFP:HDEL was diffusely distributed in *Atlastin* mutant presynaptic terminals and
concluded that the ER was extensively fragmented. To ask whether an ER membrane marker
would show similar fragmentation, we used spinning disk and Airyscan microscopy to examine
ER network structures at *Atlastin* mutant presynaptic terminals expressing the ER membrane
marker tdTomato:Sec61β. We found that tdTomato:Sec61β labeled intact ER networks in
Atlastin mutant synapses (**Figure 3A**, **Figure 3_Supplement**). Our results suggest that ER

networks exist in *Atlastin* mutant presynaptic terminals, and the previously observed defects in

synaptic function are likely due to more subtle ER structural or dynamic properties rather than

fragmentation. However, we found that the marker tdTomato:Sec61β exhibited lower

fluorescence levels in *Atlastin* mutants, as quantified by its mean intensity within the entire

presynaptic terminal in fixed preparations (**Figure 3C**). This reduction in ER membrane marker

intensity has also been observed in other HSP mutants, suggesting this is a common feature of

ER shaping mutants and could indicate changes in ER membrane composition, integrity, or

tubule thickness (Pérez-Moreno et al., 2023). We also noted that the coefficient of variation

(CoV) of tdTomato:Sec61β signal was significantly higher in *Atlastin* mutants (**Figure 3D**),

suggesting that it is more concentrated in subsynaptic structures compared to wild-type

synapses (notably the opposite of what would be expected for fragmentation).

Next, we measured the levels of tdTomato:Sec61β in various motor neuron regions to

determine if its reduced levels are specific to presynaptic terminals or a cell-wide phenotype

(**Figure 4**). Using Vglut-Gal4 to label motor neurons, we measured clusters of cell bodies in the

ventral ganglion and axon bundles extending towards body-wall muscles (**Figure 4A-B**). *Atlastin*

mutants have significantly increased tdTomato:Sec61β intensity levels in axon bundles (**Figure**

4C) and unchanged levels in cell body clusters (**Figure 4F**). We also found that the volume

occupied by tdTomato:Sec61β-labeled structures was significantly increased in axon bundles

and decreased in cell body clusters of *Atlastin* mutants. Finally, we observed a significant

decrease in the CoV of the tdTomato:Sec61β signal in both axon bundles and cell body clusters,

suggesting less concentrated ER structures (the opposite of what we observed at presynaptic

terminals). Our results suggest that the changes observed in cell body clusters, axon bundles,

and presynaptic terminals are local, indicating that the ER has neuronal region-specific

phenotypes regulated by Atlastin. These changes in tdTomato:Sec61β distribution could reflect

alterations in ER structure or ER membrane protein density.

Atlastin mutants show mild defects in synaptic ER dynamics

We hypothesized that Atlastin could regulate synaptic function by controlling the

dynamics of the ER network. We captured rapid time-lapse images of the ER labeled with

tdTomato:Sec61β in *en passant* and terminal boutons at 0.92 sec/stack (**Figure 2D-G, Figure**

5A, **Movies 7-14**). We blinded movies from controls and *Atlastin* mutants to qualitatively

categorize boutons as "dynamic" or "static", as described above (**Figure 2E**). Compared to the

89% of terminal boutons in control animals that exhibited dynamics (previously shown in **Figure**

2B, E), we found a small but significant reduction in dynamic boutons in *Atlastin* mutants (76%),

with a concomitant increase in the fraction of non-dynamic boutons (24%) (**Figure 5B**). In *en*

passant boutons, we observed similar ER dynamics in controls (62% static/38% dynamic) and

Atlastin mutants (69% static/31% dynamic). These findings indicate that *Atlastin* mutants exhibit

mild defects in ER dynamics, specifically at terminal boutons, with no significant changes in *en*

passant boutons. This suggests that Atlastin's ER tubule tethering and fusion roles are critical in

highly dynamic ER networks like those found in terminal boutons.

We performed a further comprehensive analysis of ER dynamics in *Atlastin* mutants,

examining multiple structural rearrangements, including tubule dynamics, ER budding, and the

movement of concentrated protein foci. Surprisingly, despite Atlastin being the sole homolog in

Drosophila and a key player in ER membrane fusion, we found only modest changes in ER

structure and dynamics. While we observed slight shifts in the relative frequencies of different

dynamic events (**Figure 5C-D, Figure 5A-Supplement**), none of these changes reached

statistical significance, and the overall dynamic capacity of the ER network remained intact

(**Figure 5B-Supplement**). The subtlety of these phenotypes suggests either substantial

functional redundancy with other ER-shaping proteins or the existence of compensatory

mechanisms that maintain ER organization and dynamics in neurons upon loss of *Atlastin*.

Moreover, given the numerous synaptic defects previously reported in *Atlastin* mutants, our

findings suggest that these phenotypes may arise from defects in ER function rather than gross

changes in ER structure or dynamics.

Atlastin mutant presynaptic terminals exhibit distinct BiP:sfGFP:HDEL and tdTomato:Sec61β

distributions.

Our results showing intact ER networks in *Atlastin* mutants using an ER membrane

marker sharply contrast with those from a prior study examining *Atlastin* mutants using the

luminal marker BiP:sfGFP:HDEL, where its diffuse distribution was interpreted as ER

fragmentation (Summerville et al., 2016). To further explore this discrepancy, we used Airyscan

microscopy to re-examine BiP:sfGFP:HDEL in *Atlastin* mutants. Unlike its strict localization to

the ER network in control synapses, we reproduced the observation that a significant fraction of

BiP:sfGFP:HDEL was diffusely localized in *Atlastin* mutant synapses, even at the resolution of

Airyscan microscopy. Further, the marker filled the interiors of boutons and extended to their

periphery, suggesting that it may be mislocalized to the cytosol instead of within ER fragments

(**Figure 6A-B**).

We identified two distinct ER network phenotypes in *Atlastin* mutants: NMJs with "Partial

loss" had both diffuse BiP:sfGFP:HDEL and an identifiable ER network structure, while NMJs

with "Complete loss" had a more severe phenotype with no visible ER network structures. We

blinded our images and categorized control and *Atlastin* mutant datasets as "Partial loss,"

"Complete loss," or "No phenotype" (**Figure 6C**). In *Atlastin* mutants, we found that 45% of NMJs

exhibited partial loss, 55% exhibited complete loss, and 10% had no phenotype. We observed

partial loss and complete loss phenotypes in both anterior (segment A2) and posterior (segment

A5) muscle segments, in Is (phasic) and Ib (tonic) motor neuron subtypes, and in neurons

innervating multiple different body wall muscles, suggesting that it reflects a widespread

phenomenon in NMJs of *Atlastin* mutants (**Figure 6 - Figure 6-Supplement**). Presynaptic

terminals in *Atlastin* mutants show similar mean intensities of the luminal ER marker compared

to controls, suggesting that the marker's redistribution in mutant neurons does not affect its

overall levels (**Figure 6D-E**). However, we observed a significant change in the CoV in *Atlastin*

mutants due to the redistribution of the luminal marker (**Figure 6F**). Along with our earlier finding

that the ER membrane marker, tdTomato:Sec61β, does not exhibit fragmentation, these results

suggest the possibility that the luminal ER marker is displaced to the cytosol.

Atlastin mutants may be particularly sensitized to overexpression of the BiP:sfGFP:HDEL

marker because we could not assess the co-localization of BiP:sfGFP:HDEL and

tdTomato:Sec61β, as very few larvae co-expressing these markers survived. The few that did

had grossly defective NMJ morphologies and large aggregates of both markers, even in wild-

type flies (data not shown). Further, control adult flies expressing BiP:sfGFP:HDEL were

unhealthy and could not be maintained as a stock, suggesting that this marker is toxic.

Next, we asked whether the luminal ER marker displacement phenotype is specific to all

neuronal regions or specific to presynaptic terminals. First, we examined the ER in body wall

muscles of *Atlastin* mutants by overexpressing the BiP:sfGFP:HDEL with the muscle driver

BG57-Gal4. We did not find the luminal ER marker displaced to the cytosol (**Figure 7, right**).

However, as noted in earlier studies (Lee et al., 2009), we observed a decrease in network
complexity. Next, we examined the cell body and axons of *Atlastin* mutants to determine if the
displacement of the luminal ER marker was a global or local phenotype within neurons. As
previously reported (Summerville et al., 2016), we did not observe a diffuse signal for the luminal
ER marker in cell bodies or distal axons (within ~30μm of the synapse) (**Fig 7, middle + left**).
However, we did observe ER fragmentation in some axons (**Fig 7, left**). These results indicate
that the displacement of BiP:sfGFP:HDEL in *Atlastin* mutants is a synapse-specific phenotype.
302
303 <u>The luminal ER phenotype in *Atlastin* mutants is progressive</u>
304
Atlastin dysfunction is linked to the neurodegenerative disorder HSP, in which patients
progressively lose neuronal function over time (Blackstone, 2018, 2012; Blackstone et al., 2011).
Drosophila Atlastin mutants recapitulate progressive motor dysfunction as observed in HSP
patients (Gregorio et al., 2017). We examined 1[st] instar larvae to test the hypothesis that the
partial loss phenotype may similarly progress to the complete loss phenotype through larval
development. We categorized control and *Atlastin* mutant datasets as described above (**Figure
6**). We anticipated that a progressive phenotype would decrease the proportion of NMJs showing
BiP:sfGFP:HDEL displacement and an increase in presynaptic terminals displaying no
phenotype. Indeed, we observed that the percentage of NMJs with complete loss increased from
12.5% in 1[st] instar to 55% in 3[rd] instar, while NMJs with no phenotype decreased from 38% to
10% across these developmental stages (compare **Figure 8A-C** to **Figure 6**). These results
suggest that displacement of the luminal ER marker progresses over time and that this
phenotype is initiated early in neuronal development.
318
319 *Atlastin* mutants have disrupted luminal ER protein dynamics
320
To examine ER network continuity and the dynamic properties of luminal ER proteins in
Atlastin mutants, we used fluorescence recovery after photobleaching (FRAP) to track
BiP:sfGFP:HDEL mobility at presynaptic terminals (**Figure 9A**). In this analysis, the rate and
extent of fluorescence recovery provide key insights into protein behavior: continuous ER
networks show robust recovery due to the movement of unbleached proteins across the
connected network, while fragmented ER networks fail to recover due to physical barriers
between compartments (Dayel et al., 1999; Lippincott-Schwartz et al., 2018). Further, if the

BiP:sfGFP:HDEL protein had leaked into the cytosol, we would expect rapid recovery due to the
higher diffusion rates of cytosolic proteins compared to ER-localized proteins (Dayel et al.,
1999). Our FRAP analysis examined single synaptic boutons co-expressing BiP:sfGFP:HDEL
and RFP:mCD8, which labels the neuronal plasma membrane. We photobleached *en passant*
boutons in the middle of a synaptic branch to ensure BiP:sfGFP:HDEL could recover from ER
networks surrounding the FRAP region. We recorded movies of a single NMJ slice,
simultaneously imaging both markers. After photobleaching with a 405nm laser, we captured
recovery dynamics by imaging every 10 seconds for 2 minutes (**Figure 9, Movies 15-19**).
336
We analyzed the recovery after photobleaching of RFP:mCD8 and BiP:sfGFP:HDEL to
characterize protein dynamics in *Atlastin* mutant presynaptic terminals. RFP:mCD8 served as a
control because it showed similar plasma membrane distribution patterns at synaptic boutons in
both control and *Atlastin* mutants, suggesting that plasma membrane organization was
unaffected and would exhibit comparable dynamics across all genotypes. While the recovery of
RFP:mCD8 could not be fitted to a non-linear equation due to the absence of a plateau, its similar
recovery across genotypes indicates no significant changes in its diffusion at the plasma
membrane (**Figure 9B-C**). We next analyzed BiP:sfGFP:HDEL recovery using a one-association
non-linear model, focusing first on the plateau to estimate the mobile fraction of the luminal ER
marker in the FRAP region (**Figure 9B, D**). We found that *Atlastin* mutants with partial loss of
ER networks had the least recovery, with a mobile fraction of 50.61%. By contrast, controls and
Atlastin mutants with complete loss had a higher mobile fraction of 65.73% and 66.34%,
respectively. We also measured how quickly BiP:sfGFP:HDEL recovered by calculating the time
to reach 50% of maximum recovery. *Atlastin* mutants with partial loss exhibited the slowest
BiP:sfGFP:HDEL recovery, reflected by a longer half-life (38.26 sec) compared to controls
(26.91 sec), while those with complete loss had the fastest recovery (14.62 sec). The reduced
mobile fraction in partial loss *Atlastin* mutants likely reflects discontinuous ER networks (**Figure
9B**, blue arrows). In contrast, the rapid recovery and absence of visible ER structures in complete
loss *Atlastin* mutants suggests that BiP:sfGFP:HDEL is largely displaced into the cytosol. Our
analyses indicate that Atlastin dysfunction impairs luminal ER protein diffusion and subcellular
localization. Surprisingly, these defects occur while maintaining relatively normal synaptic ER
network structure (**Figure 3-4**) and dynamics (**Figure 5**). These findings reveal a novel
pathogenic mechanism where Atlastin dysfunction primarily impacts ER protein distribution

rather than ER network morphology, potentially disrupting synaptic function through the

mislocalization of essential ER proteins.

<u>Luminal proteins are displaced to the cytosol at presynaptic terminals of *Atlastin* mutants</u>

A fundamental question emerged from our observations: does the altered

BiP:sfGFP:HDEL distribution in *Atlastin* mutants arise from ER fragmentation or genuine protein

displacement to the cytosol? Previous research shows that ER network morphology relies on a

delicate balance between ER shaping proteins; disrupting this balance may result in ER

fragmentation (Wang et al., 2016). In contrast, the displacement of luminal proteins to the cytosol

suggests changes in the ER integrity and potential activation of ER stress responses (Merighi

and Lossi, 2022). To definitively distinguish between these models, we conducted two

independent experiments to track BiP:sfGFP:HDEL sub-cellular localization at presynaptic

terminals. First, we examined whether BiP:sfGFP:HDEL in *Atlastin* mutants is sensitive to the

mild detergent digitonin, which gently permeabilizes the plasma membrane and extracts proteins

from the cytosol but not the ER (Le Gall et al., 2004). Second, we examined the direct interaction

between BiP:sfGFP:HDEL and a plasma membrane-tethered nanobody, which would only occur

if the protein was displaced to the cytosol.

We hypothesized that digitonin treatment would lead to loss of fluorescence in *Atlastin*

mutants, proportional to the fraction of the luminal ER marker protein in the cytosol (**Figure 10B-**

Supplement). We treated presynaptic terminals for 5 min with digitonin (**Figure 10A-**

Supplement) and imaged the same presynaptic terminal (expressing either cytosolic GFP or

BiP:sfGFP:HDEL) before and after digitonin treatment. We also labeled our live samples with α-

HRP antibodies to identify presynaptic terminals (Snow et al., 1987). We measured the fraction

change in mean intensity to determine the effects of digitonin treatment on cytosolic and ER

markers (**Figure 10C-Supplement**). We observed a significant loss in mean fluorescence of

cytosolic GFP following digitonin treatment. Control presynaptic terminals expressing the luminal

ER marker BiP:sfGFP:HDEL became highly fragmented, possibly due to osmolarity changes in

the cytosol, which are known to promote ER fragmentation (King et al., 2020). Despite this

fragmentation, most ER remained within the synaptic terminal, suggesting limited extraction or

permeabilization by this treatment. In half the control samples, the marker redistributed to bright

puncta and showed an unexpected increase in mean intensity. One interpretation is that digitonin

treatment allows the exchange of small molecules into the ER (Le Gall et al., 2004), potentially

altering the fluorescent properties of sfGFP. The other half of the samples had slightly reduced

mean intensity following digitonin treatment, possibly due to moderate loss of ER fragments. We

next analyzed *Atlastin* mutants categorized as "Partial loss" or "Complete loss". Digitonin

treatment yielded no significant differences between controls and *Atlastin* mutants with partial

loss. We also observed that *Atlastin* mutant synapses exhibiting partial loss had puncta following

digitonin treatment that were similar to the control. This is likely because the partial loss

presynaptic terminals have some ER network visible with the luminal marker. In contrast, we

observe a significant decrease in fluorescence intensity following digitonin treatment in *Atlastin*

mutant presynaptic terminals with complete loss, which do not exhibit visible ER networks, as

indicated by the luminal marker. These results suggest that luminal ER molecules in *Atlastin*

mutants are likely displaced to the cytosol.

From the digitonin experiments, we cannot rule out the possibility that BiP:sfGFP:HDEL

resides in small ER membrane fragments that easily diffuse out of permeabilized plasma

membranes. We hypothesized that if BiP:sfGFP:HDEL is displaced to the cytosol in *Atlastin*

mutants, it would interact with plasma membrane proteins topologically oriented toward the

cytosol. However, if BiP:sfGFP:HDEL were within a small membrane-bound ER compartment,

it could not interact with a cytosolically oriented ligand. To test this, we co-expressed

BiP:sfGFP:HDEL with a plasma membrane-tethered GFP nanobody fused to RFP (referred to

as morphotrap$_{Int,}$) (Harmansa et al., 2017)). The GFP nanobody in morphotrap$_{Int}$ faces the

cytosol, even when tethered to membranes, allowing it to interact only with cytosolic

BiP:sfGFP:HDEL. Given that the morphotrap$_{Int}$ is localized to the plasma membrane, it should

cause the relocation of cytosolic BiP:sfGFP:HDEL, but not luminal BiP:sfGFP:HDEL. Second,

since a membrane-tethered nanobody is likely to have slower diffusion than the putatively

cytosolic BiP:sfGFP:HDEL, morphotrap$_{Int}$ co-expression would be predicted to reduce cytosolic

BiP:sfGFP:HDEL dynamics, but not luminal BiP:sfGFP:HDEL dynamics (**Figure 10A**).

First, we blinded our samples and categorized control and *Atlastin* mutants as "No

phenotype", "Partial loss" and "Complete loss", as we had done previously (**Figure 6C**). We

found that *Atlastin* mutants co-expressing morphotrap$_{Int}$ and BiP:sfGFP:HDEL had 47% of NMJs

with partial loss, 3% with complete loss, and 50% with no phenotype (**Figure 10B**). This

redistribution   is   consistent   with   morphotrap$_{Int}$ binding   and   relocalizing   cytosolic

BiP:sfGFP:HDEL. Interestingly, rather than accumulating at the plasma membrane, the overall
levels of BiP:sfGFP:HDEL were reduced, suggesting that the displaced protein is targeted for
degradation (**Figure 10C**). Further, the reduction in partial and complete loss phenotypes
compared to BiP:sfGFP:HDEL expression alone (**Figure 6C vs 10B**) indicates that sequestering
BiP:sfGFP:HDEL at the plasma membrane uncovers residual ER networks masked by the
diffuse cytoplasmic signal. To test the specificity of this effect, we co-expressed
BiP:sfGFP:HDEL with RFP:mCD8, which has the same protein domain composition as
morphotrap$_{Int}$ but lacks the GFP nanobody. BiP:sfGFP:HDEL levels were unaffected in those co-
expressing RFP:mCD8 in *Atlastin* mutants and controls. We also found that 42% of NMJs had
partial loss, 50% had complete loss, and 8% had no phenotype (**Figure 10B**), similar to what we
observed when only BiP:sfGFP:HDEL was overexpressed in *Atlastin* mutants (**Figure 6C**). By
contrast, in wild-type controls, co-expression of BiP:sfGFP:HDEL with either morphotrap$_{Int}$ or
mCD8:RFP did not result in any relocalization of BiP:sfGFP:HDEL. Our results suggest that
BiP:sfGFP:HDEL protein at presynaptic terminals is displaced to the cytosol and binds to
morphotrap$_{Int}$ in *Atlastin* mutants, leading to its altered distribution and removal.

Next, we performed a FRAP experiment to test the dynamics of BiP:sfGFP:HDEL in the
presence of morphotrap$_{Int}$. First, we examined the dynamics of morphotrap$_{Int}$ itself following
photobleaching. We found that morphotrap$_{Int}$ had very slow recovery and plateaued at 20% in
wild-type and 17% in *Atlastin* mutants (**Figure 10D-E**). We hypothesized that the slow diffusion
of morphotrap$_{Int}$ would restrict the movement of any cytosolic BiP:sfGFP:HDEL in *Atlastin*
mutants. Indeed, we found that wild-type expressing morphotrap$_{Int}$ and BiP:sfGFP:HDEL had a
mobile fraction of 57%, while *Atlastin* mutants had a mobile fraction of 38% (**Figure 10F**). By
contrast, in *Atlastin* mutants co-expressing BiP:sfGFP:HDEL and non-membrane-targeted
RFP:mCD8, NMJs with partial loss had a mobile fraction of 51%, and those with complete loss
had a plateau of 66% (**Figure 9**). We hypothesize that the reduction in the mobile fraction in the
presence of morphotrap$_{Int}$ results from the binding of BiP:sfGFP:HDEL in the cytosol to
morphotrap$_{Int}$ in *Atlastin* mutants. There was a minor decrease to 57% in wild-type larvae co-
expressing BiP:sfGFP:HDEL and morphotrap$_{Int}$, while those co-expressing BiP:sfGFP:HDEL
with RFP:mCD8 plateaued at 66%. These results suggest the presence of a small cytosolic
population of BiP:sfGFP:HDEL in controls. Finally, we examined the recovery kinetics of
BiP:sfGFP:HDEL in larvae co-expressing morphotrap$_{Int}$. We found the recovery in controls is
slower, with a half-life of 53.23 s, while *Atlastin* mutants had a faster recovery with a half-life of

45.76 s. The faster half-life in *Atlastin* mutants could reflect that a small pool of BiP:sfGFP:HDEL
remains mobile, perhaps contained within a discontinuous section of the ER. Our findings
provide multiple lines of evidence that BiP:sfGFP:HDEL is displaced from the ER lumen to the
cytosol in *Atlastin* mutants, potentially reflecting disruption to ER integrity and function at
presynaptic terminals.

Discussion

467   This study revealed the architecture and dynamic behaviors of *Drosophila* neuronal ER
networks through a super-resolution *in vivo* imaging approach that can be adapted to visualize
fluorescently labeled proteins at high resolution in any larval tissue. Using this approach, we
provided the first characterization of presynaptic ER network dynamics *in vivo*, establishing a
foundation for investigating the functional significance of structural rearrangements of the ER.
Our analysis of *Atlastin* mutants uncovered an unexpected phenotype: while these mutants
maintain largely intact ER networks with only minor defects in dynamics, they exhibit
displacement of luminal ER proteins to the cytosol of presynaptic terminals but not in other
neuronal regions. These findings reveal that Atlastin serves distinct, synapse-specific functions
in maintaining both ER structure and function, providing mechanistic insight into the synaptic
deficits previously observed in *Atlastin* mutants.

ER network dynamics at presynaptic terminals

481   We observed three categories of ER dynamics at presynaptic terminals: (1) budding and
fusion of vesicular structures, (2) ER tubule dynamics, and (3) dynamics of foci that are retained
within the ER network. Disruptions in these dynamics, as seen in *Atlastin* mutants, could impair
synaptic signaling and contribute to neurodegenerative diseases linked to ER dysfunction. ER
network dynamics may depend on Atlastin-mediated ER tubule tethering and fusion to distribute
lipids and ER membrane proteins throughout the network to ensure proper distribution and
prevent fragmentation. Future work is needed to examine the specific functions of these
dynamics at presynaptic terminals, but we speculate below on some potential roles:

490   First, ER-derived vesicles could act as localized calcium signaling hubs (Burgoyne et al.,
491   2015) or serve as sites for compartmentalized local translation if they carry ribosomes (Carter et

al., 2020; Deng et al., 2021). The synaptic ER vesicle-like structures we observed are unlikely to be involved in the secretory pathway or ER-phagy (Chino and Mizushima, 2020) because we observe these vesicles re-fusing with the ER. We speculate the ER-derived vesicles observed may be REEP1-4 positive, which regulates ER vesicle formation in mammalian cells (Shibata et al., 2024). Notably, while overexpression of Atlastin can regulate REEP1-4 vesicle fusion, it is not essential for vesicle formation. Second, ER tubule dynamics may play multiple roles, such as maintaining the tension of the ER network (Espadas et al., 2019; Niu et al., 2019) and forming ER-organelle contacts (Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014). ER contacts occur at the tips and edges of ER tubules, and they have diverse functions, such as promoting mitochondrial and endosome fission (Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014). To understand how these functions are regulated at presynaptic terminals, future studies could examine ER tubule interactions with other organelles using co-localization experiments or ER-organelle sensors. Finally, we identified foci dynamics within ER networks. We hypothesize that foci emerge due to changes in the dynamics of protein clusters at the ER membrane or within the lumen. Foci dynamics occur at higher levels with the luminal ER marker in contrast to the ER membrane marker. This differential distribution suggests distinct organizational principles for luminal versus membrane proteins in presynaptic ER.

Distinct Membrane and Luminal ER Phenotypes in *Atlastin* Mutants Suggest Local Stress Responses

Using the membrane marker tdTomato:Sec61β, we discovered that *Atlastin* mutants have robust ER networks at presynaptic terminals. The relatively mild structural phenotypes stand in sharp contrast to the diffuse distribution we and others observe for the luminal marker BiP:sfGFP:HDEL, and argue against excessive ER fragmentation in *Atlastin* mutants. Despite being the sole Atlastin homolog in flies, we found that loss of Atlastin causes surprisingly subtle changes in ER structure and dynamics. However, we did observe specific defects with our FRAP experiments, which revealed *Atlastin* mutants have discontinuous ER networks. The subtle nature of these defects also suggests that other proteins may partially compensate for Atlastin's function in maintaining ER structure and dynamics at presynaptic terminals.

Considering the distinct phenotypes displayed by luminal and membrane ER markers, which one best represents the *Atlastin* mutant phenotype in the absence of a marker? We favor the hypothesis that the true *Atlastin* ER phenotype is more likely to be similar to the

tdTomato:Sec61β distribution, for the following reasons: (1) tdTomato:Sec61β-expressing
animals still show Atlastin-associated functional defects such as defective synaptic growth,
indicating that Atlastin-dependent ER functions are disrupted without dramatic defects in ER
structure; (2) The BiP:sfGFP:HDEL phenotype becomes progressively more severe throughout
larval development, suggesting this marker's displacement is a secondary effect rather than a
direct reflection of ER architecture. Future studies in other ER-shaping mutants should examine
both ER membrane and ER lumen markers, as specific defects could affect one and not the
other, as we observed in *Atlastin* mutants.

Why, then, might *Atlastin* mutants displace luminal proteins to the cytosol? A key
observation that constrains possible mechanisms is that this displacement occurs specifically at
presynaptic terminals - the luminal ER marker remains properly localized in cell bodies, axons,
and other cell types such as muscles. This spatial specificity argues against several potential
global mechanisms: First, overall disruption of ER integrity seems unlikely to explain the
phenotype; while *Atlastin* mutants show reduced levels of the ER membrane marker Sec61β
and altered network structure at presynaptic terminals, catastrophic loss of ER membrane
integrity would likely release calcium and other ER components into the cytosol, which would be
incompatible with cell survival (Wang et al., 2011). Second, general defects in protein
translocation are unlikely, since these would affect all neuronal compartments, especially the
cell body where most protein translation occurs (Glock et al., 2021; McKibbin et al., 2012). Third,
we found no evidence for apoptosis-induced ER permeability, as *Atlastin* mutant neurons lack
the characteristic debris observed after neuronal death at the NMJ (Aponte-Santiago et al., 2020;
Graf et al., 2011).

Instead, the presynaptic specificity suggests more localized mechanisms. One possibility
is compartment-specific activation of ER stress responses like UPR (Untranslated Protein
Response), ERAD (ER-associated degradation) or ERCYS (ER-to-cytosol signaling)
(Christianson et al., 2023; Igbaria et al., 2019; Lajoie and Snapp, 2020; Read and Schröder,
2021; Sicari et al., 2021; Wu and Rapoport, 2018). These pathways can actively transport
luminal proteins to the cytosol through distinct mechanisms: in UPR and ERAD, proteins are
targeted for proteasomal degradation, while in ERCYS, they can acquire novel functions in the
cytosol. Previous work has shown that *Atlastin* mutants and other HSP-linked genes have
activated UPR (Lai et al., 2014; O'Sullivan et al., 2012; Zhao et al., 2016). In the case of our

overexpressed BiP:sfGFP:HDEL marker, either pathway could explain its accumulation in the
cytosol. If UPR or ERAD is activated, BiP:sfGFP:HDEL might be continuously displaced from
the ER but accumulate when the proteasome cannot keep up with its degradation. Alternatively,
if ERCYS is activated, BiP:sfGFP:HDEL might accumulate in the cytosol as part of a regulated
signaling response. We cannot distinguish between these possibilities, but multiple lines of
evidence support the general activation of ER stress pathways in *Atlastin* mutants. First, the
phenotype progresses from 1st to 3rd instar larvae, consistent with gradually activated stress
responses. Second, synaptic terminals show a specific reduction in Sec61β levels, suggesting
local defects in protein homeostasis. Given these observations, what might trigger an ER stress
response in *Atlastin* mutants? The underlying mechanism remains to be determined, but the
progressive nature of the phenotype and the eventual failure of luminal ER protein retention
suggest a gradual deterioration of ER homeostasis at synaptic terminals. Future work will be
needed to identify the specific cellular changes that initiate this stress response.
569
ER network structure and function are compartmentalized across neuronal regions

Our findings reveal distinct requirements for Atlastin across neuronal regions. At
presynaptic terminals, loss of Atlastin leads to the displacement of luminal proteins and reduced
levels of Sec61β. This reduction in Sec61β suggests other ER membrane proteins, such as ER
tethers, could also be affected, potentially disrupting ER-dependent processes like endosome
trafficking that rely on proper ER-organelle contacts. This reduction in Sec61β mirrors findings
in *Drosophila* null mutants of Reticulon-1, another ER shaping and HSP-linked gene, in which
levels of STIM and Stukorf are also decreased on ER tubules (Pérez-Moreno et al., 2023). Yet,
a key distinction emerges as Reticulon-1 null mutants maintain normal luminal ER protein
localization, suggesting unique features of *Atlastin* mutant ER networks. These region-specific
phenotypes highlight fundamental differences in ER organization and function across neuronal
regions. Further evidence for neuronal ER compartmentalization comes from studies of
mammalian spinal axons, where Atlastin-fusion incompetent mutants showed ladder-like ER
expansion (Zhu et al., 2022). We observed similar ladder-like structures with three-way junctions
in both control and *Atlastin* mutant axons labeled with tdTomato:Sec61β, indicating that Atlastin
is not solely responsible for ER tubule fusion in axons. This neuronal-specific organization
contrasts with the long, unbranched ER tubules found in Atlastin-deficient Hela or Cos-7 cells
(Hu et al., 2009; Niu et al., 2019), suggesting neurons possess unique compensatory tubule-

fusion machinery. Future work should comprehensively examine the common and unique
defects associated with HSP gene dysfunction across neuronal compartments.

Methods

Drosophila stocks

Drosophila melanogaster was cultured on a standard medium at 22–25°C. Detailed genotypes
and sexes used for experiments are described in Table S1. Stocks used in this study
include: Vglut-Gal4 (Bloomington *Drosophila* Stock Center (BDSC_24635), BG57-Gal4 (gift
from Vivian Budnik), atl^2 (gift from James McNew), third chromosome deficiency (BDSC_7948),
UAS-BiP:sfGFP:HDEL (BDSC_64748), 20XUAS-tdTomato:Sec61β (BDSC_64746), UAS-
CD8:RFP (BDSC_27391) and UAS-morphotrap$_{Int}$ (gift from Chi-Kuang Yao).

Live imaging of luminal and membrane ER markers
1st and wandering 3rd instar larvae were dissected in Ca^{2+}-free HL3.1 (Feng et al., 2004) and
axons were severed from the central nervous system. Dissections were performed in the live
imaging slide described below. Larvae were imaged at room temperature with a 63X (n.a. 1.4)
oil immersion objective, using an Airyscan LSM 880 microscope in super-resolution mode and
Zen Black software. Raw image stacks were processed in Zen Blue or Zen Black using 3D
Airyscan processing with automatic settings.

For 3rd instar larvae, Z-stacks of soma clusters, axons or presynaptic type Ib terminals at muscle
4 in segment A5 (unless otherwise noted) were acquired to measure the levels of luminal ER
markers, to categorize ER network phenotypes, or for morphological measurements. For 1st
instar larvae, live larval fillets were stained with 1:250 dilution of 1.5 mg/ml HRP Alexa Fluor 647
(Jackson ImmunoResearch) in HL3.1 for 5 min followed by a 1 min wash. Next, Z-stacks of 1st
instar presynaptic terminals at muscle 4 in segment A3-A5 were acquired and used to categorize
ER network phenotypes.

To record the dynamics of the luminal ER marker BiP:sfGFP:HDEL, 16-bit Z-stacks of a single
bouton (8 slices with 0.5 micron spacing) were collected for 100 frames at a frame rate of 0.73
seconds/stack. To record ER dynamics with the membrane ER marker tdTomato:Sec6β, 8-bit

Z-stacks of a single bouton (20 slices with 0.1850 micron spacing) were collected for 60 frames

at a frame rate of 0.92 seconds/stack.

<u>Construction of *in vivo* imaging slide for *Drosophila* larvae</u>

To make the reusable imaging slide, Press-To-Seal Silicone Isolators (Grace Bio-labs; CQS-

13R-2.0) were mounted onto rectangular glass slides, which then were filled from their centers

to their tops with Krayden Dow Sylgard 184 Silicone (Thermo Fisher Scientific). Each slide with

a Press-To-Seal Silicone Isolator was leveled to ensure the silicone cured evenly at room

temperature. Minutien pins (Fine Science Tools) were cut with nail clippers and used to stretch

the larvae and fix it in place. Pins were pressed into the cured Sylgard 184 Silicone and a few

drops of hemolymph-like solution HL3.1 were added. Next, a coverslip was placed on top of the

dissected larvae, each corner was gently pressed down, and excess HL3.1 was removed with a

Kimwipe, thus sealing the coverslip to the silicone isolator.

<u>Fixed tissue staining</u>

Wandering 3rd instar larvae were dissected in Ca^{2+}-free HL3.1 and fixed for 17 min in Ca^{2+}-free

HL3.1 containing 4% PFA. Larvae were blocked and permeabilized overnight in PBS containing

0.25% Saponin, 2.5% normal goat serum (NGS), 2.5% bovine serum albumin (BSA), and 0.1%

sodium azide. Fixed larvae (**Figure 3**) were stained with 1:500 FluoTag®-X4 anti-RFP (#N0404,

Nanotag Biotechnologies) at 4°C for 24 hrs and with 1:100 HRP Alexa Fluor 647 at room

temperature for 2hrs. Stained larvae were mounted in ProLong Diamond Antifade Mountant

(#P36970; Thermo-Fisher Scientific, Waltham, MA, USA). Z-stacks were collected of *Drosophila*

larval NMJs in muscle 4 of segment A5 at room temperature on a Nikon Ni-E upright microscope

equipped with 100x (n.a. 1.45) oil immersion objective, a Yokogawa CSU-W1 spinning-disk

head, and an Andor iXon 897U EMCCD camera. Images were collected using Nikon Elements

AR software.

<u>FRAP experiment</u>

Imaging was conducted using a Nikon super-resolution spinning disk SoRA system in W1-

spinning disk mode, equipped with an Apo 60x Oil DIC N2 objective and set to a 12-bit depth.

We photobleached boutons in the middle of a synaptic branch to ensure BiP:sfGFP:HDEL could

recover from both the top and bottom of the photobleached region. To capture the rapid recovery

of BiP:sfGFP:HDEL, we imaged a single slice of NMJs at muscle 4 and sequentially acquired

488 and 561 channels with an exposure time of 100ms. For all NMJs, we photobleached single

boutons within a 6.8x7.8 microns region in the middle of neuronal branch. Each movie acquired

used an ND sequence acquisition consisting of three phases: (1) we acquired five frames at

maximum speed (100Mhz) before photobleaching; (2) we photobleached BiP:sfGFP:HDEL and

RFP:mCD8 or morphotrap$_{Int}$ using a 405 nm depletion laser with a dwell time of 100µs for a

duration of 2s; and (3) we imaged the recovery of both BiP:sfGFP:HDEL and RFP:mCD8 or

morphotrap$_{Int}$ with a 10s interval for 2 minutes for experiments in **Figure 9** and 5 minutes for

experiments in **Figure 10**.

Digitonin experiments

Digitonin experiments were conducted in live fillets from larvae expressing the luminal ER

marker, UAS-BiP:sfGFP:HDEL, under the control of the neuronal driver Vglut-GAL4. Fillets were

stained with 1:500 1.5 mg/ml HRP Alexa Fluor 594 (Jackson ImmunoResearch) or 1:250 HRP

Alexa Fluor 647 in HL3.1 for 5 min, followed by a 1 min HL3.1 wash. Next, pre-digitonin Z-stacks

were acquired from muscle 4 NMJs from segments A3 to A5 (both hemisegments) using the

spinning disk confocal microscope described above. Fillets were subsequently incubated with

0.05% digitonin in HL3.1 for 5 min, then rinsed with HL3.1 and re-incubated with HRP Alexa

Fluor 594 or HRP Alexa Fluor 647 for 5 min. Preps were then washed for 1 min with HL3.1.

Finally, post-digitonin Z-stacks were acquired of the same NMJs.

Image analysis

Debris, axons, and large fluorescent aggregates (sometimes observed for tdTomato:Sec61β)

were manually removed from all images before analyses of maximum projections or Z-stacks.

Neuronal masks were created by first conducting a background subtraction of the image (rolling

ball, 50 pixel radius), and then introducing a gaussian blur before thresholding using Otsu or Li

algorithms. For live image analysis, maximum intensity projections of each timepoint were

corrected for (1) photobleaching using histogram matching and (2) sample drift or muscle

contraction using the StackReg plugin (Thevenaz et al., 1998).

Presynaptic levels of ER membrane marker tdTomato:Sec61β *(**Figure 3**)*: *A* custom FIJI macro

consisting of the following steps was employed on maximum intensity projections,: (1) An NMJ

mask was created using the α-HRP neuronal marker; (2) The integrated density of the ER

membrane marker tdTomato:Sec61β was measured within the α-HRP mask; (3) The mean

intensity of the ER membrane marker at presynaptic terminals was calculated by dividing the

tdTomato:Sec61β integrated density by the area of the presynaptic terminal.

Analysis of NMJ morphology: Satellite and total bouton numbers were manually counted from

blinded maximum intensity projections of fixed and α-HRP labeled larvae. Satellite boutons were

defined as any string of 5 or fewer boutons extending from the main axis of the NMJ.

Categorization of ER dynamics: For dynamics analysis, we used the same dataset as the control

in **Figure 2**, **Figure 5**, and **Figure 5_Supplement**. Maximum intensity projection timelapse

movies were blinded and replayed 3-4 times before categorization. We identified three distinct

categories of ER dynamics in presynaptic terminals of *Drosophila* motor neurons (note that a

single bouton can exhibit multiple types of dynamics): (1) **<u>ER budding</u>** – small vesicle-like

structures emerging from the ER network and re-fusing in a different region; (2) **<u>ER tubule</u>**

<u>dynamics</u>, encompassing four subcategories as follows: (a) <u>Tubule displacement</u>, involving

tubules that changed their localization within boutons without detaching from the ER network;

(b) <u>Tubule extension</u>, where an ER tubule emerged and extended from the ER network; (c)

<u>Tubule retraction</u>, characterized by the retraction and disappearance of an ER tubule from the

ER network; and (d) <u>Tubule extension and retraction</u>, indicating an ER tubule both extending

and retracting from the ER network within the timelapse. (3) **<u>Foci dynamics</u>** – round, bright

puncta moving within the ER network and (4) **<u>static</u>** – boutons that lacked observable dynamics.

Presynaptic levels of luminal ER proteins at Drosophila *motor neurons:* A custom FIJI macro

consisting of the following steps was employed on Z-stacks: (1) We created an NMJ mask using

the luminal ER protein marker. (2) Using this mask, we measured the neuron volume and the

integrated density of the marker. (3) We calculated the mean intensity of the marker by dividing

the integrated density by the volume of the presynaptic terminal.

Categorization of luminal ER marker distribution: Images were blinded before analysis to ensure

unbiased qualitative analysis of luminal ER marker BiP:sfGFP:HDEL in controls and *Atlastin*

mutants. Maximum intensity projections were then sorted into three phenotypic categories: (1)

<u>"Partial loss"</u> - luminal ER marker localizes to ER networks but is partially displaced to the

cytosol, (2) **<u>"Complete loss"</u>** - luminal ER marker mostly displaced to the cytosol and ER

networks are not detectable, and (3) **"No phenotype"** - luminal ER marker completely localizes
to ER networks.
720

Presynaptic levels of cytosolic or luminal ER proteins pre- and post-digitonin treatment: Images
were blinded and maximum projections were created for further analysis. Neuronal outlines were
obtained by manually tracing the α-HRP-positive region in FIJI. Next, the integrated density of
UAS-GFP or UAS-BiP:sfGFP:HDEL was quantified within the manually traced neuronal outlines.
Mean intensity was calculated by dividing the integrated density by the area of the presynaptic
terminal.
727

BiP:sfGFP:HDEL levels before and after photobleaching: Images were blinded for analysis. The
20 frames corresponding to the photobleaching event were removed since this prevented the
Stackreg plugin from working correctly. Movies with uncorrectable drift were excluded. For the
analysis, we drew 3 ROIs corresponding to the region that was photobleached, a region away
from the bleach site and the background. A Fiji macro was then used to measure the mean
intensities at these 3 regions per movie. The mean intensities at the site of photobleaching of
BiP:sfGFP:HDEL, RFP:mCD8 or morphotrap$_{Int}$ were background subtracted and normalized to
the mean intensity of the non-bleached region. We also used the same approach to measure
the levels of BiP:sfGFP:HDEL before FRAP (**Figure 10C**).
737

Statistics
GraphPad Prism software was used to perform statistical analyses and generate graphs.
Statistical analyses were conducted using parametric tests (e.g. one-way ANOVA, Student's t-
test) for normally distributed data, and non-parametric tests (e.g. Kruskal-Wallis, Mann-Whitney
tests) if the data was not normally distributed. Chi-squared tests were used for categorical data.
p-values were set to *<0.05, **<0.01, ***<0.001.
744

Acknowledgements
We thank the Bloomington *Drosophila* Stock Center (Indiana University, Bloomington, IN, NIH
P40OD018537) and James McNew for fly lines, Steve Del Signore for help with image analysis,
and Bruce Goode for use of the SoRA microscope. We also thank Chi-Kuang Yao for sharing
reagents and stimulating discussion. This work was supported by NINDS grants R01 NS103967
to A.A.R., and T32 007292 and K99 NS136720-02 to M.C.Q.F.

Figure legends:

754 **Figure 1. *In vivo* imaging of ER networks in *Drosophila* motor neurons.** (**A**) The schematic setup for live imaging experiments in this study consists of a glass microscope slide with a mounted silicone isolator filled with Sylgard. Small metal pins were used to stretch the larvae (small round dots). Finally, a coverslip was placed on top of the larval fillet for imaging with an oil objective on upright or inverted microscopes. (**B**) A representative single slice of an Airyscan microscopy Z-stack of the center (top) or periphery (bottom) of the cell body of a motor neuron expressing the luminal ER marker BiP:sfGFP:HDEL. Pink dotted boxes indicate ER tubules and sheets. (**C**) Maximum projections of Z-stacks acquired with an Airyscan microscope of two different motor neuron axons expressing the luminal ER marker BiP:sfGFP:HDEL. (**D**) Representative maximum projection of a Z-stack acquired with an Airyscan microscope of a motor neuron presynaptic terminal expressing the luminal ER marker BiP:sfGFP:HDEL. The magenta outline highlights the neuronal boundary. Arrows indicate boutons containing ER baskets or ladder-like ER structures.

768 **Figure 2. Dynamics of luminal and membrane ER markers at presynaptic terminals.** (**A**) Diagram of a presynaptic terminal of *Drosophila* motor neurons highlighting *en passant* and terminal boutons with the ER in grey. (**B**, **top**) Maximum intensity projection of Airyscan Z-stacks from the beginning (t=0) and the end (t=39 sec) of a timelapse movie (**Movie S1**) of a single wild-type bouton expressing the ER membrane marker tdTomato:Sec61β. (**B, bottom**) Magnification of the region in the dotted red box of dynamics observed from 0 sec to 1.4 sec of the movie. (**C, top**) Maximum intensity projection of Airyscan Z-stacks from the beginning (t=0) and the end (t=72 sec) of a timelapse movie (**Movie S4**) of a single wild-type bouton expressing the luminal ER marker BiP:sfGFP:HDEL. (**C, bottom**) Magnification of the region in the dotted red box of dynamics observed from 0 sec to 1.4 sec of the movie. (**D**) Presynaptic terminals exhibiting ER budding, focus dynamics and various types of ER tubule dynamics (ER tubule displacement, tubule extension, and tubule retraction). (**E**) Qualitative categorization of the percentage of *en passant* and terminal boutons that are dynamic or static expressing the ER membrane marker tdTomato: Sec61β. (**F**) Qualitative categorization of the percentage of dynamic terminal boutons expressing the ER membrane marker tdTomato:Sec61β that exhibit ER budding, foci dynamics or ER tubule dynamics. (**G**) Qualitative categorization of the

784 percentage of dynamic terminal boutons expressing the ER membrane marker
785 tdTomato:Sec61β that exhibit tubule displacement, tubule extension, tubule retraction and both
786 tubule extension and retraction. The same control dataset used in E-G was used in **Figure 5**,
787 and **Figure 5_Supplement**.
788

Figure 3. *Atlastin* mutants have lower levels of the ER membrane tdTomato: Sec61β at
presynaptic terminals. (**A**) Representative maximum intensity projection of a spinning disk
confocal Z-stack from fixed larvae expressing the ER membrane marker tdTomato:Sec61β
(black) in control (**left**) and *Atlastin* mutants (**right**). The neuronal outline (derived from the α-
HRP mask) is shown in magenta. (**B**) Quantification of satellite boutons normalized to total
bouton number in control and *Atlastin* mutants. (**C**) Quantification of the intensity of the ER
membrane marker tdTomato:Sec61β, normalized to wild-type control. (**D**) Coefficient of variation
(CoV) of the ER membrane marker tdTomato:Sec61β. (**B-D**) Graphs show mean +/- s.e.m. from
30 control and 30 *Atlastin* mutant NMJs.
798

Figure 3 – Figure Supplement. *Atlastin* mutants expressing the ER membrane
tdTomato:Sec61β exhibit robust ER networks at presynaptic terminals. (**A, B**)
Representative maximum projection of a Z-stack acquired live by Airyscan microscopy of control
and *Atlastin* mutant NMJs expressing the ER membrane marker tdTomato: Sec61β at
presynaptic terminals. These images are not contrast matched because *Atlastin* mutants exhibit
lower levels of the tdTomato:Sec61β (See **Figures 3C**).
805

Figure 4. *Atlastin* mutants expressing the ER membrane tdTomato:Sec61β in axon
bundles and soma clusters have distinct phenotypes. (**A**) Representative middle slice of
axon bundles in control (top) and *Atlastin* mutants (bottom) expressing tdTomato:Sec61β. (**B**)
Representative top or middle slices of somas in control (top) and *Atlastin* mutants (bottom)
expressing tdTomato:Sec61β. (**C,F**) Quantification of the intensity of the ER membrane marker
tdTomato:Sec61β. (**D,G**) Quantification of the area of the ER membrane marker
tdTomato:Sec61β. (**E,H**) Coefficient of variation (CoV) of the ER membrane marker
tdTomato:Sec61β. For axon bundles, graphs show mean +/- s.e.m. from 29 control and 32
Atlastin mutant NMJs. For soma clusters, graphs show mean +/- s.e.m. from 26 control and 33
Atlastin mutant NMJs.

Figure 5. *Atlastin* mutants have mild ER dynamics phenotypes. (**A**) Representative images
of terminal (left) or *en passant* (right) boutons in controls (top) and *Atlastin* mutants (bottom)
(**Movies S7-14**). (**B**) Percentage of dynamic versus static boutons in controls and *Atlastin*
mutants expressing the ER membrane marker tdTomato:Sec61β, analyzed separately for
terminal and *en passant* boutons. Terminal bouton analysis included 75 control (65 dynamic, 10
static) and 83 *Atlastin* mutant boutons (63 dynamic, # static). *En passant* bouton analysis
included 29 control (11 dynamic, 18 static) and 36 *Atlastin* mutant boutons (11 dynamic, 25
static). (**C**) Distribution of ER dynamic types (budding, foci dynamics, or ER tubule dynamics) in
dynamic terminal boutons expressing tdTomato:Sec61β. (**D**) Distribution of ER dynamics types
in dynamic *en passant* boutons expressing tdTomato:Sec61β. Dynamic boutons were further
analyzed in **Figure 5_Supplement**. Data from control animals were also used in **Figure 2** and
Figure 5_Supplement.

Figure 5 – Supplement. (**A**) A qualitative categorization of the percentage of dynamic terminal
boutons in controls and *Atlastin* mutants expressing the ER membrane marker
tdTomato:Sec61β that exhibit tubule displacement, tubule extension, tubule retraction, and both
tubule extension and retraction. (**B**) A qualitative categorization of the number of dynamic
categories per terminal bouton in control and *Atlastin* mutants. The same control and *Atlastin*
mutant dataset were used in **Figure 2**, and **Figure 5**.

Figure 6. Luminal ER proteins are displaced to the cytosol in *Atlastin* mutants. Maximum
projection of a Z-stack acquired live with an Airyscan microscope of control (**A**) and *Atlastin*
mutant (**B**) NMJs expressing the ER lumen marker BiP:sfGFP:HDEL. (**B**) *Atlastin* mutants exhibit
two distinct phenotypes at presynaptic terminals that were categorized as partial loss (**B, top**) or
complete loss (**B, bottom**). Control and *Atlastin* mutant images are contrast matched. (**C**) A
qualitative categorization of presynaptic terminals in control and *Atlastin* mutants exhibiting
"partial loss", "complete loss" or "no phenotype". (**D**) Histogram of normalized mean intensities
of the ER lumen marker BiP:sfGFP:HDEL. Displacement does not strongly correlate with steady-
state levels of BiP:sfGFP:HDEL. (**E**) Quantification of the intensity of the luminal ER marker
BiP:sfGFP:HDEL. (**F**) Coefficient of variation (CoV) of the luminal ER marker BiP:sfGFP:HDEL
in neurons of control and *Atlastin* mutant. (**E, F**) Graphs show mean +/- s.e.m. from 8 control
and 15 *Atlastin* mutant NMJs.

Figure 6 – Supplement. Is and Ib motor neurons in various muscles and abdominal segments of 3rd instar *Atlastin* mutant larvae exhibit similar cytosolic displacement of BiP:sfGFP:HDEL. Maximum projection of a live Airyscan Z-stack from larvae expressing the ER lumen marker BiP:sfGFP:HDEL at muscles 12-13 (**A, left**), muscles 6-7 (**A, right**), muscle 4 in segment A2 (**B, left**) and muscle 4 in segment A4 (**B, right**).

Figure 7. Luminal ER proteins are not displaced in other neuronal compartments or cell types in *Drosophila* 3rd instar larvae. All images were acquired by live Airyscan microscopy from larvae expressing the ER lumen marker BiP:sfGFP:HDEL with Vglut-GAL4 (motor neurons) or BG57-GAL4 (body wall muscles). (**Left panel**) Maximum projection of a Z-stack of motor neuron axons. (**Middle panel**) A single slice of motor neuron cell bodies. (**Right panel**) A single slice of body wall muscles.

Figure 8. The luminal displacement phenotype in *Atlastin* mutants is progressive. (**A, B**) Maximum intensity projections of a live Airyscan Z-stack acquired of 1st instar larvae expressing the ER lumen marker BiP:sfGFP:HDEL (black) with a neuronal outline (magenta, derived from the α-HRP mask) in control (A) and *Atlastin* mutants (B). (**C**) A qualitative categorization of presynaptic terminals in control and *Atlastin* mutant 1st instar larvae expressing the ER lumen marker BiP:sfGFP:HDEL that exhibit "partial loss", "complete loss" or "no phenotype". Data are shown from 16 control NMJs and 16 *Atlastin* mutant NMJs.

Figure 9. FRAP analysis of BiP:sfGFP:HDEL recovery in *Atlastin* mutants. (**A**) Diagram of the recovery of BiP:sfGFP:HDEL in controls and *Atlastin* mutants with partial loss or complete loss. (**B**) Representative slices from movies of controls and *Atlastin* mutants with partial loss or complete loss co-expressing mCD8:RFP and BiP:sfGFP:HDEL pre-bleach, after FRAP and 2 min post-FRAP (**Movies S15-19**). (**C**) Mean recovery of mCD8:RFP in control (black squares) and *Atlastin* mutants with partial loss (empty lime green triangles) and complete loss (green triangles). (**D**) Mean recovery of BiP:sfGFP:HDEL in control (black squares) and *Atlastin* mutants with partial loss (empty lime green triangles) and complete loss (green triangles). The table contains the results from the non-linear fit one-phase association of the data for each dataset. The s.e.m. are represented by solid lines and shading: controls (n=10) have black lines with grey

shading, *Atlastin* mutants with partial loss (n=6) have lime green lines and lime green shading,

and *Atlastin* mutants with complete loss (n= 8) have green lines with light green shading.

Figure 10 – Supplement. Luminal ER proteins displaced to the cytosol are sensitive to

digitonin treatment. (**A**) Maximum projections of Z-stacks acquired live with a spinning disk

confocal of boutons expressing with Vglut-Gal4 GFP (control 1), BiP:sfGFP:HDEL (control 2),

BiP:sfGFP:HDEL in *Atlastin* mutants exhibiting partial loss and BiP:sfGFP:HDEL in *Atlastin*

mutants exhibiting complete loss. (**B**) Diagram of the expected distribution patterns before and

after digitonin treatment of GFP and BiP:sfGFP:HDEL. Note that digitonin treatment causes

partial ER fragmentation. (**C**) Fractional change of fluorescence intensity at presynaptic

terminals expressing GFP (cytosolic protein) or BiP:sfGFP:HDEL in control and *Atlastin* mutants

following digitonin treatment. Graphs show mean +/- s.e.m. from 18 control and 20 *Atlastin*

mutant NMJs. Statistical significance was only calculated between BiP:sfGFP:HDEL samples

since the baseline fluorescence and distribution of this marker was sensitive to digitonin

treatment in a manner distinct from cytosolic GFP.

Figure 10. BiP:sfGFP:HDEL in the cytosol binds morphotrap$_{Int}$ in *Atlastin* mutants. (**A**)

Diagram of the recovery of BiP:sfGFP:HDEL in controls and *Atlastin* mutants co-expressing

morphotrap$_{Int}$. (**B**) Distribution of BiP:sfGFP:HDEL phenotypes (partial loss, complete loss, or no

phenotype) in control and *Atlastin* mutant presynaptic terminals co-expressing either mCD8:RFP

or morphotrap$_{Int}$. (**C**) Intensity of BiP:sfGFP:HDEL in control and *Atlastin* mutant terminals co-

expressing either mCD8:RFP or morphotrap$_{Int}$. (**D**) Representative maximum projections of

controls (**Movies S20-22**) and *Atlastin* mutants (**Movies S23-24**) co-expressing morphotrap$_{Int}$

and BiP:sfGFP:HDEL pre-bleach, after FRAP and 5 min post-FRAP. (**E**) Mean recovery of

mCD8:RFP in control (black squares) and *Atlastin* mutants (green triangles). (**F**) Mean recovery

of BiP:sfGFP:HDEL in control (black squares) and *Atlastin* mutants (green triangles). The tables

contain the results from the non-linear fit one-phase association equation of all the datasets. The

s.e.m. are represented by solid lines and shading: controls have black lines with grey shading,

Atlastin mutants have green lines with light green shading. For 10A-B, there were 27 controls

co-expressing BiP:sfGFP:HDEL and mCD8:RFP, 28 controls co-expressing BiP:sfGFP:HDEL

and morphotrap$_{Int}$, 24 *Atlastin* mutants co-expressing BiP:sfGFP:HDEL and mCD8:RFP, and 24

Atlastin mutants co-expressing BiP:sfGFP:HDEL and morphotrap$_{Int}$. For 10 E-F, there were 16

controls co-expressing BiP:sfGFP:HDEL and morphotrap$_{Int}$ and 19 *Atlastin* mutants co-
expressing BiP:sfGFP:HDEL and morphotrap$_{Int}$.

Movies S1-3. Movies acquired at a frame rate of 0.65 frames per second of presynaptic
terminals expressing the ER membrane marker tdTomato:Sec61β. These movies are played at
a frame rate of 10 frames per second and the scale bar is 1 µm.

Movies S4-6: Movies acquired at a frame rate of 0.73 frames per second of presynaptic
terminals expressing the ER membrane marker tdTomato:Sec61β. These movies are played at
a frame rate of 10 frames per second and the scale bar is 1 µm.

Movie S7: Example movie of a static proximal bouton of controls expressing the ER membrane
marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie is played
at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S8: Example movie of a dynamic proximal bouton of controls expressing the ER
membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie
is played at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S9: Example movie of a static terminal bouton of controls expressing the ER membrane
marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie is played
at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S10: Example movie of a dynamic terminal bouton of controls expressing the ER
membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie
is played at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S11: Example movie of a static proximal bouton of *Atlastin* mutants expressing the ER
membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie
is played at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S12: Example movie of a dynamic proximal bouton of *Atlastin* mutants expressing the
ER membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The
movie is played at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S13: Example movie of a static terminal bouton of *Atlastin* mutants expressing the ER
membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie
is played at a frame rate of 5 frames per second and the scale bar is 2 µm.

Movie S14: Example movie of a dynamic terminal bouton of *Atlastin* mutants expressing the ER membrane marker tdTomato:Sec61β acquired at a frame rate of 0.92 seconds/stack. The movie is played at a frame rate of 5 frames per second and the scale bar is 2 μm.

Movie S15: Representative movie of a control NMJ expressing mCD8:RFP pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S16: Representative movie of a control NMJ expressing BiP:sfGFP:HDEL pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S17: Representative movie of an *Atlastin* mutant NMJ with partial loss expressing mCD8:RFP pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S18: Representative movie of an *Atlastin* mutant NMJ with partial loss expressing BiP:sfGFP:HDEL pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S19: Representative movie of an *Atlastin* mutant NMJ with complete loss expressing mCD8:RFP pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S20: Representative movie of an *Atlastin* mutant NMJ with complete loss expressing BiP:sfGFP:HDEL pre-bleach and after FRAP every 10 secs for 2 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S21: Representative movie of a control NMJ expressing morphotrap$_{int}$ pre-bleach and after FRAP every 10 secs for 5 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S22: Representative movie of a control NMJ expressing BiP:sfGFP:HDEL pre-bleach and after FRAP every 10 secs for 5 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S23: Representative movie of a *Atlastin* mutant NMJ expressing morphotrap$_{int}$ pre-bleach and after FRAP every 10 secs for 5 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

Movie S24: Representative movie of a *Atlastin* mutant NMJ expressing BiP:sfGFP:HDEL pre-bleach and after FRAP every 10 secs for 5 min. The movie is played at a frame rate of 5 frames per second, and the scale bar is 5 μm.

References:

996
997

Aponte-Santiago NA, Ormerod KG, Akbergenova Y, Littleton JT. 2020. Synaptic Plasticity
Induced by Differential Manipulation of Tonic and Phasic Motoneurons in *Drosophila*. *J*
Neurosci **40**:6270–6288. doi:10.1523/JNEUROSCI.0925-20.2020
Audhya A, Desai A, Oegema K. 2007. A Role for Rab5 in Structuring the Endoplasmic
Reticulum. *The Journal of Cell Biology* **178**:43–56.
Benedetti L, Fan R, Weigel AV, Moore AS, Houlihan PR, Kittisopikul M, Park G, Petruncio A,
Hubbard PM, Pang S, Xu CS, Hess HF, Saalfeld S, Rangaraju V, Clapham DE, Camilli
PD, Ryan TA, Lippincott-Schwartz J. 2024. Periodic ER-plasma membrane junctions
support long-range Ca2+ signal integration in dendrites.
doi:10.1101/2024.05.27.596121
Blackstone C. 2018. Hereditary spastic paraplegia. *Handbook of Clinical Neurology,* **Vol.**
148:633–651. doi:10.1016/B978-0-444-64076-5.00041-7
Blackstone C. 2012. Cellular Pathways of Hereditary Spastic Paraplegia. *Annu Rev Neurosci*
35:25–47. doi:10.1146/annurev-neuro-062111-150400
Blackstone C, O'Kane CJ, Reid E. 2011. Hereditary spastic paraplegias: membrane traffic and
the motor pathway. *Nat Rev Neurosci* **12**:31–42. doi:10.1038/nrn2946
Boutry M, Morais S, Stevanin G. 2019. Update on the Genetics of Spastic Paraplegias. *Curr*
Neurol Neurosci Rep **19**:18. doi:10.1007/s11910-019-0930-2
Burgoyne T, Patel S, Eden ER. 2015. Calcium signaling at ER membrane contact sites.
Biochimica et Biophysica Acta (BBA) - Molecular Cell Research, 13th European
Symposium on Calcium **1853**:2012–2017. doi:10.1016/j.bbamcr.2015.01.022
Carter SD, Hampton CM, Langlois R, Melero R, Farino ZJ, Calderon MJ, Li W, Wallace CT,
Tran NH, Grassucci RA, Siegmund SE, Pemberton J, Morgenstern TJ, Eisenman L,
Aguilar JI, Greenberg NL, Levy ES, Yi E, Mitchell WG, Rice WJ, Wigge C, Pilli J,
George EW, Aslanoglou D, Courel M, Freyberg RJ, Javitch JA, Wills ZP, Area-Gomez
E, Shiva S, Bartolini F, Volchuk A, Murray SA, Aridor M, Fish KN, Walter P, Balla T,
Fass D, Wolf SG, Watkins SC, Carazo JM, Jensen GJ, Frank J, Freyberg Z. 2020.
Ribosome-associated vesicles: A dynamic subcompartment of the endoplasmic
reticulum in secretory cells. *Sci Adv* **6**:eaay9572. doi:10.1126/sciadv.aay9572
Chino H, Mizushima N. 2020. ER-Phagy: Quality Control and Turnover of Endoplasmic
Reticulum. *Trends in Cell Biology* **30**:384–398. doi:10.1016/j.tcb.2020.02.001
Chouhan AK, Zhang J, Zinsmaier KE, Macleod GT. 2010. Presynaptic Mitochondria in
Functionally Different Motor Neurons Exhibit Similar Affinities for Ca2+ But Exert Little
Influence as Ca2+ Buffers at Nerve Firing Rates In Situ. *J Neurosci* **30**:1869–1881.
doi:10.1523/JNEUROSCI.4701-09.2010
Christianson JC, Jarosch E, Sommer T. 2023. Mechanisms of substrate processing during ER-
associated protein degradation. *Nat Rev Mol Cell Biol* 1–20. doi:10.1038/s41580-023-
00633-8
Costantini LM, Fossati M, Francolini M, Snapp EL. 2012. Assessing the Tendency of
Fluorescent Proteins to Oligomerize under Physiologic Conditions. *Traffic* **13**:643–649.
doi:10.1111/j.1600-0854.2012.01336.x
Crapart C, Scott ZC, Konno T, Sharma A, Bailey DM, Avezov E, Koslover EF. 2023. Luminal
transport rates through intact endoplasmic reticulum limit the magnitude of localized
Ca2+ signals. doi:10.1101/2023.06.23.546357

Dayel MJ, Hom EFY, Verkman AS. 1999. Diffusion of Green Fluorescent Protein in the
Aqueous-Phase Lumen of Endoplasmic Reticulum. *Biophysical Journal* **76**:2843–2851.
doi:10.1016/S0006-3495(99)77438-2
de Juan-Sanz J, Holt GT, Schreiter ER, de Juan F, Kim DS, Ryan TA. 2017. Axonal
Endoplasmic Reticulum Ca2+ Content Controls Release Probability in CNS Nerve
Terminals. *Neuron* **93**:867-881.e6. doi:10.1016/j.neuron.2017.01.010
Del Signore SJ, Mitzner MG, Silveira AM, Fai TG, Rodal AA. 2022. An approach for
quantitative mapping of synaptic periactive zone architecture and organization. *Mol Biol*
Cell mbcE22080372. doi:10.1091/mbc.E22-08-0372
Deng C, Moradi M, Reinhard S, Ji C, Jablonka S, Hennlein L, Lüningschrör P, Doose S, Sauer
M, Sendtner M. 2021. Dynamic remodeling of ribosomes and endoplasmic reticulum in
axon terminals of motoneurons. *Journal of Cell Science* **134**:jcs258785.
doi:10.1242/jcs.258785
Deshpande M, Feiger Z, Shilton AK, Luo CC, Silverman E, Rodal AA. 2016. Role of BMP
receptor traffic in synaptic growth defects in an ALS model. *MBoC* **27**:2898–2910.
doi:10.1091/mbc.E16-07-0519
Espadas J, Pendin D, Bocanegra R, Escalada A, Misticoni G, Trevisan T, Velasco del Olmo A,
Montagna A, Bova S, Ibarra B, Kuzmin PI, Bashkirov PV, Shnyrova AV, Frolov VA,
Daga A. 2019. Dynamic constriction and fission of endoplasmic reticulum membranes
by reticulon. *Nat Commun* **10**:5327. doi:10.1038/s41467-019-13327-7
Feng Y, Ueda A, Wu C-F. 2004. A modified minimal hemolymph-like solution, HL3.1, for
physiological recordings at the neuromuscular junctions of normal and mutant
Drosophila larvae. *J Neurogenet* **18**:377–402. doi:10.1080/01677060490894522
Fernandes AR, Martins JP, Gomes ER, Mendes CS, Teodoro RO. 2023. Drosophila motor
neuron boutons remodel through membrane blebbing coupled with muscle contraction.
Nat Commun **14**:3352. doi:10.1038/s41467-023-38421-9
Finsterer J, Löscher W, Quasthoff S, Wanschitz J, Auer-Grumbach M, Stevanin G. 2012.
Hereditary spastic paraplegias with autosomal dominant, recessive, X-linked, or
maternal trait of inheritance. *Journal of the Neurological Sciences* **318**:1–18.
doi:10.1016/j.jns.2012.03.025
Fowler PC, Garcia-Pardo ME, Simpson JC, O'Sullivan NC. 2019. NeurodegenERation: The
Central Role for ER Contacts in Neuronal Function and Axonopathy, Lessons From
Hereditary Spastic Paraplegias and Related Diseases. *Front Neurosci* **13**.
doi:10.3389/fnins.2019.01051
Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. 2011. ER tubules
mark sites of mitochondrial division. *Science* **334**:358–362.
doi:10.1126/science.1207385
Glock C, Biever A, Tushev G, Nassim-Assir B, Kao A, Bartnik I, tom Dieck S, Schuman EM.
2021. The translatome of neuronal cell bodies, dendrites, and axons. *Proceedings of*
the National Academy of Sciences **118**:e2113929118. doi:10.1073/pnas.2113929118
Graf ER, Heerssen HM, Wright CM, Davis GW, DiAntonio A. 2011. Stathmin is Required for
Stability of the Drosophila Neuromuscular Junction. *J Neurosci* **31**:15026–15034.
doi:10.1523/JNEUROSCI.2024-11.2011
Gregorio CD, Delgado R, Ibacache A, Sierralta J, Couve A. 2017. Drosophila Atlastin in motor
neurons is required for locomotion and presynaptic function. *J Cell Sci* **130**:3507–3516.
doi:10.1242/jcs.201657
Harmansa S, Alborelli I, Bieli D, Caussinus E, Affolter M. 2017. A nanobody-based toolset to
investigate the role of protein localization and dispersal in Drosophila. *eLife* **6**:e22549.
doi:10.7554/eLife.22549

Holcman D, Parutto P, Chambers JE, Fantham M, Young LJ, Marciniak SJ, Kaminski CF, Ron
D, Avezov E. 2018. Single particle trajectories reveal active endoplasmic reticulum
luminal flow. *Nat Cell Biol* **20**:1118–1125. doi:10.1038/s41556-018-0192-2
Hoyer MJ, Chitwood PJ, Ebmeier CC, Striepen JF, Qi RZ, Old WM, Voeltz GK. 2018. A Novel
Class of ER Membrane Proteins Regulates ER-Associated Endosome Fission. *Cell*
175:254-265.e14. doi:10.1016/j.cell.2018.08.030
Hu J, Shibata Y, Zhu P-P, Voss C, Rismanchi N, Prinz WA, Rapoport TA, Blackstone C. 2009.
A Class of Dynamin-like GTPases Involved in the Generation of the Tubular ER
Network. *Cell* **138**:549–561. doi:10.1016/j.cell.2009.05.025
Hu X, Wu F, Sun S, Yu W, Hu J. 2015. Human atlastin GTPases mediate differentiated fusion
of endoplasmic reticulum membranes. *Protein Cell* **6**:307–311. doi:10.1007/s13238-
015-0139-3
Igbaria A, Merksamer PI, Trusina A, Tilahun F, Johnson JR, Brandman O, Krogan NJ,
Weissman JS, Papa FR. 2019. Chaperone-mediated reflux of secretory proteins to the
cytosol during endoplasmic reticulum stress. *Proc Natl Acad Sci U S A* **116**:11291–
11298. doi:10.1073/pnas.1904516116
King C, Sengupta P, Seo AY, Lippincott-Schwartz J. 2020. ER membranes exhibit phase
behavior at sites of organelle contact. *Proceedings of the National Academy of*
Sciences **117**:7225–7235. doi:10.1073/pnas.1910854117
Lai Y-S, Stefano G, Brandizzi F. 2014. ER stress signaling requires RHD3, a functionally
conserved ER-shaping GTPase. *Journal of Cell Science* **127**:3227–3232.
doi:10.1242/jcs.147447
Lajoie P, Snapp EL. 2020. Size-dependent secretory protein reflux into the cytosol in
association with acute endoplasmic reticulum stress. *Traffic* **21**:419–429.
doi:10.1111/tra.12729
Le Gall S, Neuhof A, Rapoport T. 2004. The Endoplasmic Reticulum Membrane Is Permeable
to Small Molecules. *MBoC* **15**:447–455. doi:10.1091/mbc.e03-05-0325
Lee M, Paik SK, Lee M-J, Kim Y-J, Kim S, Nahm M, Oh S-J, Kim H-M, Yim J, Lee CJ, Bae YC,
Lee S. 2009. Drosophila Atlastin regulates the stability of muscle microtubules and is
required for synapse development. *Developmental Biology* **330**:250–262.
doi:10.1016/j.ydbio.2009.03.019
Lippincott-Schwartz J, Snapp EL, Phair RD. 2018. The Development and Enhancement of
FRAP as a Key Tool for Investigating Protein Dynamics. *Biophys J* **115**:1146–1155.
doi:10.1016/j.bpj.2018.08.007
Liu X, Guo X, Niu L, Li X, Sun F, Hu J, Wang X, Shen K. 2019. Atlastin-1 regulates
morphology and function of endoplasmic reticulum in dendrites. *Nat Commun* **10**:1–15.
doi:10.1038/s41467-019-08478-6
Lu L, Ladinsky MS, Kirchhausen T. 2009. Cisternal Organization of the Endoplasmic Reticulum
during Mitosis. *MBoC* **20**:3471–3480. doi:10.1091/mbc.e09-04-0327
Lü L, Niu L, Hu J. 2020. "At last in" the physiological roles of the tubular ER network. *Biophys*
Rep **6**:105–114. doi:10.1007/s41048-020-00113-y
Lu M, Tartwijk FW van, Lin JQ, Nijenhuis W, Parutto P, Fantham M, Christensen CN, Avezov
E, Holt CE, Tunnacliffe A, Holcman D, Kapitein L, Schierle GSK, Kaminski CF. 2020.
The structure and global distribution of the endoplasmic reticulum network are actively
regulated by lysosomes. *Science Advances* **6**:eabc7209. doi:10.1126/sciadv.abc7209
McKibbin C, Mares A, Piacenti M, Williams H, Roboti P, Puumalainen M, Callan AC, Lesiak-
Mieczkowska K, Linder S, Harant H, High S, Flitsch SL, Whitehead RC, Swanton E.
2012. Inhibition of protein translocation at the endoplasmic reticulum promotes

activation of the unfolded protein response. *Biochemical Journal* **442**:639–648. doi:10.1042/BJ20111220

Merighi A, Lossi L. 2022. Endoplasmic Reticulum Stress Signaling and Neuronal Cell Death. *International Journal of Molecular Sciences* **23**:15186. doi:10.3390/ijms232315186

Niu L, Ma T, Yang F, Yan B, Tang X, Yin H, Wu Q, Huang Y, Yao Z-P, Wang J, Guo Y, Hu J. 2019. Atlastin-mediated membrane tethering is critical for cargo mobility and exit from the endoplasmic reticulum. *PNAS* **116**:14029–14038. doi:10.1073/pnas.1908409116

Obara CJ, Nixon-Abell J, Moore AS, Riccio F, Hoffman DP, Shtengel G, Xu CS, Schaefer K, Pasolli HA, Masson J-B, Hess HF, Calderon CP, Blackstone C, Lippincott-Schwartz J. 2024. Motion of VAPB molecules reveals ER-mitochondria contact site subdomains. *Nature* **626**:169–176. doi:10.1038/s41586-023-06956-y

Oliva MK, Pérez-Moreno JJ, O'Shaughnessy J, Wardill TJ, O'Kane CJ. 2020. Endoplasmic Reticulum Lumenal Indicators in Drosophila Reveal Effects of HSP-Related Mutations on Endoplasmic Reticulum Calcium Dynamics. *Front Neurosci* **14**. doi:10.3389/fnins.2020.00816

Orso G, Pendin D, Liu S, Tosetto J, Moss TJ, Faust JE, Micaroni M, Egorova A, Martinuzzi A, McNew JA, Daga A. 2009. Homotypic fusion of ER membranes requires the dynamin-like GTPase Atlastin. *Nature* **460**:978–983. doi:10.1038/nature08280

O'Sullivan NC, Jahn TR, Reid E, O'Kane CJ. 2012. Reticulon-like-1, the Drosophila orthologue of the Hereditary Spastic Paraplegia gene reticulon 2, is required for organization of endoplasmic reticulum and of distal motor axons. *Hum Mol Genet* **21**:3356–3365. doi:10.1093/hmg/dds167

Özkan N, Koppers M, van Soest I, van Harten A, Jurriens D, Liv N, Klumperman J, Kapitein LC, Hoogenraad CC, Farías GG. 2021. ER - lysosome contacts at a pre-axonal region regulate axonal lysosome availability. *Nat Commun* **12**:4493. doi:10.1038/s41467-021-24713-5

Öztürk Z, O'Kane CJ, Pérez-Moreno JJ. 2020. Axonal Endoplasmic Reticulum Dynamics and Its Roles in Neurodegeneration. *Frontiers in Neuroscience* **14**:1–33. doi:10.3389/fnins.2020.00048

Perez-Alvarez A, Yin S, Schulze C, Hammer JA, Wagner W, Oertner TG. 2020. Endoplasmic reticulum visits highly active spines and prevents runaway potentiation of synapses. *Nature Communications* **11**:5083. doi:10.1038/s41467-020-18889-5

Pérez-Moreno JJ, Smith RC, Oliva MK, Gallo F, Ojha S, Müller KH, O'Kane CJ. 2023. Drosophila SPG12 ortholog, reticulon-like 1, governs presynaptic ER organization and Ca2+ dynamics. *J Cell Biol* **222**:e202112101. doi:10.1083/jcb.202112101

Raffaello A, Mammucari C, Gherardi G, Rizzuto R. 2016. Calcium at the Center of Cell Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes. *Trends in Biochemical Sciences* **41**:1035–1049. doi:10.1016/j.tibs.2016.09.001

Raiborg C, Wenzel EM, Pedersen NM, Olsvik H, Schink KO, Schultz SW, Vietri M, Nisi V, Bucci C, Brech A, Johansen T, Stenmark H. 2015. Repeated ER–endosome contacts promote endosome translocation and neurite outgrowth. *Nature* **520**:234–238. doi:10.1038/nature14359

Rao K, Stone MC, Weiner AT, Gheres KW, Zhou C, Deitcher DL, Levitan ES, Rolls MM. 2016. Spastin, atlastin, and ER relocalization are involved in axon but not dendrite regeneration. *MBoC* **27**:3245–3256. doi:10.1091/mbc.E16-05-0287

Read A, Schröder M. 2021. The Unfolded Protein Response: An Overview. *Biology (Basel)* **10**:384. doi:10.3390/biology10050384

Rismanchi N, Soderblom C, Stadler J, Zhu P-P, Blackstone C. 2008. Atlastin GTPases are required for Golgi apparatus and ER morphogenesis. *Hum Mol Genet* **17**:1591–1604. doi:10.1093/hmg/ddn046

Roos J, Hummel T, Ng N, Klämbt C, Davis GW. 2000. *Drosophila* Futsch Regulates Synaptic Microtubule Organization and Is Necessary for Synaptic Growth. *Neuron* **26**:371–382. doi:10.1016/S0896-6273(00)81170-8

Rowland AA, Chitwood PJ, Phillips MJ, Voeltz GK. 2014. ER Contact Sites Define the Position and Timing of Endosome Fission. *Cell* **159**:1027–1041. doi:10.1016/j.cell.2014.10.023

Schwarz DS, Blower MD. 2016. The endoplasmic reticulum: structure, function and response to cellular signaling. *Cell Mol Life Sci* **73**:79–94. doi:10.1007/s00018-015-2052-6

Scott ZC, Brown AI, Mogre SS, Westrate LM, Koslover EF. 2021. Diffusive search and trajectories on tubular networks: a propagator approach. *Eur Phys J E* **44**:80. doi:10.1140/epje/s10189-021-00083-0

Shibata Y, Mazur EE, Pan B, Paulo JA, Gygi SP, Chavan S, Valerio LSA, Zhang J, Rapoport TA. 2024. The membrane curvature-inducing REEP1-4 proteins generate an ER-derived vesicular compartment. *Nat Commun* **15**:8655. doi:10.1038/s41467-024-52901-6

Sicari D, Centonze FG, Pineau R, Le Reste P-J, Negroni L, Chat S, Mohtar MA, Thomas D, Gillet R, Hupp T, Chevet E, Igbaria A. 2021. Reflux of Endoplasmic Reticulum proteins to the cytosol inactivates tumor suppressors. *EMBO Rep* **22**:e51412. doi:10.15252/embr.202051412

Snow PM, Patel NH, Harrelson AL, Goodman CS. 1987. Neural-specific carbohydrate moiety shared by many surface glycoproteins in Drosophila and grasshopper embryos. *J Neurosci* **7**:4137–4144. doi:10.1523/JNEUROSCI.07-12-04137.1987

Spits M, Neefjes J. 2021. Mobile late endosomes modulate peripheral endoplasmic reticulum network architecture. *EMBO reports* **22**:e50815. doi:10.15252/embr.202050815

Subramanian K, Meyer T. 1997. Calcium-Induced Restructuring of Nuclear Envelope and Endoplasmic Reticulum Calcium Stores. *Cell* **89**:963–971. doi:10.1016/S0092-8674(00)80281-0

Summerville JB, Faust JF, Fan E, Pendin D, Daga A, Formella J, Stern M, McNew JA. 2016. The effects of ER morphology on synaptic structure and function in Drosophila melanogaster. *J Cell Sci* **129**:1635–1648. doi:10.1242/jcs.184929

Sun L, Yu F, Ullah A, Hubrack S, Daalis A, Jung P, Machaca K. 2011. Endoplasmic Reticulum Remodeling Tunes IP3-Dependent Ca2+ Release Sensitivity. *PLOS ONE* **6**:e27928. doi:10.1371/journal.pone.0027928

Thevenaz P, Ruttimann UE, Unser M. 1998. A pyramid approach to subpixel registration based on intensity. *IEEE Trans on Image Process* **7**:27–41. doi:10.1109/83.650848

Wang S, Tukachinsky H, Romano FB, Rapoport TA. 2016. Cooperation of the ER-shaping proteins atlastin, lunapark, and reticulons to generate a tubular membrane network. *eLife* **5**:e18605. doi:10.7554/eLife.18605

Wang X, Olberding KE, White C, Li C. 2011. Bcl-2 proteins regulate ER membrane permeability to luminal proteins during ER stress-induced apoptosis. *Cell Death Differ* **18**:38–47. doi:10.1038/cdd.2010.68

Wu X, Rapoport TA. 2018. Mechanistic insights into ER-associated protein degradation. *Curr Opin Cell Biol* **53**:22–28. doi:10.1016/j.ceb.2018.04.004

Wu Y, Whiteus C, Xu CS, Hayworth KJ, Weinberg RJ, Hess HF, Camilli PD. 2017. Contacts between the endoplasmic reticulum and other membranes in neurons. *PNAS* **114**:E4859–E4867. doi:10.1073/pnas.1701078114

Yalçın B, Zhao L, Stofanko M, O'Sullivan NC, Kang ZH, Roost A, Thomas MR, Zaessinger S,
Blard O, Patto AL, Sohail A, Baena V, Terasaki M, O'Kane CJ. 2017. Modeling of
axonal endoplasmic reticulum network by spastic paraplegia proteins. *eLife* **6**:1–27.
doi:10.7554/eLife.23882
Zamponi E, Meehl JB, Voeltz GK. 2022. The ER ladder is a unique morphological feature of
developing mammalian axons. *Dev Cell* **57**:1369-1382.e6.
doi:10.1016/j.devcel.2022.05.002
Zhao G, Zhu P-P, Renvoisé B, Maldonado-Báez L, Park SH, Blackstone C. 2016. Mammalian
knock out cells reveal prominent roles for atlastin GTPases in ER network morphology.
Experimental Cell Research **349**:32–44. doi:10.1016/j.yexcr.2016.09.015
Zhu P-P, Hung H-F, Batchenkova N, Nixon-Abell J, Henderson J, Zheng P, Renvoisé B, Pang
S, Xu CS, Saalfeld S, Funke J, Xie Y, Svara F, Hess HF, Blackstone C. 2022.
Transverse endoplasmic reticulum expansion in hereditary spastic paraplegia
corticospinal axons. *Hum Mol Genet* **31**:2779–2795. doi:10.1093/hmg/ddac072
Zhu P-P, Soderblom C, Tao-Cheng J-H, Stadler J, Blackstone C. 2006. SPG3A protein
atlastin-1 is enriched in growth cones and promotes axon elongation during neuronal
development. *Hum Mol Genet* **15**:1343–1353. doi:10.1093/hmg/ddl054

1251

1252

1253

A

Side view

Coverslip

Silicone isolator

Microscope Glass Slide

Top view

Microscope Glass Slide | Sylgard

Silicone isolator

B Cell body

Center

ER tubule

5µm

Periphery

Sheets

5µm

BiP:sfGFP:HDEL (ER lumen marker)

C Example #1 **Axon**

5µm

Example #2

5µm

BiP:sfGFP:HDEL (ER lumen marker)

D **NMJ**

ER "basket"

Ladder-like ER

5µm

BiP:sfGFP:HDEL (ER lumen marker)
Neuronal outline

Figure 1

Figure 2

Figure 3

A Control

NMJ #1 NMJ #2 NMJ #3

5μm 5μm 5μm

ER membrane (tdTomato:Sec61β)

B Atlastin mutant

NMJ #1 NMJ #2 NMJ #3

5μm 5μm 5μm

ER membrane (tdTomato:Sec61β)

Figure 3- Supplement

Axon

Soma cluster

A Control

5µm

Atl mutant

5µm

tdTomato:Sec61β

B Control *Top* *Middle*

5µm 5µm

Atl mutant *Top* *Middle*

5µm 5µm

tdTomato:Sec61β

C Axon bundle mean intensity — *

D Axon bundle Area (µm²) — ***

E Axon bundle CoV — *

F Soma cluster mean intensity — ns

G Soma Cluster Area (µm²) — ***

H Soma cluster CoV — *

Figure 4

Figure 5

Figure 5-Supplement

Figure 6

A

M12-13   M6-7

Control

ls
lb

3μm   3μm

Atlastin mutant

ls

lb

3μm   3μm

BiP:sfGFP:HDEL

B

M4 (segment A2)   M4 (segment A4)

Control

3μm   3μm

Atlastin mutant

3μm   3μm

BiP:sfGFP:HDEL

Figure 6- Supplement

	Body wall muscle	Motor neurons	

Figure 7

A Control 3μm

B Atlastin mutant 3μm

(Contrast enhanced)

BiP:sfGFP:HDEL
neuronal outline

C

% of NMJs

☐ Control
☐ Atlastin mutant
p-value = 0.0007

100% (No phenotype, Control)
37.5% (No phenotype, Atlastin mutant)
0 (Partial loss, Control)
50% (Partial loss, Atlastin mutant)
0 (Complete loss, Control)
12.5% (Complete loss, Atlastin mutant)

No phenotype   Partial loss   Complete loss

Figure 8

A

Wild-type
FRAP
GFP
ER network
Intermediate
BiP:sfGFP:HDEL
recovery

Partial loss Atl mutant
GFP
Fragmented ER
Slow
Bip:sfGFP:HDEL
recovery

Complete loss Atl mutant
GFP
Fast
Bip:sfGFP:HDEL
recovery

B

	Pre-bleach	FRAP	2min post-FRAP

Control — mCD8:RFP — 5µm — BiP:sfGFP:HDEL — 5µm

Partial loss Atl mutant — mCD8:RFP — 5µm — BiP:sfGFP:HDEL — 5µm

Complete loss Atl mutant — mCD8:RFP — 5µm — BiP:sfGFP:HDEL — 5µm

C

% recovery of mCD8:RFP after FRAP vs Time (secs)

- Control (n =10)
- Complete loss Atl mutant (n = 8)
- Partial loss Atl mutant (n= 6)

D

% recovery of BiP after FRAP vs Time (secs)

- Complete loss Atl mutant (n = 8)
- Control (n =10)
- Partial loss Atl mutant (n= 6)

Best-fit values (one-phase association)			
Genotypes	Control	Partial loss Atl mutant	Complete loss Atl mutant
Plateau (%)	65.73	50.61	66.34
Half Life (1/s)	26.91	38.26	14.62

p-value= <0.001

Figure 9

Figure 10-Figure Supplement 1

Figure 10