High-resolution imaging of presynaptic ER networks in _Atlastin_ mutants

Mónica C. Quiñones-Frías[1], Dina M. Ocken[1] and Avital Rodal[1,*]

[1]Department of Biology, Brandeis University, Waltham, MA

*Corresponding author:

Email: arodal@brandeis.edu

Abstract:

The endoplasmic reticulum (ER) is a continuous organelle that extends to the periphery of

neurons and regulates many neuronal functions including neurite outgrowth, neurotransmission,

and synaptic plasticity. Mutations in proteins that control ER shape are linked to the

neurodegenerative disorder Hereditary Spastic Paraplegia (HSP). However, the ultrastructure

and dynamics of the neuronal ER have been under-investigated, particularly at presynaptic

terminals. Here we developed new super-resolution and live imaging methods in _D._

melanogaster larval motor neurons to investigate ER structure at presynaptic terminals from

wild-type animals, and in null mutants of the HSP gene Atlastin. Previous studies indicated

diffuse localization of an ER lumen marker at _Atlastin_ mutant presynaptic terminals, which was

attributed to ER fragmentation. By contrast, we found using an ER membrane marker that the

ER in _Atlastin_ mutants formed robust networks. Further, our high-resolution imaging results

suggest that overexpression of luminal ER proteins in _Atlastin_ mutants causes their progressive

displacement to the cytosol at synapses, perhaps due to proteostatic stress and/or changes in

ER membrane integrity. Remarkably, these luminal ER proteins remain correctly localized in cell

bodies, axons, and other cell types such as body wall muscles, suggesting that ER tubules at

synapses have unique structural and functional characteristics. This displacement phenotype

has not been reported in numerous studies of Atlastin in non-neuronal cells, emphasizing the

importance of conducting experiments in neurons when investigating the mechanisms leading

to upper motor neuron dysfunction in HSP.

Introduction:

The endoplasmic reticulum (ER) is a continuous organelle that extends to the periphery of

neurons. The ER consists of two highly dynamic structural domains known as ER sheets (which

are enriched in cell bodies) and ER tubules (which are abundant in dendrites and axons) (Öztürk

et al., 2020; Wu et al., 2017). The ER regulates many neuronal functions including neurite

outgrowth, neurotransmission, and synaptic plasticity (de Juan-Sanz et al., 2017; Liu et al., 2019;
Oliva et al., 2020; Perez-Alvarez et al., 2020; Raffaello et al., 2016; Raiborg et al., 2015).
Increasing evidence suggests that there are structural and functional differences in ER located
at cell bodies, axons and presynaptic terminals. For example, the density of ER-organelle
contacts varies across neuronal compartments suggesting that the functions of these contacts
might be differentially regulated (Wu et al., 2017). Indeed, a recent study found that ER-
endosome contacts in cell bodies exclusively regulate the size of lysosomes (Özkan et al., 2021).
In addition, developing axons exhibit a unique ER ladder structure, whose function in axonal
biology is still unknown (Zamponi et al., 2022). However, the cell biology and ultrastructure of
these different elements of neuronal ER, particularly at presynaptic nerve terminals, have been
under-investigated. This is because most studies of neuronal ER structure have either used
conventional microscopy, which falls short of resolving individual ER tubules in small structures
like synapses, or electron microscopy (EM), which is usually restricted to thin 2D sampling, is
difficult to interpret in synaptic vesicle-rich presynaptic terminals, and does not allow examination
of live dynamics (Chouhan et al., 2010; Oliva et al., 2020; Orso et al., 2009; Pérez-Moreno et
al., 2023; Summerville et al., 2016; Yalçın et al., 2017; Zhu et al., 2022). Exploring the function
of neuronal ER structure is important because mutations in ER-shaping genes cause
neurodegenerative disorders such as Hereditary Spastic Paraplegia (HSP) and Charcot Marie
Tooth Disease (Boutry et al., 2019; Fowler et al., 2019; Öztürk et al., 2020), implying that ER
structure is uniquely essential for neuronal physiology.

An important ER tubule-shaping protein is Atlastin, a GTPase that regulates the
homotypic fusion and tethering of ER tubules to form 3-way junctions (Lü et al., 2020). In
humans, Atlastin mutations underlie one of the most common forms of autosomal dominant
HSP, highlighting the importance of examining the neuron-specific functions of Atlastin in
regulating ER structure (Finsterer et al., 2012). In a variety of experimental systems, *Atlastin*
mutants exhibit numerous neuronal phenotypes, including defects in synaptic vesicle cycling
and trafficking of neuronal cargoes (Gregorio et al., 2017), synapse development (Gregorio et
al., 2017; Lee et al., 2009; Zhu et al., 2006) and axon regeneration (Rao et al., 2016). However,
significant gaps remain in our understanding of how Atlastin-dependent changes in neuronal ER
structure and dynamics lead to these functional defects. For example, *Drosophila Atlastin*
mutants exhibit shorter ER tubule profiles in their motor neuron cell bodies by electron
microscopy (Orso et al., 2009), and largely diffuse localization of an ER lumen marker in their

presynaptic terminals, which was attributed to extensive ER fragmentation. However, this
fragmentation has not been validated by EM, or by fluorescence using ER membrane markers.
Further, ER networks in neuronal dendrites of *C. elegans Atlastin* mutants were not fragmented
but were more prone to retraction, leading to disrupted microtubule structures and mitochondria
fission (Liu et al., 2019). Finally, a ladder-like ER expansion was observed in mammalian spinal
axons (Zhu et al., 2022). In contrast to these neuronal findings, Atlastin knock-down in Hela or
Cos-7 cells leads to long and unbranched ER tubules (Hu et al., 2009, 2015; Niu et al., 2019).
Thus, loss of Atlastin leads to different ER structural phenotypes in various regions of neurons,
when observed with specific reporters, and compared to non-neuronal cell types. To resolve the
functions of this vital protein, comprehensive imaging studies are needed of ER organization in
different regions of neurons, using complementary labeling strategies.

Another important gap exists in our understanding of neuronal ER dynamics. High-
resolution imaging in non-neuronal cells has revealed transitions between ER tubules to ER
sheets (Lu et al., 2009; Sun et al., 2011), ER fragmentation (Espadas et al., 2019), tubule
extension and retraction (Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014), and
alterations in 3-way junction number (Spits and Neefjes, 2021). These dynamics have been
linked to various ER functions such as endosome or mitochondria fission (Friedman et al., 2011;
Hoyer et al., 2018; Rowland et al., 2014) and calcium signaling (Crapart et al., 2023; Sun et al.,
2011). In neurons, ER rearrangements have been suggested to alter calcium
compartmentalization (Subramanian and Meyer, 1997), local protein synthesis (Deng et al.,
2021; Schwarz and Blower, 2016), organelle contacts (Lu et al., 2020) and transport within the
ER (Scott et al., 2021). However, the types and functions of neuronal ER dynamics, particularly
in presynaptic terminals, remain largely unknown due to the challenges in imaging ER networks
over time in small compartments.

New insights into the mechanisms that regulate ER structure and ER dynamics in neurons
are crucial for understanding why and how mutations in ER shaping proteins, such as Atlastin,
predominantly affect the nervous system. Here, we addressed this gap by examining the
structure and dynamics of the ER at the *Drosophila* neuromuscular junction (NMJ) using a novel
in vivo imaging setup and super-resolution Airyscan microscopy.

Results

High resolution of the ER in *Drosophila* motor neuron cell bodies, axons, and presynaptic
terminals

At presynaptic terminals of the *Drosophila* larval NMJ, the ER is highly sensitive to fixation
(Summerville et al., 2016) which has limited imaging experiments to live samples, often acquired
on upright microscopes using water dipping objectives (Chouhan et al., 2010; Oliva et al., 2020;
O'Sullivan et al., 2012; Pérez-Moreno et al., 2023; Summerville et al., 2016; Yalçın et al., 2017).
The axial resolution limit of typical water dipping objectives is >300 nm, which is insufficient to
distinguish the densely packed ER tubules at presynaptic terminals (Espadas et al., 2019; Wu
et al., 2017). To overcome this limitation, we developed a reusable imaging slide to acquire live
super-resolution images of neuronal ER in *Drosophila* larval fillets, using high numerical aperture
oil objectives and Airyscan microscopy (**Figure 1A**). Our system provides axial resolution of
<170 nm after processing (Del Signore et al., 2022). We imaged larval fillets using this device
on both inverted and upright microscopes within 40-50 min from dissection, a time frame in which
this preparation maintains muscle potential and the capacity for evoked neurotransmitter
release, even once the axon is severed from the cell body (Feng et al., 2004).

To study the live dynamics of the ER, we used Vglut-Gal4 to express either the luminal
ER marker BiP:sfGFP:HDEL (**Figure 1**, **2B**) or the transmembrane ER marker tdTomato:Sec61β
(**Figure 2A**) in *Drosophila* 3rd instar larval motor neurons. These markers were previously shown
to co-localize at this synapse, indicating that they are comparable reporters for visualizing
synaptic ER (Summerville et al., 2016). Our new imaging method allowed more precise
visualization of ER tubules and networks labeled by these reporters in cell bodies, axons, and
presynaptic terminals compared to previously reported results (**Figure 1B-D**). As previously
observed using lower-resolution imaging, we detected a wide variety of ER tubule structures in
all neuronal compartments examined (Summerville et al., 2016), including ER tubules and ER
sheets in cell bodies, and ER tubules in axons and presynaptic terminals. In axons, ER tubules
were long and we sometimes observed a ladder-like structure consisting of pairs of
interconnected ER tubules joined by rungs or cross-bridges, similar to structures previously
described in developing axons of mammalian cultured neurons (Zamponi et al., 2022) (**Figure
1C**). At presynaptic terminals, ER tubules often appeared in a highly interconnected a basket-
like network, while other boutons had several parallel ER tubules traversing the bouton, similar
to the ladder-like structures observed in axons (**Figure 1D**). Thus, we were able to detect

previously observed structures at approximately double the resolution previously obtained,

providing us with the opportunity to visualize their detailed structure and dynamics over time.

To examine the dynamics of the ER at presynaptic terminals, we collected Z stacks of

single boutons expressing the ER membrane marker for 40 sec at 0.65 sec/stack (**Movie S1-**

S3) or the luminal ER marker for 90 sec at 0.73 sec/stack (**Movie S4-S6**). Both markers were

distributed unevenly across the ER network (**Figure 2A-2B**). In particular, the luminal ER marker

exhibited numerous discrete round foci that moved within ER tubules with fast dynamics (<1

sec). Such foci were less frequently observed with the ER membrane marker (**Figure 2A-2B**,

bottom), consistent with the model that proteins in the ER lumen and ER membrane may

differentially cluster and move within the ER (Holcman et al., 2018; Obara et al., 2022).

Conversely, static regions of high intensity (which we saw for both markers) may reflect stacked

ER tubules (Costantini et al., 2012). We focused on the ER membrane marker to define other

dynamic events, since there were fewer confounding foci, and observed (1) ER budding, in which

a fragment of ER dissociates from the network (2) ER tubule dynamics, including overall

displacement of the tubule in the XY plane, as well as lengthening (extension) and shortening

(retraction) of the tubule (**Figure 2C-F**).

To quantify the global dynamics of the ER network in neurons expressing either the ER

membrane or ER lumen markers, we calculated the Pearson correlation coefficient (PCC)

between the 1st frame of our movies with each subsequent frame (**Figure 2G**). We observed

that the luminal ER marker becomes uncorrelated at a faster rate (τ = 37.08s) than the ER

membrane marker ($\tau = 61.82s$), suggesting more rapid changes in ER dynamics. We

hypothesize the differences in the dynamics of the luminal and membrane markers are due to

the highly mobile nature of luminal ER marker foci within the ER network (**Figure 2B**). However,

despite these differences, the correlation coefficients for both ER markers were still relatively

high throughout the imaging period: The first and last frames of the luminal ER marker had a

PCC of 0.74 and the membrane ER marker had a slightly higher PCC of 0.85. Thus, the overall

structure of the ER network changes only modestly during the time-course of our movies

suggesting it is relatively stable in these neurons, which are at rest (**Figure 2A**). We also

qualitatively categorized synaptic boutons as "dynamic" or "non-dynamic". We categorized a

bouton as dynamic if we observed any changes in the structure of the ER network such as ER

tubule movement and foci dynamics. We found that 86% of boutons were dynamic while 14%

were not dynamic (**Figure 2D**). Overall, these results show that we can capture live ER dynamics

at high resolution in *Drosophila* motor neurons.

<u>*Atlastin* mutant synapses have robust ER networks but reduced levels of ER membrane markers</u>

170   We first aimed to correlate ER structure with neuronal development and function to

interrogate the potential physiological functions of presynaptic ER structure and dynamics. To

this end, we examined a mutant of the ER-shaping protein, Atlastin. Mammals have three

Atlastin homologs and Atlastin-1 is enriched in the brain, while *Drosophila* has a single Atlastin

homolog (Rismanchi et al., 2008). As in previous studies of *Atlastin* mutant phenotypes in

Drosophila, we used transheterozygotes of the *Atlastin* null mutant atl^2 and a 3^{rd} chromosome

deficiency (def) removing the *Atlastin* locus, to eliminate confounding effects from second site

mutations (Lee et al., 2009). Most atl^2/atl def mutants (from now on termed *Atlastin* mutants)

were pharate lethal with few escapers, and we observed that 3^{rd} instar larval presynaptic

terminals had a significant increase in satellite boutons (**Figure 3 Supplement A**), indicating

defects in synaptic growth and development, as previously reported (Gregorio et al., 2017; Lee

et al., 2009).

183   First, we used Airyscan microscopy to examine ER network structures at *Atlastin* mutant

presynaptic terminals expressing the ER membrane marker tdTomato:Sec61β. A previous study

used confocal microscopy of the ER lumen marker BiP:sfGFP:HDEL to examine ER structure in

Drosophila Atlastin mutant presynaptic terminals, and observed a highly diffuse distribution of

the marker, which was attributed to extensive ER fragmentation (Summerville et al., 2016). We

reasoned we would observe a similar phenotype in *Atlastin* mutants expressing an ER

membrane marker. However, *Atlastin* mutants expressing the ER membrane marker did not

exhibit diffuse or fragmented signal, and had tubular ER networks that were qualitatively similar

to controls (**Figure 3A-B**). We observed no significant difference in the density of complex

basket-like structures or simple basket-like structures labeled with tdTomato:Sec61β, despite

the smaller size of satellite boutons in *Atlastin* mutants (**Figure 3C-D**). Our results suggest that

there are indeed ER networks in *Atlastin* mutant synapses (rather than extensive fragmentation)

and that structural defects underlying the previously observed loss of synaptic functions are likely

specific to a subset of ER structural or dynamic properties. Further, our findings show that

luminal and membrane ER markers have distinct phenotypes in *Atlastin* mutants, which we
investigate further below (see **Figures 4-7)**.

200   To further explore ER architecture in the *Atlastin* mutant we generated masks of ER from
boutons with complex basket-like structures (**Figure 3E**). We found a small but significant
increase in the fraction of bouton area occupied by ER in *Atlastin* mutants compared to controls
(**Figure 3F**), suggesting that their architectures are subtly different. We also found that the
marker tdTomato:Sec61β exhibited lower fluorescence levels in *Atlastin* mutant boutons, as
quantified by its mean intensity within the segmented ER area in live preparations (**Figure 3G**),
as well as over the entire presynaptic terminal in fixed preparations (**Figure 3 Supplement – B-**
D). This reduction in ER membrane marker intensity has also been observed in other HSP
mutants, suggesting this is a common feature of ER shaping mutants, and could indicate
changes in ER membrane composition or integrity (Pérez-Moreno et al., 2023). Finally, we noted
that the Coefficient of Variation (CoV) of tdTomato:Sec61β signal was significantly higher in
Atlastin mutants (**Figure 3 Supplement E**), suggesting that it is more concentrated in
subsynaptic structures compared to wild-type synapses (notably the opposite of what would be
expected for fragmentation). Overall, our data suggests ER networks exist in *Atlastin* mutant
synapses, distinct from the diffuse distribution seen with a luminal ER marker, and that they may
have quantitative differences in architecture compared to wild-type synapses.

<u>Presynaptic terminals of *Atlastin* mutants are sensitized to the overexpression of a luminal ER</u>
<u>marker</u>
219   Our ER membrane marker results are in sharp contrast to a previous study examining the
ER structure of *Atlastin* mutants with the luminal ER marker, BiP:sfGFP:HDEL. This study
concluded from the diffuse distribution of this marker at synapses that the ER was highly
fragmented in *Atlastin* mutants (Summerville et al., 2016). Since we did not observe
fragmentation with the ER membrane marker, we used Airyscan microscopy to re-examine the
luminal ER marker BiP:sfGFP:HDEL in *Atlastin* mutants. Unlike its strict localization to the ER
network in control synapses, we reproduced the observation that a large fraction of
BiP:sfGFP:HDEL was diffusely localized in *Atlastin* mutant synapses, even at the resolution of
Airyscan microscopy. Further, the marker extended to the boundary of presynaptic terminals,
suggesting that it may be mislocalized to the cytosol (**Figure 4A-B**). We identified two distinct
ER structure phenotypes in *Atlastin* mutants: NMJs with "Phenotype A" had diffuse

BiP:sfGFP:HDEL but also had an identifiable ER network structure, while NMJs with "Phenotype B" had a more severe phenotype with no visible ER network structures. We blinded our images and categorized control and *Atlastin* mutant datasets as "Phenotype A", "Phenotype B" or no phenotype (**Figure 4C**). In *Atlastin* mutants, we found that 45% of NMJs exhibited "Phenotype A", 55% of NMJs exhibited "Phenotype B" and 10% of the NMJs had no phenotype. We observed Phenotypes A and B in both anterior (segment A2) and posterior (segment A5) muscle segments, in Is (phasic) and Ib (tonic) motor neuron subtypes, and in neurons innervating multiple different body wall muscles, suggesting that it reflects a widespread phenomenon in motor neuron synapses of *Atlastin* mutants (**Figure 4 - Figure Supplement 1**). Interestingly, we observed a shift in the distribution of presynaptic terminals with Phenotype B in *Atlastin* mutants towards lower mean intensities of the marker (**Figure 4D**). This could suggest that the luminal ER marker is either degraded or removed from presynaptic terminals, leading to lower levels. We also observed a significant change in its coefficient of variance (CoV) due to redistribution of the luminal ER marker in *Atlastin* mutants (**Figure 4F**). Together with our finding that a membrane-associated ER marker does not show fragmentation but exhibits lower intensity, these results raise the possibility that the luminal ER marker is mislocalized to the cytosol and that there may be changes to the structural integrity of the ER membrane. We were unable to assess co-localization of BiP:sfGFP:HDEL and tdTomato:Sec61β, as very few larvae co-expressing these markers survived, and the few that did exhibit grossly defective NMJ morphology and large aggregates of both markers. Further, adult flies expressing BiP:sfGFP:HDEL were unhealthy and could not be maintained as a stock, suggesting that this marker is toxic even in otherwise wild-type flies. Taken together, these results suggest that *Atlastin* mutants are particularly sensitized to overexpression of the BiP:sfGFP:HDEL marker.

To ask if the luminal ER marker displacement phenotype is specific to neurons or presynaptic terminals, we examined the ER in body wall muscles of *Atlastin* mutants and did not observe the luminal ER marker displaced to the cytosol (though we noticed reduced network complexity, as previously reported (Lee et al., 2009) (**Figure 5, right**). Next, we examined the cell body and axons of *Atlastin* mutants to determine if the displacement of the luminal ER marker was a global or local phenotype within neurons. As previously reported (Summerville et al., 2016), we did not observe a diffuse signal for the luminal ER marker in cell bodies or axons (even within ~30μm of the synapse) (**Fig 5, middle**). However, we did observe ER fragmentation

in some axons (**Fig 5, left**). These results indicate that the sensitivity of *Atlastin* mutants to

BiP:sfGFP:HDEL reflects a synapse-specific function.

<u>The luminal ER phenotype in *Atlastin* mutants is progressive</u>

Atlastin disruption is linked to the neurodegenerative disorder HSP, which is a progressive

disease in which patients lose neuronal function over time (Blackstone, 2018, 2012; Blackstone

et al., 2011). Indeed, *Drosophila Atlastin* mutants recapitulate progressive motor dysfunction as

observed in HSP patients (Gregorio et al., 2017). To test the hypothesis that Phenotype A might

similarly progress to Phenotype B, we examined 1st instar larvae and categorized control and

Atlastin mutant datasets as described above. If the phenotype is progressive, we predicted that

we would observe a lower fraction of NMJs with diffuse luminal ER marker and an increase in

the fraction of synapses with no phenotype. Indeed, we observed a decrease to 12.5% of NMJs

with Phenotype B in 1st instar compared to 55% in 3rd instar, and an increase to 38% of NMJs

with no phenotype in 1st instar compared to 10% in 3rd instar (compare **Figure 4A-C** to **Figure**

6). These results suggest that displacement of the luminal ER marker progresses over time and

that this phenotype is initiated early in neuronal development.

<u>Luminal proteins are likely located in the cytosol in *Atlastin* mutants</u>

Our results show diffuse BiP:sfGFP:HDEL in *Atlastin* mutants (even at high resolution)

compared to robust ER networks marked by tdTomato:Sec61β. This suggests that the luminal

marker is not in small ER membrane fragments as previously concluded, and instead may be

displaced to the cytosol without being associated with ER membranes. To test this hypothesis,

we treated presynaptic terminals for 5 min with the mild detergent digitonin to gently permeabilize

the plasma membrane and extract proteins from the cytosol (**Figure 7A**). We hypothesized the

digitonin treatment would lead to loss of fluorescence in *Atlastin* mutants, proportional to the

fraction of the luminal ER marker protein in the cytosol (**Figure 7B**). We imaged the same

presynaptic terminal (expressing either cytosolic GFP or BiP:sfGFP:HDEL) both before and after

digitonin treatment. To identify presynaptic terminals, we also labeled our live samples with α-

HRP antibodies (Snow et al., 1987).

We measured the fraction change in mean intensity to determine the effects of digitonin

treatment on cytosolic and ER markers (**Figure 7C**). For cytosolic GFP, we observed a

significant loss in mean fluorescence in all our samples (13/13) following digitonin treatment.

Control presynaptic terminals expressing the luminal ER marker, BiP:sfGFP:HDEL, became

highly fragmented following digitonin treatment, perhaps due to changes in the osmolarity of the

cytosol, which has been previously shown to promote ER fragmentation (King et al., 2020).

However, most of this fragmented ER did not escape the synaptic terminal, indicating that ER

fragments were not extracted or severely permeabilized with this treatment. Interestingly, in half

the control samples (9/18) the marker redistributed to bright puncta and showed an unexpected

increase in mean intensity. Digitonin treatment may allow the exchange of small molecules into

the ER (Le Gall et al., 2004), perhaps changing the fluorescent properties of sfGFP. The other

half of samples (9/18) had slightly reduced mean intensity following digitonin treatment, possibly

due to moderate loss of ER fragments. Next, we categorized and analyzed *Atlastin* mutants with

Phenotype A or Phenotype B. In *Atlastin* mutants with Phenotype A, we observed no significant

difference from controls following digitonin treatment. We observed that half (5/10) of our

samples increased their mean intensity levels following digitonin treatment while the other half

had a decrease (5/10). We also observed that *Atlastin* mutant synapses exhibiting Phenotype A

had puncta following digitonin treatment similar to control (likely because Phenotype A terminals

have some ER network visible with the luminal marker). In contrast, we observe a decrease in

fluorescence intensity following digitonin treatment in most (8/10) *Atlastin* mutant presynaptic

terminals with Phenotype B (which have almost no visible ER networks with the luminal marker).

These data suggest that the population of luminal ER molecules sensitive to digitonin in *Atlastin*

mutants is likely displaced to the cytosol (though we cannot rule out very small ER fragments).

Discussion

We introduce in this study an innovative *in vivo* imaging system to study the architecture

and dynamic behaviors of *Drosophila* neuronal ER networks. Our imaging system is versatile

and can be used to visualize genetically encoded fluorescent proteins in diverse larval tissues.

For the first time, we examined and defined the dynamic properties of ER networks at

presynaptic terminals *in vivo*, providing a foundation for future studies to identify the function of

the various ER structural rearrangements observed at presynaptic terminals. Using our high-

resolution system, we found that *Atlastin* mutant presynaptic terminals are sensitive to

overexpression of a luminal ER marker, likely resulting in its displacement to the cytosol.
However, they form robust networks with only subtle defects when labeled by an ER membrane
marker. Our results show that Atlastin plays unique and synapse-specific roles in regulating the
structure and function of ER, which may underlie its numerous previously reported deficits in
synaptic function.
333
334   **ER network dynamics at presynaptic terminals**
335
We categorized ER dynamics at presynaptic terminals as: (1) Budding and fusion of
vesicular structures, (2) ER tubule dynamics, and (3) dynamics of foci that are retained within
the ER network. Future work is needed to directly examine the function of these dynamics at
presynaptic terminals, but we speculate below on their potential functions. Neurons are highly
polarized cells and membrane trafficking pathways can have specialized functions at presynaptic
terminals (Deshpande and Rodal, 2016). ER-derived vesicles could act as localized calcium
signaling hubs (Burgoyne et al., 2015) or serve as sites for compartmentalized local translation
if they carry ribosomes (Carter et al., 2020; Deng et al., 2021). Other pathways involving the
formation of ER vesicles include ER-to-Golgi transport and ER autophagy (ER-phagy) (Chino
and Mizushima, 2020). However, the synaptic ER vesicle-like structures that we observed are
unlikely to be involved in the secretory pathway or ER-phagy because these vesicles re-fuse
with the ER. Second, ER tubule dynamics play multiple roles, such as maintaining the tension
of the ER network (Espadas et al., 2019; Niu et al., 2019) and forming ER-organelle contacts
(Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014). ER contacts occur at the tips
and edges of ER tubules, and they have diverse functions, such as promoting mitochondria and
endosome fission (Friedman et al., 2011; Hoyer et al., 2018; Rowland et al., 2014). Future
studies should perform co-localization experiments with these other organelles or generate ER-
organelle sensors to determine if ER tubule dynamics are important for these functions at
presynaptic terminals. The last type of ER dynamics we identified is focus dynamics within ER
networks. We hypothesize that foci emerge due to changes in the dynamics of protein clusters
at the ER membrane or within the lumen. Foci dynamics occur at higher levels with the luminal
ER marker in contrast to those observed with the ER membrane marker. This suggests that
proteins in the lumen are not evenly distributed compared to proteins at the ER membrane. Our
work provides a framework to shed light on the unique features of the structure and dynamics of
the presynaptic ER.

Presynaptic ER network structures are only modestly perturbed in *Atlastin* mutants but are sensitized to overexpression of luminal proteins

Using the membrane marker tdTomato:Sec61β, we discovered that *Atlastin* mutants do indeed have ER networks at presynaptic terminals. This is in sharp contrast to the diffuse distribution that we and others observe for the lumenal marker BiP:sfGFP:HDEL, and argues against excessive ER fragmentation in these mutants (since we would in that case expect to also see fragmented membranes). While we did not observe gross defects in ER network structure using the membrane marker, our quantitative analysis revealed subtle differences in the overall distribution of tubules as well as of Sec61β itself. Future analyses of dynamics at Atlastin mutant synapses may reveal additional insights into how this ER shaping protein controls synaptic transmission, signaling, and development.

Given the very different phenotypes observed with each marker, which more accurately reflects the *Atlastin* mutant phenotype in the absence of any ER marker? We favor the hypothesis that the true *Atlastin* ER phenotype is more likely to be similar to the tdTomato:Sec61β distribution, and that BiP:sfGFP:HDEL overexpression further disrupts the ER in *Atlastin* mutants, for the following reasons: (1) tdTomato:Sec61β-expressing animals still show Atlastin-associated functional defects such as defective synaptic growth, indicating that Atlastin-dependent ER functions are disrupted without dramatic defects in ER structure; (2) BiP:sfGFP:HDEL exhibits some toxicity even in wild-type adult animals. (3) The BiP:sfGFP:HDEL phenotype becomes more severe over larval development, suggesting that it is a progressive gain-of-function. Our study underscores the need to generate new endogenous ER markers in *Drosophila* to minimize the detrimental effects of protein overexpression of ER lumen or ER membrane markers. Future studies should also examine ER mutants with both ER membrane and ER lumen markers, as specific defects could affect one and not the other, as observed in *Atlastin* mutants.

Why then might *Atlastin* mutants displace luminal proteins to the cytosol upon overexpression of the BiP:sfGFP:HDEL? Several possibilities include reduced ER membrane integrity, activation of ER-associated protein degradation (ERAD), disruption of protein translocation, or induction of ER stress permeability. We outline below how these pathways

could contribute to the displacement phenotype in *Atlastin* mutants. First, *Atlastin* mutants have lower levels of the ER membrane marker Sec61β and altered ER network structures that could indicate changes in the integrity of the ER membrane. ER tubules at presynaptic terminals of *Atlastin* mutants could therefore be susceptible to rupture when luminal proteins are overexpressed, resulting in passive displacement to the cytosol. Second, luminal proteins in *Atlastin* mutants could be targeted for extrusion from the ER through ERAD (Christianson et al., 2023). However, this seems unlikely since these proteins would be targeted for degradation by the proteasome in the cytosol, and we still observe luminal proteins in presynaptic terminals of *Atlastin* mutants. Third, protein translocation defects could promote the accumulation of luminal proteins in the cytosol (McKibbin et al., 2012). However, defective protein translocation would likely result in a global displacement of luminal proteins, especially in the cell body, which is the primary site of neuronal protein translation (Glock et al., 2021). Instead, the diffuse phenotype is specific to presynaptic terminals. However, our observation that *Atlastin* mutants have reduced levels of the ER membrane marker Sec61β could reflect mild translocation defects. Nevertheless, this defect is not likely to result in total displacement of luminal ER proteins at presynaptic terminals of *Atlastin* mutants with Phenotype B. Fourth, ER stress induced by apoptosis can increase the permeability of the ER membrane (without bursting), enabling luminal ER proteins to diffuse to the cytosol (Wang et al., 2011). However, we did not find the neuronal debris that is commonly observed at muscles following neuronal apoptosis, in either first or third-instar larvae of *Atlastin* mutants (Aponte-Santiago et al., 2020; Graf et al., 2011).

Finally, it is important to note that we found that only presynaptic terminals of *Atlastin* mutants are susceptible to the over expression of the BiP:sfGFP:HDEL marker, indicating unique properties of synaptic ER compared to other neuronal compartments. Previously, a ladder-like ER expansion was observed in mammalian spinal axons of Atlastin-fusion incompetent mutants, providing another example of spatially distinct Atlastin-dependent ER network structures (Zhu et al., 2022). We observed similar ladder-like structures in both control and *Atlastin* mutant axons labeled with tdTomato:Sec61β. Interestingly, the presence of a ladder-like structure and baskets at synapses argues against the notion that Atlastin is the sole protein responsible for shaping ER tubule fusion in axons. This is in contrast to the long and unbranched ER tubules found in Atlastin-deficient Hela or Cos-7 cells (Hu et al., 2009; Niu et al., 2019), and suggests that compensatory tubule-fusion machinery might be specific to neurons. This discrepancy warrants further investigation to gain a clearer understanding of the underlying mechanisms at play. We

also found that *Atlastin* mutants have reduced Sec61β levels at the ER membrane of presynaptic
terminals. This decrease in Sec61β parallels findings in *Drosophila* mutants of Reticulon 1,
another ER shaping and HSP linked gene. This study found that in addition to Sec61β, other ER
membrane proteins, such as STIM and Stukorf, exhibited lower levels on ER tubules, suggesting
overall changes in the levels of proteins at the ER membrane (Pérez-Moreno et al., 2023).
However, unlike *Atlastin* mutants, Reticulon 1 mutants are not sensitized to the overexpression
of luminal proteins (Pérez-Moreno et al., 2023). This suggests that ER networks of *Atlastin*
mutants have unique features compared to those of other ER-shaping proteins. Future work
should comprehensively examine the common and unique defects associated with Atlastin
dysfunction across neuronal compartments.

<u>Methods</u>

<u>*Drosophila* stocks</u>
Drosophila melanogaster was cultured on a standard medium at 22–25°C. Detailed genotypes
and sexes used for experiments are described in Table S1. Stocks used in this study
include: Vglut-Gal4 (Bloomington *Drosophila* Stock Center (BDSC #: 24635), BG57-Gal4 (gift
from Vivian Budnik), *atl²* (gift from James McNew), third chromosome deficiency (BDSC #:
7948), UAS-BiP:sfGFP:HDEL (BDSC #: 24635), 20XUAS-tdTomato:Sec61β (BDSC #: 64746),
UAS-CD8:RFP (BDSC #: 27391) and UAS-GFP (gift from Griffith lab).

<u>Live imaging of luminal and membrane ER markers</u>
1st and wandering 3rd instar larvae were dissected in Ca^{2+}-free HL3.1 (Feng et al., 2004) and
axons were severed from the central nervous system. Dissections were performed in the live
imaging slide described below. Larvae were imaged at room temperature with a 63X (n.a. 1.4)
oil immersion objective, using an Airyscan LSM 880 microscope in super-resolution mode and
Zen Black software. Raw image stacks were processed in Zen Blue or Zen Black using 3D
Airyscan processing with automatic settings.

For 3rd instar larvae, Z-stacks of presynaptic type Ib terminals at muscle 4 in segment A5 (unless
otherwise noted) were acquired to measure the levels of luminal ER markers, to categorize ER
network phenotypes, or for morphological measurements. For 1st instar larvae, live larval fillets
were stained with 1/250 1.5 mg/ml HRP Alexa Fluor 647 (Jackson ImmunoResearch) in HL3.1

for 5 min followed by a 1 min wash. Next, Z-stacks of 1st instar presynaptic terminals at muscle

4 in segment A3-A5 were acquired and used to categorize ER network phenotypes.

To record the dynamics of the luminal ER marker BiP:sfGFP:HDEL, Z-stacks of a single bouton

(8 slices with 0.5 micron spacing) were collected for 100 frames at a frame rate of 0.73

seconds/stack. To record ER dynamics with the membrane ER marker tdTomato:Sec6β, Z-

stacks of a single bouton (8 slices with 0.5 micron spacing) were collected for 60 frames at a

frame rate of 0.65 seconds/stack. Imaging for longer periods led to significant photobleaching.

<u>Construction of *in vivo* imaging slide for *Drosophila* larvae</u>

To make the reusable imaging slide, Press-To-Seal Silicone Isolators (Grace Bio-labs; CQS-

13R-2.0) were mounted onto rectangular glass slides, which then were filled from their centers

to their tops with Krayden Dow Sylgard 184 Silicone (Thermo Fisher Scientific). Each slide with

a Press-To-Seal Silicone Isolator was leveled to ensure the silicone cured evenly at room

temperature. Minutien pins (Fine Science Tools) were cut with nail clippers and used to stretch

the larvae and fix it in place. Pins were pressed into the cured Sylgard 184 Silicone and a few

drops of hemolymph-like solution HL3.1 were added. Next, a coverslip was placed on top of the

dissected larvae, each corner was gently pressed down, and excess HL3.1 was removed with a

Kimwipe, thus sealing the coverslip to the silicone isolator.

<u>Fixed tissue staining</u>

Wandering 3rd instar larvae were dissected in Ca^{2+}-free HL3.1 and fixed for 17 min in Ca^{2+}-free

HL3.1 containing 4% PFA. Larvae were blocked and permeabilized overnight in PBS containing

0.25% Saponin, 2.5% normal goat serum (NGS), 2.5% bovine serum albumin (BSA), and 0.1%

sodium azide. Fixed larvae (**Figure 3 - Supplement**) were stained with 1/500 FluoTag®-X4 anti-

RFP (#N0404, Nanotag Biotechnologies) at 4°C for 24 hrs and with 1/100 HRP Alexa Fluor 647

at room temperature for 2hrs. Stained larvae were mounted in ProLong Diamond Antifade

Mountant (#P36970; Thermo-Fisher Scientific, Waltham, MA, USA). Z-stacks were collected of

Drosophila larval NMJs in muscle 4 of segment A5 with a spinning disk confocal microscope at

room temperature on a Nikon Ni-E upright microscope equipped with 100x (n.a. 1.45) oil

immersion objective, a Yokogawa CSU-W1 spinning-disk head, an Andor iXon 897U EMCCD

camera. Images were collected using Nikon Elements AR software.

<u>Digitonin experiments</u>

Digitonin experiments were conducted in live fillets from larvae expressing the luminal ER

marker, UAS-BiP:sfGFP:HDEL under the control of the neuronal driver Vglut-GAL4. Fillets were

stained with 1/500 1.5 mg/ml HRP Alexa Fluor 594 (Jackson ImmunoResearch) or 1/250 HRP

Alexa Fluor 647 in HL3.1 for 5 min, followed by a 1 min HL3.1 wash. Next, pre-digitonin Z-stacks

were acquired from muscle 4 NMJs from segments A3 to A5 (both hemisegments) using the

spinning disk confocal microscope described above. Fillets were subsequently incubated with

0.05% digitonin in HL3.1 for 5 min, then rinsed with HL3.1 and re-incubated with HRP Alexa

Fluor 594 or HRP Alexa Fluor 647 for 5 min. Preps were then washed for 1 min with HL3.1.

Finally, post-digitonin Z-stacks were acquired of the same NMJs.

<u>Image analysis</u>

Debris, axons, and large fluorescent aggregates (sometimes observed for tdTomato:Sec61β)

were manually removed from all images before analyses of maximum projections or Z-stacks.

Neuronal masks were created by first conducting a background subtraction of the image (rolling

ball, 50 pixel radius), and then introducing a gaussian blur before thresholding using Otsu or Li

algorithms. <u>For live image analysis,</u> maximum intensity projections of each timepoint were

corrected for (1) photobleaching using histogram matching and (2) sample drift or muscle

contraction using the StackReg plugin (Thevenaz et al., 1998).

Pearson correlation coefficient (PCC) for measurement of ER dynamics: To measure the PCC

of ER membrane or luminal proteins, we employed a FIJI macro consisting of the following steps:

(1) We created an NMJ mask of the neuronal outline of presynaptic terminals using the ER

membrane or luminal ER protein marker. (2) We measured the PCC between the 1st frame of

the movie and every other frame. (3) We plotted the PCC over time to measure global changes

in ER structure.

*Segmentation of presynaptic ER networks (**Figure 3C-E**):* For segmentation, time-lapse movies

of boutons with complex basket-like ER networks were selected from control and *Atlastin*

mutants. Images were blinded and maximum projections were created to segment the ER in

FIJI. We used the Trainable Weka Segmentation (TWS) machine-learning tool (Arganda-

Carreras et al., 2017) to manually annotate the first and last time-points in each movie as a

training dataset, and generated a mask of the ER throughout the movie. The mask from the first

time-point was then used for analysis. Bouton masks were created by background subtraction

and blurring of the tdTomato:Sec61β signal to quantify the fraction ER area. Finally, the mean

intensity of the ER membrane marker tdTomato:Sec61β was calculated within the ER mask.

Presynaptic levels of ER membrane marker tdTomato:Sec61β *(**Figure 3 Supplement**)*: *A*

custom FIJI macro consisting of the following steps was employed on maximum intensity

projections,: (1) An NMJ mask was created using the α-HRP neuronal marker; (2) The integrated

density of the ER membrane marker tdTomato:Sec61β was measured within the α-HRP mask;

(3) The mean intensity of the ER membrane marker at presynaptic terminals was calculated by

dividing the tdTomato:Sec61β integrated density by the area of the presynaptic terminal.

Analysis of NMJ morphology: Satellite and total bouton number were manually counted from

blinded maximum intensity projections of fixed and α-HRP labeled larvae. Satellite boutons were

defined as any string of 5 or fewer boutons extending from the main axis of the NMJ.

Categorization of ER morphology and dynamics: For analysis of tdTomato:Sec61β−labeled ER

structures (**Figure 3C-G)**, maximum intensity projection images were blinded and the following

structures were categorized: (1) Complex baskets had three or more crossing ER tubules; (2)

Simple baskets formed a single circle of ER tubules, often close to the periphery of the bouton.

The number of such structures was then normalized to the length of the NMJ arbor in the image

(excluding any arbor under 50 μm), which was measured by manually tracing and summing the

lengths of each branch of the NMJ.

For dynamics analysis (**Figure 2**), we used the same dataset as the control in **Figure 3E-G.**

Boutons with complex basket-like ER networks were selected for analysis. Maximum intensity

projection timelapse movies were blinded and replayed 3-4 times before categorization. We

identified three distinct categories of ER dynamics in presynaptic terminals of *Drosophila* motor

neurons (note that a single bouton can exhibit multiple types of dynamics): (1) **ER budding** –

small vesicle-like structures emerging from the ER network and re-fusing in a different region;

(2) **ER tubule dynamics**, encompassing four subcategories as follows: (a) Tubule

displacement, involving tubules that changed their localization within boutons without detaching

from the ER network; (b) Tubule extension, where an ER tubule emerged and extended from

the ER network; (c) <u>Tubule retraction</u>, characterized by the retraction and disappearance of an
ER tubule from the ER network; and (d) <u>Tubule extension and retraction</u>, indicating an ER tubule
both extending and retracting from the ER network within the timelapse. (3) **Foci dynamics** –
round, bright puncta moving within the ER network and (4) **no dynamics** – boutons that lacked
observable dynamics.

Presynaptic levels of luminal ER proteins at Drosophila motor neurons: A custom FIJI macro
consisting of the following steps was employed on Z-stacks: (1) We created an NMJ mask using
the luminal ER protein marker. (2) Using this mask, we measured the neuron volume and the
integrated density of the marker. (3) We calculated the mean intensity of the marker by dividing
the integrated density by the volume of the presynaptic terminal.

Categorization of luminal ER marker distribution: Images were blinded before analysis to ensure
unbiased qualitative analysis of luminal ER marker BiP:sfGFP:HDEL in controls and *Atlastin*
mutants. Maximum intensity projections were then sorted into three phenotypic categories: (1)
"Phenotype A" - luminal ER marker localizes to ER networks but is partially displaced to the
cytosol, (2) **"Phenotype B"** - luminal ER marker mostly displaced to the cytosol and ER
networks are not detectable, and (3) **"No phenotype"** - luminal ER marker completely localizes
to ER networks.

Presynaptic levels of cytosolic or luminal ER proteins pre- and post-digitonin treatment: Images
were blinded and maximum projections were created for further analysis. Neuronal outlines were
obtained by manually tracing the α-HRP-positive region in FIJI. Next, the integrated density of
UAS-GFP or UAS-BiP:sfGFP:HDEL was quantified within the manually traced neuronal outlines.
Mean intensity was calculated by dividing the integrated density by the area of the presynaptic
terminal.

<u>Statistics</u>
GraphPad Prism software was used to perform statistical analyses and generate graphs.
Statistical analyses were conducted using parametric tests (e.g. one-way ANOVA, Student's t-
test) for normally distributed data, and non-parametric tests (e.g. Kruskal-Wallis, Mann-Whitney
tests) if the data was not normally distributed. Chi-squared tests were used for categorical data.
p-values were set to *<0.05, **<0.01, ***<0.001.

Acknowledgements:

We thank the Bloomington *Drosophila* Stock Center (Indiana University, Bloomington, IN, NIH P40OD018537) and James McNew for fly lines, and Steve Del Signore for help with image analysis. This work was supported by NINDS grants R01 NS103967 to A.A.R. and T32 007292 to M.C.Q.-F.

599   **Figure legends:**

601   **Figure 1. *In vivo* imaging of ER networks in *Drosophila* motor neurons.** (**A**) Schematic of setup for live imaging experiments in this study, consisting of a glass slide with a mounted silicone isolator filled with Sylgard. Small metal pins were used to stretch the larvae (small round dots). Finally, a coverslip was placed on top of the larval fillet for imaging with an oil objective on upright or inverted microscopes. (**B**) A representative single slice of an Airyscan microscopy Z-stack of the center (top) or periphery (bottom) of the cell body of a motor neuron expressing the luminal ER marker BiP:sfGFP:HDEL. Arrows indicate ER tubules and sheets. (**C**) Maximum projections of Z-stacks acquired with an Airyscan microscope of two different motor neuron axons expressing the luminal ER marker BiP:sfGFP:HDEL. (**D**) Representative maximum projection of a Z-stack acquired with an Airyscan microscope of a motor neuron presynaptic terminal expressing the luminal ER marker BiP:sfGFP:HDEL. The magenta outline highlights the neuronal boundary. Arrows indicate boutons containing ER baskets or ladder-like ER structures.

614   **Figure 2. Dynamics of luminal and membrane ER markers at presynaptic terminals.** (**A**, **top**) Maximum intensity projection of Airyscan Z-stacks from the beginning (t=0) and the end (t=39 sec) of a timelapse movie (**Movie S1**) of a single wild-type bouton expressing the ER membrane marker tdTomato:Sec61β. (**A, bottom**) Magnification of the region in the dotted red box of dynamics observed from 0 sec to 1.4 sec of the movie. (**B, top**) Maximum intensity projection of Airyscan Z-stacks from the beginning (t=0) and the end (t=72 sec) of a timelapse movie (**Movie S4**) of a single wild-type bouton expressing the luminal ER marker BiP:sfGFP:HDEL. (**B, bottom**) Magnification of the region in the dotted red box of dynamics observed from 0 sec to 1.4 sec of the movie. (**C**) Presynaptic terminals exhibiting ER budding, focus dynamics and various types of ER tubule dynamics (ER tubule displacement, tubule

extension, and tubule retraction). (**D**) Qualitative categorization of the percentage of boutons

that are dynamic or not dynamic expressing the ER membrane marker tdTomato: Sec61β. (**E**)

Qualitative categorization of the percentage of dynamic boutons expressing the ER membrane

marker tdTomato: Sec61β that exhibit ER budding, foci dynamics or ER tubule dynamics. (**F**)

Qualitative categorization of the percentage of dynamic boutons expressing the ER membrane

marker tdTomato: Sec61β that exhibit tubule displacement, tubule extension, tubule retraction

and both tubule extension and retraction. (**G**) Pearson's Correlation Coefficient (PCC) of the first

frame (t0) with each subsequent frame (tn) in movies of boutons expressing the ER membrane

marker tdTomato: Sec61β (gray, n = 14 boutons) or ER luminal marker BiP:sfGFP:HDEL (green,

n = 10 boutons). The shaded area shows the standard deviation, and the solid black line

indicates the mean. The table shows the results of a non-linear fit of each marker's dataset.

Figure 3. *Atlastin* mutants expressing the ER membrane tdTomato:Sec61β exhibit robust

ER networks. (**A, B**) Representative maximum projection of a Z-stack acquired live by Airyscan

microscopy of control and *Atlastin* mutant NMJs expressing the ER membrane marker tdTomato:

Sec61β at presynaptic terminals. These images are not contrast matched because *Atlastin*

mutants exhibit lower levels of the tdTomato:Sec61β (See **Figures 3F** and **Figure 3 – Figure**

Supplement C). Quantification of boutons with complex (**C**) or simple (**D**) ER baskets

normalized to NMJ length. **(E)** WEKA segmentation of the ER (shown in magenta) of presynaptic

terminals. (**F**) Quantification of ER area fraction in control and *Atlastin* mutant boutons. (**G**)

Quantification of the mean intensity of the ER membrane marker tdTomato:Sec61β (within the

ER mask), normalized to wild-type control. (**D**, **E**) Graphs show mean +/- s.e.m. from 15 control

and 13 *Atlastin* mutant NMJs. **Associated with Figure 3 – Figure Supplement.**

Figure 3 – Figure Supplement. *Atlastin* mutants have lower levels of the ER membrane

tdTomato: Sec61β at presynaptic terminals. (**A**) Quantification of satellite boutons normalized

to total bouton number in control and *Atlastin* mutants. Representative maximum intensity

projection of a spinning disk confocal Z-stack from fixed larvae expressing the ER membrane

marker tdTomato:Sec61β (black) and stained with the neuronal marker HRP (magenta) in (**B**)

control and (**C**) *Atlastin* mutants. Neuronal outline is shown in blue. (**D**) Quantification of the

intensity of the ER membrane marker tdTomato:Sec61β, normalized to wild-type control. (**E**)

Coefficient of variation (CoV) of the ER membrane marker tdTomato:Sec61β. (**D**, **E**) Graphs
show mean +/- s.e.m. from 30 control and 30 *Atlastin* mutant NMJs.

Figure 4. Luminal ER proteins are displaced to the cytosol in *Atlastin* mutants. Maximum
projection of a Z-stack acquired live with an Airyscan microscope of control (**A**) and *Atlastin*
mutant (**B**) NMJs expressing the ER lumen marker BiP:sfGFP:HDEL. (**B**) *Atlastin* mutants exhibit
two distinct phenotypes at presynaptic terminals that were categorized as Phenotype A (**B, top**)
or Phenotype B (**B, bottom**). Control and *Atlastin* mutant images are contrast matched. (**C**) A
qualitative categorization of presynaptic terminals in control and *Atlastin* mutants exhibiting
"Phenotype A", "Phenotype B" or "no phenotype". (**D**) Histogram of normalized mean intensities
of the ER lumen marker BiP:sfGFP:HDEL. Displacement does not strongly correlate with steady-
state levels of BiP:sfGFP:HDEL. (**E**) Quantification of the intensity of the luminal ER marker
BiP:sfGFP:HDEL. (**F**) Coefficient of variation (CoV) of the luminal ER marker BiP:sfGFP:HDEL
in neurons of control and *Atlastin* mutant. (**E, F**) Graphs show mean +/- s.e.m. from 8 control
and 15 *Atlastin* mutant NMJs. Associated with **Figure 4 – Figure Supplement**.

Figure 4 – Figure Supplement. Is and Ib motor neurons in various muscles and abdominal
segments of 3rd instar *Atlastin* mutant larvae exhibit similar cytosolic displacement of
BiP:sfGFP:HDEL. Maximum projection of a live Airyscan Z-stack from larvae expressing the
ER lumen marker BiP:sfGFP:HDEL at muscle 12-13 (**A, left**) Muscle 6-7 (**A, right**); Muscle 4 in
segment A2 (**B, left**) and Muscle 4 in segment A4 (**B, right**).

Figure 5. Luminal ER proteins are not displaced in other neuronal compartments or cell
types in *Drosophila* 3rd instar larvae. All images were acquired by live Airyscan microscopy
from larvae expressing the ER lumen marker BiP:sfGFP:HDEL with Vglut-GAL4 (motor neurons)
or BG57-GAL4 (muscles). (**Left panel**) Maximum projection of a Z-stack of motor neuron axons.
(**Middle panel**) A single slice of motor neuron cell bodies. (**Right panel**) A single slice of body
wall muscles.

Figure 6. The luminal displacement phenotype in *Atlastin* mutants is progressive. (**A**)
Maximum intensity projections of a live Airyscan Z-stack acquired of 1st instar larvae expressing
the ER lumen marker BiP:sfGFP:HDEL (black) and stained with the neuronal marker HRP
(magenta). (**C**) A qualitative categorization of presynaptic terminals in control and *Atlastin* mutant

1st instar larvae expressing the ER lumen marker BiP:sfGFP:HDEL that exhibit "Phenotype A",

"Phenotype B" or "No phenotype". Data are shown from 16 control NMJs and 16 *Atlastin* mutant

NMJs.

Figure 7. Luminal ER proteins displaced to the cytosol are sensitive to digitonin

treatment. (**A**) Maximum projections of Z-stacks acquired live with a spinning disk confocal of

boutons expressing with Vglut-Gal4 GFP (control 1), BiP:sfGFP:HDEL (control 2),

BiP:sfGFP:HDEL in *Atlastin* mutants exhibiting Phenotype A and BiP:sfGFP:HDEL in Atlastin

mutants exhibiting Phenotype B. (**B**) Diagram of the expected distribution patterns before and

after digitonin treatment of cytosolic proteins and luminal ER proteins. Note that digitonin

treatment causes partial ER fragmentation. **(C)** Fractional change of fluorescence intensity at

presynaptic terminals expressing GFP (cytosolic protein) or BiP:sfGFP:HDEL in control and

Atlastin mutants following digitonin treatment. Graphs show mean +/- s.e.m. from 18 control and

20 *Atlastin* mutant NMJs. Statistical significance was only calculated between BiP:sfGFP:HDEL

samples since the baseline fluorescence and distribution of this marker was sensitive to digitonin

treatment in a manner distinct from cytosolic GFP.

Movies S1-3. Timelapse of maximum intensity projections from Airyscan Z-stacks of presynaptic

terminals expressing the ER membrane marker tdTomato:Sec61β. Images were acquired at a

frame rate of 0.65 frames per second and are played at a frame rate of 10 frames per second.

Scale bar is 1 µm.

Movies S4-6. Timelapse of maximum intensity projections from Airyscan Z-stacks of presynaptic

terminals expressing the ER lumen marker BiP:sfGFP:HDEL. Images were acquired at a frame

rate of 0.73 frames per second and are played at a frame rate of 10 frames per second. Scale

bar is 1 µm.

References:

Aponte-Santiago NA, Ormerod KG, Akbergenova Y, Littleton JT. 2020. Synaptic Plasticity Induced by Differential Manipulation of Tonic and Phasic Motoneurons in *Drosophila*. *J Neurosci* **40**:6270–6288. doi:10.1523/JNEUROSCI.0925-20.2020

Arganda-Carreras I, Kaynig V, Rueden C, Eliceiri KW, Schindelin J, Cardona A, Sebastian Seung H. 2017. Trainable Weka Segmentation: a machine learning tool for microscopy pixel classification. *Bioinformatics* **33**:2424–2426. doi:10.1093/bioinformatics/btx180

Blackstone C. 2018. Hereditary spastic paraplegia. *Handbook of Clinical Neurology,* **Vol. 148**:633–651. doi:10.1016/B978-0-444-64076-5.00041-7

Blackstone C. 2012. Cellular Pathways of Hereditary Spastic Paraplegia. *Annu Rev Neurosci* **35**:25–47. doi:10.1146/annurev-neuro-062111-150400

Blackstone C, O'Kane CJ, Reid E. 2011. Hereditary spastic paraplegias: membrane traffic and the motor pathway. *Nat Rev Neurosci* **12**:31–42. doi:10.1038/nrn2946

Boutry M, Morais S, Stevanin G. 2019. Update on the Genetics of Spastic Paraplegias. *Curr Neurol Neurosci Rep* **19**:18. doi:10.1007/s11910-019-0930-2

Burgoyne T, Patel S, Eden ER. 2015. Calcium signaling at ER membrane contact sites. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 13th European Symposium on Calcium **1853**:2012–2017. doi:10.1016/j.bbamcr.2015.01.022

Carter SD, Hampton CM, Langlois R, Melero R, Farino ZJ, Calderon MJ, Li W, Wallace CT, Tran NH, Grassucci RA, Siegmund SE, Pemberton J, Morgenstern TJ, Eisenman L, Aguilar JI, Greenberg NL, Levy ES, Yi E, Mitchell WG, Rice WJ, Wigge C, Pilli J, George EW, Aslanoglou D, Courel M, Freyberg RJ, Javitch JA, Wills ZP, Area-Gomez E, Shiva S, Bartolini F, Volchuk A, Murray SA, Aridor M, Fish KN, Walter P, Balla T, Fass D, Wolf SG, Watkins SC, Carazo JM, Jensen GJ, Frank J, Freyberg Z. 2020. Ribosome-associated vesicles: A dynamic subcompartment of the endoplasmic reticulum in secretory cells. *Sci Adv* **6**:eaay9572. doi:10.1126/sciadv.aay9572

Chino H, Mizushima N. 2020. ER-Phagy: Quality Control and Turnover of Endoplasmic Reticulum. *Trends in Cell Biology* **30**:384–398. doi:10.1016/j.tcb.2020.02.001

Chouhan AK, Zhang J, Zinsmaier KE, Macleod GT. 2010. Presynaptic Mitochondria in Functionally Different Motor Neurons Exhibit Similar Affinities for Ca2+ But Exert Little Influence as Ca2+ Buffers at Nerve Firing Rates In Situ. *J Neurosci* **30**:1869–1881. doi:10.1523/JNEUROSCI.4701-09.2010

Christianson JC, Jarosch E, Sommer T. 2023. Mechanisms of substrate processing during ER-associated protein degradation. *Nat Rev Mol Cell Biol* 1–20. doi:10.1038/s41580-023-00633-8

Costantini LM, Fossati M, Francolini M, Snapp EL. 2012. Assessing the Tendency of Fluorescent Proteins to Oligomerize under Physiologic Conditions. *Traffic* **13**:643–649. doi:10.1111/j.1600-0854.2012.01336.x

Crapart C, Scott ZC, Konno T, Sharma A, Bailey DM, Avezov E, Koslover EF. 2023. Luminal transport rates through intact endoplasmic reticulum limit the magnitude of localized Ca2+ signals. doi:10.1101/2023.06.23.546357

de Juan-Sanz J, Holt GT, Schreiter ER, de Juan F, Kim DS, Ryan TA. 2017. Axonal Endoplasmic Reticulum Ca2+ Content Controls Release Probability in CNS Nerve Terminals. *Neuron* **93**:867-881.e6. doi:10.1016/j.neuron.2017.01.010

Del Signore SJ, Mitzner MG, Silveira AM, Fai TG, Rodal AA. 2022. An approach for quantitative mapping of synaptic periactive zone architecture and organization. *Mol Biol Cell* mbcE22080372. doi:10.1091/mbc.E22-08-0372

Deng C, Moradi M, Reinhard S, Ji C, Jablonka S, Hennlein L, Lüningschrör P, Doose S, Sauer
M, Sendtner M. 2021. Dynamic remodeling of ribosomes and endoplasmic reticulum in
axon terminals of motoneurons. *Journal of Cell Science* **134**:jcs258785.
doi:10.1242/jcs.258785
Deshpande M, Rodal AA. 2016. The Crossroads of Synaptic Growth Signaling, Membrane
Traffic and Neurological Disease: Insights from *Drosophila*: Synaptic Signaling and
Membrane Traffic in *Drosophila*. *Traffic* **17**:87–101. doi:10.1111/tra.12345
Espadas J, Pendin D, Bocanegra R, Escalada A, Misticoni G, Trevisan T, Velasco del Olmo A,
Montagna A, Bova S, Ibarra B, Kuzmin PI, Bashkirov PV, Shnyrova AV, Frolov VA,
Daga A. 2019. Dynamic constriction and fission of endoplasmic reticulum membranes
by reticulon. *Nat Commun* **10**:5327. doi:10.1038/s41467-019-13327-7
Feng Y, Ueda A, Wu C-F. 2004. A modified minimal hemolymph-like solution, HL3.1, for
physiological recordings at the neuromuscular junctions of normal and mutant
Drosophila larvae. *J Neurogenet* **18**:377–402. doi:10.1080/01677060490894522
Finsterer J, Löscher W, Quasthoff S, Wanschitz J, Auer-Grumbach M, Stevanin G. 2012.
Hereditary spastic paraplegias with autosomal dominant, recessive, X-linked, or
maternal trait of inheritance. *Journal of the Neurological Sciences* **318**:1–18.
doi:10.1016/j.jns.2012.03.025
Fowler PC, Garcia-Pardo ME, Simpson JC, O'Sullivan NC. 2019. NeurodegenERation: The
Central Role for ER Contacts in Neuronal Function and Axonopathy, Lessons From
Hereditary Spastic Paraplegias and Related Diseases. *Front Neurosci* **13**.
doi:10.3389/fnins.2019.01051
Friedman JR, Lackner LL, West M, DiBenedetto JR, Nunnari J, Voeltz GK. 2011. ER tubules
mark sites of mitochondrial division. *Science* **334**:358–362.
doi:10.1126/science.1207385
Glock C, Biever A, Tushev G, Nassim-Assir B, Kao A, Bartnik I, tom Dieck S, Schuman EM.
2021. The translatome of neuronal cell bodies, dendrites, and axons. *Proceedings of
the National Academy of Sciences* **118**:e2113929118. doi:10.1073/pnas.2113929118
Graf ER, Heerssen HM, Wright CM, Davis GW, DiAntonio A. 2011. Stathmin is Required for
Stability of the Drosophila Neuromuscular Junction. *J Neurosci* **31**:15026–15034.
doi:10.1523/JNEUROSCI.2024-11.2011
Gregorio CD, Delgado R, Ibacache A, Sierralta J, Couve A. 2017. Drosophila Atlastin in motor
neurons is required for locomotion and presynaptic function. *J Cell Sci* **130**:3507–3516.
doi:10.1242/jcs.201657
Holcman D, Parutto P, Chambers JE, Fantham M, Young LJ, Marciniak SJ, Kaminski CF, Ron
D, Avezov E. 2018. Single particle trajectories reveal active endoplasmic reticulum
luminal flow. *Nat Cell Biol* **20**:1118–1125. doi:10.1038/s41556-018-0192-2
Hoyer MJ, Chitwood PJ, Ebmeier CC, Striepen JF, Qi RZ, Old WM, Voeltz GK. 2018. A Novel
Class of ER Membrane Proteins Regulates ER-Associated Endosome Fission. *Cell*
175:254-265.e14. doi:10.1016/j.cell.2018.08.030
Hu J, Shibata Y, Zhu P-P, Voss C, Rismanchi N, Prinz WA, Rapoport TA, Blackstone C. 2009.
A Class of Dynamin-like GTPases Involved in the Generation of the Tubular ER
Network. *Cell* **138**:549–561. doi:10.1016/j.cell.2009.05.025
Hu X, Wu F, Sun S, Yu W, Hu J. 2015. Human atlastin GTPases mediate differentiated fusion
of endoplasmic reticulum membranes. *Protein Cell* **6**:307–311. doi:10.1007/s13238-
015-0139-3
King C, Sengupta P, Seo AY, Lippincott-Schwartz J. 2020. ER membranes exhibit phase
behavior at sites of organelle contact. *Proceedings of the National Academy of
Sciences* **117**:7225–7235. doi:10.1073/pnas.1910854117

Le Gall S, Neuhof A, Rapoport T. 2004. The Endoplasmic Reticulum Membrane Is Permeable
to Small Molecules. *MBoC* **15**:447–455. doi:10.1091/mbc.e03-05-0325
Lee M, Paik SK, Lee M-J, Kim Y-J, Kim S, Nahm M, Oh S-J, Kim H-M, Yim J, Lee CJ, Bae YC,
Lee S. 2009. Drosophila Atlastin regulates the stability of muscle microtubules and is
required for synapse development. *Developmental Biology* **330**:250–262.
doi:10.1016/j.ydbio.2009.03.019
Liu X, Guo X, Niu L, Li X, Sun F, Hu J, Wang X, Shen K. 2019. Atlastin-1 regulates
morphology and function of endoplasmic reticulum in dendrites. *Nat Commun* **10**:1–15.
doi:10.1038/s41467-019-08478-6
Lu L, Ladinsky MS, Kirchhausen T. 2009. Cisternal Organization of the Endoplasmic Reticulum
during Mitosis. *MBoC* **20**:3471–3480. doi:10.1091/mbc.e09-04-0327
Lü L, Niu L, Hu J. 2020. "At last in" the physiological roles of the tubular ER network. *Biophys*
Rep **6**:105–114. doi:10.1007/s41048-020-00113-y
Lu M, Tartwijk FW van, Lin JQ, Nijenhuis W, Parutto P, Fantham M, Christensen CN, Avezov
E, Holt CE, Tunnacliffe A, Holcman D, Kapitein L, Schierle GSK, Kaminski CF. 2020.
The structure and global distribution of the endoplasmic reticulum network are actively
regulated by lysosomes. *Science Advances* **6**:eabc7209. doi:10.1126/sciadv.abc7209
McKibbin C, Mares A, Piacenti M, Williams H, Roboti P, Puumalainen M, Callan AC, Lesiak-
Mieczkowska K, Linder S, Harant H, High S, Flitsch SL, Whitehead RC, Swanton E.
2012. Inhibition of protein translocation at the endoplasmic reticulum promotes
activation of the unfolded protein response. *Biochemical Journal* **442**:639–648.
doi:10.1042/BJ20111220
Niu L, Ma T, Yang F, Yan B, Tang X, Yin H, Wu Q, Huang Y, Yao Z-P, Wang J, Guo Y, Hu J.
2019. Atlastin-mediated membrane tethering is critical for cargo mobility and exit from
the endoplasmic reticulum. *PNAS* **116**:14029–14038. doi:10.1073/pnas.1908409116
Obara CJ, Nixon-Abell J, Moore AS, Riccio F, Hoffman DP, Shtengel G, Xu CS, Schaefer K,
Pasolli HA, Masson J-B, Hess HF, Calderon CP, Blackstone C, Lippincott-Schwartz J.
2022. Motion of single molecular tethers reveals dynamic subdomains at ER-
mitochondria contact sites. doi:10.1101/2022.09.03.505525
Oliva MK, Pérez-Moreno JJ, O'Shaughnessy J, Wardill TJ, O'Kane CJ. 2020. Endoplasmic
Reticulum Lumenal Indicators in Drosophila Reveal Effects of HSP-Related Mutations
on Endoplasmic Reticulum Calcium Dynamics. *Front Neurosci* **14**.
doi:10.3389/fnins.2020.00816
Orso G, Pendin D, Liu S, Tosetto J, Moss TJ, Faust JE, Micaroni M, Egorova A, Martinuzzi A,
McNew JA, Daga A. 2009. Homotypic fusion of ER membranes requires the dynamin-
like GTPase Atlastin. *Nature* **460**:978–983. doi:10.1038/nature08280
O'Sullivan NC, Jahn TR, Reid E, O'Kane CJ. 2012. Reticulon-like-1, the Drosophila orthologue
of the Hereditary Spastic Paraplegia gene reticulon 2, is required for organization of
endoplasmic reticulum and of distal motor axons. *Hum Mol Genet* **21**:3356–3365.
doi:10.1093/hmg/dds167
Özkan N, Koppers M, van Soest I, van Harten A, Jurriens D, Liv N, Klumperman J, Kapitein
LC, Hoogenraad CC, Farías GG. 2021. ER - lysosome contacts at a pre-axonal region
regulate axonal lysosome availability. *Nat Commun* **12**:4493. doi:10.1038/s41467-021-
24713-5
Öztürk Z, O'Kane CJ, Pérez-Moreno JJ. 2020. Axonal Endoplasmic Reticulum Dynamics and
Its Roles in Neurodegeneration. *Frontiers in Neuroscience* **14**:1–33.
doi:10.3389/fnins.2020.00048

Perez-Alvarez A, Yin S, Schulze C, Hammer JA, Wagner W, Oertner TG. 2020. Endoplasmic
reticulum visits highly active spines and prevents runaway potentiation of synapses.
Nature Communications **11**:5083. doi:10.1038/s41467-020-18889-5

Pérez-Moreno JJ, Smith RC, Oliva MK, Gallo F, Ojha S, Müller KH, O'Kane CJ. 2023.
Drosophila SPG12 ortholog, reticulon-like 1, governs presynaptic ER organization and
Ca2+ dynamics. *J Cell Biol* **222**:e202112101. doi:10.1083/jcb.202112101

Raffaello A, Mammucari C, Gherardi G, Rizzuto R. 2016. Calcium at the Center of Cell
Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes.
Trends in Biochemical Sciences **41**:1035–1049. doi:10.1016/j.tibs.2016.09.001

Raiborg C, Wenzel EM, Pedersen NM, Olsvik H, Schink KO, Schultz SW, Vietri M, Nisi V,
Bucci C, Brech A, Johansen T, Stenmark H. 2015. Repeated ER–endosome contacts
promote endosome translocation and neurite outgrowth. *Nature* **520**:234–238.
doi:10.1038/nature14359

Rao K, Stone MC, Weiner AT, Gheres KW, Zhou C, Deitcher DL, Levitan ES, Rolls MM. 2016.
Spastin, atlastin, and ER relocalization are involved in axon but not dendrite
regeneration. *MBoC* **27**:3245–3256. doi:10.1091/mbc.E16-05-0287

Rismanchi N, Soderblom C, Stadler J, Zhu P-P, Blackstone C. 2008. Atlastin GTPases are
required for Golgi apparatus and ER morphogenesis. *Hum Mol Genet* **17**:1591–1604.
doi:10.1093/hmg/ddn046

Rowland AA, Chitwood PJ, Phillips MJ, Voeltz GK. 2014. ER Contact Sites Define the Position
and Timing of Endosome Fission. *Cell* **159**:1027–1041. doi:10.1016/j.cell.2014.10.023

Schwarz DS, Blower MD. 2016. The endoplasmic reticulum: structure, function and response
to cellular signaling. *Cell Mol Life Sci* **73**:79–94. doi:10.1007/s00018-015-2052-6

Scott ZC, Brown AI, Mogre SS, Westrate LM, Koslover EF. 2021. Diffusive search and
trajectories on tubular networks: a propagator approach. *Eur Phys J E* **44**:80.
doi:10.1140/epje/s10189-021-00083-0

Snow PM, Patel NH, Harrelson AL, Goodman CS. 1987. Neural-specific carbohydrate moiety
shared by many surface glycoproteins in Drosophila and grasshopper embryos. *J
Neurosci* **7**:4137–4144. doi:10.1523/JNEUROSCI.07-12-04137.1987

Spits M, Neefjes J. 2021. Mobile late endosomes modulate peripheral endoplasmic reticulum
network architecture. *EMBO reports* **22**:e50815. doi:10.15252/embr.202050815

Subramanian K, Meyer T. 1997. Calcium-Induced Restructuring of Nuclear Envelope and
Endoplasmic Reticulum Calcium Stores. *Cell* **89**:963–971. doi:10.1016/S0092-
8674(00)80281-0

Summerville JB, Faust JF, Fan E, Pendin D, Daga A, Formella J, Stern M, McNew JA. 2016.
The effects of ER morphology on synaptic structure and function in Drosophila
melanogaster. *J Cell Sci* **129**:1635–1648. doi:10.1242/jcs.184929

Sun L, Yu F, Ullah A, Hubrack S, Daalis A, Jung P, Machaca K. 2011. Endoplasmic Reticulum
Remodeling Tunes IP3-Dependent Ca2+ Release Sensitivity. *PLOS ONE* **6**:e27928.
doi:10.1371/journal.pone.0027928

Thevenaz P, Ruttimann UE, Unser M. 1998. A pyramid approach to subpixel registration
based on intensity. *IEEE Trans on Image Process* **7**:27–41. doi:10.1109/83.650848

Wang X, Olberding KE, White C, Li C. 2011. Bcl-2 proteins regulate ER membrane
permeability to luminal proteins during ER stress-induced apoptosis. *Cell Death Differ*
18:38–47. doi:10.1038/cdd.2010.68

Wu Y, Whiteus C, Xu CS, Hayworth KJ, Weinberg RJ, Hess HF, Camilli PD. 2017. Contacts
between the endoplasmic reticulum and other membranes in neurons. *PNAS*
114:E4859–E4867. doi:10.1073/pnas.1701078114

Yalçın B, Zhao L, Stofanko M, O'Sullivan NC, Kang ZH, Roost A, Thomas MR, Zaessinger S,
Blard O, Patto AL, Sohail A, Baena V, Terasaki M, O'Kane CJ. 2017. Modeling of
axonal endoplasmic reticulum network by spastic paraplegia proteins. *eLife* **6**:1–27.
doi:10.7554/eLife.23882
Zamponi E, Meehl JB, Voeltz GK. 2022. The ER ladder is a unique morphological feature of
developing mammalian axons. *Dev Cell* **57**:1369-1382.e6.
doi:10.1016/j.devcel.2022.05.002
Zhu P-P, Hung H-F, Batchenkova N, Nixon-Abell J, Henderson J, Zheng P, Renvoisé B, Pang
S, Xu CS, Saalfeld S, Funke J, Xie Y, Svara F, Hess HF, Blackstone C. 2022.
Transverse endoplasmic reticulum expansion in hereditary spastic paraplegia
corticospinal axons. *Hum Mol Genet* **31**:2779–2795. doi:10.1093/hmg/ddac072
Zhu P-P, Soderblom C, Tao-Cheng J-H, Stadler J, Blackstone C. 2006. SPG3A protein
atlastin-1 is enriched in growth cones and promotes axon elongation during neuronal
development. *Hum Mol Genet* **15**:1343–1353. doi:10.1093/hmg/ddl054

Figure 1

A

Side view

Coverslip

Silicone isolator

Glass slide

Top view

Glass slide

Sylgard

Silicone isolator

B

Cell body

Center

ER tubule

5µm

Periphery

Sheets

5µm

BiP:sfGFP:HDEL (ER lumen marker)

C

Example #1   **Axon**

5µm

Example #2

5µm

BiP:sfGFP:HDEL (ER lumen marker)

D

NMJ

ER "basket"

Ladder-like ER

5µm

BiP:sfGFP:HDEL (ER lumen marker)
Neuronal outline

Figure 2

A ER membrane marker (Sec61β)

t = 0sec   t = 39sec

2µm

0sec   0.7sec   1.4sec

0.5µm

B ER lumen (BiP:sfGFP:HDEL)

t = 0   t = 72sec

2 µm

Foci

0sec   0.7sec   1.4sec

0.5µm

C

ER budding

0s   3.9s   7.2s

0.5µm

tdTomato:Sec61β

Focus dynamics

0s   2.6s   9.1s

0.5µm

tdTomato:Sec61β

Tubule dynamics

Tubule displacement

19.5s   39s

1µm

tdTomato:Sec61β

Tubule Extension

0s   9.1s

0.5µm

tdTomato:Sec61β

Tubule retraction

0s   5.2s   7.2s

0.5µm

tdTomato:Sec61β

D % of boutons — Dynamic, Not dynamic

E % of boutons — Dynamic puncta, Foci dynamics, Tubule dynamics

F % of boutons — Displacement, Extension, Retraction, Extension AND retraction — Tubule dynamics

G PCC (t1-tn) vs Δt (seconds); ER membrane, ER lumen

	K (1/s)	Half-life (s)	Tau (s)
ER membrane	0.01618	42.85	61.82
ER lumen	0.02697	25.70	37.08

Figure 3

A Control

NMJ #1

NMJ #2

Complex ER basket

NMJ #3

Simple ER basket

ER membrane (tdTomato:Sec61β)

5µm 5µm 5µm

B Atlastin mutant

NMJ #1 NMJ #2 NMJ #3

ER membrane (tdTomato:Sec61β)

5µm 5µm 5µm

C # of complex ER baskets/ NMJ length — ns — Control, Atlastin mutant

D # of simple ER baskets/ NMJ length — ns — Control, Atlastin mutant

E Control / Atlastin mutant — 1µm — tdTomato:Sec61β, ER outline

F ER area/ bouton area — * — Control, Atlastin mutant

G tdTomato:Sec61β integrated density/ area (µm) — * — Control, Atlastin mutant

Figure 3-Supplement

Figure 4

A Control
BiP:sfGFP:HDEL

B Atlastin mutant
Phenotype A (partial loss of ER networks)
BiP:sfGFP:HDEL

Phenotype B (severe loss of ER networks)
BiP:sfGFP:HDEL

C
% of NMJs

☐ Control
☐ Atlastin mutant
p-value<0.0001

Phenotype A   Phenotype B   No phenotype

D
Relative frequency (%)

■ Control
■ Atlastin mutant (Phenotype A)
■ Atlastin mutant (Phenotype B)

BiP:sfGFP:HDEL mean intensity
100 200 300 400 500 600 700 800 900

E
BiP:sfGFP:HDEL integrated density/ NMJ volume (µm³)

ns

Control   Atlastin mutants

F
BiP:sfGFP:HDEL CoV

**

Control   Atlastin mutants

Figure 4-Supplement

Figure 5

Motor neurons

Body wall muscle

Control

Axons

Cell bodies

2μm

2μm

2μm

Genotypes: Vglut/+;UAS-BiP:sfGFP:HDEL/+

UAS-BiP:sfGFP:HDEL/+;
BG57-Gal4/+

Atlastin mutant

2μm

2μm

2μm

Genotypes: Vglut/+;UAS-BiP:sfGFP:HDEL/+; atl^2/atl def

UAS-BiP:sfGFP:HDEL/+;
atl^2/BG57-Gal4,atl def

Figure 6

Figure 7

A

	Control 1	Control 2	Atl mutant Phenotype A	Atl mutant Phenotype B

Pre-digitonin: GFP; ER lumen; ER lumen; ER lumen — 3µm

Post-digitonin — 3µm

Post-digitonin (contrast enhanced)

B

Pre-digitonin

Control: Cytosolic marker; ER lumen marker

Atl mutant *Phenotype A*; *Phenotype B* — ER lumen marker

Post-digitonin

Control: Cytosolic marker; ER lumen marker

Atl mutant *Phenotype A*; *Phenotype B* — ER lumen marker

C

$\Delta F/F$ (Pre$_{dig}$ - Post$_{dig}$)/ Pre$_{dig}$

ns

*

GFP; BiP; BiP (Atl mutant; Phenotype A); BiP (Atl mutant; Phenotype B)

Table S1 – Detailed genotypes and statistics reporting

Experiments were done at muscle 4 unless otherwise noted
All analyses were performed in Fiji
A: Pearson's correlation coefficient over time
B: Mean intensity of marker
C: CoV

Figure	Genotype/Conditions	N	Measurement	Statistical Test(s)
Figure 2D-G Dynamics of ER luminal and membrane markers Airyscan	ER lumen: Vglut/+;UAS-BiP:sfGFP:HDEL/+ ER Membrane: Vglut/+;;UAS-tdTomato:Sec61β/+	2 larvae/ 10 boutons (segment A3) 4 larvae/14 boutons (segments A2-A5)	ER dynamics A	No statistics
Figure 3C,D Complex and simple ER network structure analysis Airyscan	Control: Vglut/+;UAS-tdTomato:Sec61β/+ *Atlastin* mutant: Vglut/+;UAS-tdTomato:Sec61β/+;atl²/atl def	4 larvae/14 NMJs (segments A2-A5) 4 larvae/11 NMJs (segments A2-A5)	Qualitative analysis of ER network structure in boutons expressing tdTomato:Sec61β	Figure 3C: Unpaired Mann-Whitney test, P<0.0001 Figure 3D: Unpaired Mann-Whitney test, P<0.0001
Figure 3 F,G ER network structure bouton analysis	Control: Vglut/+;UAS-tdTomato:Sec61β/+	4 larvae/15 boutons (segments A2-A5)	ER network per bouton area and Sec61β levels B, C	Figure 3F: Unpaired Mann-Whitney test, P<0.0001

	Atlastin mutant: Vglut/+;UAS-tdTomato:Sec61β/+;atl²/atl def	4 larvae/ 13 boutons (segments A2-A5)		Figure 3G: Unpaired Mann-Whitney test, P<0.0001
Figure 3 – Figure Supplement (A)	Control: Vglut/+;UAS-tdTomato:Sec61β/+ *Atlastin* mutant: Vglut/+;UAS-tdTomato:Sec61β/+;atl²/atl def	8 larvae/ 30 NMJs (segments A2- A5) 8 larvae/ 30 NMJs (segments A2-A5)	Count of satellite boutons	Figure 3 – Figure Supplement (A): Unpaired Mann-Whitney test, P<0.0001
Figure 3 – Figure Supplement (D-E) Overall levels of ER membrane markers at nerve terminals Spinning disk	Control: Vglut/+;UAS-tdTomato:Sec61β/+ *Atlastin* mutant: Vglut/+;UAS-tdTomato:Sec61β/+;atl²/atl def	8 larva/ 30 NMJs (segments A2- A5) 8 larva/ 30 NMJs (segments A2-A5)	Sec61β levels B, C	Figure 3 – Figure Supplement (D): Unpaired Mann-Whitney test, P<0.0001 Figure 3 – Figure Supplement (E): Unpaired t-test, P<0.05
Figure 4C Categorization of ER network phenotypes in *Atlastin* mutants expressing luminal ER markers Airyscan	Control: Vglut/+;UAS-BiP:sfGFP:HDEL/+ *Atlastin* mutant: Vglut/+; UAS-BiP:sfGFP:HDEL/+;atl²/atl def	6 larvae/ 21 NMJs (Segments A2-A5) 8 larvae/ 22 NMJs (Segments A2-A5)	Categories: - Phenotype A - Phenotype B No phenotype	Chi-square, P<0.0001

Figure 4D-F	Control: Vglut/+;UAS-BiP:sfGFP:HDEL/+	2 larvae/ 8 NMJs	BiP:sfGFP:HDEL levels	Figure 4E: Unpaired Mann-Whitney test, $P<0.0001$
Overall levels of ER lumen markers at nerve terminals of *Atlastin* mutants Airyscan		(Segments A2-A5) 4 larvae/ 15 NMJs	B,C	
	Atlastin mutant: Vglut/+; UAS-BiP:sfGFP:HDEL/+;atl^2/atl def	(Segments A2-A5)		Figure 4F: Unpaired t-test, $P<0.05$
Figure 6C Categorization of ER network phenotypes in1st instar larvae *Atlastin* mutants expressing a luminal ER marker Airyscan	Control: Vglut/+;UAS-BiP:sfGFP:HDEL/+	7 larvae/ 16 NMJs (Segments A4-A5) 6 larvae/ 16NMJs (Segments A4-A5)	Categories: - Phenotype A - Phenotype B - No phenotype	Chi-square, $P<0.0001$
	Atlastin mutant: Vglut/+; UAS-BiP:sfGFP:HDEL/+;atl^2/atl def			
Figure 7C Mean intensity of luminal ER markers with digitonin treatment Spinning disk	Control1: Vglut/+;UAS-GFP/+	4 larvae/ 13NMJs (Segments A4-A5) 5 larvae/ 18 NMJs (Segments A4-A5) 5 larvae/ 20 NMJs (Segments A4-A5)	Change in fluorescence before and after digitonin treatment	Non-parametric One-way ANOVA for BiP samples only (Kruskal-Wallis test), $P<0.0001$
	Control2: Vglut/+;UAS-BiP:sfGFP:HDEL/+			
	Atlastin mutant: Vglut/+; UAS-BiP:sfGFP:HDEL/+;atl^2/atl def			