Title: Delayed localization of A-type lamins to the rupture sites in Hutchinson–

Gilford progeria syndrome

Yohei Kono [1, 2, 3], Chan-Gi Pack [4, 5], Takehiko Ichikawa [2], Arata Komatsubara [6], Stephen

A. Adam [7], Keisuke Miyazawa [2, 3], Loïc Rolas [8], Sussan Nourshargh [8], Ohad Medalia [9],

Robert D. Goldman [7], Takeshi Fukuma [2, 3], Hiroshi Kimura [1, 6], Takeshi Shimi [1, 2*]

Author Affiliation:

[1] Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology,

Yokohama, Japan.

[2] Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, Japan.

[3] Faculty of Frontier Engineering, Institute of Science and Engineering, Kanazawa

University, Kanazawa, Japan.

[4] Convergence Medicine Research Center, Asan Institute for Life Science, Asan Medical

Center, Seoul, Korea.

[5] Department of Biomedical Engineering, University of Ulsan College of Medicine, Seoul,

Korea.

[6] School of Life Science and Technology, Tokyo Institute of Technology, Yokohama,

Japan.

[7] Department of Cell and Developmental Biology, Feinberg School of Medicine,

Northwestern University, Chicago, IL.

[8] Centre for Microvascular Research, William Harvey Research Institute, Faculty of

Medicine and Dentistry, Queen Mary University of London, London, UK.

[9] Department of Biochemistry, University of Zurich, Zurich, Switzerland.

The corresponding author:

Takeshi Shimi, Ph.D., Nano Life Science Institute (WPI-NanoLSI), Kanazawa University,

Kakuma-machi, Kanazawa, 920-1192, Japan.

Phone: +81-76-234-4540; E-mail: t-shimi@staff.kanazawa-u.ac.jp

Author ORCIDs

Yohei Kono: https://orcid.org/0000-0001-8264-9417

Chan-Gi Pack: https://orcid.org/0000-0002-6578-3099

Takehiko Ichikawa: https://orcid.org/0000-0002-2438-5502

Stephen A. Adam: https://orcid.org/0000-0003-3444-7655

Keisuke Miyazawa: https://orcid.org/0000-0002-5012-8040

Loïc Rolas: https://orcid.org/0000-0001-7617-0768

Sussan Nourshargh: https://orcid.org/0000-0001-5677-1806

Ohad Medalia: https://orcid.org/0000-0003-0994-2937

Robert D. Goldman: http://orcid.org/0000-0003-0383-1435

Takeshi Fukuma: https://orcid.org/0000-0001-8971-6002

Hiroshi Kimura: https://orcid.org/0000-0003-0854-083X

Takeshi Shimi: https://orcid.org/0000-0002-9864-1820

Keywords

Lamin, HGPS, Lamina, Nuclear Envelope Rupture, Nucleus

Abstract

The nuclear lamina (NL) lines the nuclear envelope (NE) to maintain nuclear structure in metazoan cells. The major NL components, the nuclear lamins contribute to the protection against NE rupture induced by mechanical stress. Lamin A (LA) and a short form of the splicing variant lamin C (LC) are diffused from the nucleoplasm to sites of NE rupture in immortalized mouse embryonic fibroblasts (MEFs). LA localization to the rupture sites is significantly slow and weak compared to LC because of its relatively small pool in the nucleoplasm, but the precise mechanism remains unknown. In this study, we induce NE rupture in wild-type and LA/C-knockout MEFs, and Hutchinson–Gilford Progeria syndrome (HGPS) knock-in MEFs that express progerin, a LA mutant lacking the second proteolytic cleavage site, by laser microirradiation and AFM indentation. The farnesylation at the CaaX motif of unprocessed LA and the inhibition of the second proteolytic cleavage decrease the nucleoplasmic pool and slow the localization to the rupture sites in a long-time window (60-70 min) after the induction of NE rupture. Our data could explain the defective repair of NE rupture in HGPS through the farnesylation at the CaaX motif of unprocessed progerin. In addition, unique segments in LA-specific tail region were critical for inhibiting the rapid accumulation within a short-time window (3 min).

Significance Statement

Lamin A (LA) is slowly localized to sites of nuclear envelope (NE) rupture compared to lamin C (LC) because of its relatively small pool in the nucleoplasm. This study reveals that the farnesylation at the CaaX motif of unprocessed LA and the inhibition of the second proteolytic cleavage decreases the nucleoplasmic pool and slows the localization

70　to the rupture sites within a long-time window (60-70 min) after the induction of NE

71　rupture, which may explain the defective repair of NE rupture in Hutchinson–Gilford

72　Progeria syndrome (HGPS). In addition, unique segments in LA-specific tail region

73　cooperate with each other for inhibiting the rapid accumulation within a short-time

74　window (3 min), which is observed with LC.

Introduction

In eukaryotic cell nuclei, the nuclear envelope (NE) spatially separates the nuclear genome from the cytoplasm. The barrier function of the NE is due to two phospholipid bilayers of the inner and outer nuclear membranes (INM, ONM). The INM is in contact with chromatin at the nuclear side via INM proteins (1), whereas the ONM is connected with the cytoskeletal system through the linker of nucleoskeleton and cytoskeleton (LINC) complex (2), and is continuous with membranous organelles including the endoplasmic reticulum (ER) (3). The nuclear lamina (NL) is closely apposed to the INM and associates with heterochromatin to support nuclear structure. Nuclear pore complexes (NPCs) fill holes formed by the INM-ONM fusion for nucleo-cytoplasmic transport and interact with euchromatin (4).

The NL structure consists of the nuclear lamins and associated proteins (5). The lamins are type V intermediate filament proteins, which are subdivided into A-type lamins (lamins A [LA] and C [LC]) and B-type lamins (lamins B1 [LB1] and B2 [LB2]) (6, 7). LA and LC are derived from the single gene *LMNA* by alternative splicing (8), whereas LB1 and LB2 are encoded by separate genes *LMNB1* and *LMNB2*, respectively (9-12). Lamin molecules assemble into a filament with a diameter of ~3.5 nm and these filaments appear to non-randomly lay on top of each other to form meshworks of the NL with a thickness of ~13.5 nm in immortalized mouse embryonic fibroblasts (MEFs) (13, 14). LA/C and LB1 filaments associate with the outer ring structure of the NPC and are involved in regulating the distribution of the meshworks and NPCs (13-16).

The −CaaX box at the C terminus of unprocessed LA, LB1 and LB2 (pre-LA, pre-LB1 and pre-LB2, respectively) is post-translationally processed in steps, including the attachment of farnesyl group to the cysteine residue of the −CaaX box by a

farnesyltransferase (17, 18), proteolytic cleavage of three residues (−aaX) by an aaX

endopeptidase (19) and carboxyl methylation of the terminal carboxylic acid group

(−COOH) of the C-terminal cysteine residue by a carboxyl methyltransferase (20). The

maturation of LB1 and LB2 is completed by these steps. In case of the farnesylated and

carboxymethylated pre-LA, an additional 15 C-terminal residues including the

farnesylated and carboxymethylated cysteine, are cleaved off by Zinc metallopeptidase

STE24 (Zmpste24/FACE1) to produce the mature form (mature-LA) (21, 22).

The NE is locally ruptured during physiological and pathological circumstances.

Immediately after NE rupture, NE components are recruited to the rupture sites with the

endosomal sorting complex required for transport-III (ESCRT-III) complex and barrier-

to-autointegration factor (BAF) (23-25). Nuclear DNA adjacent to the rupture sites is

recognized by the DNA sensors cyclic GMP-AMP synthase (cGAS) and its downstream

signaling effector STING (23-25). The frequencies of spontaneous and forced NE rupture

are significantly increased by the depletion of lamins (26-29). LA/C promotes the

accumulation of BAF and cGAS at the rupture sites (30). Numerous mutations that have

been found throughout the *LMNA* gene cause a spectrum of human genetic disorders,

collectively called laminopathies (31). Laminopathies associated with dilated

cardiomyopathy (DCM), muscular dystrophy (MD), familial partial lipodystrophy

(FPLD) and Hutchinson–Gilford Progeria syndrome (HGPS) are frequently accompanied

by spontaneous NE rupture (32-34). In HGPS, the heterozygous transversion of a

nucleotide substitution at position 1824 T>C (G608G) leads to abnormal splicing of the

last 150 nucleotides of exon 11, resulting in the production of unprocessed HGPS mutant,

progerin (PG) (pre-PG) which lacks 50 amino acids (residues 607–656) from the C

terminus of pre-LA (Fig. 1 A). Because these residues contain the second proteolytic

cleavage site involved in pre-LA processing, the mature form (mature-PG) remains

farnesylated and carboxymethylated, leading to its permanent association with the INM

(35, 36).

Previous studies demonstrate that LA/C is recruited to the rupture sites with BAF

(30, 37, 38) but LA is significantly slow and weak in localizing to the rupture sites

compared to LC (30, 39, 40) because it is less abundant in the nucleoplasm than LC (30,

39, 40). However, a molecular mechanism for retaining nucleoplasmic LA sparse and

slowing its localization to the rupture sites still remains to be elucidated. Here, we

investigate the role of the C-terminal tail region specific for LA in this mechanism. We

previously showed that mEmerald-fused LA accumulated at the rupture sites only when

it was overexpressed at a high level (30), and the accumulation kinetics appeared to be

artifactual because it was not confirmed with endogenous LA detected by

immunofluorescence (30). Therefore, we establish the ectopic expression system for live-

cell imaging and fluorescence correlation spectroscopy (FCS) to express mEmerald-fused

LA, alternate forms of LA, and LC at a low level with the endogenous promotor in wild

type (WT), HGPS and LA/C-knockout (KO) MEFs. Our data reveal the significance of

the LA-specific tail region for retaining its small pool in the nucleoplasm and slowing its

localization to the rupture sites.

Results

PG is not only slow in localizing to the rupture sites but also slows LA localization

To investigate the involvement of the pre-LA processing in retaining small pool of LA in the nucleoplasm and slowing its localization to the rupture sites, we used immortalized WT MEFs and heterozygous knock-in MEFs with PG expression from a mutant allele of the *Lmna* gene carrying the c.1827C>T; p.Gly609Gly mutation (*Lmna*$^{G609G/+}$, G609G/+). This mutation is equivalent to the HGPS mutation in the human *LMNA* gene (41, 42). These cells stably expressed super-folder Cherry harboring two NLSs derived from SV40 large T antigen and c-Myc (NLS-sfCherry) as a NE rupture marker (30). Because spontaneous NE rupture is frequently observed in skin fibroblasts derived from a HGPS patient and mouse smooth muscle cells (SMCs) expressing PG (32, 34), we first examined if G609G/+ MEFs also exhibit misshapen nuclei, the leakage of NLS-sfCherry to the cytoplasm, the mis-localization or decreased expression of LB1, and the local accumulation of cGAS, which are all indicative of spontaneous NE rupture. WT and G609G/+ cells were stained with Hoechst 33342 for DNA and specific antibodies directed against LB1 and cGAS (Fig. S1 A). Unlike the HGPS patient cells and PG-expressing mouse SMCs, G609G/+ cells did not show these phenotypes compared to WT cells (Table S1), indicating that the HGPS mutation is not accompanied by spontaneous NE rupture in these cells. Next, to examine the effect of this mutation on recovery from forced NE rupture, single-cell compression by AFM indentation was applied to these cells, and the nuclear reentry of NLS-sfCherry was observed by time-lapse imaging before and after the induction of the rupture (Fig. 1 B) (43). Interestingly, G609G/+ cells showed a significantly slow reentry compared to the WT cells which had a ~50% recovery from the

original fluorescence intensity ~4 min after the induction of NE rupture (Fig. 1 C and D),

indicating that PG expression slows the recovery from forced NE rupture.

To determine the localization kinetics of LA and PG at the rupture sites within a

relatively long-time window, WT and G609G/+ cells were fixed at 0-10, 30-40, 60-70

and 120-130 min after the induction of NE rupture by 405-nm laser microirradiation (Fig.

1 E) (30), and then stained with Hoechst 33342 for DNA and the specific antibodies

directed against LA and PG which do not cross-react to each other (Fig. 1 A). At 0-10,

30-40, 60-70 and 120-130 min after the rupture, LA was localized to the rupture sites in

~4%, ~55%, ~77% and ~86% of WT nuclei and ~8%, ~29%, ~61% and ~68% of

G609G/+ nuclei, respectively (Fig. 1 F, G and Fig. S1 B), whereas PG was slower in

localizing to the rupture sites of G609G/+ nuclei than LA, showing ~0%, ~7%, ~32% and

~50%, respectively (Fig. 1 F, G and Fig. S1 B). These results suggest that PG expression

decelerates the localization kinetics of LA at the later time points (60 and 120 min). In

addition, both LA and PG are likely to be recruited to the rupture sites with BAF within

a relatively long-time window according to our results from BAF knockdown (KD)

experiments with short hairpin RNAs (shRNAs) (Fig. S1 C-E). WT cells expressing a

shRNA for the control (Scramble) or two shRNAs (shBAF #1 and #2) were fixed at 60-

70 min after the induction of NE rupture by laser microirradiation, and then LA

localization was determined by immunofluorescence. As expected, LA was localized to

the rupture sites in ~61%, 10% and ~16% of the control and BAF-KD#1 and #2 nuclei,

respectively (Fig. S1 D and E), indicating that BAF is required for LA localization to the

rupture sites. These results are similar to a previous finding to indicate that BAF rapidly

recruits overexpressed GFP-LA to the rupture sites (37).

PG slows LC accumulation to the rupture sites

Because the accumulation of LC to the rupture sites was much earlier (within 3 min) than the localization of LA or PG (30-120 min) (30, 39, 40) (Fig. 1 F and G) and LC depletion accelerates the cytoplasmic leakage of NLS-sfCherry at the early time points (within 10 min) (30), LC might be directly linked to the delay in retrieving NLS-sfCherry from the cytoplasm of G609G/+ cells within 10 min after the induction of forced NE rupture (Fig. 1 C and D). Therefore, PG expression could slow LC accumulation to the rupture sites. To test this idea, endogenous LA, LC and PG were directly labeled with a LA/C-specific genetically encoded probe, designed ankyrin repeat protein (DARPin)-LaA_6 (44), fused with super-folder GFP (sfGFP), sfGFP-DARPin-LA6 in WT and G609G/+ cells expressing NLS-sfCherry (30) (Fig. S2 A). Then, live-cell imaging was carried out before and after the induction of NE rupture by laser microirradiation. Because endogenous LA and PG did not localize to the rupture sites within 3 min after the induction of NE rupture (30) (Fig. 1 G), sfGFP-DARPin-LA6 accumulating at the rupture sites only represented endogenous LC within such a short-time window. Immediately after the induction of NE rupture, the accumulation of sfGFP-DARPin-LA6 was significantly weaker in G609G/+ nuclei compared to WT nuclei (Fig. S2 B and C), indicating that PG expression attenuates LC accumulation.

PG slows the diffusion of LC in the nucleoplasm

Because of the slow recruitment of LC from the nucleoplasm to the rupture sites in G609G/+ nuclei (Fig. S2 B and C), the diffusion fractions and coefficients in the nucleoplasm could be decreased in G609G/+ nuclei compared to WT nuclei. To determine them, mEmerald-LC was stably expressed at a low level from the *Rosa26* locus under the

endogenous *Lmna* promotor in WT and G609G/+ cells, and fluorescence correlation

spectroscopy (FCS) was performed using these cells (Fig. S3 A-D). The auto-correlation

curve obtained by the signal detection was fitted with fast and slow components (15, 45,

46). Interestingly, the diffusion coefficients of fast components were significantly high in

WT nuclei compared to G609G/+ nuclei (Fig. S3 D). This result indicates that PG

expression significantly decreases the diffusion coefficient of fast component despite the

increased fraction but this degree of difference is insufficient for explaining the weak

accumulation of sfGFP-DARPin-LA6 at the rupture sites in G609G/+ nuclei compared

to WT nuclei.

The nucleoplasmic pools of LA and PG are increased by the inhibition of

farnesylation

When the CaaX motifs of pre-LA and pre-PG are farnesylated (18, 36, 47), they are

retained at the NE through the hydrophobic interaction with the lipid bilayer of the INM

(47, 48). Therefore, the inhibition of farnesylation with FTI could increase their

nucleoplasmic pools, leading to their fast localization to the rupture sites. To test this

possibility, WT and G609G/+ cells were treated with DMSO for the control or

farnesyltransferase inhibitor (FTI), 3.2 μM lonafarnib for 48 h, and the expression levels

and localization of LA and PG were determined by immunoblotting using the specific

antibodies directed against LA/C, and by immunofluorescence using the LA and PG

antibodies respectively, (Fig. 2 A). For immunoblotting, pre-LA, mature-LA, pre-PG,

mature-PG, and LC were detected with the antibodies directed against LA/C (for all) and

PG (for pre- and mature-PG). For immunofluorescence, the LA antibody (for pre- and

mature-LA) and the PG antibody (for pre- and mature-PG) were used. As expected, the

levels of pre-LA and pre-PG were increased in FTI-treated cells compared to the control

cells as shown as the upper and lower band of LA and PG (Fig. 2 A). Based on ratios of

the average fluorescence intensity in the nucleoplasmic pool relative to that of the NL

(NP/NL ratios), LA and PG were significantly abundant in the nucleoplasm of FTI-treated

nuclei compared to the control nuclei (Fig. 2 B and C), indicating that LA and PG are

increased in the nucleoplasmic pool by FTI treatment. Next, these cells were fixed within

10 min and at 60-70 min after the induction of NE rupture by laser microirradiation, and

the localization of LA and PG to the rupture sites was detected by immunofluorescence.

LA was localized to the rupture sites within 10 min in ~7% and ~7% of WT and G609G/+

nuclei for the control and ~18% and ~17% of WT and G609G/+ nuclei treated with FTI,

respectively, and PG in 0% and ~37% of the control and FTI-treated G609G/+ nuclei,

respectively (Fig. S4). At 60-70 min, LA localization to the rupture sites was observed in

~59% and ~42% of the control nuclei and ~91% and ~74% of FTI-treated WT and

G609G/+ nuclei, respectively, and PG localization in ~18% and ~81% of the control and

FTI-treated G609G/+ nuclei, respectively (Fig. 2 D). These results indicate that the

inhibition of farnesylation promotes the localization of LA and PG to the rupture sites.

PG becomes diffusible in the nucleoplasm by FTI treatment

Because of the increases in nucleoplasmic pool of LA and PG by FTI treatment

(Fig. 2 B and C), the non-farnesylated forms could be highly mobile in the nucleoplasm

of FTI-treated cells. To test this idea, the diffusion coefficients of mEmerald-LA and PG

were measured by FCS (Fig. 3 A). For the FCS measurements, mEmerald-LA and PG

were stably expressed at a low level by the same system as described above with

mEmerald-LC (Fig. S3 A and B), followed by the treatment with DMSO for the control

or FTI (Fig. S4). The diffusion fractions of mEmerald-LA in the control and FTI-treated

WT and G609G/+ nuclei were all similar to each other (Fig. 3 B and C) but the diffusion

coefficient of fast component was slightly but significantly small in G609G/+ nuclei

compared to WT nuclei whereas that of slow component remained unchanged (Fig. 3 C).

These results show that PG expression slows the diffusion speed of mEmerald-LA. On

the other hand, the diffusion signals of mEmerald-PG were barely detectable in the

nucleoplasm of WT and G609G/+ nuclei for the control. Instead, the signal decays caused

by photobleaching were observed during the FCS measurements (Fig. 3 D), strongly

suggesting that these signals originate from the NL where mEmerald-PG are immobile

(15, 53). When WT and G609G/+ cells were treated with FTI, the diffusion signals were

detected by FCS as expected (Fig. 3 D and E), indicating that PG becomes diffusible in

the nucleoplasm by FTI treatment. Therefore, the nucleoplasmic pool is the likely source

of PG localized to the rupture sites (Fig. 2 C and D).

274　The second proteolytic cleavage is required for the localization of LA to the rupture

275　sites

The localization of PG to the rupture sites was significantly slow compared to LA (Fig. 1

G), and the permanent association of PG with the INM due to the absence of the second

proteolytic cleavage site could decelerate the localization kinetics (35, 36). To check this

possibility, Zmpste24, which endoproteolytically processes both the CaaX motif and the

second cleavage site (54) was knocked down by shRNAs in WT MEFs, and then the

expression levels and localization of pre- and mature-LA were determined in cells

expressing shRNAs for the control or Zmpste24 KD by immunoblotting and

immunofluorescence, respectively (Fig. S5 A). A specific antibody directed against pre-

LA used for immunoblotting and immunofluorescence reacts with pre-LA but not mature-

LA (Fig. 1 A). The C-terminal residues of pre-LA including the CaaX motif remained

partially uncleaved by Zmpste24 KD, resulting in the significant increase and decrease in

expression level of pre- and mature-LA, as shown as the upper and lower band,

respectively (Fig. S5 A). These results are consistent with the previous findings to indicate

that Zmpste24 KD inhibits the second cleavage (55). LA and pre-LA were predominantly

localized to the NL, and the NP/NL ratios remained unchanged in Zmpste24-KD cells

compared to the control cells (Fig. S5 B and C). These cells were fixed at 60-70 min after

the induction of NE rupture by laser microirradiation, and then the localization of LA and

pre-LA to the rupture sites was determined by immunofluorescence. LA was localized at

the rupture sites in ~75% of the control cells whereas LA and pre-LA were completely

absent from the rupture sites in Zmpste24-KD cells (Fig. S5 D and E), indicating that the

second cleavage is required for their localization to the rupture sites.

Inhibition of farnesylation at the CaaX motif is sufficient for the localization of LA

to the rupture sites without the second proteolytic cleavage

Zmpste24 depletion significantly increased the pre-LA expression level (56) (Fig. S5 A

and B), which is accompanied by the farnesylation at the CaaX motif (49). Therefore, the

inhibition of the farnesylation with FTI could release the pre-LA from the NE, resulting

in the fast localization of pre-LA to the rupture sites. To test this idea, the control

(Scramble) and Zmpste24-KD cells were treated with DMSO or FTI as before, and the

expression levels and localization of pre- and mature-LA were determined by

immunoblotting and immunofluorescence, respectively (Fig. 4 A and B). As expected,

Zmpste24 KD increased and decreased the pre- and mature-LA levels, respectively in

DMSO- and FTI-treated cells (Fig. 4 A). FTI treatment also had the similar effect in the

control cells but the synergetic effect with Zmpste24 KD was very subtle (Fig. 4 A). To

analyze the localization of pre- and mature-LA in those cells by immunofluorescence,

specific antibodies directed against them that do not cross-react with each other were used.

In the control cells treated with DMSO, pre-LA was under the detection limit whereas

mature-LA was predominantly localized to the NL (Fig. 4 B-D). When the control cells

were treated with FTI, pre-LA became clearly visible in the NL and mature-LA was less

prominent there instead (Fig. 4 B-D). In Zmpste24-KD cells treated with DMSO, pre-LA

was localized to the NL and mature-LA was undetectable (Fig. 4 B-D). FTI treatment

decreased pre-LA localization to the NL up to the level of the control cells treated with

FTI and mature-LA was undetectable (Fig. 4 B-D). These results were consistent with the

known effect of FTI treatment, which inhibit the farnesylation at the CaaX motif and the

following processes including the second cleavage, leading to release the pre-LA from

the NE (50). As FTI treatment increases nucleoplasmic LA levels regardless of the second

cleavage, we hypothesize that LA localizes to the rupture site more efficiently in FTI. The

localization of pre- and mature-LA to the rupture sites was determined in these cells by

immunofluorescence at 60-70 min after the induction of NE rupture by laser

microirradiation. Pre-LA was localized to the rupture sites in ~97%, ~12% and ~77% of

FTI-treated control, DMSO- and FTI-treated Zmpste24-KD nuclei, respectively (Fig. 4

E). This indicates that pre-LA without farnesylation present in the nucleoplasm

effectively localizes to the rupture sites, supporting the hypothesis. However, the

localization of mature-LA was slightly decreased in the control cells by FTI treatment

(Fig. 4 E), despite FTI treatment increases the localization of pre- and mature-LA using

antibody directed against both (Fig. 2 D), probably because its nucleoplasmic level was

not as high as pre-LA (Fig. 2 C, 4 C and D).

LA differs from LC in localization to the rupture sites through two specific

sequences in the tail region

The inhibition of farnesylation at the CaaX motif of PG significantly increased the

nucleoplasmic pool and promoted the localization to the rupture sites within 10 min after

the induction of NE rupture (Fig. S4). Interestingly, PG was not localized to the rupture

sites within the first half of the time window (0-5 min) whereas LC rapidly accumulated

(30) (Fig. S2). Therefore, we confirmed the effect of this inhibition on their NP/NL ratios

and the accumulation kinetics at the rupture sites within the short-time window (3 min)

by live-cell imaging using mEmerald-fused LA-full length (FL), LC, PG and the non-

farnesylated mutant of PG, PG-CSM that lacks the isoleucine within the CaaX motif were

expressed in LA/C-KO MEFs (Fig. 5 A). As expected, the NP/NL ratio of mEmerald-PG

was similar to that of mEmerald-LA-FL, whereas that of mEmerald-PG-CSM was

significantly high but not to the extent of mEmerald-LC (Fig. 5 B and C). mEmerald-PG

and mEmerald-PG-CSM did not accumulate at the rupture sites (Fig. 5 D), indicating that

the inhibition of the farnesylation is not sufficient for exhibiting the rapid accumulation

that was similar to that of mEmerald-LC. These results strongly suggest that the tail

regions specific for LA and PG are involved in inhibiting the rapid accumulation. In

human and mice, LC harbors identical six amino acids whereas LA has 98 and 97 amino

acids, following the residues 1-566 and 1-568 shared between LA and LC, respectively

(Fig. S6 A). Because this LC-specific amino acids do not contribute to the rapid

accumulation at the rupture sites (30), mEmerald-fusion proteins of LA truncation

mutants, LA-S575X, LA-A601X, LA-S617X, LA-F627X, LA-T644X and LA-L648X

were tested for their NP/NL ratios and the accumulation kinetics within 3 min (Fig. 5 A).

Among these mutants, mEmerald-LA-S575X has the highest NP/NL ratio (Fig. 5 B and

C), and only this mutant rapidly accumulated within 3 min (Fig. 5 D and S6 B), indicating

that the residues 575-600 is involved in inhibiting the rapid accumulation. Because the

deletion mutant, mEmerald-LAΔ26 did not accumulate at the rupture sites (Fig. 5 D),

further truncation was performed at the C-terminal side of mEmerald-LAΔ26 to produce

mEmerald-LAΔ26-L627X, mEmerald-LAΔ26-L644X and mEmerald-LAΔ26-L648X

(Fig. 5 A). The NP/NL ratio remained unchanged by these truncations compared to

mEmerald-LAΔ26 (Fig. S6 C and D) but only mEmerald-LAΔ26-L627X rapidly

accumulated (Fig. 5 E and S6 E). These results show that the residues 627-643 assists

LAΔ26 to inhibit the rapid accumulation without decreasing the nucleoplasmic pool. We

therefore designated the residues 575-600 as LA-characteristic sequence-1 (LACS1) and

the residues 627-643 as LACS2.

Discussion

A-type lamins are localized to sites of NE rupture for the repair (23, 24, 30). Our previous study shows that endogenous LC but not LA rapidly accumulates the rupture sites within 3 min after the induction of NE rupture because LC is more abundant in the nucleoplasm than LA (30, 39, 40). LA can rapidly accumulate only when LA is overexpressed at a high level to increase the nucleoplasmic pool (30, 37, 38). LC accumulation requires both the immunoglobulin-like fold (Ig-fold) domain that binds to BAF and a nuclear localization signal (NLS), and is not attributable to the LC-specific amino acids (30). In this study, we analyze the LA-specific tail region to determine a molecular mechanism for retaining its small pool in the nucleoplasm and decelerating localization kinetics at the rupture sites, and Fig. 6 illustrates our current findings following our previous study (30). The proteolytic removal of 15 amino acids containing the farnesylated C-terminal end of pre-LA by Zmpste24 increases the nucleoplasmic pool and accelerates the localization kinetics at the rupture sites within the long-time window (60-70 min) after the induction of NE rupture. In HGPS, pre-PG lacks 50 amino acids containing the second proteolytic cleavage site from the C terminus of pre-LA, leading to the permanent association with the INM, which significantly decreases the nucleoplasmic pool and decelerates the localization kinetics at the rupture sites. In contrast, FTI treatment pronouncedly increases the nucleoplasmic pools of LA and PG and promotes their localization to the rupture sites. Therefore, the processes for increasing the nucleoplasmic pool is likely to be rate-limiting steps for the localization kinetics within this time window. Additionally, LACS1 cooperates with LACS2 to inhibit the rapid accumulation at the rupture sites within the short-time window (3 min).

We use immortalized WT and G609G/+ MEFs to analyze the NP/NL ratios of

LA, pre-LA and PG, and their localization kinetics at the rupture sites. Unlike HGPS-

patient cells and mouse SMCs expressing PG (32, 34, 58), our G609G/+ cells do not

exhibit the formation of misshapen nuclei and spontaneous NE rupture, which could be

due to the immortalization process with SV40 large T antigen. In oncogene-induced

senescence of normal human fibroblasts, DNA protrusion from the nucleus, micronuclei

formation and the leakage of nuclear contents are coupled to the loss of LB1 (59-61). The

LB1 expression level could also decrease when HGPS-patient cells become prematurely

senescent (62) and PG expression levels increase with age in aortas from G609G/+ mice

whereas LB1 levels decrease (34). Therefore, spontaneous NE rupture might be induced

by the HGPS mutation because of the induction of premature senescence and rapid

cellular aging.

Instead of spontaneous NE rupture, we find that upon the induction of forced NE

rupture by AFM indentation, the nuclear reentry of NLS-sfCherry is significantly slow in

G609G/+ cells compared to WT cells, strongly suggesting that a repair process of forced

NE rupture is impaired by PG expression. The LA expression level in G609G/+ cells is

reduced compared to that in WT cells due to the efficient alternative splicing from the

mutant allele of the *Lmna* gene (41, 63), which could explain its smaller pool in the

nucleoplasm compared to WT cells to decelerate the localization kinetics at the rupture

sites. On the other hand, the LC expression level could be increased in G609G/+ cells

compared to WT cells for some reason (40, 41). However, the diffusion of the fast

component is significantly slow in the nucleoplasm of G609G/+ nuclei compared to WT

nuclei, as our results indicate. This could be explained if there is a possibility that LC is

increased in the NL and decreased in the nucleoplasm because of the strong association

with the NL by PG expression. To support this idea, LC has a significantly slow turnover

in the NL of HGPS-patient nuclei compared to normal nuclei (53, 64). These results

suggest that the alteration in expression and distribution of LA/C by PG expression impair

the repair of NE rupture.

We find that when the farnesylation at the CaaX motif of pre-LA and pre-PG is

inhibited by FTI treatment, they are increased and become diffusible in the nucleoplasm

to accelerate the localization kinetics to the rupture sites, indicating that the farnesylation

process limits the localization to the rupture sites regardless of the second cleavage of

pre-LA. Furthermore, Zmpste24 KD blocks the second cleavage, leading to the complete

inhibition of pre-LA localization to the rupture sites, and the effects of Zmpste24 KD on

the NP/NL ratios and the localization kinetics are rescued by FTI treatment. These results

strongly suggest that pre-LA is mostly farnesylated in Zmpste24 KD cells and that the

nucleoplasmic pool is little or none to localize to the rupture sites within 60 min.

Apart from the farnesylation at the CaaX motif and the inhibition of the second

cleavage of pre-LA, we find that LACS1 and LACS2 in the tail region are also involved

in slowing the localization to the rupture sites. Because the truncation of LACS1

significantly increases the NP/NL ratios, LA appears to be retained at the NL through

LACS1 compared to LC, resulting in the slow localization. It is possible that LACS1

contributes to the formation of lamin filaments in the NL and/or interacting with other

NE components including INM proteins. On the other hand, LACS2 could assist the

LACS1 function for inhibiting LA localization to the rupture sites without promoting the

retention of LA at the NL. Because the truncation of the residues 599-625 between

LACS1 and LACS2 in human increases the NP/NL ratios, a segment within these amino

acids is likely to retain its small pool in the nucleoplasm. Whereas the residues 411-553

including the NLS and the Ig-fold domain interacts with dsDNAs (67), the residues 506-

638 binds histone H3 (68). This chromatin-binding domain contains the LACSs,

suggesting that LA is partially tethered to chromatin through LACSs to slow the

localization to the rupture sites.

In HGPS, the permanent association of PG with the INM by the farnesylation at

the CaaX motif significantly reduces the nucleoplasmic pool, leading to the slow

localization to the rupture sites. Similarly, this mechanism could be also critical for their

accesses to active enhancers in the nuclear interior to affect the gene expression in HGPS

patient cells (69). Other studies show that the PG association with the INM can lead to

the increased NE instability (32, 34) but we do not observe spontaneous NE rupture in

our G609G/+ cells. Nevertheless, our findings of the slow localization kinetics of LA, LC

and PG at the rupture sites implicate pathological significance of the defective repair of

the NL. The therapeutic effect of FTI treatment appears to rescue from the repair defect

by targeting the farnesylation of PG. Therefore, the induction of spontaneous NE rupture

and the rescuing with FTI treatment need validating using the mouse model of HGPS in

future studies.

Materials and methods

Plasmid construction

Two different NLSs derived from SV40 large T antigen (NLS^{SV40}; PKKKRKV) and c-Myc (NLS^{Myc}; PAAKRVKLD) from pCDH-NLS^{SV40}-sfCherry-NLS^{Myc}-Blast (30) were ligated to pCDH-CMV-MCS-EF1-Hygro (plasmid #CD515B-1, System Biosciences) using the Ligation-Convenience Kit (Nippon Gene) and designated as pCDH-NLS^{SV40}-sfCherry-NLS^{Myc}-Hygro. The annealed oligonucleotides to express shRNAs (Zmpste24, TRCN0000366641; BAF#1, TRCN0000124958; BAF#2, TRCN0000124955; Table S2, Broad Institute) were ligated into pLKO.3-blast (30) using the Ligation-Convenience Kit. The pX459-sfCherry CRISPR-Cas9 vector were generated from pSpCas9(BB)-2A-Puro (pX459) V2.0 (Addgene plasmid # 62988 ; http://n2t.net/addgene:62988 ; RRID:Addgene_62988; a gift from Feng Zhang). The annealed oligonucleotides to express single-guide RNA (sgRNA) against the *Rosa26* target sequence (70) was ligated into pX459-sfCherry using the Ligation-Convenience Kit and designated as pX459-sfCherry-sgRosa26-1.

We cloned a 1.6 kb genomic region upstream from start codon of mouse *Lmna* as the endogenous promoter (-1407 to +249 bp from transcription start site) using PrimeSTAR HS DNA Polymerase (Takara). Primers used in this study are listed in Table S2. Mouse genomic DNA was extracted from cells by Proteinase K (Fungal; Invitrogen) digestion according to the standard protocol and used for a PCR template. The CMV promoter of mEmerald-C1 (Addgene plasmid # 53975 ; http://n2t.net/addgene:53975 ; RRID:Addgene_53975; a gift from Michael Davidson) was replaced by the *Lmna* promoter using the In-Fusion HD Cloning Kit (Clontech) and designated as pLmna-mEmerald-C1. The mouse pre-LA cDNA was amplified from pPyCAG-LA-IP (71) using the KOD One PCR Master Mix -Blue- (Toyobo), cloned into pLmna-mEmerald-C1 using the In-Fusion, and designated as pLmna-Em-LA. The mouse PG cDNA was generated by PrimeSTAR Max DNA Polymerase (Takara) to skip 150 bp, closed by In-Fusion HD, and designated as pLmna-Em-PG. The amino acid truncation mutants were generated by PCR mutagenesis using PrimeSTAR Max or PrimeSTAR HS DNA Polymerase. The LACS1 deletion mutant (Δ26) was generated using PrimeSTAR Max to skip 78 bp, and closed by In-Fusion HD. The pLmna-Em-LA and pLmna-Em-PG expression cassettes were amplified by the KOD One, fused with bGHpA amplified from LSL-Cas9-Rosa26TV

488   (Addgene plasmid # 61408 ; http://n2t.net/addgene:61408 ; RRID:Addgene_61408; a gift

489   from Feng Zhang) and *Pac* I/*Sma* I digested LSL-Cas9-Rosa26TV by In-Fusion HD, and

490   designated as pLmna-Em-LA-R26Neo, pLmna-Em-PG-R26Neo and pLmna-Em-PG-

491   CSM-R26Neo. The mouse LC cDNA was also generated by KOD One PCR and In-

492   Fusion HD and designated as pLmna-Em-LC-R26Neo.

Cell culture

MEFs isolated from WT and heterozygous c.1827C>T mutant (*Lmna*$^{G609G/+}$, G609G/+)

embryos at embryonic day 13.5 were generously gifted by Carlos López-Otín

(Universidad de Oviedo) (41, 42). Primary MEFs were cultured in modified DMEM (high

glucose; Nacalai Tesque) containing 10% FBS (qualified; Thermo Fisher), 4 mM L-

glutamine, 100 U/mL penicillin and 100 μg/mL streptomycin (Sigma-Aldrich), and 20

mM 2-[4-(2-Hydroxyethyl)-1-piperazinyl] ethanesulfonic acid (HEPES; Nacalai Tesque)

at 37°C in a humidified multi-gas incubator with 3% O_2/5% CO_2 (MCO-5M,

Sanyo/Panasonic Healthcare). Cells were immortalized by transduction with a retrovirus

expressing SV40 large T antigen from pBABE-puro largeTcDNA (Addgene plasmid #

14088 ; http://n2t.net/addgene:14088 ; RRID:Addgene_14088; a gift from William Hahn)

as previously described (13). After selection with 2 μg/ml puromycin (InvivoGen) for 4

d, immortalized WT and G609G/+ cells were cultured in growth medium without HEPES

at 37°C in a humidified chamber with 5% CO_2.

Lentiviral transduction

For lentivirus-mediated stable introduction of NLS^{SV40}-sfCherry-NLS^{Myc}, we followed

the methods that was previously described (30). Briefly, pVSV-G (PT3343-5, Clontech)

and psPAX2 (Addgene plasmid #12260; http://n2t.net/addgene:12260;

RRID:Addgene_12260; a gift from Didier Trono), together with a pCDH vector (pCDH-

NLS^{SV40}-sfCherry-NLS^{Myc}-Blast, pCDH-NLS^{SV40}-sfCherry-NLS^{Myc}-Hygro or pCDH-

sfGFP-DARPin-LA6-Hygro) in a 1:3:4 weight ratio of each plasmid was transfected into

~80% confluent 293T cells (CRL-3216, ATCC) using Lipofectamine 3000 following the

manufacturer's instructions for lentivirus production. 24 h after the transfection, the

medium was replaced with fresh medium, and the virus-containing culture supernatant

was harvested at 48 h after transfection and filtrated with a 0.45-μm syringe filter. For

virus infection, MEFs were incubated in the culture supernatants with 4 μg/mL polybrene

(Nacalai Tesque) for 24 h. Infected cells were selected by incubation in medium

containing 3 μg/mL blasticidin S or 200 μg/mL hygromycin B Gold (InvivoGen) for 4 d.

524 Live-cell imaging and NE rupture induction by laser microirradiation

Culture medium was replaced with FluoroBrite DMEM (Thermo Fisher) containing 10%

FBS, 4 mM L-glutamine, 100U/mL penicillin and 100 μg/mL streptomycin. For laser-

microirradiation and image acquisition, cells were set onto a heated stage (37°C; Tokai

Hit) with a CO_2-control system (Tokken) on a confocal microscope (FV1000, Olympus)

with a 60× PlanApo N (NA 1.4) oil lens operated by built-in software FLUOVIEW ver.

4.2 (Olympus). All live-cell images were acquired using a main scanner (4% 488-nm laser

transmission; 30% 543-nm laser transmission; 2 μs/pixel; 512×512 pixels; pinhole 100

μm; 6× zoom; ~10 s/frame). After the first image was acquired, a 2 μm diameter spot was

laser-microirradiated using a second scanner at 100% power of 405-nm laser transmission

(40 μs/pixel; approximately 780 μW) for 10 s whereas the images were acquired with

another scanner.

Fiji software (ImageJ) ver. 1.53v (National Institute of Health) was used for

measuring max intensity at the sites of NE rupture after Gaussian filtering with σ = 2.0

over time. Fluorescence intensity was normalized by each of the initial intensity in the

laser-microirradiated region.

CellProfiler ver. 3.1.9 (Broad Institute) with the NP_NL_intensity_live pipeline

(uploaded on GitHub; http://github.com/Kimura-Lab/Kono-et-al.-2023) (30) were used

for measuring mean intensity of mEmerald-LA mutants and mEmerald-LC in the NP and

the NL. Briefly, after the nuclear region was recognized by the nuclear localization of

NLS-sfCherry, 3 pixels (207 nm) and more than 10 pixels (690 nm) inside from the rim

of nuclear region were regarded as the NL and the NP, respectively, and then the mean

intensity of each section was measured. All images were processed in Photoshop ver. 24

(Adobe) for cropping and brightness/contrast adjustment.

549 Indirect immunofluorescence and microscopy

Primary antibodies used for immunofluorescence were mouse monoclonal anti-LA

(1:250; 4A58, sc-71481, Santa Cruz), rat monoclonal anti-pre-LA (1:1,000; 7G11;

MABT345, EMD Millipore; (49)), mouse monoclonal anti-mature-LA (1:8,000; 4A4-A4,

14-9848-82, Invitrogen; (72)), rabbit polyclonal anti-progerin (1:1000; generated by the

Nourshargh laboratory using a peptide immunogen of murine progerin and standard

immunization procedures; (63, 73, 74)), mouse monoclonal anti-LB1 (1:1,000; B-10; sc-

374015, Santa Cruz), rabbit monoclonal anti-cGAS (1:250; D3O8O; 21659, Cell

Signaling Technology) and rabbit monoclonal anti-BANF1/BAF (1:200; EPR7668;

ab129184, Abcam). The secondary antibodies used were Alexa Fluor 488-donkey anti-

mouse immunoglobulin G (IgG), Alexa Fluor 488-donkey anti-rabbit IgG (1:2,000;

A21202 and 1:2,000; A21206, respectively, Thermo Fisher), Cy5-donkey anti-mouse IgG

and Cy5-goat anti-rat IgG (1:2,000; 715-175-151 and 1:2,000; 712-175-153, respectively,

Jackson ImmunoResearch).

Cells were grown on 35-mm Glass Based Dishes which have a cover glass with

a grid pattern (IWAKI) and fixed after laser-microirradiation with 4% PFA (Electron

Microscopy Sciences) containing 0.1% Triton X-100 (Nacalai Tesque) for 15 min,

followed by permeabilization using 0.1% Triton X-100 in PBS for 10 min and then

blocking with Blocking One-P (Nacalai Tesque) for 30 min. Cells were incubated with

primary antibodies overnight at 4°C, washed with 10% Blocking One-P in PBS, and

incubated with secondary antibodies for 1 h at RT. DNA was stained with Hoechst 33342

(0.5 μg/mL; Thermo Fisher). For Fig. S1 A, cells were fixed with 4% PFA for 15 min,

followed by permeabilization using 0.1% Triton X-100 in PBS for 10 min and then

blocking with Blocking One-P for 30 min. After incubation with primary antibodies, the

RFP-Booster Alexa Fluor 568 (alpaca monoclonal anti-RFP Nanobody; 1:1000;

rb2AF568, ChromoTek) was used with second antibodies to amplify the signal of NLS-

sfCherry.

Confocal fluorescence images were obtained using a confocal microscope

FV1000 with a 60× PlanApo N (NA 1.4) oil lens (3.0% 405-nm laser transmission; 3.1-

11.1% 488-nm laser transmission; 5.0% 543-nm laser transmission; 1.0-4.0% 633-nm

laser transmission; 2 μs/pixel; 512×512 pixels; pinhole 100 μm). CellProfiler with the

NP_NL_intensity_fixed pipeline (uploaded on GitHub; http://github.com/Kimura-

Lab/Kono-et-al.-2023) were used for measuring mean intensity of lamins in the NP and

the NL. The nuclear region was recognized by staining DNA with Hoechst 33342.

Induction of mechanical NE rupture by atomic force microscopy (AFM)

Culture medium was replaced with Leibovitz's L-15 medium (no phenol red; Thermo

Fisher) containing 10% FBS, 2 mM L-glutamine, 100U/mL penicillin and 100 μg/mL

streptomycin. JPK NanoWizard 4 BioAFM (Bruker) incorporated into an inverted

fluorescence microscope (Eclipse Ti2, Nikon) with EM-CCD camera (iXon Ultra 888,

Andor) and light source (X-Cite XYLIS, Excelitas) was used for single-cell compression.

Cantilever (TL-CONT, Nanosensors; spring constant approximately 0.2 N/m) with a 100-

μm glass bead (BZ-01, As-One) was used for NE rupture. Cells expressing NLS-sfCherry

were set onto a heated stage (37°C) and the first fluorescent image of NLS-sfCherry was

obtained before the rupture using a TRITC filter and NIS-Elements BR (Nikon). Next,

the bead was approached on the top of the cell surface with 2 nN setpoint and then forced

down for 5 μm. After waiting for 10 s, the cantilever was retracted. Within 10 s after the

retraction, time-lapse imaging was started with 30 s exposure time every 30 s for 5 min.

Fiji software (ImageJ) was used for measuring mean intensities in the region of interest

from nucleoplasm and cytoplasm.

Chemicals

601 As a selective farnesyltransferase inhibitor (FTI), lonafarnib (SCH66336; Merck) was

602 dissolved in dimethylsulfoxide (DMSO; Nacalai Tesque) and added to the cells with one

603 daily dose of 3.2 μM for 2 days. For controls in all experiments, an equal volume of

604 vehicle (0.1% DMSO) was added.

Immunoblotting

For immunoblotting, we followed the methods that were previously described (30).

Briefly, primary antibodies used for immunoblotting were mouse monoclonal anti-LA/C

(1:5,000-20,000; 3A6-4C11, 14-9847-82, eBioscience), rat monoclonal anti-pre-LA

(1:1,000; 7G11), rabbit polyclonal anti-progerin (1:1,000; Nourshargh lab), rabbit

monoclonal anti-BANF1/BAF (1:500; EPR7668, Abcam), anti-β-Actin (1:2,000; AC-15,

sc-69879, Santa Cruz) and anti-GAPDH (1:5,000; 6C5, sc-32233, Santa Cruz). The

secondary antibodies used were anti-mouse IgG HRP-linked whole Ab sheep (1:10,000;

NA931, Amersham), anti-rabbit IgG HRP-linked F(ab')$_2$ fragment donkey (1:10,000;

NA9340, Amersham) and HRP-donkey anti-rat IgG (H+L) (1:10,000; 712-035-150,

Jackson ImmunoResearch).

Rosa26-targeted gene knock-in

The pLmna-Em-preLA-R26Neo, pLmna-Em-PG-R26Neo and pLmna-Em-LC-R26Neo

were linearized by the *Kpn* I (Takara) and purified by phenol/chloroform extraction and

ethanol precipitation. Cells were transfected with appropriate plasmids using

Lipofectamine 3000 (Thermo Fisher) by reverse transfection as previously described (30).

Briefly, the DNA-lipid complex (3 μg linearized *Rosa26* targeting vectors : 2 μg pX459-

sfCherry-sgRosa26-1 : 7.5 μL Lipofectamine 3000 in 250 μL Opti-MEM; Thermo Fisher)

was added to 1.62×10^5 cells as they were seeding onto a 35 mm dish with 2 mL of growth

medium. One day after the transfection, the medium was replaced with fresh medium.

Two days after the transfection, cells were selected by incubation in medium containing

400 μg/mL G-418 disulfate aqueous solution (Nacalai Tesque) for 7 d.

Fluorescence Correlation Spectroscopy (FCS)

FCS measurements were all performed at 25°C with an LSM780 confocal microscope

(Carl Zeiss) as previously described (15, 46, 75). FCS setups using the LSM780

microscope consisted of a continuous-wave Ar$^+$ laser (25 mW), a water-immersion

objective (C-Apochromat, ×40/1.2 NA; Carl Zeiss), and a GaAsP multichannel spectral

detector (Quasar; Carl Zeiss). Fluorescent protein of mEmerald was excited with the 488-

nm laser line with a minimal laser power (0.5%) to allow an optimal signal-to-noise ratio.

Diameter of pinhole was adjusted to 34 μm (1 Airy unit) for the 488-nm laser. Emission

fluorescence was detected at 500–550 nm in the green channel. The structure parameter

and detection volume were calibrated using Rhodamine-6G (Sigma-Aldrich) solution

with diffusion coefficient (280 μm^2/s; 25°C in water) before experiments with cells. For

FCS measurement, three positions were randomly selected inside the nucleoplasm, and

molecular diffusion was measured for 10 s five times. The first round of the five

repetitions were carried out to photobleach immobile fraction, and then the four

repetitions were used for further analysis. FCS data were analyzed with analytical

software installed in the ZEN 2012 SP5 FP1 (black) acquisition software (Carl Zeiss) (46,

75). Briefly, all measured fluorescence auto-correlation functions (FAFs) from live cells

were globally fitted by the ConfoCor3 software installed on LSM780 system using fitting

models considering three-dimensional free diffusion with a Gaussian laser profile,

including a triplet component to estimate the diffusion coefficient.

Generation of the Zmpste24- and BAF-KD MEFs

MEFs expressing shRNAs of anti-Zmpste24 and anti-BAF were generated through

lentiviral transduction. KD by shRNA expression was carried out by Lentivirus

transduction (see the section on Lentiviral transduction above) and 3 d selection with 3

µg/mL blasticidin S Gold for pLKO.3-blast.

Statistical analyses

Unless mentioned otherwise, all plots showed mean values ± SEM (error bars). Fisher's

exact test was performed to analyze the categorical data using EZR on R commander ver.

1.61 (programmed by Yoshinobu Kanda) (76). The mixed effect model fit by restricted

maximum likelihood (REML) was performed to analyze the interaction between group

and time using the *lme4* (77) and the *lmerTest* (78) packages in EZR, where cell's id was

set as a random effect, while group and time as fixed effects. The interaction between

group and time (group × time) was also set as a fixed effect. The Welch's t-test or the

Welch's one-way ANOVA followed by the Games–Howell post-hoc test was performed

to single or multiple comparison, respectively. Statistic codes for R ver. 4.2.2 and a

modified R commander for EZR 1.61 are uploaded on GitHub

(http://github.com/Kimura-Lab/Kono-et-al.-2023). Significance was determined if $P <$

0.05.

Supplemental material

Table S1 shows the phenotype characterization in WT and G609G/+ cells by immunofluorescence. Table S2 shows oligonucleotide sequences used in this study. Fig. S1 shows the immunofluorescence of WT and G609G/+ cells for phenotype characterization, and the localization of LA and PG to the rupture sites within 10 min, and at 30-40 and 120-130 min after laser microirradiation. Fig. S2 shows the protein architecture of A-type lamins and DARPin-LA6, and localization kinetics of DARPin-LA6 at the rupture sites in WT and G609G/+ cells. Fig. S3 shows WT and G609G/+ cells expressing A-type lamins under the *Lmna* promoter knocked-in to *Rosa26* locus and FCS analysis for mEmerald-LC. Fig. S4 shows the localization of LA and PG to the rupture sites within 10 min in FTI-treated WT and G609G/+ nuclei. Fig. S5 shows the immunoblotting of FTI-treated WT and G609G/+ cells expressing A-type lamins under the *Lmna* promoter knocked-in to *Rosa26* locus, and the representative image of FCS analysis. Fig. S6 shows immunoblotting, immunofluorescence and the localization of LA and pre-LA to the rupture sites in Zmpste24-KD cells. Fig. S7 shows the differences between LA and LC amino acid sequence, and dynamics of LA mutants at the rupture sites, and the NP/NL ratios of mEmerald-LA mutants by live-cell imaging.

Acknowledgments

We would like to thank Prof. Carlos López-Otín (Universidad de Oviedo) for providing us with the primary MEFs isolated from WT and *Lmna* c.1827C>T mutant (*Lmna*$^{G609G/+}$ and *Lmna*$^{G609G/G609G}$) embryos. We thank Biomaterials Analysis Division, Open Facility Center (OFC) of Tokyo Institute of Technology for nucleotide sequencing and sharing research equipment, especially Carl Zeiss ElyraS1/Airyscan/LSM780 is shared in MEXT Project Grant Number JPMXS0440200022. We also thank the Cellular Imaging Core Facility at the ConveRgence mEDIcine research cenTer (CREDIT), Asan Medical Center for technical support. The generation and validation of the anti-murine progerin antibody was funded by the Wellcome Trust (098291/Z/12/Z to S. Nourshargh) and the British Heart Foundation (FS/IBSRF/22/25121 to L. Rolas). We also acknowledge the R statistic code generated by Shinsuke Hatano (Ryukoku University). We also thank all members of the Kimura lab at Tokyo Institute of Technology and Tokuko Haraguchi (Osaka University) for advice and discussions. This work was supported by JSPS KAKENHI Grant Numbers JP20KK0158 (to T. Shimi, Y. Kono, K. Miyazawa and H. Kimura), JP20K06617 (to T. Shimi), and JP18H05527 and JP21H04764 (to H. Kimura). This work was also supported by MEXT (World Premier International Research Center Initiative [WPI]). Y. Kono was an academy scholar at the Specialized Academy for Cell Science (Ohsumi-juku) of the Tokyo Tech Organization for Fundamental Research (OFR). The authors declare no competing financial interests.

Author contributions

T. Shimi conceived the project. Y. Kono designed, performed and interpreted experiments. C.G. Pack performed FCS experiments. T. Ichikawa performed AFM experiments. A. Komatsubara performed image analysis. L. Rolas and S. Nourshargh generated, characterized, and provided the anti-murine progerin antibody. S.A. Adam, K. Miyazawa, O. Medalia and R.D. Goldman provided advice and scientific discussion. Y. Kono, C.G. Pack, T. Ichikawa and T. Shimi drafted the manuscript. S.A. Adam, R.D. Goldman, and H. Kimura edited the manuscript. T. Shimi, S.A. Adam, T. Fukuma, H. Kimura and R.D. Goldman supervised the experiments.

References

1. J. M. Zullo *et al.*, DNA sequence-dependent compartmentalization and silencing of chromatin at
the nuclear lamina. *Cell* **149**, 1474-1487 (2012).

2. M. Crisp *et al.*, Coupling of the nucleus and cytoplasm: role of the LINC complex. *J Cell Biol* **172**,
41-53 (2006).

3. G. K. Voeltz, M. M. Rolls, T. A. Rapoport, Structural organization of the endoplasmic reticulum.
EMBO Rep **3**, 944-950 (2002).

4. P. L. Paine, L. C. Moore, S. B. Horowitz, Nuclear envelope permeability. *Nature* **254**, 109-114
(1975).

5. R. Foisner, L. Gerace, Integral Membrane-Proteins of the Nuclear-Envelope Interact with Lamins
and Chromosomes, and Binding Is Modulated by Mitotic Phosphorylation. *Cell* **73**, 1267-1279
(1993).

6. L. Gerace, A. Blum, G. Blobel, Immunocytochemical localization of the major polypeptides of the
nuclear pore complex-lamina fraction. Interphase and mitotic distribution. *J Cell Biol* **79**, 546-566
(1978).

7. L. Gerace, G. Blobel, The nuclear envelope lamina is reversibly depolymerized during mitosis.
Cell **19**, 277-287 (1980).

8. F. Lin, H. J. Worman, Structural organization of the human gene encoding nuclear lamin A and
nuclear lamin C. *J Biol Chem* **268**, 16321-16326 (1993).

9. T. H. Höger, K. Zatloukal, I. Waizenegger, G. Krohne, Characterization of a second highly
conserved B-type lamin present in cells previously thought to contain only a single B-type lamin.
Chromosoma **99**, 379-390 (1990).

10. G. Biamonti *et al.*, The gene for a novel human lamin maps at a highly transcribed locus of
chromosome 19 which replicates at the onset of S-phase. *Mol Cell Biol* **12**, 3499-3506 (1992).

11. F. Lin, H. J. Worman, Structural organization of the human gene (LMNB1) encoding nuclear lamin
B1. *Genomics* **27**, 230-236 (1995).

12. H. Maeno, K. Sugimoto, N. Nakajima, Genomic structure of the mouse gene (Lmnb1) encoding
nuclear lamin B1. *Genomics* **30**, 342-346 (1995).

13. T. Shimi *et al.*, Structural organization of nuclear lamins A, C, B1, and B2 revealed by
superresolution microscopy. *Mol Biol Cell* **26**, 4075-4086 (2015).

14. Y. Turgay *et al.*, The molecular architecture of lamins in somatic cells. *Nature* **543**, 261-264 (2017).

15. T. Shimi *et al.*, The A- and B-type nuclear lamin networks: microdomains involved in chromatin
organization and transcription. *Genes Dev* **22**, 3409-3421 (2008).

16. M. Kittisopikul *et al.*, Computational analyses reveal spatial relationships between nuclear pore
complexes and specific lamins. *J Cell Biol* **220** (2021).

17.   R. J. Lutz, M. A. Trujillo, K. S. Denham, L. Wenger, M. Sinensky, Nucleoplasmic localization of
prelamin A: implications for prenylation-dependent lamin A assembly into the nuclear lamina.
Proc Natl Acad Sci U S A **89**, 3000-3004 (1992).

18.   M. Sinensky *et al.*, The processing pathway of prelamin A. *J Cell Sci* **107 (Pt 1)**, 61-67 (1994).

19.   V. L. Boyartchuk, M. N. Ashby, J. Rine, Modulation of Ras and a-factor function by carboxyl-
terminal proteolysis. *Science* **275**, 1796-1800 (1997).

20.   A. M. Winter-Vann, P. J. Casey, Post-prenylation-processing enzymes as new targets in
oncogenesis. *Nat Rev Cancer* **5**, 405-412 (2005).

21.   G. K. Leung *et al.*, Biochemical studies of Zmpste24-deficient mice. *J Biol Chem* **276**, 29051-
29058 (2001).

22.   D. P. Corrigan *et al.*, Prelamin A endoproteolytic processing in vitro by recombinant Zmpste24.
Biochem J **387**, 129-138 (2005).

23.   C. M. Denais *et al.*, Nuclear envelope rupture and repair during cancer cell migration. *Science* **352**,
353-358 (2016).

24.   M. Raab *et al.*, ESCRT III repairs nuclear envelope ruptures during cell migration to limit DNA
damage and cell death. *Science* **352**, 359-362 (2016).

25.   C. T. Halfmann *et al.*, Repair of nuclear ruptures requires barrier-to-autointegration factor. *J Cell*
Biol **218**, 2136-2149 (2019).

26.   J. D. Vargas, E. M. Hatch, D. J. Anderson, M. W. Hetzer, Transient nuclear envelope rupturing
during interphase in human cancer cells. *Nucleus* **3**, 88-100 (2012).

27.   J. Robijns *et al.*, In silico synchronization reveals regulators of nuclear ruptures in lamin A/C
deficient model cells. *Sci Rep* **6**, 30325 (2016).

28.   N. Y. Chen *et al.*, Fibroblasts lacking nuclear lamins do not have nuclear blebs or protrusions but
nevertheless have frequent nuclear membrane ruptures. *Proc Natl Acad Sci U S A* **115**, 10100-
10105 (2018).

29.   N. Y. Chen *et al.*, An absence of lamin B1 in migrating neurons causes nuclear membrane ruptures
and cell death. *Proc Natl Acad Sci U S A* **116**, 25870-25879 (2019).

30.   Y. Kono *et al.*, Nucleoplasmic lamin C rapidly accumulates at sites of nuclear envelope rupture
with BAF and cGAS. *J Cell Biol* **221** (2022).

31.   H. J. Worman, G. Bonne, "Laminopathies": a wide spectrum of human diseases. *Exp Cell Res* **313**,
2121-2133 (2007).

32.   W. H. De Vos *et al.*, Repetitive disruptions of the nuclear envelope invoke temporary loss of
cellular compartmentalization in laminopathies. *Hum Mol Genet* **20**, 4175-4186 (2011).

33.   A. J. Earle *et al.*, Mutant lamins cause nuclear envelope rupture and DNA damage in skeletal
muscle cells. *Nat Mater* **19**, 464-473 (2020).

34.   P. H. Kim *et al.*, Nuclear membrane ruptures underlie the vascular pathology in a mouse model of

788   Hutchinson-Gilford progeria syndrome. *JCI Insight* **6** (2021).

789   35.   A. De Sandre-Giovannoli *et al.*, Lamin a truncation in Hutchinson-Gilford progeria. *Science* **300**,

790   2055 (2003).

791   36.   M. Eriksson *et al.*, Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford

792   progeria syndrome. *Nature* **423**, 293-298 (2003).

793   37.   A. M. Young, A. L. Gunn, E. M. Hatch, BAF facilitates interphase nuclear membrane repair

794   through recruitment of nuclear transmembrane proteins. *Mol Biol Cell* **31**, 1551-1560 (2020).

795   38.   R. M. Sears, K. J. Roux, Mechanisms of A-Type Lamin Targeting to Nuclear Ruptures Are

796   Disrupted in LMNA- and BANF1-Associated Progerias. *Cells* **11** (2022).

797   39.   O. A. Zhironkina *et al.*, Mechanisms of nuclear lamina growth in interphase. *Histochem Cell Biol*

798   **145**, 419-432 (2016).

799   40.   S. Cho *et al.*, Progerin phosphorylation in interphase is lower and less mechanosensitive than

800   lamin-A,C in iPS-derived mesenchymal stem cells. *Nucleus* **9**, 230-245 (2018).

801   41.   I. C. Lopez-Mejia *et al.*, A conserved splicing mechanism of the LMNA gene controls premature

802   aging. *Hum Mol Genet* **20**, 4540-4555 (2011).

803   42.   F. G. Osorio *et al.*, Splicing-directed therapy in a new mouse model of human accelerated aging.

804   *Sci Transl Med* **3**, 106ra107 (2011).

805   43.   I. L. Ivanovska, M. P. Tobin, T. Bai, L. J. Dooling, D. E. Discher, Small lipid droplets are rigid

806   enough to indent a nucleus, dilute the lamina, and cause rupture. *J Cell Biol* **222** (2023).

807   44.   M. Zwerger *et al.*, Altering lamina assembly reveals lamina-dependent and -independent functions

808   for A-type lamins. *J Cell Sci* **128**, 3607-3620 (2015).

809   45.   V. Kochin *et al.*, Interphase phosphorylation of lamin A. *J Cell Sci* **127**, 2683-2696 (2014).

810   46.   T. Shimi, C. G. Pack, R. D. Goldman, Analyses of the Dynamic Properties of Nuclear Lamins by

811   Fluorescence Recovery After Photobleaching (FRAP) and Fluorescence Correlation Spectroscopy

812   (FCS). *Methods Mol Biol* **1411**, 99-111 (2016).

813   47.   D. Holtz, R. A. Tanaka, J. Hartwig, F. McKeon, The CaaX motif of lamin A functions in

814   conjunction with the nuclear localization signal to target assembly to the nuclear envelope. *Cell*

815   **59**, 969-977 (1989).

816   48.   M. H. Gelb *et al.*, Therapeutic intervention based on protein prenylation and associated

817   modifications. *Nat Chem Biol* **2**, 518-528 (2006).

818   49.   R. Lee *et al.*, Genetic studies on the functional relevance of the protein prenyltransferases in skin

819   keratinocytes. *Hum Mol Genet* **19**, 1603-1617 (2010).

820   50.   S. Y. Chang *et al.*, Inhibitors of protein geranylgeranyltransferase-I lead to prelamin A

821   accumulation in cells by inhibiting ZMPSTE24. *J Lipid Res* **53**, 1176-1182 (2012).

822   51.   H. Yao *et al.*, Targeting RAS-converting enzyme 1 overcomes senescence and improves progeria-

823   like phenotypes of ZMPSTE24 deficiency. *Aging Cell* **19**, e13200 (2020).

52.   X. Chen *et al.*, A small-molecule ICMT inhibitor delays senescence of Hutchinson-Gilford
progeria syndrome cells. *Elife* **10** (2021).

53.   K. N. Dahl *et al.*, Distinct structural and mechanical properties of the nuclear lamina in
Hutchinson-Gilford progeria syndrome. *Proc Natl Acad Sci U S A* **103**, 10271-10276 (2006).

54.   A. Quigley *et al.*, The structural basis of ZMPSTE24-dependent laminopathies. *Science* **339**, 1604-
1607 (2013).

55.   K. R. Yu *et al.*, MicroRNA-141-3p plays a role in human mesenchymal stem cell aging by directly
targeting ZMPSTE24. *J Cell Sci* **126**, 5422-5431 (2013).

56.   L. G. Fong *et al.*, Heterozygosity for Lmna deficiency eliminates the progeria-like phenotypes in
Zmpste24-deficient mice. *Proc Natl Acad Sci U S A* **101**, 18111-18116 (2004).

57.   A. Casasola *et al.*, Prelamin A processing, accumulation and distribution in normal cells and
laminopathy disorders. *Nucleus* **7**, 84-102 (2016).

58.   R. D. Goldman *et al.*, Accumulation of mutant lamin A causes progressive changes in nuclear
architecture in Hutchinson-Gilford progeria syndrome. *Proc Natl Acad Sci U S A* **101**, 8963-8968
(2004).

59.   T. Shimi *et al.*, The role of nuclear lamin B1 in cell proliferation and senescence. *Genes Dev* **25**,
2579-2593 (2011).

60.   A. Ivanov *et al.*, Lysosome-mediated processing of chromatin in senescence. *J Cell Biol* **202**, 129-
143 (2013).

61.   Z. Dou *et al.*, Autophagy mediates degradation of nuclear lamina. *Nature* **527**, 105-109 (2015).

62.   P. Scaffidi, T. Misteli, Reversal of the cellular phenotype in the premature aging disease
Hutchinson-Gilford progeria syndrome. *Nat Med* **11**, 440-445 (2005).

63.   O. Santiago-Fernández *et al.*, Development of a CRISPR/Cas9-based therapy for Hutchinson-
Gilford progeria syndrome. *Nat Med* **25**, 423-426 (2019).

64.   J. L. V. Broers *et al.*, Dynamics of the nuclear lamina as monitored by GFP-tagged A-type lamins.
J Cell Sci **112 (Pt 20)**, 3463-3475 (1999).

65.   T. Haraguchi *et al.*, Nuclear localization of barrier-to-autointegration factor is correlated with
progression of S phase in human cells. *J Cell Sci* **120**, 1967-1977 (2007).

66.   T. Haraguchi *et al.*, Live cell imaging and electron microscopy reveal dynamic processes of BAF-
directed nuclear envelope assembly. *J Cell Sci* **121**, 2540-2554 (2008).

67.   V. Stierlé *et al.*, The carboxyl-terminal region common to lamins A and C contains a DNA binding
domain. *Biochemistry* **42**, 4819-4828 (2003).

68.   A. C. Schibler, P. Jevtic, G. Pegoraro, D. L. Levy, T. Misteli, Identification of epigenetic
modulators as determinants of nuclear size and shape. *Elife* **12**, 10.1101/2022.1105.1128.493845
(2023).

69.   K. Ikegami, S. Secchia, O. Almakki, J. D. Lieb, I. P. Moskowitz, Phosphorylated Lamin A/C in

the Nuclear Interior Binds Active Enhancers Associated with Abnormal Transcription in Progeria.

Dev Cell **52**, 699-713 e611 (2020).

70.   V. T. Chu *et al.*, Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced

precise gene editing in mammalian cells. *Nat Biotechnol* **33**, 543-548 (2015).

71.   Y. Guo, Y. Kim, T. Shimi, R. D. Goldman, Y. Zheng, Concentration-dependent lamin assembly

and its roles in the localization of other nuclear proteins. *Mol Biol Cell* **25**, 1287-1297 (2014).

72.   M. Roblek *et al.*, Monoclonal antibodies specific for disease-associated point-mutants: lamin A/C

R453W and R482W. *PLoS One* **5**, e10604 (2010).

73.   B. Marcos-Ramiro *et al.*, Isoprenylcysteine Carboxylmethyltransferase-Based Therapy for

Hutchinson-Gilford Progeria Syndrome. *ACS Cent Sci* **7**, 1300-1310 (2021).

74.   A. Sánchez-López *et al.*, Cardiovascular Progerin Suppression and Lamin A Restoration Rescue

Hutchinson-Gilford Progeria Syndrome. *Circulation* **144**, 1777-1794 (2021).

75.   H. Park, S. S. Han, Y. Sako, C. G. Pack, Dynamic and unique nucleolar microenvironment revealed

by fluorescence correlation spectroscopy. *FASEB J* **29**, 837-848 (2015).

76.   Y. Kanda, Investigation of the freely available easy-to-use software 'EZR' for medical statistics.

Bone Marrow Transplant **48**, 452-458 (2013).

77.   D. Bates, M. Mächler, B. M. Bolker, S. C. Walker, Fitting Linear Mixed-Effects Models Using

lme4. *Journal of Statistical Software* **67**, 1-48 (2015).

78.   A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen, lmerTest Package: Tests in Linear Mixed

Effects Models. *Journal of Statistical Software* **82**, 1-26 (2017).

79.   M. Mirdita *et al.*, ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682

(2022).

Abbreviations

BAF, barrier-to-autointegration factor; CBD, chromatin-binding domain; cGAS, cyclic GMP-AMP synthase; DARPin, designed ankyrin repeat protein; DCM, dilated cardiomyopathy; ESCRT-III, endosomal sorting complex required for transport-III; ER, endoplasmic reticulum; FAF, fluorescence auto-correlation function; FCS, fluorescence correlation spectroscopy; FTI, farnesyltransferase inhibitor; FPLD, familial partial lipodystrophy; HGPS, Hutchinson-Gilford Progeria syndrome; Ig-fold, immunoglobulin-like fold; INM, inner nuclear membrane; KD, knockdown; KO, knockout; LA, lamin A; LACS, LA-characteristic sequence; LB1, lamin B1; LB2, lamin B2; LC, lamin C; LINC, linker of nucleoskeleton and cytoskeleton; MEF, mouse embryonic fibroblast; MD, muscular dystrophy; NE, nuclear envelope; NL, nuclear lamina; NP, nucleoplasmic pool; NPC, nuclear pore complex; ONM, outer nuclear membrane; PG, progerin; REML, restricted maximum likelihood; sfGFP, superfolder GFP; sfCherry, superfolder Cherry; shRNA, short hairpin RNA; STING, stimulator of interferon genes; WT, wild type.

 Figure legends

Figure 1. **The localization kinetics of LA and PG at the rupture sites in WT and**
G609G/+ nuclei. (A) Protein architecture of LA and PG. The coiled-coil central rod
domain (gray), the NLS (yellow), the β-strand comprising the Ig-fold domain (blue), and
the CaaX motif (orange) are shown in the diagram. Anti-LA (4A58) and anti-PG do not
cross-react with each other. Anti-pre-LA (7G11) recognizes the last amino acids that is
proteolytically cleaved off and does not react with PG. Anti-mature-LA (4A4-A4)
recognizes mature-LA but not pre-LA. Anti-LA/C (3A6-4C11) recognizes all A-type
lamin isoforms. **(B)** NE rupture was induced by AFM indentation using a cantilever with
a round tip, as shown in the diagram. **(C** and **D)** Time-lapse images of NLS-sfCherry
expressed in WT and G609G/+ cells were acquired with 30 s intervals for 5 min after the
induction of NE rupture by AFM indentation. **(C)** The fluorescence intensities in the cells
are shown with rainbow color. Bar: 10 μm. **(D)** The nuclear-to-cytoplasmic intensity
(N/C) ratios of NLS-sfCherry were measured and normalized to the initial intensities, and
plotted to monitor the nuclear reentry in the cells (means ± SEM; n = 20 cells from two
independent experiments; ***, P < 0.001 from WT by a mixed effect model). **(E)** NE
rupture was induced by 405-nm laser microirradiation with a 2-μm diameter spot at the
NE of WT and G609G/+ nuclei. **(F** and **G)** 13 to 21 of nuclei of the cells were laser-
microirradiated within 10 min and incubated for 0, 30, 60 and 120 min, followed by
fixation with 4% PFA/0.1% Triton-X 100. The fixed cells were immunostained with a
combination of the LA and PG antibodies, and the DNA was stained with Hoechst 33342.
(F) Representative confocal images of nuclei in the cells fixed within 60-70 min after the
induction of NE rupture by laser microirradiation. Magnified views of the areas indicated
with orange boxes are shown (bottom). The rupture sites are indicated with yellow
arrowheads (bottom). Bars: 5 μm (top) and 2μm (bottom). **(G)** Percentiles of the cells
with (blue) and without (gray) the localization of LA and PG to the rupture sites. The
numbers of analyzed cells from two independent experiments are indicated in the bar
charts. The line graphs show the kinetics of the percentage of the cells with localization
of LA and PG to the rupture sites.

Figure 2. **Localization of LA and PG to the rupture sites in DMSO or FTI-treated**
WT and G609G/+ nuclei. (A) Whole cell lysates from WT and G609G/+ cells treated
with 0.1% DMSO or 3.2 μM lonafarnib (FTI) were probed with anti-LA/C (3A6-4C11),
anti-PG, and anti-GAPDH (as loading control) for immunobloting. Positions of the size
standards are shown on the right. **(B** and **C)** The indicated cells were immunostained with
the LA (4A58) and PG antibodies. **(B)** Representative microscopic images. **(C)** Ratios of
the average fluorescence intensity in the nucleoplasmic pool relative to that of the NL
(NP/NL ratios) of LA and PG were measured based on immunofluorescence (n = 20 cells

from two independent experiments; ***, P < 0.001 by Welch's t-tests). **(D)** 15 to 19 of
nuclei in the cells were laser-microirradiated and incubated for 60 min, followed by
fixation with 4% PFA/0.1% Triton-X 100. The fixed cells were immunostained with a
combination of the LA and PG antibodies. Percentiles of the cells with (blue) and without
(gray) the localization of LA and PG at the rupture sites. The numbers of analyzed cells
from two independent experiments are indicated in the bar charts.
940
Figure 3. **The diffusion kinetics of LA and PG in the nucleoplasm of DMSO or FTI-**
treated WT and G609G/+ nuclei. The diffusion fractions of mEmerald-LA (Em-LA)
and mEmerald-PG (Em-PG) in DMSO- or FTI-treated WT and G609G/+ nuclei were
measured by FCS. **(A)** A representative confocal image of a WT nucleus with Em-LA
before FCS measurements (left). The yellow crosses indicate the points measured by FCS.
Bar: 10 μm. Em-LA or Em-PG molecules move in or out of the confocal volume (white-
out cylinder region in blue) at different speeds, as shown in the diagram (right). **(B)**
Representative fluorescence fluctuations of Em-LA in DMSO- (green line) and FTI-
treated (blue line) of WT and G609G/+ nuclei are plotted by count per second (CPS). **(C)**
Diffusion coefficients of fast component (plotted in black) and slow component (plotted
in gray) for Em-LA in the cells. **(D)** Representative fluorescence fluctuations of Em-PG
in DMSO- (green line) and FTI-treated (blue line) of WT and G609G/+ nuclei are plotted
by CPS. **(E)** Diffusion coefficients of fast component (plotted in black) and slow
component (plotted in gray) for Em-PG in the cells. **(C** and **E)** Mean ± SEM are indicated
on the left to plots; n = 10 cells from two independent experiments; *, P < 0.05 by a
Games–Howell test. Percentiles of these component fractions are indicated at the bottom
of the graph. N/A, not applicable because of photobleaching.
958
Figure 4. **The localization of pre- and mature-LA to the rupture sites in the control**
and Zmpste24-KD WT MEFs treated with DMSO or FTI. (A) Whole cell lysates from
the control and Zmpste24-KD cells treated with DMSO or FTI were probed with anti-
LA/C (3A6-4C11) and anti-GAPDH (as loading control). Positions of the size standards
are shown on the right. **(B)** The control and shZmpste24 cells treated with DMSO or FTI
were immunostained with a combination of anti-pre-LA (7G11) and anti-mature-LA
(4A4-A4). Bar: 5 μm. **(C** and **D)** NP/NL ratios of pre-LA (C) and mature-LA (D) in the
cells were measured based on immunofluorescence (n = 20 cells from two independent
experiments; ***, P < 0.001 by Games–Howell tests for pre-LA and by Welch's t-test for
mature-LA). **(E)** 15 to 18 of nuclei in the cells were laser-microirradiated within 10 min
and incubated for 60 min, followed by fixation with 4% PFA/0.1% Triton-X 100. The
fixed cells were immunostained with a combination of the mature-LA and pre-LA
antibodies. Percentiles of the indicated cells with (blue) and without (gray) the
localization of LA and pre-LA to NE rupture sites. The numbers of analyzed cells from

two independent experiments are indicated in the bar charts. **(C-D)** N/A, not applicable
due to below detection limit.

Figure 5. **Rapid accumulation of LA truncation and PG deletion mutants at the**
rupture sites in LA/C-KO nuclei. **(A)** Architecture of mEmerald-fused LA-full-length
(FL), PG, PG-CSM, L648X, T644X, F627X, S617X, A601X, S575X, LAΔ26, LAΔ26-
L648X, LAΔ26-L644X, LAΔ26-L627X, and LC. Terminal amino acids are numbered,
and their accumulation dynamics is summarized on the right (+, accumulated at the
rupture site; -, not accumulated). **(B** and **C)** The mEmerald-fused LA-FL, PG, PG-CSM,
L648X, T644X, F627X, S617X, A601X, S575X and LC were ectopically expressed
under the *Lmna* promoter in LA/C-KO cells. **(B)** Representative confocal images of the
indicated nuclei. Bar: 5 μm. **(C)** The NP/NL ratios were measured based on mEmerald
signals (n = 15-20 cells from two independent experiments; ‡, P<0.001 compared to
others except PG-CSM and *, P < 0.05 by a Games–Howell test). **(D** and **E)** Time-lapse
images of mEmerald-fused LA-FL, LA mutants, PG, PG deletion mutant and LC were
acquired with 10 s intervals for 3 min after the induction of NE rupture by laser-
microirradiation, and the relative intensities at the rupture sites are plotted in the graphs.
The fluorescence intensities at the rupture sites were measured and normalized to the
initial intensities (means ± SEM; n =20 cells from two independent experiments; ***, P
< 0.001 from LA-FL by a mixed effect model). **(D)** mEmerald-fused LA-FL, PG, PG-
CSM, L648X, T644X, F627X, S617X, A601X, S575X and LC. **(E)** mEmerald-fused LA-
FL, LAΔ26, LAΔ26-L648X, LAΔ26-L644X and LAΔ26-L627X. LA-FL and LC are
reproduction of D.

Figure 6. **Schematic diagram of the difference in localization to the rupture sites**
between LA, LC and PG. LA (magenta) is slowly localized to the rupture sites compared
to LC (green) because it contains two LACSs in the specific tail region (top and middle).
PG (yellow) is even slower in localizing to the rupture sites than LA because of the
permanent association with the INM. PG expression also slows the recruitment of LA and
LC to the rupture sites. When the farnesylation of PG is inhibited by FTI treatment, the
localization kinetics becomes similar to that of LA (bottom).

Table S1. **Phenotype characterization of WT and G609G/+ MEFs.**

NLS-sfCherry expressing WT and G609G/+ cells were stained with Hoechst 33342 for

DNA and RFP-Booster Alexa Fluor 568 to enhance the signal of NLS-sfCherry to detect

spontaneous NE rupture. Immunostaining with α-LB1 and α-cGAS antibodies were also

used as NE rupture markers. See Fig. S1 A for the confocal images. P values, Fisher's

exact tests between WT and G609G/+.

1010

1011 Table S2. **Oligonucleotides used in this study.**

1012 See Materials and methods. Oligonucleotides were synthesized by Sigma-Aldrich or

1013 Integrated DNA Technologies (IDT).

1014

Figure S1. **Phenotype characterization, the localization kinetics of LA and PG at the**

rupture sites in WT and G609G/+ MEFs, and the localization kinetics of LA at the

rupture sites in BAF-KD WT MEFs. (A) The phenotype of heterozygous knock-in

G609G/+ expressing NLS-sfCherry was characterized by immunofluorescence. The cells

were immunostained with anti-LB1 and anti-cGAS, and the DNA was stained with

Hoechst 33342. The fluorescence intensities of NLS-sfCherry are shown with rainbow

color. Asterisk indicates a cell with NE breakdown due to the G2-M transition and were

excluded from the analysis. Misshapen, NE-ruptured, LB1-mis-localized and cGAS-

accumulated nuclei are indicated with magenta arrowheads. **(B)** Representative confocal

images of the nuclei. Magnified views of the indicated areas by orange boxes are shown

(bottom of each row). The cells were fixed within 0-10 min (top), 30-40 min (middle) and

120-130 min (bottom) after the induction of NE rupture by laser microirradiation,

followed by fixation with 4% PFA/0.1% Triton-X 100. The fixed cells were

immunostained with anti-LA (4A58) and anti-PG. The rupture sites are indicated with

yellow arrowheads (bottom of each row). Bars: 5 μm (top of each row) and 2 μm (bottom

of each row). **(C)** Whole cell lysates from WT cells expressing shRNAs for the control

(Scramble) or two shRNAs (shBAF #1 and #2) were probed with anti-BANF1/BAF and

anti-GAPDH (as loading control) for immunostaining. Positions of the size standards are

shown on the right. **(D** and **F)** 15 to 19 of the indicated cells were laser-microirradiated

within 10 min and incubated for 60 min, followed by fixation with 4% PFA/0.1% Triton-

X 100. The fixed cells were immunostained with the LA and BAF antibodies. **(D)**

Representative confocal images of nuclei with (blue arrowheads) or without (magenta

arrowheads) the localization of LA to the rupture sites. Bar: 40 μm. **(E)** Percentiles of

cells with (blue) and without (gray) the localization of LA. The numbers of analyzed cells

from two independent experiments are indicated in the bar charts.

1040

1041 Figure S2. **Accumulation kinetics of LC at the rupture sites in WT and G609G/+**

1042 **MEFs. (A)** Protein architecture of LA, LC and PG with sfGFP-DARPin-LA6. The

summary of their accumulation at the rupture sites is indicated on the right (+, accumulated at the rupture site; -, not accumulated). The structural image of sfGFP-DARPin-LA6 was predicted by AlphaFold2 (79) and edited by the PyMOL (Schrödinger). **(B** and **C)** Time-lapse images of sfGFP-DARPin-LA6 expressed in WT and G609G/+ cells were acquired with 10 s intervals for 3 min after the induction of NE rupture by laser-microirradiation, and the relative intensities are plotted in the graph. **(B)** The dynamics of the sfGFP-DARPin-LA6 in response to NE rupture in the cells. Magnified views of the areas indicated by orange boxes are shown (the second to fifth columns). A 2-μm diameter spot at the NE was laser-microirradiated to induce NE rupture (yellow arrowheads). Bars: 5 μm (the first column) and 2 μm (the second to fifth columns). **(C)** The fluorescence intensities at the rupture sites were measured and normalized to the initial intensities (means ± SEM; n = 20 cells from two independent experiments; ***, P < 0.001 from WT by a mixed effect model).

Figure S3. **Establishment of knock-in cells expressing mEmerald-fused LA, PG and LC under the *Lmna* promoter for FCS measurements.** The DNA sequences of mEmerald-LC, mEmerald-LA and mEmerald-PG with the endogenous *Lmna* promotor were knocked-in to the *Rosa26* locus in WT and G609G/+ cells. **(A)** The schematic diagram of linearized pLmna-Em-LA-R26Neo, pLmna-Em-PG-R26Neo and pLmna-Em-LC-R26Neo knock-in vectors. **(B)** Whole cell lysates from the knocked-in cells with pLmna-Em-LA-R26Neo, pLmna-Em-PG-R26Neo and pLmna-Em-LC-R26Neo were probed with anti-LA/C (3A6-4C11) and anti-β-actin (as loading control) for immunostaining. Positions of the size standards are shown on the right. **(C-E)** The diffusion fractions and coefficients of mEmerald-LC (Em-LC) in WT and G609G/+ nuclei were measured by FCS. **(C)** A representative confocal image of a WT nucleus with mEmerald-LC before FCS measurements (top). The yellow crosses indicate the points measured by FCS. Bar: 10 μm. Em-LC molecules move in or out of the confocal volume (white-out cylinder region in blue) at different speeds, as shown in the diagram (bottom). **(D)** Diffusion coefficient of fast component (plotted in black) and slow component (plotted in gray) for Em-LC in the cells. Mean ± SEM are indicated on the left to plots; n = 10 cells from two independent experiments; ***, P < 0.001 by a Welch's t-test. Percentiles of these components are indicated at the bottom of the graph.

Figure S4. **Short-term localization of LA and PG to the rupture sites in DMSO or FTI-treated WT and G609G/+ nuclei.** 14 to 15 of nuclei in the cells were laser-microirradiated within 10 min, followed by fixation with 4% PFA/0.1% Triton-X 100. The fixed cells were immunostained with a combination of the LA and PG antibodies. Percentiles of the cells with (blue) and without (gray) the localization of LA and PG at the rupture sites. The numbers of analyzed cells from two independent experiments are

1082   indicated in the bar charts. The number of the cells in which LA and PG were localized
1083   to the rupture sites are indicated at the bottom of the graph.

1084

1085   Figure S5. **The immunoblotting of FTI-treated WT and G609G/+ knocked-in MEFs.**
1086   Whole cell lysates from the knocked-in cells with pLmna-Em-LA-R26Neo (Em-LA) and
1087   pLmna-Em-PG-R26Neo (Em-PG) treated with DMSO or FTI were probed with anti-
1088   LA/C (3A6-4C11) and anti-GAPDH (as loading control) for immunostaining. Positions
1089   of the size standards are shown on the right.

1090

Figure S6. **The localization kinetics at the rupture sites in the control and Zmpste24-**
KD WT MEFs. (A) Whole cell lysates from WT cells expressing Scramble for the
control or shZmpste24 for Zmpste24-KD were probed with anti-pre-LA (7G11), anti-
LA/C (3A6-4C11) and anti-GAPDH (as loading control) for immunobloting. Positions of
the size standards are shown on the right. **(B** and **C)** The cells were immunostained with
anti-LA (4A58) and the pre-LA (7G11) antibody. **(B)** Representative confocal images of
the nuclei. Bar: 5 µm. **(C)** NP/NL ratios of LA and pre-LA were measured based on
immunofluorescence (n = 20 cells from two independent experiments). **(D** and **E)** 18 of
nuclei in the cells were laser-microirradiated within 10 min and incubated for 60 min,
followed by fixation with 4% PFA/0.1% Triton-X 100. The fixed cells were
immunostained with the LA and pre-LA antibodies. **(D)** Representative confocal images
of the nuclei with (blue arrowheads) or without (magenta arrowheads) the localization of
LA and pre-LA to the rupture sites. Bar: 40 µm. **(E)** Percentiles of the cells with (blue)
and without (gray) the localization of LA and pre-LA. The numbers of analyzed cells
from two independent experiments are indicated in the bar charts.

1106

Figure S7. **Accumulation kinetics of LA and PG mutants at the rupture sites in LA/C-**
KO MEFs. (A) Amino acid sequences of LA- (blue) or LC-specific (red) regions from
Homo sapiens and *Mus musculus*. The amino acid homology is indicated as bars and a
gap is introduced between residue A600 and A601 in *Mus musculus* to maximize the
homology. **(B)** Dynamics of the mEmerald-LA-S575X and A601X in response to NE
rupture in LA/C-KO MEFs. **(C)** Representative images of mEmerald-LA-full-length (FL),
LA-FL without LACS1 (LAΔ26), LAΔ26-L648X, LAΔ26-T644X, LAΔ26-F627X, and
mEmerald-LC in LA/C-KO MEFs transfected for 2 d. Bar: 5 µm. **(D)** Fluorescence
intensities in the NP to the NL ratio of mEmerald-LA mutants and mEmerald-LC, as
control, were measured (n = 14-20 cells from two independent experiments; ***, P <
0.001 by a Games–Howell test). **(C** and **D)** LA-FL and LC are reproduction of Fig. 5 B
and C. **(E)** Dynamics of the mEmerald-LAΔ26-F627X and LAΔ26-T644X in response to
NE rupture in LA/C-KO MEFs. **(B** and **E)** Magnified views of the indicated areas by
orange boxes are shown (the second to fifth columns). A 2-µm diameter spot was laser-

microirradiated to induce NE rupture (yellow arrowheads). Bars: 5 μm (the first column)

and 2 μm (the second to fifth columns).

Fig. 1

A

Lamin A (LA)

Head, L1, 1A, 1B, α-helical rod, L12, 2A, L2, 2B, NLS, Ig-fold

α-LA (4A58), α-mature-LA (4A4-A4), α-pre-LA (7G11)

α-LA/C (3A6-4C11)

646, 664

Progerin (PG)

α-PG, Δ50aa, 614

B

Single-cell compression

AFM cantilever

100 μm glass bead

C

Before rupture

Time after rupture: 0 s, 60 s, 240 s

WT

G609G/+

High — Low

10 μm

D

NLS-sfCherry nucleoplasm/cytoplasm

WT
G609G/+

Time after rupture (s)

***

E

Laser microirradiation

405 nm

13~21 cells, 10 min

37°C incubation, 0, 30, 60, 120 min

PFA-fixation

F

WT — α-LA, α-PG, DNA

G609G/+ — α-LA, α-PG, DNA

60~70 (min)

5 μm

2 μm

G

Percentage and number of cells with localization

localized — not localized

α-LA WT: 0~10 (min), 30~40, 60~70, 120~130

α-LA G609G/+: 0~10, 30~40, 60~70, 120~130

α-PG G609G/+: 0~10, 30~40, 60~70, 120~130

Fig. 2

A

B

C

D

Fig. 3

Fig. 4

A

pre-LA
LA
LC
GAPDH

Scramble   shZmpste24
DMSO FTI DMSO FTI   (kD)

75
60
40
35

B

Scramble   shZmpste24
DMSO   FTI   DMSO   FTI

α-pre-LA

α-mature-LA

5 μm

C

Relative NP/NL intensity

1.4
1.2
1
0.8
0.6
0.4
0.2
0

***   ***

N/A

DMSO   FTI   DMSO   FTI
Scramble   shZmpste24
α-pre-LA

D

RelativeNP/NL intensity

1.4
1.2
1
0.8
0.6
0.4
0.2
0

***

N/A   N/A

DMSO   FTI   DMSO   FTI
Scramble   shZmpste24
α-mature-LA

E

Percentage and number of cells with localization

100%
75%
50%
25%
0%

N/A

N/A   N/A

DMSO   FTI   DMSO   FTI   DMSO   FTI   DMSO   FTI
Scramble   shZmpste24   Scramble   shZmpste24
α-pre-LA   α-mature-LA

■ localized   ■ not localized

Fig. 5

Fig. 6

Table S1. Phenotype characterization of WT and G609G/+ MEFs

	WT (n = 118 cells, 129 nuclei)	G609G/+ (n = 116 cells, 131 nuclei)	P value
DNA			0.465
normal	122 (94.6%)	120 (91.6%)	
mis-shapen	7 (5.4%)	11 (8.4%)	
NLS-sfCherry			0.102
nuclear-localized	110 (93.2%)	114 (98.3%)	
ruptured	8 (6.8%)	2 (1.7%)	
α-LB1			0.212
normal distributed	125 (96.9%)	130 (99.2%)	
mis-localized	4 (3.1%)	1 (0.8%)	
α-cGAS			1
not accumulated	129 (100%)	131 (100%)	
accumulated	0 (0%)	0 (0%)	

NLS-sfCherry expressing WT and G609G/+ cells were stained with Hoechst 33342 for DNA and RFP-Booster Alexa Fluor 568 to enhance the signal of NLS-sfCherry to detect spontaneous NE rupture. Immunostaining with α-LB1 and α-cGAS antibodies were also used as NE rupture markers. See Fig. S1 A for the confocal images. P values, Fisher's exact tests between WT and G609G/+.

Table S2. Oligonucleotides used in this study

No.	Name	Sequence (5' to 3')
1	shSte24#1-TRCN0000366641_S	CCGGCTGGATGCTCTTCCGGTTATTCTCGAGAATAACCGGAAGAGCATCCAGTTTTTG
2	shSte24#1-TRCN0000366641_AS	AATTCAAAAACTGGATGCTCTTCCGGTTATTCTCGAGAATAACCGGAAGAGCATCCAG
3	shBanf1#1-TRCN0000124958_Fw	CCGGTGGCCAGTTTCTGGTGCTAAACTCGAGTTTAGCACCAGAAACTGGCCATTTTTG
4	shBanf1#1-TRCN0000124958_Rv	AATTCAAAAATGGCCAGTTTCTGGTGCTAAACTCGAGTTTAGCACCAGAAACTGGCCA
5	shBanf1#2-TRCN0000124955_Fw	CCGGGACAAGGCTTATGTGGTCCTTCTCGAGAAGGACCACATAAGCCTTGTCTTTTTG
6	shBanf1#2-TRCN0000124955_Rv	AATTCAAAAAGACAAGGCTTATGTGGTCCTTCTCGAGAAGGACCACATAAGCCTTGTC
7	pX459V2-Fw	GAATTCTAACTAGAGCTCGCTGATCAGCCTCG
8	pX459V2-Rv	TGGGCCAGGATTCTCCTCGACGTCAC
9	if_mCherry_Fw	GAGAATCCTGGCCCAATGGTGAGCAAGGGCGAGGAGG
10	if_mCherry_Rv	CTCTAGTTAGAATTCTCACTTGTACAGCTCGTCCATGCCGCCGG
11	sgRosa26-1_s	CACCGACTCCAGTCTTTCTAGAAGA
12	sgRosa26-1_as	AAACTCTTCTAGAAAGACTGGAGTC
13	if_LA promoter_Fw	ATGCATTAGTTATTAATTGCCTCGAAAAACCAAAAAAG
14	if_LA promoter_Rv	GACCGGTAGCGCTAGCCCCGCTGTGACGGGGTCTCCATG
15	if_mLA_Fw	ATGGAGACCCCGTCACAGCGG
16	if_mLA_Rv	AACCGGTAGCGCTAGCCATGATGCTGCAGTTCTGGG
17	mLA-to-PG_Fw	CAGAGCTCCCAGAACTGCAGC
18	mLA-to-PG_Rv	TCTGGGAGCTCTGGGCTCCCGCTCCACCG
19	mLA-L648X_Fw (mature)	CACCCGCTCCTACTAGCTGGGCAACTCC
20	mLA-L648X_Rv (mature)	GGAGTTGCCCAGCTAGTAGGAGCGGGTG
21	mLA-T644X_Fw	GACAACCTAGTCTAACGCTCCTACCTC
22	mLA-T644X_Rv	GAGGTAGGAGCGTTAGACTAGGTTGTC
23	mLA-F627X_Fw	GTCACTCGAAGCTGACGCAGTGTGGGG
24	mLA-F627X_Rv	CCCCACACTGCGTCAGCTTCGAGTGAC
25	mLA-S617X_Fw	CCTCTGGCTCTTGAGCCTCCAGTGTC
26	mLA-S617X_Rv	GACACTGGAGGCTCAAGAGCCAGAGG
27	mLA-A601X_Fw	GCTGACAAGGCTTGAGGTGGAGCGGGAG
28	mLA-A601X_R2	CTCCCGCTCCACCTCAAGCCTTGTCAGCAG
29	mLA-S575X_Fw	CACTGCAGCGGCTAGGGGGACCCCGC
30	mLA-S575X_Rv	GCGGGGTCCCCCTAGCCGCTGCAGTG
31	if_mLA-del575-600_Fw	CAGCGGCGCCGGTGGAGCGGGAGCCC
32	if_mLA-del575-600_Rv	CCACCGGCGCCGCTGCAGTGGGAACCAC
33	if_Rosa26L-mLA_Fw	GTCTTTCTAGCCTTAATTAATGCCTCGAAAAACCAAAA
34	if_mLA-bGHpA-Rv	GAGGTCGACCAATTGTTACATGATGCTGCAGTTCTG
35	if_mLA-CSM-bGHpA-Rv	GAGGTCGACCAATTGTTACATGCTGCAGTTCTGGGAGCT
36	if_bGHpA_Fw	CAATTGGTCGACCTCGACCTCGACTG
37	if_Rosa26R-bGHpA_Rv	GCGGCCGCCGCGCCCACGCGTTCCCCAGCATGCC
38	if_Nsi-BGHpA_Fw	GGCAGCCGCCGCTGAATGCATCAATTGGTCGACCTCGACCTCGACTGT
39	if_mLA-to-mLC_Rv	TCAGCGGCGGCTGCCACTCACACGGTGGTGATGGAGGAGCTCTTCT

See Materials and methods. Oligonucleotides were synthesized by Sigma-Aldrich or Integrated DNA Technologies (IDT).

Fig. S1

Fig. S1

C

D

E

Percentage and number of cells with accumulation

Fig. S2

A

A-type lamins
LMNA

Rapid accumulation

Lamin A (LA) — −

sfGFP-DARPin-LA6

Head α-helical rod NLS Ig-fold

L1 L12 L2

1A 1B 2A 2B

646 | 664

Lamin C (LC) — +

Progerin (PG) — −

Δ50aa 614

B

Before rupture

Time after rupture

0 s 50 s 180 s

WT

5 μm 2 μm

G609G/+

C

***

Fig. S3

Fig. S4

0-5min localized	1	2	0	0	0	0
5-10min localized	1	3	2	5	0	11
not localized	26	23	26	25	28	19

Fig. S5

Fig. S6

Fig. S7

A

LA

```
                    ┌──────── LACS1 ────────┐                    ┌──────── LACS2 ────────┐
Homo sapiens   564 HHHGSHCSSSGDPAEYNLRSRTVLCGTCGQPADKASASGSGAQVGGPISSGSSASSVTVTRSYRSVGGSGGGSFGDNLVTRSYLLGNSSPRTQSPQNCSIM 664
                   |||.||||||||||||||||||||||||||||||||.|.||||||||||||||||||||||||.|||||||||||||||||||||||.||.|||||||
Mus musculus   566 HHRGSHCSGSGDPAEYNLRSRTVLCGTCGQPADKA-AGGAGAQVGGPISSGSSASSVTVTRSFRSVGGSGGGSFGDNLVTRSYLLGNSSPRSQSSQNCSIM 665
```
        S575X   A601X   S617X F627X   L648X
                                                                T644X

LC

```
Homo sapiens   564 HHHVSGSRR 572
                   ||.||||||
Mus musculus   566 HHRVSGSRR 574
```
        568

B

C

D

E