Title: Ecological feedback limits the spread of an adaptive β-lactamase mutant

Running title: Conditions for coexistence of β-lactamase mutants

Philip Ruelens[1*], Eline de Ridder[1,] J. Arjan G.M. de Visser[1, #], Meike Wortel[2,#]

[1] Laboratory of Genetics, Wageningen University & Research, Wageningen, The Netherlands

[2] Molecular Biology and Microbial Food Safety, Swammerdam Institute for Life Sciences,

University of Amsterdam, Amsterdam, The Netherlands

#Address correspondence to: arjan.devisser@wur.nl, m.t.wortel@uva.nl

*Present address: Philip Ruelens, Laboratory of Socioecology and Social Evolution, KU Leuven,

Leuven, Belgium

## Abstract

β-lactamases play an important role in global health problems due to antibiotic resistant bacterial infections. Understanding the spread of these enzymes may inform the development of better drug therapies. However, this is challenging, as β-lactamases affect their own selective advantage through antibiotic degradation, creating complex eco-evolutionary feedback. We investigated the effect of such feedback on the spread of bacterial strains expressing different β-lactamase enzymes conferring varying levels of resistance to the cephalosporin cefotaxime. Specifically, we conducted head-to-head competitions between two related clinically observed β-lactamase mutants, TEM-19 and TEM-52, and varied cefotaxime concentration and cell density to explore the impact of these factors on competitive fitness and coexistence. The results revealed that cell density, which modulates feedback strength, determined whether resistance alleles could replace susceptible alleles or coexist with them. Using a population dynamic model based only on measured parameters, we investigated the underlying mechanisms of these dynamics. By validating our model with experimental data, we show that it can to a large extend predict the equilibrium frequencies based on dose-dependent growth rates and antibiotic degradation activities of the individual strains. Our study emphasizes the importance of considering ecological feedback in the evolution of antibiotic resistance and the potential for predictive modelling, which could be applied in clinical contexts.

## Introduction

Antibiotic resistance is a major global health threat that poses a significant challenge to modern medicine (1). The emergence and spread of resistant bacterial strains have rendered many antibiotics ineffective, leading to longer hospital stays, and an increase in morbidity and mortality (2). β-lactamases, a family of enzymes that hydrolyze β-lactam antibiotics, are among the most common and clinically relevant antibiotic resistance determinants (3). Given their importance, the evolutionary dynamics of β-lactamase resistance have been extensively studied (4–6). A complicating factor, which has been ignored in most studies, is the potential influence of eco-evolutionary feedback due to antibiotic removal by the β-lactamases themselves on the invasion of new β-lactamase variants, which may significantly impact their competitive fitness and hence spread (7–12). A more quantitative understanding of the effect of eco-evolutionary feedback on the spread of β-lactamases is therefore required for a comprehensive understanding of their contribution to antibiotic resistance.

In population dynamics, invasion fitness is a common metric to assess whether a mutant can successfully invade and establish itself within a population (13). Negative frequency dependent selection, i.e., the benefit of a mutation decreases when it becomes more frequent, can prevent fixation and cause the coexistence of more and less resistant genotypes. In the context of antibiotic resistance, this can occur when resistance mutations decrease the environmental antibiotic concentrations, thereby benefitting less resistant mutants — a phenomenon known as cross-protection (7, 14–16). Cross-protection has been observed in high-density bacterial populations (7, 11, 14, 17, 18), including in clonal populations, where the effectiveness of an antimicrobial agent may decrease with increasing bacterial density, referred to as 'inoculum effect' (17). Cross-protection may obstruct resistance evolution and could lead to recovery of the susceptible type when antibiotic pressure diminishes. For cross-protection to lead to coexistence, it is necessary that resistance confers a fitness cost – the susceptible strain needs to grow faster without antibiotics to recover from its inhibition. This cost is often evident when antibiotic resistance determinants are located on a plasmid, but can also occur from single nucleotide polymorphisms affecting the activity of an enzyme (19). The extent to which competing bacteria expressing different antibiotic-degrading enzymes may influence each other's evolution through eco-evolutionary feedback remains unclear, because the fitness costs of a more active enzyme might be significantly lower compared to those associated with plasmid-acquired resistance, which could limit the potential for cross-protection to lead to coexistence.

In order to successfully spread, bacteria expressing β-lactamase enzymes with higher hydrolyzing
capacity must outcompete the bacteria from which they derive via mutation. Essentially, both
strains engage in a competition for limited resources, while particularly the more resistant strain
removes antibiotic from the environment through enzymatic degradation (Figure 1). When the
initial concentration of the antibiotic is above the minimum inhibitory concentration (MIC) of the
more susceptible strain, the resistant strain may rescue the susceptible strain if the antibiotic
concentration is reduced below the MIC of the susceptible strain before it is eliminated by the
antibiotic. If so, the susceptible strain can resume growth and increase in frequency when the
resistant strain exhibits a lower fitness in the absence of antibiotic (17). Population density plays
a crucial role in these competitive interactions, as high-density bacterial populations can degrade
antibiotics more rapidly, enhancing the probability of susceptible bacteria to survive until the
antibiotic concentration decreases below inhibitory concentrations. At low densities, in contrast,
resistant bacteria can grow longer without competition from susceptible bacteria, allowing their
relative frequency to increase more and/or the susceptible population to be killed entirely. The
critical density allowing survival of susceptible bacteria clearly also depends on the initial
antibiotic concentration. Notably, when the resistant strain has lower fitness than the susceptible
strain in the absence of antibiotic, conditions allowing the rescue of the susceptible bacteria by
the resistant strain may lead to the stable coexistence of both strains during the serial transfers.
88

**Figure 1. Schematic depiction of competition between resistant bacteria expressing an antibiotic-degrading**
**enzyme (pink) and susceptible bacteria (purple) in the presence of antibiotic (background shades of red).** The
enzyme inactivates the antibiotic, thereby reducing the antibiotic concentration in the environment and creating
conditions for susceptible bacteria to grow in the presence of resistant bacteria. The relative fitness of both strains
is density-dependent, as high-density bacterial populations can degrade antibiotics more rapidly, allowing
populations of susceptible bacteria to survive under otherwise lethal antibiotic concentrations.

In this study, we aim to investigate the extent to which ecological feedback caused by antibiotic
degradation can influence the evolution of resistance conferred by mutations in the naturally
occurring narrow-spectrum β-lactamase TEM-1. TEM-1 efficiently inactivates penicillins and first
generation cephalosporins, but just a few mutations are required to confer resistance to an
extended spectrum of β-lactams with serious clinical consequences (20–22). To understand how
resource competition and antibiotic breakdown affect the fitness of resistant relative to
susceptible strains, we competed two *Escherichia coli* strains expressing different extended
spectrum TEM alleles which differ by two point mutations (hereafter called β-lactamase
mutants), in the presence of varying concentrations of the third-generation cephalosporin
cefotaxime (CTX): TEM-19 (called "susceptible") and TEM-52 (called "resistant"), with low and
high activity against CTX, respectively. Moreover, we varied cell density through nutrient
availability to vary the rate of antibiotic breakdown, and therefore the level of eco-evolutionary
feedback. To quantitatively understand the experimental outcomes, we developed a population
dynamic model that interrogates how physiological parameters and ecological conditions
influence the evolutionary success of these β-lactamase mutants. Our results demonstrate a
crucial role for ecological feedback via antibiotic degradation and resource competition in
shaping the relative fitness of the competing strains and determining whether susceptible
bacteria may stably coexist alongside resistant ones or not.

**Results**

*Population density and antibiotic concentration mediate coexistence between β-lactamase*
*mutants*
To explore the extent by which antibiotic concentration and cell density affect the fitness and
possible coexistence of bacteria expressing different variants of an antibiotic-degrading enzyme,
we conducted serial passage competition experiments between two isogenic strains of
*Escherichia coli* expressing different TEM β-lactamase alleles: TEM-19 and TEM-52. These alleles
are clinically observed variations of TEM-1 β-lactamase, that differ in only two positions (TEM-19
has mutation G238S, and TEM-52 mutations E104K, M182T, and G238S), both with increased
cefotaxime (CTX) resistance (20, 23). Nevertheless, TEM-52 has an MIC for CTX that is more than

three orders of magnitude higher than that of TEM-19 ($MIC_{TEM-19}$ = 1.0 µg/ml, $MIC_{TEM-52}$ = 4,096
µg/ml; Supplementary Data 1). Thus, while both strains are able to degrade CTX more efficiently
than TEM-1, they do so at highly different rates (25), henceforth TEM-19 will be referred to as
the susceptible strain and TEM-52 as the resistant strain. To distinguish between these two
strains, the susceptible and resistant strain were chromosomally labelled with yellow (YFP) and
blue fluorescent protein (BFP), respectively. Importantly, the expression of these fluorescent
proteins did not differentially affect fitness (Supplementary Figure 1).

The maximum yield of the cultures was varied by supplementing glycerol as the sole carbon
source to a final concentration of 0.2%, 0.4%, 0.9%, or 2%. Furthermore, we investigated the
impact of three different initial antibiotic concentrations: 50, 25, or 12.5 µg/ml CTX. We
performed a total of five serial transfers using three starting ratios (1:9, 1:1 and 9:1) and eight
replicate cultures per condition and determined the fraction of resistant and susceptible bacteria
after each growth cycle by counting the number of YFP and BFP cells (cells/ml) using a flow
cytometer. We anticipated that if coexistence would occur, the different starting ratios would
converge to a single equilibrium ratio of both strains.

Our results show that the susceptible strain (TEM-19), managed to survive at antibiotic
concentrations significantly higher than its MIC in the presence of the resistant strain (TEM-52;
see Figure 2A). At an antibiotic concentration of 50 µg/ml, we observed that the average final
fraction of resistant cells decreased from 0.92 to 0.43 with increasing glycerol concentrations,
showing the effect of population density on the protection of the susceptible by resistant cells.
At lower initial antibiotic concentrations of 12.5 and 25 µg/ml, our chosen lowest population
densities were already high and higher densities hardly affected the final fraction of resistant
cells. There, we observed a gradual decline in the fraction of resistant cells over time, stabilizing
at approximately 0.21. However, we did observe an intriguing exception to this trend in
competitions initiated with a 1:9 ratio of resistant to susceptible bacteria at glycerol
concentrations of 0.2% and at 0.4% when supplemented with 50 µg/ml cefotaxime. Here, the
fraction of resistant cells surged rapidly during the initial growth cycle, overshooting the apparent
equilibrium and causing the extinction of the susceptible strain.

That the strain frequencies stabilize over the transfers, increasing to this value from a lower initial
fraction and decreasing form a higher initial fraction, suggests that a dynamic equilibrium exists.
In this equilibrium, strain frequencies still fluctuate across transfers, but in exactly the same
manner per transfer, leading to the same fraction at the beginning and the end of each transfer.
We inferred the stable equilibrium frequency of the resistant bacteria for each replicate

competition. Equilibrium values were derived by interpolating the relationship between initial
and final fractions per transfer (Figure 2B, Supplementary Information S1; see(7)). We then
employed generalized linear models to explore the influence of cell density (glycerol) and CTX
concentration, along with their interaction, on the equilibrium frequency of resistant cells. Our
findings indicate that incorporating the interaction of glycerol x CTX significantly improved the
model fit (including interaction: AICc = -174.59; excluding interaction: AICc = -168.70; Interaction
effect: $\chi_1^2 = 7.86$, p= 0.0050). The antibiotic concentration also significantly affected the
equilibrium frequency independently of glycerol (main effect antibiotic: $\chi_1^2$ = 155.567, p<0.0001),
while glycerol alone did not have a significant effect (main effect glycerol: $\chi_1^2$ = 0.450, p=0.50).
Our results thus revealed that the effect of glycerol on the equilibrium frequency was modulated
by the antibiotic concentration within the investigated range. In particular, glycerol, in the range
tested, had little impact at low CTX levels, but significantly determined equilibrium frequencies
at high antibiotic concentrations (Figure 2C).
174
175

**Figure 2. Conditions for coexistence of antibiotic-resistant and susceptible strain. (A)** Line plots showing the fraction of the resistant strain expressing TEM-52 β-lactamase in mixed cultures together with a relatively susceptible strain expressing TEM-19 β-lactamase across five serial transfers under different starting ratios, cell densities (varied by varying glycerol concentrations) and CTX concentrations. The error bars represent the standard error of the mean from eight biological replicates. Note that the error bars are not always visible due to the small standard error at that point. Dotted lines show the mean of the estimated equilibrium fractions. **(B)** Procedure to predict the equilibrium frequency of the resistant strain from experimental data by plotting the resistant frequency

every iteration during serial transfer. The equilibrium is reached when the lines connecting subsequent frequencies

cross the diagonal. The crossing of the diagonal will be lower for high densities (red) than for low densities (blue).

**(C)** Plot showing the joint impact of glycerol and CTX on the equilibrium fraction of resistant bacteria. The shaded

areas represent the smoothed 95% confidence interval around the model estimated means (depicted by the thicker

lines). Raw data means and standard errors are also depicted as dots and vertical bars.

*Physiological model predicts competitive dynamics and coexistence*

We then constructed a model that could predict the competitive dynamics and coexistence of

two strains as function of the antibiotic concentration and cell density, based on basic

physiological parameters. We expanded a previously used model with death in stationary phase

and antibiotic degradation by the more susceptible strain, two properties that we observed in

our system (7). The model parameters include the maximal growth rate, the death rate and

antibiotic clearance rate of each strain. The maximal growth rates ($0.185\pm0.00326$ $h^{-1}$ for the

resistant and $0.190\pm0.00219$ $h^{-1}$ for the susceptible strain) showed indeed a small but significant

cost of the resistance mutations, which is a necessary condition for coexistence. The death rate

of the susceptible strain ($1.54\pm0.186$ $h^{-1}$) was higher than expected, as in previous studies it was

in the order of the growth rate, while here it is an order higher (26, 27). The resistant strain

degraded the antibiotic faster than the susceptible strain, as expected. We found that this is

caused only by increased affinity (25), and the maximal degradation rate for the susceptible strain

is even higher than that of the resistant strain ($6.69*10^{-4} \pm 3.36*10^{-4}$ µg cell$^{-1}$ h$^{-1}$ versus $3.10 * 10^{-4}$

$^{4} \pm 9.15*10^{-5}$ µg cell$^{-1}$ h$^{-1}$ for the resistant strain). However, this maximal degradation rate of the

susceptible strain is never reached due to its low affinity for CTX. We estimated these parameters

in experiments (Supplementary Data 1) and used their values to parameterize our model and

predict the change in the relative frequencies of the strains at each passage (see Fig. 2A).

When accounting for only these basic properties, our model predicts that a higher initial

antibiotic concentration favors the resistant strain, while a higher cell density favors the

susceptible strain (Figure 3A). Because the dilution rate is fixed in these simulations, increasing

the cell density increases both the initial and final population size. The initial cell density affects

the dynamics, because a higher initial population size causes a faster degradation of the

antibiotic, and the final cell density affects the dynamics, because a higher final population size

allows for a longer growth period once the antibiotic is sufficiently degraded. Both these density

effects favor the more susceptible strain, which explains that the equilibrium frequency of the

resistant strain is both antibiotic and density-dependent (Figure 3A). Moreover, the range of CTX

and glycerol concentrations that allow for coexistence largely agree with the previously described

general linearized model derived from the experimental measurements (Figure 3A and B) and

shows that for the lower tested antibiotic concentrations (12.5 and 25 µg/ml) we would need
lower glycerol concentrations to see the effect of the culture density on the coexistence.

221
222

223
**Figure 3: Model predictions of the equilibrium resistant frequency. (A)** Contour plot showcasing model-projected
equilibrium frequencies of the resistant strain under a wider range of parameter values than the experimental data,
using physiological parameter measurements from both strains, across a variety of conditions. The region within the
white box corresponds to the experimentally examined range depicted in the contour plot of (B). **(B)** Contour plot,
interpolating the statistical model represented in Figure 2C, demonstrating how the equilibrium frequencies of the
resistant strain change depending on concentrations of glycerol and cefotaxime. White boxes indicate the
experimental raw data means.

*Effect of cell density on coexistence in continuous culture*
Whereas serial-passage experiments include cycles of resource supply and population size,
natural conditions often more resemble a continuous regime. Antibiotics are administrated
either intermittently, with up to several doses a day, or continuously. We modeled the
intermittent regime with a continuous culture model with antibiotics added in pulses to mirror
the clinical daily dosage administration of antibiotics. Blood cefotaxime concentrations of up to
200 µg/ml are reported (28, 29), depending on the weight of the person and the dosage, hence
we have chosen a concentration of 100 µg/ml after administration. Similar results are obtained
for continuous administration of antibiotics (see Supplementary section S.3). Under this
framework, the average population size is regulated by adjusting nutrient concentrations in the
inflow, while the actual population sizes fluctuate with the antibiotic pulses (Figure 4A).
Consequently, culture density remains relatively constant, eliminating the initial population size

variable. However, the impact of cell density on the competitive dynamics and coexistence of
antibiotic-resistant strains remains highly relevant and qualitatively similar to our previous
results. For a large range of antibiotic administration regimes, the outcome of the competition
between the susceptible and resistant strain depends on the cell density (Figure 4B).

**Figure 4. Modelling competition and coexistence under continuous nutrient input. (A)** The model parameterized
with the physiological parameters (Supplementary Data 1) and validated on the serial transfer experiments (Figure
3) is simulated in a continuous culture with antibiotic pulses (shown as shades of orange). After a pulse, the
susceptible population decreases due to death and washout, which increases the nutrient availability for the
resistant population, which therefore increases in size. The resistant cells degrade the antibiotic to a level at which
the susceptible population recovers, which reduces the nutrient availability and thereby the resistant population.
After an initial dynamic period, the fluctuations become regular and the ratio between the strains at the start of
each pulse reaches an equilibrium. **(B)** Antibiotic pulses and a constant nutrient supply leads to either a
monomorphic or a mixed population of bacteria expressing different TEM β-lactamase variants. The culture density
shows an effect on the equilibrium frequency of the resistant TEM variant. Cell densities and nutrients are modeled
as a continuous culture and CTX is modeled as regular pulses that lead to a concentration of 100 μg/ml in the culture.
The growth rates of the two strains are modelled with a logistic function that depends on the CTX concentration and
was experimentally obtained and a nutrient dependent term with an arbitrary but equal affinity.

## Discussion

The complex interplay between antibiotics and resistance mechanisms involving antibiotic
degradation vividly illustrates the concept of ecological feedback during evolution.

Understanding the evolutionary role of antibiotic degradation and other feedback mechanisms
is essential for a comprehensive understanding of the emergence and spread of antibiotic
resistance. To shed light on this topic, we explored the potential for eco-evolutionary feedback
via antibiotic degradation to modulate relative fitness and facilitate the coexistence of near-
isogenic strains with distinct levels of antibiotic resistance. Specifically, we focused on strains
carrying different alleles of the antibiotic-inactivating β-lactamase enzyme TEM-1, that differ by
just two mutations, resulting in large differences in their degradation efficiency of CTX and
variable fitness costs in the absence of the antibiotic (19). We found that both strains showed
dynamic fitness profiles that did not only depend on the antibiotic concentration, but also on the
culture density and the initial frequency of the strains. This dynamic fitness profile ultimately
dictates whether one strain dominates and becomes fixed within the population, or whether they
may stably coexist.

It is important to emphasize that in our study, when varying the culture cell density by adding
different levels of glycerol, we kept the number of generations constant, allowing us to decouple
the effect of the number of competitive generations and the culture density on competition
outcome. This deviates from the approach taken by Yurtsev et al. (2013), who demonstrated that
the initial cell density achieved by varying dilution factors can affect the equilibrium fraction.
Moreover, previous studies of coexistence via exposure protection (7, 30, 31) typically involved
coexistence between a susceptible population and a population with resistance conferred
through the introduction of an antibiotic resistance gene. This approach can introduce genetic
differences other than the resistance mechanism between the two populations, resulting in more
pronounced fitness differences. Our results demonstrate that eco-evolutionary feedback from
antibiotic degradation resulting from two single mutations can qualitatively affect the outcome
of competition, with resistant mutants compromising their own selective advantage and
frustrating the trajectory of evolution. Additionally, we show that we can predict coexistence
with a population dynamic model that does not need any fitting parameters, opening the way for
predictive modelling, as the model is not tailored to this specific experiment.

Antibiotic concentrations experienced by a bacterial population are dynamic and determined by
a multitude of factors, including physical environmental conditions such as temperature and pH,
as well as the metabolic activities of the bacteria themselves (17). Here, we have studied the
effect on the competitive success of bacteria expressing different β-lactamases, which can reduce
environmental antibiotic concentrations (17, 30, 32). Earlier modeling work examining a small set
of TEM-1 β-lactamase mutants found that the evolution of CTX resistance via mutations
enhancing CTX degradation is dependent on the administered antibiotic concentrations (33). In

another study, the coexistence of susceptible and plasmid-mediated ampicillin resistant
populations has been shown to lead to the evolution of the susceptible strain towards greater
antibiotic tolerance without significant increases in resistance, unlike monocultures where
resistance would increase (8). Our study demonstrates that exposure protection also occurs
among subtly different alleles of the TEM-1 gene, indicating the importance of eco-evolutionary
feedback interactions for the evolution of β-lactamase alleles with enhanced activity against the
β-lactam used for selection. Our findings demonstrate that predictions of evolutionary
trajectories of antibiotic resistance relying on traditional resistance measurements, such as MIC
(6), do not accurately capture the complex dynamics governing resistance evolution under all
conditions. Instead, determining relative fitness through competition experiments becomes
imperative to account for unavoidable eco-evolutionary feedback processes.
314
Our experiments revealed that in some cases, there was a marked increase in the resistant
fraction, at times even leading to the extinction of the susceptible population, especially when
the resistant fraction is initially low. This phenomenon can be attributed to the low number of
resistant cells, which keeps the antibiotic concentration at elevated levels for an extended period.
As a result, the susceptible population experiences a rapid and substantial decline that cannot be
compensated for by subsequent growth. This effect of overshooting the equilibrium, which we
observed in our experiments, has also been reported in a previous study (7), and is also replicated
in our model, underscoring its validity.
323
We modified a population dynamic model based only on very basic physiological parameters of
the two strains. Our results with this model show that we can largely understand the quantitative
outcomes of the experiments based on the properties of the individual strains, which we have
validated with experimental data. Our model's ability to predict coexistence without the need for
parameter estimation during model fitting (in contrast to previous similar approaches (e.g. (7)),
indicates that our model successfully captured the most relevant factors for predicting the effect
of ecological feedbacks in various scenarios. Importantly, our model allowed us to make
predictions beyond the laboratory setting and translate our results to more clinically relevant
conditions mimicked by a continuous culture with pulsed antibiotic administration,
demonstrating that the observed patterns hold true in more realistic scenarios.
334
The density effect on coexistence that we describe in this paper is expected to affect other
microbial interactions. Most closely, resistance conferred by other antibiotic-degrading enzymes,
including other β-lactamases and acetyltransferases capable of inactivating aminoglycosides or
chloramphenicol (34), is likely to involve similar mechanisms to those discussed here. Population

density also affects metabolic interactions: cross-feeding is only beneficial at intermediate
densities and production of compounds that are self-inhibitory can lead to coexistence
depending on the density (35, 36). Therefore, population density effects on coexistence are also
relevant to understand the evolution of microorganisms in microbial communities.
343
While our model successfully captures the feedback effect of antibiotic degradation on the
selective benefit of a more active β-lactamase allele, it has limitations compared to natural
populations and conditions. One of the limitations is that our study is limited to mutations
enhancing the activity of the β-lactamase and does not consider compensatory mutations that
can reduce the costs of resistance, which may reduce the potential for cross-protection to
frustrate the evolution of antibiotic resistance. This also means we do not consider other, more
private, mechanisms of resistance, such as target modification or decreased intake or increased
efflux of antibiotics, which will potentially come with different forms of eco-evolutionary
feedback (37, 38). Expanding our model to include spatially-structured conditions, such as those
found in biofilms, is also important. These environments are more representative of natural
conditions and may significantly influence the dynamics we have observed (11, 31, 39).
Nevertheless, our study offers valuable insights into the intricate relationship between the
dynamics of selective conditions and β-lactamase evolution, underscoring the need to consider
the complexity of the system when developing novel approaches to combat antibiotic resistance.
By acknowledging these factors, we can develop more effective strategies to prevent the
emergence and spread of antibiotic resistance, ultimately leading to improved health outcomes.

## Methods

*Strains, plasmids and medium*

The *Escherichia coli* strain used throughout this study was BW27783 (CGSC#12119), which lacks arabinose metabolizing genes and expresses *araE* from an arabinose independent promoter allowing stringent control of the arabinose-inducible promoter, pBAD (40). Plasmid pBAD332T was used as the expression vector to express TEM-1 and its alleles, TEM-19 (G238S) and TEM-52 (E104K-M182T-G238S)(41). To differentiate between the two different alleles during the coexistence experiment, BW27783 was labelled with a yellow (YFP) or blue (BFP) fluorescent marker by inserting cat-J23101-SYFP2 or cat-J23101-mTagBFP2 into chromosomal *galK* using the Quick and Easy *E. coli* Gene Deletion Kit (Gene Bridges, Heidelberg, Germany). We evaluated the fitness effect of the fluorescent proteins through pairwise competition assays. Chromosomally labeled YFP and BFP cultures were mixed and diluted at a 1:1,000 ratio in fresh M9 minimal medium(n=24). Cell frequencies were measured at t=0h and t=24h using flow cytometry. Relative fitness was calculated as the ratio of Malthusian parameters (42).

TEM alleles were amplified from previously described plasmid constructs (43), ligated into pBAD322T using T4 DNA ligase (New England Biolabs), and then electroporated into *E. coli* DH5α. After verifying successful cloning via sequencing, isolated plasmid was transformed into BW27783, BW27783-YFP and BW27783-BFP. All experiments were performed in M9 minimal media (supplemented with 0.2% Casamino acids, 1μg/mL thiamine, and 10 μg/ml tetracycline to prevent plasmid loss). Glycerol was added to the medium as carbon source to a final concentration of 0.4% unless otherwise stated. L-arabinose (0.000625% or otherwise stated) was used to express the TEM-alleles.

*Growth rate, death rate and minimum inhibitory concentration*

To determine the growth rate of BW27783 expressing TEM alleles TEM-19 or TEM-52, 200 μL M9 medium supplemented with cefotaxime and L-arabinose in a 96-well microtiter plate was inoculated at a 1:200 dilution from overnight culture in M9. The samples were subsequently covered with 50 μL mineral oil to prevent evaporation and condensation of the lid. The 96-well plate was incubated for 24h at 37 °C in a Victor3 microtiter plate reader (PerkinElmer, Waltham, MA, USA), and absorbance was measured every three minutes. The plate was shaken (double orbital) for 30 seconds before each measurement. Growth rates represent the maximum linear slope of the natural logarithm of absorbance versus time in a 2 h window and were determined using a custom python script.

The death rate of BW27783 expressing TEM-19 was estimated by performing a time-kill
experiment. Bacteria from an overnight culture grown in M9 medium supplemented with L-
arabinose were exposed to 25 μg/ml cefotaxime. After 0, 2 and 6 hours of incubation at 37°C, the
number of colony forming units per ml were determined by plating on LB agar plates and
counting the number of colonies formed after overnight incubation. The death rate was
calculated as the slope of the regression line describing the logarithm of the bacterial density as
a function of time.
401
We determined the minimum inhibitory concentration of cefotaxime of both TEM mutants using
the liquid broth microdilution method (44). From overnight culture in M9 medium, approximately
200,000 cells were transferred to 200μl of M9 medium supplemented with cefotaxime and L-
arabinose. Plates were incubated for 24 h at 37 °C with agitation. The minimum inhibitory
concentration was visually assessed and defined as the lowest cefotaxime concentration where
no growth could be observed.
408
Growth rate and minimum inhibitory concertation were determined for BW27783 carrying
pBAD322T-TEM-19 and BW27783 with pBAD322T-TEM-52. Death rate was only determined of
BW27783 expressing TEM-19.

*Estimating the in vivo cefotaxime degradation rate*
To qualitatively asses the *in vivo* degradation rate of both β-lactamase alleles, we developed a
bioassay based on the agar diffusion assay (45). In this assay, we estimated the remaining
cefotaxime concentration in medium inoculated with BW27783 carrying either pBAD322T-G238S
or pBAD322T-E104K-M182T-G238S after defined time intervals. Each strain was initially cultured
overnight in 5ml M9 medium supplemented with 10 μg/ml tetracycline and 0.000625% arabinose
at 37°C with continuous shaking at 250 rpm. After 16 hours of incubation, the OD600 of these
samples were measured in a spectrophotometer and the cells were pelleted and resuspended in
5ml M9 medium with cefotaxime but without glycerol. The initial concentration cefotaxime for
the TEM-19 mutant was 32 μg/ml and for TEM-52 4096 μg/ml. These samples were incubated at
37°C and, after 15-, 30- and 45 minutes, cell-free supernatants (CFS) were prepared and placed
on ice. The CFS was prepared by pelleting down 1.4 ml culture and filter sterilizing the
supernatants using a 0.22 μm filter. From the CFS of each mutant, an eight times two-fold
dilutions series was made in M9 medium. At each time point, the number of colony-forming units
(CFUs) was determined by collecting 1 ml culture of each mutant after 0, 15, 30 and 45 min by
plating a 10,000-fold dilution on LB-agar plates and counting the number of colonies following
incubation overnight.

In order to assess the remaining concentration of cefotaxime at different time points, a control

dilution series with known cefotaxime concentrations was also prepared. LB-agar plates were

streaked with 100 µl of overnight culture of the cefotaxime susceptible WT BW27783. Four holes

were punctured in the agar arranged in equidistant positions with a pipette tip (P100). Each hole

was filled with 100 µl from the control dilution series (in triplicate) or 100 µl from the CFS-

dilutions series of each mutant (in triplicate) and incubated at 37°C for one day. As the susceptible

WT strain cannot grow at high CTX concentrations, a zone of inhibition (ZOI) was visible on the

plates of which the diameter was measured using a ruler. A calibration curve was prepared by

plotting the known CTX concentrations of the control dilution series and its measured ZOI. The

resulting equation was subsequently used to predict the amount of remaining CTX based on the

ZOI. For each time point, the predicted concentration was divided by the average CFUs between

the current time point and the previous one. The result of this was plotted against time, resulting

in a linear relationship. The slope of this line represents the estimated degradation rate (ug cell$^{-}$

$^{1}$ h$^{-1}$)

*Coexistence experiment*

To investigate the impact of population density and antibiotics on two strains carrying either

TEM-19 or TEM-52, which were chromosomally labeled with yellow or blue fluorescent protein,

we conducted a serial passage competition experiment. The induced overnight cultures were

counted using a flow cytometer and mixed to three starting ratios (1:9, 1:1, and 9:1) in eight

biological replicates. Then, these mixed cultures were transferred to 96-well plates containing

M9 medium supplemented with glycerol (0.2, 0.43, 0.93, or 2%) and cefotaxime (50, 25, or 12.5

µg/ml) to a final volume of 200 µl. The starting population size of each condition was 200 times

less than the maximum cell number in the respective conditions. To ensure that the culture was

fully saturated, we incubated it for 48 hours at 37°C. After 48 hours of incubation with shaking,

each culture was diluted 20-fold and transferred to a new 96-well plate with freshly prepared M9

medium supplemented with glycerol and cefotaxime at the appropriate concentrations, and this

serial transfer was repeated five times. Following each growth cycle, each culture was diluted

100-fold in a new plate containing cold M9-medium before being counted using a MACSQuant10

flow cytometer (Miltenyi biotec, Bergisch Gladbach, Germany). We conducted flow cytometry

analysis with a moderate flow rate using SSC 1.4 as a trigger, and we counted a total of 20,000

events for each sample. For obtaining the equilibrium ratio from the experimental data, see

Supplementary information section S.1.

Data analysis was conducted using R statistical software (v3.6.1) accessed through RStudio (Build
576). To assess the relationship between the equilibrium fraction and antibiotic concentration
and density, a generalized linear model was applied using the 'glm' function (base R package).
Model selection was based on the second-order Akaike information criterion using 'AICc'
implemented in the 'MuMIn' package.

*Model description*

We follow the modeling approach from (7), extending their Model 7 with two additions. We
observe death in the lag phase and our susceptible strain also degrades the antibiotic. Therefore,
we have included a death term in the lag phase and degradation by the susceptible strain.
Including either of these features makes it impossible to directly calculate the equilibrium
fractions, therefore we have resorted to numerical methods to obtain the equilibrium fractions
(see supplementary Jupyter notebook).

Our model contains the following set of equations:

For $t < \tau_{lag}$:

479
$$\frac{dN_R}{dt} = 0$$

480
$$\frac{dN_S}{dt} = \begin{cases} 0, & A < MIC \\ -\gamma_D N_S, & A \geq MIC \end{cases}$$

481
$$\frac{dA}{dt} = -V_{max,R} N_R \frac{A}{A + K_{M,R}} - V_{max,S} N_S \frac{A}{A + K_{M,S}}$$

482 For $t > \tau_{lag}$:

483
$$\frac{dN_R}{dt} = \gamma_R N_R$$

484
$$\frac{dN_S}{dt} = \begin{cases} \gamma_S N_S, & A < MIC \\ -\gamma_D N_S, & A \geq MIC \end{cases}$$

485
$$\frac{dA}{dt} = -V_{max,R} N_R \frac{A}{A + K_{M,R}} - V_{max,S} N_S \frac{A}{A + K_{M,S}}$$

487 With $N_R$ the population size of the more resistant strain TEM-52, $N_S$ of the more susceptible
488 strain TEM-19 and $A$ the antibiotic concentration. For the parameters see Supplementary Table
489 S.1. See the supplementary information for the sensitivity analysis and the continuous culture
490 model.

## Acknowledgments

This study was supported by grants from the Human Frontiers in Science Program (RGP0010/2015) and the Netherlands organization for scientific research (OCENW.XS.058) to JAGMdV and an EMBO fellowship (ALTF 273-2017) to PR.

## Declaration of interests

The authors declare no competing interests.

## Appendices

**Supplementary Figure 1:** Relative competitive fitness of BW27783 carrying BFP to BW27783 carrying YFP. The statistical insignificance was assessed using the one-sample Wilcoxon non-parametric test (n=24).

**Supplementary Information:** Supporting information including Supplementary Section S1 (Equilibrium Ratio from Experimental Data), S2 (Serial Transfer Model), and S3 (Continuous Culture Model), Supplementary Tables S1, S2, and S3, as well as Supplementary Figures S1 through S4.

**Supplementary Data 1:** Source data underlying Figure 2 and Figure 3A.

**Supplementary Code:** https://github.com/WortelLab/tem1coex/

## References

1. Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, Han C, Bisignano C, Rao P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G, Kashef Hamadani BH, Kumaran EAP, McManigal B, Agarwal R, Akech S, Albertson S, Amuasi J, Andrews J, Aravkin A, Ashley E, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Bethou A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalheiro C, Castañeda-Orjuela C, Chansamouth V, Chaurasia S, Chiurchiù S, Chowdhury F, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E, Cunningham M, Darboe S, Day NPJ, De Luca M, Dokova K, Dramowski A, Dunachie SJ, Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garrett D, Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins S, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Khorana M, Kissoon N, Kobeissi E, Kostyanev T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Limmathurotsakul D, Loftus MJ, Lunn M, Ma J, Mturi N, Munera-Huertas T, Musicha P, Mussi-Pinhata MM, Nakamura T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Olivas-Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Peleg AY, Perrone C, Plakkal N, Ponce-de-Leon A, Raad M, Ramdin T, Riddell A, Roberts T, Robotham JV, Roca A, Rudd KE, Russell N, Schnall J, Scott JAG, Shivamallappa M, Sifuentes-Osornio J, Steenkeste N, Stewardson AJ, Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Turner C, Turner P, van Doorn HR, Velaphi S, Vongpradith A, Vu H, Walsh T, Waner S, Wangrangsimakul T, Wozniak T, Zheng P, Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi M. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. The Lancet 399:629–655.

2. Cosgrove SE. 2006. The Relationship between Antimicrobial Resistance and Patient Outcomes: Mortality, Length of Hospital Stay, and Health Care Costs. Clinical Infectious Diseases 42:S82–S89.

3. Bush K, Jacoby GA. 2010. Updated Functional Classification of β-Lactamases. Antimicrobial Agents and Chemotherapy 54:969–976.

4.   Barlow M, Hall BG. 2002. Predicting evolutionary potential: in vitro evolution accurately
reproduces natural evolution of the TEM β-lactamase. Genetics.

5.   Schenk MF, Zwart MP, Hwang S, Ruelens P, Severing E, Krug J, de Visser JAGM. 2022.
Population size mediates the contribution of high-rate and large-benefit mutations to
parallel evolution. Nature Ecology & Evolution.

6.   Weinreich DM. 2006. Darwinian Evolution Can Follow Only Very Few Mutational Paths to
Fitter Proteins. Science.

7.   Yurtsev EA, Chao HX, Datta MS, Artemova T, Gore J. 2013. Bacterial cheating drives the
population dynamics of cooperative antibiotic resistance plasmids. Molecular Systems
Biology 9:683.

8.   Pearl Mizrahi S, Goyal A, Gore J. 2023. Community interactions drive the evolution of
antibiotic tolerance in bacteria. Proceedings of the National Academy of Sciences
120:e2209043119.

9.   Bottery MJ, Matthews JL, Wood AJ, Johansen HK, Pitchford JW, Friman V-P. 2022. Inter-
species interactions alter antibiotic efficacy in bacterial communities. 3. ISME J 16:812–821.

10.  Medaney F, Dimitriu T, Ellis RJ, Raymond B. 2016. Live to cheat another day: bacterial
dormancy facilitates the social exploitation of β-lactamases. 3. The ISME Journal 10:778–
787.

11.  Bottery MJ, Pitchford JW, Friman V-P. 2020. Ecology and evolution of antimicrobial
resistance in bacterial communities. The ISME Journal 1–10.

12.  Amanatidou E, Matthews AC, Kuhlicke U, Neu TR, McEvoy JP, Raymond B. 2019. Biofilms
facilitate cheating and social exploitation of β-lactam resistance in Escherichia coli. 1. npj
Biofilms Microbiomes 5:1–10.

13.  Brännström Å, Johansson J, Von Festenberg N. 2013. The hitchhiker's guide to adaptive
dynamics. 3. Games 4:304–328.

14.  Vega NM, Gore J. 2014. Collective Antibiotic Resistance: Mechanisms and Implications. Curr
Opin Microbiol 21:28–34.

15.    Sorg RA, Lin L, van Doorn GS, Sorg M, Olson J, Nizet V, Veening JW. 2016. Collective

Resistance in Microbial Communities by Intracellular Antibiotic Deactivation. PLoS Biology

14.

16.    Frost I, Smith WPJ, Mitri S, Alvaro •, Millan S, Davit Y, James •, Osborne M, Pitt-Francis JM,

Maclean • R Craig, Foster KR. 2018. Cooperation, competition and antibiotic resistance in

bacterial colonies. The ISME Journal 12:1582–1593.

17.    Geyrhofer L, Ruelens P, Farr AD, Pesce D, Visser JAGM de, Brenner N. 2022. Minimal

Surviving Inoculum in Collective Antibiotic Resistance. mBio 14:e02456-22.

18.    Alan Dugatkin Ã L, Perlin M, Scott Lucas J, Atlas R. 2005. Group-beneficial traits, frequency-

dependent selection and genotypic diversity: an antibiotic resistance paradigm. Proc R Soc

B 272:79–83.

19.    Ruelens P, de Visser JAG. 2021. Choice of β-lactam resistance pathway depends critically on

initial antibiotic concentration. Antimicrobial Agents and Chemotherapy.

20.    Salverda MLM, De Visser JAGM, Barlow M. 2010. Natural evolution of TEM-1 β-lactamase:

experimental reconstruction and clinical relevance. FEMS Microbiology Reviews.

21.    Lautenbach E, Patel JB, Bilker WB, Edelstein PH, Fishman NO. 2001. Extended-Spectrum β-

Lactamase-Producing Escherichia coli and Klebsiella pneumoniae: Risk Factors for Infection

and Impact of Resistance on Outcomes. Clinical Infectious Diseases 32:1162–1171.

22.    Ramphal R, Ambrose, PG. 2006. Extended-Spectrum β-Lactamases and Clinical Outcomes:

Current Data. Clinical Infectious Diseases 42:S164–S172.

23.    Kim S, Kim J, Kang Y, Park Y, Lee B. 2004. Occurrence of Extended-Spectrum β-Lactamases

in Members of the Genus Shigella in the Republic of Korea. J Clin Microbiol 42:5264–5269.

24.    Hall BG. 2002. Predicting Evolution by In Vitro Evolution Requires Determining Evolutionary

Pathways. Antimicrobial Agents and Chemotherapy 46:3035–3038.

25.    Knies JL, Cai F, Weinreich DM. 2017. Enzyme Efficiency but Not Thermostability Drives

Cefotaxime Resistance Evolution in TEM-1 β-Lactamase. Molecular Biology and Evolution

34:1040–1054.

26. Tuomanen E, Cozens R, Tosch W, Zak O, Tomasz A. 1986. The rate of killing of Escherichia
coli by beta-lactam antibiotics is strictly proportional to the rate of bacterial growth. J Gen
Microbiol 132:1297–1304.

27. Lee AJ, Wang S, Meredith HR, Zhuang B, Dai Z, You L. 2018. Robust, linear correlations
between growth rates and β-lactam–mediated lysis rates. Proceedings of the National
Academy of Sciences of the United States of America.

28. Van Zanten ARH, Oudijk M, Nohlmans-Paulssen MKE, Van Der Meer YG, Girbes ARJ,
Polderman KH. 2007. Continuous vs. intermittent cefotaxime administration in patients
with chronic obstructive pulmonary disease and respiratory tract infections:
pharmacokinetics/pharmacodynamics, bacterial susceptibility and clinical efficacy. British
Journal of Clinical Pharmacology 63:100–109.

29. Shanmugam S, Acharya L, Mallayasamy S, Rao A, Khan SA, Thiyagu R. 2010. Study of tissue
and the plasma concentrations of cefotaxime to assess its suitability for prophylaxis in
cholecystectomy. Journal of Clinical and Diagnostic Research 4:2410–2415.

30. Dugatkin LA, Perlin M, Lucas JS, Atlas R. 2005. Group-beneficial traits, frequency-dependent
selection and genotypic diversity: an antibiotic resistance paradigm. Proc R Soc B 272:79–
83.

31. Sharma A, Wood KB. 2021. Spatial segregation and cooperation in radially expanding
microbial colonies under antibiotic stress. 10. ISME J 15:3019–3033.

32. Kim SW, Park SB, Im SP, Lee JS, Jung JW, Gong TW, Lazarte JMS, Kim J, Seo J-S, Kim J-H, Song
J-W, Jung HS, Kim GJ, Lee YJ, Lim S-K, Jung TS. 2018. Outer membrane vesicles from β-
lactam-resistant Escherichia coli enable the survival of β-lactam-susceptible E. coli in the
presence of β-lactam antibiotics. 1. Sci Rep 8:5402.

33. King ES, Maltas J, Weaver DT, Barker-Clarke R, Pelesko J, Dolson E, Scott JG. 2022. Fitness
seascapes are necessary for realistic modeling of the evolutionary response to drug therapy.
bioRxiv https://doi.org/10.1101/2022.06.10.495696.

624 34. Wright GD. 2005. Bacterial resistance to antibiotics: Enzymatic degradation and
625   modification. Advanced Drug Delivery Reviews 57:1451–1470.

626 35. Bull JJ, Harcombe WR. 2009. Population Dynamics Constrain the Cooperative Evolution of
627   Cross-Feeding. PLOS ONE 4:e4115.

628 36. Wortel MT, Bosdriesz E, Teusink B, Bruggeman FJ. 2016. Evolutionary pressures on microbial
629   metabolic strategies in the chemostat. Sci Rep 6:29503.

630 37. Blair JMA, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJV. 2015. Molecular mechanisms of
631   antibiotic resistance. 1. Nat Rev Microbiol 13:42–51.

632 38. Cartagena AJ, Taylor KL, Smith JT, Manson AL, Pierce VM, Earl AM, Bhattacharyya RP. 2023.
633   The carbapenem inoculum effect provides insight into the molecular mechanisms
634   underlying carbapenem resistance in Enterobacterales. bioRxiv
635   https://doi.org/10.1101/2023.05.23.541813.

636 39. Estrela S, Brown SP. 2018. Community interactions and spatial structure shape selection on
637   antibiotic resistant lineages. PLoS Comput Biol 14:e1006179.

638 40. Kogenaru M, Tans SJ. 2014. An improved Escherichia coli strain to host gene regulatory
639   networks involving both the AraC and LacI inducible transcription factors. J Biol Eng 8:2.

640 41. Cronan JE. 2006. A family of arabinose-inducible Escherichia coli expression vectors having
641   pBR322 copy control. Plasmid 55:152–157.

642 42. Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-term experimental evolution in
643   Escherichia coli. I. Adaptation and divergence during 2,000 generations. The American
644   Naturalist.

645 43. Schenk MF, Szendro IG, Krug J, Visser JAGM de. 2012. Quantifying the Adaptive Potential of
646   an Antibiotic Resistance Enzyme. PLOS Genetics 8:e1002783.

647 44. Wiegand I, Hilpert K, Hancock REW. 2008. Agar and broth dilution methods to determine
648   the minimal inhibitory concentration (MIC) of antimicrobial substances. Nature Protocols.

649 45. Balouiri M, Sadiki M, Ibnsouda SK. 2016. Methods for in vitro evaluating antimicrobial
650   activity: A review. J Pharm Anal 6:71–79.