Small RNAs from Mitochondrial Genome Recombination Sites are Incorporated into *T. gondii* Mitoribosomes

Sabrina Tetzlaff[1], Arne Hillebrand[1], Nikiforos Drakoulis[1], Zala Gluhic[1], Sascha Maschmann[1], Peter Lyko[2], Susann Wicke[2], Christian Schmitz-Linneweber[1]*

[1] Molecular Genetics, Humboldt University Berlin, Philippstr.13, 10115 Berlin, Germany.

[2] Biodiversity and Evolution, Humboldt University Berlin, Philippstr.13, 10115 Berlin, Germany.

* corresponding author;

Christian Schmitz-Linneweber

Humboldt-University of Berlin
· Molecular Genetics ·
Institute of Biology, Faculty of Life Sciences
Philippstr. 13, Bldg. 22
Rhoda-Erdmann-Haus
10115 Berlin
Germany
Phone: ++49-(0)30-2093-49700
www: https://www.molgen.hu-berlin.de/
email: christian.schmitz-linneweber@rz.hu-berlin.de

Running title: *Toxoplasma* small rRNAs from mitoDNA block borders

Abstract

The mitochondrial genomes of apicomplexans comprise merely three protein-coding genes, alongside a set of thirty to forty genes encoding small RNAs (sRNAs), many of which exhibit homologies to rRNA from *E. coli*. The expression status and integration of these short RNAs into ribosomes remains unclear and direct evidence for active ribosomes within apicomplexan mitochondria is still lacking. In this study, we conducted small RNA sequencing on the apicomplexan *Toxoplasma gondii* to investigate the occurrence and function of mitochondrial sRNAs. To enhance the analysis of sRNA sequencing outcomes, we also re-sequenced the *T. gondii* mitochondrial genome using an improved organelle enrichment protocol and Nanopore sequencing. It has been established previously that the *T. gondii* genome comprises 21 sequence blocks that undergo recombination among themselves but that their order is not entirely random. The enhanced coverage of the mitochondrial genome allowed us to characterize block combinations at increased resolution. Employing this refined genome for sRNA mapping, we find that many small RNAs originated from the junction sites between protein-coding blocks and rRNA sequence blocks. Surprisingly, such block border sRNAs were incorporated into polysomes together with canonical rRNA fragments and mRNAs. In conclusion, apicomplexan ribosomes are active within polysomes and are indeed assembled through the integration of sRNAs, including previously undetected sRNAs with merged mRNA-rRNA sequences. Our findings lead to the hypothesis that *T. gondii's* block-based genome organization enables the dual utilization of mitochondrial sequences as both messenger RNAs and ribosomal RNAs, potentially establishing a link between the regulation of rRNA and mRNA expression.

Introduction

Apicomplexan parasites are a group of intracellular protozoan pathogens that can cause infectious diseases, including malaria and toxoplasmosis. Toxoplasmosis is caused by *Toxoplasma gondii,* one of the most widespread human parasites, with a seroprevalence of

more than one-quarter of humans worldwide (Molan et al., 2019). For many experimental

questions, *T. gondii* remains an ideal model system for studying the molecular biology of

Apicomplexa (Szabo and Finney, 2017). Given the economic and clinical impact of *T. gondii*

and its apicomplexan relatives, there is continuous interest in distinctive biochemical and

cellular features as potential targets for therapeutic intervention.

Apicomplexan cells possess a single mitochondrion that is respirationally active and essential

for parasite survival (MacRae et al., 2012; Melo et al., 2000; Seeber et al., 1998; Vercesi et

al., 1998). With only three exceptions, all mitochondrial proteins are nuclear encoded and these

nuclear genes contribute strongly to *P. berghei* and *T. gondii* cell fitness (Bushell et al., 2017;

Sidik et al., 2016). Dozens of apicomplexan mitochondrial genomes have been sequenced

(Berná et al., 2021b). These sequences showcase the extreme reductive evolution in

apicomplexan mitochondria, setting records for the smallest mitochondrial genomes known to

date, ranging in length from 6 to 11 kb (Hikosaka et al., 2013; Oborník and Lukeš, 2015).

Mitochondrial genome organization is evolutionarily very variable, with some species having a

linear, monomeric genome (*Babesia* spp.), while others have concatenated arrays of genomes

(Hikosaka et al., 2013). The *T. gondii* mitochondrial genome is composed of 21 repetitive

sequence blocks that are organized on multiple DNA molecules of varying lengths and non-

random combinations (Berná et al., 2021a; Namasivayam et al., 2021). Apicomplexan

mitogenomes code only for three proteins (COB, COX1, COXIII) and have highly fragmented

genes for rRNAs (Feagin et al., 2012; Seeber et al., 2020). For example, in *P. falciparum*, 34

genes for rRNA fragments are scattered across the mitochondrial DNA on both strands of the

genome without any linear representation of the full-length large or small subunit rRNA (Feagin

et al., 2012). How the multitude of rRNA fragments are assembled into functional

mitoribosomes in *T. gondii* remains unknown.

Since the electron transport chain in mitochondria is essential for the survival of

apicomplexans, the expression of mitochondrial genes is likely to be essential as well. In fact,

a nuclear-encoded RNA polymerase targeted to mitochondria has been shown to be essential

in the blood stages of *Plasmodium spp.* (Ke et al., 2012; Suplick et al., 1990). Two

mitochondrially localized RNA binding proteins from the RAP (RNA-binding domain abundant

in apicomplexans) family are also essential for the survival of *P. falciparum*, although their role

in RNA metabolism has not yet been determined (Hollin et al., 2022). In addition, the depletion

of nuclear encoded mito-ribosomal proteins of *T. gondii* (Lacombe et al. 2019; Shikha et al.

2022) and *P. falciparum* (Ke et al. 2018) led to defects in the assembly of ETC complexes and

in parasite proliferation, suggesting that mitochondrial translation is important for parasite

survival. Resistance to the antimalarial drug atovaquone in *P. falciparum* and *T. gondii* has

been linked to mutations in the *cob* (*cytB*) gene of mitochondria (McFadden et al., 2000;

Srivastava et al., 1999; Syafruddin et al., 1999), further supporting the idea of active, essential

translation in apicomplexan mitochondria.

Despite the clear importance of mitogenome expression in Apicomplexa, we lack a

comprehensive catalog of transcripts and processing events. Genome-length, polycistronic

transcripts are produced in *Plasmodium* that encompass all three open reading frames (Ji et

al., 1996). The existence of full-length mRNAs was further confirmed by cDNA amplifications

(Namasivayam et al., 2021) and long read sequencing of mRNAs in both *Plasmodium* and

Toxoplasma (Lee et al., 2021; Namasivayam et al., 2021). There is no evidence of dedicated

transcription initiation for individual protein-coding genes, and it is assumed that mRNAs are

processed from the polycistronic precursor (Feagin et al., 2012; Hillebrand et al., 2018;

Rehkopf et al., 2000; Suplick et al., 1990), but the mechanism of processing and the required

machinery are unknown. Similarly, the production of rRNA fragments is achieved post-

transcriptionally by processing long precursor RNAs (Ji et al. 1996). A catalog of small RNAs,

mostly rRNA fragments, was described for *P. falciparum* based on small RNA sequencing

(Hillebrand et al., 2018), but in other apicomplexans, including *T. gondii*, our understanding of

rRNA species is limited and mostly based on predictions through sequence comparisons with

Plasmodium (Feagin et al., 2012; Namasivayam et al., 2021).

We used a slightly modified protocol of mitochondria enrichment (Esseiva et al. 2004) to

investigate the structure of the mitochondrial genome through long-read sequencing at an

unprecedented depth. In parallel, we conducted high-throughput sequencing of small RNAs

(<150 nt) and demonstrated that they are incorporated into polysomes. The combination of

DNA sequencing results and transcriptome analysis also allowed us to identify previously

undetected transcripts, many of which originate from block boundaries and represent fusions

of coding and noncoding regions.

Results

***T. gondii* organellar nucleic acids are enriched by a combination of selective membrane**

disruption and degradation of nucleo-cytosolic DNA and RNA

The *T. gondii* nuclear genome contains many insertions of mitochondrial DNA sequences

(Gjerde, 2013; Namasivayam et al., 2023; Ossorio et al., 1991). To better distinguish between

NUMTs (nuclear DNA sequences that originated from mitochondria) and true mitochondrial

sequences, it is helpful to enrich mitochondrial DNA. We modified a cell fractionation method

that takes advantage of the differential sterol content in plasma membranes and organellar

membranes (Esseiva et al., 2004; Subczynski et al., 2017). We incubated cells with the

detergent digitonin, which selectively permeabilizes sterol-rich membranes, leaving

mitochondria and other organelles intact. We used a *T. gondii* strain that constitutively

expresses GFP localized to the mitochondrial matrix to track the mitochondria during the

purification process (Figure 1A). After permeabilizing the plasma membrane with digitonin, we

treated the cells with benzonase to degrade nucleo-cytosolic nucleic acids. Following

inactivation of the benzonase, we lysed the mitochondria with high concentrations of

detergents to release the soluble content.

We observed that mitochondrial GFP remained in the pellet fraction (P in Figure 1B) with only

a minor signal in the supernatant, indicating that the mitochondria remained largely intact. In

contrast, cytosolic HSP70 largely remained in the supernatant after digitonin and benzonase

treatment, indicating that the method successfully reduced cytoplasmic proteins. Upon lysis of

the mitochondria using Triton X-100 ("Lysis" in Figure 1B), the GFP signal was predominantly

recovered in the supernatant, demonstrating efficient lysis of the mitochondria and release of

soluble components. This fraction still contains HSP70, but slight contamination with cytosol is

expected. We also found that cytosolic 18S and 26S rRNA were almost completely degraded,

indicating the high efficacy of the benzonase treatment (Figure 1C). In summary, our enhanced

protocol for organellar enrichment effectively depletes cytosolic protein and RNA components

and enriches mitochondrial macromolecules.

To assess the integrity of mitochondrial RNA following our protocol, we utilized small RNA gel

blot hybridization for two known mitochondrial rRNA fragments, SSUD and LSUF

(Namasivayam et al., 2021). Our analysis of total nucleic acids in the gel revealed effective

removal of major rRNA species by benzonase treatment (Figure 1C), while smaller

degradation products accumulated (Figure 1D lanes "Benz" and "Lysis"). In contrast, we

observed clear bands for both SSUD and LSUF, indicating their protection from benzonase

degradation. Notably, we also detected signals for both rRNA fragments in the supernatant of

the lysis fraction, confirming successful mitochondrial lysis and recovery of mitochondrial rRNA

fragments.

Figure 1: An experimental pipeline to enrich *T. gondii* mitochondrial RNAs.

A)   *T. gondii* tachyzoites were harvested and incubated in a buffer with digitonin for plasma
membrane permeabilization. Subsequently, accessible nucleic acids were digested by benzonase. After
removal of the benzonase and washing the pellet, the intact organelles were lysed by a high detergent
buffer. Soluble nucleic acids were separated by centrifugation and extracted from the supernatant.
B)   *T. gondii* cells expressing a mitochondrial-targeted GFP were subjected to organelle
enrichment. GFP was tracked by immunoblotting (5% volume of each fraction was analyzed). The GFP
signal remained in the pellet and only shifted into the supernatant after the lysis step. In contrast, much
of the cytosolic HSP70 is removed during the procedure, indicating specific enrichment of mitochondria.
C)   RNA extracted from selected fractions of the organelle enrichment protocol was analyzed by
agarose gel electrophoresis.
D)   RNA extracted from selected fractions of the organelle enrichment protocol was analyzed on a
denaturing 10% PAGE gel and blotted onto a nylon membrane. Radiolabeled DNA oligonucleotide
probes were used to detect the mitochondrial rRNA fragments SSUD (from the small subunit of the
ribosome) and LSUF (large subunit of the ribosome). Both fragments were found in pellet fractions after
digitonin treatment, where they were protected from benzonase digestion. They only shifted to the
supernatant after lysis. This demonstrates that the mitochondria stay intact during the procedure.
166

Long-read sequencing of DNA from enriched mitochondrial fractions identifies prevalent sequence block combinations

For the analysis of mitochondrial sRNAs, we needed a reference mitochondrial genome that

would represent the existing sequence block combinations. A previous sequencing study used

Oxford Nanopore sequencing technology (ONT) to identify combinations of sequence blocks

in *T. gondii* mitochondria (Namasivayam et al., 2021). To improve read depth, we performed

ONT sequencing on DNA gained from our improved organelle enrichment protocol. After

filtering against the nuclear genome, we found 86,761 reads that had similarities to the

published mitochondrial genome sequence (Namasivayam et al. 2021). This corresponds to

78.5 Mbp, which accounts for 9.2% of the total sequences that we attribute to the nucleus and

mitochondria (Tab. S1). This is a 42-fold increase in the sequencing depth of the *T. gondii*

mitochondrial genome compared to previous attempts (Tab. S1), which can be attributed to

the effectiveness of the purification process. The length of mitochondrial reads ranged from 87

nt to 17,424 nt (Tab. S2). The GC content of these reads, on average 36.4%, further supports

their mitochondrial origin (Berná et al., 2021a). We annotated the reads using published

sequence block information (Namasivayam et al., 2021) and confirmed the previously

described 21 sequence blocks, designated by letters from A-V (Namasivayam et al., 2021).

We searched the sequencing data for repeated sequence elements but found no additional

blocks, suggesting that the *T. gondii* mitochondrial genome is fully covered. We compared our

reads with published ONT reads for the *T. gondii* mitochondrial genome (Berná et al., 2021a;

Namasivayam et al., 2021). While smaller reads of our dataset are found in full within longer

reads in the published datasets, we do not find any examples for reads that would be full

matches between the dataset. This suggests that while the block content is identical between

the different sequencing results, the actual block order was reshuffled by recombination,

indicating the rapidity of mitogenome evolution in *T. gondii,* thus corroborating at a deeper

sequencing coverage conclusions so far gained from a more limited read set (Berná et al.,

2021a; Namasivayam et al., 2021).

The depth of our sequencing enabled us to quantify the genome block combinations. We first

asked whether there is a bias in sequential block succession. We observed that the rate of

occurrence of blocks varies dramatically in our dataset, with the most frequent block, J,

occurring 45,849 times and the least frequent, C, occurring 3,138 times (Figure 2A, Tab. S3).

This suggests that differences in block combinations can be expected. After counting block

combinations, we found that only a small fraction of all possible block combinations are

prevalent within the genome (Figure S1A, Tab. S4), confirming block combinations observed

previously (Namasivayam et al. 2021) and adding combination frequencies based on higher

read numbers. For example, the most frequent block combination is J-B, which occurs 19,622

times in our reads. In total we identified 84 combinations of which 52 occur less than fifty times

and make up less than 0.06% of the total number of combinations found (Tab. S4). Fourteen

blocks (R, N, M, F, L, U, I, C, K, D, H, E, T) are always found with the same preceding and

following block. The direction of the blocks in these combinations is fixed (for example, the 3'-

end of the L block is always fused to the 5'-end of the J block, never to the 3'-end of the J-

block). This indicates that the genome's flexibility is limited and that not all block combinations

are realized (see also Namasivayam et al. 2021). Variability in genome structure is caused

primarily by blocks S, O, Q, B and P, which have up to three possible neighbors, and by blocks

Fp, Kp, V, A, and J, which have more than three possible neighbors. All combinations are well

covered in our ONT results and helped to refine block borders relative to previous annotations

(Figure S2). Using the 32 high-frequency block combinations we found, we generated a map

centered on the protein-coding genes (Figure 2B). We will later show that all blocks not

encoding proteins express rRNA fragments and are thus called rRNA blocks here. The map is

not designed to represent physically existing DNA molecules, since the sequencing technology

used cannot detect branched or circular DNAs, nor are the blocks drawn to scale. However, it

helps to indicate the complexity of the nonrandom recombination events shaping the genome.

For example, when considering the positioning of the three protein-coding genes within this

map, it is conspicuous that their three 5'-ends are always flanked by blocks also encoding a

protein (Figure 2B), although it is unclear whether this is of functional relevance.

Full-length coding regions had been found previously on nanopore reads in *T. gondii*

(Namasivayam et al., 2021). In our improved representation of the mitogenome, we identified

a large number of full representations of *cob*, *coxI* and *coxIII* in our dataset (*cob*: 1,612; *coxI*:

1,404; *coxIII*: 1,487). Thus, of all reads long enough to carry one of the following ORFs, 5.1%

contain full-length coxIII, 8.6% contain full-length cob, and 11.1% full-length cox1, respectively

(Tab. S5). This may be adequate for the expression of the encoded proteins; however, we

cannot currently exclude the possibility of genomic or post-transcriptional block shuffling that

could lead to more complete open reading frames (as discussed in Berná et al., 2021b). We

next applied approaches established previously to represent biased recombination events

based on alternative block combinations (see Figure S12 in Namasivayam et al. 2021) to our

improved ONT read set. For example, block S, as part of the *coxI* coding region, was followed

in 80% of its occurrences by the rRNA block R and only in 20% by the next coding block C

(Figure 2B, Figure S1B). Similarly, block B, as part of the *coxIII* reading frame, was followed in

66% of all cases by the rRNA block U and less frequently by the terminal block of *coxIII*, which

is block M. Whether such fusions of coding and non-coding block is of functional relevance,

was unclear so far.

238
239

Figure 2 Long-read sequencing shows a specific system of block combinations.

(A) ONT DNA sequencing results were analyzed for the number of block occurrences. The gray scale
indicates how many times each of the mitochondrial sequence blocks was found in the mitochondrial
ONT reads
(B) Map of the *T. gondii* mitochondrial genome representing block combinations occuring in
mitochondrial ONT reads. The thickness of the connecting lines indicates the absolute number of
occurrences for each combination.
247

Small RNA sequencing identifies a comprehensive set of mitochondrial rRNA fragments

For *T. gondii* mitochondria, rRNA species were previously predicted based on sequence comparisons to *Plasmodium* rRNA sequences (Namasivayam et al., 2021). We used Illumina-based small RNA sequencing of *T. gondii* to identify mitochondrial rRNA fragments. We sequenced both the mitochondria-enriched fraction and the input fractions described in Figure 1. In the enriched fraction, we mostly retrieved reads for cytosolic rRNA degradation products and poor coverage of mitochondrial sequences (not shown). This is likely caused by the benzonase step in the protocol, which generates an abundance of small degradation products of cytosolic rRNA (Figure 1D; S3). In the input samples, intact full-length rRNAs are removed by selecting for small RNAs during library preparation, resulting in less prevalent cytosolic rRNA reads. We bioinformatically removed the remaining cytosolic rRNA reads as well as any reads shorter than 20 nt. Poly-A stretches at the 3'-ends were clipped as well. Finally, we mapped the remaining reads against the mitochondrial genome (Tab. S6) using the block combinations identified here by ONT sequencing (Figure 2).

The results of our mapping study show that the 5'-termini of mitochondrial small RNAs are clearly defined, making it easy to identify their transcript ends (as shown in Figure 3A). In agreement with findings from previous studies on small RNAs from *P. falciparum* (Hillebrand et al., 2018), we found that the 3'-ends of many transcripts were more variable (as seen in the example of RNA17 in Figure 3A). We identified a total of 34 small RNAs that accumulate in *T. gondii* mitochondria (Tab. S7). Among these, 11 correspond to previously predicted rRNA fragments (Namasivayam et al., 2021). Our sequencing data confirmed that these sequences are expressed and allowed us to refine the exact rRNA fragment ends (Tab. S7). Additionally, our small RNA sequencing revealed some larger differences compared to previous transcript predictions. This included a reassignment of SSUF to the opposite strand and also affected the four rRNA fragments LSUF, LSUG, LSUD, and LSUE, which had been predicted as separate transcripts of the large subunit (LSU) of the ribosome (Namasivayam et al., 2021). Our sequencing results suggest that there is accumulation of transcripts containing LSUF and

LSUG regions and LSUD and LSUE regions, respectively. Both of these transcripts were

verified via northern blotting (Figure S4). The longer transcripts were found to be much more

abundant than smaller transcripts that were also detected (Figure S4), suggesting that the

longer transcripts represent the functional rRNA fragments in *T. gondii* mitochondria. Knowing

the exact sequence of rRNA fragments is crucial for further investigations into the structure of

mitoribosomes, as it is key for predictions of secondary structures.

Figure 3: Small RNA sequencing identifies mitochondrial small RNAs.

(A)   Upper three rows: excerpt of mapping results in three strand-specific small RNA sequencing
replicates starting from total input RNA from our *T. gondii* organelle RNA preparation. Read depth was
counted by the number of reads at each position. Note that the 5'-ends of the RNAs shown here are on
the right. Lower four rows: only the terminal nucleotides of each read on the plus and minus strand of
the genome are shown. The size and position of genes below the coverage graphs are all drawn to
scale. Red bars with lowercase letters indicate positions of probes used for detecting RNAs in RNA gel
blot hybridizations shown in (B).
(B)   Equal quantities of total input RNA from *T. gondii* and from organelle preparations were loaded
onto a denaturing PAGE gel and analyzed by RNA gel blot hybridization with the probes indicated.

Discovery of previously undetected rRNA fragments

sRNAs cover most of the noncoding sequence blocks of the T. gondii *genome*

Out of the 34 small RNA fragments identified, 23 have not been described previously for *T.*

gondii (see Tab. S7 and Figure S5). 17 of the 23 are homologs of sRNA fragments in the

apicomplexans *P. falciparum* and *E. leuckarti* (marked in bold in Table S7*)*. These were named

according to their *Plasmodium* homologs (Feagin et al., 2012). Six further sRNAs are

exclusively conserved within cyst-forming Eucoccidians, the closest apicomplexan relatives of

T. gondii (last column in Table S7). We assigned numbers to these sRNAs extending the

Plasmodium nomenclature (Feagin et al. 2012). We next asked, how many of all 34 sRNAs

have homologies to rRNA from *E. coli*. We could find twelve LSU homologs and eleven SSU

homologs in accordance with previous analyses in *P. falciparum* (Feagin et al. 2012; Hillebrand

et al. 2018 ; see Tab. S8). Only for the *P. falciparum* LSU fragment LSUC and the SSU

fragment RNA12, we were unable to identify corresponding homologs in *T. gondii*.

We next analyzed the accumulation of selected RNAs previously undetected in *T. gondii* for

the sequence blocks Kp-K: Transcripts RNA5, 17, and 29, as well as the already predicted

transcripts RNA10 and SSUD are found on the minus strand (strandedness according to

(Namasivayam et al., 2021; MN077088.1 - MN077111.1). They are separated by single

nucleotides in the mitochondrial genome. In this and most other noncoding blocks, the DNA

sequence is almost fully used for transcript production (see Figure 3A, Figure S5, S6). To

validate the sequencing data, we performed RNA gel blot hybridization using probes against

RNA17, RNA29, and RNA5 (Figure 3B) and indeed detected transcripts of the expected size

(41 nt, 42 nt and 83 nt, respectively). The transcripts were more abundant in our organelle

preparations than in input samples, which indicates that they are of organellar origin and did

not originate from NUMTs. The sequencing and RNA gel blot efforts demonstrate that almost

the entire sequence of the blocks Kp-K is represented by small RNAs.

Dual use of block borders for sRNA production

Among the small RNAs identified here, there is also a class that was only detectable due to

our insights into genome block combinations. Using block combinations for mapping analysis,

we identified 15 transcripts that span two blocks, i.e. there are sRNAs at almost half of all 31

block borders. Often, we find sRNAs that share a block border sequence at one end, but differ

at the other end depending on the block combination (see Figure 4A, orange boxes). An

example of sequences shared at the 5'-end is represented by the pair RNA8 and RNA31, which

both start in block Fp but end in different blocks (Figure 4A). An example of block sharing at

the 3'-end is the RNA pair RNA1/RNA2. Both RNAs terminate at very similar positions in block

Kp but start in different blocks (Figure 4A). Using a probe for the common 3'-part in block Kp,

we detected both RNA1 and RNA2 in an RNA gel blot hybridization experiment (Figure 4A, B).

Both RNA1 and RNA2 have homology to *E. coli* LSU in their 3'-portion via block Kp, but only

RNA1 has additional homology to *E. coli* LSU via block I (Figure 4C; Feagin et al., 2012). It is

interesting to consider that RNA1 could therefore be used in two positions in the large subunit

of the ribosome. In conclusion, block combinations can lead to the expression of RNAs in *T.*

gondii that are not found in apicomplexan species with a simpler genome organization (Table

S7).

Coding regions contribute to sRNAs at block borders

In addition to the three pairs of overlapping, noncoding RNAs described above (Figure 4A), we

also found five RNAs that combine sequences from coding and non-protein-coding blocks

(Figure 5A, B). Among them, RNA16, RNA23t, and RNA34 were partially antisense to mRNAs

(Figure 5A), and we confirmed the accumulation of RNA34 by RNA gel blot hybridization

(Figure 5C). None of the three RNAs had detectable homologies to E. coli rRNA based on

simple sequence searches, but structural conservation cannot be ruled out. Mitochondrial

rRNA sequences from kinetoplastids and diplonemids show very little sequence conservation

but are still part of the mitoribosome (Valach et al. 2023; Ramrath et al. 2018), suggesting that

future analyses might uncover hidden rRNA similarities. With the exception of RNA34, the

RNAs representing a fusion of non-protein-coding regions and protein-coding regions have

homologous sequences in the mitochondrial genome of *P. falciparum* (Feagin et al. 2021). In

P. falciparum however, none of them are antisense to coding regions. Whether these sRNAs

form sRNA:mRNA interactions and whether this has functional consequences remains to be

investigated.

Two of the small RNAs contained coding sequences in the sense direction (Figure 5B). Both

were validated by RNA gel blot hybridization (Figure 5D). RNA19 contains the *coxIII* coding

sequence adjacent to the *cob* coding region by a combination of block J and block E (Figure

5E). *Cob* has two in frame start codons (Figure 5F). Using rapid amplification of cDNA ends

(RACE), we showed that the 5'-end of the *cob* mRNA is 6 nt upstream of the second start

codon and 10 nt downstream of RNA19 (Figure 5 F,G). Being only four nucleotides apart, the

3'-end of RNA19 is possibly generated during the formation of the 5'-end of *cob*. RNA19 is

conserved in other apicomplexan mitochondrial genomes (Figure 5E) but is not part of open

reading frames in *Plasmodium*. Homologies and a structural fit to the SSU rRNA of *E. coli* have

been noted for *P. falciparum* RNA19 (Figure 5E, Feagin et al. 2012). This sequence similarity

to rRNA is maintained in *T. gondii* (Figure 5E), which suggests that despite overlapping with

coxIII coding sequence at the J-E block border, RNA19 is functional. It is remarkable that this

sequence serves dual purposes for coding a protein and an rRNA. There is only low sequence

conservation of the c*oxIII* sequence used by RNA19 (Figure 5F), which might have allowed

the evolutionary acquisition of an additional use as an rRNA fragment. In addition to RNA19,

a second RNA encompassing coding sequences is RNA3, which is situated at the S-R block

border (Figure 5B). Block S is part of the *coxI* coding region - its terminal 22 nucleotides

contribute to RNA3. The similarity to E. coli 23S rRNA is, however, restricted to the noncoding

block R (Figure S7). The Block S sequence in RNA3 is therefore a 5'-extension of this rRNA,

and it remains to be determined whether this is of functional consequence for the mitochondrial

ribosome. It is noteworthy that the 5' end of RNA3 is located within a region of the 23S rRNA

secondary structure that exhibits low conservation (Feagin et al. 2012), which may suggest

that its overlap with coding regions is potentially tolerable.

In sum, these results suggest that at block borders, *T. gondii* makes dual use of several protein-

coding sequence blocks. Depending on the respective block combination, they are either part

of the protein coding sequence or can code for an rRNA fragment.

Figure 4: Noncoding RNAs at block borders.

A)   Schematic representation of three block borders that are used by different transcripts (sequences used for two transcripts are marked by orange rectangles). Transcripts are shown as wiggle lines and are named according to prior nomenclature (Namasivayam et al., 2021). Assignment of transcripts to the large and the small rRNA are indicated by color. The red bars with the lower-case letter "j" indicate the position of the probe used for detecting RNA1 and RNA2 by RNA gel blot hybridization. LSU/SSU= large/small subunit of the ribosome.

B)   RNA gel blot hybridization of RNA1 and RNA2 using RNA extracts from input and organelle-enriched fractions of the organelle enrichment protocol shown in Figure 1. Equal RNA amounts were loaded.

C)   Alignment of RNA1 (top) and RNA2 (bottom) sequences from different apicomplexan species. Sequence excerpts from *E. coli* 23S rRNA (1240-1301 for RNA1, 944-1010 for RNA2) corresponding to sections of the alignment were added manually based on previous analyses (Feagin et al. 2012). Nucleotide positions conserved in all apicomplexans are shaded in gray. The residues in the *P. falciparum* sequence suggested to replace sequence elements in the *E. coli* 23S rRNA structure (Feagin et al., 2012) are marked in bold. Residues conserved between apicomplexans and *E. coli* are marked with asterisks. Note that RNA1 and RNA2 are aligned according to their shared sequence in genomic block Kp.

Fig 5

394
395
Figure 5: Small RNAs that consist of coding and non-protein-coding sequences at block borders.

A)   Schematic representation of the genomic position of three RNAs that are partially antisense to
coding sequences.
B)   Schematic representation of the genomic position of two RNAs that partially overlap with coding
sequences.
C)   RNA gel blot hybridization of RNA34. Equal amounts of RNA extracts from input and organelle-
enriched fractions were analyzed.
D)   RNA gel blot hybridization of RNA19 and RNA3 – for details see (C)
E)   Alignment of *T. gondii* RNA19 with homologous sequences from other apicomplexan species
and from *E. coli.*
F)   Alignment of a section of the *coxIII* and *cob* sequences from different apicomplexan species.
Alignments were prepared using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/).
Nucleotides of *T. gondii* RNA19 are marked in bold in both alignments and as a lilac wiggle line above
the sequences. The sequences are interrupted by indels in *coxIII* and *cob* of other apicomplexans that
have stand-alone RNA19 sequences (e.g. *P. falciparum*). Within the *cob* sequence, the RNA19
sequence covers an annotated start codon (Alday et al., 2017, yellow box; Namasivayam et al., 2021).
However, an alternative start codon downstream has been suggested as well (red box, McFadden et al.

2000) and is in fact where sequence conservation starts. The green circle indicates the *cob* 5' transcript
terminus as determined by RACE (see G).
G)   Agarose gel analysis of 5' RACE PCR products performed to identify the *cob* 5' transcript end.
+RT and-RT indicates presence or absence of reverse transcriptase, respectively.
417

Mitochondrial small RNAs are part of a large-molecular weight complex

The apicomplexan mitochondrial ribosome with its peculiar organization has not been

characterized in any detail. There is strong evidence from genetic and pharmacological studies

that translation occurs in the mitochondria of apicomplexans (Alday et al., 2017; Lane et al.,

2018; McFadden et al., 2000; Vaidya et al., 1993). Additionally, the presence of a large

Megadalton complex containing nuclear-encoded ribosomal proteins targeted to mitochondria

has been shown by blue native gel electrophoresis (Lacombe et al., 2019). However, a

comprehensive catalog of mitoribosomal constituents is missing, and the extent to which rRNA

fragments are part of mitoribosomes remains unanswered, nor have polysomes been identified

as a direct readout for active ribosomes. To ascertain the presence of sRNAs within high-

molecular weight complexes, we fractionated enriched organellar preparations using sucrose

gradient analysis. We employed conditions known to preserve mitochondrial ribosomes (Waltz

et al., 2021a)conditions established to dissociate ribosomes (10 mM EDTA, no $MgCl_2$, 300 mM

KCl). Should sRNAs be integral to ribosomes, their dissociation would be expected to result in

a migration towards lower molecular weight fractions.

RNA extracted from the gradient fractions was initially analyzed using Ethidium bromide-

staining in denaturing polyacrylamide urea gels (Figure 6A). The paucity of detectable signals

aligns with the expected degradation of abundant cytosolic rRNAs during preparation with

benzonase. Notably, only the top fractions of the gradient (fractions 1-6) displayed bands, while

the deeper fractions showed no discernible signals, reaffirming the effectiveness of the

organellar preparation in eliminating cytosolic contaminants. The observed bands in the top

fractions may represent degradation products of prevalent rRNA species, released from

ribosomes by benzonase but persisting in the organellar preparation due to their abundance.

Subsequent to gel electrophoresis, we transferred the RNA onto membranes and hybridized

them with probes targeting seven specific sRNA species. LSUD/E and LSUF/G, along with RNA1, RNA2 and RNA3, were chosen for analysis due to their sequence similarities with E. coli 23S rRNA, as noted by Feagin et al. (2012). Similarly, SSUA and RNA19 were chosen since they resemble parts of E. coli 16S rRNA (Feagin et al., 2012). Additionally, RNA3 and RNA19 were tested as they utilize the coding sequences of coxI and coxIII, respectively. Finally, we also included RNA29, an sRNA not yet assigned to the ribosome based on sequence similarity.

We observed the peak signals for all sRNAs in the middle of the gradient, but specifically not in fractions 1-4, where free sRNAs would be expected to localize, akin to the signals observed in the ethidium bromide-stained samples (Figure 6). Secondly, all tested sRNAs were found to also migrate into the deepest gradient fractions 12-14. Collectively, these observations strongly indicate that the sRNAs in question are predominantly part of high molecular weight complexes. To evaluate the influence of ribosome-dissociative conditions on the fractionation of sRNAs in sucrose gradients, we quantified the sRNA signals in the deeper fractions (10 to 14) compared to the peak fractions (7 to 9 for LSU rRNAs and 6 to 9 for SSU rRNAs). For all sRNAs examined, a shift was observed from the deeper to the earlier fractions under dissociative conditions (Figure 6). This shift indicates that the large molecular weight complexes, with which these sRNAs are associated, are susceptible to ribosome-dissociative conditions. The most parsimonious conclusion is that all sRNAs tested are rRNAs.

We then asked whether rRNA fragments with homology to E. coli 23S versus those with homology to 16S rRNA exhibit differential signal distribution. We found that rRNAs presumed to be small subunit rRNAs, namely SSUA and RNA 19, displayed signals in fraction 6 (Figure 6D,G), a pattern not observed or less pronounced in presumed LSU rRNAs such as LSUD/E, LSUF/G, RNA1+2, and RNA3 (Figure 6B,C,E,F). The SSU-specific signal in fraction 6 persisted under both standard and dissociative conditions, implying that it represents free SSU (Figure 6D,G)). We quantified the signals in all fractions under dissociative conditions and

compared the profiles of sRNA signals across the gradient (Figure S8). This shows that under

dissociative conditions, the distribution and peaks differ markedly between the four predicted

LSU rRNAs and the two SSU sRNAs (Figure S8). The distinct distribution patterns for rRNAs

with homology to LSU versus SSU rRNA suggest the gradient allows to assign individual rRNA

fragments to the two subunits of the ribosome. Indeed, the previously unclassified RNA29

exhibited a distribution pattern very similar to LSU rRNAs (Figure 6H). It showed significantly

less signal in fraction 6 than SSUA and a distribution across the gradient similar to LSUF/G

and LSUD/E (Figure 6C,D and Figure S8). This pattern indicates that RNA29 likely belongs to

the LSU category. Unlike most other rRNAs, RNA29 also showed strong signals in the low

molecular weight fractions 1-5 (Figure 6H), hinting at a proportion of unassembled RNA29 in

our preparation. Such un-assembled rRNA was also observed for RNA2, while the cognate

RNA1 was barely detected in lower molecular weight fractions (Figure 6E). It remains to be

determined whether different assembly kinetics or degradation rates contribute to the RNA

signals in the upper parts of the gradient. Both RNA1 and RNA2 demonstrated deep

penetration into the gradient, and showed a gradient distribution consistent with the LSU

distribution pattern. Both RNA1 and RNA2 are sensitive to EDTA treatment. Together this

leads to the conclusion that RNA1 and RNA2 are both assembled into ribosomes.

We furthermore observed that despite containing sequences derived from mRNAs, both

RNA19 and RNA3 are incorporated into ribosomes (Figure 6B). For RNA19, several transcripts

were found. The largest isoform is mainly found in low-molecular weight fractions. Conversely,

the smaller RNA19 transcripts are observed to migrate into the high-molecular weight fractions,

which is EDTA-sensitive (Figure 6G). Our findings suggest that RNA19 and RNA3 are

incorporated into the ribosome.

Mitochondrial mRNAs show association with high-molecular weight complexes that are

sensitive to ribosome-dissociative conditions

We next analyzed the distribution of mitochondrial mRNAs in a sucrose gradient, under non-

dissociative and dissociative conditions. The extended length of these mRNAs facilitates their

detection using qRT-PCR, a method that is impractical for the analysis of many shorter sRNAs.

To ensure the gradient exhibited characteristics akin to those used in the sRNA distribution

analysis (Figure 6), we concurrently analyzed the sRNA LSUF/G. This confirmed that LSUF/G

migrates deep into the gradient, which is sensitive to ribosome-dissociative conditions (see

Figure 6C, 7A).

All three mRNAs migrated deep into the gradient, reaching their peak in fractions 11 to 13, as

depicted in Figure 7B-D. Upon treatment with EDTA, the mRNAs shifted towards fractions

associated with lower molecular weights. We conclude that mitochondrial mRNAs in *T. gondii*

are populated with ribosomes, which directly supports previous suggestions of active

translation in apicomplexan mitochondria.

The nuclear-encoded mitoribosomal protein L11 is found in EDTA-sensitive large

complexes

We next sought to analyze how a ribosomal protein is distributed in our sucrose gradient with

the goal to have an additional read-out for ribosomes and polysomes and to investigate

whether this mirrors the distribution of sRNAs. We chose TGRH88_003980_t1, which was in

previous *in silico* screens identified as potential mitoribosomal protein L11 (Gupta et al., 2014;

Lacombe et al., 2019). We used CRISPR/Cas9 to introduce an HA-tag to the C-terminus of

*Tg*uL11m by homologous recombination (Figure S9A,B). The tagged protein is

immunologically detected as a single band, demonstrating specificity of the tagging experiment

(Figure S9C). We next performed an immunofluorescence assay (IFA) to check the localization

*Tg*uL11m:HA. The HA-signal co-localized with the mitochondrial marker protein *Tg*Tom40 (van

Dooren et al., 2016) confirming the localization of *Tg*uL11m to *T. gondii* mitochondria (Fig 8A).

We next analyzed the distribution of *Tg*uL11m:HA in sucrose gradients and found a main peak

in fractions 9-11 with signals reaching into the deepest fractions of the gradients (Fig 8B). This

mirrors the distribution of sRNAs from the large subunits like LSUD/E (Fig 6B). Upon EDTA

treatment, the signal shifts from the deepest fractions (12-14) into lower fractions (8-10),

suggesting that we observe polysomes in fractions 11-14 and that the large subunit of the

ribosome is found in fractions 8-10.

Figure 6: Sucrose density centrifugation analysis demonstrates the association of mitochondrial small RNAs with large molecular weight complexes.

T. gondii organelle-enriched extracts were fractionated by sucrose density gradient centrifugation and subsequently analyzed by RNA gel blot hybridization. Buffer conditions during the experiment were either 30 mM MgCl$_2$, 100 mM KCl (labeled Mg^{2+} next to the blots) or 10 mM EDTA, 300 mM KCl (labeled EDTA next to the blots). (A) Prior to blotting, the gels were stained with ethidium bromide. Atop the gel images, a schematic representation illustrates the sucrose density gradient. Subsequent panels display a series of RNA gel blot hybridization assays. These assays were conducted on blots derived from the two ethidium bromide-stained gels presented in (A). Following each probe, the blot was stripped of its signal employing a denaturing buffer. The oligonucleotides in these assays targeted LSUD/E (B), LSUF/G (C), SSUA (D), RNA1 + RNA2 (E), RNA3 (F), RNA19 (G), and RNA29 (H). Percentages in square brackets encircling each blot indicate the proportion of signal detected in fraction 7-9 or 6-9 for LSU or SSU sRNAs, respectively, versus signal intensity in fraction 10-14. "I" corresponds to 10% of the lysate prior to gradient loading. The RNA names are color-coded to indicate homology: deep purple for RNAs with homology to *E. coli* rRNA from the large ribosomal subunit, light purple for those from the small subunit, and black for RNAs without previously noted homology.

Figure 7: Sucrose density centrifugation analysis of mitochondrial mRNAs.

T. gondii organelle-enriched extracts treated with standard salt concentrations or with an Mg^{2+}-depleted buffer, respectively, were fractionated by sucrose density gradient centrifugation and subsequently analyzed by RT-qPCR. Data was normalized using human ESR1 as an external spike-in control. Relative quantities per amplicon are visualized as percentage of RNA per fraction. All three mRNAs (shown in panel B-D) and one LSU rRNA fragment serving as a control (LSUF/G shown in panel A) were analyzed from a single sucrose gradient fractionation. The gradient density is shown schematically below the graphs.

Figure 8: Analysis of the putative mitoribosomal protein L11.

(A) *T. gondii* parasites expressing *Tg*uL11m:HA were subjected to immunofluorescence assays. These assays were conducted using an anti-HA antibody, visualized in the green channel. The mitochondria within these samples were identified using an antibody against the outer membrane protein Tom40, shown in the magenta channel. (B) For the analysis of HA-tagged mitochondrial ribosomal protein L11 (*Tg*uL11m:HA), sucrose density centrifugation was employed. Organelle-enriched extracts from *T. gondii* lines were either untreated (Mg^{2+}) or treated with EDTA. Subsequently, these extracts were fractionated via a sucrose gradient. The fractions obtained were analyzed using SDS-PAGE, followed by immunoblot analysis. *Tg*uL11m:HA was detected using an antibody against HA. Percentages in square brackets show the proportion of signal detected in fraction 7-9 versus signal intensity in fraction 10-14. 'I' indicates 10% of the input material for the centrifugation.

Discussion

Constraints in the recombination-active mitochondrial DNA of Toxoplasma gondii

The complexity of the *T. gondii* mitochondrial genome is puzzling, with an extensive variety of sequence block combinations and repetitions. The individual ONT reads are not repeated in the two previously published datasets (discussed in Berná et al., 2021b) and we also did not find overlap between data presented here and published previously (Namasivayam et al., 2021). Thus, the large number of block combinations identified here reinforces and elevates previous conclusions that continuous recombination shuffles the blocks (Namasivayam et al., 2021). At first sight, this might be considered a sign of evolutionary tinkering, similar to the mitochondrial idiosyncrasies found in other taxa that have been suggested to be non-adaptive,

such as the fragmented mitochondrial genomes of diplonemids (Burger and Valach, 2018) or

the repetitive and heterogeneous genomes of plant mitochondria (Kozik et al., 2019). Upon

closer scrutiny, however, the reshuffling appears limited to specific block borders and is not

random. In fact, the improved sequencing depth presented in this study revealed strong

constraints on the allowed block combinations in *T. gondii* mtDNA. Thus, a limited number of

allowed recombination sites generates the variety of actual DNA fragments sequenced. The

question remains whether the constraints outlined in the comprehensive model of

mitochondrial genome architecture presented here indicate functional features.

The purpose of genome blocks in light of small RNA production

Our small RNA sequencing results revealed that 15 small RNAs span block borders. There

are different explanations for this peculiar localization of small RNAs. One possibility is that

these RNAs are involved in the DNA recombination and replication process that occurs at block

borders. RNA molecules have been found to play important roles in DNA repair at double-

stranded breaks of nuclear DNA, both as long and small RNAs (Ohle et al., 2016; Wei et al.,

2012). However, it is currently unclear whether RNA:DNA hybrids play a role in mitochondrial

DNA recombination and repair (Allkanjari and Baldock, 2021).

Another possible explanation for the presence of small RNAs at block borders is that *T. gondii*

simply utilizes all available genomic space for RNA production, including block borders. *T.*

gondii appears to be as economical with its mitochondrial sequence space as other

apicomplexans, and it does have hardly any unused noncoding sequences in its genome. In

fact, the sequences at recombination sites could be regarded as an expansion of the

mitochondrial genome sequence space, which is not available to other apicomplexans like the

genus Plasmodium. There is evidence from recent polyA-sequencing efforts that long

transcripts spanning several short RNAs are also produced in *T. gondii* (Lee et al., 2021). This

is reminiscent of the situation in *P. falciparum*, where both strands of the entire mitochondrial

genome are transcribed into polycistronic precursor RNAs. Therefore, block borders in *T.*

gondii are likely transcribed into RNA as well. The functionality of these block border RNAs is

unclear, but the unique sequence combinations generated at the block borders give rise to

sRNAs that are not found in other apicomplexans with a simpler genome organization.

The discovery of sRNAs sharing sequences with mRNAs or being antisense to mRNAs in *T.*

gondii mitochondria is an unexpected finding of our study that was facilitated by our improved

understanding of mitochondrial genome organization. Antisense RNA is typically removed in

mitochondria from flies and humans (Pajak et al., 2019). Similar antisense degradation

mechanisms must be assumed in apicomplexans, as both strands of their genomes are

transcribed in full (Rehkopf et al., 2000). The fact that three small antisense RNAs survive

antisense RNA surveillance in *T. gondii* mitochondria is intriguing, and it remains to be

established whether RNA:RNA interactions of these sRNAs with mRNAs occur *in vivo* and

could be of regulatory impact.

Another potential regulatory link between mRNAs and sRNAs is represented by small RNAs

that utilize coding regions. *RNA19* is of particular interest because it contains a piece of *coxIII*

at the 5'-end and terminates close to the *cob* start codon, while also showing homology to

rRNA. A few cases of mRNA sequences overlapping in antisense orientation with rRNA have

been described in mammals and yeast (Coelho et al., 2002; Kermekchiev and Ivanova, 2001).

Sequences homologous to rRNAs have been found in many coding regions in sense and

antisense orientation (Mauro and Edelman, 1997). It has been suggested that this could link

ribosome production to other cellular processes by reciprocal inhibition of mRNA and rRNA

expression (Coelho et al., 2002). It is possible that *RNA19* and ribosome production could be

balanced with *cob* protein production by tuning the processing or RNA degradation of *cob*

mRNA. A quantitative analysis of RNA19 and *cob* mRNA accumulation under different

conditions could help to clarify whether there is such an inverse correlation. Overall, the

discovery of block-border sRNAs highlights the complex biogenesis of sRNAs in *T. gondii*

mitochondria and will be a starting point to understand the processing of sRNAs and their

function in general. Regarding their function, many, if not all, of the sRNAs at block borders

could be used in ribosomes as rRNA fragments, which is discussed in the next chapter.

sRNAs are incorporated into polysome-size complexes

A major unsolved problem in apicomplexan mitochondrial gene expression is the nature of the

ribosome. Recently, compelling evidence has surfaced suggesting the existence of ribosome-

sized particles, including ribosomal proteins. Using blue-native gel electrophoresis, it was

demonstrated that tagged ribosomal proteins *Tg*mS35, *Tg*bL12m, *Tg*uL3m and *Tg*uL24m

migrate as constituents of a macromolecular complex within the size range of a ribosome

(Lacombe et al., 2019; Shikha et al., 2022). Additionally, *Tg*mS35 and *Tg*uL24m coprecipitated

with mitochondria-encoded rRNA. Knockdown of *Tg*mS35, *Tg*bL35m, *Tg*bL36m and *Tg*bL28m

resulted in specific loss of activity of one respiratory complex, which is partially encoded in the

mitochondrial genome (Lacombe et al., 2019; Shikha et al., 2022). Nevertheless, it remains to

be determined whether these large complexes are actively involved in translation.

Here, we provide evidence that sRNAs with homology to rRNA are part of polysomes. This

conclusion is based on our finding that sRNAs, mRNAs and the ribosomal protein *Tg*uL11m

are all found in high-molecular weight complexes that are sensitive to ribosome-dissociative

conditions. We furthermore observed a difference in the gradient distribution of sRNAs

assigned to SSU and LSU, respectively. Based on a comparison with these two gradient

distribution patterns, we were able to assign RNA29 to LSU. In sum, our findings strongly

support the notion of active translation by mitochondrial ribosomes in tachyzoites of *T. gondii*,

and demonstrate that most if not all mitochondrial small RNAs fragments are part of the

apicomplexan mitochondrial ribosome. This includes sRNAs expressed from block borders like

the mRNA sequence-containing RNA3 and RNA19. Also, block border RNAs RNA1 and RNA2

that share their 3'-end sequence, but differ in their 5'-part are both incorporated into ribosomes.

RNA1 is of particular interest since it contains sequences homologous to two different parts of

the LSU rRNA from *E. coli* and hence could occupy two positions within a single ribosome.

RNA1 is almost exclusively found in ribosomes and polysomes, whereas RNA2 is also found

free of ribosomes - similar to RNA29. This suggests that RNA2 and RNA29 might not integrate

into ribosomes as effectively as other rRNA fragments, raising questions about the

coordination of rRNA fragment assembly. The assembly of a ribosome with numerous short

rRNA fragments and a comparable number of ribosomal proteins is undeniably complex.

Notably, longer variants of RNA19 are not incorporated into ribosomes, unlike smaller

isoforms, highlighting a connection between small RNA processing and ribosome assembly.

In the case of RNA19, processing precedes integration, contrasting with plant chloroplast

ribosomes where the 23S rRNA is split endonucleolytically at two sites post-integration (Liu et

al., 2015; Nishimura et al., 2010).

The analyses presented here mark a first step towards describing assembly steps of the

fascinating *T. gondii* ribosome. Unraveling the mechanistics of ribosome assembly will require

the identification of rRNA processing factors, which is within reach, given the excellent genetic

toolbox available for *T. gondii*. It will also eventually require understanding the ribosome 3D

structure. Within the spectrum of mitochondrial ribosome structures across species (Ramrath

et al., 2018; Saurer et al., 2023; Tobiasson et al., 2022; Waltz et al., 2021b) mitoribosomes of

T. gondii present an extreme example of ribosome diversification. The comprehensive

collection of small RNA fragments identified in our study, including those at block borders,

provides a valuable resource for future studies on the structure and function of the *T. gondii*

mitochondrial ribosome.

Conclusions

Apicomplexan mitochondrial genomes are vital for organism survival, expressing key

respiratory chain and gene expression components, particularly ribosomes. The latter were

proposed to be assembled from highly fragmented ribosomal RNAs, but whether rRNA

fragments are expressed and used in ribosomes was unclear. We adapted a protocol to enrich

T. gondii mitochondria (Esseiva et al. 2004) and used Nanopore sequencing to

comprehensively map the genome with its repeated sequence blocks. Small RNA sequencing

identified fragmented ribosomal RNAs, including some RNAs spanning block boundaries, thus

fusing protein-coding and rRNA sequences. Sucrose density gradient analysis showed that

such rRNA fragments are in polysome-size complexes. This distribution mirrored the

localization of the mitoribosomal protein L11 as well as mRNAs in sucrose gradients. sRNAs,

L11 and mRNAs shift to lower molecular weight complexes upon treatment with ribosome-

dissociative buffer. We conclude that most if not all mitochondrial sRNAs are components of

active ribosomes that assemble into polysomes. *T. gondii's* dynamic block-based genome

organization leads to usage of mitochondrial sequences in mRNA as well as rRNA contexts,

potentially linking rRNA and mRNA expression regulation.

Methods

Cultivation of host cells and *T. gondii* parasites

T. gondii parasites were cultured according to standard procedures (Jacot et al. 2020) unless

otherwise indicated. Briefly, human foreskin fibroblasts (HFF-1, ATCC SCRC-1041™) were

grown as host cells in Dulbecco's modified Eagle's medium (DMEM) (Capricorn Scientific)

supplemented with 10% fetal bovine serum (FBS) (Capricorn Scientific) and 100 µg/ml

penicillin–streptomycin (Capricorn Scientific). *T. gondii* tachyzoites were maintained in flasks

containing confluent HFF-1 cells in DMEM supplemented with 1% FBS and 100 µg/ml

penicillin–streptomycin. *T. gondii* strains were kindly provided by Frank Seeber. Experiments

were either carried out with parasites of the strain RH-ΔKUΔHX (Huynh and Carruthers 2009)

or RHpSAG1-βGal/pmt-GFP. The latter consists of RHβ1 (PMID: 8635747), which in addition

expresses eGFP N-terminally fused to the *T. gondii* mitochondrial targeting sequence S9(33-

159) (DeRocher et al., 2000).

CRISPR/Cas9 mediated endogenous tagging of *Tg*urpl11m

C-terminal tagging of TGRH88_003980_t1 was performed as described previously (Parker et

al. 2019). The sequence of a single guide RNA (sgRNA) specifically targeting the 3' end of

TGRH88_003980_t1 was integrated into pSAG1::CAS9-GFPU6::sgUPRT (Addgene plasmid

54467) using the Q5-site directed mutagenesis kit (NEB). The 3xHA epitope tag was

amplified from the pPR2-HA3 plasmid (Katris et al. 2014) and integrated into the pU6-DHFR

vector (Addgene plasmid #80329). The 3xHA-DHFR cassette was amplified from the resulting

vector by Q5 polymerase (NEB) using primers containing a 50 nt overlap homologous to the

either upstream or downstream regions of the TGRH88_003980_t1 stop codon (primers "flank

fwd *Tg*urpl11m tagging" and "flank rev *Tg*urpl11m tagging" in table S9, respectively). The

amplified donor construct was together with the pSAG1::CAS9-GFPU6::sgUPRT plasmid

encoding the Cas9 and the sgRNA co-transfected into RH-ΔKUΔHX as described previously

(Jacot et al. 2020). Parasites were grown under Pyrimethamine selection and subsequently

cloned by limiting dilution in 96 well plates (Roos et al. 1994). Genotyping was performed

directly from 96 well plates as described by (Piro et al. 2020). Proper integration of the 3xHA

tag was verified in single clones using Sanger Sequencing. Sequences of all the primers used

for cloning and genotyping are listed in table S9.

Immunofluorescence assays

Immunofluorescence assays were done as described previously (van Dooren et al., 2008).

Coverslips with confluent human foreskin fibroblasts (HFF) were infected with freshly egressed

Toxoplasma gondii parasites. The next day, cells were fixed in 3% (w/v) paraformaldehyde in

PBS for 15 min at room temperature and permeabilized in 0.25% (v/v) Triton X-100 in PBS for

10 min. Blocking was carried out in 2% (w/v) bovine serum albumin in PBS overnight. All

antibody incubation steps were done at room temperature for 1 hour. Rabbit anti-*Tg*Tom40

(van Dooren et al., 2016, 1:2000 dilution) and rat anti-HA high affinity (Sigma, 11867431001,

1:200 dilution) primary antibodies were used. Then the coverslips were washed three times in

PBS and incubated with donkey anti-rabbit CF 647 (Sigma, SAB4600177, 1:2000 dilution) and

goat anti-rat AlexaFluor 488 (Thermo Fisher Scientific, A-11006, 1:500 dilution) secondary

antibodies. Microscopy and image acquisition were performed on a DeltaVision Elite

deconvolution setup (GE Healthcare) using inverted Olympus IX71 microscope fitted with a

UPlanSApo 100x objective lens and Photometrics CoolSNAP HQ2 camera. Images were

deconvolved using SoftWoRx Suite 2.0 software, brightness and contrast were linearly

adjusted in FIJI/ImageJ (release 1.53c).

***T. gondii* organelle enrichment**

A previously established protocol to enrich *T. gondii* organelles was modified here slightly

(Esseiva et al., 2004). Freshly lysed *T. gondii* cultures from four T175 flasks (approximately

$8*10^8$ parasites) were filtered through a 3 µm pore size polycarbonate filter to remove host cell

debris and harvested by centrifugation at 1500 x *g* for 10 minutes. The parasite pellet was

briefly washed in ice-cold Dulbecco's phosphate-buffered saline (PBS) (Capricorn Scientific)

and again centrifuged for 10 minutes at 1500 x *g*. Afterwards, the parasite pellet was

resuspended in 20 mM Tris-HCl, pH 7.4, 600 mM mannitol, 5 mM MgCl$_2$, and 2 mM EDTA

containing 0.05% digitonin (Invitrogen) and incubated for 5 minutes on ice. The extract was

clarified by centrifugation at 8000 x *g* for 5 minutes at 4°C. The supernatant containing soluble

cytosolic components was discarded, and the corresponding pellet was resuspended in 20 mM

Tris-HCl, pH 7.4, 600 mM mannitol, 5 mM MgCl$_2$, and 1 mM EDTA containing benzonase (345

U/ml) (Novagen), incubated for 20 minutes at room temperature and subsequently centrifuged

at 8000 x *g* for 5 minutes at 4°C. The pellet was retained and washed in 20 mM Tris-HCl, pH

7.4, 600 mM mannitol, and 17 mM EDTA to inactivate benzonase remnants. Following a spin

at 8000 x *g* for 5 minutes at 4°C, the pellet consisted of a crude enrichment of organelles and

membranes. To study the *T. gondii* mitochondrial DNA content, the organelle pellet was directly

used for DNA isolation. For organellar RNA analysis, the organelle pellet was frozen at -80°C

for at least 24 hours. Afterwards, it was thawed on ice and resuspended in a high detergent

lysis buffer (20 mM HEPES-KOH pH 7.5, 100 mM KCl, 30 mM $MgCl_2$, 1% Igepal-CA630, 1.5%

Triton X-100, 0.5% sodium deoxycholate, 1 mM DTT, EDTA-free protease inhibitor). The lysate

was incubated during continuous end-over-end rotation at 4°C for 40 minutes and afterwards

clarified by centrifugation at 16000 x *g* for 10 minutes at 4°C. The pellet, composed of cell

debris and membranes, was discarded, and the supernatant, which contained the organelle

lysate, was resuspended in TRIzol® Reagent (Invitrogen) for RNA isolation.

SDS–PAGE and immunoblotting

To detect mitochondrial GFP in RH pSAG1-βGal/pmt-GFP parasites, samples taken during

organelle enrichment were directly diluted in Laemmli buffer (0.2 M Tris–HCl, 8% (w/v) SDS,

40% glycerol, 20% (v/v) β-mercaptoethanol, 0.005% bromophenol blue), incubated at 95°C for

5 minutes and subjected to sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–

PAGE). SDS–PAGE and immunoblotting were carried out as described previously (Kupsch et

al., 2012). For immunodetection, monoclonal primary mouse anti-GFP antibody (G1546,

Sigma–Aldrich) and polyclonal rabbit anti-heat shock protein 70 antibody (AS05 083A,

Agrisera) followed by secondary horseradish peroxidase (HRP)-labeled goat anti-mouse

antibody (ab205719, Abcam) and goat anti-rabbit antibody (ab205718, Abcam) were used.

Proteins from sucrose gradient fractions were isolated using methanol-chloroform-water

extraction (Wessel and Flügge 1984). Protein pellets were resuspended in Laemmli buffer,

heated for 5 minutes at 95°C and subjected to SDS-PAGE and immunoblotting. Primary rat

772 anti-HA antibody (11867423001, Roche) and secondary rabbit anti-rat (HRP) antibody

773 (ab6734, Abcam) were used for *Tg*uL11m:HA detection.

774 **DNA isolation and library construction**

775 DNA was isolated from organelle-enriched fractions using the GeneMATRIX Tissue DNA

776 Purification Kit (Roboklon) according to the manufacturer's protocol for cultured cells. The

777 Oxford Nanopore Technology sequencing library was prepared with 1 µg of organelle-enriched

778 DNA using the SQK-LSK108 ligation sequencing kit (Oxford Nanopore, ONT) according to the

779 manufacturer's instructions. Sequencing was performed on an R9.4.1 Spot ON Flow Cell; live-

780 basecalling was done using the ONT Guppy software package.

781 **RNA isolation and RNA Seq library construction**

782 RNA was isolated from total organelle enriched and sucrose gradient fraction samples using

783 TRIzol® Reagent (Invitrogen), followed by the Monarch™ Total RNA Miniprep Kit (New

784 England BioLabs) applying the manufacturer's protocol for TRIzol® extracted samples. Two

785 hundred nanograms of RNA was used to generate sequencing libraries using the NEBNext®

786 Multiplex Small RNA Library Prep Set for Illumina® Kit (New England BioLabs) according to

787 the manufacturer's instructions. For identification of PCR duplicates in the library preparation

788 process, the 5'-linker was modified with 8 N-nucleotides (as unique molecular identifiers,

789 UMIs). Small RNA library clean-up was carried out with AMPure XP Beads (Beckmann

790 Coulter). Sequencing was performed by Genewiz on an Illumina NovaSeq platform with a read

791 length of 150 bp for each pair.

792 **Agarose-formaldehyde gel electrophoresis**

793 RNA samples were diluted in denaturing loading buffer (Deionized formamide 62.5% (v/v),

794 formaldehyde 1.14 M, bromophenol blue 200 µg/mL, xylene cyanole 200 µg/mL, MOPS-

795 EDTA-sodium acetate) and separated on a 1% agarose gel containing 1.2% formaldehyde.

796 Cytosolic rRNAs were detected with ethidium bromide staining.

Denaturing Urea-PAGE and sRNA gel blot

RNA was separated by denaturing Urea-PAGE (10% or 12% polyacrylamide gel for total

RNA/organelle enriched RNA and RNA extracted from sucrose density gradient fractions,

respectively). Small RNA gel blotting and detection of transcripts with radiolabeled DNA oligo

probes was carried out as described previously (Loizeau et al., 2014). Sequences of the oligo

probes used in this study are listed in Tab. S9. Blot hybridization of sucrose density gradient

fractions was carried out consecutively in the order

RNA1_2,RNA19,SSUA,LSUD_E,RNA3,LSUF_G,RNA29, with removal of probes by heating

the blot to 60°C for 1 hour in 0.5% SDS.

5' RACE

T. gondii total RNA was ligated to a small RNA oligo (Rumsh, see Table S9) using T4 RNA

Ligase 1 (NEB) according to the manufacturer's instructions. Afterwards, RNA was reverse

transcribed using ProtoScript II Reverse Transcriptase (NEB) and random primers. PCR

amplification of cDNA was performed with a gene-specific *cob* primer and a primer annealing

to the ligated oligo (Rumsh1, see Table S9) using Taq DNA Polymerase (Roboklon). The

purified PCR product was sequenced by Sanger Sequencing (LGC Genomics GmbH).

Sucrose density gradient centrifugation

Ribosome fractionation by sucrose density gradient centrifugation was performed as

previously described (Waltz et al., 2021a). Briefly, *T. gondii* parasites were harvested and

organelle enrichment was performed as described above with small modifications. Salt

concentration in the digitonin and lysis buffer was either adjusted for optimal mitoribosome

stability (30 mM $MgCl_2$, 100 mM KCl) or for dissociative conditions (10 mM EDTA, 300 mM

KCl). The organelle enriched lysate was layered onto a continuous 10%-50% sucrose gradient.

Sucrose solutions (10% and 50%) were previously prepared freshly in 20 mM HEPES-KOH

pH 7.5,1 mM DTT, EDTA-free protease inhibitor and either 30 mM $MgCl_2$, 100 mM KCl or 10

mM EDTA, 300 mM KCl (dissociative conditions). A continuous sucrose gradient was formed

using BioComp Gradient Master 108. The gradient was centrifuged at 200.000xg for 2 hours

at 4°C in a S52-ST rotor (Thermo Scientific) and subsequently fractionated into 14 fractions

using the BioComp Piston Gradient Fractionator device. Fractions were resuspended directly

in TRIzol® reagent (Invitrogen). For RT-qPCR samples 1.2 ng of human estrogen receptor 1

(ESR1) RNA was introduced into each fraction as spike-in control. RNA was extracted from

fractions using Trizol-Chloroform phase separation followed by overnight precipitation with 1

volume Isopropanol.

RT-qPCR

RNA isolated from fractions was treated with TurboDNase (NEB) and afterwards subjected to

reverse transcription using ProtoScript® II Reverse Transcriptase (NEB). Synthesized cDNA

from each fraction was used as a template in qPCR. Reactions were performed as technical

triplicates using Luna® Universal qPCR Master Mix (NEB). Primers listed in Tab S9 were used

for amplification of mitochondrial cob, coxI, coxIII, LSUF/G rRNA and ESR1 spike-in for

normalization. The relative quantity of each amplicon was determined using the Pfaffl-method

(Pfaffl 2001). Relative quantities for amplicons in each fraction are presented as a percentage

of the cumulative relative quantities of the respective amplicon across all fractions. Data was

visualized using GraphPad Prism Version 10.1.0.

Bioinformatic analyses

All bioinformatic analyses except for the quantification of sequence block combinations were

performed on the Galaxy web platform (v.22.01; https://usegalaxy.eu, Afgan et al. 2018). All

reference sequences used were retrieved from GenBank (Sayers et al., 2019). Sequence

alignments were prepared using Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/)

with default settings. *E. coli* sequences were added manually according to sequence positions

established previously (Feagin et al., 2012).

Masking NUMTs in the *T. gondii* nuclear genome

To identify NUMTs in the *T. gondii* nuclear genome, mitochondrial sequence blocks

(MN077088.1-MN077111.1) were aligned against the RH-88 nuclear reference genome

(GCA_019455545.1, August 2021) using NCBI-BLASTN (v. 2.13.0) (Altschul et al., 1990)

(default parameters, except max_target_seqs = 5000). All hits obtained in the BLASTN tabular

output were manually integrated into a GFF3 file format. NUMTs in the nuclear genome were

masked with bedtools MaskFastaBed (Quinlan and Hall, 2010) based on the intervals defined

in the GFF3 file. In total, we masked 8118 sites with an average length of 92 nt representing

in total ~1% of the nuclear genome.

DNA sequencing data analysis

DNA Oxford Nanopore sequencing data were trimmed using porechop (Version 0.2.4,

https://github.com/rrwick/Porechop). As a first step, trimmed reads were mapped against the

human reference genome GRCh38.p14 (GCA_000001405.29, February 2022) using

Minimap2 (Version 2.24) (Li 2018) (PacBio/Oxford Nanopore read to reference mapping,

default settings). Mapping results were filtered using SAMtools view (Version 1.15.1) (Li et al.

2009) (-f read is unmapped). Unmapped reads were subsequently simultaneously mapped

against the *T. gondii* RH-88 nuclear reference genome masked for NUMTs, the apicoplast

genome (CM033583.1) and mitochondrial sequence blocks (MN077088.1- MN077111.1)

using Minimap2. A fraction of the reads that fit best to the mitochondrial genome still retain

shorter nuclear sequence sections and are therefore representing NUMTs. We therefore took

all reads that mapped to the mitochondrial genome in our first mapping and mapped them

against the nuclear reference genome masked for NUMTs using Minimap2 (Version 2.24). The

remaining unmapped reads were considered mitochondrial and visualized with Geneious

Prime® v. 2019.2.3. (Biomatters Ltd.). Reads were annotated providing the mitochondrial

sequence blocks (MN077088.1 - MN077111.1) and ORFs (MN077082.1 - MN077084.1) as

annotation database using the Geneious Prime® "Annotate from Database" tool with similarity

set to ≥75%.

Sequence comparisons of ONT reads found here with published ONT reads for the *T. gondii*

mitochondrial genome were performed using the NUCmer tool in the MUMmer package of

Galaxy with the following settings: minimum match length 20 and minimum alignment length

290 (shortest read in Namasivayam et al., 2021). The Show-Coords tool was used to output

alignment information. We found 1555 reads of our dataset being entirely part (defined as

≥99% read coverage) of reads from the Namasivayam et al. (2021) and 212 reads of their

dataset being fully found within reads from our dataset.

Analysis of mitochondrial DNA sequence block arrangements

For a comprehensive overview of the analysis, including the complete .Rmd report please refer

to the GitHub repository (https://github.com/Kovox91/ToxoBlocks). In brief, annotated block

IDs, indicating the directionality of each block, were exported in plain text. A custom R script,

leveraging the tidyverse library (Wickham et al. 2019), was employed for the subsequent

analysis. This script facilitated the removal of block combinations where two blocks were

distanced by more than 10 nucleotides. For two-block combinations, the script summed up all

identified combinations, filtering out those occurring less than 50 times. The results,

incorporating block orientations, were visualized using ComplexHeatmap (Gu and

Hübschmann 2022). To quantify the relative abundance of ORFs , counts for the respective

ORF combinations were tallied across all reads. The ORF frequency was then computed as

the ratio of reads containing at least one instance of the specific ORF to the total number of

reads that were at least as long as the given ORF. RNA sequencing data analysis.

Adapter and quality trimming of RNA sequencing data was performed using Cutadapt (Version

4.1; Martin, 2011). Umi-tools was used to extract UMIs (Version 1.1.2; Smith et al., 2017).

Reads were aligned to the *T. gondii* RH-88 nuclear reference genome masked for NUMTs and

the apicoplast genome (CM033583.1) using Bowtie2 (Version 2.4.5; Langmead et al., 2009)

in default settings. Mapping results were filtered using SAMtools view (Version 1.15.1; Li et al.,

2009) (-f read is unmapped). Unmapped reads were kept and mapped with Bowtie2 using

default settings against the *T. gondii* mitochondrial genome using all combinations of the

sequence blocks found in our genome sequence analysis (GenBank accession MN077088.1 -

MN077111.1, OR086910 - OR086916) as determined in our ONT DNA sequencing data.

Transcripts were determined as follows: first, most transcripts had short poly-A tails - up to 12

adenines at their 3'-ends - which is similar to the situation in *P. falciparum* (Rehkopf et al.,

2000). For the identification of the transcript 3'-ends, all polyA-tails longer than three

nucleotides were removed from reads using fastp (Chen et al., 2018). The number of mapped

3' and 5' read ends at each position in the reference was calculated using Bedtools Genome

Coverage 2.30.0 (Quinlan and Hall, 2010). The 5'-ends at the beginning of coverage plateaus

over the background were considered starting points of transcripts. For each 5' end, the

mapped 3'-end most far away but still represented by at least 10% of the reads of the 5'-

terminus and connected to the starting point by uninterrupted coverage over background was

considered the corresponding 3'-terminus. Alignments and coverage graphs were visualized

in Integrated Genome Viewer (Robinson et al., 2011). All RNA sequencing data are available

at SRA under accession number PRJNA978626.

Sequence similarity search

All newly identified transcripts were subjected to an NCBI-BLASTN homology search (v.

2.13.0; Altschul et al., 1990; default parameters, except word size=7) against the *Plasmodium*

falciparum mitochondrial genome (M76611.1) and the *Eimeria leuckarti* mitochondrial genome

(MW354691.1). Transcripts not matching the *P. falciparum* or *E. leuckarti* mitochondrial

genome were further inspected for homology to the *E. coli* rRNA operon (J01695.2) using

BLASTN.

Data access

923 The RNA and DNA sequencing data generated in this study have been submitted to the NCBI

924 BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number

925 PRJNA978626. New block annotations can be found in GenBank, OR086910 - OR086916.

Competing interests

927 The authors declare that they have no competing interests.

Acknowledgments

We are grateful to Giel van Dooren, Frank Seeber, Kai Matuschewski, and Hannes Ruwe for

critical discussion of our data and manuscript. We sincerely thank Frank Seeber for providing

the T. gondii strains. We are grateful to Giel van Dooren for providing the plasmids for

CRISPR/Cas9 tagging and for support from the microscopy facility at ANU Canberra. We thank

Svea Beier for preparing the constructs for RPL11 tagging and Florian Rösch for preparing the

vector for the ESR1-spike in amplification.

Authors' contributions

936 S. W: and C.S. designed the research; S.T., A.H., N.D., Z.G. and P.L. performed the

937 research; S.T., A.H., N.D., S.M. and S.W. analyzed the data; C.S. wrote the paper with

938 contributions from all other authors.

Funding
940 This work was supported by the German research foundation (DFG) by grant IRTG2290,

941 project B01 to C.S.

References

Alday PH, Bruzual I, Nilsen A, Pou S, Winter R, Ben Mamoun C, Riscoe MK, Doggett JS. 2017. Genetic Evidence for Cytochrome b Qi Site Inhibition by 4(1H)-Quinolone-3-Diarylethers and Antimycin in Toxoplasma gondii. *Antimicrob Agents Chemother* **61**. doi:10.1128/AAC.01866-16

Allkanjari K, Baldock RA. 2021. Beyond base excision repair: an evolving picture of mitochondrial DNA repair. *Biosci Rep* **41**. doi:10.1042/BSR20211320

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* **215**:403–410.

Berná L, Marquez P, Cabrera A, Greif G, Francia ME, Robello C. 2021a. Reevaluation of the Toxoplasma gondii and Neospora caninum genomes reveals misassembly, karyotype differences, and chromosomal rearrangements. *Genome Res* **31**:823–833.

Berná L, Rego N, Francia ME. 2021b. The Elusive Mitochondrial Genomes of Apicomplexa: Where Are We Now? *Front Microbiol* **12**:751775.

Burger G, Valach M. 2018. Perfection of eccentricity: Mitochondrial genomes of diplonemids. *IUBMB Life* **70**:1197–1206.

Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, Metcalf T, Modrzynska K, Schwach F, Martin RE, Mather MW, McFadden GI, Parts L, Rutledge GG, Vaidya AB, Wengelnik K, Rayner JC, Billker O. 2017. Functional Profiling of a Plasmodium Genome Reveals an Abundance of Essential Genes. *Cell* **170**:260–272.e8.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**:i884–i890.

Coelho PSR, Bryan AC, Kumar A, Shadel GS, Snyder M. 2002. A novel mitochondrial protein, Tar1p, is encoded on the antisense strand of the nuclear 25S rDNA. *Genes Dev* **16**:2755–2760.

DeRocher A, Hagen CB, Froehlich JE, Feagin JE, Parsons M. 2000. Analysis of targeting sequences demonstrates that trafficking to the Toxoplasma gondii plastid branches off the secretory system. *J Cell Sci* **113 (Pt 22)**:3969–3977.

Esseiva AC, Naguleswaran A, Hemphill A, Schneider A. 2004. Mitochondrial tRNA import in Toxoplasma gondii. *J Biol Chem* **279**:42363–42368.

Feagin JE, Harrell MI, Lee JC, Coe KJ, Sands BH, Cannone JJ, Tami G, Schnare MN, Gutell RR. 2012. The fragmented mitochondrial ribosomal RNAs of Plasmodium falciparum. *PLoS One* **7**:e38320.

Gjerde B. 2013. Characterisation of full-length mitochondrial copies and partial nuclear copies (numts) of the cytochrome b and cytochrome c oxidase subunit I genes of Toxoplasma gondii, Neospora caninum, Hammondia heydorni and Hammondia triffittae (Apicomplexa: Sarcocystidae). *Parasitol Res* **112**:1493–1511.

Gupta A, Shah P, Haider A, Gupta K, Siddiqi MI, Ralph SA, Habib S. 2014. Reduced ribosomes of the apicoplast and mitochondrion of Plasmodium spp. and predicted interactions with antibiotics. *Open Biol* **4**:140045.

Hikosaka K, Kita K, Tanabe K. 2013. Diversity of mitochondrial genome structure in the phylum Apicomplexa. *Mol Biochem Parasitol* **188**:26–33.

Hillebrand A, Matz JM, Almendinger M, Muller K, Matuschewski K, Schmitz-Linneweber C. 2018. Identification of clustered organellar short (cos) RNAs and of a conserved family of organellar RNA-binding proteins, the heptatricopeptide repeat proteins, in the malaria parasite. *Nucleic Acids Res.* doi:10.1093/nar/gky710

Hollin T, Abel S, Falla A, Pasaje CFA, Bhatia A, Hur M, Kirkwood JS, Saraf A, Prudhomme J, De Souza A, Florens L, Niles JC, Le Roch KG. 2022. Functional genomics of RAP proteins and their role in mitoribosome regulation in Plasmodium falciparum. *Nat Commun* **13**:1275.

Ji YE, Mericle BL, Rehkopf DH, Anderson JD, Feagin JE. 1996. The Plasmodium falciparum 6 kb element is polycistronically transcribed. *Mol Biochem Parasitol* **81**:211–223.

Ke H, Morrisey JM, Ganesan SM, Mather MW, Vaidya AB. 2012. Mitochondrial RNA polymerase is an essential enzyme in erythrocytic stages of Plasmodium falciparum. *Mol Biochem Parasitol* **185**:48–51.

Kermekchiev M, Ivanova L. 2001. Ribin, a protein encoded by a message complementary to rRNA, modulates ribosomal transcription and cell proliferation. *Mol Cell Biol* **21**:8255–8263.

Kozik A, Rowan BA, Lavelle D, Berke L, Schranz ME, Michelmore RW, Christensen AC. 2019. The alternative reality of plant mitochondrial DNA: One ring does not rule them all. *PLoS Genet*

15:e1008373.

Kupsch C, Ruwe H, Gusewski S, Tillich M, Small I, Schmitz-Linneweber C. 2012. Arabidopsis
chloroplast RNA binding proteins CP31A and CP29A associate with large transcript pools and
confer cold stress tolerance by influencing multiple chloroplast RNA processing steps. *Plant Cell*
24:4266–4280.

Lacombe A, Maclean AE, Ovciarikova J, Tottey J, Mühleip A, Fernandes P, Sheiner L. 2019.
Identification of the Toxoplasma gondii mitochondrial ribosome, and characterisation of a protein
essential for mitochondrial translation. *Mol Microbiol* **112**:1235–1252.

Lane KD, Mu J, Lu J, Windle ST, Liu A, Sun PD, Wellems TE. 2018. Selection of Plasmodium
falciparum cytochrome B mutants by putative PfNDH2 inhibitors. *Proc Natl Acad Sci U S A*
115:6285–6290.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of short
DNA sequences to the human genome. *Genome Biol* **10**:R25.

Lee VV, Judd LM, Jex AR, Holt KE, Tonkin CJ, Ralph SA. 2021. Direct Nanopore Sequencing of
mRNA Reveals Landscape of Transcript Isoforms in Apicomplexan Parasites. *mSystems* **6**.
doi:10.1128/mSystems.01081-20

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000
Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* **25**:2078–2079.

Liu J, Zhou W, Liu G, Yang C, Sun Y, Wu W, Cao S, Wang C, Hai G, Wang Z, Bock R, Huang J,
Cheng Y. 2015. The conserved endoribonuclease YbeY is required for chloroplast ribosomal RNA
processing in Arabidopsis. *Plant Physiol* **168**:205–221.

Loizeau K, Qu Y, Depp S, Fiechter V, Ruwe H, Lefebvre-Legendre L, Schmitz-Linneweber C,
Goldschmidt-Clermont M. 2014. Small RNAs reveal two target sites of the RNA-maturation factor
Mbb1 in the chloroplast of Chlamydomonas. *Nucleic Acids Res* **42**:3286–3297.

MacRae JI, Sheiner L, Nahid A, Tonkin C, Striepen B, McConville MJ. 2012. Mitochondrial metabolism
of glucose and glutamine is required for intracellular growth of Toxoplasma gondii. *Cell Host*
Microbe **12**:682–692.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet 17. doi:10.14806/ej.17.1.200.

Mauro VP, Edelman GM. 1997. rRNA-like sequences occur in diverse primary transcripts: implications
for the control of gene expression. *Proc Natl Acad Sci U S A* **94**:422–427.

McFadden DC, Tomavo S, Berry EA, Boothroyd JC. 2000. Characterization of cytochrome b from
Toxoplasma gondii and Q(o) domain mutations as a mechanism of atovaquone-resistance. *Mol*
Biochem Parasitol **108**:1–12.

Melo EJ, Attias M, De Souza W. 2000. The single mitochondrion of tachyzoites of Toxoplasma gondii.
J Struct Biol **130**:27–33.

Molan A, Nosaka K, Hunter M, Wang W. 2019. Global status of Toxoplasma gondii infection:
systematic review and prevalence snapshots. *Trop Biomed* **36**:898–925.

Namasivayam S, Baptista RP, Xiao W, Hall EM, Doggett JS, Troell K, Kissinger JC. 2021. A novel
fragmented mitochondrial genome in the protist pathogen Toxoplasma gondii and related tissue
coccidia. *Genome Res* **31**:852–865.

Namasivayam S, Sun C, Bah AB, Oberstaller J, Pierre-Louis E, Etheridge RD, Feschotte C, Pritham
EJ, Kissinger JC 2023. Massive invasion of organellar DNA drives nuclear genome evolution in
Toxoplasma. *Proc Natl Acad Sci U S A* 120:e2308569120.

Nishimura K, Ashida H, Ogawa T, Yokota A. 2010. A DEAD box protein is required for formation of a
hidden break in Arabidopsis chloroplast 23S rRNA. *Plant J* **63**:766–777.

Oborník M, Lukeš J. 2015. The Organellar Genomes of Chromera and Vitrella, the Phototrophic
Relatives of Apicomplexan Parasites. *Annu Rev Microbiol* **69**:129–144.

Ohle C, Tesorero R, Schermann G, Dobrev N, Sinning I, Fischer T. 2016. Transient RNA-DNA Hybrids
Are Required for Efficient Double-Strand Break Repair. *Cell* **167**:1001–1013.e7.

Ossorio PN, Sibley LD, Boothroyd JC. 1991. Mitochondrial-like DNA sequences flanked by direct and
inverted repeats in the nuclear genome of Toxoplasma gondii. *J Mol Biol* **222**:525–536.

Pajak A, Laine I, Clemente P, El-Fissi N, Schober FA, Maffezzini C, Calvo-Garrido J, Wibom R,
Filograna R, Dhir A, Wedell A, Freyer C, Wredenberg A. 2019. Defects of mitochondrial RNA
turnover lead to the accumulation of double-stranded RNA in vivo. *PLoS Genet* **15**:e1008240.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**:841–842.

Ramrath DJF, Niemann M, Leibundgut M, Bieri P, Prange C, Horn EK, Leitner A, Boehringer D,

Schneider A, Ban N. 2018. Evolutionary shift toward protein-based architecture in trypanosomal
mitochondrial ribosomes. *Science* **362**. doi:10.1126/science.aau7735

Rehkopf DH, Gillespie DE, Harrell MI, Feagin JE. 2000. Transcriptional mapping and RNA processing
of the Plasmodium falciparum mitochondrial mRNAs. *Mol Biochem Parasitol* **105**:91–103.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011.
Integrative genomics viewer. *Nat Biotechnol* **29**:24–26.

Saurer M, Leibundgut M, Nadimpalli HP, Scaiola A, Schönhut T, Lee RG, Siira SJ, Rackham O, Dreos
R, Lenarčič T, Kummer E, Gatfield D, Filipovska A, Ban N. 2023. Molecular basis of translation
termination at noncanonical stop codons in human mitochondria. *Science* **380**:531–536.

Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I. 2019. GenBank. *Nucleic*
Acids Res **47**:D94–D99.

Seeber F, Feagin JE, Parsons M, van Dooren GG. 2020. Chapter 11 - The apicoplast and
mitochondrion of Toxoplasma gondii In: Weiss LM, Kim K, editors. Toxoplasma Gondii (Third
Edition). Academic Press. pp. 499–545.

Seeber F, Ferguson DJ, Gross U. 1998. Toxoplasma gondii: a paraformaldehyde-insensitive
diaphorase activity acts as a specific histochemical marker for the single mitochondrion. *Exp*
Parasitol **89**:137–139.

Shikha S, Silva MF, Sheiner L. 2022. Identification and Validation of Toxoplasma gondii Mitoribosomal
Large Subunit Components. *Microorganisms* **10**. doi:10.3390/microorganisms10050863

Sidik SM, Huet D, Ganesan SM, Huynh M-H, Wang T, Nasamu AS, Thiru P, Saeij JPJ, Carruthers VB,
Niles JC, Lourido S. 2016. A Genome-wide CRISPR Screen in Toxoplasma Identifies Essential
Apicomplexan Genes. *Cell* **166**:1423–1435.e12.

Smith T, Heger A, Sudbery I. 2017. UMI-tools: modeling sequencing errors in Unique Molecular
Identifiers to improve quantification accuracy. *Genome Res* **27**:491–499.

Srivastava IK, Morrisey JM, Darrouzet E, Daldal F, Vaidya AB. 1999. Resistance mutations reveal the
atovaquone-binding domain of cytochrome b in malaria parasites. *Mol Microbiol* **33**:704–711.

Subczynski WK, Pasenkiewicz-Gierula M, Widomska J, Mainali L, Raguz M. 2017. High
Cholesterol/Low Cholesterol: Effects in Biological Membranes: A Review. *Cell Biochem Biophys*
75:369–385.

Suplick K, Morrisey J, Vaidya AB. 1990. Complex transcription from the extrachromosomal DNA
encoding mitochondrial functions of Plasmodium yoelii. *Mol Cell Biol* **10**:6381–6388.

Syafruddin D, Siregar JE, Marzuki S. 1999. Mutations in the cytochrome b gene of Plasmodium
berghei conferring resistance to atovaquone. *Mol Biochem Parasitol* **104**:185–194.

Szabo EK, Finney CAM. 2017. Toxoplasma gondii: One Organism, Multiple Models. *Trends Parasitol*
33:113–127.

Tobiasson V, Berzina I, Amunts A. 2022. Structure of a mitochondrial ribosome with fragmented rRNA
in complex with membrane-targeting elements. *Nat Commun* **13**:6132.

Vaidya AB, Lashgari MS, Pologe LG, Morrisey J. 1993. Structural features of Plasmodium cytochrome
b that may underlie susceptibility to 8-aminoquinolines and hydroxynaphthoquinones. *Mol*
Biochem Parasitol **58**:33–42.

van Dooren GG, Tomova C, Agrawal S, Humbel BM, Striepen B. 2008. Toxoplasma gondii Tic20 is
essential for apicoplast protein import. *Proc Natl Acad Sci U S A* **105**:13574–13579.

van Dooren GG, Yeoh LM, Striepen B, McFadden GI. 2016. The Import of Proteins into the
Mitochondrion of Toxoplasma gondii. *J Biol Chem* **291**:19335–19350.

Vercesi AE, Rodrigues CO, Uyemura SA, Zhong L, Moreno SN. 1998. Respiration and oxidative
phosphorylation in the apicomplexan parasite Toxoplasma gondii. *J Biol Chem* **273**:31040–31047.

Waltz F, Giegé P, Hashem Y. 2021a. Purification and Cryo-electron Microscopy Analysis of Plant
Mitochondrial Ribosomes. *Bio Protoc* **11**:e4111.

Waltz F, Salinas-Giegé T, Englmeier R, Meichel H, Soufari H, Kuhn L, Pfeffer S, Förster F, Engel BD,
Giegé P, Drouard L, Hashem Y. 2021b. How to build a ribosome from RNA fragments in
Chlamydomonas mitochondria. *Nat Commun* **12**:7176.

Wei W, Ba Z, Gao M, Wu Y, Ma Y, Amiard S, White CI, Rendtlew Danielsen JM, Yang Y-G, Qi Y.
2012. A role for small RNAs in DNA double-strand break repair. *Cell* **149**:101–112.

1110