1 Sodium channels expressed in nociceptors contribute distinctly to

2 action potential subthreshold phase, upstroke and shoulder

Phil Alexander Köster[a,b], Enrico Leipold[c], Jenny Tigerholm[b,d], Anna Maxion[a,b,e], Barbara

Namer[a,b,e], Thomas Stiehl[b,d,f*], Angelika Lampert[a,b*]

[a]University Hospital, RWTH Aachen University, Institute for Neurophysiology, Pauwelsstrasse 30, 52074 Aachen, Germany

[b]Scientific Center for Neuropathic pain Aachen SCN[AACHEN], Uniklinik RWTH Aachen University, Aachen, Germany

[c]University of Luebeck, Department of Anesthesiology and Intensive Care and CBBM-Center of Brain, Behavior and Metabolism,

Ratzeburger Allee 160, 23562 Lübeck, Germany

[d]University Hospital, RWTH Aachen University, Joint Research Center for Computational Biomedicine (JRCC), Pauwelsstrasse 30, 52074

Aachen, Germany

[e]University Hospital, RWTH Aachen University, Interdisciplinary Center for Clinical Research (IZKF), Faculty of Medicine, Research

Group Neurosciences, Pauwelsstrasse 30, 52074 Aachen, Germany

[f]University Hospital, RWTH Aachen University, Institute for Computational Biomedicine and Disease modelling with focus on phase

transitions between phenotypes, Pauwelsstrasse 30, 52074 Aachen, Germany

*Corresponding authors:

Angelika Lampert

Institute for Neurophysiology, University Hospital, RWTH Aachen University

Pauwelsstrasse 30

52074 Aachen, Germany

E-mail: alampert@ukaachen.de

ORCID: https://orcid.org/0000-0001-6319-6272

Thomas Stiehl

Institute for Computational Biomedicine and Disease Modelling with focus on phase

transitions between phenotypes, University Hospital, RWTH Aachen University

Pauwelsstrasse 19

52074 Aachen, Germany

E-mail: tstiehl@ukaachen.de

ORCID: https://orcid.org/0000-0001-9686-9197

Short running title (max. 50 characters): Sodium channel subthreshold gating

Summary (~ 40 words, present tense, refer to the work in 3rd person): Subthreshold

gating of seven sodium channels ($Na_v1.1$-3, $Na_v1.5$-8) is determined by manual patch clamp

and, together with $Na_v1.9$, integrated into a computer model of an Aδ- and a mechano-

insensitive nociceptor (CMi). Simulations reveal contribution of $Na_v1.9$ to the action potential

upstroke and shoulder and prove useful for $Na_v1.7$-related disease modelling.

1 Abstract (max. 250 words)

Voltage-gated sodium channels (VGSC) in the peripheral nervous system shape action

potentials (AP) and thereby support the detection of sensory stimuli. Most of the nine

mammalian VGSC subtypes are expressed in nociceptors, but predominantly, three are linked

to several human pain syndromes: while $Na_v1.7$ is suggested to be a (sub-)threshold channel,

$Na_v1.8$ is thought to support the fast AP upstroke. $Na_v1.9$, as it produces large persistent

currents, is attributed a role in determining the resting membrane potential.

We characterized gating of $Na_v1.1$-$Na_v1.3$ and $Na_v1.5$-$Na_v1.9$ in manual patch clamp with focus

on the AP subthreshold depolarization phase. $Na_v1.9$ exhibited the most hyperpolarized

activation while its fast inactivation resembled the depolarized inactivation of $Na_v1.8$. For some

VGSCs (e.g., $Na_v1.1$ and $Na_v1.2$), a positive correlation between ramp current and window

current was detected.

Using a modified Hodgkin-Huxley model which accounts for the time needed for inactivation

to occur, we used the acquired data to simulate two nociceptive nerve fiber types (an Aδ- and a

mechano-insensitive C-nociceptor) containing VGSC conductances according to published

human RNAseq data. Our simulations suggest that $Na_v1.9$ is supporting both the AP upstroke

and its shoulder. A reduced threshold for AP generation was induced by enhancing $Na_v1.7$

conductivity or shifting its activation to more hyperpolarized potentials, as observed in $Na_v1.7$-

related pain disorders.

Here, we provide a comprehensive, comparative functional characterization of VGSCs relevant

in nociception and describe their gating with Hodgkin-Huxley-like models, which can serve as

a tool to study their specific contributions to AP shape and sodium channel-related diseases.

1 Introduction

Next to their well-known role to initiate the upstroke of the action potential (AP), voltage-gated sodium channels (VGSC) fine-tune cellular excitability by contributing to subthreshold depolarizations and fluctuations of the membrane potential. Thus, they are involved in the detection and amplification of sensory stimuli e.g. in peripheral sensory neurons such as nociceptors (Bhattacharjee et al., 2018; Bennett et al., 2019; Goodwin and McMahon, 2021; Tian et al., 2023). Nociceptive stimuli depolarize the cell membrane of the nerve fiber ending, but if they are small, they do not initiate an AP, leading to subthreshold depolarizations. Subthreshold depolarizations directly preceding an AP are described in rats to be supported by VGSC currents sensitive to the pufferfish neurotoxin tetrodotoxin (TTX) (Enomoto et al., 2006; Vasylyev and Waxman, 2012), but the specific contribution of each channel isoform is yet to be fully understood.

To date, nine isoforms of mammalian VGSC α subunits have been described ($Na_v1.1 - Na_v1.9$), which are encoded by the genes *SCN1A-SCN5A* and *SCN8A-SCN11A* (Yu and Catterall, 2003; Catterall et al., 2005; Dib-Hajj et al., 2010). They differ in their biophysical properties, such as kinetics and voltage-dependence of activation, inactivation, and deactivation (Ahern et al., 2016; Körner and Lampert, 2020; Catterall, 2023) and are classified according to their sensitivity to TTX as either TTX-resistant (TTXr; $Na_v1.5$, $Na_v1.8$, and $Na_v1.9$) or TTX-sensitive (TTXs; all remaining isoforms) (de Lera Ruiz and Kraus, 2015).

Depending on their type and physiological function, excitable human cells express different VGSC isoforms (Catterall et al., 2005; Ahern et al., 2016). For example, $Na_v1.1 - Na_v1.3$ and $Na_v1.6$ are widely expressed in neurons of the central (CNS) and peripheral nervous system (PNS) (Whitaker et al., 2000; Vacher et al., 2008; Liang et al., 2021), while $Na_v1.7-Na_v1.9$ are expressed in PNS neurons including dorsal root ganglion (DRG) neurons, sympathetic ganglion neurons or olfactory sensory neurons (Fukuoka et al., 2008; Wang et al., 2017). The skeletal

and cardiac muscle express their own specific subtypes: $Na_v1.4$ and $Na_v1.5$, respectively

(Sheets and Hanck, 1999). $Na_v1.5$ can also be detected in PNS neurons during development or

injury (Renganathan et al., 2002) and at low levels in adults as revealed by recent

transcriptomics studies of human sensory neurons (Tavares-Ferreira et al., 2022).

The VGSC expression patterns of different excitable cell populations can overlap and are

subject to dynamic changes during development or in response to injury or sensitization.

Expression of $Na_v1.3$, for example, was described in embryonic, but not in healthy adult rat

DRG neurons. However, expression of $Na_v1.3$ and associated currents have found to be

upregulated in rodent DRG neurons after spinal cord injury (Felts et al., 1997; Hains et al.,

2003; Lampert et al., 2006).

The sensation of pain via DRG neurons is associated with the activation of their afferent C- or

Aδ-nerve fibers. Among them, the subclass of C-fibers known as "silent", "sleeping" or

mechano-insensitive (CMi) fibers is assumed to be involved in the generation of neuropathic

pain by exhibiting abnormal spontaneous activity (Namer and Handwerker, 2009; Kleggetveit

et al., 2012). Cell excitability of different fiber types and therefore their function in pain

signaling are subject to various factors of influence. Recent transcriptomic studies including

human tissue revealed distinct VGSC expression patterns in each sensory nerve fiber type

(Usoskin et al., 2015; Kupari et al., 2021; Nguyen et al., 2021; Tavares-Ferreira et al., 2022;

Jung et al., 2023; Bhuiyan et al., 2024). To date, the influence of these expression patterns on

nerve fiber excitability is yet to be fully understood. It is therefore beneficial to first investigate

the contribution of each VGSC isoform to nerve fiber excitability in isolation.

Gain-of-function variants of $Na_v1.7$ are linked to inherited pain syndromes such as

erythromelalgia (Yang et al., 2004; Lampert et al., 2010; Bennett and Woods, 2014; Brouwer

et al., 2014; Tang et al., 2015; Dib-Hajj et al., 2017) or paroxysmal extreme pain disorder

(Fertleman et al., 2006; Fertleman et al., 2007; Jarecki et al., 2008; Stępień et al., 2020). Studies

of these mutations in heterologous expression systems show a prominent shift of voltage-

dependence of activation to more hyperpolarized potentials and suggest that $Na_v1.7$ plays an

important role in the initiation of APs. Recently, however, in recordings of human sensory

neurons derived from induced pluripotent stem cells (iPSC), the contribution of $Na_v1.7$ to

subthreshold depolarizations was observed to be limited, and the channel was considered a

threshold channel rather than a subthreshold channel (Meents et al., 2019).

Functional studies in rodents and heterologous expression systems so far suggest that $Na_v1.8$ is

the main contributor to the upstroke of the AP and determines its duration by shaping the AP's

shoulder (Renganathan et al., 2001; Blair and Bean, 2002; Matsutomi et al., 2006; Han et al.,

2015a). $Na_v1.9$, on the other hand, is thought to modulate the resting membrane potential by

eliciting large persistent currents (Herzog et al., 2001; Maingret et al., 2008; Tian et al., 2023).

A contribution of $Na_v1.9$ to the AP shoulder was nicely demonstrated in enteric neurons of

$Na_v1.9$ knockout mice (Osorio et al., 2014) and observed in DRG neurons of mice carrying a

$Na_v1.9$ missense mutation (Leipold et al., 2013), while another study failed to see an influence

of $Na_v1.9$ on AP shape (Priest et al., 2005). Thus, the question of the role of $Na_v1.9$ in shaping

AP properties is left unresolved.

$Na_v1.9$ is long known to show poor heterologous expression (Vanoye et al., 2013; Goral et al.,

2015; Sizova et al., 2020). Various attempts to express $Na_v1.9$ in heterologous expression

systems yielded limited success with very few exceptions (Tate et al., 1998; Leipold et al.,

2013; Vanoye et al., 2013; Leipold et al., 2015). Although chimeric $Na_v1.9$ channels containing

the C-terminus from $Na_v1.4$ show improved expression, they exhibit altered voltage

dependence of channel activation and inactivation (Goral et al., 2015; Bothe and Lampert,

2021) and are therefore unsuitable for studying $Na_v1.9$ gating and its contribution to AP genesis.

Once the gating parameters of each channel subtype have been analyzed, their contribution to

cellular excitability can be reconstructed and investigated using mathematical and *in silico*

models. In 1952, Hodgkin and Huxley provided a mathematical description of VGSC gating

and APs, which still today is often used to model electrophysiological data and cellular

excitability (Hodgkin and Huxley, 1952; Stiles and Gray, 2021; Wang et al., 2022). However,

classic Hodgkin-Huxley models implement VGSC inactivation right at the beginning of a

voltage step stimulus and therefore do not account for the duration of protein-level

conformational changes that are assumed to occur within the channel for the so-called

inactivation particle to bind to its receptor and thereby stopping the associated sodium influx.

C-fiber excitability in particular has been modelled and used successfully to describe the

biophysical properties of this fiber type (Petersson et al., 2014; Tigerholm et al., 2014;

Tigerholm et al., 2015; Maxion et al., 2023). In most cases, only select VGSC isoforms are

included in models of excitable cells, often because of a lack of a standardized comprehensive

data set which is needed as basis for model generation. Larger studies comparing VGSC gating

in vitro in a comparable experimental setting have focused on suprathreshold gating of a subset

of VGSCs expressed in the CNS and related mutations leading e.g. to epilepsy (Thompson et

al., 2023), or on sensory neuron VGSCs with a focus on their temperature dependence

(Kriegeskorte et al., 2023).

As conclusive and complete data on the sub- and suprathreshold activity of all VGSC isoforms

are not yet available, we here conducted a standardized and comparative examination of VGSC

isoforms involved in nociception ($Na_v1.1$-1.3 and $Na_v1.5$-1.8) with consistent experimental

conditions using patch-clamp experiments in heterologous expression systems (HEK293(T)

and ND7/23 cells).

With these data, we developed a computational model which is based on the Hodgkin-Huxley

framework and parameterized using our patch-clamp data. We added $Na_v1.9$ to this model using

previously published data obtained under similar experimental conditions using transfected

ND7/23 cells (Leipold et al., 2013; Leipold et al., 2015). We implemented the VGSC isoform

proportions as reported for human CMi-fibers and Aδ-fibers (Tavares-Ferreira et al., 2022) to

investigate the contributions of individual VGSC isoforms to excitability and AP generation in

two sensory fiber types.

Materials and Methods

Cell culture and cell preparation

Cell lines

Plasmids of VGSC isoforms used in this study were either transfected stably in human

embryonic kidney cell line HEK293 (rat $Na_v1.3$ = $rNa_v1.3$, human $Na_v1.5$ = $hNa_v1.5$, mouse

$Na_v1.6$ = $mNa_v1.6$, $hNa_v1.7$) or transiently in human embryonic kidney cell line HEK293T

($hNa_v1.1$, $hNa_v1.2$) or mouse/rat hybridoma nerve cell line ND7/23 ($hNa_v1.8$).

All used cell lines were cultivated adherently under standard cell culture conditions (i.e., 37°C

and 5 % CO2) in their respective cell culture media and supplements (Table S1). Passaging for

all cell lines was performed twice per week at approximately 80% confluency using Accutase

(Sigma-Aldrich, St. Louis, Missouri, USA) with a 1:5 – 1:10 passaging rate onto cultureware

coated with Geltrex (LDEV-Free, hESC-Qualified; Gibco, Thermo Fisher Scientific, Waltham,

Massachusetts, USA). Only cells with passage numbers below 30 were used for

electrophysiological experiments. All cell lines were regularly tested negative for mycoplasma

contamination.

Plasmids

Plasmid constructs used for transient transfection were amplified using XL10-gold

ultracompetent cells (Agilent Technologies Inc., Santa Clara, California, USA) and verified by

restriction pattern analysis and commercial DNA sequencing (Eurofins Genomics GmbH,

Ebersberg, Germany). The plasmids for $hNa_v1.1$ (pCMV6-AC-$hNa_v1.1$ WT RRSSV 3' UTR,

Myc-DDK) and $hNa_v1.2$ (pCMV6-XL5-$hNa_v1.2$ WT RRSSV 3' UTR) were kindly gifted by

Frank Bosmans, Ghent University, Belgium. $hNa_v1.1$ contained the common natural variation

A1067T (allele frequency for T: 0.728). The $hNa_v1.8$ plasmid (pIRES puro3-$hNa_v1.8$ WT)

contained the V1073A variant (allele frequency for A: 0.58).

Transient transfection

HEK293T and ND7/23 cells were seeded in 35 mm petri dishes 20-24 hours prior to transient

transfection to reach a 70-90% confluency. Transfection was then performed using 3 µl jetPEI

(Polyplus-transfection, Illkirch, France), 0.25 µg reporter plasmid pMax-GFP (Lonza, Basel,

Switzerland) and 1.25 µg plasmid encoding the VGSC to be examined. Cells were incubated

for another 18-24 hours before being split onto fresh 35 mm petri dishes. About 2 hours later,

after the seeded cells attached to the cultureware, cells with a bright green fluorescence were

used for patch clamp experiments.

Electrophysiology

General patch clamp framework

Electrophysiological experiments were performed in whole-cell voltage clamp mode using an

EPC10-USB amplifier (HEKA Elektronik GmbH, Lambrecht/Pfalz, Germany). Cells were

covered in extracellular solution (ECS) containing 140 mM NaCl, 3 mM KCl, 1 mM $MgCl_2$, 1

mM $CaCl_2$, 10 mM HEPES and 20 mM Glucose. ECS was adjusted to pH 7.4 using NaOH,

while osmolarity was between 305 and 312 mosm/l. Since ND7/23 cells display endogenous

VGSC-mediated TTXs currents (Rogers et al., 2016; Lee et al., 2019), 500 nM TTX (Tocris,

Bristol, United Kingdom) was added to the ECS for measurements of the TTXr channel

$hNa_v1.8$.

Glass pipettes from borosilicate glass tubes (Biomedical Instruments, Zöllnitz, Germany) were

manufactured and fire-polished using a DMZ pipette puller (Zeitz Instruments GmbH,

Martinsried, Germany). Pipette tip resistance (R_{pip}) was between 0.8 and 2 MΩ for all

measurements. The pipette was filled with intracellular solution (ICS) containing 10 mM NaCl,

140 mM cesium fluoride (CsF), 1 mM EGTA, 10 mM HEPES and 18 mM Sucrose. ICS was

adjusted to pH 7.31 with CsOH, while osmolarity was 313 mosm/l.

Experiments were conducted at room temperature (22 ± 1°C). Cells were kept in ECS for a

maximum of 90 minutes before being discarded. Cell membrane capacitance was between 5

and 33 pF. For all measurements, seal resistance (R_{seal}) was above 0.8 GΩ. Only cells with an

initial series resistance (R_s) below 5 MΩ and a peak current amplitude above 500 pA during the

voltage dependence of activation protocol were accepted. At all times, voltage error (R_s * I)

was below 5 mV.

Data were gathered using Patchmaster Next version 1.2 (HEKA Elektronik GmbH,

Lambrecht/Pfalz, Germany). Analog signals were digitized at a sampling rate of 100 kHz

(except for AP clamping measurements, which were sampled at 20 kHz) with a 10 kHz Bessel

low-pass filter. Leak currents were measured and subtracted online (for activation, steady-state

fast inactivation (SSFI) and ramp current measurements) or offline (for AP clamping due to

technical limitations) with a *P/n* procedure (*n* = 4) preceding the respective test pulse.

Capacitive currents were cancelled, and R_s was compensated for by 77-83% for all cells. All

measurements were corrected online for a liquid junction potential of 8.2 mV.

Patch clamp protocols

For each cell, stimulation protocols were only recorded once (no technical replicates). Protocols

are indicated in the respective figures. For all protocols, the holding potential in between test

pulses (V_{hold}) was set to -120 mV.

After establishing whole-cell configuration, a current stabilization protocol was conducted for

5 minutes to ensure complete recovery, in which cells were held at holding potential and

repeatedly stimulated with test pulses to 0 mV every 10 seconds.

To investigate the voltage dependence of activation, a series of 40 ms pulses from holding

potential to a test pulse potential was elicited, with the test pulse potential increasing stepwise

from -90 mV to +40 mV in 10 mV increments (-60 mV to +70 mV for most $Na_v1.8$

measurements). The interpulse interval was 5 s. For assessment of the voltage dependence of

activation (i.e., the fraction of channels activated to an open state depending on the membrane

potential), we calculated the voltage dependent sodium conductance (G_{Na}) for each test pulse,

using the equation

4
$$G_{Na} = \frac{I_{Na}}{V_m - E_{rev}}$$

5   (Eq. 1)

with I_{Na} being the peak inward current at the respective test pulse voltage V_m, and E_{rev} being the

sodium reversal potential individually determined for each cell. To exclude effects of varying

cell size and current density, G_{Na} values were normalized to the maximum conductance $G_{Na, max}$

of that cell, before being plotted against test pulse voltages.

During the voltage dependence of SSFI protocol, after being held at holding potential, cells

were initially stimulated with a 500 ms prepulse, before being exposed to a 40 ms test pulse at

0 mV. From sweep to sweep, the prepulse voltage increased stepwise from -140 mV to -30 mV

in 10 mV increments (-110 mV to 0 mV for most $Na_v1.8$ measurements). The interpulse interval

was 10 s. For investigation of the voltage dependence of SSFI (i.e., the fraction of non-

inactivated channels available for activation depending on the preceding membrane potential),

the peak inward current for each test pulse (I_{Na}) was normalized to the maximum inward current

during inactivation measurement ($I_{Na,max}$). Normalized current values were plotted against the

respective prepulse potential.

Voltage dependence of activation conductance-voltage curves and voltage dependence of SSFI

current-voltage curves were fitted with non-linear regression using the Boltzmann equation

21
$$\frac{G_{Na}}{G_{Na,max}} \ or \ \frac{I_{Na}}{I_{Na,max}} = bottom + \frac{top - bottom}{1 + e^{\frac{V_{50} - V_m}{k}}}$$

22   (Eq. 2)

with V_{50} being the membrane potential at half-maximal channel (in)activation in mV, V_m being

the membrane voltage in mV and k being the slope factor in 1/mV. The top value of the sigmoid

curve was constrained to 1 for both activation and inactivation measurements. The bottom value

was constrained to 0 for activation measurements but was not constrained for inactivation

measurements to examine the fraction of channels remaining open after SSFI. The goodness of

fit was $R^2 > 0.9$ for all Boltzmann fits.

We equated the window current for each cell as the area under the curve (AUC) of the

superimposed activation and SSFI Boltzmann fit curves. Because SSFI Boltzmann fits do not

reach 0 values when approaching infinity voltages, the AUC was calculated with an upper limit

set 20 mV above the intersection of activation and SSFI Boltzmann fits for each cell.

For ramp current measurements, cells were stimulated with depolarizing voltage ramps from

holding potential to +20 mV at different ramp rates, namely 0.1, 0.2, 0.4, 0.6, 1.2, 2, 4, and 6

mV/ms (for ramp durations see Table S2). For each ramp rate, the measured current response

was averaged from three consecutive sweeps. The maximum inward current and the voltage at

which it occurs were measured, as well as the area under the curve (AUC) of the current

response. Because especially for $Na_v1.8$, the ramp current bell curve did not return to 0 pA at

the end of the ramp, the AUC was calculated only up to the maximum inward current.

Maximum inward current and AUC were normalized to the respective maximum inward current

from the activation protocol for each cell ($I_{Na,max}$).

To measure the VGSC current response to APs, three different pre-recorded APs (AP1 – AP3)

measured in iPSC-derived nociceptors were used as a voltage command (Blair and Bean, 2002).

The AP commands differ in their characteristics (Table 1) with e.g., subthreshold voltage slope

being between 0.2 and 2 mV/ms. These voltage stimuli included 50 ms of the respective resting

membrane potential preceding each recorded AP (Table 1) to prevent transient currents caused

by step depolarization from holding potential to initial AP voltage from interfering with the

examined AP current responses. The respective AP current response was averaged out of three

consecutive sweeps. For each stimulation, we calculated the maximum inward current and AUC

of both the total AP and of the subthreshold phase of the AP. For the latter, the slope of the

elicited current was calculated as well. Current responses for ramp and AP clamping were

normalized to the maximum inward current during activation measurement ($I_{Na,max}$).

	AP1	AP2	AP3
average resting membrane potential (mV)	-76.5	-71.9	-67.4
AP threshold voltage (mV)	-60.1	-61.2	-41
maximum AP voltage (mV)	10	11.2	59.3
voltage amplitude of AP (mV)	86.5	83.1	126.7
minimum voltage of afterhyperpolarization (mV)	-71.6	-67.7	-59.7
AP half width (ms)	3.6	2.4	3.8
time to AP peak (ms)	99.6	14.6	18
subthreshold slope (mV/ms)	0.21	1.39	1.84
upstroke slope (mV/ms)	3.31	10.64	27.1

Table 1: Properties of the Pre-Recorded Action Potentials Used as Voltage Commands. Data were gathered from current clamp measurements of iPSC-derived nociceptors using the Igor Pro software.

Sample sizes and data exclusion

We performed an a-priori analysis to estimate the number of cells needed for each experiment.

In a small pre-study with $Na_v1.3$ and $Na_v1.7$, the voltage at which the maximum inward ramp

current appeared was compared, assuming normal distribution. Using the software G*Power

version 3.1.9.7 (Heinrich-Heine-Universität, Düsseldorf, Germany) we calculated the sample

size for each VGSC to be $n = 27$ (*hedge's g** = 1.23 as a measure of the effect size *d*, two-tailed

$\alpha = 0.05$ and power $(1-\beta) = 0.95$).

We designed the study to have groups of equal size. Group size variation might occur due to

cells being excluded from analysis a-posteriori, e.g., because of measurement artifacts impeding

data analysis. Since the ramp current responses often showed a considerable drift, which made

it difficult to calculate the AUC, the following exclusion criteria were defined for ramp current

AUC measurements: cells with a drift larger than +100 pA or -500 pA as well as cells with a

drift larger than 10% of the maximum inward current during activation measurement were

excluded from analysis. For AP clamping measurements, we identified outliers using the ROUT

method with $Q = 0.1\%$ (Motulsky and Brown, 2006). Outliers were excluded from further

analysis. Beyond that, we performed no outlier analysis or elimination.

Data analysis

Data analysis for electrophysiological experiments was blinded except for $Na_v1.5$. Data

gathered by electrophysiological experiments were analysed using FitMaster version 2x90.5

(HEKA Electronik GmbH, Lambrecht, Germany) and IgorPro version 6.37 (WaveMetrics Inc.,

Portland, Oregon, USA). IgorPro analysis procedures are available upon reasonable request.

GraphPad Prism version 9.5.1 (GraphPad Software, San Diego, California, USA) was used for

Boltzmann fits and statistical testing. Graphing of data was perfomed with IgorPro, GraphPad

Prism and CorelDRAW 2017 version 19.0.0.328 (Alludo, Ottawa, Ontario, Canada).

Statistics

Data are presented as mean ± standard deviation (SD) unless stated otherwise. Error bars denote

the SD, while box plot whiskers show the 10^{th} and 90^{th} percentile. Threshold for statistical

significance was set to $p < 0.05$. Given p values are shown with 4 decimal places and

summarized as follows: ns = not significant, $* = p < 0.05$, $** = p < 0.01$, $*** = p < 0.001$, ****

$= p < 0.0001$.

Data were tested for normal distribution using the D'Agostino & Pearson omnibus K2 test.

Normal distribution for an experiment was only assumed if each group passed the test

individually. For statistical testing, groups were compared by ordinary one-way ANOVA with

Tukey-Kramer's multiple comparisons test and a single pooled variance for parametric testing

or Kruskal-Wallis test with Dunn's multiple comparison test for non-parametric testing. Values

derived from ramp current measurements (I_{max}, $U_{location\ of\ Imax}$ and AUC) were compared by

ordinary two-way ANOVA with Tukey-Kramer's multiple comparisons test and individual

variances computed for each comparison, to compare between both VGSC isoforms and an

additional factor of influence, i.e., the ramp rate. For correlations, we computed the

nonparametric Spearman correlation with a two-tailed p value (data are given as Spearman's r

with 95% CI) and graphed it with simple linear regression. Data were neither matched nor

paired.

The exact group n values are not shown in the figures for readability purposes but are indicated

in the respective tables and supplementary tables.

Computational modelling

To simulate the AP of a CMi- and an Aδ-fiber we use a modified version of the Hodgkin-

Huxley model. Compared to the original Hodgkin-Huxley model (Hodgkin and Huxley, 1952)

we introduced a delay for current decay (fast inactivation kinetics). This modification prevents

premature inactivation of sodium channels and improves the fit of the model to sodium currents

of $Na_v1.1$ - $Na_v1.3$ and $Na_v1.5$ - $Na_v1.8$ for activation voltages above 10 mV. The gating

variables and the delay of inactivation were fitted using voltage clamp data for $Na_v1.1$ - $Na_v1.3$

and $Na_v1.5$ - $Na_v1.8$ acquired at room temperature (22 ± 1°C) as well as previously published

voltage clamp data for $Na_v1.9$ (Leipold et al., 2013; Leipold et al., 2015). To fit $Na_v1.9$ kinetics,

the original Hodgkin-Huxley equations were used because they accurately captured the data.

The fits were performed with *fmincon* from the Matlab software R2021b (The MathWorks, Inc.,

Natick, Massachusetts, USA) using a multistart approach with latin hypercube sampling and a

weighted least squares cost functional. To simulate APs in a CMi-fiber and an Aδ-fiber the

ratios between the maximal conductances for the considered VGSC isoforms were chosen to

match recently published putative gene expressions quantified in human DRGs by spatial

transcriptomics (Tavares-Ferreira et al., 2022). The simulations are based on the quantifications

of gating variables at room temperature. The resting potential is -71 mV, and the model is at

rest before different stimulation currents are applied. Details, model equations and parameters

are provided in the supplementary information.

Results

Na$_v$1.5 and Na$_v$1.8 activate and fast inactivate at more hyper- and depolarized potentials than

other subtypes, respectively

To gain a comparable oversight of the biophysics of VGSC isoforms involved in nociception,

we examined Na$_v$1.1, Na$_v$1.2, Na$_v$1.3, Na$_v$1.5, Na$_v$1.6, Na$_v$1.7 and Na$_v$1.8 for their voltage

dependence of activation and steady-state fast inactivation (SSFI, Fig. 1 and 2). Of all channels

examined two stood out: Na$_v$1.5 activated and fast inactivated at more hyperpolarized potentials

than all other VGSC isoforms (Fig. 1G and 2A/B, Table 2 and S3), whereas Na$_v$1.8 was set

apart by its more depolarized activation and SSFI (Fig. 1J and 2A/B, Table 2 and S3). The

TTXs VGSCs examined activated at intermediate voltages with Na$_v$1.6 and Na$_v$1.7 being

grouped at slightly more hyperpolarized voltages than Na$_v$1.1, Na$_v$1.2 and Na$_v$1.3 (Fig. 1 and

2A/B, Table 2 and S3). Data obtained from fitting conductance-voltage or current-voltage

curves to the Boltzmann equation are summarized in Fig. 2, Table 2, and Table S3-S6.

Comparison of the Boltzmann equation fitting results showed differences between the VGSC

isoforms in all obtained parameters ($p < 0.0001$ for all ANOVA or Kruskal-Wallis tests).

	voltage dependence of activation			voltage dependence of SSFI				intersection of activation and SSFI		
						fraction of channels remaining open after SSFI (%)		Boltzmann fit curves	window current	
	$V50_{act}$ (mV)	k_{act} (1/mv)	n	$V50_{inact}$ (mV)	k_{inact} (1/mV)		n	(mV)	(AUC in mV)	n
Na$_v$1.1	-28.9 ± 4.3	8.3 ± 1.1	27	-74.3 ± 5.0	-5.6 ± 1.1	4.3% ± 5.5%	27	-56.1 ± 4.6	1.398 ± 1.054	27
Na$_v$1.2	-29.2 ± 5.6	8.0 ± 1.0	27	-77.9 ± 6.0	-5.8 ± 0.9	3.3% ± 4.5%	27	-57.7 ± 5.6	1.073 ± 0.861	27
Na$_v$1.3	-29.5 ± 4.9	7.8 ± 0.9	27	-76.2 ± 5.9	-6.7 ± 0.7	2.1% ± 3.5%	27	-54.7 ± 4.3	0.989 ± 0.661	27
Na$_v$1.5	-53.3 ± 3.5	7.8 ± 0.7	27	-104.5 ± 5.7	-5.6 ± 0.7	5.9% ± 5.7%	27	-83.1 ± 4.6	1.484 ± 1.156	27
Na$_v$1.6	-33.8 ± 4.2	7.3 ± 0.5	27	-77.9 ± 5.2	-6.9 ± 1.4	5.0% ± 3.3%	27	-56.6 ± 4.7	1.598 ± 0.608	27

Na$_v$1.7	-33.3 ± 5.6	7.9 ± 0.8	27	-84.9 ± 6.3	-6.6 ± 1.3	5.5% ± 6.2%	27	-61.6 ± 5.6	1.533 ± 1.195	27
Na$_v$1.8	-3.1 ± 4.5	11.8 ± 0.9	25	-66.6 ± 5.4	-9.3 ± 1.4	7.1% ± 5.5%	26	-38.9 ± 5.1	2.401 ± 1.079	25
Na$_v$1.9*	-52.3 ± 9.8	11.0 ± 3.2	24	-68.9 ± 10.3	-12.2 ± 4.9	22.4% ± 9.8%	17	-61.0 ± 6.3	12.952 ± 4.114	15

Table 2: **Descriptive Statistics of Parameters Obtained From Fitting Voltage Dependence of Activation and SSFI With the Boltzmann Equation.** Data are shown as mean ± SD. *Na$_v$1.9 Boltzmann values were derived from pooled previously published data from Leipold et al. 2013 and Leipold et al. 2015.

Figure 1: Conductance-Voltage or Current-Voltage Relationship of Activation and SSFI Protocols. *A* overview of all conductance-voltage curves derived from activation measurements. *B* overview of all current-voltage curves derived from SSFI measurements (voltage protocol of SSFI depicted in inlet. Note that for Na$_v$1.8 measurements, most cells were stimulated with a -110 mV to 0 mV voltage range). *C* example current traces of Na$_v$1.7 elicited by the activation protocol (voltage protocol depicted in inlet. Note that for Na$_v$1.8 measurements, most cells were stimulated with a -60 mV to +70 mV voltage range). *D-J* overlay of activation and inactivation curves for Na$_v$1.1 (*D*), Na$_v$1.2 (*E*), Na$_v$1.3 (*F*), Na$_v$1.5 (*G*), Na$_v$1.6 (*H*), Na$_v$1.7 (*I*) and Na$_v$1.8 (*J*). Data are shown as mean ± SD and superimposed Boltzmann fit curve.

Figure 2: **Activation and SSFI Parameters and Window Current.** *A* V_{50} values obtained from activation Boltzmann fits. *B* V_{50} values obtained from SSFI Boltzmann fits. *C* AUC values underneath the superimposed activation and SSFI Boltzmann fit curves, i.e., the window current. *Inlet:* schematic depiction of AUC calculation, *x* being the membrane voltage of the activation and SSFI Boltzmann fit curve intersection. *D* slope values *k* obtained from activation Boltzmann fits. *E* slope values *k* gathered from SSFI Boltzmann fits. *F* bottom values gathered from SSFI Boltzmann fits, indicating the fraction of channels remaining open after complete SSFI. Data are shown as mean ± SD. Statistical significance from multiple comparisons has not been indicated in the figure panels for readability purposes but can be consulted in Table S3-S6.

The overlap between activation and SSFI curves is often referred to as window current (Wang et al., 1996; Magistretti and Alonso, 1999; Abriel et al., 2001; Osteen et al., 2017). The intersection of activation and SSFI curves (i.e., the peak of the window current) of Na$_v$1.7 was approximately 5 mV more hyperpolarized than that for all other TTXs channels tested except for Na$_v$1.2 (Fig. 1, Table 2, Table S5). This indicates that in this heterologous expression system a higher fraction of Na$_v$1.7 channels are in an open state (i.e., activated and not yet fast inactivated) at more hyperpolarized voltages than the other assumed subthreshold channels Na$_v$1.1, Na$_v$1.3 and Na$_v$1.6, and thus could contribute to cell depolarization at subthreshold voltages.

Window current size was larger for Na$_v$1.8 than for every other VGSC examined except for

Na$_v$1.6. Among TTXs channel isoforms, Na$_v$1.6 elicited slightly higher window currents than

Na$_v$1.2 and Na$_v$1.3 but did not differ from Na$_v$1.1 and Na$_v$1.7 (Fig. 2C, Table 2, Table S5),

underlining their equal potential to support subthreshold depolarization. Concerning the

fraction of channels remaining open after SSFI, we observed a trend for Na$_v$1.2/3 and Na$_v$1.8

to smaller or larger currents, respectively (Fig. 2F, Table 2, Table S6).

To complete our data set with the last remaining VGSC essential for nociceptors, we analyzed

pooled Na$_v$1.9 data from previously published recordings (Leipold et al., 2013; Leipold et al.,

2015) (Fig. 3, Table 2). While activation of Na$_v$1.9 occurred at hyperpolarized voltages

comparable to Na$_v$1.5, SSFI voltage dependence resembled the depolarized inactivation of

Na$_v$1.8 (Fig. 3B/C, Table 2). Na$_v$1.9 displayed pronounced persistent currents (Fig. 3A) as

described previously (Herzog et al., 2001; Maingret et al., 2008). These properties enable the

formation of major window currents exceeding those of all other VGSC isoforms measured in

this study (Fig. 3E, Table 2).

Figure 3**: Summary of Na$_v$1.9 Activation and SSFI Data.** Data from Leipold et al. 2013 and Leipold et al. 2015 were pooled and analyzed concerning Na$_v$1.9 gating and window currents. *A* example current traces of Na$_v$1.9 elicited by the activation protocol. Note that cells were stimulated with a -127 mV to +63 mV voltage range. *B* overlay of activation conductance-voltage and SSFI current-voltage curves. *C* V_{50} values obtained from activation and SSFI Boltzmann fits. *D* slope values *k* obtained from activation and SSFI Boltzmann fits. *E* AUC values underneath the superimposed activation and SSFI Boltzmann fit curves, i.e., the window current. Data are shown as mean ± SD.

Ramp and window currents of only some VGSC isoforms may arise from a similar mechanism

Ramp stimuli evoked a bell-shaped inward current response for all tested VGSC isoforms (Fig.

4), as reported previously (Cummins et al., 1998; Abriel et al., 2001; Herzog et al., 2003; Power

et al., 2012; Estacion and Waxman, 2013; DeCaen et al., 2014; Han et al., 2015a; Zhang et al.,

2017). For cells expressing $Na_v1.3$, we observed two peaked ramp currents for slowly

depolarizing ramps (i.e., 0.1 mV/ms and partly 0.2 mV/ms) in six out of a total of 27 recordings,

similar to what was reported previously (Estacion and Waxman, 2013) (see Fig. S1). For the

remaining cells and higher ramp rates, a delayed current decay was observed, potentially

corresponding to $Na_v1.3$ persistent current.

Figure 4: **Example Traces of $Na_v1.7$ Ramp Currents Elicited by Slowly Depolarizing Ramp Stimuli.** Voltage protocol depicted in inlet. Scale bars apply to current traces only.

Ramp currents for all tested isoforms increased with steeper ramp stimuli, most likely due to

more channels activating before the onset of fast inactivation. As determined in post-hoc

testing, for the lowest ramp rates (i.e., ≤ 0.4 mV/ms), most VGSC isoforms did not differ in

maximum inward ramp current; only $Na_v1.8$ elicited higher maximum ramp currents than the

other VGSC isoforms (Table S7). For medium ramp rates (0.6 – 1.2 mV/ms), $Na_v1.3$ and $Na_v1.8$

exceed the other VGSC isoforms, whereas for the highest ramp rates (2 – 6 mV/ms), $Na_v1.6$

and $Na_v1.7$ gradually catch up, leaving $Na_v1.1$, $Na_v1.2$, and $Na_v1.5$ grouped at lower values

(Fig. 5A, Table S7/S8).

Figure 5: **Current-Voltage Parameters Obtained From Ramp Current Measurements.** *A* maximum inward current from ramp current measurement normalized to maximum inward current from activation measurements. *B* voltage at which the maximum inward current occurred during ramp current measurement. Data are shown as box plots with whiskers indicating the 10[th] and 90[th] percentile and dots for measurements below the 10[th] and above the 90[th] percentile.

We observed that the maximum inward current evoked by ramp stimulation was affected more

by the ramp rate than by the VGSC isoform: the difference between VGSC isoforms ($F(6, 1429)$

$= 153.0$, $p < 0.0001$) and ramp rate ($F(7, 1429) = 974.8$, $p < 0.0001$) made up for 9.7% and

71.8% of total variation, respectively (Fig. 5A, Table S7). Interaction between these terms

accounted for 3.3% of total variation ($F(42, 1429) = 7.528$, $p < 0.0001$).

Ancillary to the size of the maximum inward ramp current, we saw differences in its voltage

dependence (Fig. 5B, Table S7) between VGSC subtypes ($F(6, 1429) = 1748$, $p < 0.0001$;

82.5% of total variation), ramp rates ($F(7, 1429) = 90.15$, $p < 0.0001$; 5% of total variation) and

their interaction ($F(42, 1429) = 5.17$, $p < 0.0001$; 1.7% of total variation). Using multiple

comparison testing, VGSC isoforms showed a distribution matching the results of

activation/SSFI V_{50} results and Boltzmann fit intersections for the vast majority, with $Na_v1.5$

and $Na_v1.8$ ramp currents occurring at more hyperpolarized and depolarized voltages,

respectively, than all other TTXs ramp currents tested (Table S7/S9). TTXs channels are

grouped at intermediate values, with $Na_v1.6$ and $Na_v1.7$ shifted towards more hyperpolarized

values than $Na_v1.1$, $Na_v1.2$, and $Na_v1.3$. For steeper ramps, this effect intensified, especially for

$Na_v1.7$ and $Na_v1.3$.

As a measure of charge added to a cell during slow depolarizations such as the subthreshold

phase, we investigated the AUC underneath ramp currents (Fig. 6A, Table S7). Post-hoc testing

evidenced that $Na_v1.8$ had larger AUC values than the other VGSC isoforms for all ramp rates

(Table S10). No distinct differences in the AUC between the remaining channel isoforms were

observed for lower ramp rates (≤ 0.6 mV/ms). With rising ramp rates, the AUC of $Na_v1.3$,

$Na_v1.5$, $Na_v1.6$, and $Na_v1.7$ gradually surpasses the AUC of $Na_v1.1$ and $Na_v1.2$. The AUC is

affected by VGSC isoform ($F(6, 1180) = 95.75$, $p < 0.0001$; 15% of total variation), ramp rate

($F(7, 1180) = 281.6$, $p < 0.0001$; 51.4% of total variation) and their interaction ($F(42, 1180) =$

1.984, $p = 0.0002$; 2.2% of total variation).

Figure 6: **AUC Values From Ramp Current Measurements Correlate With Window Current Measurements for Some VGSC Isoforms.** *A* AUC values obtained from ramp current measurements, normalized to the maximum inward current from activation measurements for each cell (inlet depicts a schematic example of an AUC underneath a ramp current curve. Note that the AUC was calculated only up to the point of maximum inward current to make up for $Na_v1.8$ ramp currents being cut off at +20 mV). Data are shown as box plots with whiskers indicating the 10[th] and 90[th] percentile. *B-H* correlation between AUC values from ramp current measurements and window current values from activation/SSFI measurements. Data are shown as AUC values plotted against the window current values of the respective cell with a linear regression line for each ramp rate

Some authors refer to window current as corresponding to persistent or ramp current (Blair and

Bean, 2002; Enomoto et al., 2006; Matsutomi et al., 2006; Vasylyev and Waxman, 2012;

Estacion and Waxman, 2013), although so far we are unaware of a study investigating this

potential link. We used Spearman's rank correlation to test for a relationship between window

current and ramp current AUC (Fig. 6B-H, Table S11). For $Na_v1.1$ and $Na_v1.2$, a positive

correlation was identified for all ramp rates. While $Na_v1.5$, $Na_v1.7$ and $Na_v1.8$ exhibited a

positive correlation for most, but not all ramp rates, correlation was scarce for $Na_v1.3$ and

$Na_v1.6$. Note that smaller $Na_v1.5$ sample sizes due to the established exclusion criteria might

limit its correlation accuracy. Our results indicate that ramp and window currents correlate only

in specific cases and are thus likely to be generated by distinct mechanisms.

All tested VGSCs activate during the AP subthreshold phase except for $Na_v1.8$

We used single AP recordings from human iPSC-derived nociceptors (control cells from

Meents et al. (2019)) as voltage command to determine the contribution of each VGSC to the

phases of the AP. The three pre-recorded APs (AP1, AP2 and AP3) differ in their subthreshold

properties, their width and size of overshoot (Fig. 7A/C/E; Table 1).

The timepoint during the AP at which the maximum inward current is elicited differed between

certain VGSC isoforms for all AP commands (Fig. 7B/D/F; Table S12 and S13; $p < 0.0001$ for

all Kruskal-Wallis tests), as did the maximum inward current sizes (Fig. 8A-C, Table S12, $p <$

0.0001 for all Kruskal-Wallis tests) and the area under the current response curve (Fig. 8D-F,

Table S12; $p < 0.0001$ for all Kruskal-Wallis tests).

Figure 7: **Maximum Inward Current Occurs at Different Timepoints During Action Potential Clamping for Each VGSC Isoform.** *A/C/E* action potential waveforms (AP1 – AP3, respectively) recorded from iPSC-derived nociceptors used as voltage commands in AP clamping, with representative current traces, all from the same $Na_v1.7$ cell, charted underneath. *B/D/F* timepoint at which the maximum inward current during the action potential (AP1 – AP3, respectively) occured. *A'-F'* zoomed in depictions of figures *A-F* (excluding the current example traces). Data are shown as mean ± SD. Statistical significance from multiple comparisons has not been indicated in the figure panels for readability purposes but can be consulted in Table S13.

Figure 8: **Current and AUC Parameters Obtained From Action Potential Clamping**. *A-C* maximum inward current during the total command action potential (AP1-AP3, respectively) (inlets indicate the action potential command). *D-F* area under the current response curve to the total command action potential (AP1-AP3, respectively). *G-I* maximum inward current during the subthreshold phase of the command action potential (AP1-AP3, respectively). *J-L* area under the current response curve to the subthreshold phase of the command action potential (AP1-AP3, respectively). *M-O* slope of the inward current measured during the subthreshold phase of the command action potential (AP1-AP3, respectively). Data are normalized to the maximum inward current of activation measurement of the correspondent cell and shown as mean ± SD. Statistical significance from multiple comparisons has not been indicated in the figure panels for readability purposes but can be consulted in Table S13.

Multiple comparisons revealed that $Na_v1.8$ currents were elicited at later timepoints than all

other tested VGSC isoforms except for $Na_v1.5$, namely after the AP peak (Table S13). $Na_v1.5$

currents occurred slightly later within the AP time course as well, whereas $Na_v1.3$ leaned

towards earlier timepoints in the subthreshold phase (Fig. 7B'/D'F', Table S12 and S13).

Current sizes and AUC sizes were similarly distributed between the AP commands, with $Na_v1.8$

and $Na_v1.3$ exhibiting larger and $Na_v1.5$ exhibiting slightly lower values for maximum inward

current (Fig. 8A-F, Table S12 and S13).

The examination of the timepoint of maximum inward current during AP3 stood out since some

VGSC isoforms displayed results with a bimodal distribution (Fig. 7F'). Post-hoc data revision

revealed a transient current peak during the upstroke of the AP command in varying degrees

for $Na_v1.1$-1.3 and $Na_v1.7$, most likely caused by insufficient pipette capacitance compensation.

When this transient current exceeded the more or less evenly pronounced subthreshold influx,

the measured timepoint was shifted, forming the second cluster depicted in Fig. 7F'. For $Na_v1.8$,

which also displayed a slightly clustered distribution, no similar effect could be observed. By

excluding the measurements impeded by above mentioned transients, channel distribution

became even more distinct: $Na_v1.1$-1.3 and $Na_v1.7$ had their maximum influx during the

subthreshold phase of the AP command, considerably before $Na_v1.5$, $Na_v1.6$ and $Na_v1.8$ (Fig.

S2, Table S13).

During the subthreshold phase of the APs maximum inward current, AUC and inward current

slope differed between the VGSC isoforms ($p < 0.0001$ for all Kruskal-Wallis tests except for

AP1 subthreshold AUC ($p = 0.0008$)). On average, maximum inward currents increased with

increasing subthreshold slope of the AP commands (Fig. 8G-I, Table S12). Post-hoc testing

indicates that $Na_v1.8$ and $Na_v1.5$ elicited little to no current during the subthreshold phase, while

maximum inward current was higher for $Na_v1.3$ (AP1 and AP3), $Na_v1.7$ (AP2 and AP3), and,

for AP1, $Na_v1.6$ (Fig. 8G-I, Table S13). Especially for AP3 with the steepest subthreshold slope,

$Na_v1.3$ currents increased drastically (Fig. 8I, Table S12 and S13). The remaining TTXs

isoforms had intermediate maximum inward current responses.

A similar pattern was observed for the subthreshold phase AUC (Fig. 8J-L, Table S12 and S13).

$Na_v1.5$ and $Na_v1.8$ AUC values remained close to zero for all AP commands (note that negative

AUC values occur due to undulation of the current base rate and/or drift). $Na_v1.7$ and $Na_v1.3$

displayed larger subthreshold AUC values (the latter particularly for AP3), thereby adding a

larger charge to a cell during subthreshold depolarization and potentially alleviating initiation

of an AP.

As another measure of channel contribution to subthreshold depolarization, we determined the

slope of currents elicited during the subthreshold phase of AP clamping by each VGSC isoform

(Fig. 8M-O, Table S12 and S13). Steeper subthreshold slopes were documented for $Na_v1.7$

(AP2 and AP3), $Na_v1.6$ (AP1 and AP2), and, with emphasis on AP3, $Na_v1.3$ compared to the

other tested channels. In contrast, $Na_v1.5$ and $Na_v1.8$ subthreshold slopes were minor. The data

are in line with the assumption that $Na_v1.8$ is mainly activated during the upstroke of a

nociceptor AP, while TTXs channels (predominantly $Na_v1.7$, $Na_v1.6$, and – especially for

steeper subthreshold depolarizations – $Na_v1.3$) contribute in earlier, subthreshold phases.

In silico simulation suggests significant contribution of $Na_v1.9$ to the fast upstroke of the AP and

its shoulder

With the large data set in hand, which contains patch-clamp recordings of all VGSCs relevant

in nociceptors, we now aimed to obtain insights into the contribution of each isoform to the AP

by *in silico* experiments. Our computational model comprises the sodium currents of $Na_v1.1$ –

$Na_v1.3$, $Na_v1.5$ – $Na_v1.8$ (derived from the current trace measurements of this study), $Na_v1.9$

(derived from current trace measurements of Leipold et al. (2013) and Leipold et al. (2015))

and a generic potassium and leak current (see Methods and Supplements).

In order to generate an adequate fit of the recorded current traces of the different VGSCs, we

implemented a modification to the original Hodgkin-Huxley model: we introduced a waiting

state after pulse onset due to which the fast inactivation gate (denoted as h in the Hodgkin-

Huxley framework) is delayed (see Methods and Supplements). This waiting state could mimic

the time during which VGSC may open and are not ready to inactivate yet. This modification

led to improved trace fitting for $Na_v1.1$-1.3 and $Na_v1.5$-1.8, as can be seen in the reduced

deviation from the original trace (see Supplements). The Hodgkin-Huxley-like models of each

sodium channel subtype were integrated into a non-spatial model of an Aδ-fiber and a CMi-

fiber, which were used to simulate APs in response to current injections (see Methods and

Supplements).

To model CMi-fibers (Fig. 9/S3) and Aδ-fibers (Fig. 10/S4), we accounted for the fiber-specific

expression patterns of VGSCs quantified in Tavares-Ferreira et al. (2022), as these sequencing

results provide well accessible, reliable human expression data of all voltage-gated sodium

channels in sensory neurons.

Figure 9: **Computer Simulations of APs in a CMi-fiber.** The pie charts indicate the abundance of different VGSC isoforms. APs are triggered by an injection of 30 μA/cm² which persists until the end of the simulation. *A* The upper left panel shows the AP for the relative abundance of VGSC isoforms quantified in spatial gene expression. The initial condition of the simulation is the resting state of the fiber. The other panels show changes of the AP resulting from removal of Na$_v$1.9 (upper right), Na$_v$1.8 (middle left), Na$_v$1.7 (middle right) and Na$_v$1.1-1.3, 1.5 & 1.6 (bottom left). The VGSC isoforms are coded by the colors specified in the bottom right panel. *B/C* VGSC isoform contributions to the simulated AP depicted in the upper left panel of *A*. The upper panels of *B* and *C* show the membrane potential during the initial 3 ms (*B*) and 1.5 ms (*C*) of the simulated AP. The middle panels show the relative contribution of the different sodium currents, the potassium current and the leak current over the course of the AP, plotted as stacked individual currents normalized to total current at each time point. The bottom panels show VGSC isoform contributions to the total sodium current, plotted as stacked individual VGSC isoform currents normalized to the total sodium current at each time point. The black line visualizes the shape of the AP. The color coding is identical to that in *A*.

Figure 10: **Computer Simulations of APs in a Aδ-fiber.** The pie charts indicate the abundance of different VGSC isoforms. APs are triggered by an injection of 30 µA/cm² which persists until the end of the simulation. *A* The upper left panel shows the AP for the relative abundance of VGSC isoforms quantified in spatial gene expression. The initial condition of the simulation is the resting state of the fiber. The other panels show changes of the AP resulting from removal of Na$_v$1.9 (upper right), Na$_v$1.8 (middle left), Na$_v$1.7 (middle right) and Na$_v$1.1-1.3, 1.5 & 1.6 (bottom left). The VGSC isoforms are coded by the colors specified in the bottom right panel. *B/C* VGSC isoform contributions to the simulated AP depicted in the upper left panel of *A*. The upper panels of *B* and *C* show the membrane potential during the initial 3 ms (*B*) and 1.5 ms (*C*) of the simulated AP. The middle panels show the relative contribution of the different sodium currents, the potassium current and the leak current over the course of the AP, plotted as stacked individual currents normalized to total current at each time point. The bottom panels show VGSC isoform contributions to the total sodium current, plotted as stacked individual VGSC isoform currents normalized to the total sodium current at each time point. The black line visualizes the shape of the AP. The color coding is identical to that in *A*.

The maximal total voltage-gated sodium, potassium and leak conductivities were chosen to

recapitulate the experimentally observed AP morphology. In the simulations, the APs were

triggered by current injection of 30 µA/cm² (Fig. 9/10) or 40 µA/cm² (Fig. S3/S4) throughout

the duration of the experiment. Both fiber types revealed an AP with overshoot and a shoulder

(stronger for CMi-fibers) upon stimulation with 30 µA/cm² (Fig. 9A/10A) and 40 µA/cm² (Fig.

S3A/S4A).

To assess the role of a specific VGSC isoform, we removed it from the model and observed the

resulting AP changes (Fig. 9A/10A/S3A/S4A/S5). Notably, the removal of an isoform resulted

in a reduction of the total sodium conductance.

The most striking finding of our simulations is the strong contribution of $Na_v1.9$ to the fast

upstroke of the AP as well as to the formation of a shoulder in both fiber types (Fig. 9B/C, 10

B/C, S3B/C, S4B/C, S5, S6). Omission of $Na_v1.9$ at low stimulation intensities aborted AP

generation, very similar to omission of $Na_v1.7$ (Fig. 9A/10A). The latter simulates *loss-of-*

function as it may occur for $Na_v1.7$ in patients suffering from chronic insensitivity to pain

(Lampert et al., 2014). While removal of $Na_v1.8$ from the model resulted in AP waveforms that

failed to overshoot, this intervention did not affect formation of the AP shoulder, suggesting

that other subtypes such as $Na_v1.9$ may contribute to shoulder formation (Fig. 9A/10A).

To investigate this further, we reduced the conductance of $Na_v1.9$ gradually in our CMi-model,

while increasing that of the other sodium channels, to keep the overall VGSC conductance

constant (Fig. S6). Reducing $Na_v1.9$ gradually flattened the shoulder; the fast upstroke of the

AP turned less steep and subthreshold depolarizations following the AP disappeared.

Additionally, the afterhyperpolarization was reduced, which may reflect a weaker activation of

voltage-gated potassium channels due to the vanishing of the shoulder. Removal of $Na_v1.1$ –

1.3, $Na_v1.5$ and $Na_v1.6$ (at the same time) had no significant impact on the AP of the CMi-fiber

but reduced the overshoot in the Aδ-fiber (Fig. 9A/10A/S3A/S4A/S5).

Fig. 9B/C and 10B/C show the relative contributions of the different VGSC isoforms to the

simulated APs of the CMi-fiber and the Aδ-fiber, respectively. In both fiber types the relative

contribution of $Na_v1.7$ is maximal right before or at the beginning of the fast upstroke of the

AP (Fig. 9C, 10C, S3C, S4C), in line with reports on $Na_v1.7$ as subthreshold or threshold

channel (Rush et al., 2007; Vasylyev and Waxman, 2012; Meents et al., 2019). The largest

relative (Fig. 9B/C, 10B/C), but also absolute (Fig. S7) conduction by far in our two models is

mediated by $Na_v1.9$. Most of its relative contribution occurs during subthreshold

depolarizations, then during the fast upstroke of the AP (relative and absolute), and, with a

prominent second peak (relative and absolute), during the falling phase of the AP coinciding

with the shoulder.

$Na_v1.8$, on the other hand, shows largest conductivity during the upstroke, the peak and the

falling phase of the AP (Fig. 9B/C, 10B/C, S3B/C, S4B/C, S7), but when the AP shoulder

occurs, it seems to be already partially inactivated. The relative and absolute contribution of

$Na_v1.8$ and $Na_v1.7$ to the AP, especially during the peak, seems to be flipped between the two

fiber models investigated here: while in the CMi-fiber model $Na_v1.7$ contributes more, it is

$Na_v1.8$ in the Aδ-fiber model. The other channels show only minor contributions during the

course of the AP: $Na_v1.5$ shows some activity during subthreshold depolarizations and a small

peak during the shoulder in the CMi-fiber model, while $Na_v1.6$ is active at the AP threshold in

both models. $Na_v1.6$ is continuously active throughout the AP in the Aδ model (Fig. 9B/C,

10B/C, S3B/C, S4B/C, S7).

Gain-of-function mutations of $Na_v1.7$ linked to inherited pain syndromes often modify channel

gating such that during an AP more channels are activated. Erythromelalgia mutations in $Na_v1.7$

lead to hyperpolarized activation of the channel, inducing hyperexcitability when

overexpressed in rodent sensory neurons (Körner and Lampert, 2020) and enhance spontaneous

activity in human stem cell derived nociceptors (Mis et al., 2019). We used our computational

model to simulate *gain-of-function* in two ways: 1) by increasing the maximal conductance of

$Na_v1.7$ and comparing it to an increased $Na_v1.8$ conductance and 2) by shifting activation of

$Na_v1.7$ to more hyperpolarized potentials.

When we increased $Na_v1.7$ conduction by 5 times in our computer model, both, the CMi-fiber

model and the Aδ-fiber model showed persistent firing in response to an ongoing current

injection of 1 $\mu A/cm^2$ (Fig. 11A/12A). Interestingly, enhancing $Na_v1.8$ conduction by 11 times

(revealing similar overall conductance as for a five-fold increased $Na_v1.7$ conductance), was

unable to induce ongoing AP firing (Fig. 11B/12B). This finding supports the concept that the

specific properties of $Na_v1.7$ and not the increase of the total sodium conductance trigger the

repetitive activity observed in Fig. 11A and 12A.

Figure 11: **Computational Disease Modelling in a CMi-fiber.** *A* Repetitive depolarizations resulting from increasing the maximal conductance for Na$_v$1.7 to 500% of the wildtype value. The current injection of 1 µA/cm² persists until the end of the simulation. *B* An increase of the maximal conductance for Na$_v$1.8 to 1100% of the wildtype value results in an isolated AP. The total sodium conductances in *A* and *B* are approximately the same. The current injections of 30 µA/cm² and 40 µA/cm² persist until the end of the simulations. The pie charts indicate the abundance of different VGSC isoforms. *C* Current injections of 10 µA/cm² and 15 µA/cm² trigger no APs in the simulated wildtype CMi (isoform abundances as in Fig 9A, upper left panel). *D* If the activation of Na$_v$1.7 is shifted by 5 mV to hyperpolarized potentials, a current injection of 15 µA/cm² but not of 10 µA/cm² triggers an AP. *E* If the activation of Na$_v$1.7 is shifted by 10 mV to hyperpolarized potentials, current injections of 10 µA/cm² and 15 µA/cm² trigger an AP. The current injections in *C-E* persist until the

Figure 12: **Computational Disease Modelling in an Aδ fiber.** *A* Repetitive depolarizations resulting from increasing the maximal conductance for Na$_v$1.7 to 500% of the wildtype value. The current injection of 1 µA/cm² persists until the end of the simulation. *B* An increase of the maximal conductance for Na$_v$1.8 to 800% of the wildtype value results in an isolated AP. The total sodium conductances in *A* and *B* are approximately the same. The current injections of 30 µA/cm² and 40 µA/cm² persist until the end of the simulations. The pie charts indicate the abundance of different VGSC isoforms. *C* Current injections of 10 µA/cm² and 15 µA/cm² trigger no APs in the simulated wildtype CMi (isoform abundances as in Fig 10A, upper left panel). *D* If the activation of Na$_v$1.7 is shifted by 5 mV to hyperpolarized potentials, a current injection of 15 µA/cm² but not of 10 µA/cm² triggers an AP. *E* If the activation of Na$_v$1.7 is shifted by 10 mV to hyperpolarized potentials, current injections of 10 µA/cm² and 15 µA/cm² trigger an AP. The current injections in *C-E* persist until the

With unshifted $Na_v1.7$ activation, subthreshold stimulations of 10 or 15 $\mu A/cm^2$ did not evoke

an AP in the CMi-fiber model (Fig. 11C), and neither did subthreshold stimulations of 8 or 10

$\mu A/cm^2$ in the Aδ-fiber model (Fig. 12C). When $Na_v1.7$ activation was shifted to hyperpolarized

potentials by only 5 mV – which is often observed in erythromelagia mutations (e.g., Lampert

et al. (2009)) – an AP was initiated by small stimulations, showing a reduced firing threshold

(Fig. 11D/12D). When shifted by 10 mV, weak stimulations even induced an unstable

membrane potential leading to several strong, repetitive depolarizations (Fig. 11E/12E).

Taken together, our simulations including Hodgkin-Huxley-like models of all nociceptive

sodium channels suggest a contribution of $Na_v1.9$ to the fast upstroke of the AP and its shoulder

formation and proved useful in modelling excitability changes associated with $Na_v1.7$-linked

chronic pain syndromes such as erythromelalgia.

Discussion

In this study, we provide a detailed, comparative investigation of the gating properties of all

VGSCs relevant in nociceptors with focus on subthreshold depolarizations and propose

computational models to predict their contributions to AP electrogenesis in two types of sensory

neurons: CMi-fibers and Aδ-fibers. The models suggest that $Na_v1.9$ may actively contribute to

AP generation and formation of its shoulder in these neurons.

VGSC isoforms are characterized by their distinct gating behavior *in vitro* and *in silico*

The TTXr channels measured in this study, i.e., $Na_v1.5$ and $Na_v1.8$, showed divergent gating

properties with shifted voltage dependence of activation, SSFI and ramp currents toward

hyperpolarized and depolarized voltages, respectively. While $Na_v1.9$ activation occurred at very

hyperpolarized voltages, voltage dependence of $Na_v1.9$ SSFI resembled the depolarized

inactivation of $Na_v1.8$. Out of the remaining TTXs channels, which all had their voltage

dependence of activation, SSFI and ramp currents at subthreshold voltages, especially $Na_v1.3$

and $Na_v1.7$ elicited prominent currents in ramp and AP clamping. During the subthreshold

phases of AP clamping, both displayed large AUC values and steeper current slopes compared

to the other VGSCs tested. For $Na_v1.7$, we observed slightly more hyperpolarized voltage

dependences for both, SSFI and ramp currents, than for other TTXs channel isoforms,

potentially enabling $Na_v1.7$ to charge a cell at subthreshold or threshold voltages.

Our experiments emphasized the importance of $Na_v1.7$ as AP initiator, also since knocking out

or enhancing $Na_v1.7$ *in silico* led to omitted or multiple APs, respectively. This resembles

putative *loss-* and *gain-of-function* pathophysiology that is attributed to $Na_v1.7$ channelopathies

(Dib-Hajj et al., 2005; Cox et al., 2006; Faber et al., 2012; Bennett et al., 2019; McDermott et

al., 2019). Shifting activation of $Na_v1.7$ to more hyperpolarized potentials in models for CMi-

or Aδ-fibers revealed the experimentally observed threshold reduction, suggesting the high

potential of our simulations for disease modelling.

Our findings confirmed $Na_v1.8$ contribution to the AP upstroke (Renganathan et al., 2001;

Matsutomi et al., 2006; Rush et al., 2007; Vasylyev and Waxman, 2012), as it activates and

fast-inactivates at more depolarized potentials, elicits large ramp currents at more depolarized

potentials and showed little to no subthreshold activity in AP clamping compared to the other

VGSCs tested. Interestingly, in our simulations including $Na_v1.9$, the contribution of $Na_v1.8$ to

the shoulder was minor: knocking out $Na_v1.8$ in the *in silico* experiments left APs in the CMi-

model almost unchanged, while in the Aδ-model the overshoot was reduced. This is in line with

findings from $Na_v1.8$ knockout DRGs, where maximum AP voltages were lower compared to

wildtype DRGs (Renganathan et al., 2001; Harty and Waxman, 2007). Yet, the initiation of the

AP was not impeded when $Na_v1.8$ was omitted from the established model.

The strongest difference between the two fibers modelled in this study is the inverse relative

contribution of $Na_v1.7$ compared to $Na_v1.8$ during the AP: for the CMi-fiber model, $Na_v1.7$ has

the larger current and conductivity, whereas for the Aδ-fiber model it is $Na_v1.8$. In an earlier

modelling study, we investigated the contribution of $Na_v1.7$, $Na_v1.8$ and $Na_v1.9$ to the

excitability of a peripheral C-fiber, to reproduce axonal spike propagation speed and activity-

dependent slowing as recorded in microneurography experiments (Tigerholm et al., 2014;

Tigerholm et al., 2015). We showed that the speed of two APs following each other depends

on the influx of sodium through $Na_v1.7$ relative to $Na_v1.8$ current. Thus, fiber specific

biophysical properties may result from the specific contribution of the sodium channels

expressed in their membrane.

Our simulations suggest that $Na_v1.9$, in addition to its activation during rest, strongly

contributes to the fast upstroke and is a major determinant of the shoulder of the AP. It

represents by far the largest current and conductivity in our model of all integrated channels.

The findings are in line with current-clamp recordings on enteric neurons from $Na_v1.9$ knockout

mice which show no shoulder in APs from mice lacking $Na_v1.9$ (Osorio et al., 2014). An early

report on the sensory neuron AP properties of those knockout mice revealed no differences in

their AP width (Priest et al., 2005). The latter study did not differentiate between sensory neuron

subtypes, yet in the realm of recent transcriptomic advances we know that $Na_v1.9$ expression

differs strongly between subgroups (Tavares-Ferreira et al., 2022). It is also known that not all

sensory neuron subtypes display a shoulder in their AP (Körner and Lampert, 2022). Thus,

pooling the groups may have hidden the effect of $Na_v1.9$ on the AP shoulder in this study. As

$Na_v1.9$ channelopathies are linked to both gain- and loss-of-pain syndromes (Leipold et al.,

2013; Huang et al., 2017), clarification of the fiber type-specific role of $Na_v1.9$ during AP

generation could help to understand their complex pathophysiology.

We documented large $Na_v1.3$ currents during clamping of an AP with steep subthreshold

voltage slope (AP3), and – consistent with literature – during steeper voltage ramps (Cummins

et al., 2001). This might play a role in neuropathic pain, as – though only marginally expressed

in C- and Aδ-fibers – some studies suggest that $Na_v1.3$ is upregulated in spinal cord injuries or

peripheral nerve injuries in rodents and therefore increases excitability (Hains et al., 2003;

Hains et al., 2004; Lampert et al., 2006). Other studies present more ambiguous or even

contradictory results, e.g., $Na_v1.3$ knockout mice did not show lowered pain thresholds after

nerve injury than wildtype mice (Lindia et al., 2005; Nassar et al., 2006).

$Na_v1.6$ stood out to some extent, as it showed higher maximum inward currents and steeper

current slopes during the subthreshold phase of AP clamping than other TTXs VGSC isoforms.

Furthermore, $Na_v1.6$ current was larger in the Aδ-fiber than in the CMi-fiber during the course

of the AP. Yet, omitting $Na_v1.1$, $Na_v1.2$ and $Na_v1.6$ during the *in silico* testing had no noticeable

effect on AP morphology, likely because of their minor share in the spatial transcriptomics-

derived expression patterns used in our model (Tavares-Ferreira et al., 2022). $Na_v1.6$ has been

linked to resurgent currents (O'Brien and Meisler, 2013), and these currents seem to be

enhanced in A-fibers compared to C-fibers (Klinger et al., 2012). This is suggested as a reason

why sea-anemone toxin ATX-II led to painful sensations by enhancing resurgent currents in A-

fibers, but not in C-fibers (Klinger et al., 2012).

$Na_v1.5$ played no major role during subthreshold AP clamping and in contributing to the

simulated AP, while it is also only marginally detected in C- and Aδ-fibers (Tavares-Ferreira

et al., 2022). Due to its hyperpolarized activation and SSFI, it may be responding in cells with

more hyperpolarized membrane potential. During development, $Na_v1.5$ is the third TTXr

conductance present and may have a more prominent role than observed in the two fiber types

described here (reflecting the transcriptomic expression pattern of adult DRG donors). The role

of $Na_v1.5$ during development, injury and regeneration needs to be investigated in more detail

in follow-up studies (Renganathan et al., 2002; Tseng et al., 2010; Sousounis et al., 2020).

Correlation between ramp current and window current is channel subtype-specific

Slowly depolarizing voltage ramps induce ramp currents, mimicking the subthreshold phase of

an AP. Ramp currents of wild-type channels have been examined for all VGSC isoforms

individually (Cummins et al., 1998; Abriel et al., 2001; Herzog et al., 2003; El-Bizri et al., 2011;

Power et al., 2012; Estacion and Waxman, 2013; DeCaen et al., 2014; Han et al., 2015a; Han

et al., 2015b; Zhang et al., 2017), but not yet in direct comparison, establishing experimenter-

to-experimenter or lab-to-lab variations. Hereby, we provide an extensive and highly

comparable data set of VGSC ramp currents.

While maximum ramp current size and AUC are predominantly influenced by the steepness of

the inducing voltage ramp stimulus, VGSC subtypes rather influence the voltage dependence

of ramp current responses and hence co-determine in which AP phase they contribute to its

formation.

All tested TTXs channel isoforms elicited their peak ramp current at subthreshold voltages,

enabling them to contribute to cell discharge prior to the AP threshold being exceeded. Of these

subtypes, $Na_v1.3$, $Na_v1.6$ and $Na_v1.7$ elicited the highest maximum ramp currents,

corroborating their role as key players in subthreshold depolarization.

In VGSC literature, ramp and window currents were often assumed to describe a similar gating

mechanism and were sometimes even used interchangeably (Wang et al., 1996; Magistretti and

Alonso, 1999; Abriel et al., 2001; Blair and Bean, 2002; Enomoto et al., 2006; Matsutomi et

al., 2006; Vasylyev and Waxman, 2012; Estacion and Waxman, 2013; Osteen et al., 2017). We

here tested this hypothesis and showed that $Na_v1.1$ and $Na_v1.2$ (partly also $Na_v1.5$, $Na_v1.7$ and

$Na_v1.8$) showed a positive correlation between ramp current and window currents, suggesting

a common underlying mechanism. Yet, because for $Na_v1.3$ and $Na_v1.6$ correlation was mostly

non-existent, we suppose that additional factors such as persistent current play into ramp current

formation, as suggested previously (Estacion and Waxman, 2013).

Expression systems affect *in vitro* VGSC measurements

Heterologous expression systems bear their own interpretation pitfalls, as they do not resemble

the complex environment of primary neurons, e.g., regarding cell morphology, intercellular

contacts, or membrane protein interactions. VGSC isoforms were shown to have different

gating properties when expressed in different expression systems (Cummins et al., 2001; Rush

et al., 2006a; Tan and Soderlund, 2009; Zhang et al., 2017).

In this study, we used both HEK and ND7/23 cells for electrophysiological VGSC examination.

Multiple differences in their respective cell characteristics lead to subtype gating differences

among each other having to be analyzed carefully. While endogenous $Na_v1.6$ and $Na_v1.7$

currents in ND7/23 can easily be suppressed by adding TTX to the extracellular solution while

measuring TTXr isoforms (Rogers et al., 2016; Lee et al., 2019), altered VGSC gating

properties by endogenous expression of β1 and β3 subunits in ND7/23 cells cannot be

eliminated (John et al., 2004; Lee et al., 2019). However, also HEK cells themselves express

β1A subunits endogenously (Moran et al., 2000). Therefore, comparisons between VGSC α

subunit measurements have to be treated carefully nonetheless and additional examination of α

subunits in presence of varying β subunits is encouraged. Furthermore, other cellular proteins

such as fibroblast growth factor homologous factors, microtubule-associated proteins or

aquaporins have been shown to modify VGSC gating and/or current density (Wittmack et al.,

2004; Rush et al., 2006b; Zhang and Verkman, 2010; O'Brien et al., 2012; Vanoye et al., 2013).

Recent comparative studies of $Na_v1.2$ and $Na_v1.6$ variants also highlight the influence of

alternative splicing on channel gating (Thompson et al., 2020; Vanoye et al., 2024).

$Na_v1.8$ expression in HEK is challenging and resulting currents are low (Gladwell et al., 1998;

John et al., 2004). As in our hands, expression of $Na_v1.8$ in HEK293T cells yielded little to no

current, we used ND7/23 cells, equivalently to the $Na_v1.9$ measurements integrated into this

study (Leipold et al., 2013; Leipold et al., 2015). Biophysical characteristics of $Na_v1.8$ currents

were shown to be similar between when expressed in ND7/23 cells or recorded from native

DRG TTXr currents (John et al., 2004).

Even non-native neuronal cell lines such as ND7/23 do not resemble a comprehensive primary

neuronal environment (Yin et al., 2016). Studies in expression systems closer to human

peripheral sensory neurons such as rodent or human DRG neurons could give an even better

insight into the channels biophysics *in vivo*. Moreover, nociceptors derived from human

induced pluripotent stem cells offer the possibility to investigate ion channels in an adaption of

their native environment (Meents et al., 2019; Mis et al., 2019; Namer et al., 2019). This could

also make up for the fact that we included neither co-expression of β-subunits into our

measurements nor did we check for channel dimerization, both of which have shown to alter

biophysical properties of VGSCs (Brackenbury and Isom, 2011; Chahine and O'Leary, 2011;

Clatot et al., 2017; Rühlmann et al., 2020).

Albeit our observations allow unprecedented comparison between isoforms as the experiments

were conducted under the so far best possible comparable conditions, it may be that differing

expression systems do not model certain isoform specificities.

Biophysical differences such as larger persistent or ramp currents can occur between VGSC

isoforms from different mammalian species (Browne et al., 2009; Tan and Soderlund, 2009;

Han et al., 2015a). For $Na_v1.3$ and $Na_v1.6$ measurements in this study, we used rodent channel

isoforms. Hence, complementary measurements with human channel isoforms are desirable.

In this study as well as the studies providing the implemented $Na_v1.9$ data, intracellular solution

contained CsF which is widely used in patch clamp experiments to enhance high seal resistances

for stable and long-lasting patch clamp configurations (Kostyuk et al., 1975; Fernandez et al.,

1984). Yet, usage of CsF has been shown to affect the gating of VGSCs in a subtype-specific

manner. For example, the voltage dependence of activation and inactivation of $Na_v1.9$ is shifted

by about 15-20 mV to more hyperpolarized potentials in presence of CsF (Rugiero et al., 2003;

Coste et al., 2004). The gating of $Na_v1.3$ and $Na_v1.7$ is also affected by intracellular fluoride,

albeit to a lesser extent (Chen et al., 2000; Meadows et al., 2002; Jarecki et al., 2008). While

this does not necessarily limit subtype comparability within our *in vitro* measurements, which

all included CsF, comparison to CsF-free experiments such as measurements in primary

neurons have to be performed cautiously. Even minor shifts in channel biophysics can

profoundly impact complex and therefore vulnerable computational models. Although the

model established in this study provides valuable insights into the AP contribution of each

channel, reissuing the model with data from CsF-free measurements would be desirable to

increase its predictive power.

Limitations of *in silico* modelling

Computational models are simplifications of the real world and therefore must be handled

cautiously, especially when model complexity suggests the opposite. We estimated the

distribution of the maximal sodium conductances per VGSC subunit by using mRNA

expression data from a human DRG spatial transcriptomics study (Tavares-Ferreira et al.,

2022). While this approach is closest to a quantitative measure currently accessible from human

DRGs, mRNA expression levels do not always indicate that the respective channel is also

translated to protein, trafficked to the cell surface and fully functional. Even though correlation

between mRNA expression, presence of VGSCs in the cell membrane and biophysical

properties has been suggested (Thériault and Chahine, 2014), alternative splicing, correct

folding and association with additional proteins or membrane turnover are just some factors

that may impact channel translation from RNA to functional protein in the membrane

(Buccitelli and Selbach, 2020). It is possible that mRNA levels do not directly relate to the

amount of protein produced in the cells (Lindhout et al., 2020), and that we are overestimating

or underestimating the conductance of the VGSCs modelled in this study. As proteomics of

human DRGs, especially for VGSCs, is not yet available as a basis for our modelling, we used

the published transcriptomics data to estimate the conductivity of the VGSC subtypes integrated

in our simulations.

A promising solution to these problems are single-cell transcriptomics combined with

functional analysis, such as Patch-seq, which offer the ability to link expression data directly to

cell physiology (Lipovsek et al., 2021; Körner et al., 2022) or single cell proteomics, which are

currently developed also for human nociceptors. Furthermore, since expression data are

gathered at the cell soma, only limited assumptions can be made for small nerve fibers, e.g., in

the human skin.

Several studies suggested the possibility of neuronal channel composition changing in

compensation for *loss-* or *gain-of-function* mutations in VGSC isoforms, e.g., TTXs currents

being upregulated in DRG neurons of $Na_v1.8$ knockout mice (Akopian et al., 1999;

Renganathan et al., 2000). Compensatory changes could also involve the expression of other

ion channel types, e.g., calcium channels, or pathways indirectly affecting excitability. When

modelling the effect of VGSC *loss-* or *gain-of-function* mutations *in silico*, these compensatory

changes should be considered, as soon as their extent is determined in precedent studies.

VGSC data are usually gathered at room temperature, as it was done in this study. However,

temperature dependence of VGSCs varies for each channel isoform and recorded voltages

(Kriegeskorte et al., 2023). Thus, most computational models published in the recent years

including the one established in this study do not accurately account for temperature, since this

would require additional experimental data and re-parametrization of the models (Almog and

Korngreen, 2016).

For fitting the gating variables of each channel subtype, we used current traces recorded at

several potentials. At voltages below -40 mV it often occurred that multiple parameter

combinations described the currents successfully, creating some uncertainty (unidentifiability).

Particularly in the case of $Na_v1.9$ this may impact on simulation results.

Our computational model describes ion channels based on modified Hodgkin-Huxley dynamics

and neglects spatial aspects. To further study the VGSC influence on AP generation, the VGSC

isoforms should be implemented in a morphologically detailed C-fiber or Aδ-fiber model

(Tigerholm et al., 2014; Tigerholm et al., 2019).

VGSC isoform characterization can be the basis for assessing potential drug targets

In this study we provide a broadly based groundwork for future comparative studies of VGSC

isoforms, their biophysics and interaction with each other. By feeding our measurements into a

computational model, we generated an *in silico* tool to test hypotheses, investigate the

contribution of each nociceptive sodium channel to AP genesis and model pharmacology and

effect of disease-related mutations. This may also prove useful when developing new potential

analgesics such as specific channel blockers or optimizing therapy schemes for chronic pain.

Additional information

Data availability

All data gathered in this study and all analysis procedures used are available from the

corresponding authors upon request.

Disclosures/Competing interests

The authors declare no competing financial interests and no conflicts of interest with the

contents of this article. A.L. had an unrelated research agreement with Grunenthal. A.L.

receives counseling fees from Grunenthal.

Acknowledgements

We thank Brigitte Hoch, Petra Hautvast, and Raya A. Bott for their excellent technical support

(cell culture, molecular biology, electrophysiology). We thank Jannis Körner, Sophia

Kriegeskorte and Jannis E. Meents for their thoughtful discussions as well as Jannis E. Meents

for providing the AP stimuli used in this study.

CRediT author contributions

P. A. K.: Conceptualization, Methodology, Investigation, Software, Validation, Formal

analysis, Data curation, Visualization, Writing - Original draft. E. L.: Resources, Data curation,

Writing - Review & Editing. J. T.: Writing - Review & Editing. A. M.: Writing - Review &

Editing. B. N.: Writing - Review & Editing. T. S.: Resources, Methodology, Investigation,

Software, Validation, Formal analysis, Data curation, Visualization, Writing - Review &

Editing. A. L.: Supervision, Project administration, Funding acquisition, Resources,

Conceptualization, Methodology, Validation, Data curation, Writing - Review & Editing.

Funding/Grants

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation 363055819/GRK2415 Mechanobiology of 3D epithelial tissues (ME3T) to A.L.;

368482240/GRK2416, MultiSenses-MultiScales to A.L., LA2740/6-1 to A.L.; NA 970/5-1 to

B.N.).   A.L. (IZKF TN1-1/IA 532001) and B.N. were supported by a grant from the

Interdisciplinary Center for Clinical Research within the Faculty of Medicine at the RWTH

Aachen University.

Abbreviations

The following abbreviations are used: *AP* action potential, *AUC* area under the curve, *CMi*

mechano-insensitive C-fiber, *CNS* central nervous system, *CsF* cesium fluoride, *DMEM*

Dulbecco's modified Eagle medium, *DRG* dorsal root ganglion, *ECS* extracellular solution, E_{rev}

reversal potential, *FBS* fetal bovine serum, G_{Na} sodium conductance, $G_{Na,max}$ maximum sodium

conductance, *G418* Geneticin disulphate, *ICS* intracellular solution, I_{Na} inward sodium current,

$I_{Na,max}$ maximum inward sodium current, *iPSC* induced pluripotent stem cells, *k* slope factor,

PNS peripheral nervous system, R_{pip} pipette tip resistance, R_s series resistance, R_{seal} seal

resistance, *SD* standard deviation, *SSFI* steady-state fast inactivation, *TTX* tetrodotoxin, *TTXr*

TTX-resistant, *TTXs* TTX-sensitive, *VGSC* voltage-gated sodium channel, V_{hold} holding

potential, V_m membrane potential, V_{50} membrane potential at half-maximal channel

(in)activation.

References

Abriel, H., C. Cabo, X.H. Wehrens, I. Rivolta, H.K. Motoike, M. Memmi, C. Napolitano, S.G. Priori, and
R.S. Kass. 2001. Novel arrhythmogenic mechanism revealed by a long-QT syndrome mutation
in the cardiac Na(+) channel. *Circ Res*. 88:740-745. doi: 10.1161/hh0701.089668.
Ahern, C.A., J. Payandeh, F. Bosmans, and B. Chanda. 2016. The hitchhiker's guide to the voltage-
gated sodium channel galaxy. *J Gen Physiol*. 147:1-24. doi: 10.1085/jgp.201511492.
Akopian, A.N., V. Souslova, S. England, K. Okuse, N. Ogata, J. Ure, A. Smith, B.J. Kerr, S.B. McMahon,
S. Boyce, R. Hill, L.C. Stanfa, A.H. Dickenson, and J.N. Wood. 1999. The tetrodotoxin-resistant
sodium channel SNS has a specialized function in pain pathways. *Nat Neurosci*. 2:541-548.
doi: 10.1038/9195.
Almog, M., and A. Korngreen. 2016. Is realistic neuronal modeling realistic? *J Neurophysiol*. 116:2180-
2209. doi: 10.1152/jn.00360.2016.
Bennett, D.L., A.J. Clark, J. Huang, S.G. Waxman, and S.D. Dib-Hajj. 2019. The Role of Voltage-Gated
Sodium Channels in Pain Signaling. *Physiol Rev*. 99:1079-1151. doi:
10.1152/physrev.00052.2017.
Bennett, D.L., and C.G. Woods. 2014. Painful and painless channelopathies. *Lancet Neurol*. 13:587-
599. doi: 10.1016/s1474-4422(14)70024-9.

Bhattacharjee, A., Y. Xiao, Z. Pei, and T.R. Cummins. 2018. Specialized Sodium Channels in Pain
Transmission. *In*. Oxford University Press, The Oxford Handbook of Neuronal Ion Channels.
doi: 10.1093/oxfordhb/9780190669164.013.3.
Bhuiyan, S.A., M. Xu, L. Yang, E. Semizoglou, P. Bhatia, K.I. Pantaleo, I. Tochitsky, A. Jain, B. Erdogan,
S. Blair, V. Cat, J.M. Mwirigi, I. Sankaranarayanan, D. Tavares-Ferreira, U. Green, L.A.
McIlvried, B.A. Copits, Z. Bertels, J.S. Del Rosario, A.J. Widman, R.A. Slivicki, J. Yi, R. Sharif-
Naeini, C.J. Woolf, J.K. Lennerz, J.L. Whited, T.J. Price, W.G.I. Robert, and W. Renthal. 2024.
Harmonized cross-species cell atlases of trigeminal and dorsal root ganglia. *Sci Adv*.
10:eadj9173. doi: 10.1126/sciadv.adj9173.
Blair, N.T., and B.P. Bean. 2002. Roles of tetrodotoxin (TTX)-sensitive Na+ current, TTX-resistant Na+
current, and Ca2+ current in the action potentials of nociceptive sensory neurons. *J Neurosci*.
22:10277-10290. doi: 10.1523/JNEUROSCI.22-23-10277.2002. .
Bothe, S.N., and A. Lampert. 2021. The insecticide deltamethrin enhances sodium channel slow
inactivation of human Nav1.9, Nav1.8 and Nav1.7. *Toxicol Appl Pharmacol*. 428:115676. doi:
10.1016/j.taap.2021.115676.
Brackenbury, W.J., and L.L. Isom. 2011. Na Channel β Subunits: Overachievers of the Ion Channel
Family. *Front Pharmacol*. 2:53. doi: 10.3389/fphar.2011.00053.
Brouwer, B.A., I.S. Merkies, M.M. Gerrits, S.G. Waxman, J.G. Hoeijmakers, and C.G. Faber. 2014.
Painful neuropathies: the emerging role of sodium channelopathies. *J Peripher Nerv Syst*.
19:53-65. doi: 10.1111/jns5.12071.
Browne, L.E., J.J. Clare, and D. Wray. 2009. Functional and pharmacological properties of human and
rat NaV1.8 channels. *Neuropharmacology*. 56:905-914. doi:
10.1016/j.neuropharm.2009.01.018.
Buccitelli, C., and M. Selbach. 2020. mRNAs, proteins and the emerging principles of gene expression
control. *Nat Rev Genet*. 21:630-644. doi: 10.1038/s41576-020-0258-4.
Catterall, W.A. 2023. Voltage gated sodium and calcium channels: Discovery, structure, function, and
Pharmacology. *Channels*. 17:2281714. doi: 10.1080/19336950.2023.2281714.
Catterall, W.A., A.L. Goldin, and S.G. Waxman. 2005. International Union of Pharmacology. XLVII.
Nomenclature and structure-function relationships of voltage-gated sodium channels.
Pharmacol Rev. 57:397-409. doi: 10.1124/pr.57.4.4.
Chahine, M., and M.E. O'Leary. 2011. Regulatory Role of Voltage-Gated Na Channel β Subunits in
Sensory Neurons. *Front Pharmacol*. 2:70. doi: 10.3389/fphar.2011.00070.
Chen, Y.H., T.J. Dale, M.A. Romanos, W.R. Whitaker, X.M. Xie, and J.J. Clare. 2000. Cloning,
distribution and functional analysis of the type III sodium channel from human brain. *Eur J*
Neurosci. 12:4281-4289. doi:
Clatot, J., M. Hoshi, X. Wan, H. Liu, A. Jain, K. Shinlapawittayatorn, C. Marionneau, E. Ficker, T. Ha,
and I. Deschênes. 2017. Voltage-gated sodium channels assemble and gate as dimers. *Nat*
Commun. 8:2077. doi: 10.1038/s41467-017-02262-0.
Coste, B., N. Osorio, F. Padilla, M. Crest, and P. Delmas. 2004. Gating and modulation of presumptive
NaV1.9 channels in enteric and spinal sensory neurons. *Mol Cell Neurosci*. 26:123-134. doi:
10.1016/j.mcn.2004.01.015.
Cox, J.J., F. Reimann, A.K. Nicholas, G. Thornton, E. Roberts, K. Springell, G. Karbani, H. Jafri, J.
Mannan, Y. Raashid, L. Al-Gazali, H. Hamamy, E.M. Valente, S. Gorman, R. Williams, D.P.
McHale, J.N. Wood, F.M. Gribble, and C.G. Woods. 2006. An SCN9A channelopathy causes
congenital inability to experience pain. *Nature*. 444:894-898. doi: 10.1038/nature05413.
Cummins, T.R., F. Aglieco, M. Renganathan, R.I. Herzog, S.D. Dib-Hajj, and S.G. Waxman. 2001.
Nav1.3 sodium channels: rapid repriming and slow closed-state inactivation display
quantitative differences after expression in a mammalian cell line and in spinal sensory
neurons. *J Neurosci*. 21:5952-5961. doi: 10.1523/JNEUROSCI.21-16-05952.2001.
Cummins, T.R., J.R. Howe, and S.G. Waxman. 1998. Slow closed-state inactivation: a novel
mechanism underlying ramp currents in cells expressing the hNE/PN1 sodium channel. *J*
Neurosci. 18:9607-9619. doi: 10.1523/JNEUROSCI.18-23-09607.1998.

de Lera Ruiz, M., and R.L. Kraus. 2015. Voltage-Gated Sodium Channels: Structure, Function,
Pharmacology, and Clinical Indications. *J Med Chem*. 58:7093-7118. doi: 10.1021/jm501981g.
DeCaen, P.G., Y. Takahashi, T.A. Krulwich, M. Ito, and D.E. Clapham. 2014. Ionic selectivity and
thermal adaptations within the voltage-gated sodium channel family of alkaliphilic Bacillus.
eLife. 3:e04387. doi: 10.7554/eLife.04387.
Dib-Hajj, S.D., T.R. Cummins, J.A. Black, and S.G. Waxman. 2010. Sodium channels in normal and
pathological pain. *Annu Rev Neurosci*. 33:325-347. doi: 10.1146/annurev-neuro-060909-
153234.
Dib-Hajj, S.D., P. Geha, and S.G. Waxman. 2017. Sodium channels in pain disorders: pathophysiology
and prospects for treatment. *Pain*. 158 Suppl 1:S97-s107. doi:
10.1097/j.pain.0000000000000854.
Dib-Hajj, S.D., A.M. Rush, T.R. Cummins, F.M. Hisama, S. Novella, L. Tyrrell, L. Marshall, and S.G.
Waxman. 2005. Gain-of-function mutation in Nav1.7 in familial erythromelalgia induces
bursting of sensory neurons. *Brain*. 128:1847-1854. doi: 10.1093/brain/awh514.
Eberhardt, E., S. Havlicek, D. Schmidt, Andrea S. Link, C. Neacsu, Z. Kohl, M. Hampl, Andreas M. Kist,
A. Klinger, C. Nau, J. Schüttler, C. Alzheimer, J. Winkler, B. Namer, B. Winner, and A. Lampert.
2015. Pattern of Functional TTX-Resistant Sodium Channels Reveals a Developmental Stage
of Human iPSC- and ESC-Derived Nociceptors. *Stem Cell Reports*. 5:305-313. doi:
10.1016/j.stemcr.2015.07.010.
El-Bizri, N., K.M. Kahlig, J.C. Shyrock, J.A.L. George, L. Belardinelli, and S. Rajamani. 2011. Ranolazine
block of human Nav1.4 sodium channels and paramyotonia congenita mutants. *Channels*.
5:161-172. doi: 10.4161/chan.5.2.14851.
Enomoto, A., J.M. Han, C.-F. Hsiao, N. Wu, and S.H. Chandler. 2006. Participation of sodium currents
in burst generation and control of membrane excitability in mesencephalic trigeminal
neurons. *J Neurosci*. 26:3412-3422. doi: 10.1523/JNEUROSCI.5274-05.2006.
Estacion, M., and S.G. Waxman. 2013. The response of NaV1.3 sodium channels to ramp stimuli:
multiple components and mechanisms. *J Neurophysiol*. 109:306-314. doi:
10.1152/jn.00438.2012.
Faber, C.G., J.G. Hoeijmakers, H.S. Ahn, X. Cheng, C. Han, J.S. Choi, M. Estacion, G. Lauria, E.K.
Vanhoutte, M.M. Gerrits, S. Dib-Hajj, J.P. Drenth, S.G. Waxman, and I.S. Merkies. 2012. Gain
of function Nanu1.7 mutations in idiopathic small fiber neuropathy. *Ann Neurol*. 71:26-39.
doi: 10.1002/ana.22485.
Felts, P.A., S. Yokoyama, S. Dib-Hajj, J.A. Black, and S.G. Waxman. 1997. Sodium channel alpha-
subunit mRNAs I, II, III, NaG, Na6 and hNE (PN1): different expression patterns in developing
rat nervous system. *Brain Res Mol Brain Res*. 45:71-82. doi: 10.1016/s0169-328x(96)00241-0.
Fernandez, J.M., A.P. Fox, and S. Krasne. 1984. Membrane patches and whole-cell membranes: a
comparison of electrical properties in rat clonal pituitary (GH3) cells. *J Physiol*. 356:565-585.
doi: 10.1113/jphysiol.1984.sp015483.
Fertleman, C.R., M.D. Baker, K.A. Parker, S. Moffatt, F.V. Elmslie, B. Abrahamsen, J. Ostman, N.
Klugbauer, J.N. Wood, R.M. Gardiner, and M. Rees. 2006. SCN9A mutations in paroxysmal
extreme pain disorder: allelic variants underlie distinct channel defects and phenotypes.
Neuron. 52:767-774. doi: 10.1016/j.neuron.2006.10.006.
Fertleman, C.R., C.D. Ferrie, J. Aicardi, N.A. Bednarek, O. Eeg-Olofsson, F.V. Elmslie, D.A. Griesemer, F.
Goutières, M. Kirkpatrick, I.N. Malmros, M. Pollitzer, M. Rossiter, E. Roulet-Perez, R.
Schubert, V.V. Smith, H. Testard, V. Wong, and J.B. Stephenson. 2007. Paroxysmal extreme
pain disorder (previously familial rectal pain syndrome). *Neurology*. 69:586-595. doi:
10.1212/01.wnl.0000268065.16865.5f.
Fukuoka, T., K. Kobayashi, H. Yamanaka, K. Obata, Y. Dai, and K. Noguchi. 2008. Comparative study of
the distribution of the alpha-subunits of voltage-gated sodium channels in normal and
axotomized rat dorsal root ganglion neurons. *J Comp Neurol*. 510:188-206. doi:
10.1002/cne.21786.

Gladwell, Z., D. Trezise, C. Plumpton, X. Xie, and S. Tate. 1998. Expression and functional analysis of recombinant rat sensory neuron specific Na+ channels in a mammalian cell line. *J Physiol*. 513:141P. doi:

Goodwin, G., and S.B. McMahon. 2021. The physiological function of different voltage-gated sodium channels in pain. *Nat Rev Neurosci*. 22:263-274. doi: 10.1038/s41583-021-00444-w.

Goral, R.O., E. Leipold, E. Nematian-Ardestani, and S.H. Heinemann. 2015. Heterologous expression of NaV1.9 chimeras in various cell systems. *Pflugers Arch*. 467:2423-2435. doi: 10.1007/s00424-015-1709-1.

Hains, B.C., J.P. Klein, C.Y. Saab, M.J. Craner, J.A. Black, and S.G. Waxman. 2003. Upregulation of sodium channel Nav1.3 and functional involvement in neuronal hyperexcitability associated with central neuropathic pain after spinal cord injury. *J Neurosci*. 23:8881-8892. doi: 10.1523/jneurosci.23-26-08881.2003.

Hains, B.C., C.Y. Saab, J.P. Klein, M.J. Craner, and S.G. Waxman. 2004. Altered sodium channel expression in second-order spinal sensory neurons contributes to pain after peripheral nerve injury. *J Neurosci*. 24:4832-4839. doi: 10.1523/jneurosci.0300-04.2004.

Han, C., M. Estacion, J. Huang, D. Vasylyev, P. Zhao, S.D. Dib-Hajj, and S.G. Waxman. 2015a. Human Na(v)1.8: enhanced persistent and ramp currents contribute to distinct firing properties of human DRG neurons. *J Neurophysiol*. 113:3172-3185. doi: 10.1152/jn.00113.2015.

Han, C., Y. Yang, B.T.A. de Greef, J.G.J. Hoeijmakers, M.M. Gerrits, C. Verhamme, J. Qu, G. Lauria, I.S.J. Merkies, C.G. Faber, S.D. Dib-Hajj, and S.G. Waxman. 2015b. The Domain II S4-S5 Linker in Nav1.9: A Missense Mutation Enhances Activation, Impairs Fast Inactivation, and Produces Human Painful Neuropathy. *NeuroMolecular Med*. 17:158-169. doi: 10.1007/s12017-015-8347-9.

Harty, T.P., and S.G. Waxman. 2007. Inactivation Properties of Sodium Channel Nav1.8 Maintain Action Potential Amplitude in Small DRG Neurons in the Context of Depolarization. *Molecular Pain*. 3:1744-8069-1743-1712. doi: 10.1186/1744-8069-3-12.

Herzog, R.I., T.R. Cummins, F. Ghassemi, S.D. Dib-Hajj, and S.G. Waxman. 2003. Distinct repriming and closed-state inactivation kinetics of Nav1.6 and Nav1.7 sodium channels in mouse spinal sensory neurons. *J Physiol*. 551:741-750. doi: 10.1113/jphysiol.2003.047357.

Herzog, R.I., T.R. Cummins, and S.G. Waxman. 2001. Persistent TTX-resistant Na+ current affects resting potential and response to depolarization in simulated spinal sensory neurons. *J Neurophysiol*. 86:1351-1364. doi: 10.1152/jn.2001.86.3.1351.

Hodgkin, A.L., and A.F. Huxley. 1952. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol*. 117:500-544. doi: 10.1113/jphysiol.1952.sp004764.

Huang, J., C.G. Vanoye, A. Cutts, Y.P. Goldberg, S.D. Dib-Hajj, C.J. Cohen, S.G. Waxman, and A.L. George, Jr. 2017. Sodium channel NaV1.9 mutations associated with insensitivity to pain dampen neuronal excitability. *The Journal of clinical investigation*. 127:2805-2814. doi: 10.1172/JCI92373.

Jarecki, B.W., P.L. Sheets, J.O. Jackson, 2nd, and T.R. Cummins. 2008. Paroxysmal extreme pain disorder mutations within the D3/S4-S5 linker of Nav1.7 cause moderate destabilization of fast inactivation. *J Physiol*. 586:4137-4153. doi: 10.1113/jphysiol.2008.154906.

John, V.H., M.J. Main, A.J. Powell, Z.M. Gladwell, C. Hick, H.S. Sidhu, J.J. Clare, S. Tate, and D.J. Trezise. 2004. Heterologous expression and functional analysis of rat NaV1.8 (SNS) voltage-gated sodium channels in the dorsal root ganglion neuroblastoma cell line ND7–23. *Neuropharmacology*. 46:425-438. doi: https://doi.org/10.1016/j.neuropharm.2003.09.018.

Jung, M., M. Dourado, J. Maksymetz, A. Jacobson, B.I. Laufer, M. Baca, O. Foreman, D.H. Hackos, L. Riol-Blanco, and J.S. Kaminker. 2023. Cross-species transcriptomic atlas of dorsal root ganglia reveals species-specific programs for sensory function. *Nat Commun*. 14:366. doi: 10.1038/s41467-023-36014-0.

Kleggetveit, I.P., B. Namer, R. Schmidt, T. Helås, M. Rückel, K. Ørstavik, M. Schmelz, and E. Jørum. 2012. High spontaneous activity of C-nociceptors in painful polyneuropathy. *Pain*. 153:2040-2047. doi: 10.1016/j.pain.2012.05.017.

Klinger, A.B., M. Eberhardt, A.S. Link, B. Namer, L.K. Kutsche, E.T. Schuy, R. Sittl, T. Hoffmann, C. Alzheimer, T. Huth, R.W. Carr, and A. Lampert. 2012. Sea-anemone toxin ATX-II elicits A-fiber-dependent pain and enhances resurgent and persistent sodium currents in large sensory neurons. *Mol Pain*. 8:69. doi: 10.1186/1744-8069-8-69.

Klugbauer, N., L. Lacinova, V. Flockerzi, and F. Hofmann. 1995. Structure and functional expression of a new member of the tetrodotoxin-sensitive voltage-activated sodium channel family from human neuroendocrine cells. *Embo j*. 14:1084-1090. doi: 10.1002/j.1460-2075.1995.tb07091.x.

Körner, J., N. Haag, I. Kurth, and A. Lampert. 2022. Genetics meets function in sodium channel-related pain disorders. *Neuroforum*. 28:67-75. doi: doi:10.1515/nf-2021-0035.

Körner, J., and A. Lampert. 2020. Sodium channels. *In* The Senses - A Comprehensive Reference. B. Fritzsch, editor. Academic Press. 120-141.

Körner, J., and A. Lampert. 2022. Functional subgroups of rat and human sensory neurons: a systematic review of electrophysiological properties. *Pflugers Arch*. 474:367-385. doi: 10.1007/s00424-021-02656-6.

Körner, J., J. Meents, J.P. Machtens, and A. Lampert. 2018. β1 subunit stabilises sodium channel Nav1.7 against mechanical stress. *J Physiol*. 596:2433-2445. doi: 10.1113/jp275905.

Kostyuk, P.G., O.A. Krishtal, and V.I. Pidoplichko. 1975. Effect of internal fluoride and phosphate on membrane currents during intracellular dialysis of nerve cells. *Nature*. 257:691-693. doi: 10.1038/257691a0.

Kriegeskorte, S., R. Bott, M. Hampl, A. Korngreen, R. Hausmann, and A. Lampert. 2023. Cold and warmth intensify pain-linked sodium channel gating effects and persistent currents. *J Gen Physiol*. 155. doi: 10.1085/jgp.202213312.

Kupari, J., D. Usoskin, M. Parisien, D. Lou, Y. Hu, M. Fatt, P. Lönnerberg, M. Spångberg, B. Eriksson, N. Barkas, P.V. Kharchenko, K. Loré, S. Khoury, L. Diatchenko, and P. Ernfors. 2021. Single cell transcriptomics of primate sensory neurons identifies cell types associated with chronic pain. *Nat Commun*. 12:1510. doi: 10.1038/s41467-021-21725-z.

Laezza, F., A. Lampert, M.A. Kozel, B.R. Gerber, A.M. Rush, J.M. Nerbonne, S.G. Waxman, S.D. Dib-Hajj, and D.M. Ornitz. 2009. FGF14 N-terminal splice variants differentially modulate Nav1.2 and Nav1.6-encoded sodium channels. *Mol Cell Neurosci*. 42:90-101. doi: 10.1016/j.mcn.2009.05.007.

Lampert, A., S.D. Dib-Hajj, E.M. Eastman, L. Tyrrell, Z. Lin, Y. Yang, and S.G. Waxman. 2009. Erythromelalgia mutation L823R shifts activation and inactivation of threshold sodium channel Nav1.7 to hyperpolarized potentials. *Biochem Biophys Res Commun*. 390:319-324. doi: 10.1016/j.bbrc.2009.09.121.

Lampert, A., M. Eberhardt, and S.G. Waxman. 2014. Altered sodium channel gating as molecular basis for pain: contribution of activation, inactivation, and resurgent currents. *Handb Exp Pharmacol*. 221:91-110. doi: 10.1007/978-3-642-41588-3_5.

Lampert, A., B.C. Hains, and S.G. Waxman. 2006. Upregulation of persistent and ramp sodium current in dorsal horn neurons after spinal cord injury. *Exp Brain Res*. 174:660-666. doi: 10.1007/s00221-006-0511-x.

Lampert, A., A.O. O'Reilly, P. Reeh, and A. Leffler. 2010. Sodium channelopathies and pain. *Pflugers Arch*. 460:249-263. doi: 10.1007/s00424-009-0779-3.

Lee, J., S. Kim, H.-m. Kim, H.J. Kim, and F.H. Yu. 2019. NaV1.6 and NaV1.7 channels are major endogenous voltage-gated sodium channels in ND7/23 cells. *PloS one*. 14:e0221156. doi: 10.1371/journal.pone.0221156.

Leipold, E., A. Hanson-Kahn, M. Frick, P. Gong, J.A. Bernstein, M. Voigt, I. Katona, R. Oliver Goral, J. Altmüller, P. Nürnberg, J. Weis, C.A. Hübner, S.H. Heinemann, and I. Kurth. 2015. Cold-

aggravated pain in humans caused by a hyperactive NaV1.9 channel mutant. *Nat Commun*. 6:10049. doi: 10.1038/ncomms10049.

Leipold, E., L. Liebmann, G.C. Korenke, T. Heinrich, S. Gießelmann, J. Baets, M. Ebbinghaus, R.O. Goral, T. Stödberg, J.C. Hennings, M. Bergmann, J. Altmüller, H. Thiele, A. Wetzel, P. Nürnberg, V. Timmerman, P. De Jonghe, R. Blum, H.-G. Schaible, J. Weis, S.H. Heinemann, C.A. Hübner, and I. Kurth. 2013. A de novo gain-of-function mutation in SCN11A causes loss of pain perception. *Nature Genetics*. 45:1399. doi: 10.1038/ng.2767.

Liang, L., S. Fazel Darbandi, S. Pochareddy, F.O. Gulden, M.C. Gilson, B.K. Sheppard, A. Sahagun, J.Y. An, D.M. Werling, J.L.R. Rubenstein, N. Sestan, K.J. Bender, and S.J. Sanders. 2021. Developmental dynamics of voltage-gated sodium channel isoform expression in the human and mouse brain. *Genome Med*. 13:135. doi: 10.1186/s13073-021-00949-0.

Lindhout, F.W., R. Kooistra, S. Portegies, L.J. Herstel, R. Stucchi, B.L. Snoek, A.M. Altelaar, H.D. MacGillavry, C.J. Wierenga, and C.C. Hoogenraad. 2020. Quantitative mapping of transcriptome and proteome dynamics during polarization of human iPSC-derived neurons. *Elife*. 9. doi: 10.7554/eLife.58124.

Lindia, J.A., M.G. Köhler, W.J. Martin, and C. Abbadie. 2005. Relationship between sodium channel NaV1.3 expression and neuropathic pain behavior in rats. *Pain*. 117:145-153. doi: 10.1016/j.pain.2005.05.027.

Lipovsek, M., C. Bardy, C.R. Cadwell, K. Hadley, D. Kobak, and S.J. Tripathy. 2021. Patch-seq: Past, Present, and Future. *J Neurosci*. 41:937-946. doi: 10.1523/jneurosci.1653-20.2020.

Magistretti, J., and A. Alonso. 1999. Biophysical properties and slow voltage-dependent inactivation of a sustained sodium current in entorhinal cortex layer-II principal neurons: a whole-cell and single-channel study. *J Gen Physiol*. 114:491-509. doi: 10.1085/jgp.114.4.491.

Maingret, F., B. Coste, F. Padilla, N. Clerc, M. Crest, S.M. Korogod, and P. Delmas. 2008. Inflammatory mediators increase Nav1.9 current and excitability in nociceptors through a coincident detection mechanism. *J Gen Physiol*. 131:211-225. doi: 10.1085/jgp.200709935.

Matsutomi, T., C. Nakamoto, T. Zheng, J. Kakimura, and N. Ogata. 2006. Multiple types of Na(+) currents mediate action potential electrogenesis in small neurons of mouse dorsal root ganglia. *Pflugers Arch*. 453:83-96. doi: 10.1007/s00424-006-0104-3.

Maxion, A., E. Kutafina, M.F. Dohrn, P. Sacré, A. Lampert, J. Tigerholm, and B. Namer. 2023. A modelling study to dissect the potential role of voltage-gated ion channels in activity-dependent conduction velocity changes as identified in small fiber neuropathy patients. *Front Comput Neurosci*. 17:1265958. doi: 10.3389/fncom.2023.1265958.

McDermott, L.A., G.A. Weir, A.C. Themistocleous, A.R. Segerdahl, I. Blesneac, G. Baskozos, A.J. Clark, V. Millar, L.J. Peck, D. Ebner, I. Tracey, J. Serra, and D.L. Bennett. 2019. Defining the Functional Role of NaV1.7 in Human Nociception. *Neuron*. 101:905-919.e908. doi: 10.1016/j.neuron.2019.01.047.

Meadows, L.S., Y.H. Chen, A.J. Powell, J.J. Clare, and D.S. Ragsdale. 2002. Functional modulation of human brain Nav1.3 sodium channels, expressed in mammalian cells, by auxiliary beta 1, beta 2 and beta 3 subunits. *Neuroscience*. 114:745-753. doi: 10.1016/s0306-4522(02)00242-7.

Meents, J.E., E. Bressan, S. Sontag, A. Foerster, P. Hautvast, C. Rosseler, M. Hampl, H. Schuler, R. Goetzke, T.K.C. Le, I.P. Kleggetveit, K. Le Cann, C. Kerth, A.M. Rush, M. Rogers, Z. Kohl, M. Schmelz, W. Wagner, E. Jorum, B. Namer, B. Winner, M. Zenke, and A. Lampert. 2019. The role of Nav1.7 in human nociceptors: insights from human induced pluripotent stem cell-derived sensory neurons of erythromelalgia patients. *Pain*. 160:1327-1341. doi: 10.1097/j.pain.0000000000001511.

Mis, M.A., Y. Yang, B.S. Tanaka, C. Gomis-Perez, S. Liu, F. Dib-Hajj, T. Adi, R. Garcia-Milian, B.R. Schulman, S.D. Dib-Hajj, and S.G. Waxman. 2019. Resilience to Pain: A Peripheral Component Identified Using Induced Pluripotent Stem Cells and Dynamic Clamp. *J Neurosci*. 39:382-392. doi: 10.1523/jneurosci.2433-18.2018.

Moran, O., M. Nizzari, and F. Conti. 2000. Endogenous expression of the β1A sodium channel subunit
in HEK-293 cells. *FEBS Letters*. 473:132-134. doi: https://doi.org/10.1016/S0014-
5793(00)01518-0.
Motulsky, H.J., and R.E. Brown. 2006. Detecting outliers when fitting data with nonlinear regression -
a new method based on robust nonlinear regression and the false discovery rate. *BMC*
Bioinformatics. 7:123. doi: 10.1186/1471-2105-7-123.
Namer, B., and H.O. Handwerker. 2009. Translational nociceptor research as guide to human pain
perceptions and pathophysiology. *Exp Brain Res*. 196:163-172. doi: 10.1007/s00221-009-
1777-6.
Namer, B., D. Schmidt, E. Eberhardt, M. Maroni, E. Dorfmeister, I.P. Kleggetveit, L. Kaluza, J. Meents,
A. Gerlach, Z. Lin, A. Winterpacht, E. Dragicevic, Z. Kohl, J. Schuttler, I. Kurth, T. Warncke, E.
Jorum, B. Winner, and A. Lampert. 2019. Pain relief in a neuropathy patient by lacosamide:
Proof of principle of clinical translation from patient-specific iPS cell-derived nociceptors.
EBioMedicine. 39:401-408. doi: 10.1016/j.ebiom.2018.11.042.
Nassar, M.A., M.D. Baker, A. Levato, R. Ingram, G. Mallucci, S.B. McMahon, and J.N. Wood. 2006.
Nerve injury induces robust allodynia and ectopic discharges in Nav1.3 null mutant mice. *Mol*
Pain. 2:33. doi: 10.1186/1744-8069-2-33.
Nguyen, M.Q., L.J. von Buchholtz, A.N. Reker, N.J. Ryba, and S. Davidson. 2021. Single-nucleus
transcriptomic analysis of human dorsal root ganglion neurons. *Elife*. 10. doi:
10.7554/eLife.71752.
O'Brien, J.E., and M.H. Meisler. 2013. Sodium channel SCN8A (Nav1.6): properties and de novo
mutations in epileptic encephalopathy and intellectual disability. *Front Genet*. 4:213. doi:
10.3389/fgene.2013.00213.
O'Brien, J.E., L.M. Sharkey, C.N. Vallianatos, C. Han, J.C. Blossom, T. Yu, S.G. Waxman, S.D. Dib-Hajj,
and M.H. Meisler. 2012. Interaction of voltage-gated sodium channel Nav1.6 (SCN8A) with
microtubule-associated protein Map1b. *J Biol Chem*. 287:18459-18466. doi:
10.1074/jbc.M111.336024.
Osorio, N., S. Korogod, and P. Delmas. 2014. Specialized functions of Nav1.5 and Nav1.9 channels in
electrogenesis of myenteric neurons in intact mouse ganglia. *J Neurosci*. 34:5233-5244. doi:
10.1523/jneurosci.0057-14.2014.
Osteen, J.D., K. Sampson, V. Iyer, D. Julius, and F. Bosmans. 2017. Pharmacology of the Nav1.1
domain IV voltage sensor reveals coupling between inactivation gating processes. *Proc Natl*
Acad Sci U S A. 114:6836-6841. doi: 10.1073/pnas.1621263114.
Petersson, M.E., O. Obreja, A. Lampert, R.W. Carr, M. Schmelz, and E. Fransén. 2014. Differential
axonal conduction patterns of mechano-sensitive and mechano-insensitive nociceptors--a
combined experimental and modelling study. *PloS one*. 9:e103556. doi:
10.1371/journal.pone.0103556.
Power, K.E., K.P. Carlin, and B. Fedirchuk. 2012. Modulation of voltage-gated sodium channels
hyperpolarizes the voltage threshold for activation in spinal motoneurones. *Exp Brain Res*.
217:311-322. doi: 10.1007/s00221-011-2994-3.
Priest, B.T., B.A. Murphy, J.A. Lindia, C. Diaz, C. Abbadie, A.M. Ritter, P. Liberator, L.M. Iyer, S.F. Kash,
M.G. Kohler, G.J. Kaczorowski, D.E. MacIntyre, and W.J. Martin. 2005. Contribution of the
tetrodotoxin-resistant voltage-gated sodium channel Na$_V$1.9 to sensory
transmission and nociceptive behavior. *Proceedings of the National Academy of Sciences*.
102:9382-9387. doi: doi:10.1073/pnas.0501549102.
Renganathan, M., T.R. Cummins, W.N. Hormuzdiar, and S.G. Waxman. 2000. alpha-SNS produces the
slow TTX-resistant sodium current in large cutaneous afferent DRG neurons. *J Neurophysiol*.
84:710-718. doi: 10.1152/jn.2000.84.2.710.
Renganathan, M., T.R. Cummins, and S.G. Waxman. 2001. Contribution of Na(v)1.8 sodium channels
to action potential electrogenesis in DRG neurons. *J Neurophysiol*. 86:629-640. doi:
10.1152/jn.2001.86.2.629.

Renganathan, M., S. Dib-Hajj, and S.G. Waxman. 2002. Nav1.5 underlies the 'third TTX-R sodium
current' in rat small DRG neurons. *Brain Res Mol Brain Res*. 106:70-82. doi: 10.1016/S0169-
328X(02)00411-4.
Rogers, M., N. Zidar, D. Kikelj, and R.W. Kirby. 2016. Characterization of Endogenous Sodium
Channels in the ND7-23 Neuroblastoma Cell Line: Implications for Use as a Heterologous Ion
Channel Expression System Suitable for Automated Patch Clamp Screening. *Assay Drug Dev*
Technol. 14:109-130. doi: 10.1089/adt.2016.704.
Rugiero, F., M. Mistry, D. Sage, J.A. Black, S.G. Waxman, M. Crest, N. Clerc, P. Delmas, and M. Gola.
2003. Selective expression of a persistent tetrodotoxin-resistant Na+ current and NaV1.9
subunit in myenteric sensory neurons. *J Neurosci*. 23:2715-2725. doi: 10.1523/jneurosci.23-
07-02715.2003.
Rühlmann, A.H., J. Körner, R. Hausmann, N. Bebrivenski, C. Neuhof, S. Detro-Dassen, P. Hautvast, C.A.
Benasolo, J. Meents, J.P. Machtens, G. Schmalzing, and A. Lampert. 2020. Uncoupling sodium
channel dimers restores the phenotype of a pain-linked Na(v) 1.7 channel mutation. *Br J*
Pharmacol. 177:4481-4496. doi: 10.1111/bph.15196.
Rush, A.M., T.R. Cummins, and S.G. Waxman. 2007. Multiple sodium channels and their roles in
electrogenesis within dorsal root ganglion neurons. *J Physiol*. 579:1-14. doi:
10.1113/jphysiol.2006.121483.
Rush, A.M., S.D. Dib-Hajj, S. Liu, T.R. Cummins, J.A. Black, and S.G. Waxman. 2006a. A single sodium
channel mutation produces hyper- or hypoexcitability in different types of neurons. *Proc Natl*
Acad Sci U S A. 103:8245-8250. doi: 10.1073/pnas.0602813103.
Rush, A.M., E.K. Wittmack, L. Tyrrell, J.A. Black, S.D. Dib-Hajj, and S.G. Waxman. 2006b. Differential
modulation of sodium channel Na(v)1.6 by two members of the fibroblast growth factor
homologous factor 2 subfamily. *Eur J Neurosci*. 23:2551-2562. doi: 10.1111/j.1460-
9568.2006.04789.x.
Sheets, M.F., and D.A. Hanck. 1999. Gating of skeletal and cardiac muscle sodium channels in
mammalian cells. *J Physiol*. 514 (Pt 2):425-436. doi: 10.1111/j.1469-7793.1999.425ae.x.
Sizova, D.V., J. Huang, E.J. Akin, M. Estacion, C. Gomis-Perez, S.G. Waxman, and S.D. Dib-Hajj. 2020. A
49-residue sequence motif in the C terminus of Nav1.9 regulates trafficking of the channel to
the plasma membrane. *J Biol Chem*. 295:1077-1090. doi: 10.1074/jbc.RA119.011424.
Sousounis, K., B. Erdogan, M. Levin, and J. Whited. 2020. Precise control of ion channel and gap
junction expression is required for patterning of the regenerating axolotl limb. *Int J Dev Biol*.
52. doi: 10.1387/ijdb.200114jw.
Stępień, A., D. Sałacińska, J. Staszewski, M. Durka-Kęsy, and J. Dobrogowski. 2020. Paroxysmal
extreme pain disorder in family with c.3892G > T (p.Val1298Phe) in the SCN9A gene mutation
- case report. *BMC Neurol*. 20:182. doi: 10.1186/s12883-020-01770-9.
Stiles, P.J., and C.G. Gray. 2021. Improved Hodgkin–Huxley type model for neural action potentials.
Eur Biophys J. 50:819-828. doi: 10.1007/s00249-021-01547-z.
Tan, J., and D.M. Soderlund. 2009. Human and rat Nav1.3 voltage-gated sodium channels differ in
inactivation properties and sensitivity to the pyrethroid insecticide tefluthrin.
Neurotoxicology. 30:81-89. doi: 10.1016/j.neuro.2008.10.008.
Tang, Z., Z. Chen, B. Tang, and H. Jiang. 2015. Primary erythromelalgia: a review. *Orphanet J Rare Dis*.
10:127. doi: 10.1186/s13023-015-0347-1.
Tate, S., S. Benn, C. Hick, D. Trezise, V. John, R.J. Mannion, M. Costigan, C. Plumpton, D. Grose, Z.
Gladwell, G. Kendall, K. Dale, C. Bountra, and C.J. Woolf. 1998. Two sodium channels
contribute to the TTX-R sodium current in primary sensory neurons. *Nature Neuroscience*.
1:653-655. doi: 10.1038/3652.
Tavares-Ferreira, D., S. Shiers, P.R. Ray, A. Wangzhou, V. Jeevakumar, I. Sankaranarayanan, A.M.
Cervantes, J.C. Reese, A. Chamessian, B.A. Copits, P.M. Dougherty, R.W.t. Gereau, M.D.
Burton, G. Dussor, and T.J. Price. 2022. Spatial transcriptomics of dorsal root ganglia
identifies molecular signatures of human nociceptors. *Sci Transl Med*. 14:eabj8186. doi:
10.1126/scitranslmed.abj8186.

Thériault, O., and M. Chahine. 2014. Correlation of the electrophysiological profiles and sodium channel transcripts of individual rat dorsal root ganglia neurons. *Front Cell Neurosci*. 8:285. doi: 10.3389/fncel.2014.00285.

Thompson, C.H., R. Ben-Shalom, K.J. Bender, and A.L. George. 2020. Alternative splicing potentiates dysfunction of early-onset epileptic encephalopathy SCN2A variants. *J Gen Physiol*. 152. doi: 10.1085/jgp.201912442.

Thompson, C.H., F. Potet, T.V. Abramova, J.-M. DeKeyser, N.F. Ghabra, C.G. Vanoye, J.J. Millichap, and A.L. George, Jr. 2023. Epilepsy-associated SCN2A (NaV1.2) variants exhibit diverse and complex functional properties. *J Gen Physiol*. 155. doi: 10.1085/jgp.202313375.

Tian, J., A.G. Bavencoffe, M.X. Zhu, and E.T. Walters. 2023. Readiness of nociceptor cell bodies to generate spontaneous activity results from background activity of diverse ion channels and high input resistance. *Pain*. doi: 10.1097/j.pain.0000000000003091.

Tigerholm, J., M.E. Petersson, O. Obreja, E. Eberhardt, B. Namer, C. Weidner, A. Lampert, R.W. Carr, M. Schmelz, and E. Fransén. 2015. C-fiber recovery cycle supernormality depends on ion concentration and ion channel permeability. *Biophys J*. 108:1057-1071. doi: 10.1016/j.bpj.2014.12.034.

Tigerholm, J., M.E. Petersson, O. Obreja, A. Lampert, R. Carr, M. Schmelz, and E. Fransén. 2014. Modeling activity-dependent changes of axonal spike conduction in primary afferent C-nociceptors. *J Neurophysiol*. 111:1721-1735. doi: 10.1152/jn.00777.2012.

Tigerholm, J., A.H. Poulsen, O.K. Andersen, and C.D. Mørch. 2019. From Perception Threshold to Ion Channels-A Computational Study. *Biophys J*. 117:281-295. doi: 10.1016/j.bpj.2019.04.041.

Tseng, A.S., W.S. Beane, J.M. Lemire, A. Masi, and M. Levin. 2010. Induction of vertebrate regeneration by a transient sodium current. *J Neurosci*. 30:13192-13200. doi: 10.1523/jneurosci.3315-10.2010.

Usoskin, D., A. Furlan, S. Islam, H. Abdo, P. Lönnerberg, D. Lou, J. Hjerling-Leffler, J. Haeggström, O. Kharchenko, P.V. Kharchenko, S. Linnarsson, and P. Ernfors. 2015. Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat Neurosci*. 18:145-153. doi: 10.1038/nn.3881.

Vacher, H., D.P. Mohapatra, and J.S. Trimmer. 2008. Localization and targeting of voltage-dependent ion channels in mammalian central neurons. *Physiol Rev*. 88:1407-1447. doi: 10.1152/physrev.00002.2008.

Vanoye, C.G., T.V. Abramova, J.M. DeKeyser, N.F. Ghabra, M.J. Oudin, C.B. Burge, I. Helbig, C.H. Thompson, and A.L. George, Jr. 2024. Molecular and cellular context influences SCN8A variant function. *JCI Insight*. 9. doi: 10.1172/jci.insight.177530.

Vanoye, C.G., J.D. Kunic, G.R. Ehring, and A.L. George, Jr. 2013. Mechanism of sodium channel NaV1.9 potentiation by G-protein signaling. *J Gen Physiol*. 141:193-202. doi: 10.1085/jgp.201210919.

Vasylyev, D.V., and S.G. Waxman. 2012. Membrane properties and electrogenesis in the distal axons of small dorsal root ganglion neurons in vitro. *J Neurophysiol*. 108:729-740. doi: 10.1152/jn.00091.2012.

Wang, D.W., K. Yazawa, A.L. George, and P.B. Bennett. 1996. Characterization of human cardiac Na+ channel mutations in the congenital long QT syndrome. *Proc Natl Acad Sci U S A*. 93:13200-13205. doi: 10.1073/pnas.93.23.13200.

Wang, J., S.W. Ou, and Y.J. Wang. 2017. Distribution and function of voltage-gated sodium channels in the nervous system. *Channels (Austin)*. 11:534-554. doi: 10.1080/19336950.2017.1380758.

Wang, T., Y. Wang, J. Shen, L. Wang, and L. Cao. 2022. Predicting Spike Features of Hodgkin-Huxley-Type Neurons With Simple Artificial Neural Network. *Front Comput Neurosci*. 15. doi: 10.3389/fncom.2021.800875.

Whitaker, W.R., J.J. Clare, A.J. Powell, Y.H. Chen, R.L. Faull, and P.C. Emson. 2000. Distribution of voltage-gated sodium channel alpha-subunit and beta-subunit mRNAs in human hippocampal formation, cortex, and cerebellum. *J Comp Neurol*. 422:123-139. doi: 10.1002/(sici)1096-9861(20000619)422:1<123::aid-cne8>3.0.co;2-x.

Wittmack, E.K., A.M. Rush, M.J. Craner, M. Goldfarb, S.G. Waxman, and S.D. Dib-Hajj. 2004.
Fibroblast growth factor homologous factor 2B: association with Nav1.6 and selective
colocalization at nodes of Ranvier of dorsal root axons. *J Neurosci*. 24:6765-6775. doi:
10.1523/jneurosci.1628-04.2004.
Yang, Y., Y. Wang, S. Li, Z. Xu, H. Li, L. Ma, J. Fan, D. Bu, B. Liu, Z. Fan, G. Wu, J. Jin, B. Ding, X. Zhu, and
Y. Shen. 2004. Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients
with primary erythermalgia. *J Med Genet*. 41:171-174. doi: 10.1136/jmg.2003.012153.
Yin, K., G.J. Baillie, and I. Vetter. 2016. Neuronal cell lines as model dorsal root ganglion neurons: A
transcriptomic comparison. *Mol Pain*. 12. doi: 10.1177/1744806916646111.
Yu, F.H., and W.A. Catterall. 2003. Overview of the voltage-gated sodium channel family. *Genome*
Biol. 4:207. doi: 10.1186/gb-2003-4-3-207.
Zhang, H., and A.S. Verkman. 2010. Aquaporin-1 tunes pain perception by interaction with Na(v)1.8
Na+ channels in dorsal root ganglion neurons. *J Biol Chem*. 285:5896-5906. doi:
10.1074/jbc.M109.090233.
Zhang, X., B.T. Priest, I. Belfer, and M.S. Gold. 2017. Voltage-gated Na(+) currents in human dorsal
root ganglion neurons. *Elife*. 6. doi: 10.7554/eLife.23235.

Na$_v$1.1

Na$_v$1.2

Na$_v$1.3

Na$_v$1.5

Na$_v$1.6

Na$_v$1.7

Na$_v$1.8

I/I$_{max}$

G/G$_{max}$

I/I_{max}

G/G_{max}

0.2 mV/ms
0.4 mV/ms
0.6 mV/ms
1.2 mV/ms
2 mV/ms
4 mV/ms
6 mV/ms

U location of Imax(ramp)

voltage command
ramp current

I max(ramp)

Na_v1.1   Na_v1.2   Na_v1.3   Na_v1.5   Na_v1.6   Na_v1.7   Na_v1.8

Na_v1.1
Na_v1.2
Na_v1.3
Na_v1.5
Na_v1.6
Na_v1.7
Na_v1.8

ramp current

AUC

$I_{max(ramp)}$

Na$_v$1.1 Na$_v$1.2 Na$_v$1.3 Na$_v$1.5 Na$_v$1.6 Na$_v$1.7 Na$_v$1.8

Na$_v$1.1 Na$_v$1.2 Na$_v$1.3 Na$_v$1.5

Na$_v$1.6 Na$_v$1.7 Na$_v$1.8

CMi: current injection 30 µA/cm²

Aδ fiber: current injection 30 µA/cm²

Aδ fiber

Modified expression level

A — Nav1.7 conductivity 5 times increased

B — Nav1.8 conductivity 8 times increased

Shifted activation of Nav1.7

C — Wildtype Nav1.7

D — Shift by 5 mV

E — Shift by 10 mV