Recurrent circuits encode visual center-surround computations in the mouse superior colliculus

Peng Cui[1*], Kuisong Song[2*], Dimitris Mariatos-Metaxas[1], Arturo G. Isla[3,4], Teresa Femenia[2], Iakovos Lazaridis[5], Konstantinos Meletis[1], Arvind Kumar[1,6] and Andreas A. Kardamakis[1,2,#]

[1]Department of Neuroscience, Karolinska Institute, Stockholm, Sweden

[2]Instituto de Neurociencias CSIC-UMH, San Juan de Alicante, Spain

[3]Tecnológico de Monterrey, Escuela de Medicina y Ciencias de la Salud, México

[4]Department of Neurobiology, Care Sciences and Society, Karolinska Institute, Sweden

[5]McGovern Institute, Massachusetts Institute of Technology, Cambridge, USA

[6]Division of Computational Science and Technology. School of Electrical Engineering and Computer Science. KTH Royal Institute of Technology, Stockholm, Sweden

[*] These authors contributed equally to the work

[#]Corresponding author

Send correspondence to lead contact:

Andreas Kardamakis
Instituto de Neurociencias CSIC-UMH
Avenida Santiago Ramón y Cajál s/n
San Juan de Alicante, 03015
Spain

Email:
akardamakis@umh.es
andreas.kardamakis@ki.se

Abstract

The interaction of retinal center-surround processes with early visual subcortical and cortical circuits forms the basis for saliency computations. The subcortical circuit mechanisms contributing to surround modulation, however, remain unclear. We developed an optogenetic approach to functionally delineate visual center and surround zones of individual neurons in the superficial layers of the superior colliculus (SCs). We demonstrate that local circuit interactions within the SCs are sufficient to generate surround suppression and that surround network activation leads to decreases in center excitation rather than increases in synaptic inhibition. Using a range of complementary methods, we reveal the identity of a SCs-based circuit with two key interacting circuit motifs, recurrent excitation and feedback inhibition, capable of modulating levels of center excitation when the visual surround is active. We propose that subcortical and cortical visual circuits are both

40 recurrent-based and have evolved to independently perform surround modulation, thus enabling
41 parallel saliency computations across multiple levels.
42
Introduction
Enabling neurons to compare central inputs with their immediate surround represents a fundamental
and conserved computation [1,2]. Termed 'surround modulation' and usually suppressive, it is
widespread across sensory modalities such as vision [3], hearing [4], somatosensation [5], and olfaction
[6]. A typical example found across species is during size tuning of a visual neuron; the firing response
initially increases when a stimulus is centered on its receptive field [7,8], but then decreases as the
stimulus expands beyond this area [9–12]. Such observations have been measured throughout the
visual thalamocortical system - in the lateral geniculate nucleus [13], primary visual cortex 'V1' [14] and
extrastriate cortex [15] - and retinotectal systems in the premammalian optic tectum [9,16,17] or superior
colliculus 'SC' [8,10]. While center-surround interactions are known to originate in the retina, which has
been extensively studied [18–21], little is known about the extent to which they are inherited by later
stages of the visual hierarchy and how extraretinal circuits in the SC and V1 can actively contribute
to surround modulation.
56
This process relies on the interplay between excitatory cells releasing glutamate and inhibitory cells
releasing GABA (γ-aminobutyric acid). As intuition would suggest, surround suppression can be
mediated by increases of levels of GABAergic inhibition, however, different neural systems may
assume different mechanisms [22]. For example, surround suppression in the primary visual cortex
(V1) is driven by decreases in levels of excitation rather than increases in synaptic inhibition [23].
Determining this mechanism, along with its underlying circuit motifs, is essential for uncovering the
minimal circuitry underlying surround network interactions. This insight brings us one step closer to
interpreting how cortical signals can interact with SC circuitry for saliency computations [24–26] or
visuospatial behavior like target selection in overt [27,28] and covert attention [29,30].
66
Here, we demonstrate that early visual circuits within the superficial layers of the superior colliculus
(SCs) contain the sufficient circuitry to generate center-surround interactions in absence from any
extrinsic input. To reveal the circuit mechanism, we devised a functional receptive field mapping
approach that uses optogenetics and whole-cell electrophysiology to delineate and selectively
activate center and surround zones of single neurons. Through this approach, we found that robust
surround suppression is mediated by decreases in levels of excitation in absence of any increase in
synaptic inhibition. We subsequently identified the underlying functional circuit motifs between local
excitatory (*E*) and inhibitory (*I*) neurons using cell-type-specific trans-synaptic mapping. Large-scale
computational modelling revealed the minimal circuit connections among those identified motifs,
revealing the importance of local recurrent excitatory interactions critical for generating visual
surround suppression in the SCs. Finally, we observed functional decreases in the E/I balance in the

surround when the center was active by using *in vivo* cfos induction with optogenetics, thus
confirming a key model prediction. In sum, our results suggest that phylogenetically older early visual
circuits, based in the SCs, are capable of generating surround suppression independently of cortical
saliency computations, while still sharing common circuit strategies in their diverse implementation.
82

Results

Visual surround network activation modulates center excitability in the SCs

To establish the mechanism and circuit basis of midbrain visual surround modulation, we sought to
recapitulate 'classical' center and surround interactions using an *ex vivo* experimental setting by
substituting visual stimulus patterns with optogenetically-induced retinotopic impulse patterns in
midbrain sections of the SCs (Fig. 1a-c). By leveraging the aligned topographic correspondence of
the retinotectal pathway [31], we generated an input-dependent visual field. We made it light-excitable
by injecting an adeno-associated virus (AAV) intravitreally that expressed ChR2 in retinal ganglion
cells (RGCs) of both wild-type C57BL/6NCrl ($n = 22$, $N = 14$) and vGAT-Cre ($n = 7$, $N = 5$) mice (*see*
top panel, Fig. 1a). To ensure sufficient expression of ChR2 across the SCs, pan-retinal transfection
was an essential prerequisite for trial inclusion (Bottom left, Fig. 1a); partial expressions were not
taken into consideration to minimize the risk of false positive surround classification that could arise
from lack of ChR2 expression. We mapped the receptive fields of single SCs neurons ($n = 29$, $N = $
19), referred herein to as center zones (C), by detecting subregions that triggered excitatory
postsynaptic currents (EPSCs) in response to site-specific light patterns (Fig. 1b-c), which would
activate the retinotopically arranged ChR2-expressing RGC axonal terminals in coronal SC sections
(Bottom right, Fig. 1a).

Given the *a priori* unknown perimeter of each neuron's center zone, the surround zone (S) can exhibit
significant spatial variability and extend over a wide area (Fig. 1b-c). To accurately cover each
neuron's center zone and ensure a significant surround area, it was essential to create a sufficiently
large field of view (FOV) that could accommodate their distinct morphologies. To do this, we used
one-photon widefield optics via a low magnification objective (16×) while using an in-house built
dual-focus visual inspection system to selectively magnify or demagnify the optical path (Fig. 1b).
Demagnification (by 0.5×) captured a large field of view of > 1mm^2 that incorporated most of the SCs
(magenta box in lower right of Fig. 1a), thus allowing for patterned micro-optostimulation of the entire
visual layer. The other path diverted the light for magnification (by 2×), thereby achieving sufficient
single cell resolution to visually target our neurons of interest for whole-cell recordings (Fig. 1b). This
functional receptive field mapping approach allowed us to delineate spatial patterns defining center
and surround regions of single neurons, which were subsequently used to selectively activate
separately or in sequences during recordings.

Figure 1c illustrates the center zone of a vGAT+ horizontal cell, which was determined by measuring
evoked EPSCs after optostimulation (5 impulses at 10 Hz) of a 10 × 10 grid covering the FOV. The
areas that evoked EPSC amplitudes exceeding 10-20% of the peak value (*see* Supplementary Fig.
S1) collectively formed an image pattern that were classified as the center, which usually
encompassed the soma and the majority of the dendritic processes (Bottom left, Fig. 1c; *see* also
Supplementary Fig. S1 and S4). Selective activation of this center zone always triggered short-
latency compound EPSCs (mean amplitude: 120.8 ± 15.5 pA) characterized by prominent
multimodal peaks (*see* arrowheads in the magenta trace in Fig. 1d), suggesting the presence of local
recurrent excitatory-excitatory connectivity in addition to monosynaptic excitatory RGC input (Fig. 2-
3).

Fig. 1: Visual surround network activation modulates center excitability in SCs cell-types.
(a) Top: Procedure showing expression of ChR2 in RGCs using an intravitreal viral injection of the vector
rAAV2-hsyn-hChR2(H134R)-EYFP (in WT mice, *N=14*) and unilateral labeling of contralateral SCs
inhibitory neurons with the vector rAAV5-FLEX-tdTomato (in this case vGAT-Cre mice were used, *N=5*).
Bottom left: Inverted fluorescent image of whole retinal mount confirming homogeneous transfection of
RGCs with ChR2-EYFP. Scale bar: 1200 μm. Insets show enlarged aspects of the optic disc (1) and
individual RGCs (2). Scale bar: 25 μm. Bottom right: Coronal midbrain section with ChR2-expressing
RGC axonal terminals seen occupying the SCs layers.
(b) A dual magnification setup for functional receptive field mapping of single cells. This enabled the
simultaneous visualization of a large field of view (FOV) encompassing the SCs layer, and single cell
resolution (SCR) for identifying center excitation zones using whole-cell recordings and patterned
optostimulation via a digital mirror device (DMD). *See* also experimental procedures and Fig. S1-S4.
(c) Top left: a representative example of this approach applied to an inhibitory horizontal SCs neuron
stained intracellularly with Neurobiotin (NB). Scale bar: 100 μm. The neuron responds with excitatory
postsynaptic currents (EPSCs, shown in magenta) to a 20 Hz 5-light-pulse train delivered pseudo-
randomly. Each trace comes from a single optostimulated rectangular area of 120×100 μm (shown on
the right). Bottom left: reconstructed neuron is shown with the denoted excitation zone (dotted line), which
is referred to as center, whereas the non-responsive areas as surround (S) zone. Whole-field
optostimulation is referred to as 'global' (G).

(d) Synaptic modulation between the center and surround zone of the neuron shown in (c). Top: Temporal patterns underlying surround and center interaction. Bottom: Individual traces of synaptic inward and outward currents evoked in response to center stimulation when recording at reversal equilibriums for GABA$_A$-chloride mediated inhibition and glutamatergic excitation (-65 and 0 mV, respectively). Arrowheads show EPSC recurrence in an individual trace (darker shade). Scale bar: 100 μm. Full traces in Fig. S2.

(e) Voltage traces revealing the strong modulation that the surround exerts on center activity in a stellate SCs interneuron. Voltage responses during surround followed by center (green traces) at two membrane holding potentials (-20 mV and -65 mV, top and middle, respectively) or during center alone (magenta). Bottom: Current traces when holding at 0 mV using the same condition.

(f) Plot of the inhibitory conductance as a function of the excitatory conductance in the neuron shown in (c) and (d). Arrow shows direction of process when neuron is activated.

(g-h) Comparison of synaptic excitatory and inhibitory conductances between conditions: center only (C_1) and center after surround (C_s). Color coding corresponds to morphologically or genetically identified SCs neuron cell-types. Orange and red arrows depict data points from neurons shown in (d) and (e), respectively.

(i) Modulation of center excitation plotted as a function of inhibitory conductance that is present at the time of center optostimulation (t_c).

(j) Schematic showing how the same level of response suppression can be achieved by either decrease in excitatory conductance or by increase in the inhibitory conductance. The gray colormap indicates a 2D input-output transfer function of a neuron as a function of normalized input excitation and inhibition. The two curved lines indicate iso-firing rate contours.

Statistical comparisons were performed as paired *t*-tests followed by Wilcoxon rank-signed tests. *Abbreviations: c.*, contralateral; *WT*, wild-type; *RGC*, retinal ganglion cell; *VIS,* primary visual cortex; *RSP*, retrosplenial cortex; *SCs*, superior colliculus superficial layer; *SCi*, superior colliculus intermediate layer; *SCd*, superior colliculus deep layer; *PAG*, periaqueductal gray matter.

Optostimulation of the center zone (control traces in magenta) always resulted in an excitatory phase quickly followed by an inhibitory one (Fig. 1d-f). To induce center-surround interactions, we first activated the surround using three different durations and onsets (*see* dotted box in Fig. 1d). Light stimulation intensity was held constant for both the center and surround zones. The surround-stimulation phases were terminated 20 ms before center-stimulation onset to ensure adequate time for the build-up of surround-induced activity while allowing for signal synchronization with the center-induced activity (cyan traces in Fig. 1d; *see* also Fig. S3d). Obviously, within these experimental settings, simultaneous center and surround activation, akin to whole-field (global) stimulation (Fig. 1b and 2i), would mask any discernible effect the surround would have on the center. Comparing the magenta and cyan traces in Fig. 1d reveals a distinct and consistent decline in center responsiveness after surround activity, seen by a significant reduction in center-evoked EPSC amplitudes (~50% decrease in this recorded cell) along with a concurrent decrease in the number of multimodal peaks. In line with previous studies [9,32], these evoked EPSCs were rapidly followed (mean peak latency ± s.e.m.: 5.9 ± 0.8 ms) by a feedforward inhibitory component (mean amplitude ± s.e.m.: 24.3 ± 5.0 pA) observed at the equilibrium potential for glutamate-mediated excitation (0 mV, bottom of Fig. 1d), and here we observe that it occurred regardless of the presence of a preceding surround.

The degree of surround modulation varied across different SCs neurons. Based on the intensity of the reduction in center EPSC amplitude, the suppression was classified as mild, moderate or strong (Fig. 1d-e; strong: > 60%, *n* = 9/24; moderate: 20-60%, *n* = 10/24; mild: < 20%, *n* = 5/24). Figure 1e

shows the voltage and current traces of a stellate SCs interneuron that was subject to a strong
surround influence (*see* Supplementary Fig. S2 and S4). When its surround was activated first, visual
input arriving to the neuron's center zone (top and middle traces) was completely suppressed by an
outward current pulling its membrane potential toward the reversal for chloride-mediated $GABA_A$ (γ-
aminobutyric acid type A) inhibition (~69 mV; *see* bottom traces in Fig. 1e). Notably, the precise
relative timing between surround (t_s) and center stimulation onsets (i.e., -70, -40, or -25 ms) or
surround stimulation duration (5, 20, or 50 ms) had little impact on the amplitude variance of
surround-triggered inhibitory postsynaptic currents (*see* arrow, Fig. 1d) or on the relative impact of
center modulation. This mainly stems from latency differences in signaling between center and
surround, with surround *E* and *I* conductances peaking at average latencies of 20 and 28 ms,
whereas center conductances peaked at 9 and 15 ms, respectively (Supplementary Fig. 3d).

Surround network activation typically generated a variable combination of excitation and inhibition in
terms of amplitude and timing. However, during the onset of center stimulation (t_c), only the inhibitory
components persisted while any remaining excitation was quenched due to the relatively faster
transmission of synaptic excitation (*see* Supplementary Fig. 2). This can be seen by the non-null
values of surround-triggered outward currents when holding at 0 mV at t_c (*see* arrow, Fig. 1d and 1i).
As expected, removal of surround input via pharmacological blockade of $GABA_A$ receptors using
bath-applied gabazine (GBZ) transformed any input into seizure-like activity while saturating
subsequent center response profiles (bottom traces in Fig. S2a).

To test how surround signaling can account for the observed modulation of center excitability, we
calculated the theoretical contributions of excitatory (G_e) and inhibitory (G_i) synaptic conductances
[33,34] during center only (magenta) and center-surround interaction (cyan) in Figure 1f. Here, the
relationship between inhibition and excitation of the recorded neuron plotted in Figure 1d is shown
during these two conditions (*see* also Fig. S3). Notably, consecutive center stimuli (C_1-C_2) reveal
that inhibition can increase to nearly double the levels while simultaneously maintaining twice the
levels of excitation compared to the significantly decreased levels evoked by surround-center
stimulation (S-C_s). This result suggests that center-surround interactions reduce the persistence of
both types of inputs. This was true for the majority of SCs interneuron cell types, with over half of
them being classifiable based on either morphology (i.e., horizontal, wide-field, narrow-field, or
stellate; for details, *see* [35]) or genotype (e.g., vGAT+; *see* Fig. S4 for quantification). We found that
peak excitation (G_e: mean amplitudes of C_1: 1.7 ± 0.3 nS and C_s: 0.6 ± 0.1 nS with a mean percentage
decrease of 62.3 ± 4.7%) and peak total conductance (G_t: mean amplitudes of C_1: 2.1 ± 0.3 nS and
C_s: 1.2 ± 0.2 nS with an average percentage decrease of 40.6 ± 5.3%) of the center response was
significantly reduced following surround network activation in all these recordings (Fig. 1g and Fig.
S3), while no significant accompanying increase was observed on average to the center inhibition
at peak G_t (Fig. 1h; mean amplitudes of C_1: 0.53 ± 0.08 nS and C_s: 0.55 ± 0.09 nS).

There are two explanations for this weakening of the center excitatory conductance: (i) inhibitory
conductances persisting at t_c, which might act linearly or non-linearly, and/or (ii) a lesser number of
active inputs arising from local glutamatergic interneurons contributing to the center response.
Although a weak, yet statistically significant, correlation was observed between the inhibitory
conductance at t_c (mean amplitude ± s.e.m.: 0.46 ± 0.08 nS, p = 0.004, n = 24) and the reduction in
the center peak G_e (Fig. 1i), the low slope value indicates that it can only account for a partial
explanation of the observed decrease in center excitation. Hence, both center and surround zones
reciprocally affect each other through synergistic recurrent interactions among local populations of
excitatory and inhibitory neurons, aiming to reduce the overall levels of excitation in the system (Fig.
1j).

SCs space defined by local recurrent excitatory and inhibitory circuit motifs
At the network level, this reduction of center activity can be achieved by: 1) direct activation of long-
range inhibitory neurons in the surround, 2) indirect activation of inhibitory neurons in the center via
excitatory neurons in the surround, or 3) a combination of both mechanisms. To reveal the underlying
circuit motifs, we performed retrograde trans-synaptic labeling from either vGluT2 or vGAT neurons
to identify their monosynaptic and presynaptic sources of excitatory and inhibitory input in the SCs.
Considering their comparable proportions (i.e., 40% vGluT2 and 60% vGAT; see [36]) and their
intermingling within the SCs, we opted for genetically restricted approaches to accurately target and
classify these neurons based on their neurotransmitter.

To trace local afferent input to excitatory and inhibitory neurons restricted within the boundaries of
the SCs layer, an AAV-DIO-TVA-V5-RG was microinjected (50 nL) into the SCs of adult (older than
58 days) vGluT2- and vGAT-Cre mice (Fig. 2a and Fig.S5). This Cre-dependent AAV helper vector
was designed to express the TVA receptor together with the engineered rabies glycoprotein (RG;
see [37,38] exclusively in either excitatory or inhibitory neurons depending on the mouse line used. The
injection of the helper virus was delivered into the SCs layer to ensure that neurons expressing the
RG would be located within the SCs layer, thereby minimizing spillover into the intermediate layers
(SCi) as indicated by the green box in the coronal midbrain section of Figure 2b. Any cases where
helper expression extended beyond the boundaries of the SCs were excluded from the analysis.
Three weeks later, the enhanced green fluorescent protein (EGFP)-expressing EnvA-coated and G-
deleted rabies (Rb; see [39] viral vector was injected, targeting the same location in the SCs as the
helper virus injection (Fig. 2a-b).

Neurons that were transfected with the Cre-dependent helper virus, as detected by V5
immunohistochemistry, which would also express Rb-EGFP were referred to as 'starter' neurons
(Top, Fig. 2c). Therefore, in experiments involving vGluT2-Cre animals (*N* = 3), vGluT2+ excitatory

cells served as starter neurons, while in experiments with vGAT-Cre animals ($N = 3$), vGAT+
inhibitory cells were the starter neurons. On the other hand, cells labeled solely with Rb-EGFP,
without the presence of the V5 tag, were classified as input neurons, providing monosynaptic input
to the starter neurons (Fig. 2c). Rather than performing a whole-brain input analysis in the SCs (*see*
[40], we focused on shedding light on the local connectivity between vGluT2+ and vGAT+ neurons in
the SCs. We quantified the total number of presynaptic input neurons (Fig. 2c; *see* also Fig. 2g) and
examined the spatial distribution (Fig. 2e-f) onto each other within the SCs, as well as long-range
inputs arising from both retinas.
278

Fig. 2: Connectivity between local excitatory and inhibitory motifs in SCs space
(a) The helper virus AAV-DIO-TVA-V5-RG expresses the rabies glycoprotein (RG), and a fusion of the
TVA receptor with the V5 tag. The RG-deleted rabies virus Rb, pseudotyped with EnvA, expresses
EGFP. The AAV helper virus was first injected into vGluT2-Cre (*N=3*) and vGAT-Cre (*N=3*) mice and 3
weeks later the Rb-EGFP was delivered in the same location in the SCs.
(b) Left: coronal section of the midbrain of a vGAT-Cre animal with emphasis on initial injection site where
expression of the V5 tag (in yellow) should be restricted to and not beyond the SCs layer. Subsequently,
EGFP-expressing neurons are a result of Rb-transfection (magenta). Scale bar: 300 µm. Insets below
show colocalization of the enlarged area (green box). Scale bar: 100 µm. Right: Contralateral and
ipsilateral retinas showing retinotopy of presynaptic Rb-EGFP RGCs (black). Counterstaining with s-opsin
(light blue) reveals retinal orientation. Dorsal, D; anterior, A; ventral, V; posterior, P. Scale bars: 1000 µm.
(c) Top: starter neurons are identified based on the co-expression of the V5 tag and Rb-EGFP. Input
neurons only express Rb-EGFP and provide monosynaptic and presynaptic input to the starter neurons.
Bottom: Input convergence shows the ratio of input-to-starter neurons arising from: i) within the SCs, ii)
contralateral (*c.*), and iii) ipsilateral (*i.*) retina.
(d) RNAscope ISH for vGAT (for vGluT2, *see* Fig. S5) revealed the identity of Rb-EGFP input neurons
while confirming starter neuron cell-types. White arrows indicate starter neurons, whereas red and blue
show excitatory (*E*) and inhibitory (*I*) input neurons, respectively. Scale bars: 50 µm.
(e) Schematic depicting the dorsoventral and mediolateral location of starter and input neurons in five
separate anterior-posterior (AP) coronal sections of the SCs layer in a vGAT-Cre animal.
(f) Contour plot showing the spatial distribution of *E* and *I* input neurons to *I* starter neurons in the AP and
medio-lateral ML dimensions across the SCs (for case shown in **e**).
(g) (Left to right) Schematic of all possible motifs between input and starter neurons. The next three
panels show the quantitative relationship between the density of input neurons and starter neurons for *E*
and *I* cell types across the three anatomical dimensions (AP, ML and DV). Solid lines depict data obtained

from excitatory starters averaged from vGluT2-Cre animals (*N*=3), whereas dashed lines show data obtained from inhibitory starters averaged from vGAT-Cre animals (*N*=3). *See* Fig. S5e for the distributions obtained from individual animals.

(h) Quantitative cell-type-specific input-to-starter ratios (referred to as *input convergence*) reveal feedforward (i.e., *E-I* and *I-E*) and recurrent connectivity patterns (i.e., *E-E* and *I-I*). Comparison of means performed using a *t*-test.

(i) Excitatory and inhibitory current recordings (at -65 and 0 mV) from SCs neurons (*n*=5) were performed in response to center (C) and global (G) optostimulation conditions of ChR2-expressing RGC axonal terminals. Bath application of 1 µM TTX and 100 µM 4-AP isolate monosynaptic RGC-triggered excitation while simultaneously abolishing recurrent excitation and recruited feedforward inhibition. Mean evoked EPSC amplitudes ± s.e.m.: from left to right, for Ctrl: 0.40 ± 0.08 nA; for Tx: 0.02 ± 0.01 nA; for Ctrl: 0.29 ± 0.08 nA; for Tx: 0.010 ± 0.004 nA. RM one-way ANOVA, *p* = 0.0072; Tukey's multiple comparisons test, *p*(global-Ctrl *v.s.* global-Tx) = 0.0382, *p*(global-Ctrl *v.s.* center-Tx) = 0.0308, *p*(center-Ctrl *v.s.* center-Tx) = 0.0821. Washout not shown for clarity.

Abbreviations: HPF, hippocampal formation; *Ctrl*, control condition; and *Tx,* treated condition.

We found that vGluT2+ and vGAT+ starter neurons received a similar number of inputs from ipsilateral SCs interneurons (mean ± s.e.m.: 9.5 ± 2 and 10.2 ± 2.8 cells, respectively) but also long-range inputs from both contralateral (2.9 ± 0.3 and 2.8 ± 0.7 cells, respectively) and ipsilateral RGCs (0.07 ± 0.01 and 0.03 ± 0.02 cells, respectively). As expected, the retinotopic pattern is observed (right of Fig. 2b), with contralateral RGC input mainly originating in central regions and ipsilateral RGC input restricted to peripheral regions of the retina, reflecting our systematic viral targeting of the middle region along the mediolateral (ML) and anteroposterior (AP) axes of the SCs (Fig. 2a and 4a). To determine whether the Rb-EGFP+ input cell types were excitatory or inhibitory, we performed RNAscope *in situ* hybridization (ISH) targeting the vesicular GABA transporter (vGAT) or the vesicular glutamate transporter 2 (vGluT2; Fig. 2b and 2d). Neurons that were negative for vGAT, were positive for vGluT2 and vice versa. Figure S5 shows that co-staining Rb-EGFP+ cells with vGluT2 and vGAT probes did not display colocalization, while nearly all detected input cells (269 out of 272 in *N* = 2 animals) were labeled by only one of these two probes.

The relative spatial distributions of vGluT2+ and vGAT+ starter and input neurons were determined by counting the neurons expressing V5 and/or RV-EGFP and mapping their location onto the Allen Reference Atlas coordinate system across the SC layer (Fig. 2e-f and Supplementary Fig. S5e; *see* Methods for details). In Figure 2e, a representative example of a vGAT-Cre animal is shown, illustrating a small number of concentrated inhibitory starter neurons (V5+ and Rb-EGFP+; depicted as yellow dots) together with a broader distribution of input neurons (Rb-EGFP+), which were further classified based on their neurotransmitter type as excitatory (red dots) or inhibitory (blue dots). For all animals, we determined the mediolateral (ML), anteroposterior (AP), and dorsoventral (DV) distribution using this approach for each starter and input neuron, as well as their respective cell type classification (Supplementary Fig. 5e).

The Gaussian kernel-based density plots in Figure 2f reveal the horizontal distribution patterns along the ML and AP dimensions (*see* Methods) of the representative example shown in Figure 2e. Inhibitory starter neurons (left panel in Fig. 2f) were observed to occupy a confined retinotopic area

within the SCs, influenced monosynaptically by a broader excitatory area (right panel in Fig. 2f), as
well as an even broader inhibitory area (middle panel), thereby, receiving visual input from RGCs
located across larger retinal regions. At peak density, the inhibitory starters were found to receive
synaptic input from more than double as many inhibitory input neurons establishing a motif of
recurrent inhibition (Fig. 2f-g).

Using this approach applied in all animals ($N = 6$), we found that every possible motifs between input
and starter neurons existed (Fig. 2g-h and Supplementary Fig. 5e). In Figure 2g, color-coded solid
lines represent average spatial distributions using excitatory neurons as starters, while color-coded
dashed lines correspond to those from inhibitory starters (*see* Fig. S5e for individual traces). As
expected, input neurons always display higher normalized density distribution compared to their
starter counterparts. Notably, while *I-E* and *I-I* input-to-starter connections span a broader range on
the ML, AP and DV axes, a stable E/I ratio is maintained throughout the SCs, with no identifiable
asymmetries arising from a specific connection. By contrast to excitatory neurons, inhibitory inputs
exhibited stronger connections with both inhibitory starter neurons (78.8 ± 4.6%, n = 3) and excitatory
starter neurons (69.2 ± 2.3%, n = 3). The remaining proportion was accounted for by excitatory input
neurons. Notably, about one third of excitatory input neurons establish monosynaptic contact with
excitatory starter neurons (second column in Fig. 2g), thereby, establishing a local recurrent
excitatory network that can be also triggered by monosynaptic long-range RGC input (Fig. 2b).

Figure 2i demonstrates the efficacy of the recurrent excitatory connection (*E-E)* by recording SC
neurons (*n = 5*) in response to optostimulation of RGC terminals within their identified center zones
in the presence of 1 µM TTX and 100 µM 4-AP (washout *not shown* for clarity). Evoked EPSC
amplitudes were drastically reduced by about ~90%, with a small component persisting resulting
from monosynaptic RGC input (Fig. 2i, mean amplitudes ± s.e.m.: for Ctrl: 405.5 ± 81.1 pA; for Tx:
22.3 ± 11.0 pA). This observation suggests the existence of additional center-dependent excitatory
input (Fig. 2d, 2g-h and S5e), which may serve to amplify incoming visual signals. Furthermore, the
lack of observed outward currents (traces held at 0 mV) during this treated condition (Tx) indicates
the removal of recruited di- or oligo-synaptic inhibition, thereby exposing the indirect functional
impact of local *I-E* and *I-I* inputs (Fig. 2i). As expected, the similar response profiles observed
between center-specific and whole-field (global) optostimulation indicate that most of the *E-E* input
is generated within the neuron's center zone.

Local recurrent excitation and feedback inhibition drives visual surround suppression
To isolate the contribution of different connections (*E-E, E-I, I-I,* and *I-E*) to the center-surround
interactions in Figure 1d-j, we constructed a large-scale model consisting of 12,800 spiking neurons
to simulate the SCs layer. The model was constrained by the measured input convergence ratios
and spatial distributions (Fig. 2g-h, Supplementary Fig. 5e) summarized in the cartoon in Figure 3a.

To quantify these network interactions, we used a spiking neuron model with conductance-based
synapses, which were connected to each other based on physical distance according to a Gaussian
connectivity kernel (*see* Table 3 and Supplementary Fig. S4e). Because we were interested in the
contribution of different connections, we kept the connection probability and distance-dependent
connectivity kernel fixed and systematically varied the connection strengths (*see* Table 3-7). For all
synaptic strengths the network remained in an inhibition-dominated activity regime in which inhibitory
neurons displayed a higher background activity than excitatory neurons [41].

To mimic center zone activation, we triggered activity in 314 excitatory and 314 inhibitory model
neurons in four conditions (Fig. 3b-d) and measured the evoked excitatory synaptic conductance in
a randomly chosen neuron from the center region (Fig. 3e). First, the center neurons alone were
stimulated (*see* no.1 in Fig. 3b, top row in Fig. 3c and first column in Fig. 3d). As with our experimental
recordings (*see* Fig. 1d), model center response patterns typically demonstrated multiple peaks,
often characterized by a dual peak profile (green trace in Fig. 3e). The first peak is provided
monosynaptically by glutamatergic RGC input and followed by a second peak that indicates that
center stimulation engaged activity in neighboring unstimulated excitatory neurons. As expected, the
second peak depended mainly on the strength of recurrent excitatory connection (*E-E*) and was also
found to covary with the strength of the *I-E* connection (Fig. 4f).

When center stimulation was immediately followed by a surround input (*see* #4 in Fig. 3b, bottom
row in Fig. 3c and fourth column in Fig. 3d), the peak of the secondary response was determined by
the mutual coupling between excitatory and inhibitory neurons (blue traces in Fig. 3e). In line with
our results presented in Fig. 1, the initial surround-induced inhibitory conductance persisted after the
removal of surround inputs, effectively attenuating the secondary response to below threshold
(yellow line in Fig. 3e). This occurred because fewer *E* and *I* neurons could respond beyond
threshold to the center input (bottom row in Fig. 3c). This was further supported by the observation
that secondary response attenuation was more pronounced when center input coincided with the
presence of surround input (red traces in Fig. 3e). Even though center and global responses are
comparable in amplitude (Fig. 2i), confirmation of this prediction was not possible experimentally as
center responses would prevail during the simultaneous optogenetic activation of center and
surround (*see* #3 in Fig. 3b and third column in Fig. 3d).

Fig. 3: Model predicts recurrent networks a key feature in visual surround modulation

(a) Schematic of the network model using E/I connectivity patterns shown in Fig. 2g. 6400 excitatory and 6400 inhibitory neurons were placed on a square lattice which was folded as a torus to avoid boundary effects (*see* **c**).

(b) Raster of spiking activity with four stimulus conditions that are marked with vertical numbered lines that simulate experimental conditions shown in Fig. 1. Conditions *1:* center only before surround, *2*: surround only, *3*: center during surround, and *4*: center after surround)

(c) Expanded network response of conditions 1 (top) and 4 (bottom) shown in (b).

(d) Spatial arrangement of center and surround network response profiles during the 4 different stimulus impulse conditions. The square schematically shows the region in which neurons were placed on a square grid. Excitatory (top) and inhibitory neurons (bottom) within the white circle received inputs corresponding to the center stimulation. All the neurons in white square but outside the inner circle received inputs corresponding to the surround network. Each subplot shows activity measured within a bin of 5 ms.

(e) Time courses of excitatory conductance of a center neuron. Each trace is an average of 20 trials and shading indicated 25%-75% quartiles. The first peak is generated with low-variance monosynaptic activation of direct retinal input whereas the second peak is a network effect generated by balanced recurrent excitation and recurrent inhibition. The connectivity parameters for these six examples are marked in panel **(f)** with Latin numerals.

(f) Normalized center conductance during and after surround stimulation. Each colormap shows normalized change in excitatory conductance (*see* Methods) in a center neuron (shown in panel e) as a function of recurrent excitation (top: weak connection; bottom: strong connection) and mutual connectivity weights between excitation and inhibition (x-axis: inhibition of excitatory neurons; y-axis: excitation of inhibitory neurons).

To quantify the contribution of individual circuit motifs, we normalized the center response during and after surround input (left and right column in Fig. 3f) by the peak when only center input was delivered (Fig. 3f). As expected, any increase in the *I-E* connection increased its attenuation, whereas *E-I* connectivity did not have a strong effect on the attenuation of the secondary peak. Notably, increase in the strength of recurrent excitation also increased the attenuation of the secondary excitatory conductance peak (compare top and bottom rows in Fig. 3f). The effect of surround-triggered inhibition on center excitation was larger when center input arrived in the presence of an active surround (left column in Fig. 3e). Overall these results demonstrate the

reduction in excitatory conductance during surround suppression requires both recurrent *E-E* and *I-*
E connections.

Center activation lowers E/I balance of active SCs neurons in the surround

Results obtained from the circuit mapping (Fig. 2c and 2g-h) corroborated with the simulated network
(Fig. 3e–f) indicate that the balance between excitation and inhibition (E/I) should shift to lower
values in the surround region when the center is stimulated. To test this prediction, we intravitreally
injected wild-type C57BL/6NCrl mice with an AVV2-ChR2-EYFP (*N* = 5) or AAV2-mCherry (*N* = 3,
for control) to express ChR2 (or a control protein) in RGC axonal terminals (top, Fig. 4a). The
successful activation of SCs neurons following *in vivo* optostimulation of RGC terminals via an
implanted fiber optic unilaterally was confirmed through *cFos* labeling, a surrogate marker of neural
activity (Fig. 4a-b). The animals were maintained in a dark chamber to eliminate the potential
influence of natural visually driven activity.

Figure 4b shows the distribution of active SCs cells in coronal sections (top is control, bottom is
ChR2). After control optostimulation, the presence of a sparse population of spontaneously active
neurons located could be seen reflecting baseline activity at approximate equidistant locations along
the mediolateral axis with a higher concentration near the midline (top image in Fig. 4b).
Optostimulation of ChR2-expressing RGC axonal terminals (bottom image in Fig. 4b; *see* Methods)
revealed a significant increase in *cFos* expression under the implant (top, Fig. 4c), with higher overall
levels observed across both hemispheres (bottom, Fig. 4c). Furthermore, during control the spatial
distribution of active neurons across the mediolateral axis maintained a balance across the
hemispheres, which shifted in the case of ChR2-expressing RGC input activation (bottom, Fig. 4b).

Shifts in E/I balance were determined by quantifying the proportion of inhibitory neurons among the
total activated cells. To do this, we defined the center region as the area located directly under the
implant (C), along with two distal surround zones located more than 400 μm from the center: a medial
surround zone (SM) and a lateral surround zone (SL). Figure 4d (top) shows a subset of *cFos* positive
neurons in the center zone that were immunoreactive for the neurotransmitter GABA revealing their
inhibitory origin (*see* also Supplementary Fig. S7). Notably, optostimulation drives a balance of
excitation and inhibition in the center with a population of putative glutamatergic neurons (i.e.,
negative for GABA) that matches the population size of active inhibitory neurons (C: 0.49 ± 0.07, *N*
= 4; mean ± s.e.m.). The fraction of colocalized neurons, however, shows a significant increase in
the ratio of active GABAergic neurons in both surround zones compared to the center (SM: 0.70 ±
0.13, SL: 0.73 ± 0.14, *N* = 4; mean ± s.e.m.). This finding supports the notion that there will be fewer
recurrent excitatory neurons active to amplify visual signals in the surround when the center is active,
and vice versa.

495
496
Fig. 4: Functional identification of active SCs neurons reveals that center activation lowers
E/I balance in the surround

(a) Pipeline that we used to measure *in vivo* cFos expression induced by local optostimulation of RGC
axonal terminals in the SCs. Top left: As in Fig. 1a, expression of ChR2 in RGCs was achieved using an
intravitreal viral injection of the vector rAAV2-hsyn-hChR2(H134R)-EYFP (in WT mice, *N=5*) or rAAV2-
hsyn-mCherry for control (in WT mice, *N=3*). Bottom left: schematic illustrating assay performed in a dark
box. Right panels: cartoon of the optic fiber placement site shown in a coronal (top) and a horizontal
(bottom) midbrain sections.
(b) Top panels: Coronal midbrain sections of the SCs centered around the implant for c-*Fos* induction
following local optostimulation. Yellow, c-*Fos* immunostaining; magenta, control (top) or ChR2 (bottom);
cyan, GABA immunostaining. Scale bar represents 500 µm. *Abbrev*: C, center; SM (surround medial);
SL (surround lateral). Bottom panels: The location (top) and distribution (bottom) of c-*Fos* positive
neurons across the hemispheres of the SCs. Blue circles are control; Yellow circles are ChR2-induced.
The central zone (C) is defined by the implant tip, which defines the origin of the x-axis. Shading is
standard error.
(c) Comparison of the total cell count of cFos+ cells induced in SCs neurons between control (in blue,
N=3) and during activation of ChR2-expressing RGC axonal terminals (in yellow, *N=5*) across the entire
SCs layer and center zone, respectively.
(d) Top: Grayscale image taken from the center region showing *cFos* (left) and GABA immunostaining
(right). Blue arrows show colocalization. Bottom: Ratio of GABA+ to cFos+ cells across the three zones
(C, SM and SL) compared between the two conditions (ChR2-induced, *N=4*; and control, *N=3*). Statistical
testing was done using Student's *t*-test (*, *p < 0.05*; ***, *p < 0.0001*; *ns*, no significance). Bars show
standard error of the mean.
(e) Comparison of the distribution between activated and connected neurons. Plot of activated neurons
(in yellow) was obtained from the bottom right panel in **(b)**, whereas the plot of connected neurons (in
red) was obtained using cell-type-specific transsynaptic retrograde mapping shown in Fig. 2 and Fig. S5.
(f) Qualitative comparison of the effect with (top) and without (bottom) the recurrent *E-E* connection on
the distribution of active neurons in response to center stimulation using the same model parameters
used in Fig. 3. The square schematically shows the region in which neurons were placed on a square
grid (as in Fig. 3d). Inner circle is the site of center stimulation, whereas the surround is located between
the inner white square excluding the area occupied by the white circle. Probability of spiking (estimated

from 20 trials) of a neuron when center alone was stimulated. The color bar ranges from 0 to 1 and shows
the probability of spiking when the center is stimulated.
Abbreviations: C, center; SM, surround medial; SL, surround lateral.
531

Figure 4e compares the spatial distribution between activated neurons (data from Fig. 4b) and
connected neurons (data from Fig. 2g and Supplementary Fig. S5e). Considering the unlikely event
of activated fibers of passages targeting areas other than the center [42], which may contribute to this
discrepancy, two plausible reasons can be identified: 1) activation of retinotopically-aligned recurrent
networks and 2) activation of upstream circuits connected to the SCs arising, for example, from the
V1 and parabigeminal nucleus (Fig. 4e). To explore whether local recurrent excitation can account
for the observed drop in the surround E/I balance mediated, we tested the efficacy of the connectivity
footprint by increasing the intensity of the model's center stimulation while utilizing the same
parameters as depicted in Fig. 3 (refer to Table 5 for details). When recurrent excitatory connections
were active (top row in Fig. 4f), center stimulation triggered neuronal activity in the surround that was
not directly stimulated as indicated by the non-zero firing probability in neurons outside the white
circle depicting the model 'center' (Fig. 4f). Recruited responses in inhibitory neurons spread out
more than in excitatory neurons indicating the spatial discrepancy between the *E-E* and *E-I*
connections, which we found to be independent of center size. This qualitative description obtained
by the model simulation could explain shifts in surround E/I without the exclusive reliance on long-
range horizontal connections.

Discussion

This study offers mechanistic insight into how the SCs computes visual saliency through recurrent-
based center-surround interactions, capable of operating independently yet in parallel with other
extraretinal brain regions like V1. Targeted optogenetic activation of center and surround zones in
individual neurons allowed us to test the presence and circuit implementation of surround
modulation, using the SCs as a conserved model of early visual processing. Our experiments show
that a selective decrease in center excitation occurs when the visual surround is active (Fig. 1); a
potential computation contributing to midbrain target selection where a current central input becomes
the next saccade's surround input. Given our experimental settings, this result proves that evoked
changes in synaptic excitatory conductance of varying magnitude (usually moderate or strong) are
due to local interactions placed within the midbrain SCs itself. This finding suggests that the main
extraretinal circuits (i.e., SCs and V1) belonging to divergent subcortical and cortical pathways can
perform surround suppression independently. This implies that a stimulus can undergo selective
suppression in one pathway (e.g. thalamocortical) while concurrently being preferred in the other
(e.g. retinotectal). Understanding how phylogenetically older and neocortical circuits can cooperate
or compete is important for comprehending the versatility in pre-attentive visual processing.

As in cortical circuits [43], the SCs also contains excitatory and inhibitory neurons that are spatially
intermingled, a feature that complicates the unravelling of their function. By combining intracellular
measurements (Fig. 1) and connectivity analysis (Fig. 2) with a computational model (Fig. 3), we
revealed two main local circuit motifs: recurrent excitation (*E-E*) and feedback inhibition (*I-E*) that
contribute to surround suppression in distinct ways. *E-E* connections amplify responses in a specific
retinotopic 'center' whereas *I-E* performs a dual function. It overlaps with center-specific *E-E*, thereby
modulating the responses of the same center neurons subject to *E-E*, while simultaneously, reaching
slightly beyond this zone, curbing the responsiveness of neighboring surround neurons outside the
center (Fig. 2–3). In natural conditions, however, every input should be accompanied by some
surround input since one neuron's center will be another neuron's surround. We show that in such
a case, these recurrent interactions conspire to reduce the persistence of input in the absence of
top-down input. Further research is required to establish whether multiple classes of inhibitory
interneurons are involved in specific computations in the SCs, drawing parallels to findings in V1
where the potential role of somatostatin-positive [44] in coalition with recurrent excitatory connections
[45] are involved in cortical surround suppression.

The question of whether these networks process all visual streams uniformly, with each stream
conveyed by distinct RGC subtypes (of which more than 30 are known, *see* [49], can be addressed in
future by using our approach presented in this study. For instance, can some inputs bypass this
circuitry, evade surround suppression, and avoid engaging *I-E* subcircuits? Can they selectively
drive surround facilitatory connections for contour integration across the visual layer [50]? This can be
addressed by optostimulating specific RGC subtypes using opsins expressed in specialized Cre-
lines (e.g., Kcng4-Cre for targeting alpha RGCs, *see* [51]), instead of a panretinal approach. This
strategy can be taken beyond sensory systems, interrogating the presence and nature of center-
surround interactions in modular architectures within motor systems that would be technically
challenging to address *in vivo*. The functional receptive field mapping approach designed in this
study can provide key insight into local circuit function and underscores the need for developing
approaches that link such local processes to global network function.

Visual saliency computations in the SCs provide cortical circuits the flexibility to modulate SCs
activity either by amplifying or suppressing with retinotopic correspondence (*see* white arrow in Fig.
2b, *see also* [27]). This could allow them to engage with SCs recurrent networks and modulate local
center-surround dynamics in a top-down manner. For instance, by granting selective access for
pyramidal neurons in V1 to interact with *E-E* connections, responses can be biased to specific
incoming retinal inputs from certain areas of the visual field, or in contrast, they can be suppressed
by activating *I-E* connections. Importantly, these phylogenetically conserved circuits in the SCs can
also influence cortical activity via tecto-thalamic pathways [52], thereby influencing visual attention [30,53].
This highlights the ability of SCs activity to modulate neural activity in both upstream and downstream

targets, potentially influencing higher-order functions like visuospatial perception and behavioral
selection.
607
Taken together, our results show that selective decrease in center excitation when the visual
surround is active is consistent with models that involve amplification in recurrent networks [46], but
inconsistent with models that are dominated by feedback inhibition [47]. In the latter, transient increase
in inhibition would account for reduction in overall excitation and inhibition levels – an outcome that
our measurements did not support. Computational models of trial-by-trial variability suggest that,
while both increases in inhibition or decreases in recurrent excitation can similarly affect the average
output response, the latter is more likely to minimize variability across trials [48]. We propose that SCs
circuitry has evolved to mitigate this variability where surround inputs are continuously transformed
into center inputs during visual exploration.
617

Acknowledgements

We thank Drs. Michael Haüsser, Abdel El Manira and Julien Bouvier for comments on the
manuscript. We thank Dr. Sten Grillner for his support throughout this study. We thank Drs. Mats
Hellström and Sias Jordaan for technical assistance. A.A.K. acknowledges funding from the Swedish
Research Council (Starting Grant in Medicine no. 2016-03134 and Project Grant in Medicine no.
2020-02927), CIDEGENT funding from the Valencian Community in Spain (no. 2021-028), Strategic
research area neuroscience StratNeuro (Start-up program 2017), the Swedish Brain Foundation
(Hjärnfonden no. FO2018-0254), the Hedlunds Foundation (no. M-2017-0691) the Olle Engvist
Foundation (no. 196-0148) and the Stiftelsen Kronprinsessan Margeratas Arbetnämnd för
synskadade (nos. 2019-027 and 2020-182).

Author Contributions

630
P.C. performed electrophysiological recordings, viral injections, optogenetic experiments and data
analysis. K.S. performed neuroanatomical experiments, RNAscope *in situ* hybridization, viral
injections, fiber implantations and data analysis. D.M.M. performed *in vivo* cFos induction
experiments, visualization of neuroanatomy, and data analysis. A.G.I. performed
electrophysiological recordings. I.L. and T.F. contributed to neuroanatomy and *in vivo* experiments.
A.A.K. performed the conductance analysis. I.L., K.M., T.F., and A.A.K contributed to the
conceptualization and experimental study design. K.M. contributed vectors and resources for
visualization of neuroanatomical data. A.K. contributed to the computational study design and
performed the numerical experiments. A.A.K. designed the figures, supervised and funded the study.
A.A.K. wrote the manuscript with help from A.K. All authors reviewed and commented on the
manuscript.
642

Lead contact

644 Further information and requests for sources should be directed to and will be fulfilled by the lead
645 contact, Andreas Kardamakis (akardamakis@umh.es).
646

Material and Methods

Animal strains

Experimental procedures were approved by the Stockholm municipal committee for animal experiments and the Karolinska Institute in Sweden (N9179-2017 and N6747-2019 to A.A.K.), as well as, the Committee on Animal Research at the Universidad Miguel Hernández and carried out in compliance with the Research Council of Spain (CSIC), the Generalitat Valenciana (GVA) and European regulations (2022/VSC/PEA/0236 to A.A.K.). Both male and female vGAT-Cre (Jackson: Slc32a1tm2(cre)Lowl), vGluT2-Cre (Jackson: Slc17a6tm2(cre)Lowl) and wild-type mice (C57BL/6N; Charles River) were used throughout the experiments. Mice were kept on a 12-h day/night cycle under stable temperature (21 ± 1 °C) and air humidity (50–65%) with freely available food and water. The exact numbers of cells (n) and animals (N) used for each experiment are reported in the *Results* section and corresponding figure legend.

Viral injections

The complete list of viruses we used is provided in Supplementary Table 1. Injections were performed in 2–5 months old wild-type mice and transgenic mice under isoflurane anesthesia and injected with buprenorphine post-surgery. Viral titers of adeno-associated viruses (AAV) and modified rabies virus were between 1×10^{12} and 1×10^{13} viral genomes per milliliter (vg/mL). To express ChR2 in the axons of RGCs in the superficial layer of the right hemisphere SC, we injected 1000 nL into the vitreous bodies of the left retinas using the following viral vectors:

1.rAAV2/hsyn-hChR2(H134R)-EYFP-WPRE-PA (UNC Vector Core),

2.rAAV2/hsyn-hChR2(H134R)-mcherry-WPRE-PA (UNC Vector Core),

3.rAAV2-EF1α-DIO-hChR2(H134R)-EYFP (UNC Vector Core), or

4. ssAAV-2/2-shortCAG-dlox-hChR2(H134R)_EYFP(rev)-dlox-WPRE-hGHp(A) (VVF, Zurich).

We used the vector rAAV2/hsyn-mcherry (UNC Vector Core) as in our control group for the intravitreal injections. To identify GABAergic neurons in the SCs, the vector rAAV5-CAG-FLEX-tdTomato (400 nL, UNC Vector Core) was injected into two collicular sites in order to maximize expression: 1) the rostral SCs (from bregma: anteroposterior (AP) -3.6 mm; right (R) -0.8 mm; dorsoventral (DV) -1.6 mm from the surface of the skull), and 2) the caudal SCs (AP -4.2 mm; R -1.8 mm; DV -1.5 mm) respectively on the same day.

For the rabies virus tracing, 50 nL of helper virus AAV5-EF1a-DIO-TVA-V5-t2A-RG was injected into the right SCs (AP -3.9 mm; R -0.8 mm; DV -1.05 to -1.15 mm from the surface of the

brain) in vGAT-Cre or vGluT2-Cre mice, while taking great care to maintain viral spread to within
the SCs layer. Three weeks later, 150 nL of the engineered rabies virus EnvA-coated G-deleted
rabies virus with EGFP (EnvA-ΔG-Rb-EGFP) was injected into the same location.

For secondary injections, fiber optic implantations and electrophysiological recordings, two
to three weeks were allowed to pass after the initial injection. Viral suspensions were delivered at
500 nL/min and 100 nL/min for intravitreal and brain injections, respectively. A puncture was made
2 mm posterior to the corneal limbus using a sterile 30-gauge (G) needle after pupil dilation was
achieved with 0.5% Alcaine. After which 1000 nL vitreous was aspirated manually using a Hamilton
microliter syringe. The glass capillary was then inserted through the same puncture by using a
microliter syringe (1.5 mm volume) into the eye, and the tip was angled toward the vitreous humor
and back of the eye, taking care to avoid any damage to the retina or lens. The capillary remained
in the eye for 2 min in order to prevent reflux of the viral suspensions. Brain injections were
performed by using a digital stereotaxic frame combined with a Quintessential Stereotaxic injector
(Stoelting) via a glass capillary. The capillary firstly reached 0.2 mm deeper than the target location
to give space for the viral suspensions, then was slowly retracted to the target location to complete
the injections. Afterwards, the capillary was retracted by 0.2 mm again and remained in position for
8 min. During the injections, all the mice were kept on a heating pad at 37 °C with a subcutaneous
injection of buprenorphine while maintaining the eyes were moisturized with Viscotears gel.

Patch electrophysiology in acute midbrain slices

The animals were deeply anaesthetized by using a combination of pentobarbital injected
intraperitoneally. To ensure rapid cooling of the midbrain, we perfused the animals transcardially
with an ice-cold cutting solution, containing the following (in mM): NaCl 40, KCl 2.5, NaH_2PO_4 1.25,
$NaHCO_3$ 26, glucose 20, sucrose 37.5, HEPES 20, NMDG 46.5, Na L-ascorbic acid 5, $CaCl_2$ 0.5
and $MgCl_2$ 5 (pH 7.3 with HCl). The brain was rapidly dissected out after decapitation. Coronal
midbrain slices (400 µm) were prepared by using a vibratome (VT1000S, Leica, Germany) in the
cutting solution and incubated in the same solution at 35 °C for 12 min. The sections were
subsequently maintained at room temperature in recording solution that contained the following (in
mM): NaCl 124, KCl 2.5, NaH_2PO_4 1.25, $NaHCO_3$ 26, glucose 20, $CaCl_2$ 2 and $MgCl_2$ 1 (pH 7.3).
Cutting and recording solutions were continuously infused with carbogen (95% O_2 and 5% CO_2)
throughout the procedure.

Whole-cell patch clamp recordings were either performed on random neurons in the SCs of
wild-type mice or tdTomato-expressing neurons of vGAT-Cre mice at room temperature using
pCLAMP 11 (Molecular Devices, USA). Midbrain slices were continuously perfused with
oxygenated recording solution at a rate of 2 mL/min during recording and visualized under an
upright CleverScope motorized microscope (Micro Control Instruments, UK) equipped with a Dodt
gradient contrast system (DGC-2), an oil condenser (U-AAC, Olympus, Japan) and two
Chameleon3 cameras (Teledyne FLIR, USA). Whole-cell recordings were obtained under visual

control using a 16× (0.80 NA) water dipping objective (CFI75 LWD 16X W, Nikon, Japan) combined with 2× c-mount extender (EX2C, Computar, USA) and c-mount adapters (U-TV1X-2 and U-CMAD3, Olympus, Japan) with Axon MultiClamp 700B amplifier and Axon Digidata 1320A digitizer (Molecular Devices, USA). Signals were sampled at 20 kHz. Pipettes used for recording were pulled from borosilicate glass capillaries (article no. 1403542, Hilgenberg, Germany) by a P-97 Micropipette Puller (Sutter Instrument, USA). Patch pipettes (5–8 MΩ) were filled with a potassium gluconate based intracellular solution contained (in mM): K D-gluconate 125, EGTA 1, KCl 10, HEPES 10, ATP-Mg 4, GTP-Na 0.3, phosphocreatine 10, $CaCl_2$ 0.1 and QX314 3 (pH 7.2 with KOH; 290 mOsm). Due to the long recording times (usually one hour), the access resistance was evaluated every 10 min to ensure that initial values (only under ~30 MΩ were considered) did not increase throughout the recording session.

ChR2-expressing axons of retinal ganglion cells were optically stimulated with 470 nm light from a high-power collimated LED source (Mightex, Canada) connected to a digital mirror device (Polygon1000, Mightex, Canada) by a 3 mm-core liquid lightguide (Newport, USA). The stimulation patterns were triggered via Master-8 pulse stimulator (A.M.P.I, Israel) and designed via the software PolyScan2 (Mightex, Canada). TdTomato-expressing vGAT-positive neurons in the SCs were detected with 540 nm light from a high-power collimated LED source (Mightex, Canada) and an RFP filter set (Olympus, Japan). The 540 nm LED was connected to a 495 nm dichroic filter cube (Mightex, Canada) with the 470 nm LED during the transgenic animal experiments. Both LEDs were directed via a four channel BioLED light source control module (Mightex, Canada).

For membrane property measurements, a series of 1000 ms negative and positive injected current steps from -30 pA to +40 pA in 5 pA increments were delivered through the recording electrode under current-clamped conditions to each recorded neuron.

Functional receptive field mapping of individual neurons

We combined whole-cell electrophysiology with a digital mirror device (DMD) based dual camera system to map the receptive fields of individual neurons. Optogenetic stimulation patterns were created and performed during the recordings by the DMD (Polygon1000, Mightex, Canada). The 16× objective combined with the 2× front tube optics (Mightex, Canada) and 0.5× single photo port tube lens camera adapter (Olympus, Japan) enabled a large workable field-of-view (~1.2 × 1 mm working area). With this customized visual inspection system with this dual camera approach, whole-cell patch-clamp recordings were performed using the high magnification light pathway whereas optostimulations patterns could be performed using the low magnification pathway that enables the large field of view. All neurons were voltage-clamped at −65 and 0 mV to measure excitatory postsynaptic currents (EPSCs) and inhibitory postsynaptic currents (IPSCs) respectively. In most neurons, synaptic currents were also recorded at intermediate potentials (at -45 mV), when recordings were sufficiently stable.

For control experiments, whole-field stimulation patterns were delivered by 10 Hz trains of 8
stimuli (5 ms duration), followed by a single test stimulus at 5 s time interval, and each trace included
four repetitive sweeps. For the center-surround experiments, SCs neurons were subjected to the
following patterns: 'GridScan', 'Center-Center', 'Surround-Surround' and 'Surround-Center'. Center
(C) and Surround (S) areas were determined by using the GridScan protocol, which was delivered
10 Hz trains of 5 light pulses of 5 ms duration in 100 non-adjacent grids arranged in a 10 × 10 spatial
configuration across the FOV (0.1 × 0.07 mm for each one). After the end of the GridScan, we
promptly calculated the maximum EPSC and IPSC amplitudes for each grid after it was normalized
and filtered using custom-written MATLAB (MathWorks). Excitatory postsynaptic current amplitudes
arising from each grid location were thresholded on the basis of their evoked amplitude. Grid
locations that yielded ≥ 10% of the peak EPSC amplitude for the recorded neuron, were considered
to be part of the center zone, whereas locations that did not pass this threshold were considered to
be part of the surround zone. To trigger center-surround interactions, the surround was
optostimulated by a single pulse that was then followed by a two or five-pulse train stimulation
delivered to the center. We experimented with a range of different surround stimulation durations (5,
20, 50 ms) and with a range of surround-center delays (0, 10, 20, 50 ms) to explore the parameter
space. Data was obtained using intermediate values for most of the trials, thereby also avoiding
technical issues or stimulation artifacts.
770

Drugs used in electrophysiology

Drugs were bath applied and an overall duration of at least 5 min was allowed before testing their
effects, which was followed by a washout phase of at least 10 minutes. Unless otherwise indicated,
drugs were perfused during patch-clamp recordings under the given conditions using a two-gauge
peristaltic pump (Pretech Instruments, Sweden). The complete list of drugs used is provided in
Supplementary Table 2.
777

Estimation of synaptic conductances

To deconvolve inhibitory and excitatory synaptic conductance, recordings were performed during
'center-center', 'surround-surround', 'surround-center', and 'global' optostimulation. A Cesium-based
QX314 solution was used in the recording pipette to improve intrinsic conductance isolation and
block action potential discharge conditions (Fig. 1 and Supplementary Fig. 2-3). Evoked synaptic
currents used in this analysis were obtained by averaging values from 5 to 10 sweeps for each
recorded cell. Due to the long recording times (usually one hour), the access resistance was
evaluated every 10 min to ensure that initial values (only under ~30 MΩ were considered) did not
increase throughout the recording session. This procedure was repeated for two to three different
potentials (−65, −45, and 0 mV) held with different levels of current injection. Customized Matlab

scripts were written to estimate the total, excitatory and inhibitory synaptic conductances based on
the equations used in Wehr and Zador (2004; *see also* Haider et al., 2015).

Neuronal reconstruction

To identify the dendritic fields from the recorded neurons, we used 0.3% Neurobiotin (Vector
Laboratories, no.VESP-1120-50) in the intracellular solution during whole-cell recordings. We post
fixed the midbrain slices and then used PBS to wash them. To visualize the neurons, we stained
them with streptavidin Cy5 (1:400, Jackson ImmunoResearch, no. 016-170-084) and imaged them
using confocal microscopy with a 20× objective (LSM-800, Zeiss, Germany). The images were
processed with ImageJ software, and the neurons were reconstructed by using the plugin Simple
Neurite tracer. The dendritic areas of each labeled neuron were calculated by tracing a convex
polygon around the outermost tips of the dendritic field. The SC cell types were then classified on
the basis of their geometry [35].

Fiber optic implantation and cFos activation

Two weeks after the intravitreal injection, a 200 µm diameter fiber optic (Thorlabs, FG200UEA) was
implanted into the right SCs (from bregma: AP -3.6 mm; R -1.0 mm; DV -1.0 mm from the brain
surface). The mice recovered for at least one week before the cFos induction experiment. The mice
were habituated to handling in an open field area (60 cm × 60 cm × 60 cm; W×D×H) and the
connected optic fiber for at least 3 days before proceeding with the experiment. To minimize stress,
bedding from the cage was placed in the box. Optogenetic stimulation was not initiated for 90 min
after the mice were placed in the dark box to avoid any prior visual effects on the observed cFos
expression. The RGC axonal terminals in the SCs were stimulated with a train of 1 s square pulses
of 40 Hz frequency (10 ms ON and 15 ms OFF) every 10 sec for a total of 40 min. After the
stimulation period, the mice were left undisturbed for 60-70 min to ensure sufficient cFos protein
expression. A maximum of 90 min after the end of the stimulation period was allowed before the
mice were euthanized with pentobarbital and perfused with ice-cold 4% paraformaldehyde (PFA) in
0.01 M (1×) phosphate-buffered saline (PBS) buffer. The brain and retinas were subsequently
dissected, and post fixed in 4% PFA overnight at 4°C.

Immunofluorescent staining and *in situ* hybridization

Mice were anesthetized by pentobarbital and perfused with 1× PBS and 4% PFA. The brains (and
eyes, to confirm pan-retina transfection) were harvested and fixed in PFA at 4 °C overnight, then
washed with PBS for 3 times. The brain was placed in 15% sucrose solution for 24 h, then
transferred into 30% sucrose solution for 24–72 h. The brains were subsequently embedded in

optimal cutting temperature compound (OCT), frozen with dry ice and kept at -20 °C for short-term
storage. The cryostat sectioning was performed on a Epredia NX70 cryostat. The brains, between
the thalamus and brainstem, were sectioned in 14 µm thickness, about 80% (12 to 15) sections in
the SCs were collected on glass slides, and every 6th section was sampled for the following staining
(the interval is about 0.105 mm). The other parts of the brain were sectioned in 40 µm thickness
and stored in 1× PBS with 0.01% sodium azide.

The fluorescence *in situ* hybridization was performed with RNAscope Multiplex Fluorescent
Reagent Kit v2 (ACD, Cat. No. 323100). The probes for vGluT2 (Mm-Slc17a6-C2, ACD, Cat. No.
319171-C2) and vGAT (Mm-Slc32a1, ACD, Cat. No. 319191) were used to identify the cell types.
The manufacturer's protocol was followed. Co-staining of vGAT and vGluT2 *in situ* hybridizations
was also performed for control experiment, and the results were shown in Supplementary Fig. 5.

Immediately following the *in situ* hybridization, the brain sections were incubated in chicken
anti-V5 primary antibody (Abcam, ab9113) in 1:750 at room temperature overnight, then washed
with 1× TBST (0.3% TritonX-100), and incubated in donkey anti-chicken Cy3 secondary antibody
(Jackson ImmunoResearch, no. 703-165-155) in 1:1000. All the antibodies were diluted with 1×
TBST (0.3% TritonX-100, 3% BSA, 0.01% sodium azide). After PBS washing and DAPI staining,
the sections were mounted with an antifade mounting medium (VWR, BSBT AR1109). For the cFos
staining, rat or rabbit anti-cFos primary antibodies (Synaptic Systems, no. 226-017 or no. 226-003,
1:500) and GABA primary antibody (Sigma-Aldrich, A2052, 1:2000) were used in 1× PBST (0.3%
TritonX-100, 5% donkey serum, 0.01% sodium azide) at room temperature overnight, then the
sections were stained with Cy3 donkey anti-rabbit (Jackson ImmunoResearch, no. 711-165-152) or
AF647 donkey anti-rat (Jackson ImmunoResearch, no. 712-605-153). The same following steps
are previously described.

The retinas were first removed from the eye in the perfused animals and then flattened with
the help of four incisions, and finally washed in 1x PBS for 3 times. For the retina immunostaining,
the retina was first treated with pre-heated (about 70 °C) 1× antigen retrieval citrate buffer (Sigma,
C9999), then followed the same protocol as with the brain section described above. Rabbit anti-S-
opsin primary antibody (Sigma, AB5407) and donkey anti-rabbit Cy3 secondary antibody were both
used in 1:1000 to visualize the short-wave photoreceptors, which reveals retinal orientation.

Network model

We used an integrate and fire model to simulate the dynamics of neurons. The subthreshold
membrane potential (V_m) of each neurons was described by the following differential equation:

$$C_m dV_m/dt = g_L (V_m - E_L) + I^{bkg}_{syn} + I^{net}_{syn} + I^{stim}_{syn},$$

where C_m is the membrane capacitance, g_L is the leak conductance, I^{net}_{syn} is the total synaptic input

from neurons within the network, I^{bkg}_{syn} is the background input from other networks, I^{stim}_{syn} is input

corresponding to center or surround inputs (*see* Supplementary Table 3). When the membrane

potential reached V_{th}, spike was elicited and the V_m was reset to E_L for t_{ref} millisecond. In the network

all neurons were identical. Neuron parameters are provided in Supplementary Table 4.

Synaptic inputs were modelled as conductance transients with following dynamics:

864                         $g_{syn} = J_{syn}t/tau_{syn} \; exp(-(t-(t_{spk}+ d_{syn}))/tau)$ *when $t-t_{spk} >0$ else* $g_{syn} = 0,$

where t_{spk} is the spike time, d_{syn} is the synaptic delay, J_{syn} *{syn: ee, ei, ie, ii, ex, ix)* is the amplitude

of the postsynaptic potential (PSP), and *tau_{syn}* is the synaptic time constant. In our model all

synapses of a particular type (external, excitatory, or inhibitory) have the same parameters (*see*

Supplementary Table 5). In our model we fixed the amplitude (J_{syn}) of the external excitatory (J_{ex}),

inhibitory (J_{ix}) synapses and J_{ii} (inh. → inh.). The value of J_{ee} (exc. → exc.), J_{ei} (exc. → inh.), and

J_{ie} (inh. → exc.) were systematically varied. Synapse parameters are provided in Supplementary

Table 5.

We considered a network with 6400 excitatory and 6400 inhibitory neurons. Neurons were

placed on an 80 × 80 grid. Because we have the same population size for the two neuron types,

both types of neurons were placed on the same size grid. Neurons were connected based on their

spatial distance. To avoid boundary effects, the 80 × 80 grid was folded as a torus. The connection

probability decreased as a function of distance according to a Gaussian function. The standard

deviation of the Gaussian was set to 16 and 20 for excitatory and inhibitory synapses, respectively.

This was based on our experimental data which showed that inhibitory neurons have a slightly

bigger extent of their spatial connectivity than that of excitatory synapses (Fig. 2f). 16 and 20 grid

points constitute 2% and 2.5% of our network space. The area of the Gaussian function (*i.e.* the

total number of connections sent out by a neuron) was decided based on the input convergence we

measured in our experiments (Fig. 2g-h). Network connectivity parameters are provided in

Supplementary Table 6.

Each neuron received Poisson type excitatory and inhibitory spiking inputs to mimic inputs

from outside the network and to obtain a low firing rate (< 1 Hz) background activity. *See*

Supplementary Table 7 for parameters.

To model a short flash like input to the center region, we selected 314 excitatory and 314

inhibitory neurons, located in a circle (radius = 10 grid points) at the center of our network model.

These neurons received a single synchronous spike from 10 presynaptic neurons (from the retina).

Each spike elicited 1.1 mV depolarization. Each neuron received such a synchronous spike event

at the stimulus onset (*Tcenter*). Each individual neuron received the synchronous spike event at a

time *Tcenter* + *r,* where *r* is a random number drawn from a uniform distribution (U[0,1]). Center

stimulus was presented to the neurons in three conditions: center only, center stimulus just at the

end of the surround stimulus (*see* below) and center while the surround stimulus was still on. *See*
Supplementary Table 3 for more details.

To mimic the surround stimulus, we selected 3286 E-neurons and 3286 I-neurons from a
square region in the network with a side of 60 grid points. The center neurons (see above) were
located at the center of this square. For the surround stimulus we excluded the center neurons. The
surround stimulus was injected in the selected neurons in the form of uncorrelated Poisson type
spike trains. See Supplementary Table 3 for more details.

The network model was simulated using the simulator NEST [54].

Statistical analysis and data plotting

For each mouse, 10–16 sections were stained and visualized. Confocal images (20×) were used
for the manual scoring of co-expression of vGAT/vGluT2, V5 and RV-GFP. Whole-brain images
(10×) were acquired with a fluorescent microscope (Leica DM6000B) and a digital camera
(Hamamatsu Orca-FLASH 4.0 C11440). They were manually registered to a bregma AP coordinate
(from bregma -3.2 to -4.9), and subsequently aligned to the Allen Reference Atlas with the
WholeBrain R package [37,55]. Every detected Rb-EGFP positive neuron was registered, and the
coordinates were used for the following analysis. As the main source of monosynaptic input and
local input, only the RV-GFP positive cells in the SCs were included in the calculation. For the
comparison between mice, the number of input neurons were normalized to the total number of
detected Rb-EGFP positive neurons in SCs. All the density estimations were done using a Gaussian
kernel and were then plotted in R.

To minimize the effects of M-L differences of the injection sites between animals, the
centroid of the starter neurons was first computed using k-means clustering analysis, which then
allowed us to plot the relative M-L coordinate for each neuron. As for the depth estimation, the
coordinates of the SC surface were first obtained with the WholeBrain R package, then the depth
of each neuron was calculated after subtracting the difference. In both, M-L and depth density plots,
only 3 central brain sections with the highest starter density were included according to the AP
density plot.

After confocal imaging the cFos fluorescence was quantified by a 3d object counter plug-in
of Fiji. Colocalized cFos+ and GABA+ cells were manually counted in three areas in SCs (C, SM
and SL; *see* Fig. 4). The statistical analysis and visualization were done in Python (code available
upon request).

We used three model stimulation protocols: center only (center), center stimulus just at the
end of the surround stimulus (center after surround) and center while the surround stimulus was
still on (center with surround). Each paradigm was repeated 20 times. For each stimulus
presentation, we recorded excitatory conductance from a single neuron. We estimated the across

trial mean and variance of the conductance input to the recorded neuron. Finally, the results were rendered as the mean and variance of the excitatory conductance at specific time points after the onset of the stimulus in all three stimulus paradigms. In addition, we recorded spiking activity from all the neurons and estimated the firing rate of neurons. The network model was simulated using the simulator NEST. All differential equations were integrated using Runga-Kutta method with a time step of 0.1 ms.

Electrophysiology data analyses were performed using Clampfit 11 (Molecular Devices, USA), GraphPad Prism (GraphPad Software 7.00, USA), ImageJ (1.53c, NIH, USA), RStudio (R 3.6.1) or MATLAB (R2018b, MathWorks, USA). Data were all presented as mean values ± s.e.m. Before performing paired *t*-test or one-way ANOVA analysis, the normality and the homogeneity of variance were first evaluated using Kolmogorov–Smirnov's test and Brown–Forsythe's test. If the *p* value of Kolmogorov–Smirnov's test was less than 0.05, Wilcoxon's signed-rank test would be used instead of paired *t*-test. If *p* value of Brown–Forsythe's test was less than 0.05, Friedman test and Kruskal–Wallis test would be used instead of a repeated-measures one-way ANOVA (RM one-way ANOVA) and ordinary one-way ANOVA, respectively. *p* values were corrected for deviation using the Geisser–Greenhouse correction. Once the *p* value of a one-way ANOVA analysis was less than 0.05, Tukey's multiple comparisons test would be performed. *F* and df (df1, df2) are degrees of freedom for one-way ANOVA and paired *t*-test, respectively. *p* values less than 0.05 were considered significant with asterisks in figures denoting as follows: $*p < 0.05$, $**p < 0.01$, $***P < 0.001$, $****p < 0.0001$, *ns*, no significance.

Supplementary Table 1. Viruses used in this study

Name	Type	Titer	Source
rAAV2/hsyn-hChR2(H134R)-EYFP-WPRE-PA	AAV2	5.6×10^{12} vg/mL	UNC Vector Core
rAAV2/hsyn-hChR2(H134R)-mcherry-WPRE-PA	AAV2	5.1×10^{12} vg/mL	UNC Vector Core
rAAV2-EF1α-DIO-hChR2(H134R)-EYFP	AAV2	4.4×10^{12} vg/mL	UNC Vector Core

ssAAV-2/2-shortCAG-dlox-hChR2(H134R)_EYFP(rev)-dlox-WPRE-hGHp(A)	AAV2	1.4×10^{12} vg/mL	ZNZ Viral Vector Facility	
rAAV2/hsyn-mcherry	AAV2	5.3×10^{12} vg/mL	UNC Vector Core	
rAAV5-CAG-FLEX-tdTomato	AAV5	7.8×10^{12} vg/mL	UNC Vector Core	
AAV5-EF1a-DIO-TVA-V5-t2A-RG	AAV5	1.2×10^{12} vg/mL	DMC laboratory	
EnvA-ΔG-Rb-EGFP	Modified Rabies	1×10^{10} vg/mL	DMC laboratory	

954 **Supplementary Table 2. Drugs used in electrophysiology experiments**

Compound	Drug name	Concentration	Source
Gabazine (SR 95531)	6-Imino-3-(4-methoxyphenyl)-1(6*H*)-pyridazinebutanoic acid hydrobromide	10 µM	Tocris Bioscience, USA
TTX	(4*R*,4a*R*,5*R*,7*S*,9*S*,10*S*,10a*R*,11*S*,12*S*)-Octahydro-12-(hydroxymethyl)-2-imino-5,9:7,10a-dimethano-10a*H*-[1,3]dioxocino[6,5-*d*]pyrimidine-4,7,10,11,12-pentol	1 µM	Tocris Bioscience, USA
4-AP	4-Aminopyridine	100 µM	Sigma-Aldrich
QX314	*N*-(2,6-Dimethylphenylcarbamoylmethyl)triethylammonium bromide	3 mM	Tocris Bioscience, USA

960 **Supplementary Table 3. Stimulus input**

Type of input	Stimulated neurons	Spike input	Stimulus PSP	Repeats
Center Input	All neurons (exc. and inh.) in a circle with radius 10 grid points. 314 exc. neurons and 314 inh. neurons.	Each neuron received 10 synchronous spikes corresponding to the presentation of a flash like center stimulus. Arrival time of each synchronous spike event was taken from a uniform distribution U[0,1] ms.	1.03 mV at -54 mV to exc. neurons 0.86 mV at -54 mV to exc. neurons	20
Surround input	All neurons (exc. and inh.) within a square of size 60 grid point centered at the center of the network. Neurons part of the center stimulus were excluded. 3286 exc. neurons and 3286 inh. neurons.	Poisson type spiking input with a spike rate of 100 Hz. Surround stimulus was presented for 200 ms.	2.58 mV at -54 mV	20
Surround input	All neurons (exc. and inh.) within a square of size 60 grid point centered at the center of the network. Neurons part of the center stimulus were excluded. 3286 exc. neurons and 3286 inh. neurons.	Poisson type spiking input with a spike rate of 50:10:130 Hz. Surround stimulus was presented for 200 ms.	2.58 mV at -54 mV	20

Supplementary Table 4. Neuron parameters

Parameter	Value	Description
C_m	200 pF	Membrane capacitance
g_L	20 nS	Leak conductance
V_{th}	-54 mV	Spike threshold
E_L	-70 mV	Reversal potential for leak conductance
t_{ref}	2 ms	Refractory time

Supplementary Table 5. Synapse parameters

Parameter	Value	Description	Notes
J_{ex}	0.63 mV	EPSP amplitude for connection from external excitatory inputs	EPSP amplitude was measured at a holding potential of -54 mV
J_{ix}	-0.3 mV	IPSP amplitude for connection from external inhibitory inputs	IPSP amplitude was measured at a holding potential of -54 mV
J_{ee}	{0.13, 0.26} mV	EPSP amplitude for exc. → exc. connection	EPSP amplitude was measured at a holding potential of -54 mV
J_{ei}	{0.11, 0.21, 0.32, 0.42, 0.53, 0.63} mV	EPSP amplitude for exc. → inh. connection	EPSP amplitude was measured at a holding potential of -54 mV
J_{ie}	{-0.14, -0.17, -0.19, -0.22, -0.28, -0.3} mV	IPSP amplitude for inh. → exc. connection	IPSP amplitude was measured at a holding potential of -54 mV
J_{ii}	0.055 mV	IPSP amplitude for inh. → inh. connection	IPSP amplitude was measured at a holding potential of -54 mV
tau_e	1.0 ms	Time constant of excitatory synapses	This was the same for both exc. → exc., exc. → inh. synapses and external excitatory input synapses

tau_i	3.0 ms	Time constant of inhibitory synapses	This was the same for both inh. → exc., inh. → inh. synapses and external inhibitory input synapses
d_{ee}	0.2 ms	Synaptic delay for exc. → exc. synapses	
d_{ei}	0.1 ms	Synaptic delay for exc. → inh. synapses	
d_{ie}	0.1 ms	Synaptic delay for inh. → exc. synapses	
d_{ii}	0.1 ms	Synaptic delay for inh. → inh. synapses	

970 **Supplementary Table 6. Network connectivity**

Parameter	Value	Description	Notes
N_e	6400	Excitatory neurons	
N_i	6400	Inhibitory neurons	
Excitatory grid size	80 × 80	Neurons were placed on a regular grid	
Inhibitory grid size	80 × 80	Neurons were placed on a regular grid	
C_{ee}	0.135	Connection probability between exc. to exc. neurons. This number refers to the integral of the connectivity kernel	Based on our experimental data (*See* Fig. 2g)
C_{ei}	0.112	Connection probability between exc. to inh neurons. This number refers to the integral of the connectivity kernel	Based on our experimental data (*See* Fig. 2g)
C_{ie}	0.315	Connection probability between inh to exc. neurons. This number refers to the integral of the connectivity kernel	Based on our experimental data (*See* Fig. 2g)
C_{ii}	0.3375	Connection probability between inh. to inh. neurons. This number refers to the integral of the connectivity kernel	Based on our experimental data (*See* Fig. 2g)
S_e	16 grid points 2% of the network size	Standard deviation of Gaussian used to estimate distance dependent connectivity from exc. to exc. and exc. to inh. neurons.	

S_i	20 grid points 2.5% of the network size	Standard deviation of Gaussian used to estimate distance dependent connectivity from inh. to exc. and inh. to inh. neurons.	

971
972

Supplementary Table 7. External Input

Parameter	Value	Description
nu_exc	1000 Hz	Poisson type spike trains were injected in all neurons
nu_inh	600 Hz	Poisson type spike trains were injected in all neurons

Data and code availability

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request. Data reported in this paper and all original code are available and will be shared by the lead contact upon request.

979
980

REFERENCES

1. Churchland, P. S. & Sejnowski, T. J. Neural Representation and Neural Computation. *Philos. Perspect.* **4**, 343–382 (1990).

2. Grossberg, S. The quantized geometry of visual space: The coherent computation of depth, form, and lightness. *Behav. Brain Sci.* **6**, 625–657 (1983).

3. Angelucci, A. *et al.* Circuits and Mechanisms for Surround Modulation in Visual Cortex. *Annu. Rev. Neurosci.* **40**, 425–451 (2017).

4. Knudsen, E. I. & Konishi, M. Center-Surround Organization of Auditory Receptive Fields in the Owl. *Science* **202**, 778–780 (1978).

5. Sachdev, R., Krause, M. & Mazer, J. Surround suppression and sparse coding in visual and barrel cortices. *Front. Neural Circuits* **6**, (2012).

6. Olsen, S. R. & Wilson, R. I. Lateral presynaptic inhibition mediates gain control in an olfactory circuit. *Nature* **452**, 956–960 (2008).

7. Hubel, D. H. & Wiesel, T. N. Receptive fields of single neurones in the cat's striate cortex. *J.*

Physiol. **148**, 574–591 (1959).

8. Schiller, P. H. & Koerner, F. Discharge characteristics of single units in superior colliculus of

the alert rhesus monkey. *J. Neurophysiol.* **34**, 920–936 (1971).

9. Kardamakis, A. A., Saitoh, K. & Grillner, S. Tectal microcircuit generating visual selection

commands on gaze-controlling neurons. *Proc. Natl. Acad. Sci.* **112**, E1956–E1965 (2015).

10. Kasai, M. & Isa, T. Imaging population dynamics of surround suppression in the superior

colliculus. *Eur. J. Neurosci.* **44**, 2543–2556 (2016).

11. McIlwain, J. T. & Buser, P. Receptive fields of single cells in the cat's superior colliculus. *Exp.*

Brain Res. **5**, 314–325 (1968).

12. Schiller, P. H. & Stryker, M. Single-unit recording and stimulation in superior colliculus of the

alert rhesus monkey. *J. Neurophysiol.* **35**, 915–924 (1972).

13. Bonin, V., Mante, V. & Carandini, M. The Suppressive Field of Neurons in Lateral Geniculate

Nucleus. *J. Neurosci.* **25**, 10844–10856 (2005).

14. Jones, H. E., Grieve, K. L., Wang, W. & Sillito, A. M. Surround suppression in primate V1. *J.*

Neurophysiol. **86**, 2011–2028 (2001).

15. Born, R. T. & Bradley, D. C. Structure and Function of Visual Area Mt. *Annu. Rev. Neurosci.*

28, 157–189 (2005).

16. Fernandes, A. M. *et al.* Neural circuitry for stimulus selection in the zebrafish visual system.

Neuron **109**, 805-822.e6 (2021).

17. Kardamakis, A. A., Pérez-Fernández, J. & Grillner, S. Spatiotemporal interplay between

multisensory excitation and recruited inhibition in the lamprey optic tectum. *eLife* **5**, e16472

(2016).

18. Flores-Herr, N., Protti, D. A. & Wässle, H. Synaptic currents generating the inhibitory surround

of ganglion cells in the mammalian retina. *J. Neurosci. Off. J. Soc. Neurosci.* **21**, 4852–4863

(2001).

19. Kim, Y. J. *et al.* Origins of direction selectivity in the primate retina. *Nat. Commun.* **13**, 2862

(2022).

20. Solomon, S. G., Lee, B. B. & Sun, H. Suppressive Surrounds and Contrast Gain in

Magnocellular-Pathway Retinal Ganglion Cells of Macaque. *J. Neurosci.* **26**, 8715–8726

(2006).

21. Zhang, Y., Kim, I.-J., Sanes, J. R. & Meister, M. The most numerous ganglion cell type of the

mouse retina is a selective feature detector. *Proc. Natl. Acad. Sci.* **109**, E2391–E2398 (2012).

22. Carandini, M. & Heeger, D. J. Normalization as a canonical neural computation. *Nat. Rev.*

Neurosci. **13**, 51–62 (2012).

23. Sato, T. K., Haider, B., Häusser, M. & Carandini, M. An excitatory basis for divisive

normalization in visual cortex. *Nat. Neurosci.* **19**, 568–570 (2016).

24. Veale, R., Hafed, Z. M. & Yoshida, M. How is visual salience computed in the brain? Insights

from behaviour, neurobiology and modelling. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, 20160113

(2017).

25. Zhang, X., Zhaoping, L., Zhou, T. & Fang, F. Neural Activities in V1 Create a Bottom-Up

Saliency Map. *Neuron* **73**, 183–192 (2012).

26. Zhaoping, L. From the optic tectum to the primary visual cortex: migration through evolution of

the saliency map for exogenous attentional guidance. *Curr. Opin. Neurobiol.* **40**, 94–102

(2016).

27. Ahmadlou, M., Tafreshiha, A. & Heimel, J. A. Visual Cortex Limits Pop-Out in the Superior

Colliculus of Awake Mice. *Cereb. Cortex* **27**, 5772–5783 (2017).

28. McPeek, R. M. & Keller, E. L. Deficits in saccade target selection after inactivation of superior

colliculus. *Nat. Neurosci.* **7**, 757–763 (2004).

29. Wang, L., Herman, J. P. & Krauzlis, R. J. Neuronal modulation in the mouse superior colliculus

during covert visual selective attention. *Sci. Rep.* **12**, 2482 (2022).

1046 30. Wang, L. & Krauzlis, R. J. Visual Selective Attention in Mice. *Curr. Biol.* **28**, 676-685.e4

1047   (2018).

1048 31. McLaughlin, T. & O'Leary, D. D. M. Molecular gradients and development of retinotopic

1049   maps. *Annu. Rev. Neurosci.* **28**, 327–355 (2005).

1050 32. Phongphanphanee, P. *et al.* Distinct local circuit properties of the superficial and intermediate

1051   layers of the rodent superior colliculus. *Eur. J. Neurosci.* **40**, 2329–2343 (2014).

1052 33. Haider, B., Häusser, M. & Carandini, M. Inhibition dominates sensory responses in the awake

1053   cortex. *Nature* **493**, 97–100 (2013).

1054 34. Wehr, M. & Zador, A. M. Balanced inhibition underlies tuning and sharpens spike timing in

1055   auditory cortex. *Nature* **426**, 442–446 (2003).

1056 35. Gale, S. D. & Murphy, G. J. Distinct Representation and Distribution of Visual Information by

1057   Specific Cell Types in Mouse Superficial Superior Colliculus. *J. Neurosci.* **34**, 13458–13471

1058   (2014).

1059 36. Liu, Y. *et al.* Mapping visual functions onto molecular cell types in the mouse superior

1060   colliculus. *Neuron* **111**, 1876-1886.e5 (2023).

1061 37. Ährlund-Richter, S. *et al.* A whole-brain atlas of monosynaptic input targeting four different

1062   cell types in the medial prefrontal cortex of the mouse. *Nat. Neurosci.* **22**, 657–668 (2019).

1063 38. Lazaridis, I. *et al.* A hypothalamus-habenula circuit controls aversion. *Mol. Psychiatry* **24**,

1064   1351–1368 (2019).

1065 39. Wickersham, I. R. *et al.* Monosynaptic restriction of transsynaptic tracing from single,

1066   genetically targeted neurons. *Neuron* **53**, 639–647 (2007).

1067 40. Reinhard, K. *et al.* A projection specific logic to sampling visual inputs in mouse superior

1068   colliculus. *eLife* **8**, e50697 (2019).

1069 41. Gehr, C., Sibille, J. & Kremkow, J. Retinal input integration in excitatory and inhibitory

1070   neurons in the mouse superior colliculus in vivo. 2023.04.07.536092 Preprint at

1071   https://doi.org/10.1101/2023.04.07.536092 (2023).

42. Hong, Y. K., Kim, I.-J. & Sanes, J. R. Stereotyped axonal arbors of retinal ganglion cell subsets

in the mouse superior colliculus. *J. Comp. Neurol.* **519**, 1691–1711 (2011).

43. Niell, C. M. & Scanziani, M. How Cortical Circuits Implement Cortical Computations: Mouse

Visual Cortex as a Model. *Annu. Rev. Neurosci.* **44**, 517–546 (2021).

44. Adesnik, H., Bruns, W., Taniguchi, H., Huang, Z. J. & Scanziani, M. A neural circuit for spatial

summation in visual cortex. *Nature* **490**, 226–231 (2012).

45. O'Rawe, J. F. *et al.* Excitation creates a distributed pattern of cortical suppression due to varied

recurrent input. *Neuron* (2023) doi:10.1016/j.neuron.2023.09.010.

46. Douglas, R. J., Koch, C., Mahowald, M., Martin, K. A. & Suarez, H. H. Recurrent excitation in

neocortical circuits. *Science* **269**, 981–985 (1995).

47. Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**, 51–

58 (2013).

48. Guo, L. & Kumar, A. Role of interneuron subtypes in controlling trial-by-trial output variability

in the neocortex. *Commun. Biol.* **6**, 1–13 (2023).

49. Baden, T. *et al.* The functional diversity of retinal ganglion cells in the mouse. *Nature* **529**,

345–350 (2016).

50. Kapadia, M. K., Ito, M., Gilbert, C. D. & Westheimer, G. Improvement in visual sensitivity by

changes in local context: parallel studies in human observers and in V1 of alert monkeys.

Neuron **15**, 843–856 (1995).

51. Krieger, B., Qiao, M., Rousso, D. L., Sanes, J. R. & Meister, M. Four alpha ganglion cell types

in mouse retina: Function, structure, and molecular signatures. *PLoS ONE* **12**, e0180091 (2017).

52. Brenner, J. M., Beltramo, R., Gerfen, C. R., Ruediger, S. & Scanziani, M. A genetically defined

tecto-thalamic pathway drives a system of superior-colliculus-dependent visual cortices.

Neuron **111**, 2247-2257.e7 (2023).

53. Zénon, A. & Krauzlis, R. J. Attention deficits without cortical neuronal deficits. *Nature* **489**,

434–437 (2012).

54. Fardet, T. *et al.* NEST 2.20.0. (2020) doi:10.5281/zenodo.3605514.

55. Fürth, D. *et al.* An interactive framework for whole-brain maps at cellular resolution. *Nat.*

Neurosci. **21**, 139–149 (2018).

1101

SUPPLEMENTARY FIGURES AND FIGURE LEGENDS (S1-S7)

Fig. S1: Mapping excitatory and inhibitory zones of SCs neurons

(a-b) A wide-field (WF) SCs neuron responding with the neuron responding with excitatory and inhibitory postsynaptic currents (EPSCs measured at -65 mV shown in **a**; IPSCs measured at 0 mV shown in **b**) in response to a 20 Hz 5-light-pulse train delivered pseudo-randomly as a 120×100 µm grid optostimulation. Bottom panels: Response maps superimposed with the reconstructed morphology of the intracellularly stained WF neuron with Neurobiotin during whole-cell recording. Heat maps with darker shades coding for stronger responses in amplitudes.

(c) Distance from soma as a function of normalized postsynaptic amplitude. Red for excitatory; Blue for inhibitory. Vertical dashed line indicates threshold applied at a normalized value of 0.2.

(d) Center and surround zone of the WF zone determined by following thresholding approach applied to the excitatory PSCs shown in **(c)**. Areas occupying a larger value compared to threshold are considered to be part of the center zone.

1121

1122

1123

Fig. S2: Three different examples of synaptic modulation between center and surround zones.
(a) Top: Individual current traces held at reverse potentials for chloride-mediated GABAergic
inhibition (-65 mV) and glutamate-mediated ionotropic excitation (0 mV) that span continuously
surround and center optostimulation of the neuron (2 pulses separated by 20 ms) shown in Fig. 1d.
Here, surround network activation can be seen to effectively reduce the center EPSC amplitudes
(blue rectangle) when comparing to center only responses (in magenta). Evoked responses to
surround stimulation of all durations (5, 20 and 50 ms; shades of green) are also seen to drive
bistable excitatory and inhibitory responses. (Bottom) Voltage-clamp (VC) recordings in response to
surround optostimulation after bath application of 10 µM Gabazine (GBZ), thereby removing the
effect of GABAergic inhibition. Same color-coding scheme as in Fig. 1. Time stamps are centered
around the onset of the first center pulse.
(b) (Top) Synaptic current recordings performed in voltage clamp of a stellate neuron shown in Fig.
1e that is subject to the 20 Hz optostimulation (2 pulses). Center EPSCs shown in the rectangle are
truncated (dotted line). (Bottom) Same neuron recorded in current clamp (CC) reveals the shunting
effect that surround activation has on center responses (gray rectangle) when recording at -65 mV
and at a depolarized membrane potential of -20 mV. This is attributed to chloride current triggered
by the surround (revealed when held at 0 mV in VC; top traces).
(c) Current traces from a vGAT+ SCs neuron in response to 20 Hz optostimulation (5 center pulses)
that shows signs of surround suppression without the direct effect of synaptic inhibition. Top and
middle traces: Current recordings performed at -65 mV and 0 mV reveals that this neuron does not
receive direct inhibition from the surround yet its center response is attenuated suggesting that it
receives less excitatory input. Bottom: Enlarged view of current traces found in blue rectangle in
middle panel focusing on first and second paired pulse center stimulations. Surround stimulation has
a significant effect on the first center pulse by reducing its amplitude by 50% (***, $p < 0.0001$;
Student's t-test).

1149
1150
Fig. S3: Center-surround dynamics are defined by transient excitatory and inhibitory
conductances.
(a) Total synaptic conductance (G_T, black line) triggered by center only (C_1) and surround-center (C_s)
conditions is decomposed into excitatory (G_e, red) and inhibitory (G_i, blue) conductances.
Conductances belong to recording shown in Fig. 1f, which are estimated by Wehr and Zador (2003)
by using synaptic current recordings performed at -65 mV and 0 mV. Time stamps t_c, t_e and t_T
symbolize the time onset for center stimulation, the time instant when peak excitation and maximal
total conductance is reached, respectively. Solid lines are with center only condition and dashed
lines are when center follows surround optostimulation. Red arrow indicates non-zero inhibitory
conductance evoked by surround network activation at the time of t_c.
(b) Comparison of total synaptic conductance G_T between C_1 and C_s conditions. Corresponding
excitatory and inhibitory conductances at peak G_T are shown in Fig. 1g-h. Color coding corresponds
to morphologically or genetically identified SCs neuron cell-types (*see* Fig. 1g).
(c) Full time course of excitatory and inhibitory conductance during both stimulation conditions, C_1
and C_s. *Legends:* C-C (paired pulse center optostimulation) and S-C refers to surround followed by
center optostimulation after a 20 ms refractory period.
(d) Latency to peak inhibition as a function of latency to peak excitation. On average, when
comparing the center (magenta) and surround (cyan) conditions, it appears that there is an
approximately 20 ms delay. This suggests that the signals originating from the surround region
interact with the center responses with this time difference. Curves are a gaussian probability
distribution function fitted to the data points.

1172
1173
Fig. S4: Identification of SCs cell types used for electrophysiology analysis
(a) Reconstructed morphologies from various subtypes of SCs neurons that were recorded and
presented in Fig. 1, including wide-field, narrow-field, stellate and horizontal neurons, that were
stained with Neurobiotin.
(b-c) Quantification and classification of cell types according to morphology (i.e. horizontal, stellate,
etc.) and genotype (i.e. vGAT+/- or random type). A total of $n = 41$ were performed of which a subset
of $n = 24$ were included in the analysis shown in Fig. 1.
(d) Relationship between dendritic field and the neuron's corresponding retinorecipient center zone.
Individual center (C) zones typically exceed the area of the dendritic zone. Note that this correlation
was performed for recorded neurons where morphological data were successfully obtained.

Fig. S5: Classification of vGluT2 and vGAT cell types and their distribution across the SCs.
(a) Coexpression of vGAT and vGluT2 *in situ* hybridization in the SCs using RNAscope ISH
performed to reveal the identity of input neurons. White arrows indicate Rb-EGFP+ cells that are
positive for vGluT2 and negative for vGAT, suggesting that these input neurons are excitatory. By
contrast blue arrows show Rb-EGFP+ cells that are inhibitory since they are positive for vGAT and
negative for vGluT2. Scale bar: 25 µm.
(b) The proportion of vGluT2+ (red, n = 70) and vGAT+ (blue, n = 199) input cells that colocalize
with Rb-EGFP. Unidentified neurons shown in gray (n = 3/272). N = 1 animal for each group, 3
sections were used per animal. Total number of counted neurons is: 272.
(c) RNAscope ISH for identity confirmation of starter neurons (*see* white arrows). Top row:
Colocalization of vGlut2 detected in cells that express Rb-EGFP and the Cre-inducible V5 tag
expressed in vGluT2-Cre mouse line, confirm starter neurons are excitatory. Bottom row: The same
approach applied in a vGAT-Cre. Colocalization Rb-EGFP+ with V5+ but negative for vGluT2,
suggests that the starter neuron is putatively inhibitory. Scale bar: 25 µm.
(d) the distribution of starter neurons in each animal, shown as proportion in different sublayers. The
cell count for starter neurons of each animal are: vGLUT2 #1: 52; #2: 16; #3: 79. vGAT #1: 5; #2:
17; #3: 106.) SCig: SC, motor related, intermediate gray layer, SCop: superior colliculus, optic layer,
SCsg: SC, superficial gray layer; SCzo: SC, zonal layer.
(e) Normalized density plot of starter and input neurons along anterior-posterior axis (AP; Left),
medio-lateral axis (ML; Center) or depth from SC surface (Right). Distribution of excitatory and
inhibitory neurons are shown in the top and bottom rows, respectively. The ML density plot is
computed on the relative coordinates to the k-means centroid of starter cells. Dashed lines represent
individual animals and solid lines show the average.

1209
Fig. S6 Complete raster of spiking network simulation (supplemental to Fig. 3b)
(a) Entire time course of network simulation illustrating the raster of spiking activity of excitatory (red
dots) and inhibitory SCs neurons (blue points) during the four stimulus conditions shown in Fig. 3.
Vertical green lines show the time onset of 'center *only'* stimulation; black horizontal lines show
'surround' stimulation; vertical red and blue lines indicate the time onset of 'center *during* surround'
and 'center *after* surround'. A sub-region of this trial containing all simulation conditions has been
selected (orange shaded region) and shown in Fig. 3b-c.
(b) Spiking frequency of model inhibitory neurons as a function of the spiking frequency of excitatory
neurons. The inverse relationship is correlated of the peak amplitude ratio of the 'center *after*
surround' divided by the 'center *before* surround' response (*see* color map).
1220
1221

1222
1223
Fig. S7: Functional shifts in surround network E/I balance (supplemental to Fig. 4d)
In vivo cFos expression resulting from local optostimulaon of RGC axonal terminals in two regions
of interest: 1) center zone defined as the area under the tip of the optic fiber implanted in the SCs
(top row), and 2) lateral zone relative to the center referred to as 'surround lateral' (SL; bottom row).
From left to right columns: a) merged image, b) *c*Fos staining, c) GABA immunostaining (arrows
show colocalization), and d) ChR2-expressing retinal ganglion axonal terminals. ChR2 shown in
magenta; *c*Fos in yellow; GABA in cyan. A higher ratio of active inhibitory neurons is found to be
engaged in the surround after center zone stimulation (*see* Fig. 4d). Scale bar is 50 μm.