**Uncovering the mosaic evolution of carnivoran skeletal systems**

Chris J. Law[1,2], Leslea J. Hlusko[3], Z. Jack Tseng[4]

[1]Department of Integrative Biology, University of Texas

[2]Burke Museum and Department of Washington, University of Washington

[3]National Research Center on Human Evolution (CENIEH), Burgos, Spain

[4]Department of Integrative Biology, University of California Berkeley

**Abstract**

The diversity of vertebrate skeletal forms is often attributed to adaptations to distinct ecological

factors such as diet, locomotion, and sensory environment. Although the adaptive evolution of

cranial, appendicular, and vertebral skeletal systems is well studied in vertebrates,

comprehensive investigations of all skeletal components are rarely performed simultaneously.

Consequently, we know little of how modes of evolution differ among skeletal components. We

collected 103 linear measurements from 208 osteological museum specimens to capture skeletal

variation across the major components of the skeletal systems of 119 extant carnivores. With

these data, we tested if ecological and phylogenetic effects led to distinct modes of evolution

among the cranial, appendicular, and vertebral skeletal regions by fitting multivariate

evolutionary models. We found mosaic evolution in which only the mandible, hindlimb, and

posterior region of the vertebral column showed evidence of adaptation towards ecological

regimes whereas the remaining skeletal components followed clade-specific evolutionary shifts.

We hypothesize that the decoupled evolution of individual skeletal components may have led to

the origination of distinct adaptive zones and morphologies among extant carnivoran families

that reflect phylogenetic hierarchies. Overall, our work highlights the importance of examining

multiple skeletal components simultaneously in ecomorphological analyses. Ongoing work

integrating the fossil and paleoenvironmental record will further govern carnivoran diversity we

see today and reveal the complexity of evolutionary processes in multicomponent systems.

Key words: adaptive landscape; Carnivora; ecomorphology; macroevolution; Ornstein-

Uhlenbeck modeling; phylogenetic comparative methods

**Introduction**

The diversity of animal forms is one of the most salient patterns across the tree of life.

For example, within mammals, morphological innovations in the cranial, appendicular, and axial

skeletal systems facilitate the incredible diversity found today, ranging from flying bats with

winged forelimbs to the biggest animals to have ever lived on earth. Many researchers have

examined how variation in the skull [1–6], limbs [7–10], or vertebrae [11–15] serve as crucial

adaptations to their evolution. These skeletal systems (skull, limbs, and vertebrae) are

traditionally independently examined and have rarely been investigated simultaneously even

though these anatomical regions comprise a single, functionally integrated system to support

movement, sensation, and other life functions. When considered wholistically, the observed

variation across the different components of organismal anatomy are generally explained by

multitudinous factors, some that are potentially incongruous. While this evolutionary push-and-

pull between anatomical regions may characterize the process of evolution, the hypothesis can

only be tested when the different components of the skeleton are explored simultaneously rather

than piecemeal. Such anatomically integrated studies are critical to our understanding of the role

of developmental and/or functional integration in canalizing macroevolutionary trajectories [16–

18].

Here, we use carnivorans as a model to investigate how ecological and phylogenetic

factors correspond with evolutionary changes in the cranial, appendicular, and axial skeletal

systems simultaneously. Carnivorans (bears, cats, dogs, seals, weasels, and their relatives) are an

especially productive model system to examine skeletal evolution because of their high species

richness and vast geographic distribution across most biomes in all continents and oceans, factors

that have led to consequently broad ecological diversity in locomotor traits (e.g., arboreal,

cursorial, fossorial, aquatic) and feeding adaptations ranging from herbivory to generalists to

hypercarnivory.

Components of carnivoran skeletal systems are well studied individually. In the skull,

craniomandibular diversity is influenced by several ecological factors and phylogeny [19–23].

Recently, decoupled modes of evolution were found within the skull: cranial shape follows

clade-specific evolutionary shifts suggesting phylogenetic constraints, whereas mandibular shape

evolution is linked to broad dietary regimes [6]. In the appendicular skeleton, researchers

examined ecomorphological divergence between the hindlimbs, which are adapted primarily for

locomotion, and the forelimb, which are adapted for multiple functions ranging from running to

grappling prey to manipulating objects [24–27]. Additionally, more recent work using

phylogenetic comparative methods found that scaling and phylogeny exhibit stronger effects on

limb skeletal evolution than do ecological parameters [28–30]. In contrast to craniomandibular

and appendicular ecomorphology, research on the axial skeleton is in nascent stages. Initial

research has found that distinct regions of the vertebral column are under different evolutionary

pressures. For example, the anterior region exhibits low disparity due to phylogenetic constraints

or ecological conservatism, whereas the posterior region exhibits higher disparity that may be

due to adaptations to various locomotor ecologies [12,31]. In contrast to these anatomically-

localized studies, researchers examining proxies of the overall body plan report that the evolution

of traits like body mass, skeletal size, and body shape often follow a Brownian motion model or

clade-based shift model rather than being associated with ecological regimes [23,32,33].

76           Although well studied individually, simultaneous investigation of skulls, limbs,

vertebrae, and overall body plan are rarely conducted, likely because of the enormous amount of

data that would need to be collected and the complexity of the multivariate analyses required.

However, most researchers would likely agree that a more comprehensive approach to skeletal

evolution is essential to elucidate the complexity of skeletal evolution. The search for system-

level trends and variations is further obscured by the disparate methods employed to test the

effects of ecology and phylogeny on different skeletal systems by different researchers. In this

study, we address both of these issues, combining anatomical data from across the entire skeleton

and applying the same analytical methods. Here, we investigate the mosaic evolution of

carnivoran skeletons by creating a new skeletal trait dataset that encompasses all major

components of the skeletal system. We then tested the ecological and phylogenetic effects

influencing the modes of evolution of the various skeletal components using multivariate

generalized evolutionary models.

**Methods**

*Skeletal and ecological traits*

We collected 103 linear measurements from 208 osteological specimens to capture the

skeletal morphology of 119 extant carnivorans (Fig. S1; Table S1). This data set includes seven

cranial traits, seven mandibular traits, 13 forelimb traits, 13 hindlimb traits, and seven traits in

third cervical, fifth cervical, first thoracic, middle thoracic, diaphragmatic thoracic, last thoracic,

first lumbar, middle lumbar, and last lumbar vertebrae. For these analyses, we focus only on e

male specimens in order to remove the complicating factor of sexual dimorphism as carnivorans

exhibit differing degrees of sexual dimorphism [34,35]. To remove the effects of scaling, we

calculated log shape ratios by dividing each skeletal trait by the geometric mean of all 103 traits

as our metric of size. We then used individual principal component analyses (PCAs) to reduce

the dimension of each skeletal component (i.e., cranium, mandible, forelimb, hindlimb, and each

of the nine vertebrae) and retained a number of PC axes that corresponded to at least 90% of the

explained variance. We also conducted a PCA on the entire dataset as our proxy of the whole-

skeleton phenome and retained the first six PC axes (~75% of explained variance) for subsequent

analyses. We classified the 119 carnivoran species into distinct locomotor modes, hunting

behaviors, and dietary regimes following (Law 2021).

*Phylogenetic comparative methods*

We tested whether each skeletal component evolved as adaptation to diet, hunting

behavior, or locomotor mode or exhibited clade-specific evolutionary shifts by fitting

multivariate generalized evolutionary models on the retained PC axes of each individual skeletal

component [36–38]. For the adaptive ecological models, we fit three multivariate multi-optima

Ornstein-Uhlenbeck models (i.e., $mvOUM_{diet}$ , $mvOUM_{hunting\ behavior}$, and $mvOUM_{locomotion}$) to test

if dietary, hunting behavioral, or locomotor regimes influenced the evolution of each set of PC

axes using mvMORPH v1.1.4 [38]. All three models were fit across 500 stochastically mapped

trees to account for uncertainty in phylogenetic topology and ancestral character states (see

electronic supplementary materials for full methods). We also calculated the phylogenetic half-

lives (ln(2)/alpha) of the best supported adaptive ecological model, defined as the time it takes

for a trait to evolve halfway toward its new expected optimum after a regime shift [36]. A short

phylogenetic half-life relative to the age of Carnivora (48.2 myr) would suggest that skeletal

traits are strongly pulled toward distinct ecological peaks across the adaptive landscape. For the

clade-based model, we fit a multi-optima OU model (mvOUM$_{phyloEM}$) without *a priori* ecological

groupings with the PhylogeneticEM v1.4.0 [39]. We also fit a single-rate multivariate Brownian

motion model (mvBM1) and a single-optimum OU model (mvOU1) as null models. Relative

support of models was assessed through computation of small sample-corrected Akaike weights

(AICcW).

Preliminary results revealed that phenotypic differences between pinnipeds (i.e., seals

and sea lions) and terrestrial carnivorans are often the greatest source of variation for most

skeletal components. These results are unsurprising considering pinnipeds exhibit derived

morphologies that enable them to be fully aquatic. Therefore, we repeated the above model

fitting using a reduced dataset with no pinnipeds. Results of the full dataset that includes

pinnipeds are presented in the electronic supplementary material.

**Results and Discussion**

We found mosaic evolution of the carnivoran skeleton in which ecology and phylogeny

have differing influences on the evolutionary mode of the various skeletal components.

Consistent with recent results, the carnivoran skull exhibited decoupled modes of evolution

between the cranium and mandible. In the cranium, the clade-specific shift model exhibited

overwhelmingly greater support (mvOUM$_{phyloEM}$; AICcW > 0.99) compared to adaptive

ecological models based on diet, hunting behavior, and locomotion (Fig. 1; Table S2). We found

eight evolutionary shifts in cranial morphology that correspond to carnivoran clades (Fig. S2A).

In contrast, the adaptive ecological model based on diet was the best supported model

(mvOUM$_{diet}$; AICcW = 0.96) for the mandible with a short phylogenetic half-life of 2.52 myr

(Fig. 1; Table S2). These results are congruent with previous findings revealing that mandibular

shape is evolutionarily labile with respect to dietary evolution whereas cranial shape is

partitioned between families rather than between dietary groups [6]. Decoupling between the

cranium and mandible may be explained by their functions. Diet is often found to have had a

strong influence on the evolution of the mandible because of its direct mechanical role in feeding

[3,40–44]. However, in addition to feeding, the multiple sensory functions that the cranium also

performs also influence its evolution [45–47]; therefore, cranial shape may be more constrained

in its ability to evolve to match dietary demands alone.

The appendicular system exhibited decoupled modes of evolution between forelimbs and

hindlimbs. The clade-specific shift model (mvOUM$_{phyloEM}$; AICcW > 0.99) was best supported

in the forelimb (Fig. 1; Table S2). Seven evolutionary shifts in forelimb evolution occur

primarily along familial branches (Fig. S2B), indicating that the complexity and variation of

carnivoran forelimb adaptations cannot be captured effectively by dietary, hunting behavioral, or

locomotor categories. Instead, these shifts suggest that clade-specific adaptations enabled the

diversity of forelimb skeletons for tasks such as grappling or manipulating prey, swimming, or

digging [24–27,30,48,49]. For example, most felids use their prehensile forelimbs to ambush and

subdue prey, most canids and hyaenids pounce and pursue prey, and some mustelids use their

powerful forelimbs to dig out prey while other, more derived mustelids (i.e., weasels and

polecats) pursue prey in tight crevices and burrows [50]. In contrast, the adaptive ecological

model based on locomotor mode was the best supported model (mvOUM$_{\text{locomotion}}$; AICcW =

0.83) in the hindlimb with a short phylogenetic half-life of 5.05 myr (Fig. 1; Table S2),

supporting hypotheses that the hindlimb is adapted primarily for locomotion as typically found in

quadrupedal mammals [51]. Together, this body of work is in line with findings that non-

cursorial carnivorans exhibit weaker integration between the limbs [30] and supports the

hypothesis of functional divergence between the forelimbs and hindlimbs of carnivorans.

169           The axial skeleton exhibits distinct modes of evolution between the anterior and posterior

regions of the vertebral column: clade-specific shift or single-peak OU models tended to be the

best supported model for cervical and most thoracic vertebrae, whereas adaptive ecological

models based on hunting behavior or locomotor mode were the best supported model for the last

thoracic and all lumbar vertebrae (Fig. 1; Fig. S2C–F; Table S2). Our findings strengthen the

coalescing hypothesis that anterior vertebrae exhibit lower disparity, higher evolutionary

constraints, and more subtle adaptations to locomotion whereas posterior vertebrae exhibit the

opposite pattern in carnivorans [31] and broadly across mammals [13]. We posit that high

evolutionary constraints of the anterior vertebrae are associated with clade-specific shifts in the

cervical and most thoracic vertebrae. In contrast, relaxed evolutionary constraints of the posterior

vertebrae facilitate the evolution of disparate lumbar vertebrae across the entire carnivoran order.

These disparate vertebrae serve as adaptations to diverse locomotor modes or hunting behaviors

based on the mobility of the posterior backbone and irrespective of clade origins. The short

phylogenetic half-lives (1.47–5.12 myr; Fig. 1) further suggests strong pulls towards these

different adaptive optima. More broadly, this increased mobility of the lumbar region over

evolutionary time is hypothesized to be an evolutionary innovation for the evolution of extant

mammals [13,52,53].

Overall, we elucidate the mosaic evolution of the carnivoran skeleton, finding that

different skeletal components exhibit distinct modes of evolution. Our results suggest that

different methodologies and taxonomic samples do not necessarily explain previously reported

region-specific macroevolutionary patterns; rather, complexity in explanatory factors of skeletal

diversity is a key feature of Carnivora. The ability of individual skeletal components to adapt to

specific ecological factors independently from each other may have contributed to the clade's

*hierarchical* [54,55] evolution. As previously hypothesized [23,32], the restriction of carnassial

shear to the P4/m1 pair may have been the key innovation that facilitated the initial carnivoran

diversification early in the clade's evolutionary history. Subsequent evolution led to the

continual partitioning between clades, resulting in the origination of extant carnivoran families as

discrete phylogenetic clusters that occupy different adaptive zones [56] with distinct

morphologies including body size and shape [33,57], various components of the skeleton [e.g.,

skull 6]; Fig. 1), as well as the overall skeletal phenome (Fig. 2). Within-clade variation then

arises to reflect resource partitioning among ecologically similar taxa, leading to adaptations in

morphologies such as the mandible, hindlimb, and posterior region of the vertebral column (Fig.

1). These traits were strongly pulled toward distinct ecological peaks across the adaptive

landscape as revealed by their short phylogenetic half-lives (1.47–5.12 myr) relative to the age of

Carnivora itself (48.2 myr).

Our research statistically revealed the mosaic evolution of carnivoran skeletons. These

distinct evolutionary modes demonstrate the importance of examining multiple skeletal

components in ecomorphological analyses. Nevertheless, key questions remain: 1) what spurred

the evolutionary transitions towards the evolutionary shifts or adaptations of the various skeletal

components? 2) when in the 55 million years of carnivoran evolutionary history did these

evolutionary events occur? And lastly, 3) what developmental and genetic phenomena underlie

the dissociation of various skeletal elements? Ongoing work integrating the fossil and

paleoenvironmental record will further elucidate the carnivoran diversity we see today and reveal

the complexity of evolutionary processes in multicomponent systems.

**Acknowledgements**

We are grateful to the staff and collections at the American Museum of Natural History,

California Academy of Sciences, Field Museum of Natural History, Natural History Museum of

Los Angeles County, Museum of Vertebrate Zoology, Natural History Museum London, San

Diego Natural History Museum, Texas Vertebrate Paleontology Collection, National Museum of

Natural History, and Burke Museum of Natural History and Culture.

**Funding**

Funding was supported by the National Science Foundation (DBI–2128146). CJL was also

supported by a University of Texas Early Career Provost Fellowship and Stengl-Wyer

Endowment Grant (SWG-22-02). L.J.H. was also supported by the European Research Council

(ERC) under the European Union's Horizon 2020 research and innovation program (Tied2Teeth,

grant agreement n° 101054659).

**Fig. 1.** Diagram of the skeletal components and their best-fitting evolutionary model. AICcW are

in parentheses. See Table S2 for full AICc table. $t_{1/2}$ = phylogenetic half-life calculated as ln(2)/

α for adaptive ecological models. Myr = million years. C3 = third cervical vertebrae; C5 = fifth

cervical vertebrae; T1 = first thoracic vertebrae; midT = middle thoracic vertebrae; diaT =

diaphragmatic thoracic vertebrae; lastT = last thoracic vertebrae; L1 = first lumbar vertebrae;

midL = middle lumbar vertebrae; lastL = last lumbar vertebrae.

**Fig. 2.** A) Phylomorphospace and B) clade-specific evolutionary shifts of the full skeletal dataset

across terrestrial carnivorans identified by PhylogeneticEM. Shifts are calculated from PCs 1-6

(74.7% of explained variance) and are represented as pink circles. Branches on the phylogenies

are colored according to each regime.

References

1. Dumont ER, Dumont E, Davalos LM, Goldberg A, Santana SE, Santana SE, Rex K, Voigt CC. 2012 Morphological innovation, diversification and invasion of a new adaptive zone. *Proceedings of the Royal Society B: Biological Sciences* **279**, 1797–1805. (doi:10.1098/rspb.2011.2005)

2. Zelditch ML, Ye J, Mitchell JS, Swiderski DL. 2017 Rare ecomorphological convergence on a complex adaptive landscape: Body size and diet mediate evolution of jaw shape in squirrels (Sciuridae). *Evolution* **71**, 633–649. (doi:10.1111/evo.13168)

3. Grossnickle DM. 2020 Feeding ecology has a stronger evolutionary influence on functional morphology than on body mass in mammals. *Evolution* **74**, 610–628. (doi:10.1111/evo.13929)

4. Arbour JH, Curtis AA, Santana SE. 2019 Signatures of echolocation and dietary ecology in the adaptive evolution of skull shape in bats. *Nat Commun* **10**, 2036. (doi:10.1038/s41467-019-09951-y)

5. Goswami A *et al.* 2022 Attenuated evolution of mammals through the Cenozoic. *Science* **378**, 377–383. (doi:10.1126/science.abm7525)

6. Law CJ, Blackwell EA, Curtis AA, Dickinson E, Hartstone-Rose A, Santana SE. 2022 Decoupled evolution of the cranium and mandible in carnivoran mammals. *Evolution* **76**, 2959–2974. (doi:10.1111/evo.14578)

7. Patel BA. 2010 Functional morphology of cercopithecoid primate metacarpals. *J. Hum. Evol.* **58**, 320–337. (doi:10.1016/j.jhevol.2010.01.001)

8. Kilbourne BM. 2017 Selective regimes and functional anatomy in the mustelid forelimb: Diversification toward specializations for climbing, digging, and swimming. *Ecol Evol* **66**, 2369–12. (doi:10.1002/ece3.3407)

9. Grossnickle DM, Chen M, Wauer JGA, Pevsner SK, Weaver LN, Meng QJ, Liu D, Zhang YG, Luo ZX. 2020 Incomplete convergence of gliding-mammal skeletons. *Evolution* **74**, 2662–2680. (doi:10.1111/evo.14094)

10. Rickman J, Burtner AE, Linden TJ, Santana SE, Law CJ. 2023 Size And Locomotor Ecology Have Differing Effects on the External and Internal Morphologies of Squirrel (Rodentia: Sciuridae) Limb Bones. *Integr. Org. Biol.* **5**, obad017. (doi:10.1093/iob/obad017)

11. Pierce SE, Clack JA, Hutchinson JR. 2011 Comparative axial morphology in pinnipeds and its correlation with aquatic locomotory behaviour. *J Anat* **219**, 502–514. (doi:10.1111/j.1469-7580.2011.01406.x)

12. Randau M, Cuff AR, Hutchinson JR, Pierce SE, Goswami A. 2017 Regional differentiation of felid vertebral column evolution: a study of 3D shape trajectories. *Org Divers Evol* **17**, 1–15. (doi:10.1007/s13127-016-0304-4)

13. Jones KE, Benitez L, Angielczyk KD, Pierce SE. 2018 Adaptation and constraint in the
evolution of the mammalian backbone. *Bmc Evol Biol* **18**, 172. (doi:10.1186/s12862-018-1282-
2)

14. Martín-Serra A, Pérez-Ramos A, Pastor FJ, Velasco D, Figueirido B. 2021 Phenotypic
integration in the carnivoran backbone and the evolution of functional differentiation in
metameric structures. *Evolution Letters* , 1–14. (doi:10.1002/evl3.224)

15. Linden TJ, Burtner AE, Rickman J, McFeely A, Santana SE, Law CJ. 2023 Scaling patterns
of body plans differ among squirrel ecotypes. *PeerJ* **11**, e14800. (doi:10.7717/peerj.14800)

16. Goswami A, Randau M, Polly PD, Weisbecker V, Bennett CV, Hautier L, Sánchez-Villagra
MR. 2016 Do Developmental Constraints and High Integration Limit the Evolution of the
Marsupial Oral Apparatus? *Integr. Comp. Biol.* **56**, 404–415. (doi:10.1093/icb/icw039)

17. Goswami A, Weisbecker V, Sánchez-Villagra MR. 2009 Developmental modularity and the
marsupial–placental dichotomy. *J. Exp. Zoöl. Part B: Mol. Dev. Evol.* **312B**, 186–195.
(doi:10.1002/jez.b.21283)

18. Goswami A, Smaers JB, Soligo C, Polly PD. 2014 The macroevolutionary consequences of
phenotypic integration: from development to deep time. *Philos. Trans. R. Soc. B: Biol. Sci.* **369**,
20130254. (doi:10.1098/rstb.2013.0254)

19. Radinsky LB. 1981 Evolution of skull shape in Carnivores .1. Representative modern
Carnivores. *Biol J Linn Soc* **15**, 369–388. (doi:10.1111/j.1095-8312.1981.tb00770.x)

20. Valkenburgh BV. 2007 Deja vu: the evolution of feeding morphologies in the Carnivora.
*Amer. Zool.* **47**, 147–163. (doi:10.1093/icb/icm016)

21. Law CJ, Duran E, Hung N, Richards E, Santillan I, Mehta RS. 2018 Effects of diet on cranial
morphology and biting ability in musteloid mammals. *J. Evol. Biol.* **31**, 1918–1931.
(doi:10.1111/jeb.13385)

22. Tseng ZJ, Flynn JJ. 2018 Structure-function covariation with nonfeeding ecological variables
influences evolution of feeding specialization in Carnivora. *Sci Adv* **4**, eaao5441.
(doi:10.1126/sciadv.aao5441)

23. Slater GJ, Friscia AR. 2019 Hierarchy in adaptive radiation: A case study using the
Carnivora (Mammalia). *Evolution* **73**, 524–539. (doi:10.1111/evo.13689)

24. Valkenburgh BV. 1985 Locomotor diversity within past and present guilds of large predatory
mammals. *Paleobiology* **11**, 406–428. (doi:10.1017/s0094837300011702)

25. Valkenburgh BV. 1987 Skeletal indicators of locomotor behavior in living and extinct
carnivores. *J Vertebr Paleontol* **7**, 162–182. (doi:10.1080/02724634.1987.10011651)

26. Iwaniuk AN, Pellis SM, Whishaw IQ. 1999 The relationship between forelimb morphology and behaviour in North American carnivores (Carnivora). *Can J Zool* **77**, 1064–1074. (doi:10.1139/z99-082)

27. Samuels JX, Meachen JA, Sakai SA. 2013 Postcranial morphology and the locomotor habits of living and extinct carnivorans. *J Morpho* **274**, 121–146. (doi:10.1002/jmor.20077)

28. Martín-Serra A, Figueirido B, Palmqvist P. 2014 A Three-Dimensional Analysis of Morphological Evolution and Locomotor Performance of the Carnivoran Forelimb. *Plos One* **9**, e85574. (doi:10.1371/journal.pone.0085574)

29. Martín-Serra A, Figueirido B, Palmqvist P. 2014 A three-dimensional analysis of the morphological evolution and locomotor behaviour of the carnivoran hind limb. *BMC Evol Biol* **14**, 1–13. (doi:10.1186/1471-2148-14-129)

30. Martín-Serra A, Figueirido B, Pérez-Claros JA, Palmqvist P. 2015 Patterns of morphological integration in the appendicular skeleton of mammalian carnivores. *Evolution* **69**, 321–340. (doi:10.1111/evo.12566)

31. Figueirido B, Martín-Serra A, Pérez-Ramos A, Velasco D, Pastor FJ, Benson RJ. 2021 Serial disparity in the carnivoran backbone unveils a complex adaptive role in metameric evolution. *Communications Biology* , 1–15. (doi:10.1038/s42003-021-02346-0)

32. Slater GJ. 2022 Topographically distinct adaptive landscapes for teeth, skeletons, and size explain the adaptive radiation of Carnivora (Mammalia). *Evolution* (doi:10.1111/evo.14577)

33. Law CJ. 2021 Ecological drivers of carnivoran body shape evolution. *Am Nat* **198**, 406–420. (doi:10.5061/dryad.pg4f4qrpm)

34. Law CJ. 2019 Solitary meat-eaters: solitary, carnivorous carnivorans exhibit the highest degree of sexual size dimorphism. *Sci Rep-uk* **9**, 15344. (doi:10.1038/s41598-019-51943-x)

35. Morris JS, Carrier DR. 2016 Sexual selection on skeletal shape in Carnivora. *Evolution* **70**, 767–780. (doi:10.1111/evo.12904)

36. Hansen TF. 1997 Stabilizing selection and the comparative analysis of adaptation. *Evolution* **51**, 1341–1351. (doi:10.1111/j.1558-5646.1997.tb01457.x)

37. Butler MA, King AA. 2004 Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *Am Nat* **164**, 683–695. (doi:10.1086/426002)

38. Clavel J, Escarguel G, Merceron G. 2015 mv morph: an R package for fitting multivariate evolutionary models to morphometric data. *Methods Ecol Evol* **6**, 1311–1319. (doi:10.1111/2041-210x.12420)

39. Bastide P, Ané C, Robin S, Mariadassou M. 2018 Inference of Adaptive Shifts for
Multivariate Correlated Traits. *Syst. Biol.* **67**, 662–680. (doi:10.1093/sysbio/syy005)

40. Meloro C, Raia P, Piras P, Barbera C, OHiggins P. 2008 The shape of the mandibular corpus
in large fissiped carnivores: allometry, function and phylogeny. *Zool J Linn Soc* **154**, 832–845.
(doi:10.1111/j.1096-3642.2008.00429.x)

41. Figueirido B, Serrano-Alarcón FJ, Slater GJ, Palmqvist P. 2010 Shape at the cross-roads:
homoplasy and history in the evolution of the carnivoran skull towards herbivory. *J. Evol. Biol.*
**23**, 2579–2594. (doi:10.1111/j.1420-9101.2010.02117.x)

42. Figueirido B, Tseng ZJ, Martín-Serra A. 2013 Skull shape evolution in durophagous
carnivorans. *Evolution* **67**, 1975–1993. (doi:10.1111/evo.12059)

43. Prevosti FJ, Turazzini GF, Ercoli MD, Hingst-Zaher E. 2011 Mandible shape in marsupial
and placental carnivorous mammals: a morphological comparative study using geometric
morphometrics. *Zool J Linn Soc* **164**, 836–855. (doi:10.1111/j.1096-3642.2011.00785.x)

44. Morales-García NM, Gill PG, Janis CM, Rayfield EJ. 2021 Jaw shape and mechanical
advantage are indicative of diet in Mesozoic mammals. *Communications Biology* , 1–14.
(doi:10.1038/s42003-021-01757-3)

45. Finarelli JA, Flynn JJ. 2009 Brain-size evolution and sociality in Carnivora. *Proc National*
*Acad Sci* **106**, 9345–9349. (doi:10.1073/pnas.0901780106)

46. Curtis AA, Valkenburgh BV. 2014 Beyond the Sniffer: Frontal Sinuses in Carnivora. *Anat.*
*Rec.* **297**, 2047–2064. (doi:10.1002/ar.23025)

47. Bird DJ, Murphy WJ, Fox-Rosales L, Hamid I, Eagle RA, Valkenburgh BV. 2018 Olfaction
written in bone: cribriform plate size parallels olfactory receptor gene repertoires in Mammalia.
*Proc. Biol. Sci.* **285**, 20180100–9. (doi:10.1098/rspb.2018.0100)

48. Meachen-Samuels J, Valkenburgh BV. 2009 Forelimb indicators of prey-size preference in
the Felidae. *J Morpho* **270**, 729–744. (doi:10.1002/jmor.10712)

49. Michaud M, Veron G, Fabre A. 2020 Phenotypic integration in feliform carnivores:
Covariation patterns and disparity in hypercarnivores versus generalists. *Evolution* **74**, 2681–
2702. (doi:10.1111/evo.14112)

50. Wilson DE, Mittermeier RA. 2014 *Handbook of the Mammals of the World: Volume 4. Sea*
*Mammals*. Lynx Edicions.

51. Polly PD. 2007 Limbs in mammalian evolution. In *Fins into Limbs Evolution, Development,*
*and Transformation* (ed BK Hall), pp. 245–268. Fins into Limbs Evolution, Development, and
Transformation.

52. Jones KE, Angielczyk KD, Polly PD, Head JJ, Fernandez V, Lungmus JK, Tulga S, Pierce
SE. 2018 Fossils reveal the complex evolutionary history of the mammalian regionalized spine.
*Science* **361**, 1249–1252. (doi:10.1126/science.aar3126)

53. Jones KE, Angielczyk KD, Pierce SE. 2019 Stepwise shifts underlie evolutionary trends in
morphological complexity of the mammalian vertebral column. *Nat Commun* **10**, 5071.
(doi:10.1038/s41467-019-13026-3)

54. Simpson GG. 1944 *Tempo and Mode in Evolution*. Columbia University Press.

55. Simpson GG. 1955 *Major features of evolution*. Columbia University Press.

56. Humphreys AM, Barraclough TG. 2014 The evolutionary reality of higher taxa in mammals.
*Proc Royal Soc B Biological Sci* **281**, 20132750. (doi:10.1098/rspb.2013.2750)

57. Law CJ. 2019 Evolutionary shifts in extant mustelid (Mustelidae: Carnivora) cranial shape,
body size and body shape coincide with the Mid-Miocene Climate Transition. *Biol. Lett.* **15**,
20190155–6. (doi:10.1098/rsbl.2019.0155)

397