Uncovering the mosaic evolution of carnivoran skeletal systems

Chris J. Law[1,2], Leslea J. Hlusko[3], Z. Jack Tseng[4]

[1]Department of Integrative Biology, University of Texas

[2]Burke Museum and Department of Biology, University of Washington

[3]National Research Center on Human Evolution (CENIEH), Burgos, Spain

[4]Department of Integrative Biology, University of California Berkeley

Abstract

The diversity of vertebrate skeletons is often attributed to adaptations to distinct ecological

factors such as diet, locomotion, and sensory environment. Although the adaptive evolution of

cranial, appendicular, and vertebral skeletal systems is well studied in vertebrates,

comprehensive investigations of all skeletal components simultaneously are rarely performed.

Consequently, we know little of how modes of evolution differ among skeletal components.

Here, we tested if ecological and phylogenetic effects led to distinct modes of evolution among

the cranial, appendicular, and vertebral regions in extant carnivoran skeletons. Using multivariate

evolutionary models, we found mosaic evolution in which only the mandible, hindlimb, and

posterior region of the vertebral column showed evidence of adaptation towards ecological

regimes whereas the remaining skeletal components reflect clade-specific evolutionary shifts.

We hypothesize that the decoupled evolution of individual skeletal components may have led to

the origination of distinct adaptive zones and morphologies among extant carnivoran families

that reflect phylogenetic hierarchies. Overall, our work highlights the importance of examining

multiple skeletal components simultaneously in ecomorphological analyses. Ongoing work

integrating the fossil and paleoenvironmental record will further clarify deep-time drivers that

govern carnivoran diversity we see today and reveal the complexity of evolutionary processes in

multicomponent systems.

Key words: adaptive landscape; Carnivora; ecomorphology; macroevolution; Ornstein-

Uhlenbeck modeling; phylogenetic comparative methods

Introduction

The diversity of animal forms is one of the most salient patterns across the tree of life. In

mammals, morphological innovations in the cranial, appendicular, and axial skeletal systems

facilitate the incredible diversity found today, ranging from bats with winged forelimbs to the

biggest animals to have ever lived on earth. Many researchers have examined how variation in

the skull [1–6], limbs [7–10], or vertebrae [11–15] serve as crucial adaptations to their evolution.

These skeletal systems are traditionally examined independently and are rarely investigated

simultaneously even though these anatomical regions comprise a single, functionally integrated

system to support movement, sensation, and other life functions. When considered wholistically,

the observed variation across the different components of organismal anatomy is generally

explained by multitudinous factors, some that are potentially incongruous [16–18]. While this

evolutionary push-and-pull between anatomical regions may characterize the process of

evolution, the hypothesis can only be tested when the different skeletal components are explored

simultaneously rather than piecemeal. Simultaneous investigation of integrated components is

critical to our understanding of the role of developmental and/or functional integration in

canalizing macroevolutionary trajectories [19–21]. Here, we use carnivorans to investigate how

ecological and phylogenetic factors correspond to evolutionary changes in the cranial,

appendicular, and axial skeletal systems. Carnivorans (bears, cats, dogs, seals, and their relatives)

are a productive model system to examine skeletal evolution because of their high species

richness and vast distribution across most biomes in all continents and oceans, along with broad

ecological diversity in locomotor traits and feeding adaptations.

Components of carnivoran skeletal systems are well studied individually. In the skull,

craniomandibular diversity is influenced by several ecological factors and phylogeny [22–26].

The skull exhibits decoupled evolutionary modes: cranial shape follows clade-specific

evolutionary shifts, whereas mandibular shape evolution is linked to broad dietary regimes

[6,27]. In the appendicular skeleton, ecomorphological divergence exists between the hindlimbs,

which are adapted primarily for locomotion, and the forelimb, which are adapted for multiple

functions ranging from running to grappling prey to manipulating objects [28–31]. Additionally,

more recent work using phylogenetic comparative methods found that scaling and phylogeny

exhibit stronger effects on limb evolution than do ecological parameters [32–34]. In contrast to

craniomandibular and appendicular ecomorphology, research on the axial skeleton is in nascent

stages. Initial research indicates that distinct regions of the vertebral column are under different

evolutionary pressures. The anterior region exhibits low disparity due to phylogenetic constraints

or ecological conservatism, whereas the posterior region exhibits higher disparity that may be

due to adaptations to various locomotor ecologies [12,35]. In contrast to these morphologically-

localized studies, analyses of the evolution of whole-body traits like body mass, skeletal size,

and body shape often follow a Brownian motion model or clade-based shift model rather than

being associated with ecological regimes [26,36,37].

Compared to skeletal system-specific findings, simultaneous investigation of skulls,

limbs, vertebrae, and overall body plan are rarely conducted, likely because of the enormous

amount of data that would need to be collected and the complexity of the multivariate analyses

required. However, a more comprehensive approach to quantifying skeletal evolution is essential

to elucidate its complexity more fully. The search for system-level trends and variations is

further obscured by the disparate methods employed to test the effects of ecology and phylogeny

on different skeletal systems by different researchers. In this study, we address both issues in our

investigation of the mosaic evolution of carnivoran skeletons by creating a new phenomic dataset

that encompasses all major components of the skeletal system and using a unified set of

multivariate evolutionary models to test the ecological and phylogenetic effects influencing the

modes of evolution of these skeletal components.

Methods

Skeletal and ecological traits

We collected 103 linear measurements to capture the skeletal morphology of 119

carnivoran species (208 osteological specimens; Fig. S1; Table S1). This dataset includes seven

cranial traits, seven mandibular traits, 13 forelimb traits, 13 hindlimb traits, and seven traits in

third cervical, fifth cervical, first thoracic, middle thoracic, diaphragmatic thoracic, last thoracic,

first lumbar, middle lumbar, and last lumbar vertebrae. Because carnivorans exhibit differing

degrees of sexual dimorphism [38,39], we use only male specimens. To remove size effects, we

calculated log shape ratios by dividing each skeletal trait by the geometric mean of all 103 traits

[40,41]. We then used principal component analyses (PCAs) to reduce the dimension of each

skeletal component (i.e., cranium, mandible, forelimb, hindlimb, and each of the nine vertebrae)

and retained a number of PC axes that corresponded to >90% of the explained variance. We also

conducted a PCA on the entire dataset as our proxy of the whole-skeleton phenome and retained

the first six PC axes (~75% of explained variance) for subsequent analyses. We classified the

119 carnivoran species into distinct locomotor modes, hunting behaviors, and dietary regimes

following [37].

Phylogenetic comparative methods

We tested whether each skeletal component evolved as adaptation to specific ecological

regimes or exhibited clade-specific evolutionary shifts by fitting multivariate evolutionary

models on the retained PC axes of each skeletal component [42–44]. For the adaptive ecological

models, we fit three multivariate multi-optima Ornstein-Uhlenbeck models (i.e., mvOUM$_{diet}$,

mvOUM$_{hunting,}$ and mvOUM$_{locomotion}$) to test if dietary, hunting behavioral, or locomotor regimes

influenced the evolution of each skeletal component using mvMORPH [44]. The models were fit

across 500 stochastically mapped trees to account for uncertainty in phylogenetic topology and

ancestral character states (see electronic supplementary materials). We also calculated the

phylogenetic half-lives of the best supported adaptive ecological model [42]. A short

phylogenetic half-life relative to the age of Carnivora (48.2 myr) would suggest that skeletal

traits are strongly pulled toward distinct ecological optima across the adaptive landscape. For the

clade-based model, we fit a multi-optima OU model (mvOUM$_{phyloEM}$) without *a priori* ecological

regimes with PhylogeneticEM [45]. We also fit a single-rate multivariate Brownian motion

model (mvBM1) and a single-optimum OU model (mvOU1). We assessed the relative support of

models using small sample-corrected Akaike weights (AICcW). Lastly, we assessed the

covariation among skeletal components using partial least squares with geomorph [46].

Preliminary results revealed that phenotypic differences between pinnipeds (i.e., seals

and sea lions) and terrestrial carnivorans are often the greatest source of variation for most

skeletal components. These results are unsurprising considering pinnipeds exhibit derived

morphologies that enable them to be fully aquatic. Therefore, we repeated our analyses using a

reduced dataset with no pinnipeds. Results of the full dataset with pinnipeds are presented in the

electronic supplementary material.

Results and Discussion

We found mosaic evolution of the carnivoran skeleton in which ecology and phylogeny

have differing influences on the evolutionary mode of the various skeletal components.

Consistent with [6,27], the cranium and mandible exhibited decoupled evolutionary modes. In

the cranium, the clade-specific shift model exhibited overwhelmingly greater support

($mvOUM_{phyloEM}$; AICcW>0.99) compared to adaptive ecological models (Fig. 1; Table S2). We

found eight evolutionary shifts in cranial morphology that correspond to carnivoran clades (Fig.

2A). In contrast, the adaptive dietary model was the best supported model ($mvOUM_{diet}$;

AICcW=0.96) for the mandible with a short phylogenetic half-life of 2.52 myr (Fig. 1; Fig. S2B;

Table S2; see Supplementary Results for optima distribution in phylomorphospace). These

results are congruent with findings revealing that mandibular shape is evolutionarily labile with

respect to dietary evolution whereas cranial shape is partitioned among families rather than

among dietary groups [6]. Despite their covariation (r = 0.73; Table S3), decoupled evolutionary

modes between the cranium and mandible may be explained by their functions. Diet is often

found to have had a strong influence on mandibular evolution because of its direct role in feeding

[3,47–51]. In contrast, the cranium has multiple sensory functions in addition to feeding that

influence its evolution [52–54], and therefore, the signal from dietary adaptations in its

morphology may be obscured.

The appendicular system exhibited decoupled evolutionary modes between forelimbs and

hindlimbs. The forelimb was best supported by the mvOUM$_{phyloEM}$ model (AICcW>0.99; Fig. 1;

Table S2). Seven shifts in forelimb evolution occur primarily along familial branches (Fig. 2B),

indicating that the complexity and variation of carnivoran forelimb morphology cannot be

captured effectively by dietary, hunting behavioral, or locomotor categories. Instead, these shifts

suggest that clade-specific adaptations enabled the diversity of forelimb skeletons for tasks such

as grappling or manipulating prey, swimming, or digging [28–31,34,55,56]. For example, most

felids use their prehensile forelimbs to ambush and subdue prey, most canids and hyaenids

pounce and pursue prey, and some mustelids use their powerful forelimbs to dig out prey while

other more derived mustelids (i.e., weasels) pursue prey in tight crevices and burrows [57]. In

contrast, the hindlimb was best supported by the mvOUM$_{locomotion}$ model (AICcW=0.83) in the

hindlimb with a short phylogenetic half-life of 5.05 myr (Fig. 1; Table S2), supporting

hypotheses that the hindlimb is adapted primarily for locomotion as typically found in

quadrupedal mammals [58]. Although the forelimb and hindlimb covaries (r = 0.87; Table S3),

previous work found that this integration is weaker than expected in carnivorans that do not

specialize in cursoriality [34]. This work together supports the hypothesis of functional

divergence between the forelimbs and hindlimbs of carnivorans.

The axial skeleton exhibits distinct evolutionary modes between the anterior and

posterior regions of the vertebral column: cervical and most thoracic vertebrae tended to be best

supported by clade-specific shift or single-peak OU models, whereas the last thoracic and all

lumbar vertebrae were best supported by mvOUM$_{hunting}$ or mvOUM$_{locomotion}$ models (Fig. 1; Fig.

2C–F; Table S2). Our findings strengthen the coalescing hypothesis that anterior vertebrae

exhibit lower disparity, higher evolutionary constraints, and more subtle adaptations to

locomotion whereas posterior vertebrae exhibit the opposite patterns in carnivorans [35] and

broadly across mammals [13]. We posit that high evolutionary constraints of the anterior

vertebrae are associated with clade-specific shifts in the cervical and most thoracic vertebrae.

Importantly, subtle adaptations in these anterior vertebrae could be masked by many-to-one or

one-to-many mappings, making it difficult to uncover the form-function associations with

evolutionary models [59]. In contrast, relaxed evolutionary constraints of the posterior vertebrae

facilitate the evolution of disparate lumbar vertebrae across the entire carnivoran order. These

disparate vertebrae adapt to diverse locomotor modes or hunting behaviors based on the mobility

of the posterior backbone and irrespective of clade origins. The short phylogenetic half-lives

(1.47–5.12 myr) further suggests strong pulls towards these different adaptive optima. More

broadly, this increased mobility of the lumbar region over evolutionary time is hypothesized to

be an innovation characterizing crown mammals [13,60,61]. Correspondingly, the posterior

vertebrae are tightly integrated ($r = 0.84–0.96$; Table S3).

Lastly, we found that the clade-specific shift model ($mvOUM_{phyloEM}$; AICcW > 0.99) best

described the overall skeletal phenome (Table S2), a pattern that is consistent with previous

investigations of whole-body proxies such as body size and body shape [26,36,37]. The

mammalian body plan is comprised of cranial, axial, and appendicular components; therefore, its

multidimensionality transcends one-to-one mapping relationships between morphology and

ecological function. Instead, individual skeletal components within distinct body plans can adapt

to specific ecological factors independently from each other, enabling species with distinct body

plans to exhibit similar ecological or functional regimes and vice versa.

Overall, we elucidate the mosaic evolution of the carnivoran skeleton, finding that

different skeletal components exhibit distinct modes of evolution. Our results suggest that

different methodologies and taxonomic samples do not necessarily explain previously reported

region-specific macroevolutionary patterns; rather, complexity in explanatory factors of skeletal

diversity is a key feature of Carnivora. The ability of individual skeletal components to adapt to

specific ecological factors independently from each other may have contributed to the clade's

hierarchical [62,63] evolution. As previously hypothesized [26,36], the restriction of carnassial

shear to the P4/m1 pair may have been the key innovation that facilitated the initial carnivoran

diversification early in the clade's evolutionary history. Subsequent evolution led to the

continual partitioning between clades, resulting in the origination of extant carnivoran families as

discrete phylogenetic clusters that occupy different adaptive zones [64] with distinct

morphologies including body size and shape [37,65] and various components of the skeleton

([6]; Fig. 2). Within-clade variation then arises to reflect resource partitioning among

ecologically similar taxa, leading to adaptations in morphologies such as the mandible, hindlimb,

and posterior region of the vertebral column (Fig. 1). These traits were strongly pulled toward

distinct ecological peaks across the adaptive landscape as revealed by their short phylogenetic

half-lives (1.47–5.12 myr) relative to the clade's age (48.2 myr).

Our research statistically revealed the mosaic evolution of carnivoran skeletons. These

distinct evolutionary modes demonstrate the importance of examining multiple skeletal

components in ecomorphological analyses. Nevertheless, key questions remain: What spurred

the evolutionary transitions towards the evolutionary shifts or adaptations of the various skeletal

components? When in the 55 million years of carnivoran evolutionary history did these

evolutionary events occur? And what developmental and genetic phenomena underlie the

evolutionary dissociation of various skeletal elements? Ongoing work integrating the fossil and

paleoenvironmental record will further elucidate the carnivoran diversity we see today and reveal

the complexity of evolutionary processes in multicomponent systems.

Acknowledgements

We are grateful to the staff and collections at the American Museum of Natural History,

California Academy of Sciences, Field Museum of Natural History, Natural History Museum of

Los Angeles County, Museum of Vertebrate Zoology, Natural History Museum London, San

Diego Natural History Museum, Texas Vertebrate Paleontology Collection, National Museum of

Natural History, and Burke Museum of Natural History and Culture.

Funding

Funding was supported by the National Science Foundation (DBI–2128146), a University of

Texas Early Career Provost Fellowship and Stengl-Wyer Endowment Grant (SWG-22-02) to

CJL, and the European Research Council (Tied2Teeth, grant agreement n° 101054659) to LJH.

Fig. 1. Diagram of the skeletal components and their best-fitting evolutionary model on *Lontra*

canadensis. AICcW are in parentheses. See Table S2 for full AICc table. diaT = diaphragmatic

thoracic vertebrae

Fig. 2. Clade-specific evolutionary shifts in skeletal components across terrestrial carnivorans

identified by PhylogeneticEM. Shifts are represented as pink circles, and branches on the

phylogenies are colored according to each regime.

References

1. Dumont ER, Davalos LM, Goldberg A, Santana SE, Rex K, Voigt CC. 2012 Morphological innovation, diversification and invasion of a new adaptive zone. *Proceedings of the Royal Society B: Biological Sciences* **279**, 1797–1805. (doi:10.1098/rspb.2011.2005)

2. Zelditch ML, Ye J, Mitchell JS, Swiderski DL. 2017 Rare ecomorphological convergence on a complex adaptive landscape: Body size and diet mediate evolution of jaw shape in squirrels (Sciuridae). *Evolution* **71**, 633–649. (doi:10.1111/evo.13168)

3. Grossnickle DM. 2020 Feeding ecology has a stronger evolutionary influence on functional morphology than on body mass in mammals. *Evolution* **74**, 610–628. (doi:10.1111/evo.13929)

4. Arbour JH, Curtis AA, Santana SE. 2019 Signatures of echolocation and dietary ecology in the adaptive evolution of skull shape in bats. *Nat Commun* **10**, 2036. (doi:10.1038/s41467-019-09951-y)

5. Goswami A *et al.* 2022 Attenuated evolution of mammals through the Cenozoic. *Science* **378**, 377–383. (doi:10.1126/science.abm7525)

6. Law CJ, Blackwell EA, Curtis AA, Dickinson E, Hartstone Rose A, Santana SE. 2022 Decoupled evolution of the cranium and mandible in carnivoran mammals. *Evolution* **76**, 2959–2974. (doi:10.1111/evo.14578)

7. Patel BA. 2010 Functional morphology of cercopithecoid primate metacarpals. *J. Hum. Evol.* **58**, 320–337. (doi:10.1016/j.jhevol.2010.01.001)

8. Kilbourne BM. 2017 Selective regimes and functional anatomy in the mustelid forelimb: Diversification toward specializations for climbing, digging, and swimming. *Ecol Evol* **66**, 2369–12. (doi:10.1002/ece3.3407)

9. Grossnickle DM, Chen M, Wauer JGA, Pevsner SK, Weaver LN, Meng QJ, Liu D, Zhang YG, Luo ZX. 2020 Incomplete convergence of gliding mammal skeletons. *Evolution* **74**, 2662–2680. (doi:10.1111/evo.14094)

10. Rickman J, Burtner AE, Linden TJ, Santana SE, Law CJ. 2023 Size And Locomotor Ecology Have Differing Effects on the External and Internal Morphologies of Squirrel (Rodentia: Sciuridae) Limb Bones. *Integr. Org. Biol.* **5**, obad017. (doi:10.1093/iob/obad017)

11. Pierce SE, Clack JA, Hutchinson JR. 2011 Comparative axial morphology in pinnipeds and its correlation with aquatic locomotory behaviour. *J Anat* **219**, 502–514. (doi:10.1111/j.1469-7580.2011.01406.x)

12. Randau M, Cuff AR, Hutchinson JR, Pierce SE, Goswami A. 2017 Regional differentiation
of felid vertebral column evolution: a study of 3D shape trajectories. *Org Divers Evol* **17**, 1–15.
(doi:10.1007/s13127-016-0304-4)

13. Jones KE, Benitez L, Angielczyk KD, Pierce SE. 2018 Adaptation and constraint in the
evolution of the mammalian backbone. *Bmc Evol Biol* **18**, 172. (doi:10.1186/s12862-018-1282-
2)

14. Martín-Serra A, Pérez-Ramos A, Pastor FJ, Velasco D, Figueirido B. 2021 Phenotypic
integration in the carnivoran backbone and the evolution of functional differentiation in
metameric structures. *Evolution Letters* , 1–14. (doi:10.1002/evl3.224)

15. Linden TJ, Burtner AE, Rickman J, McFeely A, Santana SE, Law CJ. 2023 Scaling patterns
of body plans differ among squirrel ecotypes. *PeerJ* **11**, e14800. (doi:10.7717/peerj.14800)

16. Friedman ST, Price SA, Wainwright PC. 2021 The Effect of Locomotion Mode on Body
Shape Evolution in Teleost Fishes. *Integr Org Biology* **3**, obab016. (doi:10.1093/iob/obab016)

17. Law CJ. 2021 Evolutionary and morphological patterns underlying carnivoran body shape
diversity. *Evolution* **75**, 365–375. (doi:10.1111/evo.14143)

18. Bergmann PJ, Irschick DJ. 2009 Alternate pathways of body shape evolution translate into
common patterns of locomotor evolution in two clades of lizards. *Evolution* **64**, 1569–1582.
(doi:10.1111/j.1558-5646.2011.01491.x)

19. Goswami A, Randau M, Polly PD, Weisbecker V, Bennett CV, Hautier L, Sánchez-Villagra
MR. 2016 Do Developmental Constraints and High Integration Limit the Evolution of the
Marsupial Oral Apparatus? *Integr. Comp. Biol.* **56**, 404–415. (doi:10.1093/icb/icw039)

20. Goswami A, Weisbecker V, Sánchez␣Villagra MR. 2009 Developmental modularity and the
marsupial–placental dichotomy. *J. Exp. Zoo␣l. Part B: Mol. Dev. Evol.* **312B**, 186–195.
(doi:10.1002/jez.b.21283)

21. Goswami A, Smaers JB, Soligo C, Polly PD. 2014 The macroevolutionary consequences of
phenotypic integration: from development to deep time. *Philos. Trans. R. Soc. B: Biol. Sci.* **369**,
20130254. (doi:10.1098/rstb.2013.0254)

22. Radinsky LB. 1981 Evolution of skull shape in Carnivores .1. Representative modern
Carnivores. *Biol J Linn Soc* **15**, 369–388. (doi:10.1111/j.1095-8312.1981.tb00770.x)

23. Valkenburgh BV. 2007 Deja vu: the evolution of feeding morphologies in the Carnivora.
Amer. Zool. **47**, 147–163. (doi:10.1093/icb/icm016)

24. Law CJ, Duran E, Hung N, Richards E, Santillan I, Mehta RS. 2018 Effects of diet on cranial
morphology and biting ability in musteloid mammals. *J. Evol. Biol.* **31**, 1918–1931.
(doi:10.1111/jeb.13385)

25. Tseng ZJ, Flynn JJ. 2018 Structure-function covariation with nonfeeding ecological variables
influences evolution of feeding specialization in Carnivora. *Sci Adv* **4**, eaao5441.
(doi:10.1126/sciadv.aao5441)

26. Slater GJ, Friscia AR. 2019 Hierarchy in adaptive radiation: A case study using the
Carnivora (Mammalia). *Evolution* **73**, 524–539. (doi:10.1111/evo.13689)

27. Figueirido B, MacLeod N, Krieger J, Renzi MD, Pérez-Claros JA, Palmqvist P. 2011
Constraint and adaptation in the evolution of carnivoran skull shape. *Paleobiology* **37**, 490–518.
(doi:10.1666/09062.1)

28. Valkenburgh BV. 1985 Locomotor diversity within past and present guilds of large predatory
mammals. *Paleobiology* **11**, 406–428. (doi:10.1017/s0094837300011702)

29. Valkenburgh BV. 1987 Skeletal indicators of locomotor behavior in living and extinct
carnivores. *J Vertebr Paleontol* **7**, 162–182. (doi:10.1080/02724634.1987.10011651)

30. Iwaniuk AN, Pellis SM, Whishaw IQ. 1999 The relationship between forelimb morphology
and behaviour in North American carnivores (Carnivora). *Can J Zool* **77**, 1064–1074.
(doi:10.1139/z99-082)

31. Samuels JX, Meachen JA, Sakai SA. 2013 Postcranial morphology and the locomotor habits
of living and extinct carnivorans. *J Morpho* **274**, 121–146. (doi:10.1002/jmor.20077)

32. Martín-Serra A, Figueirido B, Palmqvist P. 2014 A Three-Dimensional Analysis of
Morphological Evolution and Locomotor Performance of the Carnivoran Forelimb. *Plos One* **9**,
e85574. (doi:10.1371/journal.pone.0085574)

33. Martín-Serra A, Figueirido B, Palmqvist P. 2014 A three-dimensional analysis of the
morphological evolution and locomotor behaviour of the carnivoran hind limb. *BMC Evol Biol*
14, 1–13. (doi:10.1186/1471-2148-14-129)

34. Martín-Serra A, Figueirido B, Pérez-Claros JA, Palmqvist P. 2015 Patterns of morphological
integration in the appendicular skeleton of mammalian carnivores. *Evolution* **69**, 321–340.
(doi:10.1111/evo.12566)

35. Figueirido B, Martín-Serra A, Pérez-Ramos A, Velasco D, Pastor FJ, Benson RJ. 2021 Serial
disparity in the carnivoran backbone unveils a complex adaptive role in metameric evolution.
Communications Biology , 1–15. (doi:10.1038/s42003-021-02346-0)

36. Slater GJ. 2022 Topographically distinct adaptive landscapes for teeth, skeletons, and size
explain the adaptive radiation of Carnivora (Mammalia). *Evolution* (doi:10.1111/evo.14577)

37. Law CJ. 2021 Ecological drivers of carnivoran body shape evolution. *Am Nat* **198**, 406–420.
(doi:10.5061/dryad.pg4f4qrpm)

38. Law CJ. 2019 Solitary meat-eaters: solitary, carnivorous carnivorans exhibit the highest
degree of sexual size dimorphism. *Sci Rep-uk* **9**, 15344. (doi:10.1038/s41598-019-51943-x)

39. Morris JS, Carrier DR. 2016 Sexual selection on skeletal shape in Carnivora. *Evolution* **70**,
767–780. (doi:10.1111/evo.12904)

40. Claude J. 2013 Log-shape ratios, Procrustes superimposition, elliptic Fourier analysis: three
worked examples in R. *Hystrix, the Italian Journal of Mammalogy* **24**, 94–102.

41. Mosimann JE. 1970 Size Allometry: Size and Shape Variables with Characterizations of the
Lognormal and Generalized Gamma Distributions. *J. Am. Stat. Assoc.* **65**, 930–945.
(doi:10.1080/01621459.1970.10481136)

42. Hansen TF. 1997 Stabilizing selection and the comparative analysis of adaptation. *Evolution*
51, 1341–1351. (doi:10.1111/j.1558-5646.1997.tb01457.x)

43. Butler MA, King AA. 2004 Phylogenetic comparative analysis: a modeling approach for
adaptive evolution. *Am Nat* **164**, 683–695. (doi:10.1086/426002)

44. Clavel J, Escarguel G, Merceron G. 2015 mv morph: an R package for fitting multivariate
evolutionary models to morphometric data. *Methods Ecol Evol* **6**, 1311–1319.
(doi:10.1111/2041-210x.12420)

45. Bastide P, Ané C, Robin S, Mariadassou M. 2018 Inference of Adaptive Shifts for
Multivariate Correlated Traits. *Syst. Biol.* **67**, 662–680. (doi:10.1093/sysbio/syy005)

46. Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. 2021 geomorph v4.0 and gmShiny:
Enhanced analytics and a new graphical interface for a comprehensive morphometric experience.
Methods Ecol Evol **12**, 2355–2363. (doi:10.1111/2041-210x.13723)

47. Meloro C, Raia P, Piras P, Barbera C, OHiggins P. 2008 The shape of the mandibular corpus
in large fissiped carnivores: allometry, function and phylogeny. *Zool J Linn Soc* **154**, 832–845.
(doi:10.1111/j.1096-3642.2008.00429.x)

48. Figueirido B, Serrano-Alarcón FJ, Slater GJ, Palmqvist P. 2010 Shape at the cross☐roads:
homoplasy and history in the evolution of the carnivoran skull towards herbivory. *J. Evol. Biol.*
23, 2579–2594. (doi:10.1111/j.1420-9101.2010.02117.x)

49. Figueirido B, Tseng ZJ, Martín-Serra A. 2013 Skull shape evolution in durophagous
carnivorans. *Evolution* **67**, 1975–1993. (doi:10.1111/evo.12059)

50. Prevosti FJ, Turazzini GF, Ercoli MD, Hingst-Zaher E. 2011 Mandible shape in marsupial
and placental carnivorous mammals: a morphological comparative study using geometric
morphometrics. *Zool J Linn Soc* **164**, 836–855. (doi:10.1111/j.1096-3642.2011.00785.x)

51. Morales-García NM, Gill PG, Janis CM, Rayfield EJ. 2021 Jaw shape and mechanical advantage are indicative of diet in Mesozoic mammals. *Communications Biology* , 1–14. (doi:10.1038/s42003-021-01757-3)

52. Finarelli JA, Flynn JJ. 2009 Brain-size evolution and sociality in Carnivora. *Proc National Acad Sci* **106**, 9345–9349. (doi:10.1073/pnas.0901780106)

53. Curtis AA, Valkenburgh BV. 2014 Beyond the Sniffer: Frontal Sinuses in Carnivora. *Anat. Rec.* **297**, 2047–2064. (doi:10.1002/ar.23025)

54. Bird DJ, Murphy WJ, Fox-Rosales L, Hamid I, Eagle RA, Valkenburgh BV. 2018 Olfaction written in bone: cribriform plate size parallels olfactory receptor gene repertoires in Mammalia. *Proc. Biol. Sci.* **285**, 20180100–9. (doi:10.1098/rspb.2018.0100)

55. Meachen-Samuels J, Valkenburgh BV. 2009 Forelimb indicators of prey-size preference in the Felidae. *J Morpho* **270**, 729–744. (doi:10.1002/jmor.10712)

56. Michaud M, Veron G, Fabre A. 2020 Phenotypic integration in feliform carnivores: Covariation patterns and disparity in hypercarnivores versus generalists. *Evolution* **74**, 2681–2702. (doi:10.1111/evo.14112)

57. Wilson DE, Mittermeier RA. 2014 *Handbook of the Mammals of the World: Volume 4. Sea Mammals*. Lynx Edicions. See http://books.google.com/books/content?id=_8tgwgEACAAJ&;printsec=frontcover&img=1&zoom=1&source=gbs_api.

58. Polly PD. 2007 Limbs in mammalian evolution. In *Fins into Limbs Evolution, Development, and Transformation* (ed BK Hall), pp. 245–268. Fins into Limbs Evolution, Development, and Transformation.

59. Simon MN, Moen DS. 2023 Bridging Performance and Adaptive Landscapes to Understand Long-Term Functional Evolution. *Physiol. Biochem. Zoo□l.* **96**, 304–320. (doi:10.1086/725416)

60. Jones KE, Angielczyk KD, Polly PD, Head JJ, Fernandez V, Lungmus JK, Tulga S, Pierce SE. 2018 Fossils reveal the complex evolutionary history of the mammalian regionalized spine. *Science* **361**, 1249–1252. (doi:10.1126/science.aar3126)

61. Jones KE, Angielczyk KD, Pierce SE. 2019 Stepwise shifts underlie evolutionary trends in morphological complexity of the mammalian vertebral column. *Nat Commun* **10**, 5071. (doi:10.1038/s41467-019-13026-3)

62. Simpson GG. 1944 *Tempo and Mode in Evolution*. Columbia University Press. See http://books.google.com/books?id=f5qucNebo-kC&;printsec=frontcover&dq=intitle:Tempo+and+mode+in+evolution&hl=&cd=1&source=gbs_api.

63. Simpson GG. 1955 *Major features of evolution*. Columbia University Press. See
http://krishikosh.egranth.ac.in/handle/1/2034793.

64. Humphreys AM, Barraclough TG. 2014 The evolutionary reality of higher taxa in mammals.
Proc Royal Soc B Biological Sci **281**, 20132750. (doi:10.1098/rspb.2013.2750)

65. Law CJ. 2019 Evolutionary shifts in extant mustelid (Mustelidae: Carnivora) cranial shape,
body size and body shape coincide with the Mid-Miocene Climate Transition. *Biol. Lett.* **15**,
20190155–6. (doi:10.1098/rsbl.2019.0155)

406