Serial colonization shapes genetic variation and defines conservation units in Asian
elephants

Anubhab Khan*[1,2,&], Maitreya Sil*[1,3], Tarsh Thekaekara[1,4], Kritika M. Garg[5,6], Ishani
Sinha[1], Rupsy Khurana[7], Raman Sukumar[7,&], Uma Ramakrishnan[1,&]

[1] National Centre for Biological Sciences, TIFR, GKVK Campus, Bangalore, India
[2] School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow,
Glasgow, UK
[3] National Institute of Science Education and Research, Bhubaneshwar, India
[4] The Shola Trust, Gudalur, India
[5] Centre for Interdisciplinary Archaeological Research, Ashoka University, Sonepat,
India
6 Department of Biology, Ashoka University, Sonepat, India
[7] Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India
*These authors contributed equally
[&] Corresponding authors: anubhabkhan@gmail.com, rsuku@iisc.ac.in,
uramakri@ncbs.res.in

Keywords: Genomics, genetic load, extinction, megaherbivores, South Asia

Abstract

Asian elephants (*Elephas maximus*) are the largest extant terrestrial megaherbivores native to Asia, with 60% of their wild population found in India. Despite ecological and cultural importance, their population genetic structure and diversity, demographic history, and ensuing implications for management/conservation remain understudied. We analysed 34 whole genomes (between 11X - 32X) from most known elephant landscapes in India and identified five management/conservation units corresponding to elephants in Northern (Northwestern/Northeastern) India, Central India and three in Southern India. Our genetic data reveal signatures of serial colonisation, and a dilution of genetic diversity from north to south of India. The Northern populations diverged from other populations more than 70,000 years ago, have higher genetic diversity, and low inbreeding/high effective size (Pi = 0.0016±0.0001; $F_{ROH> 1MB}$= 0.09±0.03). Two of three populations in Southern India (South of Palghat Gap: SPG, and South of Shencottah Gap:SSG) have low diversity and are inbred, with very low effective population sizes compared to current census sizes (Pi = 0.0014±0.00009 and 0.0015±0.0001; $F_{ROH> 1MB}$= 0.25±0.09 and 0.17±0.02). Analyses of genetic load reveals purging of potentially high-effect insertion/deletion (indel) deleterious alleles in the Southern populations and potential dilution of all deleterious alleles from north to south in India. However, despite dilution and purging for the damaging mutation load in Southern India, the load that remains is homozygous. High homozygosity of deleterious alleles, coupled with low neutral genetic diversity make these populations (SPG and SSG) high priority for conservation attention. Most surprisingly, our study suggests that patterns of genetic diversity and genetic load can correspond to geographic signatures of serial founding events, even in large, highly mobile, endangered mammals.

Introduction

Megaherbivores are ecological engineers responsible for maintaining several ecosystem functions (Owen-Smith 1987, Waldram et al. 2008, Coverdale et al. 2016, Berzaghi et al. 2023). They are uniquely responsible for nutrient distribution, seed dispersal and germination, and vegetation/habitat modification (Le Roux et al. 2018, Harich et al. 2016, Sekar et al. 2017). Megaherbivores have been especially impacted negatively in the Anthropocene across their range (Enquist et al. 2020). In Asia, megaherbivores have lost 56% (for gaur, *Bos gaurus*) to near 100% (for Javan rhino, *Rhinoceros sondaicus*) of their historic range (Mahmood et al. 2021). Conservation genomics suggests that they are more inbred, and lack genetic diversity compared to their small-bodied counterparts (Brüniche-Olsen et al. 2018). While megaherbivore genomes are being extensively studied in African landscapes, such investigations for Asian counterparts have been lagging, leading to challenges in identification of potential genetic threats relevant to their conservation and management.

Asian elephants (*Elephas maximus*) are charismatic megaherbivores distributed across South and Southeast Asia and are culturally important across the globe (Sukumar 2011). They are found in a variety of natural ecosystems from tropical evergreen forests through deciduous forest to grasslands at various elevations. India harbours at least 60% of the population of wild Asian elephants (Sukumar 2011, Menon and Tiwari 2019). Increase in human footprint and land use change over the past two centuries has impacted elephants significantly, resulting in population isolation even at regional scales. Today, Asian elephant habitats are fragmented and interspersed with farmland, human settlement, commercial plantations and linear transport infrastructure throughout their range (Liu et al., 2017; Padalia et al., 2020) leading to extensive and, often, intense human-elephant conflicts (Sukumar, 1989, 2003; Gubbi et al., 2014). We might expect low genetic diversity and high impacts of isolation under such conditions. However, camera trap data (Srinivasaiah et al., 2022; Srinivasaiah et al., 2021) and radiotelemetry studies (Baskaran et al. 1995, Venkataraman et al. 2005, Sukumar 2003, Sukumar et al. 2003) reveal that elephants have annual home ranges of several hundred square kilometres in India, often encompassing dense human habitation. Such movement through human-

dominated areas might offset impacts of fragmentation, and the associated loss of
genetic variation and inbreeding (Parida et al. 2022). Clearly, understanding
elephant phylogeographic history and current population genetic structuring, coupled
with possible effects of recent fragmentation could be useful for elephant
conservation and management.

Earlier phylogeographic studies of Asian elephants using mitochondrial DNA point
toward a tangled history of demographic change, range expansion/contraction, as
well as population admixture possibly mediated by climate change (glacial-
interglacial cycles) during the Pleistocene (Fernando et al. 2000, Fleischer at al.
2001, Vidya et al., 2009). These studies showed the presence of two distinct
mitochondrial DNA clades (alpha and beta) of Asian elephants, which underwent
allopatric separation likely in glacial refugia in Myanmar and Sri Lanka around 1.9
million years ago. Using six (nuclear) microsatellites and a mitochondrial region,
Vidya et al. (2005b) reported four Indian elephant population genetic clusters, one
each in Northwest-Northeastern India and Central India, and two in Southern India
separated by the Palghat Gap in the Western Ghats mountain range, broadly
corresponding to their regional population distributions. However, a more recent
study by De et al. (2021) suggests only three major genetic clusters corresponding to
Northwestern India, Northeastern India, and a combined Southern and Central
Indian population (with the Northern (NW&NE) population appearing to be admixed).
The patterns of genetic diversity also varied across different ecoregions. Vidya et al.
(2005b) showed that the Southern Indian populations harboured lower mitochondrial
genetic diversity (haplotype diversity) compared to other populations, while
according to De et al. (2021), the Northeastern Indian populations showed low
heterozygosity. The only genome-wide study of elephants globally (Palkopoulou et
al., 2018), includes poor range-wide sampling of Asian elephants, and suggests old
divergences (circa 20-30 kya) between populations in Southern India across the
Palghat Gap, as well as between the Northern and the Southern populations. Given
the limited genomic data and analyses within Asian elephants, it is difficult to
understand demographic history, population genetic structure and, consequently,
conservation priorities.

To address these gaps, we used whole genome sequences of wild-caught Asian

elephants from all four regions and most landscapes within these in India,

encompassing known biogeographic barriers, to assess their population structure,

demographic history and genetic variation. Our analyses help infer the predominant

factors that have shaped the observed genetic patterns across the subcontinent.

Further, we use this information to propose population management units (MU) and

speculate on future genetic issues in elephant conservation.

Results and Discussion

Population structure

We used 28 blood samples collected from wild-born captive elephants of known

origin from almost all known elephant landscapes in India (Supplementary Table 1,

Parida et al. 2022) and re-sequenced whole genomes (Figure 1). We merged this

data with six previously published elephant whole genome sequencing data of which

4 were from India. A set of 2,675,655 SNP markers revealed distinct population

structure within elephant landscapes in India. A principal component analysis (PCA),

for understanding the clustering of genetic variation, revealed that populations

separate from the south to north direction along PC1 axis (Figure 2a). Elephants

from Northwestern India and Northeastern India (NW&NE) form a single cluster

(Figures 2a, b and c). Populations from Central India (CI) and NW&NE cluster

together along the PC axes 1 and 2 but resolve into separate clusters along PC axis

3 and in the ADMIXTURE plot (Figures 2a,b,c,d) with the most optimum support for

five population clusters (Northwest-Northeast, Central and three clusters in Southern

India (namely, North of the Palghat Gap (NPG), South of the Palghat Gap (SPG) and

South of Shencottah Gap (SSG)), EvalAdmix, Supplementary Figures 1 and 2).

Northwest and Northeast Indian (NW&NE) populations together, henceforth, are

referred to as the Northern Indian population while the population in the south in the

Western Ghats (NPG, SPG and SSG) are collectively referred to as the Southern

Indian population, unless the subregional populations are being described. The

admixture graph that includes a Bornean elephant (ERR2260499) as an outgroup

with branch lengths qualitatively adjusted to the drift parameter (Figure 2e,

Supplementary Figure 6) suggests a deep divergence between the Northern and the
Southern Indian elephants. Additionally, we observe strong drift parameter values for
the two populations South of Palghat Gap potentially indicating low effective
population sizes. The phylogenetic (Neighbour Joining or NJ) tree supports the
above observed patterns with a minor difference, where the Central population
seems to be nested within the Northern population cluster (Supplementary Figure 5).

Our results support previous conclusions that the Northwest and Northeast Indian
populations of elephants collectively are different from other Indian populations
(Vidya et al. 2005a,b; De et al. 2021) and that the two major northern river systems,
namely, the Ganges and the Brahmaputra, have acted as potential barriers to gene
flow (Figures 1, 2). Previous studies suggested that the Brahmaputra river acts as an
incomplete barrier, with female-led family groups not venturing to cross except
perhaps at the upper reaches, but not a barrier to adult male elephants (Vidya et al.,
2005b). Largely consistent with these studies, we observe one individual (Rami –
female) in the Central Indian samples had admixture with the Northern populations,
while another elephant (Bolanath – male) from the Central Indian population
dispersed northward across the Ganges river before it was rescued recently from a
location close to the Northeastern population (Figure 1, Supplementary Table 1).
Anecdotal information suggests male elephants swim across the Brahmaputra (RS
and AK independent personal observations), suggesting it is not a complete barrier
to dispersal and movement. Additionally, the Northern Indian elephant samples have
ancestries found in all other clusters (Figure 2d), which could be due to incomplete
lineage sorting. While geographically proximate, we find that Central Indian
populations are genetically distinct as suggested previously (Vidya et al. 2005b).
Consistent with movement ecology inferences (Koirala et al. 2016) the elephants in
the Northwestern population (De et al. 2021) are connected to the Northeastern
Indian elephants. This is a large landscape running west to east along the Himalayan
foothills, though elephant habitat connectivity here is fragile (Koirala et al. 2016) or
even completely disrupted at places in recent times (Sukumar 2011, Menon and
Tiwari 2019).

Our data and analyses allow identification of a novel genetic cluster in Southern
India and suggest three genetically differentiated populations regionally. Along the

Western Ghats in the south, certain breaks or passes divide the elephant population,

with the Palghat Gap being the most prominent barrier to elephant dispersal. Further

south, the Shencottah Gap also acts as a previously unknown impediment to

elephant movement (though anecdotal information suggests that elephants moved

across this gap until a few decades ago). Genetic differentiation of SPG and SSG

(also see Supplementary Figures 3 and 4) could be due to founder effects and

inbreeding combined with recent isolation and small population size (Khan et al.

2022). Gene flow between north and south of the Shencottah Gap may have

reduced recently (compared to across the Palghat Gap) largely due to a railway line,

a highway and associated development along these transportation infrastructures.

While nested biogeographic implications of these gaps on population structure and

phylogeography has been highlighted for smaller species (e.g. montane birds, Robin

et al., 2015; bush frogs, Vijaykumar et al., 2017, geckoes, Chaitanya et al., 2019)

and some mammals (e.g. Lion Tailed Macaque, Ram et al., 2015), that they result in

such deeply divergent lineages in a large, highly mobile mammal such as the

elephant is surprising. Alternatively, founding events followed by minimal gene flow

might have resulted in the observed patterns of divergence. Additionally, elephant

numbers and densities South of the Shencottah Gap have always remained small

(on the order of 100 individuals). Such small populations are subject to genetic drift,

which could accentuate the observed patterns. Pairwise F_{ST} supports these

inferences (Supplementary Table 2) and we find no significant gene flow between

the clusters based on F3 statistics (Supplementary Table 3). F3 statistics tests

whether a target population is an admixture of two test populations. If the F3 value is

negative the target population is admixed between the two test populations signifying

geneflow between the two populations. For Indian elephants we do not find a

significant negative F3 value for any combination of three populations

(Supplementary Table 3).

Interestingly, the results obtained from the haplotype network analysis based on the

mitogenome paints a slightly different picture (Supplementary Figure 7). Similar to

the nuclear genome, the Northwestern population was embedded within the

Northeastern population. However, the Central Indian population showed closer

affinity to the Southern populations, unlike the results obtained from the nuclear data,

and more in line with some previous studies (Vidya et al., 2009, De et al., 2021).

Such discordance between mitochondrial and nuclear datasets in phylogeographic
interpretations has been reported previously too (Toews and Brelsford 2012). Since
elephants form matrilineal herds, individuals in a herd should share their
mitochondrial haplotype. It is possible that herds of matrilineally-related elephants
from the Northern population colonized Central India (CI) and North of Palghat Gap,
and subsequently founded the other southern populations, leading to the observed
patterns.

Demographic History

We investigated recent demographic history for clusters with more than nine
samples (Northern and NPG) using GONE (Santiago et al. 2020) and show both
these populations have undergone a recent bottleneck around 1500-1000 years ago
(Figure 3a). Since the taming of elephants in the subcontinent during Harappan
times, at least four millennia ago, there has been regular exploitation of wild
individuals for military and domestic use (Sukumar 2011). We do not have estimates
of the actual extent of exploitation (in terms of annual offtakes) but can only
qualitatively infer the levels of exploitation from the stocks of elephants in captivity.
Historical accounts suggest that the armies of ancient kingdoms and republics in the
north (the Gangetic basin) maintained several thousand captive elephants from as
early as the 3[rd] century BCE until late Mughal times in the early 17[th] century CE,
suggestive of overexploitation of wild populations for nearly two centuries, until the
invention and use of gunpowder in warfare rendered the war elephant irrelevant
(Sukumar 2011). This is the first report of a bottleneck in Asian elephant populations
in recent historical times, potentially mediated by human exploitation. Our results
also suggest that both populations (Northern and NPG) may have started recovering
from the bottlenecks around 300-500 years ago. However, the modelled "recovery"
could also be an artefact due to our under-sampling, or population structure. The
recent recovery-like pattern may be observed due to recent and fine-scale population
structure in the Northern cluster (Sukumar 2011, Menon and Tiwari 2019), which
would reduce shared linkage disequilibrium (LD) blocks between populations, giving
an illusion of population expansion (Novo et al., 2023). Often, population structure
can dictate the models of demographic history (Mazet et al. 2016).

Additionally, we find signatures of population bottlenecks around 100,000 years ago

(Supplementary Figure 11). However, further investigation with hPSMC (Cahill et al.

2016) and fastsimcoal 2 (Excoffier et al. 2021) suggests that this coincides with

populations differentiating from each other (Figure 3 b,c). hPSMC results suggest

that the Northern elephant population separated from all other populations about

70,000-100,000 years ago while *fastsimcoal* suggests it split from the Southern

Indian populations 134-245 Kya and from the Central Indian (CI) cluster 102-174

Kya. The hPSMC results further suggest that the Central Indian elephants diverged

from the rest around 50,000-80,000 years ago, while the three Southern Indian

populations diverged from each other only around 20,000-30,000 years ago.

However, *fastsimcoal2* supports a model with Southern Indian populations splitting

from each other between 72-100 Kya. The serial reduction of historic effective

population size from north to south indicates a potential serial founding of

populations of elephants in India along this direction (Slatkin & Excoffier 2012).

Overall, our results emphasise the antiquity of the Northern populations of elephants

consistent with Vidya et al. (2005b, 2009). Palkopoulou et al. (2018) had suggested

a more recent divergence which could be due to the differences in the mutation rates

used for making inferences (4.06e-8 vs 5.3e-9; Prado et al. 2023). Vidya et al. (2009)

made inferences on phylogeography based solely on mitochondrial DNA sequences,

which are expected to provide older estimates of divergence times even up to an

order of magnitude greater (Zheng et al. 2011). Discordance between mitochondrial

and nuclear genome estimates of phylogeography has been attributed to several

reasons include sex-biased dispersal, faster sorting rate of mtDNA than nuDNA and

differences in the geographical range or numbers of hybridizing populations (Peters

et al. 2012, Toews and Brelsford 2012); thus, inferences of phylogeny and

divergence times from mtDNA alone are best avoided (Weins et al. 2010). However,

consistent with Palkopoulou et al. (2018) who used whole genomes, we find that the

NPG and SPG populations in the south diverged from each other only around 20,000

years ago, or the time of the Last Glacial Maximum when southern India, in

particular, the Western Ghats, was more arid (Sukumar et al. 1993, Rajagopalan et

al. 1997). These findings further support the recognition of five elephant

management units in India, emphasising their antiquity and unique evolutionary

histories.

Genetic Diversity

To measure how the management units compare with each other, we estimated the genetic diversity as the number of average pairwise differences in sequences (pi) within each cluster. We find that the clusters that are from Southern India (SPG, SSG and NPG) have lower average nucleotide diversity than the Northern (NW&NE) Indian and Central Indian (CI) clusters. Interestingly, there is no visible difference in nucleotide diversity within the Southern and between the Northern and Central clusters (Figure 4a) (Supplementary Table 4). This suggests that all the clusters within Southern India have a similar number of haplotypes and that the Central and Northern cluster have a similar number of haplotypes. Expectedly, the only significant statistical differences were observed between Northern and the Southern clusters, and Central and the Southern clusters, respectively (Supplementary Table 5). About 75% of the discovered SNPs are shared, or present in all populations (Supplementary Table 6). However, the number of heterozygous sites encountered per Mb is the highest for the Northern population while it is lowest for the southernmost population (SSG) (Figure 4b). We find no difference in heterozygous SNV per Mb for NPG and CI (Supplementary Table 5). Overall, our results suggest that most populations of elephants have similar nucleotide diversity but, in the population South of the Shencottah Gap (SSG), similar nucleotides are often paired together, indicating lower effective population size, while in the populations in Northern India most of the nucleotides pair with a different one, indicative of higher effective size. These patterns are consistent with signatures of serial founding, where it is predicted that the heterozygosity of each successive population decreases in comparison to the parent population and is proportional to the effective population size of the founders (Slatkin & Excoffier 2012).

Genetic variation is related to effective population size which, in turn, is dependent not only on the census population size of a species but other variables such as the adult sex ratio (Frankel and Soulé 1981). Although we cannot speculate on the historical population sizes of Asian elephants across the various regions we have investigated, a cursory look at the recent population sizes of the five management/conservation units we have inferred from the genomic data shows that census population size is not indicative of the observed genetic variation (pi and

heterozygosity) (Supplementary Table 7). For instance, the genetic variation of the

Northeastern (NE) population is much higher than that of the Southern population to

the North of the Palghat Gap (NPG). Both these clusters have similar census

population size on the order c. 10,000 individuals. Central Indian elephants have

higher variation than Southern populations even though the former has distinctly

lower census population size of c.3000 elephants (Supplementary Table 7). This

might be indicative of recent bottlenecks in Central India, as heterozygosity decays

slowly, while the Southern populations, especially those South of the Palghat Gap

may have had historically smaller populations. The highly female-biased sex ratios in

Southern populations, especially those to the South of the Palghat Gap, in recent

historical times (1970s-1990s) from selective poaching of male elephants for ivory

(Sukumar 1989, Ramakrishnan et al. 1998), could have decreased the effective

population size (compared to census size). Elephants to the South of Shencottah

Gap were also connected to the SPG population through movement of males until

the early 1980s according to anecdotal evidence (Gangadharan et al. 2011,

Johnsingh et al. 2009). Overall, our results are thus also consistent with a serial

dilution of variation that could be the result of sequential colonization (Hellenthal et

al., 2008; Pierce et al., 2014; Pless et al., 2022) from north to south: from Northern to

Central and Northern to Southern, and then from North of Palghat Gap to South of

Shencottah Gap, in that order.

We further test this by estimating F_{ROH}, the proportion of the genome in homozygous

stretches, with longer stretches indicative of recent bottlenecks and/or inbreeding

(Ceballos et al. 2018). We observe that individuals from South of the Shencottah

Gap (SSG) have high average $F_{ROH>0.1Mb}$ of 0.4 (40% of the genome is in

homozygous stretches) while individuals from Northern India have the least average

$F_{ROH>0.1Mb}$ of 0.2 (20% of the genome is in homozygous stretches, Figure 5a). There

is no significant difference between individuals from SPG, NPG and CI populations

(average $F_{ROH>0.1Mb}$ is 0.25). Longer homozygous stretches are indicative of recent

inbreeding (Curik et al., 2014; Sumreddee, 2021). ROH longer than 1Mb and 10Mb

shows few differences between populations, although the Northern Indian population

has the least recent inbreeding. Additionally, the SPG and Northern populations

have lower variance in F_{ROH} compared to other populations, potentially indicating

gene flow and connectivity within these large landscapes, which will result in lower

variance in parental relatedness. The results suggest that despite inbreeding
avoidance in elephants (Sukumar 2003), founder effects and drift led to highly inbred
individuals.

While the F_{ROH} analyses suggest individual inbreeding, a high average F_{ROH} in a
population need not result in inbreeding depression, as the homozygous segments
may differ between individuals within a population. To measure the consequences of
mating between individuals within each population, we measured the percent
genome that is identical by descent (IBD) shared between pairs of individuals in
each population (Figure 5b). Elephants in the Northern Indian cluster (NW&NE)
share very few IBD stretches of the genome (on an average 1.6% in more than
10Mb long and 6.6% in more than 0.1Mb long IBD stretches) with each other, while
the two individuals we had from South of Shencottah Gap (SSG) shared about 43%
of their genome in stretches longer than 10Mb and about 52% of their genome in
stretches longer than 0.1Mb. Apart from the one outlier pair in the SPG sample set,
we observe negligible differences in the SPG, NPG and CI populations with regards
to the proportion IBD stretches of genome shared between pairs of individuals. On
average these three clusters share more than 10% of their genome in IBD longer
than 10Mb (NPG: 8%, SPG: 14%, CI:12%) and 19% of their genome in IBD longer
than 0.1Mb (NPG: 15%, SPG: 25%, CI:17%). These results indicate inbreeding
depression in the future in SSG, SPG, CI populations, due to inbreeding in upcoming
generations. Since the present individuals already share large IBD stretches of the
genome, impacts on fitness will be maximal, especially if these individuals have
higher numbers of deleterious alleles. However, since there are a few outlier pairs
with lower amounts of shared genomic IBD stretches (Figure 5b), there is hope that
maintaining gene flow within these population clusters could sustain extant genetic
variation in these populations into the future.

Genetic load

We attempted to understand whether individuals, from certain populations, that have
high homozygosity could suffer fitness effects due to inbreeding depression by
estimating genetic load, or putative damaging mutations. This depends on the

number of derived deleterious mutations harboured across the genome, how many
are homozygous, and the putative magnitude of their fitness effects (indel Loss of
function > Loss of function > missense mutation) (Barton & Zeng 2018; Khan et al.
2021). We normalised this by derived SNPs in neutral/synonymous/intergenic
regions, to account for possible missingness/uneven coverage across genomes. As
expected, populations with lowest genetic variation (SSG) had fewer LOF mutations
(6a, x axis). This could be because of the serial founding demographic history from
north to south (see Figure 1e). However, LOF mutations in SSG were not entirely a
subset of those in the Northern population, potentially supportive of the emergence
of *de-novo* mutations in the SSG population along with differential influence of drift
and selection due to the long history of isolation between the two populations. The
SSG population has the highest homozygosity for the LOF mutations. Interestingly,
SPG also had fewer homozygous LOF mutations (Figure 6a), potentially due to
higher effective population size and *de-novo* mutation accumulation due to historical
isolation.

To better understand whether elephant populations such as SSG are endangered by
imminent inbreeding depression, we compared number and homozygosity of indel
LOFs between all five populations. The total indel LOFs did not vary considerably
between populations (x axis, Figure 6b). The observation that all populations have a
similar indel LOF load (compared to LOF load, Figure 6a, x axis) suggests that they
are under strong purifying selection, and that these putative high negative effect
mutations could have already been purged from these populations. In contrast, the
homozygous indel LOFs remain higher in SSG and SPG. This suggests that despite
purging, elephants to South of the Palghat Gap could experience negative fitness
consequences of indel LOFs first. This population may be threatened with inbreeding
depression, or lower fitness of individuals, ensuing negative feedback on population
growth rates, and a potential for further increase in homozygosity due to inbreeding
in the future generations (Robinson et al. 2023). However, the Asian elephant in
Borneo, despite its extremely narrow genetic base (Goosens et al. 2016), possibly
the outcome of a severe bottleneck during colonization of the island in the last glacial
maximum (Sharma et al. 2018), has recovered and maintains a population of about
2000 individuals. Also, African elephant populations (such as at Addo National Park,
South Africa) grew rapidly with no apparent deleterious effects after a severe

bottleneck about a century ago (Whitehouse and Hall-Martin 2000, Whitehouse and

Harley 2001). Genetically small populations are expected to be especially efficient at

purging strongly deleterious alleles (Kyriazis et al. 2023, Khan et al. 2021), this does

seem to be happening in the isolated population to South of the Shencottah Gap

leading to fewer number of deleterious alleles there. But the question that remains is

whether these endangered populations with long generation times and low

population growth rates can tolerate high genetic load.

Southern populations (NPG, SPG and SSG) have fewer derived missense mutations

than the Central (CI) and the Northern populations (Supplementary Figure 8) but the

populations South of Palghat Gap (SPG and SSG) have higher homozygous

missense mutation load than the NPG, CI and Northern populations, again

potentially due to drift and inbreeding. The fewer number of mildly deleterious

missense alleles in the Southern Indian population to the North of Palghat Gap

(NPG) compared to CI and Northern is counterintuitive, as current literature suggests

that larger populations (Supplementary Table 7) host high numbers of mildly

deleterious alleles (Kyriazis et al. 2023, Khan et al. 2021). We suggest serial dilution

of missense variants combined with a long isolation from the larger and connected

Northern Indian population may have led to restricted immigration of mild impact

deleterious alleles in the NPG population while the LOF are purged out equally well

in the large NPG, CI and Northern populations. Additionally, the low homozygosity in

the NPG, CI and Northern populations indicates a potential for inbreeding depression

(Robinson et al. 2023). Overall, we observe that the populations in Northern India

have high numbers of deleterious alleles but the effect of these alleles are masked

due to heterozygosity. Although, the Southern populations have fewer deleterious

alleles, they have a high chance of expressing them due to high homozygosity. For

example, in the southernmost population (SSG) more than 40% of the LOF

mutations are homozygous while in the northernmost population (NW&NE) about

20% of the LOF mutations are homozygous. Thus, serial founding in elephants

seems to increase the realized genetic load in every successive founding event.

While the exact fitness effects of mutations, and the distribution of fitness effects is

unknown in most wild species, we tried to better understand the possible fitness

effects of load. The LOF mutations were annotated using the FUSIL database

(Cacheiro et al. 2020). The database creates knock-out mutations of specific genes
in mice and evaluates if specific knock-out mutations are lethal, sub-viable (only
12.5% individuals survive) or harmless. Among the genes harbouring LOF mutations
in Indian elephants, 892 were represented in the FUSIL database. Out of these 892
genes, 188 LOF mutation were found to be homozygous lethal in mice and 112
mutations were found to lead to a sub-viable phenotype in mice. Thus, about 34% of
the LOF mutations, when homozygous, produce a deleterious phenotype in mice.
This highlights the risks the elephants face in near future. While the mutations
causing lethal phenotypes are expected to be purged out faster, the ones with sub-
viable phenotype may pose the real threat and lead to a slow decline in elephant
populations.

Functionally, the derived missense mutations are related to sensory perception,
detection of chemical stimulus and especially olfaction (Supplementary Figure 9).
However, there is an abundance of olfactory genes in the elephant genome (Reddy
et al. 2015) and thus there are more opportunities for mutations. Additionally, several
olfactory genes can be pseudogenes. However, since we also polarized the alleles
with the Bornean elephant genome, the missense mutations can be confidently
classified to be recently derived and maybe potentially deleterious. The loss-of-
function mutations mostly affect protein, ions and nucleic acid binding abilities along
with transferases and transporters (Supplementary Figure 10). Although inbred
individuals are more homozygous for these mutations and maybe expected to be
severely affected, it is possible that sociality and herd living allows reduced sensory
abilities in individuals. However, this needs further investigation.

Conservation implications

Overall, our results suggest five management units for Asian elephants in India.
Elephants in the Himalayan foothills from the Northwest to the Northeast of India are
a single genetic cluster that diverged from other Indian populations more than 70,000
years ago. This makes the Northern (NW&NE) cluster the oldest and, hence, the
most evolutionarily unique population.

Elephant to South of the Shencottah Gap need priority conservation attention to
better understand census sizes, connectivity and genomic variation. Conservation

and management action should focus on protecting remaining habitats, minimise

unnatural deaths, examine the feasibility of restoring the link between NPG and

SSG, and serious considerations about translocations into and out of this population.

More genetic sampling of the SSG population as well as studies on inbreeding and

possible associated phenotypes will be important to understand future fitness

trajectories here.

Populations to the North and South of the Palghat Gap are distinct management

units, and animals should be moved across this biogeographic divide only after very

careful biological evaluations of the consequences. Further, given the high genetic

load in these two management units, connectivity must be maintained within each of

these two major landscapes across the gap.

The Ganges and Brahmaputra rivers in the north function as biogeographic divides,

though incomplete, making elephants south of these rivers in Central India a unique

management/conservation unit. Detailed and geographically widespread genetic

studies in central India are important to better understand this management unit, in

particular in the recent context of large-scale dispersals.

Overall, elephants across India face significant challenges because of a suite of

factors, mainly negative human impacts, and we hope that conservation efforts can

be bolstered by recognition of their unique evolutionary history. Using whole genome

data (Weins et al. 2010, Toews and Brelsford 2012) as opposed to solely

mitochondrial DNA (Vidya et al. 2009) can allow a better understanding of how Asian

elephant evolution was shaped by Pleistocene climate cycles, and will respond to

future climate change in the subcontinent (Kanagaraj et al. 2019). Previously

researchers have suggested mixing lineages for conservation (provided risks of

outbreeding depression are evaluated (Ralls et al. 2018), or assumptions can be

made to ignore outbreeding depression (Powell 2023)). We find deep divergences

between the populations of elephants. We recommend that maintaining historic

connectivity between populations to allow natural dispersal between populations will

be a better approach than translocation for conservation until risks of geneflow can

be evaluated with empirical evidence.

While the impact of genetic variation on survival of small and isolated populations
continues to be investigated, several recent studies highlight purging of deleterious
allele load (Khan et al. 2021, Robinson et al. 2023 and the references therein). This
study on Indian elephants reveals that while the number of deleterious alleles is low
in small populations (such as SSG), the realized load is high because of higher
homozygosity. Additionally, our data demonstrate that populations can have low (but
unmasked) deleterious allele load due to their unique demographic histories.
Regardless, populations with homozygous deleterious mutation load should have
high priority for conservation. The exact phenotypic and demographic trajectories of
inbreeding depression and extinction will depend on the specific load, and maybe
difficult to generalise across populations and species.

Our approach of using high-throughput genomic data allowed for unique and
important conservation insights. Elephants in India present one of the first clear
examples of serial founding of a large landscape at a subcontinental scale, with
decreasing founder size from the source. Our data allows us to detect recent
declines, potentially mediated by historic elephant captures on a large scale. Further,
elephants provide an excellent system to better understand the interplay between
genetic load and inbreeding in a set of five populations, a rare set up in endangered
species. While on-ground conservation challenges for elephants remain mitigation of
human-associated conflict and human infrastructure associated mortality,
conservation genomics insights provide long-term conceptual guidance for future
survival of these populations.

Methods

Sample collection: State Forest Department databases of captive elephants were
examined from 6 states, to locate individual elephants that were captured or rescued
from all of the known four major disjunct populations across India. Within these
populations, individuals representing habitats across major barriers such as the
Palghat and the Shencottah Gaps in Southern India, and the Ganges and the
Brahmaputra rivers in Northern (Northwestern and Northeastern) India were also
located. Blood samples and exact capture locations were collected from 28 such
individuals and included in our final dataset (see Supplementary Table 1). Out of
those samples, 12 were collected from the Southern eco-region, four were collected

from the Central eco-region, three were collected from the Northwestern eco-region,

and lastly nine were collected from the Northeastern eco-region. Out of the 12

samples collected from the Southern eco-region, six were collected from North of

Palghat Gap, four from South of Palghat Gap (but North of Shencottah Gap), and

two from South of Shencottah Gap. We additionally included genomic data from five

more individuals obtained from online sources — one individual each from Borneo,

Myanmar and Northeastern India, and three individuals from Southern India (two

North of Palghat Gap and one South of Palghat Gap) (See Supplementary Table 1).

DNA extraction, library preparation and sequencing

Genomic DNA was extracted from blood samples using Qiagen DNeasy Blood &

Tissue Kit. The library preparation and whole genome resequencing was carried out

at Medgenome INC. The DNA extraction and sequencing were carried out following

Khan et al., 2020.

Variant calling and filtering

The raw sequencing reads were trimmed using the default settings in TrimGalore-

0.4.5 (https://github.com/FelixKrueger/TrimGalore). The trimmed reads were

mapped to the *Elephas maximus* reference genome

(https://dnazoo.s3.wasabisys.com/index.html?prefix=Elephas_maximus/) using

default settings of BWA *mem* (https://github.com/lh3/bwa). The mapped reads were

converted to binary format and sorted using Samtools-1.9 (Li et al. 2009). The PCR

and optical duplicates were marked using Picardtools MarkDuplicates

(http://broadinstitute.github.io/picard) or Samtools-markdup. SNPs were identified

using Strelka variant caller 2.9.10 (Saunders et al. 2012) for only the chromosomal

scaffolds (scaffolds named HiC_scaffold_1 to HiC_scaffold_28). The variants were

filtered using VcftoolsV13 (Danecek et al. 2011). We removed indels and retained

only SNP loci with minimum phred scaled base quality 30, minimum genotype quality

30, a minimum minor allele count of 3, did not deviate from Hardy-Weinberg

equilibrium with chi square p value of 0.05. All sites with the FILTER flag other than

PASS were removed. Sites that showed mean depth across individuals more than

the 97.5th percentile and less than 2.5th percentile or were missing in more than 20%

individuals, were removed. We identified the sex chromosome scaffolds as

described in Armstrong et al. (2021) and removed the sex chromosome scaffold

identified as HiC_scaffold_28.

Population genetic structure:

PCA

We first employed PCA to partition the data along their main axes of variation. The

PCA was carried out using the *--pca* function in Plink (Purcell et al. 2007) based on

the final filtered variants. The resulting eigenvectors were plotted in R using ggplot2.

ADMIXTURE

Thereafter, a maximum likelihood-based method, ADMIXTURE (Alexander et al.

2009) was employed to investigate the population genomic structure of Asian

elephants. We estimated the number of clusters across *K* values ranging from 1 to 6,

based on the number of clusters obtained from PCA. The evaluation of the most

optimal number of clusters was carried out using EvalAdmix (Garcia☐Erill et al.

2020). EvalAdmix estimates the correlation of the residual matrices of the individuals

from the Admixture analyses. The resulting correlation matrices were plotted in R.

qpgraph

We assigned a population to each individual based on the results from admixture.

We converted the .vcf file to .ped format using VCFtools and then .ped to .bed format

using Plink. Employed the *find_graphs* function in ADMIXTOOLS2

(https://github.com/uqrmaie1/admixtools) to automatically find the optimum graphs

using various values for the number of admixture events. The model with zero

admixture had the best statistical support as tested using the protocol suggested in

ADMIXTOOLS2 with functions *qpgraph_resample_multi* and *compare_fits*.

Demographic history

Recent Demographic History

We used the LD based GONE (Santiago et al. 2020) for estimating recent

demography history up to a couple of hundred generations ago. We used the default

settings of the parameters and set *PHASE* as 0. We plotted the results assuming a

generation time of 31 years (Palkopoulou et al. 2018). This is a multiple sample

analysis and requires several samples for accurate estimates. Hence, we performed

this analysis only for the NW&NE and the NPG cluster which have at least nine

samples each. The results from this analysis better predict recent events but are not

very reliable for estimating events older than a hundred generations (Santiago et al.

2020).

Divergence time

Divergence time was estimated by contrasting various models using *fastsimcoal2.7*

(Excoffier et al. 2021) and hPSMC (Cahill et al. 2016).

We filtered the autosomal SNPs to remove sites that are in LD with r^2 value of 0.5.

This retained 80,684 SNPs. We estimated folded site frequency spectrum using

easySFS (https://github.com/isaacovercast/easySFS) to subsample 9,4,6,4,2 diploid

individuals from the NW&NE, CI, NPG, SPG and SSG populations using the options

-a --proj 18,8,12,8,4. This allowed us to retain maximum number of non-missing

SNPs for each of these populations while maximising the number of individuals

retained. We tested scenarios presented in Figures 7a and 7b with constant

population size and growth rate. We model two scenarios of divergence for both

models in Figures 7a and 7b, one with advent of agriculture (7000 ya) as the

maximum time of divergence and another with minimum divergence time around the

last glacial maxima (17,500ya, Jones et al. 2017). We estimated parameters using

fastsimcoal2.7 using options *-n 500,000 -m -M 0.001 -l 30 -L 60* as described in

Armstrong et al. (2021). We then selected the best model using the best AIC values

(Supplementary Table 8). We estimated confidence intervals using bootstrap. For

this, we randomly sampled 80% of the loci from the dataset 10 times and estimated

parameter with initial values declared using *-initvalues* and fastsimcoal2.7 options *-n*

500,000 -m -M 0.001 -l 30 -L 60.

We also estimate divergence times between populations using hPSMC (Cahill et al.

2016). We used a single individual selected at random from each population

(Supplementary Table 1). We created a pseudo-diploid individual using samples

from two populations for each scaffold except for the sex chromosome scaffold

(https://github.com/jacahill/hPSMC). Then performed PSMC with the default settings.

We plotted the results assuming a mutation rate of 5.3*10^-9/base/generation (Prado

et al. 2023) and a generation time of 31 years. The time point where the effective

population size estimated from the pseudo-diploid individual rises exponentially is

the point where the two haplotypes do not coalesce and hence signify population

divergence.

Genetic diversity

Nucleotide diversity: pi

We randomly subsampled four individuals from NW&NE, CI, NPG and SPG sample

sets and retained the SSG dataset (n=2) as is. We estimated the average number of

pairwise differences per site for each population cluster as described in Wang et al.

(2020). Briefly, we set the function *dosaf* to 1 ANGSD (Korneliussen et al. 2014) to

estimate allele frequency likelihood for each site. The we used the used the function

realSFS and estimated a folded site frequency spectrum. We set the function

doThetas to 1 and estimated pi per site and used the *ThetaStat* function the

summarise the pi value for each scaffold. We used only the chromosomal scaffolds

from this summary and divided the "tP" values by the number of sites for each

scaffold to obtain the average pi per site. Furthermore, we statistically compared the

significance of the estimated values between populations (see Supplementary Table

4).

Heterozygous SNV encounter rate

We estimated the number of heterozygous SNPs for each individual using the

vcfstats function of RTGtools (Cleary et al. 2015,

https://github.com/RealTimeGenomics/rtg-tools). We then divided the number of

heterozygous sites by the total number of sites genotyped for the individual and

multiplied by 10^6 to obtain SNV encounter rate/Mb.

Inbreeding

FROH

We estimated runs of homozygosity (ROH) using the *roh* function in BCFtools

version 1.3 (Narasimhan et al. 2016) as described previously (Shukla et al. 2023).

We classified the ROH stretches into three size classes: the more than 0.1 Mb (100

Kb) stretches show cumulative inbreeding due to old and recent bottlenecks, the

more than 1Mb stretches show cumulative inbreeding in the recent past and the

more than 10Mb stretches show recent inbreeding. FROH was estimated as

described previously (Khan et al. 2021).

Shared IBD stretches

We used IBDseq (Browning & Browning, 2013) to detect the stretches of genome

that are identical-by-descent in pairs of individuals. For each pair of individuals in a

population, we summed the lengths of IBD stretches longer than 0.1Mb and 10Mb

and divided by the total autosomal length.

Genetic load

Filtering variant sites

We filtered the raw variants obtained from Strelka variant caller 2.9.10 again in

VCFtools using the parameters with minimum phred scaled base quality 30,

minimum genotype quality 30, minor allele count 1, FILTER flag as PASS. We also

removed sited with $-0.5>Fis>0.95$ as described previously (Khan et al. 2021) and

removed sites missing in more than 20% of the individuals. We filtered out the sex

chromosomes as well. We call this set of loci as deleterious_allele_set.

Identifying ancestral alleles

We defined the ancestral allele as the allele present most commonly in sister species

of Asiatic elephants. For this we used the genome of African elephant (genbank

accession: GCA_000001905.1) and dugong (genbank accession: GCA_015147995).

We converted these assemblies to 100bp long FASTQ reads and mapped to the

Asiatic elephant reference genome and removed reads mapping to multiple regions

as described in Khan et al. (2021). We converted the mapped reads to consensus

fasta files and estimated the majority allele at a locus. We further filtered these

alleles with the genome of Bornean elephant (SRA accession: ERR2260499). Any

loci where the Bornean elephant was not homozygous for the ancestral allele was

removed from the analysis thus ensuring that the derived alleles are new to the

population which is an important assumption for the identification of deleterious

alleles (Khan 2023).

Derived indel load

We further filtered the loci in the deleterious_allele_set using the *--keep-only-indels*

tag in VCFtools. We annotated this set of loci with the Asiatic elephant genome

annotation (https://dnazoo.s3.wasabisys.com/index.html?prefix=Elephas_maximus/),

consisting of 26246 coding regions, using Ensembl Variant Effect Predictor (VEP,

McLaren et al. 2016). For indels, the ancestral state cannot be determined using

distant species. Hence, the indel allele that was homozygous in the Bornean

elephant was referred to as the ancestral allele and any indel site where the Bornean

elephant was not homozygous, was removed from the analysis. We then estimated

the number of indels predicted to cause transcript_ablation, splice_donor_variant,

splice_acceptor_variant, stop_gained, frameshift_variant, inframe_insertion,

inframe_deletion, splice_region_variant were classified as LoF causing indels. We

also counted the derived indels in the intergenic regions of the genome. We divided

the number of LoF indels with the number of intergenic indels to control for

differences in depth leading to missingness in the data.

Derived LoF load

We filtered the loci in the deleterious_allele_set to remove indels and that had mean

depth across individuals more than the 97.5th percentile and less than 2.5th

percentile. This set was annotated on VEP and LoF mutations were identified as

described earlier. We then counted the number of derived LoF alleles and divided

them with the number of synonymous mutations.

Derived missense load

We used the same set of loci used for estimating the LoF load but chose those set of

loci that caused non-synonymous changes in the genome and followed the same

procedure as described for the LoF load.

Acknowledgements

This work was supported by a Department of Biotechnology grant [BT/PR23223/BCE/8/1490/2018] awarded to RS and UR. Permissions for sample collection were granted in letters C-66011/02/2019 dated 09/07/2019 (West Bengal), 1729/5-6 dated 09/12/2020 (Uttarakhand), 4838/2019-CWW/WL10 dated 06/10/2019 (Kerala), PCCF(WL)/E2/CR-79/2018-19 dated 26/05/2018 (Karnataka), WL/FG31/Technical Committee/2019 dated 26/08/2019 (Assam) and CWL/G/173/2018-19/Pt.VII/1695-702 dated 10/10/2019 (Arunachal Pradesh). We would like to thank the respective forest veterinary doctors of all the states, particularly Dr. Aditi Sharma, Dr. Arun Zachariah and Dr. Shweta Mandal. RS was supported as National Science Chair by Science and Engineering Research Board (SERB), Government of India. KMG is supported from the DBT-Ramalingaswami Fellowship (No. BT/HRD/35/02/2006). Gratitude to NCBS IT team support during data analysis; NCBS data cluster used is supported under project no. 12-R&D-TFR-5.04-0900, Department of Atomic Energy, Government of India). VV Robin and Ansil BR provided helpful comments on drafts of the manuscript. Gratitude to field assistants for their continuous presence in the field and assisting in sample collection for this work.

Data availability

All sequencing data have been deposited in BioProject SUB13810487 in NCBI.

References

Alexander, D.H., Novembre, J. and Lange, K., 2009. Fast model-based estimation of ancestry in unrelated individuals. Genome research, 19(9), pp.1655-1664.

Armstrong, E.E., Khan, A., Taylor, R.W., Gouy, A., Greenbaum, G., Thiéry, A., Kang, J.T., Redondo, S.A., Prost, S., Barsh, G. and Kaelin, C., 2021. Recent evolutionary history of tigers highlights contrasting roles of genetic drift and selection. *Molecular Biology and Evolution*, *38*(6), pp.2366-2379.

Baskaran, N., Balasubramanian, M., Swaminathan, S. and Desai, A.A., 1995. Home range of elephants in the Nilgiri Biosphere Reserve, south India. A week with elephants. Bombay Natural History Society and Oxford University Press, Bombay, pp.296-313.

Barton, H.J. and Zeng, K., 2018. New methods for inferring the distribution of fitness effects for INDELs and SNPs. *Molecular Biology and Evolution*, *35*(6), pp.1536-1546.

Berzaghi, F., Bretagnolle, F., Durand-Bessart, C. and Blake, S., 2023. Megaherbivores modify forest structure and increase carbon stocks through multiple pathways. *Proceedings of the National Academy of Sciences*, *120*(5), p.e2201832120.

Browning, B.L. and Browning, S.R., 2013. Detecting identity by descent and estimating genotype error rates in sequence data. The American Journal of Human Genetics, 93(5), pp.840-851

Brüniche-Olsen, A., Kellner, K.F., Anderson, C.J. and DeWoody, J.A., 2018. Runs of homozygosity have utility in mammalian conservation and evolutionary studies. *Conservation Genetics*, *19*(6), pp.1295-1307.

Cacheiro, P., Muñoz-Fuentes, V., Murray, S.A., Dickinson, M.E., Bucan, M., Nutter, L.M., Peterson, K.A., Haselimashhadi, H., Flenniken, A.M., Morgan, H. and Westerberg, H., 2020. Human and mouse essentiality screens as a resource for disease gene discovery. *Nature communications*, *11*(1), p.655.

Cahill, J.A., Soares, A.E., Green, R.E. and Shapiro, B., 2016. Inferring species
divergence times using pairwise sequential Markovian coalescent modelling and low-
coverage genomic data. Philosophical Transactions of the Royal Society B:
Biological Sciences, 371(1699), p.20150138.

Ceballos, F.C., Joshi, P.K., Clark, D.W., Ramsay, M. and Wilson, J.F., 2018. Runs of
homozygosity: windows into population history and trait architecture. *Nature Reviews
Genetics*, *19*(4), pp.220-234.

Chaitanya, R., Giri, V.B., Deepak, V., Datta-Roy, A., Murthy, B.H.C.K. and Karanth,
P., 2019. Diversification in the mountains: a generic reappraisal of the Western
Ghats endemic gecko genus Dravidogecko Smith, 1933 (Squamata: Gekkonidae)
with descriptions of six new species. Zootaxa, 4688(1), pp.1-56.

Cleary, J.G., Braithwaite, R., Gaastra, K., Hilbush, B.S., Inglis, S., Irvine, S.A.,
Jackson, A., Littin, R., Rathod, M., Ware, D. and Zook, J.M., 2015. Comparing
variant call files for performance benchmarking of next-generation sequencing
variant calling pipelines. *BioRxiv*, p.023754.

Coverdale, T.C., Kartzinel, T.R., Grabowski, K.L., Shriver, R.K., Hassan, A.A.,
Goheen, J.R., Palmer, T.M. and Pringle, R.M., 2016. Elephants in the understory:
opposing direct and indirect effects of consumption and ecosystem engineering by
megaherbivores. Ecology, 97(11), pp.3219-3230.

Curik, I., Ferenčaković, M. and Sölkner, J., 2014. Inbreeding and runs of
homozygosity: A possible solution to an old problem. Livestock Science, 166, pp.26-
34.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A.,
Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T. and McVean, G., 2011. The
variant call format and VCFtools. Bioinformatics, 27(15), pp.2156-2158.

De, R., Sharma, R., Davidar, P., Arumugam, N., Sedhupathy, A., Puyravaud, J.P.,
Selvan, K.M., Rahim, P.A., Udayraj, S., Parida, J. and Digal, D.K., 2021. Pan-India
population genetics signifies the importance of habitat connectivity for wild Asian
elephant conservation. Global Ecology and Conservation, 32, p.e01888.

Enquist, B.J., Abraham, A.J., Harfoot, M.B., Malhi, Y. and Doughty, C.E., 2020. The
megabiota are disproportionately important for biosphere functioning. *Nature*
Communications, *11*(1), p.699

Excoffier, L., Marchi, N., Marques, D.A., Matthey-Doret, R., Gouy, A. and Sousa,
V.C., 2021. fastsimcoal2: demographic inference under complex evolutionary
scenarios. *Bioinformatics*, *37*(24), pp.4882-4885.

Frankel, O.H. and Soulé, M.E. 1981. Conservation and evolution. Cambridge
University Press, Cambridge, U.K.

Fernando, P., Pfrender, M.E., Encalada, S.E. and Lande, R., 2000. Mitochondrial
DNA variation, phylogeography and population structure of the Asian
elephant. *Heredity*, *84*(3), pp.362-372.

Fleischer, R.C., Perry, E.A., Muralidharan, K., Stevens, E.E. and Wemmer, C.M.,
2001. Phylogeography of the Asian elephant (Elephas maximus) based on
mitochondrial DNA. *Evolution*, *55*(9), pp.1882-1892.

Gangadharan, A., Vaidyanathan, S. and Ram, S., 2011. Identifying critical areas for
a landscape-level wildlife corridor in the southern Western Ghats. *Foundation for*
Ecological Research, Advocacy and Learning (FERAL), Pondicherry.

Goossens, B., Sharma, R., Othman, N., Kun-Rodrigues, C., Sakong, R., Ancrenaz,
M., Ambu, L.N., Jue, N.K., O'Neill, R.J., Bruford, M.W. and Chikhi, L., 2016. Habitat
fragmentation and genetic diversity in natural populations of the Bornean elephant:
Implications for conservation. *Biological Conservation*, *196*, pp.80-92.

Gubbi, S., Swaminath, M.H., Poornesha, H.C., Bhat, R. and Raghunath, R., 2014.
An elephantine challenge: human–elephant conflict distribution in the largest Asian
elephant population, southern India. Biodiversity and conservation, 23, pp.633-647.

Harich, F.K., Treydte, A.C., Ogutu, J.O., Roberts, J.E., Savini, C., Bauer, J.M. and
Savini, T., 2016. Seed dispersal potential of Asian elephants. Acta Oecologica, 77,
pp.144-151.

Hedrick, P.W., 1999. Perspective: highly variable loci and their interpretation in
evolution and conservation. Evolution, 53(2), pp.313-318.

Hellenthal, G., Auton, A. and Falush, D., 2008. Inferring human colonization history
using a copying model. PLoS genetics, 4(5), p.e1000078.

Johnsingh, A.J.T., Sathyakumar, S. and Sunderraj, S.W., 1991. Ariankavu Pass, a
lost elephant corridor in South India. *Environmental Conservation*, *18*(4), pp.368-
368.

Jones, R.A., Williams, J.W. and Jackson, S.T., 2017. Vegetation history since the
last glacial maximum in the Ozark highlands (USA): A new record from Cupola Pond,
Missouri. *Quaternary Science Reviews*, *170*, pp.174-187.

Kanagaraj, R., Araujo, M.B., Barman, R., Davidar, P., De, R., Digal, D.K., Gopi, G.V.,
Johnsingh, A.J.T., Kakati, K., Kramer-Schadt, S. and Lamichhane, B.R., 2019.
Predicting range shifts of Asian elephants under global change. *Diversity and*
Distributions, *25*(5), pp.822-838.

Khan, A., Krishna, S.M., Ramakrishnan, U. and Das, R., 2022. Recapitulating whole
genome based population genetic structure for Indian wild tigers through an ancestry
informative marker panel. Heredity, 128(2), pp.88-96.

Khan, A., Patel, K., Shukla, H., Viswanathan, A., van der Valk, T., Borthakur, U.,
Nigam, P., Zachariah, A., Jhala, Y.V., Kardos, M. and Ramakrishnan, U., 2021.
Genomic evidence for inbreeding depression and purging of deleterious genetic
variation in Indian tigers. *Proceedings of the National Academy of Sciences*, *118*(49),
p.e2023018118.

Khan, A., Patel, K., Bhattacharjee, S., Sharma, S., Chugani, A.N., Sivaraman, K.,
Hosawad, V., Sahu, Y.K., Reddy, G.V. and Ramakrishnan, U., 2020. Are shed hair
genomes the most effective noninvasive resource for estimating relationships in the
wild?. *Ecology and Evolution*, *10*(11), pp.4583-4594.

Khan, A., 2023. The year of the tiger and the year of tiger genomes!. *Molecular*
Ecology Resources, *23*(2), pp.327-329.

Koirala, R.K., Ji, W., Aryal, A., Rothman, J. and Raubenheimer, D., 2016. Dispersal
and ranging patterns of the Asian Elephant (Elephas maximus) in relation to their
interactions with humans in Nepal. Ethology Ecology & Evolution, 28(2), pp.221-231.

Korneliussen, T.S., Albrechtsen, A. and Nielsen, R., 2014. ANGSD: analysis of next
generation sequencing data. *BMC bioinformatics*, *15*(1), pp.1-13.

Kyriazis, C.C., Beichman, A.C., Brzeski, K.E., Hoy, S.R., Peterson, R.O., Vucetich,
J.A., Vucetich, L.M., Lohmueller, K.E. and Wayne, R.K., 2023. Genomic
underpinnings of population persistence in Isle Royale moose. Molecular biology and
evolution, 40(2), p.msad021.

Le Roux, E., Kerley, G.I. and Cromsigt, J.P., 2018. Megaherbivores modify trophic
cascades triggered by fear of predation in an African savanna ecosystem. *Current*
Biology, *28*(15), pp.2493-2499.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R. and 1000 Genome Project Data Processing Subgroup,
2009. The sequence alignment/map format and SAMtools. bioinformatics, 25(16),
pp.2078-2079.

Liu, P., Wen, H., Harich, F.K., He, C., Wang, L., Guo, X., Zhao, J., Luo, A., Yang, H.,
Sun, X. and Yu, Y., 2017. Conflict between conservation and development: cash
forest encroachment in Asian elephant distributions. Scientific Reports, 7(1), p.6404.

Mahmood, T., Vu, T.T., Campos-Arceiz, A., Akrim, F., Andleeb, S., Farooq, M.,
Hamid, A., Munawar, N., Waseem, M., Hussain, A. and Fatima, H., 2021. Historical
and current distribution ranges and loss of mega-herbivores and carnivores of Asia.
PeerJ, *9*, p.e10738.

Mazet, O., Rodríguez, W., Grusea, S., Boitard, S. and Chikhi, L., 2016. On the
importance of being structured: instantaneous coalescence rates and human
evolution—lessons for ancestral population size inference? Heredity, 116(4), pp.362-
371.

McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R., Thormann, A., Flicek, P.
and Cunningham, F., 2016. The ensembl variant effect predictor. Genome biology,
17(1), pp.1-14.

Menon, V. and Tiwari, S.K., 2019. Population status of Asian elephants Elephas
maximus and key threats. *International Zoo Yearbook*, *53*(1), pp.17-30.

Menotti-Raymond, M. and O'Brien, S.J., 1993. Dating the genetic bottleneck of the
African cheetah. *Proceedings of the National Academy of Sciences, 90*(8), pp.3172-
3176.

Narasimhan, V., Danecek, P., Scally, A., Xue, Y., Tyler-Smith, C. and Durbin, R.,
2016. BCFtools/RoH: a hidden Markov model approach for detecting autozygosity
from next-generation sequencing data. Bioinformatics, 32(11), pp.1749-1751.

Novo, I., Ordás, P., Moraga, N., Santiago, E., Quesada, H. and Caballero, A., 2023.
Impact of population structure in the estimation of recent historical effective
population size by the software GONE. Genetics Selection Evolution, 55(1), p.86.

Owen-Smith, N., 1987. Pleistocene extinctions: the pivotal role of megaherbivores.
Paleobiology, 13(3), pp.351-362.

Padalia, H., Ghosh, S., Reddy, C.S., Nandy, S., Singh, S. and Kumar, A.S., 2019.
Assessment of historical forest cover loss and fragmentation in Asian elephant
ranges in India. Environmental Monitoring and Assessment, 191, pp.1-13.

Palkopoulou, E., Lipson, M., Mallick, S., Nielsen, S., Rohland, N., Baleka, S.,
Karpinski, E., Ivancevic, A.M., To, T.H., Kortschak, R.D. and Raison, J.M., 2018. A
comprehensive genomic history of extinct and living elephants. Proceedings of the
National Academy of Sciences, 115(11), pp.E2566-E2574.

Parida, J., Sharma, R., De, R., Kalam, T., Sedhupathy, A., Digal, D.K., Reddy, P.A.,
Goyal, S.P., Puyravaud, J.P. and Davidar, P., 2022. Genetic characterisation of
fragmented Asian elephant populations with one recent extinction in its eastern-
central Indian range. Ecological Genetics and Genomics, 24, p.100132.

Peters, J.L., Bolender, K.A. and Pearce, J.M., 2012. Behavioural vs. molecular
sources of conflict between nuclear and mitochondrial DNA: the role of male‐biased
dispersal in a Holarctic sea duck. *Molecular Ecology, 21*(14), pp.3562-3575.

Pierce, A.A., Zalucki, M.P., Bangura, M., Udawatta, M., Kronforst, M.R., Altizer, S.,
Haeger, J.F. and de Roode, J.C., 2014. Serial founder effects and genetic
differentiation during worldwide range expansion of monarch butterflies. Proceedings
of the Royal Society B: Biological Sciences, 281(1797), p.20142230.

Pless, E., Powell, J.R., Seger, K.R., Ellis, B. and Gloria-Soria, A., 2022. Evidence for
serial founder events during the colonization of North America by the yellow fever
mosquito, Aedes aegypti. Ecology and Evolution, 12(5), p.e8896.

Powell, D.M., 2023. Losing the forest for the tree? On the wisdom of subpopulation
management. *Zoo Biology*, *42*(5), pp.591-604.

Prado, N.A., Armstrong, E.E., Brown, J.L., Goldenberg, S.Z., Leimgruber, P.,
Pearson, V.R., Maldonado, J.E. and Campana, M.G., 2023. Genomic Resources for
Asian (Elephas maximus) and African Savannah Elephant (Loxodonta africana)
Conservation and Health Research. *bioRxiv,* pp.2023-02.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D.,
Maller, J., Sklar, P., De Bakker, P.I., Daly, M.J. and Sham, P.C., 2007. PLINK: a tool
set for whole-genome association and population-based linkage analyses. The
American journal of human genetics, 81(3), pp.559-575.

Rajagopalan, G., Sukumar, R., Ramesh, R., Pant, R.K. and Rajagopalan, G., 1997.
Late Quaternary vegetational and climatic changes from tropical peats in southern
India–an extended record up to 40,000 years BP. *Current Science*, pp.60-63. .

Ralls, K., Ballou, J.D., Dudash, M.R., Eldridge, M.D., Fenster, C.B., Lacy, R.C.,
Sunnucks, P. and Frankham, R., 2018. Call for a paradigm shift in the genetic
management of fragmented populations. *Conservation Letters*, *11*(2), p.e12412.

Ram, M.S., Marne, M., Gaur, A., Kumara, H.N., Singh, M., Kumar, A. and Umapathy,
G., 2015. Pre-historic and recent vicariance events shape genetic structure and
diversity in endangered lion-tailed macaque in the Western Ghats: implications for
conservation. PloS one, 10(11), p.e0142597.

Robin, V.V., Vishnudas, C.K., Gupta, P. and Ramakrishnan, U., 2015. Deep and
wide valleys drive nested phylogeographic patterns across a montane bird
community. Proceedings of the Royal Society B: Biological Sciences, 282(1810),
p.20150861.

Robinson, J., Kyriazis, C.C., Yuan, S.C. and Lohmueller, K.E., 2023. Deleterious
variation in natural populations and implications for conservation genetics. Annual
review of animal biosciences, 11, pp.93-114.

Santiago, E., Novo, I., Pardiñas, A.F., Saura, M., Wang, J. and Caballero, A., 2020.
Recent demographic history inferred by high-resolution analysis of linkage
disequilibrium. Molecular Biology and Evolution, 37(12), pp.3642-3653.

Saunders, C.T., Wong, W.S., Swamy, S., Becq, J., Murray, L.J. and Cheetham, R.K.,
2012. Strelka: accurate somatic small-variant calling from sequenced tumor–normal
sample pairs. Bioinformatics, 28(14), pp.1811-1817.

Sekar, N., Lee, Chia-Lo and Sukumar, R. (2017). Functional non-redundancy of
elephants in a disturbed tropical forest. Conservation Biology
DOI: 10.1111/cobi.12907

Sharma, R., Goossens, B., Heller, R., Rasteiro, R., Othman, N., Bruford, M.W. and
Chikhi, L., 2018. Genetic analyses favour an ancient and natural origin of elephants
on Borneo. *Scientific Reports*, *8*(1), p.880.

Slatkin, M. and Excoffier, L., 2012. Serial founder effects during range expansion: a
spatial analog of genetic drift. *Genetics*, *191*(1), pp.171-181.

Srinivasaiah NM, Vaidyanathan S, Sukumar R, Sinha A. (2021). Elephants on the
move: Implications for human–elephant interactions. In: *The Crisis of Climate*
Change: Weather Report. (eds. Agarwal R and Goyal O). Routledge, Abingdon,
United Kingdom and New York, pp. 92-103

Srinivasaiah NM, Vaidyanathan S, Sukumar R, Sinha A. (2022). The rurban
elephant: Behavioural ecology of Asian elephants in response to large-scale land
use change in a human-dominated landscape in peri-urban southern India. In: *New*
Forms of Urban Agriculture: An Urban Ecology Perspective (eds. Diehl J A and Kaur
H). Springer Nature, Singapore, pp. 289-31

Sukumar, R. 1989. The Asian elephant: ecology and management. Cambridge
University Press, Cambridge, U.K.

Sukumar, R., Ramakrishnan, U. and Santosh, J.A., 1998, November. Impact of
poaching on an Asian elephant population in Periyar, southern India: a model of
demography and tusk harvest. In *Animal Conservation forum* (Vol. 1, No. 4, pp. 281-
291). Cambridge University Press.

Sukumar, R., Ramesh, R., Pant, R.K. and Rajagopalan, G., 1993. A δ13C record of

late Quaternary climate change from tropical peats in southern

India. *Nature*, *364*(6439), pp.703-706.

Sukumar, R. 2003. The Living Elephants: Evolutionary Ecology, Behavior and

Conservation. Oxford University Press, New York

Sukumar, R., 2011. *The story of Asia's elephants*. Mumbai: MARG Foundation.

Shukla, H., Suryamohan, K., Khan, A., Mohan, K., Perumal, R.C., Mathew, O.K.,

Menon, R., Dixon, M.D., Muraleedharan, M., Kuriakose, B. and Michael, S., 2023.

Near-chromosomal de novo assembly of Bengal tiger genome reveals genetic

hallmarks of apex predation. GigaScience, 12, p.giac112.

Sumreddee, P., Hay, E.H., Toghiani, S., Roberts, A., Aggrey, S.E. and Rekaya, R.,

2021. Grid search approach to discriminate between old and recent inbreeding using

phenotypic, pedigree and genomic information. BMC genomics, 22, pp.1-17

Terhorst, J., Kamm, J.A. and Song, Y.S., 2017. Robust and scalable inference of

population history from hundreds of unphased whole genomes. Nature genetics,

49(2), pp.303-309.

Toews, D.P. and Brelsford, A., 2012. The biogeography of mitochondrial and nuclear

discordance in animals. *Molecular ecology*, *21*(16), pp.3907-3930.

Venkataraman, A.B., Saandeep, R., Baskaran, N., Roy, M., Madhivanan, A. and

Sukumar, R., 2005. Using satellite telemetry to mitigate elephant–human conflict: an

experiment in northern West Bengal, India. Current Science, pp.1827-1831.

Vidya, T.N.C., Fernando, P., Melnick, D.J. and Sukumar, R. 2005a. Population

differentiation within and among Asian elephant (Elephas maximus) populations in

southern India. Heredity, 94(1), pp.71-80.

Vidya, T.N.C., Fernando, P., Melnick, D.J. and Sukumar, R., 2005b. Population
genetic structure and conservation of Asian elephant (*Elephas maximus*) across
India. Animal Conservation, 8: 377-388.

Vidya, T.N.C., Sukumar, R. and Melnick, D.J., 2009. Range-wide mtDNA
phylogeography yields insights into the origins of Asian elephants. Proceedings of
the Royal Society B: Biological Sciences, 276(1658), pp.893-902.

Vijayakumar, S.P., Menezes, R.C., Jayarajan, A. and Shanker, K., 2016. Glaciations,
gradients, and geography: multiple drivers of diversification of bush frogs in the
Western Ghats Escarpment. Proceedings of the Royal Society B: Biological
Sciences, 283(1836), p.20161011.

Whitehouse, A.M. and Hall-Martin, A.J., 2000. Elephants in Addo Elephant National
Park, South Africa: reconstruction of the population's history. Oryx, 34(1), pp.46-55.

Whitehouse, A.M. and Harley, E.H., 2001. Post-bottleneck genetic diversity of
elephant populations in South Africa, revealed using microsatellite
analysis. *Molecular ecology*, *10*(9), pp.2139-2149.

Waldram, M.S., Bond, W.J. and Stock, W.D., 2008. Ecological engineering by a
mega-grazer: white rhino impacts on a South African savanna. Ecosystems, 11,
pp.101-112.

Wiens, J.J., Kuczynski, C.A. and Stephens, P.R., 2010. Discordant mitochondrial
and nuclear gene phylogenies in emydid turtles: implications for speciation and
conservation. *Biological Journal of the Linnean Society*, *99*(2), pp.445-461.

Wang, X., Bernhardsson, C. and Ingvarsson, P.K., 2020. Demography and natural
selection have shaped genetic variation in the widely distributed conifer Norway
spruce (Picea abies). *Genome biology and evolution*, *12*(2), pp.3803-3817.

Zheng, Y., Peng, R., Kuro-o, M. and Zeng, X., 2011. Exploring patterns and extent of
bias in estimating divergence time from mitochondrial DNA sequence data in a
particular lineage: a case study of salamanders (Order Caudata). *Molecular Biology*
and Evolution, *28*(9), pp.2521-2535.

1006

1007

1008

1009 **Figure 1: Sampling locations and potential geographic barriers to elephant**

1010 **dispersal. SSG represents south of Shencottah Gap, SPG represents South of**

1011 **Palghat Gap but north of Shencottah Gap, NPG is North of Palghat Gap, CI**

1012 **represents the Central Indian landscape, NW is north-western India, NE is**

1013 **north-eastern India.**

1014

1015

Figure 2: Population structure based on 2.7 million SNPs (a,b,c) PCA, (d)

ADMIXTURE plot at K=5, and (e) qpgraph with no admixture events with the

branch length qualitatively adjusted to the drift parameter. NW&NE (n=12, sky

red) indicates samples from Northeastern India (n=9) and Northern India (n=3),

CI (n=5, dark blue) from Central India, NPG (n=9, orange) from North of Palghat

Gap, SPG (n=4, green) from South of Palghat Gap but North of Shencottah

Gap, and SSG (n=2, yellow) from South of Shencottah Gap.

 Figure 3: Demographic history (a) recent demographic history within in the last

 100 generations inferred from GONE (Santiago et al, 2020) (b) most likely

 demographic model obtained from fastsimcoal2 (Excoffier et al. 2021) (c)

 population divergence estimated from hPSMC (Cahill et al. 2016). The point

 where population sizes start increasing exponentially in hPSMC plot is the

 time of population divergence. The colours used to represent the populations

 remain the same as earlier, Northwest and Northeast (sky blue), Central India

 (dark blue), North of Palghat Gap (orange), South of Palghat Gap (green) and

 South of Shencottah Gap (yellow).

1036

Figure 4: Genetic diversity revealed as (a) boxplots of pairwise nucleotide
1037

differences per site (pi) in the populations (error bars represent variance in
1038

mean pi/site across scaffolds), and (b) heterozygous SNV encounter rate per
1039

Mb of the genome. The populations are represented, as earlier, the different
1040

colours include Northwest and Northeast (red), Central India (dark blue), North
1041

of Palghat Gap (orange), South of Palghat Gap (green) and South of
1042

Shencottah Gap (yellow).
1043

1044

Figure 5: Present and potential future inbreeding (a) inbreeding measured as
1045

FROH of individuals in each population based on ROH stretches longer than
1046

0.1 Mb (mustard), 1 Mb (blue) and 10 Mb (pink) of genome and (b) identical by
1047

descent (IBD) stretches of the genome longer than 0.1Mb vs 10Mb shared
1048

between pairs of individuals in each population. The populations are
1049

represented, as earlier, the different colours include Northwest and Northeast
1050

(red), Central India (dark blue), North of Palghat Gap (orange), South of Palghat
1051

Gap (green) and South of Shencottah Gap (yellow).
1052

1053

1054

Figure 6: Mutation load measured as a function of homozygosity vs number of
derived (a) loss-of-function (LoF) mutations and (b) indel mutations. The
number of derived deleterious alleles/number of derived neutral alleles is a
proxy for number of deleterious alleles. The error bars indicate standard
deviations. The populations are represented, as earlier, the different colours
include Northwest and Northeast (red), Central (dark blue), North of Palghat
Gap (orange), South of Palghat Gap (green) and South of Shencottah Gap
(yellow).

1063

1064 **Figure7 (a, b). Models tested on fastsimcoal2.7 using scenarios of recent**
1065 **divergence (less than 7,000 ya) and old divergence (more than 17,500 ya).**

1066

a

a