The RecBC complex protects single-stranded DNA gaps during lesion bypass

Gaëlle Philippin[1], Eric Le Cam[2], Élodie Chrabaszcz[1], Maialen Iturralde[1], Mauro Modesti[3], Pauline Dupaigne[2], Vincent Pagès[1,#,*] and Luisa Laureti[1,#,*]

[1]Cancer Research Center of Marseille: Team DNA Damage and Genome Instability | CNRS, Aix Marseille Univ, Inserm, Institut Paoli-Calmettes, Marseille, France

[2]Genome Integrity and Cancers, UMR 9019 CNRS, Université-Paris-Saclay, Gustave Roussy, F-94805 Villejuif Cedex, France.

[3]Cancer Research Center of Marseille: Team Homologous Recombination, NHEJ and Maintenance of Genomic Integrity | CNRS, Aix Marseille Univ, Inserm, Institut Paoli-Calmettes, Marseille, France

Co-last authors

*To whom correspondence should be addressed:

Tel: + 33 486 97 73 84

Fax: + 33 486 97 74 99

Email: vincent.pages@cnrs.fr

Correspondence may also be addressed to: luisa.laureti@cnrs.fr

ABSTRACT

Following encounter with an unrepaired DNA lesion, replication is halted and can restart downstream of the lesion leading to the formation of a single-stranded DNA (ssDNA) gap. To complete replication, this ssDNA gap is filled in by one of the two lesion tolerance pathways: the error-prone Translesion Synthesis (TLS) or the error-free Homology Directed Gap Repair (HDGR). In the present work, we evidence a new role for the RecBC complex distinct from its canonical function in homologous recombination at DNA double strand breaks: RecBC (independently of the RecD subunit) is required to protect the nascent DNA, in order to promote efficient lesion bypass. In the absence of RecBC, our data indicate that the nuclease ExoI can access and degrade the nascent DNA, affecting both TLS and HDGR

mechanism. We show that the recruitment of RecBC becomes particularly important at strong blocking lesions, when post-replicatively ssDNA gaps persist and are covered by the single-stranded DNA binding protein (SSB). This protective role of RecBC is reminiscent of the role of BRCA2 in protecting the nascent DNA in human cells, highlighting once again the evolutionary conservation of DNA replication mechanisms across all living organisms.

INTRODUCTION

DNA replication is constantly challenged by different types of roadblocks that can transiently slow down or stall replication progression. Unrepaired DNA lesions are one of these major roadblocks all organisms have to face. Several works in *Escherichia coli* have demonstrated that in the presence of unrepaired lesions, the progression of the replicative helicase DnaB is not halted (1–3) and replication can restart ahead of the lesion (1, 4, 5), resulting in lesion skipping and the formation of a single-stranded DNA (ssDNA) gap (6–8). In order to complete replication, the ssDNA gap is filled in by one of the two lesion tolerance pathways, identified both in prokaryotes and eukaryotes: 1) Translesion Synthesis (TLS), which employs specialized low-fidelity DNA polymerases that insert a nucleotide directly opposite to the lesion, with the potential to introduce mutations; 2) Homology Directed Gap Repair (HDGR), which uses the information of the sister chromatid to bypass the lesion in a non-mutagenic way through homologous recombination mechanisms (9–11). Regulation of these two pathways is fundamental as it defines the level of mutagenesis during lesion bypass. Our laboratory has previously shown that several factors contribute to regulate the balance between HDGR and TLS: the nature of the DNA lesion (12, 13), the amount of TLS polymerases present in the cell (14) and the density of lesions (15). Very recently, we have also shown that the processing of the ssDNA gap during the presynaptic phase of the RecF pathway plays a role in the regulation of lesion tolerance pathways (16) and that ssDNA gaps persistence at blocking lesions contributes to recruit TLS polymerases (17).

Interestingly, we have previously observed that loss of the *recB* gene has an impact on lesion tolerance pathways, as it leads to a strong decrease in TLS events, and this independently of the nuclease and the RecA loading activities of RecB (13, 18). RecB is part of the RecBCD complex, also known as exonuclease V, a heterotrimeric complex that plays a role in several aspects of cell metabolism and genome maintenance, such as DNA double-strand

break repair, conjugation, chromosomal segregation, degradation of foreign DNA and completion of DNA replication (19–21). In particular, RecBCD is the key factor in homologous recombination pathway at double stranded DNA breaks, also referred to as the RecBCD pathway (22–25). The complex is composed of three subunits: RecB is the main subunit, containing a DNA-dependent ATPase activity, a 3'->5' DNA helicase activity, the nuclease active site and the RecA interacting domain (26). RecC itself does not possess any catalytic activities, but it does bind ssDNA, and can stimulate the ATPase and helicase activities of the RecB protein, with which it forms a complex of 1:1 stoichiometry (27). The main role of RecC is to recognize a specific sequence, the χ (Chi) site (28–30), that participates in the regulation of the complex. RecD is the smallest subunit and possesses a ssDNA-dependent ATPase activity and 5'->3' DNA helicase activity (31, 32). The other functions of RecD are to regulate the nuclease activity of RecB, to increase the processivity of the complex and to negatively regulate RecA loading (33–36). The preferred substrate of the RecBCD complex is a blunt or nearly blunt double stranded DNA ends: the RecB subunit binds to the 3' end while the RecD subunit associates with the 5' extremity (37–39). Individually, each subunit is a poor ATPase, helicase or nuclease, but together they form one of the most processive helicase/nuclease machine (39–41).

Initially, the complex was thought to contain only RecB and RecC subunits (42), and RecD was identified later as the subunit regulating the nuclease activity and containing a helicase activity (43–45). Whereas *recD* null mutants display normal cell viability, resistance to UV light and other DNA damaging agents, and are recombinant proficient, mutations in *recB* or *recC* genes strongly affect cell viability, reduce conjugal recombination 100- to 1000-fold, and sensitize cells to DNA-damaging agents (19). This mild phenotype of the recD mutant is due to the fact that in the *recD* deficient strain, the nuclease activity can be substituted by the nuclease RecJ (45) and that RecA is constitutively loaded, independently of the χ site (34). Since the single *recB* and *recC* mutant strains resulted in identical phenotypes, this suggested that the minimal complex was RecBC. However, *in vitro* RecBC and RecBCD complex do not possess the same enzymatic properties: several works showed that compared to the potent and processive nuclease/helicase RecBCD, RecBC has no exonucleolytic activity on a linear substrate and its endonucleolytic and helicase activities are strongly inhibited. Yet, the DNA-dependent ATPase activity of RecBC is preserved, in particular in the presence of a ssDNA substrate (26, 31, 46). Therefore, it is possible that the two complexes exist in the cell with

different physiological functions, but up to now, no specific cellular role has been associated to the sole RecBC complex.

In this work, we show that the RecBC complex is required to protect the nascent DNA when a persistent ssDNA gap is formed after lesion skipping and repriming. We propose that the RecBC complex, by binding to the SSB-covered ssDNA gap, disorganizes the ssDNA region in a way that will prevent the action of the nuclease ExoI. In the absence of RecBC, ExoI action on the nascent DNA turns out to strongly inhibit TLS polymerases and also to affect HGDR mechanism. Therefore, the RecBC complex is necessary to enable efficient lesion bypass, a new function independent of its catalytic activities and its canonical role in DNA double strand break repair.

MATERIAL AND METHODS

Bacterial strains and growth conditions

All *E. coli* strains used in this work are listed in Supplementary Table 1. They are derivatives of strains FBG151 and FBG152 (12), that carry a plasmid that allows the expression of the *int–xis* genes after IPTG induction. Strains were grown on solid and liquid Lysogeny Broth (LB) medium. Gene disruptions were achieved by the one-step PCR method (47) or by P1 transduction using the Keio collection (48). Following the site-specific recombination reaction, the lesion is located either in the lagging strand (FBG151 derived strains) or in the leading strand (FBG152 derived strains). The mutant alleles of *exoI* ($exoI^{A183V}$) and *recB* ($recB^{F64A}$, $recB^{Y420AF422AR423A}$) have been obtained by CRISPR-Cas9 technology adapted for *E. coli* according to (49). These strains have been verified by sequencing (the entire modified gene). Antibiotics were used at the following concentrations: ampicillin 50 or 100 µg/ml; tetracycline 10 µg/ml, kanamycin 100 µg/ml, chloramphenicol 30 µg/ml. When necessary, IPTG and X-Gal were added to the medium at 0.2mM and 80 µg/ml, respectively.

Plasmids

pVP135 expresses the integrase and excisionase (*int–xis*) genes from phage lambda under the control of a *trc* promoter that has been weakened by mutations in the -35 and the -10 region. Transcription from Ptrc is regulated by the *lac* repressor, supplied by a copy of *lacI^q* on the plasmid. The vector has been modified as previously described (12).

pVP146 is derived from pACYC184 plasmid where the chloramphenicol resistance gene has been deleted. This vector, which carries only the tetracycline resistance gene, serves as an internal control for transformation efficiency.

pVP141-144 and pGP9 are derived from pLDR9-attL-lacZ as described in (12). pLL1 and pLL7 are derived from pVP141 as previously described (13). pAP1, pAP2 and pGP22 are derived from pVP143. All these plasmid vectors contain the following characteristics: the ampicillin resistance gene, the R6K replication origin that allows plasmid replication only if the recipient strain carries the *pir* gene, and the 5' end of the *lacZ* gene in fusion with the *attL* site-specific recombination site of phage lambda. The P'3 site of *att*L has been mutated (AATCATTAT to AAT̲T̲ATTAT) to avoid the excision of the plasmid once integrated. These plasmids are produced in strain EC100D pir-116 (from Epicentre Biotechnologies, cat# EC6P0950H) in which the pir-116 allele supports higher copy number of R6K origin plasmids. Vectors carrying a single lesion for integration were constructed as previously described following the gap-duplex method (12). A 15-mer oligonucleotide 5'-ATCACCGGC̲GCCACA-3' containing or not a single G-AAF adduct (underlined) in the *Nar*I site was inserted into the gapped-duplex pLL1/7 (to measure HDGR) or into the gapped-duplexes pVP141-142 or pVP143-144 to score respectively for TLS0 Pol V-dependent and for TLS-2 Pol II-dependent. A 13-mer oligonucleotide, 5'-GAAGACCT̲GCAGG, containing no lesion or a dG-BaP(-) lesion (underlined) was inserted into the gapped-duplex pVP143/pGP9 to measure Pol IV TLS events. A 14-mer oligonucleotide 5'-ATACCCGG̲GACATC-3' containing no lesion or a single G-AF or a 1hpG adduct (underlined) was inserted into the gapped-duplex pAP1-AP2 to measure TLS like events. A 20-mer oligonucleotide 5'-CTACCT̲GTGGACGGCTGCGA-3' containing or not the N^2-furfuryl adduct was inserted into the gapped duplex pAP1-pGP22 to measure TLS like events. All these new batches of plasmid constructions are carefully quantified and validated by integration in the parental strain.

pVP148 plasmid expresses both *polB* and *umuD'C* loci under their own promoters. It is derived from pRW134 (50), a pGB2 vector that carries *umuD'C*, in which the spectinomycin resistance gene was replaced by the chloramphenicol resistance gene. The *polB* gene was added by subcloning a SalI-BamHI fragment from pYG787 (51). UmuD' is the cleaved version of UmuD that forms the heterotrimeric complex (UmuD')$_2$UmuC, *i.e.*, the active form of Pol V (52).

pLL70 is a plasmid expressing *exoI* gene under its own promoter. It was obtained by removing the *umuDC* gene and its promoter from pVP145 by PstI/BamHI digestion and cloning *exoI*

gene+promoter by PstI/BamHI. pVP145 is derived from pRW134 (50), in which the spectinomycin resistance gene was replaced by the chloramphenicol resistance gene.

Integration protocol: monitoring HDGR and TLS events

The protocol for lesion integration assay is described in details in (18, 53). Cells were diluted and plated before the first cell division using the automatic serial diluter and plater EasySpiral Dilute (Interscience) and were counted using the Scan 1200 automatic colony counter (Interscience).

For every integration experiment, we used the lesion versus non-lesion plasmid constructions, each plasmid solution containing an equal amount of pVP146 plasmid as internal control. Following the integration of the pLL1/7 vector (AAF lesion), sectored blue/white colonies represent HDGR events. Following integration of the vectors pVP141/142 (TLS0 for G-AAF), pVP143/144 (TLS-2 for G-AAF), pVP143/pGP9 (BaP lesion), pAP1-AP2 (G-AF or 1hpG lesions), pAP1-pGP22 (N^2-furfuryl lesion) sectored blue/white colonies represent TLS events. The relative integration efficiencies of lesion-carrying vectors compared with their lesion-free homologues, normalized by the transformation efficiency of pVP146 plasmid in the same electroporation experiment, allow the overall rate of lesion tolerance to be measured (which corresponds to the cell survival in the presence of a single lesion). In the parental strain one lesion gives about 100% cell survival. For the G-AAF lesion we combine the results obtained with the different plasmids to measure HDGR and TLS events. Briefly, the mean of the absolute values (blue colonies/total colonies) of HDGR and TLS events were multiplied by the mean of the cell survival obtained with all the plasmids construction for a given lesion. Indeed, whether TLS or HDGR was measured, cell survival will not change for a given lesion. The value of DCL was calculated by subtracting the HDGR and TLS values to the total cell survival.

Tolerance events (Y axis) represent the percentage of cells able to survive in presence of the integrated lesion compared to the lesion-free control. The data in every graph represent the average and standard deviation of at least three independent experiments of a lesion inserted in the leading (or in the lagging) orientation. Statistical analyses were done using GraphPad Prism applying an unpaired *t*-test.

Sequence alignment

We used Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) for multiple protein sequence alignment to identify the seven conserved motifs specific of SF1 helicases in RecB. For the alignment, we used the Rep and UvrD helicase from *E. coli*, the PcrA helicase from *Staphylococcus aureus* and *Bacillus stearothermophilus* similar to (54, 55).

RecBC protein expression and purification

The protocol for RecBC expression and purification has been adapted from (36). Briefly, *E. coli* RecBC complex was purified from BL21 DE3 Δ*recBD* strain harboring B13 (pETduet-His$_6$-TEVsite-recB, Amp^R) and B3 (pRSFduet-recC, Kan^R) plasmids. The cells were induced with 1mM IPTG at mid-log phase (0.6 OD$_{600}$) and grown for 4h at 37°C. The lysate was purified on HisTrap HP Ni-Sepharose (Cytiva), then dialyzed overnight with TEV protease to cleave the His-tag. The His-tag was removed by a second step on HisTrap HP column. The complex was further purified using HiTrap Heparin-Sepharose (Cytiva), then HiTrap Q FF Sepharose (Cytiva) before continuing with MonoQ (Cytiva) anion exchange chromatography. The protein was dialyzed overnight in 20 mM Tris-HCl pH8, 1mM EDTA pH 8, 1mM DTT, 50mM NaCl, 10% glycerol and flash-frozen. The expression and purification of the RecB^{F64A}C mutant was obtained in the same way.

Preparation of the gapped DNA substrates

Plasmid with a 5.3 kb ssDNA gap: 3 pmol of a 50nt oligonucleotide (VP683, TCGAGCTGCGCAAGGATAGGTCGAATTTTCTCATTTTCCGCCAGCAGTCC) was 5' radiolabelled using 3.3 pmol of γ-^{32}P-ATP (Perkin Elmer) and 10 units of PNK (NEB) in a total volume of 25μl PNK buffer 1X. The reaction was incubated during 30 minutes at 37°C and inactivated during 20 minutes at 65°C. Then it was purified using a MicroSpin G-25 column (Cytiva). The oligonucleotides (60nM) were hybridized to φX174 ssDNA plasmids (5.3 kb, NEB) (20nM) in 50 μl buffer containing 10 mM Tris-HCl pH8, 1mM EDTA pH8 and 50mM NaCl. The reaction was incubated in water at 75°C and let cool down to room temperature. Substrates were checked on 0.8% agarose gel.

Plasmid with a 2.5 kb ssDNA gap: 10 μM VP 1394 oligonucleotides (ACGATTAACCCTGATACCAA) were phosphorylated with 1mM ATP (Sigma) and 10 units PNK (NEB) in 50 μl PNK buffer 1X (30' at 37°C, inactivation 20' at 65°C). A 2.8 kb dsDNA fragment

was obtained by PCR with Q5 Polymerase (NEB) using phosphorylated VP 1394, VP 1361 (ACGCCCTGCATACGAAAAGA) and φX174 as template. PCR products were digested by ExoI (NEB) to remove excess of oligos and purified using Qiagen PCR purification kit. The phosphorylated DNA strand was then specifically digested with lambda exonuclease (NEB); the ssDNA strands obtained were checked on 0.8% agarose gel. They were 5'-radiolabelled using γ-^{32}P-ATP and purified using a MicroSpin G-25 column. Then they were hybridized to φX174 ssDNA plasmids (ratio 1 :1) in 50 µl buffer containing 10 mM Tris-HCl pH8 and 50 mM NaCl. The reaction was incubated in water at 80°C and let cool down to room temperature. Substrates were checked on 0,8% agarose gel.

This substrate (non radio-labelled) was used also for TEM analysis.

Electrophoretic mobility shift assays

The RecBC complex (and the mutant allele RecB^{F64A}C) was freshly expressed and purified since the RecBC complex proved to have a reduced shelf life as also mentioned by (46). The SSB protein was provided by BioAcademia, and the truncated SSB ΔC8 protein was kindly provided by Michael Cox (University of Wisconsin-Madison).

When using the 5.3 kb ssDNA gap substrate, 4 fmol of radiolabelled DNA substrates were first incubated for 20' at RT with increasing concentrations of SSB (0-11-22-43 nM) in 12 µl binding buffer (20 mM Tris-HCl pH8, 100 µg/ml BSA-Ac, 1mM DTT and 1mM Mg-Ac). Then 160 nM RecBC (or RecBC buffer as a negative control) was added to the reaction and incubated for 20' at RT. 1X loading dye was added and the samples were loaded on 0.6% agarose gel, running at 4°C. The gel was dried for 1h30 at 60°C, exposed overnight to a Fuji screen and scanned with a Typhoon (Cytiva).

The same reaction conditions were used in Fig 5D except for SSB and SSB ΔC8 concentration (78 nM).

When using the 2.5 kb ssDNA gap substrate 15 fmol radiolabelled DNA substrates were first incubated for 20' at RT with SSB (0-130-260-520 nM) in 10 µl binding buffer. Then 400 nM RecBC (or RecBC buffer as a negative control) was added to the reaction and incubated for 20' at RT. 1X loading dye was added and the samples were loaded on 0.6% agarose gel, running at 4°C. The gel was dried for 1h30 at 60°C, exposed overnight to a Fuji screen and scanned

with a Typhoon (Cytiva). The same reaction conditions were used in Fig 5C with 433 nM SSB and 500 nM RecBC/RecBC[F64A].

Transmission Electron Microscopy (TEM)

The DNA substrate containing a 2.5 kb ssDNA gap (18 µM nucleotides for ssDNA part) was first incubated in non-saturating conditions with 200 nM SSB protein during 10' at RT in a binding buffer containing 10mM Tris-HCl pH 7.5, 50 mM NaCl, 2 mM $MgCl_2$ and 1mM DTT. The resulting SSB-ssDNA gap complex was then incubated with 100 nM RecBC protein in the same binding buffer during 10' at RT.

Complexes formed with or without RecBC were quickly diluted 40 times in the binding buffer then spread onto the TEM grids as following: 5 µL were immediately deposited for 1 minute on a 600-mesh copper grid coated with a thin carbon film, preactivated by glow-discharge in the presence of amylamine (Sigma-Aldrich) in a homemade device as previously described (56). The grids were washed with 0.2% (w/vol) aqueous uranyl acetate (Merck) and dried with ashless filter paper (VWR). The grids were directly subjected to TEM analysis in filtered annular dark-field mode with a Zeiss 902 microscope using Veletta high-resolution CCD camera and iTEM software (Olympus, Soft Imaging Solutions). TEM images were acquired at a magnification of x 140000.

RESULTS

We have previously shown that in the presence of a UV photoproduct lesion (*i.e.* TT6-4), deletion of *recB* gene affects lesion tolerance pathways, decreasing HDGR mechanisms and strongly inhibiting the TLS pathway (13, 18). We have also shown that this role in lesion bypass is independent of the nuclease and the RecA loading activities of RecB and more interestingly, that the RecB contribution to lesion bypass does not involve processing of a DNA double strand break that could potentially arise in the presence of a ssDNA gap (18). In this study, we further clarify this unexpected role of RecB.

To monitor lesion tolerance mechanisms, we took advantage of our previously described genetic assay (see Suppl Fig 1A) that allows us to insert a single replication-blocking lesion into the chromosome of *E. coli,* either on the lagging or in the leading strand orientation regarding the replication fork direction*.* TLS or HDGR events are visualized and measured by a colorimetric assay based on the *lacZ* reporter gene using specific lesion integration plasmids (Suppl Fig 1B and 1C) (12, 13, 53). This assay had allowed us also to reveal an additional lesion tolerance strategy, that we named Damaged Chromatid Loss (DCL), which promotes cell proliferation by replicating only the undamaged chromatid at the expense of losing the damaged chromatid (13) (see Suppl Fig 1C). In the present study, we mainly used the chemical guanine adduct N-2-acetylaminofluorene (G-AAF) positioned in the *Nar*I hotspot sequence: in this configuration the lesion can be bypassed either by the TLS polymerase Pol V in an error-free manner (indicated as TLS0 events), or by the TLS polymerase Pol II, inducing a -2 frameshift (indicated as TLS-2 events) (57).

The *E. coli* strains used are deficient for the mismatch repair system (*mutS*) to prevent correction of the genetic markers of the integrated plasmids, as well as for nucleotide excision repair (*uvrA*), to avoid excision of the lesion and to focus on lesion tolerance events.

RecBC complex but not RecD participates in lesion bypass

Since RecB is part of the RecBCD complex, we decided to test whether all the subunits of the complex were involved in lesion bypass. We inactivated separately *recB, recC* and *recD* genes and measured tolerance events (*i.e.* HDGR, TLS and DCL) in the presence of a G-AAF lesion. Inactivation of *recB* gene resulted in a decrease of HDGR, loss of Pol II TLS events (from 2% to <0.4%) and an increase of DCL events (Fig 1), confirming the previous results obtained

with the UV lesion (13, 18). A *recC* deficient strain behaved like the *recB* strain for lesion tolerance events (Fig 1). On the other hand, no decrease in HDGR or TLS was measured upon inactivation of *recD* (Fig 1), indicating that only the RecBC complex is required for lesion bypass. RecD subunit is involved in the regulation of the nuclease activity of RecB (33, 35). Since we have already shown that the nuclease dead allele of RecB (RecB[D1080A]) has no effect on HDGR or TLS (18), it is not surprising that RecD subunit is not required.

Further confirmation that RecBC, without RecD, is involved in lesion bypass came using a plasmid construction containing two lesions on opposite strands: this configuration induces a two-fold increase in TLS events due to the generation of overlapping ssDNA gaps that prevent HDGR (15). Deletion of the *recB* gene had a strong effect on TLS levels, while the deletion of *recD* gene or the nuclease dead allele of *recB* behaved in a manner similar to the parental strain (See Suppl Figure 2).

Exonuclease screen in the *recB* deficient strain

In the absence of RecBC, HDGR mechanism is decreased and TLS is severely impaired. From these data, we hypothesized that RecBC could be involved in the protection of the nascent DNA blocked at the lesion site from nuclease degradation, similar to what has been described in human cells where BRCA2, together with RAD51, protects the replication fork against the MRE11 nuclease attack (58, 59). Degradation of the 3' end of the nascent DNA is detrimental for TLS polymerases since their cognate substrate (*i.e.* the 3' end blocked at the lesion site) is missing. In addition, important degradation of the nascent DNA will also impact the efficiency of HDGR mechanism. To corroborate our hypothesis, we performed a genetic screen to identify the putative nuclease(s) responsible for this degradation in the absence of *recB*. We looked for candidate genes that encode exonucleases with a 3'->5' polarity that preferentially degrade single-stranded DNA and that have a role in DNA metabolism and genome maintenance. Based on the literature (see reviews (60, 61)), we chose to test ExoI (also known as XonA/SbcB), ExoVII, ExoX and SbcCD.

We inactivated each exonuclease individually in the *recB* deficient strain and measured Pol II dependent TLS events in the presence of a G-AAF lesion in order to see whether we could restore the level of the parental strain. From this first screen, ExoI and SbcCD emerged as good candidates, since deletion of *exoI* or *sbcCD* in the *recB* strain recovered fully or partially the level of TLS-2 (Fig 2A).

ExoI prevents efficient lesion bypass in the absence of RecBC

Previously, we have shown that the deletion of *recB* affects lesion bypass of all three TLS polymerases (18). As mentioned above, when positioned in the *Nar*I hotspot sequence, the G-AAF lesion can be bypassed by Pol II, but also by Pol V. However, under physiological conditions, Pol V bypass is very low (<0.5%) (12). Hence, to assess the role of ExoI and SbcCD on Pol V TLS events, we used a plasmid expressing both the active form of Pol V (UmuD'C in which UmuD is already cleaved (52, 62)) and Pol II under their own promoters (14). The expression level of both TLS polymerases is 3-5 times higher than in the parental strain (due to plasmid copy number), which results in an increase of Pol II TLS events (from 2% to 12%) and of Pol V TLS events (from 0.5% to more than 20%) (Fig 2B). These conditions allow a better dynamic range to monitor the effect of the nucleases on both Pol V and Pol II-mediated TLS. As expected, deletion of the *recB* gene strongly decreased the level of both Pol II and Pol V dependent TLS events (Fig 2B). While deletion of the nuclease *sbcCD* in the *recB* strain did not restore TLS levels, the double mutant *exoI recB* fully or partially recovered Pol II and Pol V dependent TLS events, respectively (Fig 2B).

Finally, we tested the effect of the two nucleases on the TLS polymerase Pol IV in the presence of the benzo(a)pyrene (BaP) lesion (63). Only deletion of *exoI* in the *recB* deficient strain was able to almost fully restore the TLS level of the parental strain (Fig 2C), thus pointing ExoI as the major nuclease acting in the absence of RecBC.

Since loss of RecBC also affects HDGR mechanism (Fig 1), we measured HDGR level in an *exoI recB* strain in the presence of a G-AAF lesion and observed full recovery of HDGR level similar to the parental strain (Suppl Fig 3A). Hence, from these data it appears that the defect in lesion bypass observed in the absence of RecBC can be largely alleviated by inactivation of ExoI.

RecBC protects from ExoI exonuclease

ExoI is a Mg^{2+}-dependent exonuclease that rapidly processes ssDNA in the 3'->5' direction (64). It is a highly processive nuclease whose activity is stimulated by the single stranded DNA binding protein (SSB) (65). ExoI plays several roles in DNA metabolism and DNA repair, in particular in mismatch repair, recombination and completion of DNA replication (21, 66–68). Hence, we pursued the characterization of the role of ExoI in lesion bypass. As expected, complementation experiments in the *exoI recB* strain, using a plasmid carrying a

functional *exoI* gene under its native promoter, resulted in low levels of Pol II TLS events in the presence of a G-AAF lesion similar to the one observed in a *recB* deficient strain (Fig 2A: *exoI recB*+p*ExoI*).

To reinforce our hypothesis that ExoI is the exonuclease preventing efficient lesion bypass in the absence of RecBC, we used the well-described mutant allele *sbcB15* (*exoIA183V*), which is inactivated in its nuclease active site (68). We introduced the point mutation in the *exoI* gene by CRISPR-Cas9 technology in the *recB* deficient strain and measured the level of Pol II TLS in the presence of a G-AAF lesion. We observed full recovery of Pol II TLS, similar to the parental strain (Fig 2A: *exoIA183V recB*)).

Taken together these data indicate that in the absence of RecBC, ExoI is the major exonuclease that negatively affects lesion bypass. Given that its exonuclease catalytic activity is required, we surmise that it acts by degrading the nascent DNA that presents a 3' extremity blocked at the lesion site. We wondered whether in the presence of RecBC, ExoI could still access to the 3' end of the nascent DNA, thus affecting lesion bypass. We inactivated *exoI* gene in the parental strain and measured Pol II TLS events and HDGR in the presence of a G-AAF lesion. No change in the level of Pol II TLS events, nor in HDGR events were observed compared to the parental strain (Fig 2A and Suppl Fig 3A) suggesting that in the presence of RecBC, ExoI has no effect on lesion bypass.

ExoI and the TLS polymerases are in competition for the same substrate

Based on these genetic data, we proposed that the RecBC complex protects the nascent DNA blocked at the lesion site from the attack of the ExoI nuclease, in order to allow efficient lesion bypass, either by TLS polymerases or by HDGR mechanism. In the absence of RecBC, ExoI can access the 3' end of the nascent DNA, degrading it and affecting lesion bypass. According to this model, in the absence of RecBC, the action of ExoI would be in competition with lesion tolerance pathways, in particular with the TLS polymerases since their cognate substrate (the 3' end) is the same. Therefore, overexpression of the TLS polymerases should partially compensate for the absence of RecBC. Indeed, we can see that TLS events in a *recB* strain are much higher when Pol II and Pol V are overexpressed compared to the *recB* strain with basal level of Pol II and Pol V (Fig 2A and 2B). Further confirmation came from testing both Pol II and Pol V TLS levels in a strain deficient for both *recB* and *lexA*: in such strain, constitutive SOS induction leads to a strong increase in the expression level of TLS

polymerases that in turn lead to a 15-fold increase of both TLS events (TLS-2 and TLS0) compared to a *recB* strain (Suppl Fig 3B).

RecBC needs to bind to the DNA to protect the nascent DNA

RecB is a 134 kDa protein composed of two parts, the N-terminal region belonging to the superfamily 1 (SF1) helicase and the C-terminal region containing the nuclease activity (33, 69) and the RecA interaction motif (34, 70). The two regions are connected by a 70 amino acid linker (see Fig 3A). We previously showed that neither the nuclease activity nor the RecA loading activity of RecB were required for lesion bypass. Indeed, the *recB*[D1080A] mutant allele behaved as the parental strain for both TLS and HDGR pathways and the expression of the *recA730* allele, that does not require a mediator to load itself on ssDNA, did not restore the level of lesion tolerance pathways in the absence of RecB (see (18) and Fig 3B). Therefore, we decided to investigate whether the SF1 helicase core of the protein could be involved in the protection of the nascent DNA.

SF1 helicases are composed of a four-subdomain structure (1A, 1B, 2A, 2B) that contains seven conserved motifs (54, 55) (See Figure 3A): motif I contains the Walker A box responsible for binding the triphosphate tail of ATP; motif II is the "DEAD" box motif responsible for Mg+ binding, required for ATP hydrolysis; motif III, IV and VI are involved in ATP hydrolysis and in coupling ATP hydrolysis to DNA translocation (54, 71, 72); motif Ia, III and V were shown to contain residues involved in DNA binding and ssDNA translocation (55, 72, 73).

In *E. coli* the UvrD and Rep helicases are representative of this SF1 family. We performed multiple protein sequence comparison with these two helicases as well as other well studied bacterial SF1 helicases and identified in RecB all the characteristic SF1 motifs and most of the conserved amino acids responsible for the binding of NTPs or DNA (see Fig 3A). We hypothesized that RecBC needs to bind DNA in order to protect it from ExoI degradation. Hence, we mutated specific amino acid residues of RecB presumably involved in DNA binding and monitor the effect on lesion bypass. Based on several crystal structure studies, residues in Motif Ia and III seemed to be the most critical for ssDNA binding (55, 72, 73), therefore we generated the mutant allele *recB*[F64A], in which the phenylalanine 64 of Motif Ia is mutated to alanine, and the mutant allele *recB*[YFR-AAA], in which tyrosine 420, phenylalanine 422 and arginine 423 of Motif III have all been mutated to alanine (see Fig 3A). We monitored Pol II

TLS events in these mutant alleles in the presence of a G-AAF lesion: TLS decreased to a level similar to the *recB* deficient strain (Fig 3B), indicating that these residues are required for the protective role of RecBC. We corroborated these results using the plasmid that overexpresses Pol II in the *recB^YFR-AAA* strain. In this mutant, despite the increased level of Pol II, the level of TLS-2 events remains low, similar to the *recB* deficient strain expressing the same plasmid (Fig 3C), strongly supporting the hypothesis that RecBC needs to interact with the DNA.

RecBC is required in the presence of strong blocking lesions

We have shown the requirement for RecBC in the presence of strong blocking lesions such as G-AAF and UV lesions (13, 18), when a ssDNA gap is formed downstream of the lesion. When replication is stalled by a less blocking lesion, bypass can occur without the need for replication to restart downstream of the lesion: in this situation, no ssDNA gap will form as the lesion will be rapidly bypassed by the replicative polymerase or a TLS polymerase. Therefore, we asked whether in this situation RecBC was still required to protect the nascent DNA. For this purpose, we used three less blocking lesions, the amino fluorene (AF) (74), the hydroxypropyl (1hpG) (75) and the N2-G-furfuryl (17). As shown in Fig 4, TLS rate at these lesions is higher than for the strong blocking lesions such as the G-AAF or the UV photoproducts (12). In the absence of RecBC, less than two-fold decrease in lesion bypass was observed for the three lesions (Fig 4), while we have measured a 7-fold decrease in the absence of RecBC when stronger blocking lesions were used (see (18) and Fig 2). These data indicated that RecBC requirement is more critical in the presence of a persistent ssDNA gap: when the 3' end of the nascent DNA is exposed longer, its protection becomes critical to allow TLS and HDGR mechanisms.

RecBC complex binds to ssDNA gap in the presence of SSB

Our genetic data have shown that the amino acids presumably responsible for the DNA binding are necessary for efficient lesion bypass (Fig 3B and 3C). In order to confirm and support our model, we performed *in vitro* assays to test whether RecBC was indeed binding to a ssDNA gap.

We constructed two circular gapped DNA substrates using the phage phiX174 (ϕX174), with two different ssDNA gaps of 5.3 kb and 2.5 kb (Fig 5A). In parallel, we expressed and

purified the RecBC complex according to (36) (Suppl Fig. 4A). As mentioned before, the RecBC complex was shown to lack exonuclease activity while still preserving a weak helicase activity (46, 76, 77). Our purified RecBC complex was indeed not able to degrade a linear blunt DNA substrate but we observed a weak helicase activity with the same DNA substrate (Suppl Fig 4B and 4C).

We then performed electrophoretic mobility shift assays using our gapped substrates. When using the RecBC complex alone, no shift was observed with either substrate (Fig 5B). However, when we added SSB to the reaction, the RecBC complex produced an additional band shift with both DNA substrates (Fig 5B). The binding of RecBC in the presence of SSB is specific to the ssDNA gap, since no binding was observed with a circular ssDNA plasmid even in the presence of SSB (Suppl Fig 4D). To further confirm that RecBC was binding to the ssDNA gap, we performed the same electrophoretic mobility shift assay using the RecB^{F64A} allele that is supposed to be defective for DNA binding and that showed to be impaired in Pol II TLS bypass *in vivo* (Fig 3B). The RecB^{F64A}C complex was unable to produce an additional shift in the presence of SSB, indicating that the complex could not bind the ssDNA gap as expected (Fig 5C).

The acidic tip of SSB is not required for RecBC binding to DNA gaps

To characterize the cooperative binding of RecBC with SSB, we asked whether there was a direct interaction between RecBC and SSB. Indeed, bacterial SSBs are known to be a hub for protein recruitment, in particular their C-terminal tail contains several amino acids (also called the "acidic tip") responsible for the binding of several proteins involved in recombination, repair and replication (78–83). We performed mobility shift assay, using a well-described truncated form of SSB (SSBΔ8) that lacks the last 8 amino acids, thus affecting most of protein-protein interactions (84), but not the DNA binding capacity (78). In the presence of the SSBΔ8, the RecBC complex was still able to bind to the substrate (Fig 5D), suggesting no direct interaction between the two proteins via this acidic tip. However, we cannot completely rule out the possibility that RecBC does interact with SSB, in particular with its intrinsically disordered linker (IDL) domain since it has been shown recently that this part of the protein might also be involved in protein-protein interactions (82, 83, 85). Our data showing that RecBC cannot bind to a circular ssDNA plasmid even in the presence of SSB (Suppl. Fig 4D) seems to support that no direct binding between RecBC and SSB is required. It

appears therefore that it is more a structural conformation created by the binding of SSB to the ssDNA gap that is required for RecBC binding.

RecBC binding disrupts the SSB-covered ssDNA gap structure

To protect the nascent DNA, we hypothesized that the RecBC complex binds to the double strand-single stranded (ds-ss) DNA junction at the lesion site. To address this question, we performed transmission electron microscopy (TEM) imaging using our gapped DNA substrate (2.5kb ssDNA gap) (Fig 6A) and our purified proteins. Incubation of non-saturating concentrations of SSB, similar to what was used for the previous *in vitro* electrophoretic mobility shift assays (Fig 5), resulted in SSB binding, partially covering the ssDNA gap of the substrate and creating a nicely folded SSB filament, as expected (Fig 6B). Surprisingly, the addition of RecBC to the reaction induced tangled and disorganized structures of the SSB-covered ssDNA substrate (Fig 6C and 6D). This result suggests that instead of binding to the ds-ssDNA junction, RecBC binds internally to the ssDNA gap, thus changing the dynamic and the structure of the SSB filament. These entangled structures are probably not a suitable substrate for the nuclease ExoI, thus preventing its action and preserving the nascent DNA.

DISCUSSION

This study unveils a cellular role for the RecBC complex, without the RecD subunit, in protecting the nascent DNA to allow efficient lesion bypass in the presence of a single-stranded DNA gap. Indeed, in the absence of RecBC lesion bypass is severely compromised, most likely because degradation of the nascent DNA prevents the action of the TLS polymerases and also affects the efficiency of the HDGR mechanism. While we have recently shown that the ssDNA gap needs to be enlarged on its 5' end by the action of the nuclease RecJ and the helicase RecQ before the mediator proteins recF-O-R load RecA (16), the present study shows that the 3' end of the gap needs to be protected from nucleolytic degradation operated by the exonuclease ExoI in order to allow an efficient lesion bypass and complete the replication of a damaged DNA.

RecBC protects the 3' end of persistent ssDNA gaps

Based on our *in vivo* and *in vitro* results, we propose a model (Fig 7) in which RecBC is required to protect the nascent DNA when a ssDNA gap is formed and persists. This situation is typical of a strong roadblock when replication needs to restart downstream in order to continue replication, creating a persistent ssDNA gap that will be later filled in by one of the two lesion tolerance pathways. The ssDNA gap is immediately covered by the SSB protein that protects the ssDNA and orchestrates the recruitment of several proteins to preserve genome integrity. We propose that, once SSB covers the ssDNA gap, RecBC complex(es) binds to the SSB filament and disrupts its organized folding, creating tangled structures that protects the nascent DNA from the action of the nuclease ExoI, allowing an efficient lesion bypass. On the other hand, when replication is stalled by a less blocking lesion that can be easily bypassed by either the replicative polymerase or a TLS polymerase, the ssDNA created by the DNA unwinding is only transient and RecBC protection is not necessary.

SSB is a constitutive component of the replisome, associating with the ssDNA on the lagging strand during Okazaki fragment formation or ssDNA created during the DNA helicase unwinding. However, the SSB binding lifetime to these "physiological" ssDNA gaps and the SSB binding lifetime to the ssDNA gaps formed upon lesion encountering are probably different, as recently suggested by Loparo's group (86). The same group also suggested that the mobility change in SSB molecules upon replication perturbation could act as a molecular switch to enrich proteins at the lesion site, as it is the case for Pol IV TLS polymerase (17, 86). Hence, this difference in SSB dynamics could explain why RecBC is specifically recruited to protect the nascent DNA of the ssDNA gap formed in the presence of strong blocking lesions and not in the presence of a less blocking lesion that generates more transient gaps. Our *in vitro* reconstitution experiments indicate no direct interaction between SSB and RecBC, corroborating the idea that the key for RecBC recruitment is more the dynamics and the mobility change of SSB.

Degradation of nascent DNA strongly affects lesion tolerance pathways

In the absence of RecBC, the SSB filament folding onto the ssDNA gap is not altered and therefore the nascent DNA is deprotected. We propose that in this situation ExoI can access the nascent DNA (probably with the help of a still to be identified helicase) and degrade it, thus strongly inhibiting TLS polymerases access/activity since ExoI and the TLS polymerases are in competition for the same substrate (the 3' end). In line with this, overexpression of the

TLS polymerases can partially outcompete the action of ExoI. Degradation of the nascent DNA also affects HDGR leading to an increase in damage chromatid loss events (DCL, Fig 1). However, high levels of HDGR still persist in the absence of RecBC, suggesting that not all the nascent DNA is degraded by ExoI, or that the action of ExoI is also kinetically in competition with the RecA filament formation. Indeed, it is likely that once the recombinogenic filament is formed, it will engage in homology search and strand invasion and ExoI will not be able to access the nascent DNA. This is corroborated by the phenotype of the double mutant *recF recB* in a previous study (13) where we observed a strong loss of HDGR and cell survival (similar to a *recA* deficient strain). Delaying the formation of the RecA filament (in the absence of its mediator RecF) in addition to the loss of RecBC will favor ExoI activity, thus degradation of the nascent DNA will strongly impact HDGR mechanisms.

In the last decade, growing evidence indicate that the nascent DNA must be protected against uncontrolled and excessive nucleolytic degradation, and failure to do so might severely compromise the completion of DNA and the stability of the genome (87, 88). Surprisingly, proteins usually involved in homologous recombination pathway, such as RAD51 and BRCA1/2 in humans, came to the fore as guardians of the nascent DNA against the attack of specific nucleases (59, 89, 90). This function is distinct from their canonical function in homologous recombination. Similarly, in this study we have shown that RecBC, that is part of the RecBCD complex involved in homologous recombination repair in *E. coli* and is one of the mediators of RecA, plays a new and unexpected role in lesion bypass, independent of its canonical catalytic activities. This new finding indicates once more how the mechanisms of DNA replication are evolutionary conserved in all living organisms. It will be interesting to investigate in human cells whether lesion bypass may also be compromised in cells deficient for the so-called fork protection when a ssDNA gap is formed.

DATA AVAILABILITY

All relevant data are within the manuscript or in the Supplementary files and are available on request.

FUNDING

This work was supported by Fondation pour la Recherche Médicale [Equipe FRM-EQU201903007797] https://www.frm.org (VP), by ANR (17-CE12-0015) and ERM-Université Paris-Saclay (ELC and PD). Funding for open access charge : Fondation pour la Recherche Médicale. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

ACKNOWLEDGEMENTS

We thank Jean-Hugues Guervilly for critical reading of the manuscript. We thank Dave Wigley for kindly providing us the plasmids for RecBC expression/purification and Michael Cox for kindly providing us the SSBΔ8 protein.

CONFLICT OF INTEREST

none

REFERENCES

1. Yeeles,J.T.P. and Marians,K.J. (2013) Dynamics of leading-strand lesion skipping by the replisome. *Molecular Cell* **52**, 855-865.
2. Beattie,T.R., Kapadia,N., Nicolas,E., Uphoff,S., Wollman,A.J., Leake,M.C. and Reyes-Lamothe,R. (2017) Frequent exchange of the DNA polymerase during bacterial chromosome replication. *Elife* **6**, e21763.
3. Yang,W., Seidman,M.M., Rupp,W.D. and Gao,Y. (2019) Replisome structure suggests mechanism for continuous fork progression and post-replication repair. *DNA Repair (Amst)* **81**, 102658.
4. Heller,R.C. and Marians,K.J. (2006) Replication fork reactivation downstream of a blocked nascent leading strand. *Nature* **439**, 557-562.
5. Yeeles,J.T.P. and Marians,K.J. (2011) The Escherichia coli replisome is inherently DNA damage tolerant. *Science (New York, N.Y.)* **334**, 235-238.
6. Henrikus,S.S., Wood,E.A., McDonald,J.P., Cox,M.M., Woodgate,R., Goodman,M.F., van Oijen,A.M. and Robinson,A. (2018) DNA polymerase IV primarily operates outside of DNA replication forks in Escherichia coli *PLoS genetics* **14**, e1007161.
7. Quinet,A., Tirman,S., Cybulla,E., Meroni,A. and Vindigni,A. (2021) To skip or not to skip: choosing repriming to tolerate DNA damage. *Mol Cell*
8. Pham,P., Shao,Y., Cox,M.M. and Goodman,M.F. (2022) Genomic landscape of single-stranded DNA gapped intermediates in Escherichia coli. *Nucleic Acids Res* **50**, 937-951.
9. Branzei,D. and Psakhye,I. (2016) DNA damage tolerance. *Current opinion in cell biology* **40**, 137-144.
10. Henrikus,S.S., van Oijen,A.M. and Robinson,A. (2018) Specialised DNA polymerases in Escherichia coli: roles within multiple pathways *Current genetics* **64**, 1189-1196.
11. Ashour,M.E. and Mosammaparast,N. (2021) Mechanisms of damage tolerance and repair during DNA replication *Nucleic Acids Research* **49**, 3033-3047.

12. Pagès,V., Mazón,G., Naiman,K., Philippin,G. and Fuchs,R.P. (2012) Monitoring bypass of single replication-blocking lesions by damage avoidance in the Escherichia coli chromosome *Nucleic acids research* **40**, 9036-9043.

13. Laureti,L., Demol,J., Fuchs,R.P. and Pagès,V. (2015) Bacterial proliferation: keep dividing and don't mind the gap *PLoS genetics* **11**, e1005757.

14. Naiman,K., Philippin,G., Fuchs,R.P. and Pagès,V. (2014) Chronology in lesion tolerance gives priority to genetic variability. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 5526-5531.

15. Chrabaszcz,É., Laureti,L. and Pagès,V. (2018) DNA lesions proximity modulates damage tolerance pathways in Escherichia coli *Nucleic acids research* **46**, 4004-4012.

16. Laureti,L., Lee,L., Philippin,G., Kahi,M. and Pagès,V. (2022) Single strand gap repair: The presynaptic phase plays a pivotal role in modulating lesion tolerance pathways. *PLoS Genet* **18**, e1010238.

17. Chang,S., Thrall,E.S., Laureti,L., Piatt,S.C., Pagès,V. and Loparo,J.J. (2022) Compartmentalization of the replication fork by single-stranded DNA-binding protein regulates translesion synthesis. *Nat Struct Mol Biol* **29**, 932-941.

18. Laureti,L., Lee,L., Philippin,G. and Pagès,V. (2017) A non-catalytic role of RecBCD in homology directed gap repair and translesion synthesis. *Nucleic Acids Research* **45**, 5877-5886.

19. Arnold,D.A. and Kowalczykowski,S.C. (2001) RecBCD Helicase/Nuclease *Encyclopedia of life sciences* 1-6.

20. Dillingham,M.S. and Kowalczykowski,S.C. (2008) RecBCD enzyme and the repair of double-stranded DNA breaks. *Microbiology and molecular biology reviews : MMBR* **72**, 642-671.

21. Wendel,B.M., Courcelle,C.T. and Courcelle,J. (2014) Completion of DNA replication in Escherichia coli. *Proceedings of the National Academy of Sciences* **111**, 16454-16459.

22. Kowalczykowski,S.C., Dixon,D.A., Eggleston,A.K., Lauder,S.D. and Rehrauer,W.M. (1994) Biochemistry of homologous recombination in Escherichia coli. *Microbiological reviews* **58**, 401-465.

23. Kuzminov,A. (1999) Recombinational repair of DNA damage in Escherichia coli and bacteriophage lambda. *Microbiology and molecular biology reviews : MMBR* **63**, 751-813.

24. Spies,M. and Kowalczykowski,S.C. (2005) Homologous recombination by RecBCD and RecF pathways *The bacterial chromosome* 389-403.

25. Amundsen,S.K. and Smith,G.R. (2023) RecBCD enzyme: mechanistic insights from mutants of a complex helicase-nuclease. *Microbiol Mol Biol Rev* e0004123.

26. Hickson,I.D., Robson,C.N., Atkinson,K.E., Hutton,L. and Emmerson,P.T. (1985) Reconstitution of RecBC DNase activity from purified Escherichia coli RecB and RecC proteins. *Journal of Biological Chemistry* **260**, 1224-1229.

27. Masterson,C., Boehmer,P.E., McDonald,F., Chaudhuri,S., Hickson,I.D. and Emmerson,P.T. (1992) Reconstitution of the activities of the RecBCD holoenzyme of Escherichia coli from the purified subunits. *Journal of Biological Chemistry* **267**, 13564-13572.

28. Handa,N., Ohashi,S., Kusano,K. and Kobayashi,I. (1997) Chi-star, a chi-related 11-mer sequence partially active in an E. coli recC1004 strain. *Genes Cells* **2**, 525-536.

29. Arnold,D.A., Handa,N., Kobayashi,I. and Kowalczykowski,S.C. (2000) A novel, 11 nucleotide variant of chi, chi*: one of a class of sequences defining the Escherichia coli recombination hotspot chi. *J Mol Biol* **300**, 469-479.

30. Handa,N., Yang,L., Dillingham,M.S., Kobayashi,I., Wigley,D.B. and Kowalczykowski,S.C. (2012) Molecular determinants responsible for recognition of the

single-stranded DNA regulatory sequence, χ, by RecBCD enzyme. *Proc Natl Acad Sci U S A* **109**, 8901-8906.

31. Amundsen,S.K., Neiman,A.M., Thibodeaux,S.M. and Smith,G.R. (1990) Genetic dissection of the biochemical activities of RecBCD enzyme. *Genetics* **126**, 25-40.

32. Chamberlin,M. and Julin,D.A. (1996) Interactions of the RecBCD Enzyme from Escherichia coliand Its Subunits with DNA, Elucidated from the Kinetics of ATP and DNA Hydrolysis with Oligothymidine Substrates † *Biochemistry* **35**, 15949-15961.

33. Yu,M., Souaya,J. and Julin,D.A. (1998) Identification of the nuclease active site in the multifunctional RecBCD enzyme by creation of a chimeric enzyme. *Journal of Molecular Biology* **283**, 797-808.

34. Churchill,J.J., Anderson,D.G. and Kowalczykowski,S.C. (1999) The RecBC enzyme loads RecA protein onto ssDNA asymmetrically and independently of χ, resulting in constitutive recombination activation *Genes & development* **13**, 901-911.

35. Amundsen,S.K., Taylor,A.F. and Smith,G.R. (2000) The RecD subunit of the Escherichia coli RecBCD enzyme inhibits RecA loading, homologous recombination, and DNA repair. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 7399-7404.

36. Wilkinson,M., Chaban,Y. and Wigley,D.B. (2016) Mechanism for nuclease regulation in RecBCD *Elife* **5**, e18227.

37. Taylor,A.F. and Smith,G.R. (1985) Substrate specificity of the DNA unwinding activity of the RecBC enzyme of Escherichia coli. *Journal of Molecular Biology* **185**, 431-443.

38. Taylor,A.F. and Smith,G.R. (1995) Monomeric RecBCD Enzyme Binds and Unwinds DNA (∗) *Journal of Biological Chemistry* **270**, 24451-24458.

39. Singleton,M.R., Dillingham,M.S., Gaudier,M., Kowalczykowski,S.C. and Wigley,D.B. (2004) Crystal structure of RecBCD enzyme reveals a machine for processing DNA breaks. *Nature* **432**, 187-193.

40. Dillingham,M.S., Spies,M. and Kowalczykowski,S.C. (2003) RecBCD enzyme is a bipolar DNA helicase. *Nature* **423**, 893-897.

41. Taylor,A.F. and Smith,G.R. (2003) RecBCD enzyme is a DNA helicase with fast and slow motors of opposite polarity. *Nature* **423**, 889-893.

42. Emmerson,P.T. (1968) Recombination deficient mutants of Escherichia coli K12 that map between thy A and argA. *Genetics* **60**, 19-30.

43. Biek,D.P. and Cohen,S.N. (1986) Identification and characterization of recD, a gene affecting plasmid maintenance and recombination in Escherichia coli. *Journal of bacteriology* **167**, 594-603.

44. Amundsen,S.K., Taylor,A.F., Chaudhury,A.M. and Smith,G.R. (1986) recD: the gene for an essential third subunit of exonuclease V. *Proceedings of the National Academy of Sciences of the United States of America* **83**, 5558-5562.

45. Lovett,S.T., Luisi-DeLuca,C. and kolodner,R.D. (1988) The genetic dependence of recombination in recD mutants of Escherichia coli. *Genetics* **120**, 37-45.

46. Palas,K.M. and Kushner,S.R. (1990) Biochemical and physical characterization of exonuclease V from Escherichia coli. Comparison of the catalytic activities of the RecBC and RecBCD enzymes. *Journal of Biological Chemistry* **265**, 3447-3454.

47. Datsenko,K.A. and Wanner,B.L. (2000) One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 6640-6645.

48. Baba,T., Ara,T., Hasegawa,M., Takai,Y., Okumura,Y., Baba,M., Datsenko,K.A., Tomita,M., Wanner,B.L. and Mori,H. (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection *Molecular systems biology* **2**, 2006.0008.

49. Reisch,C.R. and Prather,K.L.J. (2015) The no-SCAR (Scarless Cas9 Assisted Recombineering) system for genome editing in Escherichia coli. *Scientific reports* **5**, 15096.

50. Ennis,D.G., Levine,A.S., Koch,W.H. and Woodgate,R. (1995) Analysis of recA mutants with altered SOS functions. *Mutation research* **336**, 39-48.

51. Kokubo,K., Yamada,M., Kanke,Y. and Nohmi,T. (2005) Roles of replicative and specialized DNA polymerases in frameshift mutagenesis: mutability of Salmonella typhimurium strains lacking one or all of SOS-inducible DNA polymerases to 26 chemicals. *DNA Repair (Amst)* **4**, 1160-1171.

52. Reuven,N.B., Arad,G., Maor-Shoshani,A. and Livneh,Z. (1999) The mutagenesis protein UmuC is a DNA polymerase activated by UmuD′, RecA, and SSB and is specialized for translesion replication *Journal of Biological Chemistry* **274**, 31763-31766.

53. Pagès,V. and Fuchs,R.P. (2018) Inserting Site-Specific DNA Lesions into Whole Genomes. *Methods Mol Biol* **1672**, 107-118.

54. Subramanya,H.S., Bird,L.E., Brannigan,J.A. and Wigley,D.B. (1996) Crystal structure of a DExx box DNA helicase. *Nature* **384**, 379-383.

55. Korolev,S., Hsieh,J., Gauss,G.H., Lohman,T.M. and Waksman,G. (1997) Major domain swiveling revealed by the crystal structures of complexes of E. coli Rep helicase bound to single-stranded DNA and ADP. *Cell* **90**, 635-647.

56. Beloin,C., Jeusset,J., Revet,B., Mirambeau,G., Le Hégarat,F. and Le Cam,E. (2003) Contribution of DNA conformation and topology in right-handed DNA wrapping by the Bacillus subtilis LrpC protein. *J Biol Chem* **278**, 5333-5342.

57. Pagès,V. and Fuchs,R.P.P. (2002) How DNA lesions are turned into mutations within cells? *Oncogene* **21**, 8957-8966.

58. Hashimoto,Y., Ray Chaudhuri,A., Lopes,M. and Costanzo,V. (2010) Rad51 protects nascent DNA from Mre11-dependent degradation and promotes continuous DNA synthesis. *Nat Struct Mol Biol* **17**, 1305-1311.

59. Schlacher,K., Christ,N., Siaud,N., Egashira,A., Wu,H. and Jasin,M. (2011) Double-Strand Break Repair-Independent Role for BRCA2 in Blocking Stalled Replication Fork Degradation by MRE11 *Cell* **145**, 529-542.

60. Shevelev,I.V. and Hübscher,U. (2002) The 3′–5′ exonucleases *Nature reviews Molecular cell biology* **3**, 364-376.

61. Lovett,S.T. (2011) The DNA Exonucleases of Escherichia coli *EcoSal Plus* **4**,

62. Schlacher,K., Leslie,K., Wyman,C., Woodgate,R., Cox,M.M. and Goodman,M.F. (2005) DNA polymerase V and RecA protein, a minimal mutasome. *Molecular Cell* **17**, 561-572.

63. Napolitano,R., Janel-Bintz,R., Wagner,J. and Fuchs,R.P. (2000) All three SOS-inducible DNA polymerases (Pol II, Pol IV and Pol V) are involved in induced mutagenesis. *The EMBO journal* **19**, 6259-6265.

64. Korada,S.K.C., Johns,T.D., Smith,C.E., Jones,N.D., McCabe,K.A. and Bell,C.E. (2013) Crystal structures of Escherichia coli exonuclease I in complex with single-stranded DNA provide insights into the mechanism of processive digestion. *Nucleic Acids Research* **41**, 5887-5897.

65. Lu,D., Myers,A.R., George,N.P. and keck,J.L. (2011) Mechanism of Exonuclease I stimulation by the single-stranded DNA-binding protein. *Nucleic Acids Research* **39**, 6536-6545.

66. Viswanathan,M. and Lovett,S.T. (1998) Single-strand DNA-specific exonucleases in Escherichia coli. Roles in repair and mutation avoidance. *Genetics* **149**, 7-16.

67. Burdett,V., Baitinger,C., Viswanathan,M., Lovett,S.T. and Modrich,P. (2001) In vivo requirement for RecJ, ExoVII, ExoI, and ExoX in methyl-directed mismatch repair. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 6765-6770.

68. Thoms,B., Borchers,I. and Wackernagel,W. (2008) Effects of Single-Strand DNases ExoI, RecJ, ExoVII, and SbcCD on Homologous Recombination of recBCD+ Strains of Escherichia coli and Roles of SbcB15 and XonA2 ExoI Mutant Enzymes *Journal of bacteriology*

69. Yu,M., Souaya,J. and Julin,D.A. (1998) The 30-kDa C-terminal domain of the RecB protein is critical for the nuclease activity, but not the helicase activity, of the RecBCD enzyme from Escherichia coli. *Proc Natl Acad Sci U S A* **95**, 981-986.

70. Spies,M. and Kowalczykowski,S.C. (2006) The RecA Binding Locus of RecBCD Is a General Domain for Recruitment of DNA Strand Exchange Proteins *Molecular Cell* **21**, 573-580.

71. Brosh,R.M. and Matson,S.W. (1996) A partially functional DNA helicase II mutant defective in forming stable binary complexes with ATP and DNA. A role for helicase motif III. *The Journal of biological chemistry* **271**, 25360-25368.

72. Dillingham,M.S., Soultanas,P. and Wigley,D.B. (1999) Site-directed mutagenesis of motif III in PcrA helicase reveals a role in coupling ATP hydrolysis to strand separation. *Nucleic Acids Research* **27**, 3310-3317.

73. Dillingham,M.S., Soultanas,P., Wiley,P., Webb,M.R. and Wigley,D.B. (2001) Defining the roles of individual residues in the single-stranded DNA binding site of PcrA helicase. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 8381-8387.

74. Belguise-Valladier,P. and Fuchs,R.P. (1995) N-2-aminofluorene and N-2 acetylaminofluorene adducts: the local sequence context of an adduct and its chemical structure determine its replication properties. *J Mol Biol* **249**, 903-913.

75. Mazon,G., Philippin,G., Cadet,J., Gasparutto,D., Modesti,M. and Fuchs,R.P. (2010) Alkyltransferase-like protein (eATL) prevents mismatch repair-mediated toxicity induced by O6-alkylguanine adducts in Escherichia coli. *Proceedings of the National Academy of Sciences* **107**, 18050-18055.

76. Bianco,P.R. and Kowalczykowski,S.C. (2000) Translocation step size and mechanism of the RecBC DNA helicase. *Nature* **405**, 368-372.

77. Wu,C.G., Bradford,C. and Lohman,T.M. (2010) Escherichia coli RecBC helicase has two translocase activities controlled by a single ATPase motor. *Nat Struct Mol Biol* **17**, 1210-1217.

78. Curth,U., Genschel,J., Urbanke,C. and Greipel,J. (1996) In vitro and in vivo function of the C-terminus of Escherichia coli single-stranded DNA binding protein. *Nucleic Acids Res* **24**, 2706-2711.

79. Shereda,R.D., Kozlov,A.G., Lohman,T.M., Cox,M.M. and Keck,J.L. (2008) SSB as an organizer/mobilizer of genome maintenance complexes *Critical reviews in biochemistry and molecular biology* **43**, 289-318.

80. Costes,A., Lecointe,F., McGovern,S., Quevillon-Cheruel,S. and polard,P. (2010) The C-terminal domain of the bacterial SSB protein acts as a DNA maintenance hub at active chromosome replication forks. *PLoS Genetics* **6**, e1001238.

81. Bianco,P.R. (2017) The tale of SSB *Progress in biophysics and molecular biology* **127**, 111-118.

82. Antony,E. and Lohman,T.M. (2019) Dynamics of E. coli single stranded DNA binding (SSB) protein-DNA complexes. *Semin Cell Dev Biol* **86**, 102-111.

83. Bianco,P.R. (2021) The mechanism of action of the SSB interactome reveals it is the first OB-fold family of genome guardians in prokaryotes. *Protein Sci* **30**, 1757-1775.

84. Shereda,R.D., Bernstein,D.A. and Keck,J.L. (2007) A central role for SSB in Escherichia coli RecQ DNA helicase function *Journal of Biological Chemistry* **282**, 19247-19258.

85. Bianco,P.R., Pottinger,S., Tan,H.Y., Nguyenduc,T., Rex,K. and Varshney,U. (2017) The IDL of E. coli SSB links ssDNA and protein binding by mediating protein-protein interactions. *Protein Sci* **26**, 227-241.

86. Thrall,E.S., Piatt,S.C., Chang,S. and Loparo,J.J. (2022) Replication stalling activates SSB for recruitment of DNA damage tolerance factors. *Proc Natl Acad Sci U S A* **119**, e2208875119.

87. Pasero,P. and Vindigni,A. (2017) Nucleases Acting at Stalled Forks: How to Reboot the Replication Program with a Few Shortcuts *Annual review of genetics* **51**, 477-499.

88. Patel,D.R. and Weiss,R.S. (2018) A tough row to hoe: when replication forks encounter DNA damage. *Biochemical Society transactions*

89. Costanzo,V. (2011) Brca2, Rad51 and Mre11: Performing balancing acts on replication forks *DNA Repair* **10**, 1060-1065.

90. Kolinjivadi,A.M., Sannino,V., de Antoni,A., Técher,H., Baldi,G. and Costanzo,V. (2017) Moonlighting at replication forks: a new life for homologous recombination proteins BRCA1, BRCA2 and RAD51 *FEBS letters*

FIGURES and LEGENDS

Figure 1. The RecBC complex, without RecD, is involved in lesion bypass.

The graph represents the partition of lesion tolerance pathways, that are Homology Directed Gap Repair (HDGR), Translesion Synthesis (TLS) and Damaged Chromatid Loss (DCL), as well as cell survival in the presence of a G-AAF lesion in the mutants of the three subunits of the RecBCD complex (for more details see Material & Methods section). The lesion was inserted either in the lagging (lag) or in the leading (lead) strand. For TLS-2 events, the data for lagging and leading have been pooled together since no difference has been observed. The blue and orange asterisks represent the statistical significance for HDGR and TLS-2 events, respectively, for every mutant strain compared to the parental strain.

The data in every graph represent the average and standard deviation of at least three independent experiments. Statistical analyses have been performed with Prism using an unpaired *t*-test. * p < 0,05; ** p < 0,005; *** p < 0,0005.

NB: The TLS data for the parental and *recB* strains have been already published in (18).

Figure 2. Exonuclease screen. A) The graph represents the percentage of Pol II TLS events in the presence of a G-AAF lesion for several exonuclease mutant strains combined with *recB* deletion. The orange asterisks represent the statistical significance for every exonuclease mutant strain compared to *recB* deficient strain. pExoI is a plasmid expressing *exoI* gene under its own promoter. ExoIA183V is the nuclease dead allele of ExoI (also known as *sbcB15*). **B)** The graph represents the percentage of Pol II and Pol V TLS events in the presence of a G-AAF lesion. All the strains contain the pVP148 plasmid that expresses *polB* and *umuD'C* genes under their native promoters. The orange and green asterisks represent the statistical significance for the mutant strains compared to *recB* deficient strain. **C)** The graph represents the percentage of Pol IV TLS events in the presence of a BaP lesion. The violet asterisks represent the statistical significance for the mutant strains compared to *recB* deficient strain. The data in every graph represent the average and standard deviation of more than four independent experiments obtained in the lagging or in the leading orientation. Statistical analyses have been performed with Prism using an unpaired *t*-test: * p < 0,05; ** p < 0,005; *** p < 0,0005.

NB: The data for the parental and *recB* strains in A) and C) panels have been already published in (18).

Figure 3. Involvement of RecB domains. A) Domain organization of RecB protein: the N-term SF1 helicase core region (residue 1-928) is linked by a 70 aa linker to the C-term nuclease core (residue 998-1180). The SF1 helicase core is composed of two domains (1 and 2), each divided in two subdomains (A and B) and seven conserved motifs (I, Ia, II, III, IV, V, VI). The amino acid positions of the *E. coli* motifs are indicated in brackets. S.A stands for *Staphylococcus aureus* and B.S for *Bacillus stearothermophilus*. In red are indicated the amino acid residues that we have mutated by CRISPR-Cas9 technology. **B)** The graph represents the Pol II TLS events in the presence of a G-AAF lesion of the DNA binding mutant alleles of RecB. **C)** The graph represents the Pol II TLS events in the presence of a G-AAF lesion of the DNA binding mutant RecB$^{YFR-AAA}$ in the presence of the plasmid overexpressing Pol II (pVP148). The orange asterisks represent the statistical significance compared to parental strain. The data represent the average and standard deviation of more than four independent experiments obtained in the lagging or in the leading orientation. Statistical analyses have been performed with Prism using an unpaired *t*-test. *** $p < 0,0005$. YFR-AAA is a mutant in the amino acid residues Y420A, F422A, R423A.

NB: The data for the parental, *recB* and *recB*^{D1080A} strains have been already published in (18).

Figure 4: RecBC is required for efficient lesion bypass only in the presence of strong-replication blocking lesions. The graph represents the tolerance events TLS like in the presence of less blocking lesions due to DNA pol III or to a TLS polymerase bypass. The data represent the average and standard deviation of more than four independent experiments obtained in the lagging or in the leading orientation.

Figure 5: RecBC complex binds to a ssDNA gap in the presence of SSB

A) DNA substrates used for the electrophoretic mobility shift assays. The circular ssDNA of the phage φX174 was used to obtained a 5.3kb ssDNA gap or a 2.5kb ssDNA gap. The 5'of the ds-ss junction of the substrates have been radiolabeled with ^{32}P. **B)** Mobility shift assays on agarose gel using both DNA substrates in the presence of RecBC complex and with (or without) increased concentrations of SSB (see Material and Methods for more details). In the absence of SSB, no band shift is observed for RecBC alone. **C)** Mobility shift assay on agarose gel using the 5.3kb DNA gap in the presence of the RecB^{F64A}C allele. **D)** Mobility shift assay on agarose gel using the 5.3kb DNA gap in the presence of the truncated form of SSB (SSBΔ8).

Figure 6: Transmission Electron Microscopy analysis of RecBC-SSB-ssDNA gap complexes in darkfield imaging mode. A) Control DNA substrate containing a 2.5 kb ssDNA gap. ssDNA is collapsed under these spreading conditions. **B)** A non-saturating SSB concentration (200 nM) was incubated with 18 mM (in nucleotides) of a 2.5 kb ssDNA gap substrate to allow ssDNA unfolding without being completely covered by SSB. **C and D)** RecBC (100 nM) was added to the reaction. RecBC interacts with the SSB partially covered ssDNA and induces a local condensation and restructuring of the ssDNA. Complexes are more or less condensed making unable to distinguish between SSB and RecBC proteins. The scale bar corresponds to 100 nm. The red rectangles indicate the region of the substrate as depicted in the cartoon.

Figure 7: Model of RecBC complex protecting the nascent DNA in the presence of a ssDNA gap. The replicative DNA polymerase is temporarily blocked at a DNA lesion : i) when the lesion can be easily bypassed by the replicative DNA polymerase itself or by a TLS polymerase at the fork, the SSB-covered ssDNA formed by the DNA unwinding is transient and the RecBC complex is not needed for protection; ii) in the presence of a strong blocking lesion, replication restarts downstream creating a persistent SSB-covered ssDNA gap: RecBC is recruited and binds to the SSB-covered ssDNA gap by restructuring and condensing the ssDNA to protect the nascent DNA against nuclease attack. This will allow efficient lesion bypass by one of the two lesion tolerance pathways. In the absence of RecBC, ExoI can access the 3' end of the nascent DNA (potentially with the help of a helicase) degrading the nascent DNA. This will impact lesion bypass, in particular the action of the TLS polymerases, but also the HDGR mechanism.