Structure and evolution of Alanine/Serine Decarboxylases and the engineering of theanine production

Hao Wang[a,1], Biying Zhu[b,1], Siming Qiao[b], Chunxia Dong[b], Xiaochun Wan[b], Weimin Gong[a,*], Zhaoliang Zhang[b,*]

[a]Department of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[b]State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, Hefei, Anhui 230036, China.

[1]These authors contributed equally to this work.

* To whom correspondence should be addressed. Email: zhlzhang@ahau.edu.cn or wgong@ustc.edu.cn.

Author contributions: W.G, Z.Z and X.W designed research; H.W, B.Z, S.Q and C.D performed research; H.W and B.Z analyzed data; H.W, B.Z, W.G, and Z.Z wrote the paper.

Competing interest statement: The authors declare no competing financial interests.

Classification: Biochemistry (major), Plant Sciences (minor)

Keywords: alanine decarboxylase, serine decarboxylase, substrate specificity, theanine, *Camellia sinensis*

Abstract

Ethylamine (EA), the precursor of theanine biosynthesis, is synthesized from alanine decarboxylation by alanine decarboxylase (AlaDC) in tea plants. AlaDC evolves from serine decarboxylase (SerDC) through neofunctionalization and has lower catalytic activity. However, lacking structure information hinders the understanding of the evolution of substrate specificity and catalytic activity. In this study, we solved the X-ray crystal structures of AlaDC from *Camellia sinensis* (CsAlaDC) and SerDC from *Arabidopsis thaliana* (AtSerDC). Tyr^{341} of AtSerDC or the corresponding Tyr^{336} of CsAlaDC is essential for their enzymatic activity. Tyr^{111} of AtSerDC and the corresponding Phe^{106} of CsAlaDC determine their substrate specificity. Both CsAlaDC and AtSerDC have a distinctive zinc finger and have not been identified in any other Group II PLP-dependent amino acid decarboxylases. Based on the structural comparisons, we conducted mutation screen of CsAlaDC. The results indicated that the mutation of L110F or P114A in the CsAlaDC dimerization interface significantly improved the catalytic activity by 110% and 59%, respectively. Combining a double mutant of CsAlaDC$^{L110F/P114A}$ with theanine synthetase increased theanine production 672% in an *in vitro* system. This study provides the structural basis for the substrate selectivity and catalytic activity of CsAlaDC and AtSerDC and provides a route to more efficient biosynthesis of theanine.

Introduction

Tea is one of the most popular natural non-alcoholic beverages consumed worldwide. About 2 billion cups of tea are consumed worldwide daily (1). Theanine (γ-glutamylethylamide) is a unique non-protein amino acid and the most abundant free amino acid in tea plants (*Camellia sinensis*). It accounts for more than 50% of the total free amino acids and approximately 1-2% of the dry weight of the new shoots of tea plants (2). Theanine is the secondary metabolite conferring the umami taste of tea infusion and also balances the astringency and bitterness of tea infusion caused by catechins and caffeine (3). It has also many health-promoting functions, including neuroprotective effects, enhancement of immune functions, and potential anti-obesity capabilities, among others (2, 4-9). Therefore, theanine content is highly correlated with green tea quality (10).

Theanine is synthesized from ethylamine (EA) and glutamate (Glu) by theanine synthetase (11, 12). Importantly, the large amount of theanine biosynthesis is determined by the high availability of EA in tea plants (13). Therefore, the evolution of EA biosynthesis in tea plants provided the basis for theanine biosynthesis and the formation of tea quality. EA is synthesized from alanine decarboxylation by alanine decarboxylase (CsAlaDC) (14) (Figure 1A). Indeed, CsAlaDC expression level and catalytic activity largely determine the theanine accumulation level in tea plants (15). As a novel gene, *CsAlaDC* had not been reported in other plant species before it was identified in tea plants. Previous studies indicated that *AlaDC* originated from the serine decarboxylase gene (*SerDC*) by gene duplication in tea plants (16). However, despite

sharing highly conserved amino acid sequences, CsAlaDC and CsSerDC specifically

catalyze the decarboxylation of alanine and serine, respectively (16) (Figure 1A and B).

In addition, CsAlaDC exhibited a much lower enzymatic activity compared with

CsSerDC (16). However, the structural basis and key amino acids underlying the

evolution of the substrate specificity and enzymatic activity of CsAlaDC are unknown.

SerDC belongs to the Group II pyridoxal-5-phosphate (PLP)-dependent

decarboxylase superfamily (17). These Group II amino acid decarboxylases produce

many bioactive secondary metabolites and signaling molecules (18-20). In plants,

SerDC catalyzes the biosynthesis of ethanolamine from serine, which was first

functionally characterized in *Arabidopsis thaliana* (21) (Figure 1A). Ethanolamine is

an important metabolite in the synthesis of phosphatidylethanolamine and

phosphatidylcholine, two main phospholipids involved in maintaining eukaryotic

membrane structures in eukaryotic cell membranes (22-24). Furthermore, the study has

shown that SerDC is essential in the embryogenesis of *Arabidopsis* (25). SerDC is of

vital importance for plant growth and development, but the mechanism of substrate

recognition and catalytic activity of SerDC also remains unclear.

In this work, we attempt to understand the mechanism of functional divergence of

plant AlaDC and SerDC by analyzing their structural characteristics. Here, we obtained

the X-ray crystal structure of CsAlaDC and AtSerDC. According to the crystal structure,

we found a distinctive zinc finger structure located at both CsAlaDC and AtSerDC,

which has not been identified in any other Group II PLP-dependent amino acid

decarboxylases that have been previously characterized. By comparing the substrate

binding pockets, we identified Phe[106] of CsAlaDC and Tyr[111] of AtSerDC as the crucial

sites for substrate specificity. By conducting mutation screening based on the protein

structures, we identified the amino acids repressing the catalytic activity and discovered

that CsAlaDC[L110F/P114A] exhibited a 2.3-fold increase in catalytic activity compared to

that of CsAlaDC and enhanced the engineering of theanine production *in vitro*.

Results

Enzymatic properties of CsAlaDC, AtSerDC, and CsSerDC

CsAlaDC originates from CsSerDC by gene duplication and neofunctionalization in tea

plants, but they catalyze different metabolic processes (Figure 1A). We performed a

multiple sequence alignment of the amino acid sequences of the 6 SerDCs from

kiwifruit (*Actinidia chinensis*), grape (*Vitis vinifera*), coffee (*Coffea eugenioides*),

cocoa (*Theobroma cacao*), Arabidopsis and tea plants and CsAlaDC (Figure 1B). The

results showed that the amino acid sequences of CsAlaDC and SerDCs were highly

conserved, but CsAlaDC has amino acid mutations in highly conserved regions

compared with SerDCs.

Next, to verify the substrate specificity and enzyme activity, we conducted enzyme

activity detection and enzyme kinetics analysis for CsAlaDC, CsSerDC and AtSerDC.

The 5′ truncated *CsAlaDC* and *AtSerDC* were inserted into the pET22b and pET28a

expression vectors to generate recombinant plasmids pET22b-*CsAlaDC* and pET28a-

AtSerDC, correspondingly. Additionally, the full-length protein of CsSerDC was

inserted into the pET28a expression vector to generate the recombinant plasmid

pET28a-*CsSerDC*. Subsequently, the recombinant proteins were expressed in

Escherichia coli and purified via nickel affinity chromatography. The purified proteins

were examined through SDS-PAGE analysis (Figure 2A), supplemented by size-

exclusion chromatography assessments (Supplementary figure 1A), it seems that the

protein manifests in an oligomeric configuration within the solution.

To verify the substrate specificity of CsAlaDC, AtSerDC, and CsSerDC, enzyme

activity assays were conducted using Ala and Ser as substrates, followed by product

identification via UPLC analysis. The results showed that CsAlaDC can effectively

catalyze the decarboxylation of alanine to generate EA, whereas its catalytic efficacy

towards serine was relatively inferior. Conversely, AtSerDC and CsSerDC selectively

catalyzed serine decarboxylation to yield ethanolamine but did not exhibit activity

towards alanine (Figure 2B). These findings confirmed that CsAlaDC and SerDCs do

not have promiscuous decarboxylase activity on Ala and Ser.

The kinetic properties of CsAlaDC, AtSerDC, and CsSerDC were determined

through the use of corresponding substrates (Figure 2C), and the results are presented

in Table 1. The kinetic parameter Km of CsAlaDC was determined to be 1.215 mM,

which is similar to that of AtSerDC and CsSerDC at 1.522 mM and 2.364 mM,

respectively. However, AtSerDC has a $Vmax$ of 7.053 μmo L^{-1} s^{-1}, which is 4-fold that

of CsAlaDC's $Vmax$ of 1.709 μmo L^{-1} s^{-1}; while CsSerDC has a $Vmax$ of 9.031 μmo L^{-}

^{1} s^{-1}, which is 5.3-fold that of CsAlaDC. The findings suggest that the enzymatic

catalytic efficiency of CsSerDC and AtSerDC is approximately triple compared to that

of CsAlaDC, and the $Vmax$ of CsAlaDC is considerably lower than that of both

CsSerDC and AtSerDC. The observed disparity in $Vmax$ suggests that CsAlaDC and

SerDCs may exhibit discrepancies in substrate-to-product conversion rates under

saturated substrate conditions. It is plausible that SerDCs possess a higher turnover rate,

facilitating the rapid conversion of substrate to product and subsequent release of the

enzyme. Another possibility is that SerDCs demonstrate enhanced stability, thereby

maintaining superior catalytic activity even at higher substrate concentrations,

consequently leading to a higher *Vmax* value. Due to the similarity in catalytic activity

between CsSerDC and AtSerDC, we chose the more representative AtSerDC for further

analysis.

The overall structures of CsAlaDC and AtSerDC

To enhance our comprehension of CsAlaDC and AtSerDC, we conducted structural

analyses of these two proteins. Through optimization of crystallization conditions, we

successfully determined the crystal structures of CsAlaDC, CsAlaDC-EA, and

AtSerDC at resolutions of 2.50 Å, 2.60 Å and 2.85 Å, respectively (Supplementary

table 1).

The overall structure of CsAlaDC or AtSerDC is homodimeric with two subunits

exhibiting an asymmetrical arrangement (Figure 3A and 3D). The monomer of

CsAlaDC or AtSerDC is divided into three distinct structural domains: an N-terminal

domain (N-terminal -104 aa in CsAlaDC, N-terminal -109 aa in AtSerDC), a large

domain (105-364 aa in CsAlaDC, 110-369 aa in AtSerDC), and a small C-terminal

domain (365 aa-C-terminus in CsAlaDC, 370-C-terminus as in AtSerDC), colored light

pink, khaki, and sky blue, respectively. Compared to the full-length protein, the N-

terminal structures of CsAlaDC and AtSerDC were truncated by 60 and 65 amino acid

residues, respectively. The N-terminal domain of the truncated protein contains a long

α-helix, which is connected to the large domain through a long loop. The helix of one

subunit is antiparallel to the corresponding helix in the neighboring subunit, forming a

clamp. Therefore, the N-terminal domain may not be an independent folding unit and

is likely only stable in dimers. The large domain contains a seven-stranded mixed β-

sheet surrounded by eight α-helices, while the C-terminal domain is composed of a

three-stranded antiparallel β-sheet and three α-helices (Supplementary figure 1). The

catalytic site of the enzyme is positioned within a superficial crevice at the junction of

two subunits forming a dimeric arrangement. The amino acid residues derived from

both subunits participate in the binding of the PLP, effectively anchoring it to the active

site. In the catalytic site, one monomer accommodates PLP, while the other monomer

also contributes to PLP binding and enzymatic function (Figure 3A and 3D).

Notably, our investigation has revealed the presence of a distinctive zinc finger

structure (as depicted in Figure 3A and 3D) located at the C-terminus of CsAlaDC and

AtSerDC. This structure is composed of a loop structure spanning 17 amino acid

residues, wherein coordination of Zn^{2+} is facilitated by three Cys residues and one His

residue. Importantly, this particular configuration is exclusive to the two proteins under

examination and has not been identified in any other Group II PLP-dependent amino

acid decarboxylases that have been previously characterized (26, 27).

The crystal structures of CsAlaDC-EA complex and AtSerDC were obtained and

further analyzed. In the former complex, PLP bound to Lys[309] via a Schiff base linkage

(internal aldimine form), and the pyridine moiety of PLP is positioned between the

imidazole ring of His[196] and the methyl group of Ala[279] in a parallel orientation to the

imidazole ring. The carboxylic group of Asp[277] stabilizes the N1 of PLP by a salt bridge

interaction, providing the latter with the strong electrophilicity necessary to stabilize

carbon ion intermediates during enzymatic catalysis. The O3 forms hydrogen bonds

with Thr[247] and Lys[309]. Additionally, His[308], Gly[169], Thr[170], Lys[309], and Ser[347]*("*"

denotes the amino acids on adjacent subunits) establish a hydrogen bonding network

with the phosphate group of PLP, consequently fortifying its attachment to the protein

(Figure 3C). Similarly, in the latter structure, the pyridine ring of PLP was sandwiched

between the imidazole ring of His[201] and the methyl group of Ala[284]. The N1 of PLP

formed a salt bridge with Asp[282], while the O3 formed hydrogen bonds with Thr[252]. The

phosphate group of PLP established hydrogen bonding interactions with Lys[314], His[313],

Gly[174], Thr[175], and Ser[352]* (Figure 3E). The aforementioned amino acid residues are

observed across various Group II PLP-dependent amino acid decarboxylases and

exhibit considerable stability (see further discussion below), suggesting that they could

share common features in terms of catalytic mechanisms.

Factors affecting CsAlaDC and AtSerDC activity

The Group II PLP-dependent amino acid decarboxylases are dimeric enzymes with

each monomer containing an active site located within a shallow cavity (28). A flexible

loop of amino acids originating from one monomer extends into the active site of the

other monomer and plays a crucial role in catalysis. The catalytic loop in AtSerDC,

consisting of amino acid residues 328-341, is well-structured and exhibits clear electron

density. However, the corresponding loop in CsAlaDC is disordered (Figure 3F). The

aforementioned loop harbors a conserved Tyr residue that is believed to function as a proton donor of the carbanion in the catalytic process (29). In particular, CsAlaDC Tyr[336] and AtSerDC Tyr[341] correspond to that Tyr residue (Supplementary figure 2). Upon superimposing the active site residues of CsAlaDC apo and CsAlaDC-EA complex, it was observed that the hydroxyl group of Tyr[336]* in the CsAlaDC-EA complex exhibits a substantial 60-degree deviation along the Cα-Cβ bond (Figure 3G). To assess the function of the Tyr, we substituted the corresponding Tyr residues in CsAlaDC and AtSerDC with Phe. The resulting CsAlaDC[Y336F] and AtSerDC[Y341F] mutants were then exposed to Ala and Ser, respectively. We measured the production of EA or ethanolamine in the reaction mixtures. Our findings indicate that these mutants catalyze abortive decarboxylation, as evidenced by the absence of detectable EA and only a small amount of ethanolamine observed in the reaction mixture (Figure 3H). This result suggested this Tyr is required for the catalytic activity of CsAlaDC and AtSerDC.

Identification of key amino acids for the substrate specificity

The superposition of amino acid residues in the substrate binding pocket of CsAlaDC and AtSerDC revealed that all residues are identical, except for the substitution of Tyr at position 111 in AtSerDC with Phe at position 106 in CsAlaDC (Figure 4A). This observation suggests a potential role for this specific residue in determining the selective binding of the appropriate substrate.

To gain further insights into the functional role of this residue (Tyr[111] in AtSerDC or Phe[106] in CsAlaDC) in selective binding of the appropriate substrate, we identified 563 potential serine decarboxylases in Embryophyta using the amino acid sequences of

AtSerDC (Supplementary figure 3A). By comparing the amino acid sequences of these

serine decarboxylases, we observed that the corresponding residues to Tyr^{111} or Phe^{106}

are located within a conserved motif comprising 9 amino acids (Figure 4B). Within this

motif, the first two residues, Y (Tyrosine) and P (Proline) are completely conserved

across all 563 proteins. However, the third residue where Tyr^{111} in AtSerDC or Phe^{106}

in CsAlaDC is positioned exhibits variability among Y, T (Threonine), F

(Phenylalanine), V (Valine), A (Alanine), I (Isoleucine), L (Leucine), or other amino

acids (Figure 4B). Remarkably similar to Tyr^{111}, this residue is predominantly Y in 83.7%

of these homologs; conversely, F was found to be present in only 2.3% of these

homologs within plant species belonging to Poales, Asterales, Ranunculales, Ericales,

Caryophyllales and Nymphaeales orders (Figure 4C-4D).

To verify the role of Tyr^{111} in AtSerDC or Phe^{106} in CsAlaDC in the selective

binding of the appropriate substrate, we introduced mutations in Tyr^{111} to Phe

($AtSerDC^{Y111F}$) and Phe^{106} to Tyr ($CsAlaDC^{F106Y}$), followed by *in vitro* enzymatic

activity assays. The results revealed that $AtSerDC^{Y111F}$ acquired alanine decarboxylase

activity, whereas $CsAlaDC^{F106Y}$ completely lost its alanine decarboxylase activity

(Figure 4E). Additionally, we generated other mutations of Tyr^{111} in AtSerDC, including

$AtSerDC^{Y111A}$, $AtSerDC^{Y111I}$, $AtSerDC^{Y111L}$, $AtSerDC^{Y111V}$ and $AtSerDC^{Y111W}$. Among

these mutants, only $AtSerDC^{Y111I}$ exhibited a marginal level of alanine decarboxylase

activity (Figure 4E). These findings indicate the essential role of Phe^{106} in the selective

binding of alanine for CsAlaDC.

On the other side, both CsAlaDC and $CsAlaDC^{F106Y}$ exhibit a very low level of

serine decarboxylase activity (Figure 4F). Moreover, compared with AtSerDC, the serine decarboxylase activity of AtSerDC^{Y111F} was about 30% of AtSerDC; AtSerDC^{Y111I} retained lower than 5% of serine decarboxylase activity of AtSerDC; while other mutations, including AtSerDC^{Y111A}, AtSerDC^{Y111L}, AtSerDC^{Y111V} and AtSerDC^{Y111W} abolished the serine decarboxylase activity (Figure 4F). These results suggested the Tyr^{111} of AtSerDC is also important for the selective binding of serine for AtSerDC.

To further verify that Phe^{106} of CsAlaDC and Tyr^{111} of AtSerDC were key amino acid residues determining the selective binding of substrates *in planta*, we employed the *Nicotiana benthamiana* transient expression system. To this end, *A. tumefaciens* strain GV3101 (pSoup-p19), carrying recombinant plasmid, was infiltrated into leaves of 5-week-old *N. benthamiana* plants, and the pCAMBIA1305 empty vector was used as the control (EV). Relative mRNA levels were detected (Supplementary figure 4), indicating that they had been successfully overexpressed in tobacco. Here, we found that a high level of EA was produced in the CsAlaDC-expressing tobacco leaves; while no EA product was detected in tobacco leaves infiltrated with the mutant CsAlaDC^{F106Y}. In addition, we did not detect EA products in the AtSerDC-expressing tobacco leaves, while a high level of EA was detected in tobacco leaves infiltrated with mutant AtSerDC^{Y111F}. As anticipated, EA was not detected in tobacco leaves infiltrated with EV (Figure 4G and H). These results further verified the critical role of Phe^{106} in the

selective binding of alanine for CsAlaDC.

Key amino acids for the evolution of CsAlaDC enzymatic activity

CsAlaDC and AtSerDC have a high sequence similarity of 74.5% and a nearly identical

structure with an RMSD (root mean square deviation) of only 0.77Å for monomer

structures. However, the two enzymes catalyze different amino acid decarboxylation

reactions, and AtSerDC exhibits significantly higher $Vmax$ than CsAlaDC (Figure 2B

and C). In light of this observation, we postulated a hypothesis: EA, generated via Ala

decarboxylation by CsAlaDC, can be toxic and harmful to plants if accumulated

excessively. Thus, during the evolution of plant serine decarboxylase into alanine

decarboxylase, the enzyme has evolved not just to alter the substrate preference but also

to reduce catalytic activity to control EA production within a suitable range. Based on

this hypothesis, we suggest that mutating specific amino acids in CsAlaDC to those

corresponding amino acids in SerDCs could enhance its activity, and the results could

provide insights into the evolution of CsAlaDC enzymatic activity.

Dimerization is essential to AlaDC/SerDC activities because the active site is

composed of residues from two monomers. To identify the amino acids repressing the

enzymatic activity of CsAlaDC during evolution from SerDC, we analyzed the crystal

structures and the amino acids at the dimer interface between CsAlaDC and AtSerDC.

This analysis revealed that the amino acids at the dimer interface at positions 66, 97,

110, 114, 116, 117, 122, 315 and 345 are different in CsAlaDC and SerDCs (Figure 1B).

Therefore, we mutated these amino acids of CsAlaDC into those in the corresponding

positions of AtSerDC and CsSerDC, and performed enzyme activity assays (Figure 5A).

The results demonstrated that CsAlaDC[L110F] and CsAlaDC[P114A] exhibited significantly

enhanced enzyme activity compared to the wild-type CsAlaDC, with a 2.1-fold and

1.59-fold increase, respectively. Furthermore, the catalytic activity of the

CsAlaDC[L110F/P114A] double mutant exhibits a remarkable 2.3-fold increase compared to

that of the wild-type protein (Figure 5B). These findings suggested a critical role of

Leu[110] and Pro[114] of CsAlaDC in the evolution of enzymatic activity and provided a

basis to improve CsAlaDC activity. It is possible that these amino acid residues could

potentially augment the hydrophobic nature of the protein dimer interface.

In Vitro **synthesis of theanine**

The biosynthetic pathway of theanine in tea plants comprises two consecutive

enzymatic steps: alanine decarboxylase facilitates the decarboxylation of alanine to

generate EA, while theanine synthetase catalyzes the condensation reaction between

EA and Glu to synthesize theanine. By conducting mutation screens, we discovered a

CsAlaDC mutant protein (L110F/P114A) that exhibited a 2.3-fold higher catalytic

activity compared to the wild-type protein. Subsequently, we employed an *in vitro*

theanine synthesis system utilizing CsAlaDC and either glutamine synthetase (PsGS

[P*seudomonas syringae* pv. syringae]) or gamma-glutamate methylamine ligase

(MmGMAS [*Methylovorus mays*]) which have the ability to synthesize theanine from

EA and Glu (30, 31).

The results illustrated the successful synthesis of theanine using CsAlaDC in

conjunction with two theanine synthetases, employing Ala and Glu as substrates (Figure

5C). The theanine content generated by the combination of CsAlaDC[L110F] and

MmGMAS, as well as CsAlaDC$^{L110F/P114A}$ and MmGMAS, was 4.57-fold and 6.72-fold

higher than the content produced in the combination of wild-type CsAlaDC and

MmGMAS, respectively. Similarly, when combined with the PsGS, comparable

outcomes are observed as well. The theanine content resulting from the combination of

CsAlaDC^{L110F} and PsGS, as well as CsAlaDC$^{L110F/P114A}$ and PsGS, exhibit

enhancements of 1.62-fold and 4.33-fold compared to the wild-type protein

combination, respectively (Figure 5C). Hence, the utilization of CsAlaDC$^{L110F/P114A}$

could effectively enhance the theanine production yield and thus, holds potential for

large-scale engineering production of theanine.

Discussion

The common and distinctive structural characteristics of CsAlaDC and AtSerDC

compared with other amino acid decarboxylases

CsAlaDC and AtSerDC are Group II PLP-dependent amino acid decarboxylases, which

exhibit numerous structural features common to other amino acid decarboxylases. The

Dali search (32) revealed that AtSerDC and CsAlaDC share structural similarities with

7CIG (MetDC [*Streptomyces sp.* 590]), 7ERV (HisDC1 [*Photobacterium*

phosphoreum]), 6KHO (TrpDC [*Oryza sativa Japonica* Group]), 4E1O (HisDC2

[*Homo sapiens*]), 6JY1 (AspDC [*Methanocaldococcus jannaschii*]), and 5GP4 (GluDC

[*Levilactobacillus brevis*]). The monomers of these amino acid decarboxylases all

consist of three characteristic domains: the C-terminal domain, the large domain and

the N-terminal domain (Supplementary figure 5). The amino acid residues that bind to

PLP cofactors are conserved across multiple enzymes (Supplementary figure 5 and 6).

Out of the eight proteins analyzed, a total of thirteen amino acid residues were found to

be conserved across all sequences (Glu^{143}, Glu^{171}, Lys^{201}, Gly^{245}, Thr^{247}, Asp^{253}, His^{275},

Asp^{277}, Ala^{279}, Pro^{286}, Ser^{302}, Lys^{309}, and Tyr^{336} in CsAlaDC). Some of them are situated

within the active center and play a role in stabilizing PLP or facilitating catalytic

reactions. Conversely, other residues reside outside the active center and their function

remains unclear (Supplementary figure 6).

Group II PLP-dependent amino acid decarboxylases are characterized by the

existence of a highly flexible loop, which is of great significance for the catalytic

mechanism of decarboxylase (33). The loops are located at the dimer interface and

extend to the active sites of other monomers in a closed conformation. Prior research

has established that the conserved amino acid residue Tyr within the loop plays a crucial

role in catalysis by donating protons to carbanions of quinone intermediates that arise

following decarboxylation (34, 35). Our experiments demonstrate that the substitution

of the corresponding Tyr with Phe in the loop renders the protein inactive. Through the

application of circular dichroism, we have corroborated that the stability of the protein

is not compromised by mutations (Supplementary figure 7A, B, D, E). Both the mutant

and wild-type proteins manifest absorption bands at the 420 nm wavelength, signifying

the formation of a Schiff base between PLP and the lysine residues found at the active

site. Given that PLP has been incorporated during the protein purification stage, the

absorbance observed for both the mutant and wild-type proteins at 420 nm is consistent

when compared at equivalent concentrations (Supplementary figure 8). This

corroboration implies that the protein mutation does not interfere with the binding to

PLP. Consequently, the inactivation observed in the mutant proteins is not a consequence of instability incited by corresponding mutations, nor changes in the binding affinity between the protein and PLP induced by these mutations.

We observed that the substitution of Phe[106] for Tyr in CsAlaDC rendered CsAlaDC inactive, while the substitution of Tyr[111] for Phe in AtSerDC enabled alanine decarboxylase activity, and the alteration in the activity of the mutant protein bears no correlation to protein stability and affinity for PLP (Supplementary figure 7A, C, D, F and Supplementary figure 8), suggesting that the amino acid in the position is critical for determining substrate specificity. Integrative crystallographic structure of the two enzymes, we speculate the hydrophilic nature of the Lys residue in AtSerDC predisposes the active site to preferentially accommodate the hydrophilic amino acid Ser as its substrate. In contrast, the equivalent position in CsAlaDC is occupied by Phe, an amino acid lacking the hydroxyl group. This substitution enhances the hydrophobic nature of the substrate-binding pocket. Consequently, CsAlaDC demonstrates a unique predilection, selectively binding Ala (an amino acid with comparatively hydrophobic properties) as its preferred substrate.

The monomeric configuration of CsAlaDC and AtSerDC is akin to that of other Group II PLP-dependent amino acid decarboxylases, except for a conspicuous dissimilarity. Unlike other Group II PLP-dependent amino acid decarboxylases, the C-terminal of CsAlaDC and AtSerDC harbors an evident zinc finger structure. We propose that this structure could potentially influence enzyme stability. To test this hypothesis, we truncated the zinc finger structure from both proteins and expressed them. Our

findings indicate that, following the excision of the zinc finger structure, CsAlaDC

became insoluble, while AtSerDC displayed a similar trait to a certain extent

(Supplementary figure 9). Hence, we can establish that the zinc finger structure indeed

exerts an influence on the enzyme's stability. The potential for additional functions

necessitates further investigation.

Evolution of AlaDC and SerDC

The structure of CsAlaDC seems like with AtSerDC, however, there are differences in

substrate specificity. To clarify the relationship between CsAlaDC and other SerDCs,

we analyzed serine decarboxylase-like proteins in Embryophyta that are highly

homologous to CsAlaDC and constructed phylogenetic trees to gain insight into their

evolutionary relationships (Supplementary figure 3). Based on our experimental

findings and evolutionary evidence, we have identified a conserved YPX motif at the

substrate binding pocket. For most plants, the residue X in this motif is predominantly

Tyr, which corresponds to the serine decarboxylase protein. However, other residues

such as Thr, Phe, Val, Ala, and Ile can also occupy this position (Figure 4B and C).

Interestingly, serine decarboxylase-like proteins containing YPF motifs, which have the

potential to function as alanine decarboxylases, were found to be distributed throughout

the phylogenetic tree, including Asteroideae, Ericales, Chenopodiaceae, Poaceae, and

Ranunculales (Supplementary figure 3B). However, these proteins were absent in some

more recent species. This observation implies that the emergence of alanine

decarboxylase in these particular species could be attributed to convergent evolution.

Moreover, we have observed the enrichment of serine decarboxylase-like proteins

that possess a YPT motif within the Fabales (Supplementary figure 3B). Prior research has demonstrated that XP_004496485-1, which possesses a YPT motif, displays catalytic efficacy toward the processes of decarboxylation and oxidative deamination of Phe, Met, Leu, and Trp (36). Given that the YPT motif is highly conserved and widely distributed in Fabales, serine decarboxylase-like proteins bearing the YPT motif may have developed a unique substrate specificity in Fabales, beyond their conventional decarboxylation functions, as exemplified by XP_004496485-1 protein mentioned above. We speculate that these proteins may be capable of catalyzing other reactions, potentially involving non-protein amino acids or other substances as substrates.

Applied to improve the synthesis of theanine

Theanine is an important indicator of green tea quality. Therefore, improving the synthesis of theanine in tea plants is the focus of research. In this study, through crystal structure analysis and mutation verification, we found that the catalytic activity of CsAlaDC$^{L110F/P114A}$ is 2.3 times higher than that of the wild-type protein, resulting in a more abundant synthesis of theanine *in vitro*. This gives us great inspiration to improve the synthesis of EA in tea plants by gene editing, thus increasing the content of theanine in tea plants (Figure 5).

Theanine is highly demanded, by the market, due to its health effects and medicinal value, and as a constituent in food, cosmetics and other fields. To meet the market demand, a variety of methods have been used to acquire theanine, with the main methods including direct extraction, chemical synthesis, biotransformation (microbial

fermentation) and plant cell culture. Based on the findings of this study, site-directed

mutagenesis can be employed to modify enzymes involved in theanine synthesis. This

modification enhances the capacity of bacteria, yeast, model plants, and other

organisms to synthesize theanine, thereby facilitating its application in industrial

theanine production.

Conclusions

In conclusion, our structural and functional analyses have significantly advanced

understanding of the substrate-specific activities of alanine and serine decarboxylases,

typified by CsAlaDC and AtSerDC. Critical amino acid residues responsible for

substrate selection were identified-Tyr111 in AtSerDC and Phe106 in CsAlaDC-

highlighting pivotal roles in enzyme specificity. The engineered CsAlaDC mutant

(L110F/P114A) not only displayed enhanced catalytic efficiency but also substantially

improved L-theanine yield in a synthetic biosynthesis setup with PsGS or MmGMAS.

Our research expanded the repertoire of potential alanine decarboxylases through the

discovery of 13 homologous enzyme candidates across embryophytic species and

uncovered a special motif present in serine protease-like proteins within Fabale,

suggesting a potential divergence in substrate specificity and catalytic functions. These

insights lay the groundwork for the development of industrial biocatalytic processes,

promising to elevate the production of L-theanine and supporting innovation within the

tea industry.

Materials and Methods

Plants materials.

Tobacco (*Nicotiana benthamiana*) plants were grown in a controlled chamber, under a 16-h light and 8-h dark photoperiod at 25 °C. Leaves of 5-week-old tobacco plants were used for transient transformation, mediated by *Agrobacterium tumefaciens* strain GV3101.

Gene cloning and protein expression

The coding sequences of AtSerDC and CsAlaDC, amino acid residues 66-482 and 61-478, respectively, were amplified using PCR and then ligated into the *Nde* I and *Xho* I restriction sites of the pET-28a and pET-22b vector, respectively, providing the recombinant vector pET-28a-*AtSerDC* and pET-22b-*CsAlaDC*. The cDNAs encoding CsSerDC were amplified using PCR and then ligated into the *Nde* I and *Xho* I restriction sites of the pET-28a vector. Gene-specific primers were listed in Supplementary table 2. The recombinant plasmid was transformed into *E. coli* BL21 (DE3) competent cells. Positive transformants were grown in a 5 mL LB medium containing 30 μg/mL kanamycin or 50 μg/mL ampicillin at 37 °C overnight and then subcultured into an 800 mL LB medium containing the corresponding antibiotic. Protein expression was induced by the addition of 0.2 mM isopropyl-β-d-thiogalactoside (IPTG) for 20 h at 16 °C when the optical density (OD) at 600 nm reached 0.6-0.8, harvested by centrifugation at 4 °C and 4,000 rpm for 30 min.

Protein production and crystallization

The cell pellet was suspended in 30 mL of lysis buffer (20 mM HEPES, pH 7.5, 200 mM NaCl, 0.1 mM PLP), disrupted by High-Pressure Homogenizer, and then

centrifuged at 16,000 rpm for 30 min at 4 °C to remove the cell debris. The supernatants

were purified with a Ni-Agarose resin column followed by size-exclusion

chromatography. Before crystallization, purified proteins were concentrated at 10

mg/mL. Crystallization conditions were screened by the sitting-drop vapor diffusion

method using the reservoir solutions supplied in commercially available screening kits

(Crystal Screen, Crystal Screen 2, PEGRx 1, 2, and SaltRx 1, 2). A droplet made by

mixing 1.0 μL of purified AtSerDC or CsAlaDC (10 mg/mL) with an equal volume of

a reservoir solution was equilibrated against 100 μL of the reservoir solution at 16°C.

The crystal of AtSerDC was obtained using buffer pH 8.0 containing 20% (w/v)

PEG400 as a precipitate and 0.2 M $CaCl_2$. The crystal of the CsAlaDC-EA complex

was obtained at pH 7.5 containing 2.6 M sodium acetate. The crystal of CsAlaDC was

obtained at pH 6.0 containing 3.5 M sodium formate.

Data collection and processing

Crystals were grown by sitting-drop vapor diffusion method at 16 °C. The volume of

the reservoir solution was 100 μL and the drop volume was 2 μL, containing 1 μL of

protein sample and 1 μL of reservoir solution. The reservoir solution of AtSerDC

contained 0.1 M Tris-HCl (pH 8.0), 0.2 M $CaCl_2$, and 20% PEG 400. The reservoir

solution of CsAlaDC contained 0.1 M HEPES (pH 7.5), 2.6 M sodium acetate or 0.1 M

Bis-Tris (pH 6.0), and 3.5 M sodium formate. Crystals grew in 3-5 days using a protein

concentration of 10 mg/mL. Diffraction data were collected at Shanghai Synchrotron

Radiation Facility (China). The collected data sets were indexed, integrated, and scaled

using the HKL3000 software package. The structure of AtSerDC was solved by

molecular replacement using the structure of HisDC (PDB code: 7ERV [*Photobacterium phosphoreum*]) as the model, and utilizing AtSerDC as a molecular displacement model for both CsAlaDC and the CsAlaDC-EA complex. The AtSerDC, CsAlaDC, and CsAlaDC-EA complex exhibit resolutions of 2.85 Å, 2.50 Å, and 2.60 Å, respectively. The statistics for data collection and processing are summarized in Supplementary table 1.

Enzyme activity assays

Decarboxylase activity was measured by detecting products (EA or ethanolamine) in Waters Acquit ultraperformance liquid chromatography (UPLC) system (16, 37). The 100 μL reaction mixture, containing 20 mM substrate (Ala or Ser), 100 mM potassium phosphate, 0.1 mM PLP, and 0.025 mM purified enzyme, was prepared and incubated at standard conditions (45 °C and pH 8.0 for CsAlaDC, 40 °C and pH 8.0 for AtSerDC for 10 min). Then, the reaction was stopped with 20 μL of 10% trichloroacetic acid. The product was derivatized with 6-aminoquinolyl-N-hydroxy-succinimidyl carbamate (AQC) and subjected to analysis by UPLC. All enzymatic assays were performed in triplicate.

The detection methodology for theanine production remains consistent with the aforementioned approach. The 100 μL reaction mixture, containing 20 mM Ala, 45 mM Glu, 50 mM HEPES (pH 7.5), 0.1 mM PLP, 30 mM $MgCl_2$, 10 mM ATP, 0.03 mM PsGS/MmGMAS, and 0.025 mM CsAlaDC/CsAlaDC [L110F]/CsAlaDC [L110F/P114A], was prepared and incubated at standard conditions for 1 h. Subsequently, the reaction was terminated via immersion of the reaction vessel in a metal bath at 96 °C for 3 min (31).

Theanine was derivatized with AQC and subjected to analysis by UPLC. All enzymatic

assays were performed in triplicate.

Site-directed mutagenesis

Site-directed mutagenesis experiment was conducted using a PCR method from the

wild-type construct pET-28a-*AtSerDC* and pET-22b-*CsAlaDC*, respectively. *Dpn* I

endonucleases were used to digest the parental DNA template. The reaction mixture

was used to transform *E. coli* DH5α competent cells and the plasmids from positive

strains were extracted to *E. coli* BL21 (DE3) for protein expression, purification, and

analysis of the enzymatic activity.

In vivo* enzyme activity assay in *N. benthamiana

The amplified PCR products were fused to the plant expression vector, pCAMBIA1305.

Linearization was conducted by restriction digest with *Spe* I and *Bam*H I. The

recombinant colonies were selected for PCR validation on the appropriate antibiotics

plate. After validation, the plasmids were electroporated into *Agrobacterium*

tumefaciens strain GV3101(pSoup-p19). Empty pCAMBIA1305 vector, with an intron

containing the GFP gene, was treated as the control.

Agrobacterium transient expression assays were performed on 5-week-old *N.*

benthamiana plants. *Agrobacterium tumefaciens* strain GV3101 (pSoup-p19), carrying

the above-described vectors, were cultured in Luria Bertani (LB) medium, containing

appropriate antibiotics, at 28 °C. When the absorbance of bacteria colonies reached

OD_{600} = 0.6-0.8, bacterial cells were collected and resuspended in MMA solution (10

mM $MgCl_2$, 10 mM 2-(N-morpholino) ethane sulfonic acid (MES), pH 5.6). After the

OD$_{600}$ of the resuspended bacterial solution was adjusted to approx. 1.0, acetosyringone

(AS) was added with a final concentration of 200 μM, and this solution was then

incubated, at room temperature, for at least 3 h in darkness. Next, cell suspensions were

infiltrated into *N. benthamiana* leaves with a needle-free syringe. These *N.*

benthamiana leaves were then collected 3 days post infiltration, frozen in liquid

nitrogen and stored at -80 °C. Internal EA in *N. benthamiana* leaves was extracted as

previously described (18), and then the solvent was subjected to gas chromatography-

mass spectrometry GS-MS system.

Transcript level analysis in *N. benthamiana*

Total RNA was isolated from samples using the RNAprep Pure Plant Kit (Tiangen,

Beijing, China), according to the manufacturer's protocol. The cDNAs were

synthesized using TransScript One-Step gDNA Removal and cDNA Synthesis

SuperMix Kit (TransGen Biotech, Beijing, China). The qRT-PCR assays were

performed, as previously described (38). Primers used for qRT-PCR assays were listed

in Supplementary table 3 and the qRT-PCR was run on a Bio-Rad CFX96TM RT PCR

detection system and CFX Manager Software. Each reaction reagent (20 μL) contained

0.4 μL forward and reverse primers (10 μM), 2 μL cDNA (200±5 ng/μL), 10 μL SYBR

Green Supermix (Vazyme, Nanjing, China) and 6.2 μL double-distilled water. Reaction

was performed by a two-step method: 95 °C for 5 min; 40 cycles of 95 °C for 10 s; and

60 °C for 30 s. The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene was

used for internal normalization in each RT-PCR, and the $2^{-\Delta\Delta CT}$ method (39) was used

to calculate the relative gene expression. All samples were performed with three

replicates.

Multiple sequence alignment

544   MUSCLE was used to generate the protein multiple sequence alignment of amino acid

545   decarboxylases with default settings (40). ESPript 3.x was used to display the multiple

546   sequence alignment (41).

Phylogenetic analysis

The AtSerDC was used as a query for searching its homologs in Embryophyta using

BLASTP (42) (version 2.13.0). To fetch homologs of AtSerDC in distant species, we

performed a two-round blast. We first search for the best hit in the given order under

Embryophyta. Then we blast each species under a provided order using the order best

hit as a query and as many as 10 hits were kept. The sequences with e values <= 0.001

were removed. We further dropped sequences with lengths less than 200 or larger than

800 amino acids. Then multiple sequence alignment was performed via clustalo

(version 1.2.4) with default settings on these filtered sequences (43). The NJ algorithm

implemented in clustalw (version 2.1) was adopted to build the phylogenetic tree (44).

The R package ggtree (version 3.2.1) and universalmotif (version 1.12.4) were used for

tree and motif visualization (45).

Acknowledgments

560   We thank the team of beamline BL18U1 in the Shanghai Synchrotron Radiation Facility

561   for diffraction data collection.

Funding

563   This study is supported by grants from the Ministry of Science and Technology of China

(2019YFA0904100 to W.G, 2022YFF1003103 to X.W) and the Natural Science Foundation of China (T2221005 to W.G, 32072624 to Z.Z).

Data availability

The structures of AtSerDC, CsAlaDC-EA complex and CsAlaDC have been deposited in the Protein Data Bank (PDB) under accession numbers 8JG7, 8JIK and 8JIJ respectively.

References

1. Drew, L. (2019) The growth of tea. *Nature* **566**, S2-S4.

2. Juneja, L. R., Chu, D. C., Okubo, T., Nagato, Y., Yokogoshi, H. (1999) L-theanine - a unique amino acid of green tea and its relaxation effect in humans. *Trends Food Sci. Technol.* **10**, 425-425.

3. Lin, J. Z. *et al*. (2022) Effect of the presence of stem on quality of oolong tea. *Foods* **11**, 3439.

4. Haskell, C. F., Kennedy, D. O., Milne, A. L., Wesnes, K. A., Scholey, A. B. (2008) The effects of L-theanine, caffeine and their combination on cognition and mood. *Biol. Psychol.* **77**, 113-122.

5. Kakuda, T. *et al*. (2000) Protective effect of gamma-glutamylethylamide (theanine) on ischemic delayed neuronal death in gerbils. *Neuro. Sci. Lett.* **289**, 189-192.

6. Liu, Q., Duan, H., Luan, J., Yagasaki, K., Zhang, G. (2009) Effects of theanine on growth of human lung cancer and leukemia cells as well as migration and invasion of human lung cancer cells. *Cytotechnology* **59**, 211-217.

7. Lu, K. *et al*. (2004) The acute effects of L-theanine in comparison with alprazolam on anticipatory anxiety in humans. *Hum. Psychopharmacol.* **19**, 457-465.

8. Takagi, Y. *et al*. (2010) Combined administration of (L)-cystine and (L)-theanine enhances immune functions and protects against influenza virus infection in aged mice. *J. Vet. Med. Sci.* **72**, 157-165.

9. Zheng, G., Sayama, K., Okubo, T., Juneja, L. R., Oguni, I. (2004) Anti-obesity effects of

three major components of green tea, catechins, caffeine and theanine, in mice. *In Vivo* **18**, 55-62.

10. Lin, S. *et al*. (2023) Theanine metabolism and transport in tea plants (*Camellia sinensis* L.): advances and perspectives. *Crit. Rev. Biotechnol*. **43**, 327-341.

11. Deng, W., Ogita, S., Ashihara, H. (2008) Biosynthesis of theanine (γ-ethylamino-l-glutamic acid) in seedlings of *Camellia sinensis*. *Phytochem. Lett.* **1**, 115-119.

12. Deng, W.-W., Ogita, S., Ashihara, H. (2009) Ethylamine content and theanine biosynthesis in different organs of camellia sinensis seedlings. *Z. Naturforsch. C* **64**, 906-906.

13. Cheng, S. *et al*. (2019) Differential accumulation of specialized metabolite L-theanine in green and albino-induced yellow tea (*Camellia sinensis*) leaves. *Food Chem.* 276, 93-100.

14. Takeo, T. (1974) L-alanine as a precursor of ethylamine in *Camellia sinensis*. *Phytochem.* **13**, 1401-1406.

15. Zhu, B. *et al*. (2021) CsAlaDC and CsTSI work coordinately to determine theanine biosynthesis in tea plants (*Camellia sinensis* L.) and confer high levels of theanine accumulation in a non-tea plant. *Plant Biotechnol. J.* **19**, 2395-2397.

16. Bai, P. *et al*. (2021) Biochemical characterization of specific Alanine Decarboxylase (ADC) and its ancestral enzyme Serine Decarboxylase (SDC) in tea plants (*Camellia sinensis*). *BMC Biotechnol.* **21**,17.

17. Han, S. W., Shin, J. S. (2022) Aromatic L-amino acid decarboxylases: mechanistic features and microbial applications. *Appl. Microbiol. Biotechnol.* **106**, 4445-4458.

18. Berger, M., Gray, J. A., Roth, B. L. (2009) The expanded biology of serotonin. *Annu. Rev. Med.* **60**, 355-366.

19. Khan, M. Z., Nawaz, W. (2016) The emerging roles of human trace amines and human trace amine-associated receptors (hTAARs) in central nervous system. *Biomed. Pharmacother.* **83**, 439-449.

20. Ng, J., Papandreou, A., Heales, S. J., Kurian, M. A.   (2015) Monoamine neurotransmitter disorders--clinical advances and future perspectives. *Nat. Rev. Neurol.* **11**, 567-584.

21. Rontein, D. *et al*. (2001) Plants synthesize ethanolamine by direct decarboxylation of serine using a pyridoxal phosphate enzyme. *J. Biol. Chem.* **276**, 35523-35529.

22. Gibellini, F., Smith, T. K. (2010) The Kennedy pathway-De novo synthesis of

phosphatidylethanolamine and phosphatidylcholine. *IUBMB Life* **62**, 414-428.

23. Mudd, S. H., Datko, A. H. (1989) Synthesis of ethanolamine and its regulation in *Lemna*

paucicostata. Plant Physiol. **91**, 587-597.

24. Zinser, E. *et al.* (1991) Phospholipid synthesis and lipid composition of subcellular

membranes in the unicellular eukaryote *Saccharomyces cerevisiae. J. Bacteriol.* **173**,

2026-2034.

25. Yunus, I. S., Liu, Y.-c., Nakamura, Y. (2016) The importance of SERINE

DECARBOXYLASE1 (SDC1) and ethanolamine biosynthesis during embryogenesis of

Arabidopsis thaliana. Plant J. **88**, 559-569.

26. Komori, H. et al. (2012) Structural Study Reveals That Ser-354 Determines Substrate

Specificity on Human Histidine Decarboxylase* *J Biol Chem.* **287**, 29175-83.

27. Huang, J. et al. (2018) Lactobacillus brevis CGMCC 1306 glutamate decarboxylase:

Crystal structure and functional analysis. *Biochem Biophys Res Co.* **503**, 1703-1709

28. Paiardini, A., Giardina, G., Rossignoli, G., Voltattorni, C. B., Bertoldi, M. (2017) New

insights emerging from recent investigations on human group II pyridoxal 5'-phosphate

decarboxylases. *Curr. Med. Chem.* **24**, 226-244.

29. Torrens-Spence, M. P. *et al.* (2020) Structural basis for divergent and convergent evolution

of catalytic machineries in plant aromatic amino acid decarboxylase proteins. *Proc. Natl.*

Acad. Sci. U. S. A. **117**, 10806-10817.

30. Hagihara, R., Ohno, S., Hayashi, M., Tabata, K., Endo, H. (2021) Production of L-theanine

by *Escherichia coli* in the absence of supplemental ethylamine. *Appl. Environ. Microbiol.*

87, e00031-21.

31. Yamamoto, S., Wakayama, M. & Tachiki, T. (2007) Characterization of theanine-forming

enzyme from *Methylovorus mays* no. 9 in respect to utilization of theanine production.

Biosci. Biotechnol. Biochem. **71**, 545-552.

32. Holm, L. (2022) Dali server: structural unification of protein families. *Nucleic. Acids Res.*

50, W210-W215.

33. Okawa, A. *et al.* (2021) Structural basis for substrate specificity of L-methionine

decarboxylase. *Protein Sci.* **30**, 663-677.

34. Fenalti, G. *et al.* (2007) GABA production by glutamic acid decarboxylase is regulated by

a dynamic catalytic loop. *Nat. Struct. Mol. Biol.* **14**, 280-286.

35. Gayathri, S. C., Manoj, N. (2019) Structural insights into the mechanism of internal

aldimine formation and catalytic loop dynamics in an archaeal Group II decarboxylase. *J.*

Struct. Biol. **208**, 137-151.

36. Torrens-Spence, M. P., von Guggenberg, R., Lazear, M., Ding, H., Li, J. (2014) Diverse

functional evolution of serine decarboxylases: identification of two novel acetaldehyde

synthases that uses hydrophobic amino acids as substrates. *BMC Plant Biol.* **14**, 247.

37. Qu, L., Gu, S., Zhang, J., Zhao, C., Deng, X. (2021) Determination of 18 amino acids in

three different kinds of milk powder by ultra performance liquid chromatography coupled

with pre-column derivatization. *Chin. J. Chromatogr.* **39**, 472-477.

38. Madeira, F. *et al.* (2022) Search and sequence analysis tools services from EMBL-EBI in

2022. *Nucleic Acids Res.* **50**, W276-W279.

39. Dong C. *et al.* (2020) Theanine transporters identified in tea plants (*Camellia*

sinensis L.). *Plant J.* **101**, 57-70.

40. Schmittgen T. D. *et al.* (2000) Quantitative reverse transcription-polymerase chain

reaction to study mRNA decay: comparison of endpoint and real-time methods. *Anal.*

Biochem. **285**, 194–204.

41. Robert X., Gouet P. (2014) Deciphering key features in protein structures with the new

ENDscript server. *Nucleic Acids Res.* **42**, W320-W324.

42. Camacho, C. *et al.* (2009) BLAST+: architecture and applications. *BMC Bioinf.* **10**, 421.

43. Sievers, F., Higgins, D. G. (2018) Clustal Omega for making accurate alignments of many

protein sequences. *Protein Sci.* **27**, 135-145.

44. Larkin, M. A. *et al.* (2007) Clustal W and clustal X version 2.0. *Bioinformatics.* **23**, 2947-

2948.

45. Yu, G., Lam, T. T.-Y., Zhu, H., Guan, Y. (2018) Two methods for mapping and visualizing

associated data on phylogeny using ggtree. *Mol. Biol. Evol.* **35**, 3041-3043.

Figures and table

Figure 1. Metabolic pathways and sequence analysis of CsAlaDC and SerDCs in

plants. (**A**) The decarboxylation of serine and alanine in plants. SerDC, serine

decarboxylase; AlaDC, alanine decarboxylase. (**B**) Multiple alignment of the amino

acid sequences of AlaDC and SerDCs. The amino acid sequences of the 6 SerDCs from

kiwifruit (*Actinidia chinensis*), grape (*Vitis vinifera*), coffee (*Coffea eugenioides*),

cocoa (*Theobroma cacao*) and Arabidopsis. Primary (100%), secondary (80%), and

687   tertiary (60%) conserved percent of similar amino acid residues were shaded in deep

688   blue, light blue and cheer red, respectively. "*" indicated amino acid residues mutated

689   only in CsAlaDC.

Figure 2. Purification and characterization of CsAlaDC, AtSerDC, and CsSerDC.

(**A**) Identification of CsAlaDC, AtSerDC, and CsSerDC using SDS-PAGE. (**B**)

Detection of enzyme activities of CsAlaDC, AtSerDC, and CsSerDC by UPLC. (**C**)

Reaction rates of substrates with different concentrations catalyzed by CsAlaDC,

AtSerDC, and CsSerDC.

Figure 3. Crystal structures of CsAlaDC and AtSerDC. (**A**) Dimer structure of

CsAlaDC. The color display of the N-terminal domain, large domain, and C-terminal

domains of chain A is shown in light pink, khaki and sky blue, respectively. Chain B is

shown in spring green. The PLP molecule is shown as a sphere model. The zinc finger

structure at the C-terminus of CsAlaDC is indicated by the red box. The gray spheres

represent zinc ions, while the red dotted line depicts the coordination bonds formed by

zinc ions with cysteine and histidine. (**B**) The 2Fo-Fc electron density maps of K309-

PLP-EA (contoured at 1σ level). The PLP is shown in violet, the K309 is shown in

spring green, and the EA is shown in lightblue. (**C**) Active center of the CsAlaDC-EA

complex, with hydrogen bonds denoted by black dotted lines. "*" denotes the amino

acids on adjacent subunits. (**D**) Dimer structure of AtSerDC. The color display of the

N-terminal domain, large domain, and C-terminal domains of chain A is shown in light

pink, khaki and sky blue, respectively. Chain B is shown in cyan. The PLP molecule is

shown as a sphere model. The zinc finger structure at the C-terminus of AtSerDC is

indicated by the red box. The gray spheres represent zinc ions, while the red dotted line

depicts the coordination bonds formed by zinc ions with cysteine and histidine. (**E**)

Active center of the AtSerDC, with hydrogen bonds denoted by black dotted lines. "*"

denotes the amino acids on adjacent subunits. (**F**) The monomers of CsAlaDC and

AtSerDC are superimposed. CsAlaDC is depicted in spring green, while AtSerDC is

shown in plum. The conserved amino acid catalytic loop is indicated by the red box.

(**G**) Amino acid residues of the active center in CsAlaDC apo and CsAlaDC-EA

complex are superimposed. CsAlaDC apo is shown in floral white, while CsAlaDC-EA

complex is shown in spring green. (**H**) The relative activity of wild-type CsAlaDC and

its Y336F mutant (left), as well as wild-type AtSerDC and its Y341F mutant (right) is

shown.

Figure 4. Key amino acid residues for substrate recognition. (A) Superposition of

substrate binding pocket amino acid residues in CsAlaDC and AtSerDC. The amino

acid residues of CsAlaDC are shown in spring green, the amino acid residues of

AtSerDC are shown in plum, with the substrate specificity-related amino acid residue

highlighted in a red ellipse. **(B)** Active-site-lining amino acid residues of SDC

homologs from Embryophyta were identified. The height of each amino acid is scaled

proportionally to the amount of information content (measured in bits). The first line

depicts the conserved motif in all SerDC homologs from Embryophyta, whereas lines

2-5 represent the conserved motifs based on the variable third amino acid residue. **(C)**

Histogram showing the distribution of the number of key motifs. **(D)** Histogram

showing the number of key motifs in different plant orders. (**E**) Relative enzyme

activities of wild-type CsAlaDC and mutant protein CsAlaDC^{F106Y} against Ala

substrate (columns 1 and 2), and enzyme activities of wild-type AtSerDC and various

AtSerDC mutant proteins against Ala substrate (columns 3-9) are presented. The

percentage graph shows the relative activity of each protein compared to wild-type

CsAlaDC activity (taken as a 100% benchmark). (**F**) Relative enzyme activities of wild-

type AtSerDC and AtSerDC mutant proteins (columns 1-7) against Ser substrates, and

enzyme activities of wild-type CsAlaDC and mutant protein CsAlaDC^{F106Y} against Ser

substrates(columns 8, 9) were measured. The percentage graph shows the relative

activity of each protein compared to the wild-type AtSerDC (taken as a 100%

benchmark). Three independent experiments were conducted. (**G**) The EA contents of

AtSerDC and its mutant AtSerDC^{Y111F} in *N. benthamiana*. (**H**) The EA contents of

CsAlaDC and its mutant CsAlaDC^{F106Y} in *N. benthamiana*. The significance of the

difference ($P<0.05$) was labeled with different letters according to Duncan's multiple

range test.

Figure 5. Mutations enhance CsAlaDC enzyme activity and theanine synthesis *in*

***vitro*. (A)** Relative enzyme activities of CsAlaDC mutant proteins against Ala substrate.

(B) Relative enzyme activities of CsAlaDC^{L110F}, CsAlaDC^{P114A}, and

CsAlaDC$^{L110F/P114A}$ against Ala substrate. **(C)** Histogram showing the relative content

of theanine resulting from different combinations of alanine decarboxylase and

theanine synthetase. Three independent experiments were conducted.

758 **Table 1.** Kinetic parameters of CsAlaDC, AtSerDC and CsSerDC

Enzyme	Substrate	Km (mM)	$Vmax$ (μmol L^{-1} s^{-1})	$Kcat$ (s^{-1})	$Kcat/Km$ (s^{-1} M^{-1})
CsAlaDC	Alanine	1.215 ± 0.237	1.709 ± 0.072	0.068	55.7
AtSerDC	Serine	1.522 ± 0.297	7.053 ± 0.297	0.281	184.6
CsSerDC	Serine	2.364 ± 0.431	9.031 ± 0.520	0.361	152.7

Supplementary figure 1. Purification of CsAlaDC, AtSerDC, and CsSerDC and

Crystal structures of CsAlaDC and AtSerDC. (**A**) Comparison of elution profiles of

CsAlaDC, AtSerDC and CsSerDC. (**B**) Monomer structure of CsAlaDC. (**C**) Monomer

structure of AtSerDC. The color display of the N-terminal domain, large domain, and

C-terminal domains is shown in light pink, khaki and sky blue, respectively. The PLP

molecule is shown as a sphere model. The gray spheres represent zinc ions.

Supplementary figure 2. Catalytic mechanisms and conformational changes of CsAlaDC. After the transaldimation of the internal aldimine within CsAlaDC, resulting in the release of the active-site residue Lys309, the PLP amino acid external aldimine undergoes decarboxylation, leading to the removal of the α-carboxyl group as CO2. This process generates a quinonoid intermediate that is stabilized by the delocalization of paired electrons (1,2). The carbanion at Cα is subsequently protonated by the acidic p-hydroxyl group of Tyr336* located on the large loop, facilitated by its neighboring residue His196, which is situated on the small loop (3,4). Simultaneously, the internal aldimine LLP309 in CsAlaDC is restored, resulting in the release of the product (5).

Supplementary figure 3. Evolutionary analysis of CsAlaDC in Embryophyta. (A)

The presented diagram depicts an evolutionary tree of CsAlaDC, which is devoid of a

root and solely portrays the topological structure of the tree without including distance

information. The color of the inner ring corresponds to various orders, while the outer

ring's leaf nodes are colored based on the motif types that the sequence exhibits. **(B)**

Diversity of serine decarboxylase-like proteins in Embryophyta (196 species). Colored

scatter spots on the right side of leaf nodes correspond to the respective motifs shown

in Figure A.

Supplementary figure 4. The relative mRNA levels of AtSerDC and its mutant Y111F

(*A*), CsAlaDC and its mutant F106Y (*B*) in *N. benthamiana* leaves were measured by

two primers. WT, wild type of *N. benthamiana*; EV, empty vector control; *NbGAPDH*

was used as an internal control. Data represent mean ± SD (n=3). The significance of

the difference (*P*<0.05) was labeled with different letters according to Duncan's

multiple range test.

Supplementary figure 5. Structures of HisDC2, MetDC, TrpDC, AspDC, HisDC1

and GluDC. (**A**) The Overall Structures of HisDC2, MetDC, TryDC, AspDC, HisDC1

and GluDC. Chain A is shown in khaki, chain B is shown in cyan. (**B**) Amino acid

residues in the substrate binding pocket of HisDC2, MetDC, TryDC, AspDC, HisDC1

and GluDC. The amino acid residues in chain A are shown in khaki, and the amino acid

residues in chain B are shown in cyan.

Supplementary figure 6. Multiple sequence alignment of CsAlaDC, AtSerDC, MetDC,

HisDC1, TrpDC, HisDC2, TyrDC, and GluDC were generated using MUSCLE and

visualized with ESPript 3.x. Conserved amino acid residues in all eight proteins are

highlighted with red backgrounds. The magenta box marks key amino acid residues

involved in substrate recognition for CsAlaDC and AtSerDC. The Lys residue

covalently bound to the PLP cofactor is denoted by a red star, while the green triangle

indicates the Tyr residue associated with enzymatic activity. Amino acid residues

involved in the CsAlaDC substrate binding pocket are marked with blue circles.

Supplementary figure 7. Circular Dichroism Spectra of proteins. (**A**) Circular Dichroism Spectra of CsAlaDC (WT). (**B**) Circular Dichroism Spectra of CsAlaDC^{Y336F}. (**C**) Circular Dichroism Spectra of CsAlaDC^{F106Y}. (**D**) Circular Dichroism Spectra of AtSerDC (WT). (**E**) Circular Dichroism Spectra of AtSerDC^{Y341F}. (**F**) Circular Dichroism Spectra of AtSerDC^{Y111F}.

Supplementary figure 8. Absorption Spectra of different proteins. (**A**) Absorption

Spectra of CsAlaDC (WT), CsAlaDC^{Y336F} and CsAlaD^{F106Y}. (**B**) Absorption Spectra of

AtSerDC (WT), AtSerDC^{Y341F} and AtSerDC^{Y111F}.

832 **Supplementary figure 9. Purification of truncated CsAlaDC and AtSerDC**. Lane 1

833 represents marker. The precipitated and eluted samples of AtSerDC with a truncated

834 zinc finger structure are shown in lanes 2 and 3 respectively, the precipitated and eluted

835 samples of CsAlaDC with a truncated zinc finger structure are illustrated in lanes 4 and

836 5 respectively.

Supplementary tables

Supplementary table 1. Data collection and refinement statistics

	CsAlaDC	CsAlaDC-EA complex	AtSerDC
Data Collection			
Space Group	*P3121*	*P3121*	*C2*
Cell Dimensions			
a, b, c (Å)	163.5,163.5, 50.2	164.0,164.0,50.2	115.4,149.5,139.1
α, β, γ (°)	α=β=90, γ=120	α=β=90, γ=120	α=γ=90, β=109.2
Wavelength (Å)	0.979	0.979	0.979
Resolution (Å)	50-2.50	50-2.60	50-2.85
Completeness (%)	99.3 (99.9)	99.6 (100.0)	99.6 (98.1)
Redundancy	5.4 (5.1)	13.0 (12.5)	5.9 (5.0)
$<I>/<\sigma I>$	9.3 (1.9)	18.9 (4.6)	10.6 (2)
R_{merge}	0.128 (0.603)	0.156 (0.562)	0.173 (0.551)
Refinement			
R_{work}/ R_{free}	0.173/0.222	0.172/0.223	0.214/0.262
No. reflections	25175	22813	49543
B-facator ($Å^2$)			
Protein	43.3	48.3	57.6
PLP	29.5	31.5	52.8
Zn^{2+}	58.7	43.8	71.6
Ca^{2+}	-	55.5	-
Ethylamine	-	27.5	-
Glycerol	-	-	51.7
Water	38.5	40.9	-
R.m.s. deviations			
Bond lengths (Å)	0.008	0.008	0.007
Bond angles (°)	1.531	1.601	1.410
No. of atoms			
Protein	3292	3265	3250
PLP	15	15	60
Zn^{2+}	2	1	4
Ca^{2+}	-	1	-
Ethylamine	-	5	-
Glycerol	-	-	24
Water	115	60	-
Ramachandran statistics (%)			
Favoured	94.01	93.58	91.52
Allowed	5.74	5.68	7.61
Outliers	0.25	0.74	0.87

$R_{merge}=\sum hkl\sum i|Ii(hkl)-<I(hkl)>|/\sum hkl\sum i\ Ii(hkl)$, where$<I(hkl)>$is the main value of I(hkl).

$R_{work}=\sum||Fobs|-|Fcalc||/\sum|Fobs|$, where Fobs and Fcalc are observed and calculated structure factors.

The free R factor was calculated using 5% of reflections ommitted from the refinement.

845   Highest resolution shell is shown in parenthesis

848   **Supplementary table 2.** Primers used for gene cloning.

Name	Forward primers (5'-3')	Reverse primers (5'-3')
CsAlaDC	GGAATTCCATATGACAACAAGCCTCACCATCACGG	GATCTCGAGTCATTTATGAAGATCACAATCACAATTCTCACTTC
AtSerDC	GGAATTCCATATGACCACCAGCCTGGCCG	GATCTCGAGTTATTTATGGGCCGGACAAATGCAATTATTG
CsSerDC	ATGGTGGGAAGTGTTGGGGT	TCACTTGTGTAGTGCACAAAGACAGT

851   **Supplementary table 3.** Primers used for real-time PCR.

Name	Forward primers (5'-3')	Reverse primers (5'-3')
CsAlaDC	CACTGTGATGGGGCTCTGTT	TGTTATCTGGACACCGCACG
AtSerDC	TCACTCGCTGTAACGGAACC	AACTGACCAAGCGCACCATA
NbGAPDH	AGCTCAAGGGAATTCTCGATG	AACCTTAACCATGTCATCTCCC