Structure and evolution of Alanine/Serine Decarboxylases and the engineering of theanine production

Hao Wang[a,1], Biying Zhu[b,1], Siming Qiao[b], Chunxia Dong[b], Xiaochun Wan[b], Weimin Gong[a,*], Zhaoliang Zhang[b,*]

[a]Department of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[b]State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, Hefei, Anhui 230036, China.

[1]These authors contributed equally to this work.

* To whom correspondence should be addressed. Email: zhlzhang@ahau.edu.cn or wgong@ustc.edu.cn.

Author contributions: W.G, Z.Z and X.W designed research; H.W, B.Z, S.Q and C.D performed research; H.W and B.Z analyzed data; H.W, B.Z, W.G, and Z.Z wrote the paper.

Competing interest statement: The authors declare no competing financial interests.

Classification: Biochemistry (major), Plant Sciences (minor)

Keywords: alanine decarboxylase, serine decarboxylase, substrate specificity, theanine, *Camellia sinensis*

Abstract

Ethylamine (EA), the precursor of theanine biosynthesis, is synthesized from alanine decarboxylation by Alanine Decarboxylase (AlaDC) in tea plants. AlaDC evolves from Serine Decarboxylase (SerDC) through neofunctionalization and has lower catalytic activity. However, lacking structure information hinders the understanding of the evolution of substrate specificity and catalytic activity. In this study, we solved the X-ray crystal structures of AlaDC from *Camellia sinensis* (CsAlaDC) and SerDC from *Arabidopsis thaliana* (AtSerDC). Tyr[341] of AtSerDC or the corresponding Tyr[336] of CsAlaDC is essential for their enzymatic activity. Tyr[111] of AtSerDC and the corresponding Phe[106] of CsAlaDC determine their substrate specificity. Both CsAlaDC and AtSerDC have a distinctive zinc finger that contributes to the activity and has not been identified in any other Group II PLP-dependent amino acid decarboxylases. Based on the structure comparison and evolution analysis AlaDC or SerDC led us to perform mutation screen of CsAlaDC. The results indicated that the mutation of L110F or P114A in the CsAlaDC dimerization interface significantly improved the catalytic activity by 110% and 59%, respectively. Combining double mutant CsAlaDC[L110F/P114A] with theanine synthetase increased 672% of theanine production in an *in vitro* system. This study provides the structural basis for the substrate selectivity and catalytic activity of CsAlaDC and AtSerDC and sheds light on a new direction for the efficient industrial synthesis of theanine.

Introduction

Tea is one of the most popular natural non-alcoholic beverages consumed worldwide. About 2 billion cups of tea are consumed worldwide daily (1). The popularity of tea is determined by its favorable flavor and numerous health benefits (2-4). The flavor and health-beneficial effects of tea are conferred by the abundant secondary metabolites, including catechins, caffeine, theanine, volatiles, etc (5). Theanine (γ-glutamylethylamide) is a unique non-protein amino acid and the most abundant free amino acid in tea plants (*Camellia sinensis*). It accounts for more than 50% of the total free amino acids and approximately 1-2% of the dry weight of the new shoots of tea plants (6). Theanine is the secondary metabolite conferring the umami taste of tea infusion and also balances the astringency and bitterness of tea infusion caused by catechins and caffeine (7). It has also many health-promoting functions, such as reducing stress and anxiety, improving mood and cognition, protecting the nervous system, etc (6, 8-13). Therefore, theanine content is highly correlated with green tea quality (14).

Theanine is synthesized from ethylamine (EA) and glutamate (Glu) by theanine synthetase (15, 16). Importantly, the large amount of theanine biosynthesis is determined by the high availability of EA in tea plants (17). Therefore, the evolution of EA biosynthesis in tea plants provided basis for theanine biosyntheis and the formation of tea quality. EA is synthesized from alanine decarboxylation by alanine decarboxylase (CsAlaDC) (18) (Figure 1A). Indeed, CsAlaDC expression level and catalytic activity largely determine the theanine accumulation level in tea plants (19). As a novel gene,

CsAlaDC had not been reported in other plant species before it was identified in tea

plants. Previous studies indicated that *AlaDC* originated from the serine decarboxylase

gene (*SerDC*) by gene duplication in tea plants (20). However, despite sharing highly

conserved amino acid sequences, CsAlaDC and CsSerDC specifically catalyze the

decarboxylation of alanine and serine, respectively (20) (Figure 1A, 1C). In addition,

CsAlaDC exhibited a much lower enzymatic activity compared with CsSerDC (20).

However, the structural basis and key amino acids underlying the evolution of the

substrate specificity and enzymatic activity of CsAlaDC are unknown.

SerDC belongs to the Group II pyridoxal-5-phosphate (PLP)-dependent

decarboxylase superfamily (21). These Group II amino acid decarboxylases produce

many bioactive secondary metabolites and signaling molecules (22-24). In plants,

SerDC catalyzes the biosynthesis of ethanolamine from serine, which was first

functionally characterized in *Arabidopsis thaliana* (25) (Figure 1A). Ethanolamine is

an important metabolite in the synthesis of phosphatidylethanolamine and

phosphatidylcholine, two main phospholipids involved in maintaining eukaryotic

membrane structures in eukaryotic cell membranes (26-28). Furthermore, the study has

shown that SerDC is essential in the embryogenesis of *Arabidopsis* (29). SerDC is of

vital importance for plant growth and development, but the mechanism of substrate

recognition and catalytic activity of SerDC also remains unclear.

In this work, we attempt to understand the mechanism of functional divergence of

plant AlaDC and SerDC by analyzing their structural characteristics. Here, we obtained

the X-ray crystal structure of CsAlaDC and AtSerDC. According to the crystal structure,

we found a distinctive zinc finger structure located at both CsAlaDC and AtSerDC,

which has not been identified in any other Group II PLP-dependent amino acid

decarboxylases that have been previously characterized. By comparing the substrate

binding pockets, we identified Phe[106] of CsAlaDC and Tyr[111] of AtSerDC as the crucial

sites for substrate specificity. By conducting mutation screening based on the protein

structures, we identified the amino acids repressing the catalytic activity and discovered

that CsAlaDC[L110F/P114A] exhibited a 2.3-fold increase in catalytic activity compared to

that of CsAlaDC and enhanced the engineering of theanine production *in vitro*.

Results

Enzymatic properties of CsAlaDC, AtSerDC, and CsSerDC

CsAlaDC originates from CsSerDC by gene duplication and neofunctionalization in tea

plants, but they catalyze different metabolic processes (Figure 1A). A phylogenetic tree

was constructed to further reveal the evolutionary relationship between CsAlaDC and

its homologs in *Algal*, *Pteridohyta*, *Bryophyta*, *Monocots* and *Dicots* (Figure 1B).

Phylogenetic analysis indicated that CsAlaDC is homologous with SerDCs in Dicots

plants. As expected, CsSerDC was closer to AtSerDC, which implies that they shared

similar functions. However, CsAlaDC is relatively distant from CsSerDC. Amino acid

sequence similarity analysis also showed that CsSerDC and AtSerDC had higher

smilarity than that between CsAlaDC and CsSerDC (Supplementary table 1). Next, we

performed a multiple sequence alignment of the amino acid sequences of the 6 SerDCs

from kiwifriut (*Actinidia chinensis*), grape (*Vitis vinifera*), coffee (*Coffea eugenioides*),

cocoa (*Theobroma cacao*), Arabidopsis and tea plants and CsAlaDC (Figure 1C). The results showed that the amino acid sequences of CsAlaDC and SerDCs were highly conserved, but CsAlaDC has amino acid mutations in highly conserved regions compared with SerDCs.

Next, to verify the substrate specificity and enzyme activity, we conducted enzyme activity detection and enzyme kinetics analysis for CsAlaDC, CsSerDC and AtSerDC. The 5′ truncated *CsAlaDC* and *AtSerDC* were constructed into the pET22b and pET28a expression vectors to generate recombinant plasmids pET22b-*CsAlaDC* and pET28a-*AtSerDC*, correspondingly. Additionally, the full-length protein of CsSerDC was inserted into the pET28a expression vector to generate recombinant plasmid pET28a-*CsSerDC*. Subsequently, the recombinant proteins were expressed in *Escherichia coli* and purified via nickel affinity chromatography. The purified proteins were examined through SDS-PAGE analysis (Figure 2A). To investigate the substrate specificity of CsAlaDC, AtSerDC, and CsSerDC, enzyme activity assays were conducted using Ala and Ser as substrates, followed by product identification via UPLC analysis. The results showed that CsAlaDC can effectively catalyzed the decarboxylation of alanine to generate EA, whereas its catalytic efficacy towards serine was relatively inferior. Conversely, AtSerDC and CsSerDC selectively catalyzed serine decarboxylation to yield ethanolamine but did not exhibit activity towards alanine (Figure 2B). These findings confirmed that CsAlaDC and SerDCs do not have promiscuous decarboxylase activity on Ala and Ser.

The kinetic properties of CsAlaDC, AtSerDC, and CsSerDC were determined

through the use of corresponding substrates (Figure 2C), and the results are presented

in Table 1. The kinetic parameter Km of CsAlaDC was determined to be 1.215 mM,

which is similar to that of AtSerDC and CsSerDC at 1.522 mM and 2.364 mM,

respectively. However, AtSerDC has a Vmax of 7.053 μmol/L/s, which is 4-fold that of

CsAlaDC's Vmax of 1.709 μmol/L/s; while CsSerDC has a Vmax of 9.031 μmol/L/s,

which is 5.3-fold that of CsAlaDC. These findings indicated that the catalytic efficiency

of CsAlaDC is considerably lower than that of both CsSerDC and AtSerDC. The

observed disparity in Vmax suggests that CsAlaDC and SerDCs may exhibit

discrepancies in substrate-to-product conversion rates under saturated substrate

conditions. It is plausible that SerDCs possess a higher turnover rate, facilitating the

rapid conversion of substrate to product and subsequent release of the enzyme. Another

possibility is that SerDCs demonstrate enhanced stability, thereby maintaining superior

catalytic activity even at higher substrate concentrations, consequently leading to a

higher Vmax value. Due to the similarity in catalytic activity between CsSerDC and

AtSerDC, we chose the more representative AtSerDC for further analysis.

The overall structures of CsAlaDC and AtSerDC

To enhance our comprehension of CsAlaDC and AtSerDC, we conducted structural

analyses of these two proteins. Through optimization of crystallization conditions, we

successfully determined the crystal structures of CsAlaDC, CsAlaDC-EA, and

AtSerDC at resolutions of 2.50 Å, 2.60 Å and 2.85 Å, respectively (Supplementary

table 2).

The overall structure of CsAlaDC and AtSerDC is homodimeric with two subunits

exhibiting an asymmetrical arrangement (Figure 3A and 3D). The monomer of

CsAlaDC or AtSerDC is divided into three distinct structural domains: an N-terminal

domain (N-terminal -104 aa in CsAlaDC, N-terminal -109 aa in AtSerDC), a large

domain (105-364 aa in CsAlaDC, 110-369 aa in AtSerDC), and a small C-terminal

domain (365 aa-C-terminus in CsAlaDC, 370-C-terminus as in AtSerDC), colored light

pink, khaki, and sky blue, respectively. Compared to the full-length protein, the N-

terminal structures of CsAlaDC and AtSerDC were truncated by 60 and 65 amino acid

residues, respectively. The N-terminal domain of the truncated protein contains a long

α-helix, which is connected to the large domain through a long loop. The large domain

contains a seven-stranded mixed β-sheet surrounded by eight α-helices, while the C-

terminal domain is composed of a three-stranded antiparallel β-sheet and three α-helices

(Supplementary figure 1). The catalytic site of the enzyme is positioned within a

superficial crevice at the junction of two subunits forming a dimeric arrangement. The

amino acid residues derived from both subunits participate in the binding of the PLP,

effectively anchoring it to the active site. In the catalytic site, one monomer

accommodates PLP, while the other monomer also contributes to PLP binding and

enzymatic function (Figure 3A and 3D).

Notably, our investigation has revealed the presence of a distinctive zinc finger

structure (as depicted in Figure 3A and 3D) located at the C-terminus of CsAlaDC and

AtSerDC. This structure is composed of a loop structure spanning 17 amino acid

residues, wherein coordination of Zn^{2+} is facilitated by three Cys residues and one His

residue. Importantly, this particular configuration is exclusive to the two proteins under

examination and has not been identified in any other Group II PLP-dependent amino

acid decarboxylases that have been previously characterized (30, 31).

The crystal structures of CsAlaDC-EA complex and AtSerDC were obtained and

further analyzed. In the former complex, PLP bound to Lys[309] via a Schiff base linkage

(internal aldimine form), and the pyridine moiety of PLP is positioned between the

imidazole ring of His[196] and the methyl group of Ala[279] in a parallel orientation to the

imidazole ring. The carboxylic group of Asp[277] stabilizes the N1 of PLP by a salt bridge

interaction, providing the latter with the strong electrophilicity necessary to stabilize

carbon ion intermediates during enzymatic catalysis. The O3 forms hydrogen bonds

with Thr[247] and Lys[309]. Additionally, His[308], Gly[169], Thr[170], Lys[309], and Ser[347]*("*"

denotes the amino acids on adjacent subunits) establish a hydrogen bonding network

with the phosphate group of PLP, consequently fortifying its attachment to the protein.

(Figure 3C). Similarly, in the latter structure, the pyridine ring of PLP was sandwiched

between the imidazole ring of His[201] and the methyl group of Ala[284]. The N1 of PLP

formed a salt bridge with Asp[282], while the O3 formed hydrogen bonds with Thr[252]. The

phosphate group of PLP established hydrogen bonding interactions with Lys[314], His[313],

Gly[174], Thr[175], and Ser[352]* (Figure 3E). The aforementioned amino acid residues are

observed across various Group II PLP-dependent amino acid decarboxylases and

exhibit considerable stability (see further discussion below), suggesting that they could

share common features in terms of catalytic mechanisms.

Factors affecting CsAlaDC and AtSerDC activity

The Group II PLP-dependent amino acid decarboxylases are dimeric enzymes with

each monomer containing an active site located within a shallow cavity (32). A flexible

loop of amino acids originating from one monomer extends into the active site of the

other monomer and plays a crucial role in catalysis. The catalytic loop in CsAlaDC,

consisting of amino acid residues 328-341, is well-structured and exhibits clear electron

density. However, the corresponding loop in AtSerDC is disordered (Figure 3F). The

aforementioned loop harbors a conserved Tyr residue that is believed to function as a

proton donor of the carbanion in the catalytic process (33). In particular, CsAlaDC

Tyr^{336} and AtSerDC Tyr^{341} correspond to that Tyr residue (Supplementary figure 2).

Upon superimposing the active site residues of CsAlaDC apo and CsAlaDC-EA

complex, it was observed that the hydroxyl group of $Tyr^{336}*$ in the CsAlaDC-EA

complex exhibits a substantial 60-degree deviation along the $C\alpha$-$C\beta$ bond (Figure 3G).

To assess the function of the Tyr, we substituted the corresponding Tyr residues in

CsAlaDC and AtSerDC with Phe. The resulting $CsAlaDC^{Y336F}$ and $AtSerDC^{Y341F}$

mutants were then exposed to L-alanine and L-serine, respectively. We measured the

production of EA or ethanolamine in the reaction mixtures. Our findings indicate that

these mutants catalyze abortive decarboxylation, as opposed to wild-type CsAlaDC and

AtSerDC, as evidenced by the absence of detectable EA and only a small amount of

ethanolamine observed in the reaction mixture (Figure 3H). This result suggested this

Tyr is required for the catalytic activity of CsAlaDC and AtSerDC.

Identification of key amino acids for the substrate specificity

Through the superposition of the amino acid residues present in the substrate binding

pocket of CsAlaDC and AtSerDC, it was observed that the residues were identical,

except for Phe at position 106 in CsAlaDC (Figure 4A). In AtSerDC, this position is

occupied by Tyr at 111, implying that this specific amino acid residue is related to

substrate specificity.

To obtain more insights into the role of specific amino acid residue in substrate

specificity, we utilized amino acid sequences of AtSerDC to identify all potential serine

decarboxylases in *Embryophyta* and constructed phylogenetic trees (Supplementary

figure 3). We further identified conserved motifs in these SerDC homologs in

Embryophyta (Figure 4B). Importantly, the Phe^{106} of CsAlaDC or Tyr^{111} of AtSerDC

locates in a conserved motif (Figure 4B). In this conserved motif, the first two amino

acids Y and P are 100% conserved, but the third amino acid is variable and can be Y, T,

F, V, A, I, or other amino acids (Figure 4B and C); interestingly, the Phe^{106} of CsAlaDC

and Tyr^{111} of AtSerDC locates at this variable amino acid position. For these first three

amino acids, the YPY was identified in 83.7% of the SerDC homologs, while the YPF

motif was found in 2.3% of these SerDC homologs (Figure 4C). These analyses

provided more insights into the role of Phe^{106} of CsAlaDC and Tyr^{111} of AtSerDC in the

substrate specificity.

To verify the role of Phe^{106} of CsAlaDC or Tyr^{111} of AtSerDC in substrate specificity,

we mutated Phe[106] of CsAlaDC into Tyr (CsAlaDC[F106Y]) and Tyr[111] of AtSerDC into

Phe (AtSerDC[Y111F]) and tested enzymatic activity *in vitro*. The results showed that

CsAlaDC[F106Y] totally lost the alanine decarboxylase activity, and AtSerDC[Y111F] gained

alanine decarboxylase activity (Figure 4E). We also performed other mutations of Tyr[111]

of AtSerDC, including AtSerDC[Y111A], AtSerDC[Y111I], AtSerDC[Y111L], AtSerDC[Y111V] and

AtSerDC[Y111W]. Within these mutations, only AtSerDC[Y111I] exhibited a low level of

alanine decarboxylase activity (Figure 4E). These results verified that the Phe[106] of

CsAlaDC is required for the alanine decarboxylase activity from serine decarboxylase

activity.

On the other side, both CsAlaDC and CsAlaDC[F106Y] exhibit a very low level of

serine decarboxylase activity (Figure 4F). Moreover, compared with AtSerDC, the

serine decarboxylase activity of AtSerDC[Y111F] was about 30% of AtSerDC;

AtSerDC[Y111I] retained lower than 5% of serine decarboxylase activity of AtSerDC;

while other mutations, including AtSerDC[Y111A], AtSerDC[Y111L], AtSerDC[Y111V] and

AtSerDC[Y111W] abolished the serine decarboxylase activity (Figure 4F). These results

suggested the Tyr[111] of AtSerDC is also important for the substrate specificity of

AtSerDC.

To further verify that Phe[106] of CsAlaDC and Tyr111 of AtSerDC were key amino

acid residues determining its substrate recognition *in planta*, *Nicotiana benthamiana*

transient expression system was carried out. To this end, *A. tumefaciens* strain GV3101

(pSoup-p19), carrying recombinant plasmid, was infiltrated into leaves of 5-week-old

N. benthamiana plants, and the pCAMBIA1305 empty vector was used as the control

(EV). Relative mRNA levels were detected (Supplementary figure 4), indicating that

they had been successfully overexpressed in tobacco. Here, we found that a high level

of EA was produced in the CsAlaDC-expressing tobacco leaves; while no EA product

was detected in tobacco leaves infiltrated with the mutant CsAlaDC^{F106Y}. In addition,

we did not detect EA products in the AtSerDC-expressing tobacco leaves, while a high

level of EA was detected in tobacco leaves infiltrated with mutant AtSerDC^{Y111F}. As

anticipated, EA was not detected in tobacco leaves infiltrated with EV (Figure 4G and

H). These results further verified the critical role of Phe^{106} in the substrate specificity

of CsAlaDC.

Key amino acids for the evolution of CsAlaDC enzymatic activity

CsAlaDC and AtSerDC have a high sequence similarity of 74.5% and a nearly identical

structure with an RMSD (root mean square deviation) of only 0.77Å for monomer

structures. However, the two enzymes catalyze different amino acid decarboxylation

reactions, and AtSerDC exhibits significantly higher activity than CsAlaDC (Figure 2B

and C). To elucidate this observation, we proposed a prediction test: EA, generated via

Ala decarboxylation by CsAlaDC, can be toxic and harmful to plants if accumulated

excessively. Thus, during the evolution of plant serine decarboxylase into alanine

decarboxylase, the enzyme has evolved not just to alter the substrate preference but also

to reduce catalytic activity to control EA production within a suitable range. Based on

this hypothesis, we suggest that mutating specific amino acids in CsAlaDC to those

corresponding amino acids in SerDCs could enhance its activity, and the results could

provide insights into the evolution of CsAlaDC enzymatic activity.

Dimerization is essential to AlaDC/SerDC activities because the active site is

composed of residues from two monomers. To identify the amino acids repressing the

enzymatic activity of CsAlaDC during evolution from SerDC, we analyzed the crystal

structures and the amino acids at the dimer interface between CsAlaDC and AtSerDC.

This analysis revealed that the amino acid residues at positions 66, 97, 110, 114, 116,

117, 122, 315 and 345 are different in CsAlaDC and SerDCs (Figure 1C). Therefore,

we mutated these amino acids of CsAlaDC into those in the corresponding positions of

AtSerDC and CsSerDC, and performed enzyme activity assays (Figure 5A). The results

demonstrated that CsAlaDC^{L110F} and CsAlaDC^{P114A} exhibited significantly enhanced

enzyme activity compared to the wild-type CsAlaDC, with a 2.1-fold and 1.59-fold

increase, respectively. Furthermore, the catalytic activity of the CsAlaDC$^{L110F/P114A}$

double mutant exhibits a remarkable 2.3-fold increase compared to that of the wild-

type protein (Figure 5B). These findings suggested a critical role of Leu^{110} and Pro^{114}

of CsAlaDC in the evolution of enzymatic activity and provided a basis to improve

CsAlaDC activity. It is possible that these amino acid residues could potentially

augment the hydrophobic nature of the protein dimer interface.

***In Vitro* synthesis of theanine**

The biosynthetic pathway of theanine in tea plants comprises two sequential enzymatic

steps: alanine decarboxylase mediates alanine decarboxylation to yield EA, which is

subsequently utilized by theanine synthetase to effectuate the condensation reaction

with glutamate to synthesize theanine. By conducting mutation screening, we discovered a CsAlaDC mutant protein (L110F/P114A) that exhibited a 2.3-fold higher catalytic activity compared to the wild-type protein. Subsequently, we employed an *in vitro* theanine synthesis system utilizing CsAlaDC and either glutamine synthetase (PtGS [P*seudomonas syringae* pv. syringae]) or gamma-glutamate methylamine ligase (GMAS [*Methylovorus mays*]) which have the activity to synthesize theanine from EA and Glu (34, 35).

The results illustrated the successful synthesis of theanine using CsAlaDC in conjunction with two theanine synthetases, employing Ala and glutamate as substrates (Figure 5C). The theanine content generated by the combination of CsAlaDC^{L110F} and GMAS, as well as CsAlaDC$^{L110F/P114A}$ and GMAS, was 4.57-fold and 6.72-fold higher than the content produced in the combination of wild-type CsAlaDC and GMAS, respectively. Similarly, when combined with the PsGS, comparable outcomes are observed as well. The theanine content resulting from the combination of CsAlaDC^{L110F} and PsGS, as well as CsAlaDC$^{L110F/P114A}$ and PsGS, exhibit enhancements of 1.62-fold and 4.33-fold compared to the wild-type protein combination, respectively (Figure 5C). Hence, the utilization of CsAlaDC$^{L110F/P114A}$ could effectively enhance the theanine production yield and thus, holds potential for large-scale engineering production of theanine.

Discussion

Comparison of CsAlaDC and AtSerDC with other decarboxylases

The Dali search (36) revealed that AtSerDC and CsAlaDC exhibit structural similarity

to several other decarboxylases, including 7CIG (MetDC [*Streptomyces sp.* 590]),

7ERV (HisDC1 [*Photobacterium phosphoreum*]), 6KHO (TrpDC [*Oryza sativa*

Japonica Group]), 4E1O (HisDC2 [*Homo sapiens*]), 6JY1 (AspDC

[*Methanocaldococcus jannaschii*]), and 5GP4 (GluDC [*Levilactobacillus brevis*]). The

monomers of these amino acid decarboxylases all consist of three characteristic

domains (Figure 3 and Supplementary figure 6). The C-terminal domain, also known

as the small domain, is composed of three antiparallel β-sheets and three α-helices,

facing toward the opposite monomers. The large domain contains seven stranded β-

sheets surrounded by eight α-helices, exhibiting characteristics typical of Type I

enzymes. This domain includes a conserved PLP-binding active center and amino acid

residues that are conserved across species. The N-terminal domain consists of elongated

α-helices and is connected to the large domain via a loop. The helix of one subunit is

antiparallel to the corresponding helix in the neighboring subunit, forming a clamp.

Therefore, the N-terminal domain may not be an independent folding unit and is likely

only stable in dimers.

The monomeric structure of CsAlaDC and AtSerDC exhibits numerous structural

features common to other Group II PLP-dependent amino acid decarboxylases, and the

amino acid residues that bind to PLP cofactors are conserved across multiple enzymes

(Supplementary figure 5 and 6). Out of the eight proteins analyzed, a total of thirteen

amino acid residues were found to be conserved across all sequences (Glu^{143}, Glu^{171},

Lys^{201}, Gly^{245}, Thr^{247}, Asp^{253}, His^{275}, Asp^{277}, Ala^{279}, Pro^{286}, Ser^{302}, Lys^{309}, and Tyr^{336} in

CsAlaDC). Some of them are situated within the active center and play a role in

stabilizing PLP or facilitating catalytic reactions. Conversely, other residues reside

outside the active center and their function remains unclear. (Supplementary figure 5).

PLP is bound to conserved lysine residues in the active center through the Schiff base.

The carboxylic group of conserved Asp residues stabilizes protonated pyridine nitrogen

of PLP by electrostatic action, providing the latter with strong electrophilicity to

stabilize carbanion intermediates during enzymatic catalysis. Another conserved His

residue is situated laterally to the PLP, while on the opposite side, there is a conserved

Ala residue. The oxygen atom at position 3 of PLP is stabilized by hydrogen bonding

with the Thr residue. The phosphate moiety of PLP is additionally secured to the protein

through an elongated system of hydrogen bonding. The α5 helix located within the large

domain exhibits evolutionary conservation in Group II enzymes, facilitating interaction

with the PLP phosphate moiety and conferring appropriate stabilization thereof.

The superimposition of the crystal structures of CsAlaDC and AtSerDC with those

of related enzymes suggests that residue Phe106 in CsAlaDC (corresponding to Tyr in

AtSerDC and Glu in MetDC) may play a role in determining substrate specificity. This

hypothesis was tested via mutagenesis experiments, which revealed that substitution of

Phe^{106} for Tyr rendered CsAlaDC inactive, while substitution of Tyr^{111} for Phe in

AtSerDC enabled alanine decarboxylase activity. Upon contemplation of these

experimental outcomes and informed by the elucidated crystal structures of both

enzymes, we speculate the hydrophilic nature of the Lys residue in AtSerDC

predisposes the active site to preferentially accommodate the hydrophilic amino acid

Ser as its substrate. In contrast, the analogous position in CsAlaDC is occupied by Phe,

an amino acid devoid of the hydroxyl moiety present in Lys, which augments the

hydrophobic character of the substrate-binding site. This structural nuance portends a

predisposition for CsAlaDC to select the comparatively hydrophobic amino acid

alanine as its suitable substrate. Likewise, the substitution of Glu^{64} for Ala in MetDC

reduced enzyme activity by approximately 50-fold compared to the wild-type (37),

bolstering the conclusion that the amino acid residue at this site influences the substrate

specificity of the enzyme.

Group II amino acid decarboxylase is characterized by the existence of highly

flexible catalytic rings, which are of great significance for the catalytic mechanism of

decarboxylase (33). The catalytic rings are located at the dimer interface and extend to

the active sites of other monomers in a closed conformation. In the structure of

CsAlaDC, the catalytic ring is orderly and allocated, while in the structure of AtSerDC,

it is disordered and lacks electron density (Figure 3F). Prior research has established

that the conserved amino acid residue Tyr within the ring plays a crucial role in catalysis

by donating protons to carbanions of quinone intermediates that arise following

decarboxylation (38, 39). Our experiments demonstrate that the substitution of the

corresponding Tyr with Phe in the loop renders the protein inactive. Furthermore,

alterations in the structure of CsAlaDC when combined with EA provide further

evidence supporting this conclusion. Specifically, the structure of CsAlaDC-EA

complex displayed a significant conformational change in Tyr^{336}* located in the

catalytic site (as depicted in Figure 3G). Comparing the structure of CsAlaDC apo, the

CsAlaDC-EA complex exhibited a notable 60-degree shift in the p-hydroxyl group of

Tyr^{336}* along the Cα-Cβ bond.

The monomeric configuration of CsAlaDC and AtSerDC is akin to that of other

Group II amino acid decarboxylases, except for a conspicuous dissimilarity. Unlike

other Group II amino acid decarboxylases, the C-terminal of CsAlaDC and AtSerDC

harbors an evident zinc finger structure. We propose that this structure could potentially

influence enzyme stability. To test this hypothesis, we truncated the zinc finger structure

from both proteins and expressed them. Our findings indicate that, following the

excision of the zinc finger structure, CsAlaDC became insoluble, while AtSerDC

displayed a similar trait to a certain extent.

Evolution of AlaDC and SerDC

The sequence identity between CsAlaDC and AtSerDC is 74.5%, indicating a

significant degree of homology. Moreover, the two proteins exhibit a high level of

monomer structural similarity. Notably, they differ in substrate specificity. Using the

crystal structure of CsAlaDC and AtSerDC as a basis, we identified pivotal amino acid

residues that govern the substrate specificity of serine or alanine decarboxylase. We

observed that sequences proximal to these sites exhibit a high degree of conservation

in both proteins and possess characteristic motifs (Figure 4B). Moreover, we analyzed

serine decarboxylase-like proteins in *Embryophyta* that are highly homologous to

CsAlaDC and constructed phylogenetic trees to gain insight into their evolutionary

relationships. Based on our experimental findings and evolutionary evidence, we have

identified a conserved YPX motif at the substrate binding pocket. For most plants, the

residue X in this motif is predominantly Tyr, which corresponds to the serine

decarboxylase protein. However, other residues such as Thr, Phe, Val, Ala, and Ile can

also occupy this position (Figure 4B and C). Interestingly, serine decarboxylase-like

proteins containing YPF motifs, which have the potential to function as alanine

decarboxylases, were found to be distributed throughout the phylogenetic tree,

including *Asteroideae*, *Ericales*, *Chenopodiaceae*, *Poaceae*, and *Ranunculales*

(Supplementary figure 3B). However, these proteins were absent in some more recent

species. This observation implies that the emergence of alanine decarboxylase in these

particular species could be attributed to convergent evolution.

Moreover, we have observed the enrichment of serine decarboxylase-like proteins

that possess a YPT motif within the *Fabales*. Prior research has demonstrated that

XP_004496485-1, which possesses a YPT motif, displays catalytic efficacy toward the

processes of decarboxylation and oxidative deamination of Phe, Met, Leu, and Trp (40).

Given that the YPT motif is highly conserved and widely distributed in *Fabales*, serine

decarboxylase-like proteins bearing the YPT motif may have developed a unique

substrate specificity in *Fabales*, beyond their conventional decarboxylation functions,

as exemplified by XP_004496485-1 protein mentioned above. We speculate that these

proteins may be capable of catalyzing other reactions, potentially involving non-protein

amino acids or other substances as substrates.

Applied to improve the synthesis of theanine

Theanine is an important indicator of green tea quality. Therefore, improving the

synthesis of theanine in tea plants is the focus of research. In this study, through crystal

structure analysis and mutation verification, we found that the catalytic activity of

CsAlaDC$^{L110F/P114A}$ is 2.3 times higher than that of the wild-type protein, resulting in a

more abundant synthesis of theanine *in vitro*. This gives us great inspiration to improve

the synthesis of EA in tea plants by gene editing, thus increasing the content of theanine

in tea plants (Figure 5).

Theanine is highly demanded, by the market, due to its health effects and medicinal

value, and as a constituent in food, cosmetics and in other fields. To meet the market

demand, a variety of methods have been used to acquire theanine, with the main

methods including direct extraction, chemical synthesis, biotransformation (microbial

fermentation) and plant cell culture. Based on this study, we can use mutation

technology to modify engineering bacteria and improve the possibility of synthesis of

theanine in bacteria, yeast, model plants, etc, promoting its use for industrial theanine

production.

Conclusions

In conclusion, our structural and functional analyses have significantly advanced

understanding of the substrate-specific activities of alanine and serine decarboxylases,

typified by CsAlaDC and AtSerDC. Critical amino acid residues responsible for substrate selection were identified—Tyr111 in AtSerDC and Phe106 in CsAlaDC—highlighting pivotal roles in enzyme specificity. The engineered CsAlaDC mutant (L110F/P114A) not only displayed enhanced catalytic efficiency but also substantially improved L-theanine yield in a synthetic biosynthesis setup with PsGS or GMAS. Our research expanded the repertoire of potential alanine decarboxylases through the discovery of 13 homologous enzyme candidates across embryophytic species and uncovered a special motif present in serine protease-like proteins within *Fabale*, suggesting a potential divergence in substrate specificity and catalytic functions. These insights lay the groundwork for the development of industrial biocatalytic processes, promising to elevate the production of L-theanine and supporting innovation within the tea industry.

Materials and Methods

Plants materials.

Tobacco (*Nicotiana benthamiana*) plants were grown in a controlled chamber, under a 16-h light and 8-h dark photoperiod at 25 °C. Leaves of 5-week-old tobacco plants were used for transient transformation, mediated by *Agrobacterium tumefaciens* strain GV3101.

Gene cloning and protein expression

The coding sequences of AtSerDC and CsAlaDC, amino acid residues 66-482 and 61-

478, respectively, were amplified using PCR and then ligated into the *Nde* I and *Xho* I

restriction sites of the pET-28a and pET-22b vector, respectively, providing the

recombinant vector pET-28a-*AtSerDC* and pET-22b-*CsAlaDC*. The cDNAs encoding

CsSerDC were amplified using PCR and then ligated into the *Nde* I and *Xho* I restriction

sites of the pET-28a vector. Gene-specific primers were listed in Supplementary table

3. The recombinant plasmid was transformed into *E. coli* BL21 (DE3) competent cells.

Positive transformants were grown in a 5 mL LB medium containing 30 μg/mL

kanamycin or 50 μg/mL ampicillin at 37 °C overnight and then subcultured into an 800

mL LB medium containing the corresponding antibiotic. Protein expression was

induced by the addition of 0.2 mM isopropyl-β-d-thiogalactoside (IPTG) for 20 h at

16 °C when the optical density (OD) at 600 nm reached 0.6-0.8, harvested by

centrifugation at 4 °C and 4,000 rpm for 30 min.

Protein production and crystallization

The cell pellet was suspended in 30 mL of lysis buffer (20 mM HEPES, pH 7.5, 200

mM NaCl, 0.1 mM PLP), disrupted by High-Pressure Homogenizer, and then

centrifuged at 16,000 rpm for 30 min at 4 °C to remove the cell debris. The supernatants

were purified with a Ni-Agarose resin column followed by size-exclusion

chromatography. Before crystallization, purified proteins were concentrated at 10

mg/mL. Crystallization conditions were screened by the sitting-drop vapor diffusion

method using the reservoir solutions supplied in commercially available screening kits

(Crystal Screen, Crystal Screen 2, PEGRx 1, 2, and SaltRx 1, 2). A droplet made by

mixing 1.0 µL of purified AtSerDC or CsAlaDC (10 mg/mL) with an equal volume of

a reservoir solution was equilibrated against 100 µL of the reservoir solution at 16°C.

The crystal of AtSerDC was obtained using buffer pH 8.0 containing 20% (w/v)

PEG400 as a precipitate and 0.2 M $CaCl_2$. The crystal of the CsAlaDC-EA complex

was obtained at pH 7.5 containing 2.6 M sodium acetate. The crystal of CsAlaDC was

obtained at pH 6.0 containing 3.5 M sodium formate.

Data collection and processing

Crystals were grown by sitting-drop vapor diffusion method at 16 °C. The volume of

the reservoir solution was 100 µL and the drop volume was 2 µL, containing 1 µL of

protein sample and 1 µL of reservoir solution. The reservoir solution of AtSerDC

contained 0.1 M Tris-HCl (pH 8.0), 0.2 M $CaCl_2$, and 20% PEG 400. The reservoir

solution of CsAlaDC contained 0.1 M HEPES (pH 7.5), 2.6 M sodium acetate or 0.1 M

Bis-Tris (pH 6.0), and 3.5 M sodium formate. Crystals grew in 3-5 days using a protein

concentration of 10 mg/mL. Diffraction data were collected at Shanghai Synchrotron

Radiation Facility (China). The collected data sets were indexed, integrated, and scaled

using the HKL3000 software package. The structure of AtSerDC was solved by

molecular replacement using the structure of HisDC (PDB code: 7ERV

[*Photobacterium phosphoreum*]) as the model, and utilizing AtSerDC as a molecular

displacement model for both CsAlaDC and the CsAlaDC-EA complex. The AtSerDC,

CsAlaDC, and CsAlaDC-EA complex exhibit resolutions of 2.85 Å, 2.50 Å, and 2.60

Å, respectively. The statistics for data collection and processing are summarized in

Supplementary table 2.

Enzyme activity assays

Decarboxylase activity was measured by detecting products (EA or ethanolamine) in

Waters Acquit ultraperformance liquid chromatography (UPLC) system (20, 41). The

100 μL reaction mixture, containing 20 mM substrate (Ala or Ser), 100 mM potassium

phosphate, 0.1 mM PLP, and 0.025 mM purified enzyme, was prepared and incubated

at standard conditions (45 °C and pH 8.0 for CsAlaDC, 40 °C and pH 8.0 for AtSerDC

for 10 min). Then, the reaction was stopped with 20 μL of 10% trichloroacetic acid. The

product was derivatized with 6-aminoquinolyl-N-hydroxy-succinimidyl carbamate

(AQC) and subjected to analysis by UPLC. All enzymatic assays were performed in

triplicate.

The detection methodology for theanine production remains consistent with the

aforementioned approach. The 100 μL reaction mixture, containing 20 mM Ala, 45 mM

Glu, 50 mM HEPES (pH 7.5), 0.1 mM PLP, 30 mM $MgCl_2$, 10 mM ATP, 0.03 mM

PsGS/GMAS, and 0.025 mM CsAlaDC/CsAlaDC [L110F]/CsAlaDC [L110F/P114A], was

prepared and incubated at standard conditions for 1 h. Subsequently, the reaction was

terminated via immersion of the reaction vessel in a metal bath at 96 °C for 3 min (35).

Theanine was derivatized with AQC and subjected to analysis by UPLC. All enzymatic

assays were performed in triplicate.

Site-directed mutagenesis

Site-directed mutagenesis experiment was conducted using a PCR method from the

wild-type construct pET-28a-*AtSerDC* and pET-22b-*CsAlaDC*, respectively. *Dpn* I

endonucleases were used to digest the parental DNA template. The reaction mixture

was used to transform *E. coli* DH5α competent cells and the plasmids from positive

strains were extracted to *E. coli* BL21 (DE3) for protein expression, purification, and

analyzing the enzymatic activity.

In vivo* enzyme activity assay in *N. benthamiana

The amplified PCR products were fused to the plant expression vector, pCAMBIA1305.

Linearization was conducted by restriction digest with *Spe* I and *Bam*H I. The

recombinant colonies were selected for PCR validation on the appropriate antibiotics

plate. After validation, the plasmids were electroporated into *Agrobacterium*

tumefaciens strain GV3101(pSoup-p19). Empty pCAMBIA1305 vector, with an intron

containing the GFP gene, was treated as the control.

Agrobacterium transient expression assays were performed on 5-week-old *N.*

benthamiana plants. *Agrobacterium tumefaciens* strain GV3101 (pSoup-p19), carrying

the above-described vectors, were cultured in Luria Bertani (LB) medium, containing

appropriate antibiotics, at 28 °C. When the absorbance of bacteria colonies reached

OD_{600} = 0.6-0.8, bacterial cells were collected and resuspended in MMA solution (10

mM $MgCl_2$, 10 mM 2-(N-morpholino) ethane sulfonic acid (MES), pH 5.6). After the

OD_{600} of the resuspended bacterial solution was adjusted to approx. 1.0, acetosyringone

(AS) was added with a final concentration of 200 μM, and this solution was then

incubated, at room temperature, for at least 3 h in darkness. Next, cell suspensions were

infiltrated into *N. benthamiana* leaves with a needle-free syringe. These *N.*

benthamiana leaves were then collected 3 days post infiltration, frozen in liquid

nitrogen and stored at -80 °C. Internal EA in *N. benthamiana* leaves was extracted as

previously described (22), and then the solvent was subjected to gas chromatography-

mass spectrometry GS-MS system.

Transcript level analysis in *N. benthamiana*

Total RNA was isolated from samples using the RNAprep Pure Plant Kit (Tiangen,

Beijing, China), according to the manufacturer's protocol. The cDNAs were

synthesized using TransScript One-Step gDNA Removal and cDNA Synthesis

SuperMix Kit (TransGen Biotech, Beijing, China). The qRT-PCR assays were

performed, as previously described (42). Primers used for qRT-PCR assays were listed

in Supplementary table 4 and the qRT-PCR was run on a Bio-Rad CFX96TM RT PCR

detection system and CFX Manager Software. Each reaction reagent (20 μL) contained

0.4 μL forward and reverse primers (10 μM), 2 μL cDNA (200±5 ng/μL), 10 μL SYBR

Green Supermix (Vazyme, Nanjing, China) and 6.2 μL double-distilled water. Reaction

was performed by a two-step method: 95 °C for 5 min; 40 cycles of 95 °C for 10 s; and

60 °C for 30 s. The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene was

used for internal normalization in each RT-PCR, and the $2^{-\Delta\Delta CT}$ method (43) was used

to calculate the relative gene expression. All samples were performed with three

replicates.

Multiple sequence alignment

586   MUSCLE was used to generate the protein multiple sequence alignment of amino acid

587   decarboxylases with default settings (44). ESPript 3.x was used to display the multiple

588   sequence alignment (45).

Phylogenetic analysis

The CsAlaDC was used as a query for searching its homologs in *Embryophyta* using

BLASTP (46) (version 2.13.0). To fetch homologs of CsAlaDC in distant species, we

performed a two-round blast. We first search for the best hit in the given order under

Embryophyta. Then we blast each species under a provided order using the order best

hit as a query and as many as 10 hits were kept. The sequences with e values <= 0.001

were removed. We further dropped sequences with lengths less than 200 or larger than

800 amino acids. Then multiple sequence alignment was performed via clustalo

(version 1.2.4) with default settings on these filtered sequences (47). The NJ algorithm

implemented in clustalw (version 2.1) was adopted to build the phylogenetic tree (48).

The R package ggtree (version 3.2.1) and universalmotif (version 1.12.4) were used for

tree and motif visualization (49).

Acknowledgments

We thank the team of beamline BL18U1 in the Shanghai Synchrotron Radiation Facility

for diffraction data collection.

Funding

This study is supported by grants from the Ministry of Science and Technology of China

(2019YFA0904100 to W.G, 2022YFF1003103 to X.W) and the Natural Science

Foundation of China (T2221005 to W.G, 32072624 to Z.Z).

Data availability

The structures of AtSerDC, CsAlaDC-EA complex and CsAlaDC have been deposited

in the Protein Data Bank (PDB) under accession numbers 8JG7, 8JIK and 8JIJ

respectively.

References

1.   Drew, L. (2019) The growth of tea. *Nature* **566**, S2-S4.

2.   Wheeler, D. S., Wheeler, W. J. (2004) The medicinal chemistry of tea. *Drug Dev. Res.* **61**,
45-65.

3.   Xia, E. H. *et al*. (2017) The tea tree genome provides insights into tea flavor and
independent evolution of caffeine biosynthesis. *Mol. Plant* **10**, 866-877.

4.   Yang, C. S., Hong, J. G. (2013) Prevention of Chronic Diseases by Tea: Possible
Mechanisms and Human Relevanc. *Annu. Rev. Nut.* **33**, 161-181.

5.   Xia, E. *et al*. (2020) The reference genome of tea plant and resequencing of 81 diverse
accessions provide insights into its genome evolution and adaptation. *Mol. Plant* **13**, 1013-
1026.

6.   Juneja, L. R., Chu, D. C., Okubo, T., Nagato, Y., Yokogoshi, H.   (1999) L-theanine - a
unique amino acid of green tea and its relaxation effect in humans. *Trends Food Sci.*

Technol. **10**, 425-425.

7.   Lin, J. Z. *et al*. (2022) Effect of the presence of stem on quality of oolong tea. *Foods* **11**, 3439.

8.   Haskell, C. F., Kennedy, D. O., Milne, A. L., Wesnes, K. A., Scholey, A. B. (2008) The effects of L-theanine, caffeine and their combination on cognition and mood. *Biol. Psychol.* **77**, 113-122.

9.   Kakuda, T. *et al*. (2000) Protective effect of gamma-glutamylethylamide (theanine) on ischemic delayed neuronal death in gerbils. *Neuro. Sci. Lett.* **289**, 189-192.

10.   Liu, Q., Duan, H., Luan, J., Yagasaki, K., Zhang, G. (2009) Effects of theanine on growth of human lung cancer and leukemia cells as well as migration and invasion of human lung cancer cells. *Cytotechnology* **59**, 211-217.

11.   Lu, K. *et al*. (2004) The acute effects of L-theanine in comparison with alprazolam on anticipatory anxiety in humans. *Hum. Psychopharmacol.* **19**, 457-465.

12.   Takagi, Y. *et al*. (2010) Combined administration of (L)-cystine and (L)-theanine enhances immune functions and protects against influenza virus infection in aged mice. *J. Vet. Med. Sci.* **72**, 157-165.

13.   Zheng, G., Sayama, K., Okubo, T., Juneja, L. R., Oguni, I. (2004) Anti-obesity effects of three major components of green tea, catechins, caffeine and theanine, in mice. *In Vivo* **18**, 55-62.

14.   Lin, S. *et al*. (2023) Theanine metabolism and transport in tea plants (*Camellia sinensis* L.): advances and perspectives. *Crit. Rev. Biotechnol*. **43**, 327-341.

15.   Deng, W., Ogita, S., Ashihara, H. (2008) Biosynthesis of theanine (γ-ethylamino-l-glutamic acid) in seedlings of *Camellia sinensis*. *Phytochem. Lett.* **1**, 115-119.

16.   Deng, W.-W., Ogita, S., Ashihara, H. (2009) Ethylamine content and theanine biosynthesis in different organs of camellia sinensis seedlings. *Z. Naturforsch. C* **64**, 906-906.

17.   Cheng, S. *et al*. (2019) Differential accumulation of specialized metabolite L-theanine in green and albino-induced yellow tea (*Camellia sinensis*) leaves. *Food Chem.* 276, 93-100.

18.   Takeo, T. (1974) L-alanine as a precursor of ethylamine in *Camellia sinensis*. *Phytochem.* **13**, 1401-1406.

19. Zhu, B. *et al*. (2021) CsAlaDC and CsTSI work coordinately to determine theanine

biosynthesis in tea plants (*Camellia sinensis* L.) and confer high levels of theanine

accumulation in a non-tea plant. *Plant Biotechnol. J.* **19**, 2395-2397.

20. Bai, P. *et al*. (2021) Biochemical characterization of specific Alanine Decarboxylase

(ADC) and its ancestral enzyme Serine Decarboxylase (SDC) in tea plants (*Camellia*

sinensis). *BMC Biotechnol.* **21**,17.

21. Han, S. W., Shin, J. S. (2022) Aromatic L-amino acid decarboxylases: mechanistic

features and microbial applications. *Appl. Microbiol. Biotechnol.* **106**, 4445-4458.

22. Berger, M., Gray, J. A., Roth, B. L. (2009) The expanded biology of serotonin. *Annu. Rev.*

Med. **60**, 355-366.

23. Khan, M. Z., Nawaz, W. (2016) The emerging roles of human trace amines and human

trace amine-associated receptors (hTAARs) in central nervous system. *Biomed.*

Pharmacother. **83**, 439-449.

24. Ng, J., Papandreou, A., Heales, S. J., Kurian, M. A.   (2015) Monoamine neurotransmitter

disorders--clinical advances and future perspectives. *Nat. Rev. Neurol.* **11**, 567-584.

25. Rontein, D. *et al*. (2001) Plants synthesize ethanolamine by direct decarboxylation of

serine using a pyridoxal phosphate enzyme. *J. Biol. Chem.* **276**, 35523-35529.

26. Gibellini, F., Smith, T. K. (2010) The Kennedy pathway-De novo synthesis of

phosphatidylethanolamine and phosphatidylcholine. *IUBMB Life* **62**, 414-428.

27. Mudd, S. H., Datko, A. H. (1989) Synthesis of ethanolamine and its regulation in *Lemna*

paucicostata. *Plant Physiol.* **91**, 587-597.

28. Zinser, E. *et al*. (1991) Phospholipid synthesis and lipid composition of subcellular

membranes in the unicellular eukaryote *Saccharomyces cerevisiae*. *J. Bacteriol.* **173**,

2026-2034.

29. Yunus, I. S., Liu, Y.-c., Nakamura, Y. (2016) The importance of SERINE

DECARBOXYLASE1 (SDC1) and ethanolamine biosynthesis during embryogenesis of

Arabidopsis thaliana. *Plant J.* **88**, 559-569.

30. Komori, H. et al. (2012) Structural Study Reveals That Ser-354 Determines Substrate

Specificity on Human Histidine Decarboxylase* *J Biol Chem.* **287**, 29175-83.

31. Huang, J. et al. (2018) Lactobacillus brevis CGMCC 1306 glutamate decarboxylase:
Crystal structure and functional analysis. *Biochem Biophys Res Co*. **503**, 1703-1709

32. Paiardini, A., Giardina, G., Rossignoli, G., Voltattorni, C. B., Bertoldi, M. (2017) New
insights emerging from recent investigations on human group II pyridoxal 5'-phosphate
decarboxylases. *Curr. Med. Chem.* **24**, 226-244.

33. Torrens-Spence, M. P. *et al*. (2020) Structural basis for divergent and convergent evolution
of catalytic machineries in plant aromatic amino acid decarboxylase proteins. *Proc. Natl.*
Acad. Sci. U. S. A. **117**, 10806-10817.

34. Hagihara, R., Ohno, S., Hayashi, M., Tabata, K., Endo, H. (2021) Production of L-theanine
by *Escherichia coli* in the absence of supplemental ethylamine. *Appl. Environ. Microbiol.*
87, e00031-21.

35. Yamamoto, S., Wakayama, M. & Tachiki, T. (2007) Characterization of theanine-forming
enzyme from *Methylovorus mays* no. 9 in respect to utilization of theanine production.
Biosci. Biotechnol. Biochem. **71**, 545-552.

36. Holm, L. (2022) Dali server: structural unification of protein families. *Nucleic. Acids Res.*
50, W210-W215.

37. Okawa, A. *et al*. (2021) Structural basis for substrate specificity of L-methionine
decarboxylase. *Protein Sci.* **30**, 663-677.

38. Fenalti, G. *et al*. (2007) GABA production by glutamic acid decarboxylase is regulated by
a dynamic catalytic loop. *Nat. Struct. Mol. Biol.* **14**, 280-286.

39. Gayathri, S. C., Manoj, N. (2019) Structural insights into the mechanism of internal
aldimine formation and catalytic loop dynamics in an archaeal Group II decarboxylase. *J.*
Struct. Biol. **208**, 137-151.

40. Torrens-Spence, M. P., von Guggenberg, R., Lazear, M., Ding, H., Li, J. (2014) Diverse
functional evolution of serine decarboxylases: identification of two novel acetaldehyde
synthases that uses hydrophobic amino acids as substrates. *BMC Plant Biol.* **14**, 247.

41. Qu, L., Gu, S., Zhang, J., Zhao, C., Deng, X. (2021) Determination of 18 amino acids in
three different kinds of milk powder by ultra performance liquid chromatography coupled
with pre-column derivatization. *Chin. J. Chromatogr.* **39**, 472-477.

42. Madeira, F. *et al*. (2022) Search and sequence analysis tools services from EMBL-EBI in

2022. *Nucleic Acids Res.* **50**, W276-W279.

43. Dong C. *et al*. (2020) Theanine transporters identified in tea plants (*Camellia*

sinensis L.). *Plant J.* **101**, 57-70.

44. Schmittgen T. D. *et al*. (2000) Quantitative reverse transcription-polymerase chain

reaction to study mRNA decay: comparison of endpoint and real-time methods. *Anal.*

Biochem. **285**, 194–204.

45. Robert X., Gouet P. (2014) Deciphering key features in protein structures with the new

ENDscript server. *Nucleic Acids Res.* **42**, W320-W324.

46. Camacho, C. *et al*. (2009) BLAST+: architecture and applications. *BMC Bioinf.* **10**, 421.

47. Sievers, F., Higgins, D. G. (2018) Clustal Omega for making accurate alignments of many

protein sequences. *Protein Sci.* **27**, 135-145.

48. Larkin, M. A. *et al*. (2007) Clustal W and clustal X version 2.0. *Bioinformatics.* **23**, 2947-

2948.

49. Yu, G., Lam, T. T.-Y., Zhu, H., Guan, Y. (2018) Two methods for mapping and visualizing

associated data on phylogeny using ggtree. *Mol. Biol. Evol.* **35**, 3041-3043.

Figures and table

Figure 1. Metabolic pathways and phylogeny analysis of SerDC and AlaDC in plants. (**A**) The decarboxylation of serine and alanine in plants. SerDC, serine decarboxylase; AlaDC, alanine decarboxylase. (**B**) Phylogenetic analysis of the SerDC homologous proteins in Algal, Pteridohyta, Bryophyta, Dicots and Monocots. The

phylogenetic tree was performed using the neighbor-joining method in MAGA 7.0

software and bootstrap values based on 1000 replicates were displayed at branch points.

These proteins were represented by their GenBank accession number. (**C**) Multiple

alignment of the amino acid sequences of AlaDC and SerDCs. Primary (100%),

secondary (80%), and tertiary (60%) conserved percent of similar amino acid residues

were shaded in deep blue, light blue and cheer red, respectively. The multiple alignment

was constructed by MAGA 7.0 software by ClustalW method, and GeneDOC software

was used to visualize multiple sequence alignments. "*" indicated amino acid residues

mutated only in CsAlaDC.

Figure 2. Purification and characterization of CsAlaDC, AtSerDC, and CsSerDC.

(**A**) Identification of CsAlaDC, AtSerDC, and CsSerDC using SDS-PAGE. (**B**)

Detection of enzyme activities of CsAlaDC, AtSerDC, and CsSerDC by UPLC. (**C**)

Reaction rates of substrates with different concentrations catalyzed by CsAlaDC,

AtSerDC, and CsSerDC.

Figure 3. Crystal structures of CsAlaDC and AtSerDC. (**A**) Dimer structure of

CsAlaDC. The color display of the N-terminal domain, large domain, and C-terminal

domains of chain A is shown in light pink, khaki and sky blue, respectively. Chain B is

shown in spring green. The PLP molecule is shown as a sphere model. The zinc finger

structure at the C-terminus of CsAlaDC is indicated by the red box. The gray spheres

represent zinc ions, while the red dotted line depicts the coordination bonds formed by

zinc ions with cysteine and histidine. (**B**) The 2Fo-Fc electron density maps of K309-

PLP-EA (contoured at 1σ level). The PLP is shown in violet, the K309 is shown in

spring green, and the EA is shown in lightblue. (**C**) Active center of the CsAlaDC-EA

complex, with hydrogen bonds denoted by black dotted lines. "*" denotes the amino

acids on adjacent subunits. (**D**) Dimer structure of AtSerDC. The color display of the

N-terminal domain, large domain, and C-terminal domains of chain A is shown in light

pink, khaki and sky blue, respectively. Chain B is shown in cyan. The PLP molecule is

shown as a sphere model. The zinc finger structure at the C-terminus of AtSerDC is

indicated by the red box. The gray spheres represent zinc ions, while the red dotted line

depicts the coordination bonds formed by zinc ions with cysteine and histidine. (**E**)

Active center of the AtSerDC, with hydrogen bonds denoted by black dotted lines. "*"

denotes the amino acids on adjacent subunits. (**F**) The monomers of CsAlaDC and

AtSerDC are superimposed. CsAlaDC is depicted in spring green, while AtSerDC is

shown in plum. The conserved amino acid catalytic ring is indicated by the red box. (**G**)

Amino acid residues of the active center in CsAlaDC apo and CsAlaDC-EA complex

are superimposed. CsAlaDC apo is shown in floral white, while CsAlaDC-EA complex

is shown in spring green. (**H**) The relative activity of wild-type CsAlaDC and its Y336F

mutant (left), as well as wild-type AtSerDC and its Y341F mutant (right) is shown.

Figure 4. Key amino acid residues for substrate recognition. (**A**) Superposition of substrate binding pocket amino acid residues in CsAlaDC and AtSerDC. The amino acid residues of CsAlaDC are shown in spring green, the amino acid residues of AtSerDC are shown in plum, with the substrate specificity-related amino acid residue highlighted in a red ellipse. (**B**) Active-site-lining residues from *Embryophyta* AlaDC were identified. The height of the residue label displays the relative amino acid frequency. (**C**) Histogram showing the distribution of the number of key motifs. (**D**) Histogram showing the number of key motifs in different plant orders. (**E**) Relative enzyme activities of wild-type CsAlaDC and mutant protein CsAlaDC[F106Y] against Ala substrate (columns 1 and 2), and enzyme activities of wild-type AtSerDC and various

AtSerDC mutant proteins against Ala substrate (columns 3-9) are presented. The

percentage graph shows the relative activity of each protein compared to wild-type

CsAlaDC activity (taken as a 100% benchmark). (**F**) Relative enzyme activities of wild-

type AtSerDC and AtSerDC mutant proteins (columns 1-7) against Ser substrates, and

enzyme activities of wild-type CsAlaDC and mutant protein CsAlaDC^{F106Y} against Ser

substrates(columns 8, 9) were measured. The percentage graph shows the relative

activity of each protein compared to the wild-type AtSerDC (taken as a 100%

benchmark). Three independent experiments were conducted. (**G**) The EA contents of

AtSerDC and its mutant AtSerDC^{Y111F} in *N. benthamiana*. (**H**) The EA contents of

CsAlaDC and its mutant CsAlaDC^{F106Y} in *N. benthamiana*. The significance of the

difference ($P<0.05$) was labeled with different letters according to Duncan's multiple

range test.

Figure 5. Mutations enhance CsAlaDC enzyme activity and theanine synthesis *in vitro*. (A) Relative enzyme activities of CsAlaDC mutant proteins against Ala substrate.

(B) Relative enzyme activities of CsAlaDC^{L110F}, CsAlaDC^{P114A}, and CsAlaDC$^{L110F/P114A}$ against Ala substrate. **(C)** Histogram showing the relative content of theanine resulting from different combinations of alanine decarboxylase and theanine synthetase. Three independent experiments were conducted.

818 **Table 1.** Kinetic parameters of CsAlaDC, AtSerDC and CsSerDC

Enzyme	Substrate	Km (mM)	$Vmax$ (μmol/L/s)
CsAlaDC	Alanine	1.215 ± 0.237	1.709 ± 0.072
AtSerDC	Serine	1.522 ± 0.297	7.053 ± 0.297
CsSerDC	Serine	2.364 ± 0.431	9.031 ± 0.520

Supplementary figure 1. Crystal structures of CsAlaDC and AtSerDC. (**A**) Monomer structure of CsAlaDC. (**B**) Monomer structure of AtSerDC. The color display of the N-terminal domain, large domain, and C-terminal domains is shown in light pink, khaki and sky blue, respectively. The PLP molecule is shown as a sphere model. The gray spheres represent zinc ions.

Supplementary figure 2. Catalytic mechanisms and conformational changes of CsAlaDC. After the transaldimation of the internal aldimine within CsAlaDC, resulting in the release of the active-site residue Lys309, the PLP amino acid external aldimine undergoes decarboxylation, leading to the removal of the α-carboxyl group as CO2. This process generates a quinonoid intermediate that is stabilized by the delocalization of paired electrons (1,2). The carbanion at Cα is subsequently protonated by the acidic p-hydroxyl group of Tyr336* located on the large loop, facilitated by its neighboring residue His196, which is situated on the small loop (3,4). Simultaneously, the internal aldimine LLP309 in CsAlaDC is restored, resulting in the release of the product (5).

Supplementary figure 3. Evolutionary analysis of CsAlaDC in *Embryophyta*. (**A**)

The presented diagram depicts an evolutionary tree of CsAlaDC, which is devoid of a

root and solely portrays the topological structure of the tree without including distance

information. The color of the inner ring corresponds to various orders, while the outer

ring's leaf nodes are colored based on the motif types that the sequence exhibits. (**B**)

Diversity of serine decarboxylase-like proteins in *Embryophyta* (196 species). Colored

scatter spots on the right side of leaf nodes correspond to the respective motifs shown

in Figure A.

Supplementary figure 4. The relative mRNA levels of AtSerDC and its mutant Y111F

(*A*), CsAlaDC and its mutant F106Y (*B*) in *N. benthamiana* leaves were measured by

two primers. WT, wild type of *N. benthamiana*; EV, empty vector control; *NbGAPDH*

was used as an internal control. Data represent mean ± SD (n=3). The significance of

the difference (*P*<0.05) was labeled with different letters according to Duncan's

multiple range test.

Supplementary figure 5. Multiple sequence alignment of CsAlaDC, AtSerDC, MetDC,

HisDC1, TrpDC, HisDC2, TyrDC, and GluDC were generated using MUSCLE and

visualized with ESPript 3.x. Conserved amino acid residues in all eight proteins are

highlighted with red backgrounds. The magenta box marks key amino acid residues

involved in substrate recognition for CsAlaDC and AtSerDC. The Lys residue

covalently bound to the PLP cofactor is denoted by a red star, while the green triangle

indicates the Tyr residue associated with enzymatic activity. Amino acid residues

involved in the CsAlaDC substrate binding pocket are marked with blue circles.

Supplementary figure 6. Structures of HisDC2, MetDC, TrpDC, AspDC, HisDC1

and GluDC. (**A**) The Overall Structures of HisDC2, MetDC, TryDC, AspDC, HisDC1

and GluDC. Chain A is shown in khaki, chain B is shown in cyan. (**B**) Amino acid

residues in the substrate binding pocket of HisDC2, MetDC, TryDC, AspDC, HisDC1

and GluDC. The amino acid residues in chain A are shown in khaki, and the amino acid

residues in chain B are shown in cyan.

Supplementary tables

Supplementary table 1. Amino acid sequence similarity matrix of the SerDC homologous protein

Accession number	CsAlaDC	CsSerDC	PSS19488.1	PSS17472.1	XP_010108654.1	XP_027091704.1	XP_027148923.1	XP_007019463.1	XP_034676506.1	AtSerDC	XP_002266398.1	RVW94436.1	RVW37111.1	PWZ22363.1	PWZ25657.1	XP_008678685.1	NP_001403584.1	XP_015689218.1	XP_051217297.1	XP_047064312.1	KAF7016478.1	KAF7009155.1	GBG58923.1	XP_001689659.2	OAE25238.1	XP_024373856.1	XP_024371327.1	XP_002976233.2	XP_002987141.2
CsAlaDC	100	84.92	87.16	79.92	85.69	81.1	80.63	85.34	86.45	84.92	86.65	80.4	66.8	80.12	75.9	80.81	80.73	77.65	82.29	82.05	74.12	73.11	57.84	68.46	74.76	77.41	68.72	80.71	80.54
CsSerDC	84.92	100	86.04	78.03	87.02	86.14	85.07	88.1	90.12	86.21	89.92	83.17	69.69	83.56	77.43	82.34	84.99	81.2	83.2	82.79	76.56	76.51	58	66.54	76.53	78.63	71.05	80.97	81.26
PSS19488.1	87.16	86.04	100	87.5	89.12	88	87.48	88.26	90.53	86.86	90.53	83.77	69.29	85.86	79.69	86.13	86.13	84.76	86.67	86.25	78.84	79.51	61.27	69.06	72.01	79.22	67.83	82.3	82.3
PSS17472.1	79.92	78.03	87.5	100	80.45	80.45	77.73	80.16	81.99	79.38	81.99	76.99	75.54	74.9	72.61	76.12	76.12	72.44	77.48	71.67	71.67	72.35	53.73	67.01	67.22	73.35	64.84	75.82	75.98
XP_010108654.1	85.69	87.02	89.12	80.45	100	88.2	87.48	89.78	91.63	87.73	91.43	85.97	70.84	85.74	79.47	84.57	84.87	84.85	85.66	85.66	73.45	75.73	60.53	67.05	75.8	73.35	67.98	81.25	81.25
XP_027091704.1	81.1	86.14	88	80.45	88.2	100	99.4	88.29	88.4	86	88.65	82.14	74.5	85.29	78.33	82.32	82.97	83.69	82.24	81.15	73.45	81.4	64.38	67.57	76.69	79.81	71.71	81.51	81.71
XP_027148923.1	80.63	85.07	87.48	77.73	87.48	99.4	100	87.57	87.87	85.49	88.12	81.66	74.06	84.98	78.25	83.2	82.48	83.6	81.55	81.75	73.06	81.21	63.46	67.18	75.98	79.36	70.69	78.82	79.02
XP_007019463.1	85.34	88.1	88.26	80.16	89.78	88.29	87.57	100	91.65	87.9	91.45	84.72	70.96	85.1	77.49	85.14	85.14	85.57	85.37	85.37	77.7	81.54	58.29	68.15	79.66	78.9	71.4	82.29	82.09
XP_034676506.1	86.45	90.12	90.53	81.99	91.63	88.4	87.87	91.65	100	87.12	99.59	84.33	77.18	84.33	78.24	85.19	85.14	84.79	85.19	85.19	78.24	78.61	60.33	66.73	74.21	79.92	72.89	83.57	83.16
AtSerDC	84.92	86.21	86.86	79.38	87.73	86	85.49	87.9	87.12	100	87.65	81.26	67.95	82.24	77.07	82	82.48	84.79	83.71	83.88	78.87	81	61.02	66.34	72.63	77.42	69.6	80.98	81.02
XP_002266398.1	86.65	89.92	90.53	81.99	91.43	88.65	88.12	91.45	99.59	87.65	100	92.93	76.98	83.43	77.34	82.27	84.44	84.79	85.04	85.04	77.7	78.61	60.16	66.73	77.06	78.1	72.89	82.96	82.96
RVW94436.1	80.4	83.17	83.77	76.99	85.97	82.14	81.66	84.72	84.33	81.26	92.93	100	70.79	70.79	71.48	75.92	79.17	79.48	79.52	73.08	73.94	62.12	56.41	62.12	72.37	72.12	68.14	78.02	77.62
RVW37111.1	66.8	69.69	69.29	75.54	70.84	74.5	74.06	70.96	77.18	67.95	76.98	70.79	100	64.64	67.98	65.89	64.73	64.85	68.27	68.27	60.07	60.36	46.62	53.3	59.67	65.35	59.58	65.03	64.64
PWZ22363.1	80.12	83.56	85.86	74.9	85.74	85.29	84.98	85.1	84.33	82.24	83.43	70.79	64.64	100	89.35	95.14	90.71	90.02	90.02	90.71	78.73	79.05	64.32	70.3	71.53	79.35	68.5	80.72	80.64
PWZ25657.1	75.9	77.43	79.69	72.61	79.47	78.33	78.25	77.49	78.24	77.07	77.34	71.48	67.98	89.35	100	93.86	83.78	82.96	82.38	82.38	77.9	77.9	64.32	64.09	75.85	72.92	64.33	76.23	76.04
XP_008678685.1	80.81	82.34	86.13	76.12	84.57	82.32	83.2	85.14	85.19	82	82.27	75.92	65.89	95.14	93.86	100	89.23	88.25	87.76	87.76	78.35	79.86	60.36	68.1	75.85	77.39	67.87	81.12	80.92
NP_001403584.1	80.73	84.99	86.13	76.12	84.87	82.97	82.48	85.14	85.14	82.48	84.44	79.17	64.73	90.71	83.78	89.23	100	92.54	92.54	88.8	79.52	79.66	60.36	69.11	75.75	77.46	72.4	82.24	82.04
XP_015689218.1	77.65	81.2	84.76	72.44	84.85	83.69	83.6	85.57	84.79	84.79	84.79	79.48	64.85	90.02	82.96	88.25	92.54	100	87.17	88.8	78.35	78.35	61.49	69.57	73.46	72.08	68.5	79.28	79.28
XP_051217297.1	82.29	83.2	86.67	77.48	85.66	82.24	81.55	85.37	85.19	83.71	85.04	79.52	68.27	90.02	82.38	87.76	88.8	87.17	100	98.93	87.12	83.37	61.56	68.39	73.67	76.86	67.88	82.24	82.04
XP_047064312.1	82.05	82.79	86.25	71.67	85.66	81.15	81.75	85.37	85.19	83.88	85.04	73.08	68.27	90.71	82.38	87.76	88.8	88.8	98.93	100	83.02	82.79	60.93	63.81	73.67	77.46	68.06	82.55	82.34
KAF7016478.1	74.12	76.56	78.84	71.67	73.45	73.06	73.06	77.7	78.46	78.87	77.7	73.08	60.07	78.73	78.74	78.35	79.52	78.39	83.4	83.02	100	90.69	63.98	63.47	69.5	70.6	70.9	75.7	75.51
KAF7009155.1	73.11	76.51	79.51	72.35	75.73	81.4	81.21	81.54	78.61	81	78.61	73.94	60.36	79.05	77.9	79.86	79.66	78.35	83.37	82.79	90.69	100	63.5	64.29	69.91	70.9	68.02	75.46	75.28
GBG58923.1	57.84	58	61.27	53.73	60.53	64.38	64.71	58.29	60.33	61.02	60.16	56.41	46.62	64.32	62.61	60.36	62.61	61.49	61.56	60.93	63.98	63.5	100	53.95	62.66	62.63	64.11	61.98	62.63
XP_001689659.2	68.46	66.54	69.06	67.01	67.05	67.57	67.18	68.15	66.73	66.34	66.73	62.12	53.3	70.3	64.09	68.1	69.11	69.57	68.39	63.81	63.47	64.29	53.95	100	67.92	65.79	61.89	70.56	70.16
OAE25238.1	74.76	76.53	72.01	67.22	75.8	76.69	75.98	79.66	74.21	72.63	77.06	72.37	59.67	71.53	75.85	75.75	75.75	73.46	73.67	72.52	69.5	69.91	62.46	67.92	100	83.21	75.17	79.92	79.73
XP_024373856.1	77.41	78.63	79.22	73.35	78.89	79.81	79.36	78.9	79.92	77.42	78.1	72.12	65.35	79.35	72.92	77.39	77.46	72.08	76.86	77.46	70.6	70.9	66.1	65.79	83.21	100	81.15	78.46	78.27
XP_024371327.1	68.72	71.05	67.83	64.84	67.98	71.71	72.49	71.4	72.89	69.6	72.89	68.14	59.58	68.5	64.33	67.87	72.4	68.5	67.88	68.06	69.4	68.02	64.11	61.89	75.17	81.15	100	72.53	72.53
XP_002976233.2	80.71	80.97	82.3	75.82	81.25	81.51	78.82	82.29	83.57	80.98	83.37	76.8	65.03	80.72	76.23	81.12	82.24	79.28	82.24	82.55	75.7	75.46	61.98	70.56	79.92	78.46	72.53	100	99.79
XP_002987141.2	80.54	81.26	82.3	75.98	81.25	81.71	79.02	82.09	83.16	81.02	82.96	76.4	64.64	80.64	76.04	80.92	82.04	79.28	82.04	82.34	75.51	75.28	62.63	70.16	78.27	78.27	72.53	99.79	100

 Supplementary table 2. Data collection and refinement statistics

	CsAlaDC	CsAlaDC-EA complex	AtSerDC
Data Collection			
Space Group	*P3121*	*P3121*	*C2*
Cell Dimensions			
a, b, c (Å)	163.5,163.5, 50.2	164.0,164.0,50.2	115.4,149.5,139.1
α, β, γ (°)	α=β=90, γ=120	α=β=90, γ=120	α=γ=90, β=109.2
Wavelength (Å)	0.979	0.979	0.979
Resolution (Å)	50-2.50	50-2.60	50-2.85
Completeness (%)	99.3 (99.9)	99.6 (100.0)	99.6 (98.1)
Redundancy	5.4 (5.1)	13.0 (12.5)	5.9 (5.0)
$<I>/<\sigma I>$	9.3 (1.9)	18.9 (4.6)	10.6 (2)
R_{merge}	0.128 (0.603)	0.156 (0.562)	0.173 (0.551)
Refinement			
R_{work}/ R_{free}	0.173/0.222	0.172/0.223	0.214/0.262
No. reflections	25175	22813	49543
B-facator ($Å^2$)			
Protein	43.3	48.3	57.6
PLP	29.5	31.5	52.8
Zn^{2+}	58.7	43.8	71.6
Ca^{2+}	-	55.5	-
Ethylamine	-	27.5	-
Glycerol	-	-	51.7
Water	38.5	40.9	-
R.m.s. deviations			
Bond lengths (Å)	0.008	0.008	0.007
Bond angles (°)	1.531	1.601	1.410
No. of atoms			
Protein	3292	3265	3250
PLP	15	15	60
Zn^{2+}	2	1	4
Ca^{2+}	-	1	-
Ethylamine	-	5	-
Glycerol	-	-	24
Water	115	60	-
Ramachandran statistics (%)			
Favoured	94.01	93.58	91.52
Allowed	5.74	5.68	7.61
Outliers	0.25	0.74	0.87

$R_{merge} = \sum hkl\sum i|Ii(hkl)-<I(hkl)>|/\sum hkl\sum i\ Ii(hkl)$, where $<I(hkl)>$ is the main value of I(hkl).

$R_{work} = \sum ||Fobs|-|Fcalc||/\sum |Fobs|$, where Fobs and Fcalc are observed and calculated structure
factors.

The free R factor was calculated using 5% of reflections ommitted from the refinement.

Highest resolution shell is shown in parenthesis

Supplementary table 3. Primers used for gene cloning.

Name	Forward primers (5'-3')	Reverse primers (5'-3')
CsAlaDC	GGAATTCCATATGACAACAAGCCTCAC CATCACGG	GATCTCGAGTCATTTATGAAGATCACAA TCACAATTCTCACTTC
AtSerDC	GGAATTCCATATGACCACCAGCCTGGC CG	GATCTCGAGTTATTTATGGGCCGGACA AATGCAATTATTG
CsSerDC	ATGGTGGGAAGTGTTGGGGT	TCACTTGTGTAGTGCACAAAGACAGT

886   **Supplementary table 4.** Primers used for real-time PCR.

Name	Forward primers (5'-3')	Reverse primers (5'-3')
CsAlaDC	CACTGTGATGGGGCTCTGTT	TGTTATCTGGACACCGCACG
AtSerDC	TCACTCGCTGTAACGGAACC	AACTGACCAAGCGCACCATA
NbGAPDH	AGCTCAAGGGAATTCTCGATG	AACCTTAACCATGTCATCTCCC