Phylogeny structures species' interactions in experimental ecological communities

Paula Lemos-Costa[1], Zachary R. Miller[1], Stefano Allesina[1]

[chicago]Department of Ecology & Evolution, University of Chicago
[uiuc]Department of Plant Biology, University of Illinois

Abstract

The advent of molecular phylogenetics provided a new perspective on the structure and function of ecological communities. In particular, the hypothesis that traits responsible for species' interactions are largely determined by shared evolutionary history has suggested the possibility of connecting the phylogeny of ecological communities to their functioning. However, statistical tests of this link have yielded mixed results. Here we propose a novel framework to test whether phylogeny influences the patterns of coexistence and abundance of species assemblages, and apply it to analyze data from large biodiversity-ecosystem functioning experiments. In our approach, phylogenetic trees are used to parametrize species' interactions, which in turn determine the abundance of species in a specified assemblage. We use a maximum likelihood-based approach to score models parameterized with a given phylogenetic tree. To test whether evolutionary history structures interactions, we fit and score ensembles of randomized trees, allowing us to determine if phylogenetic information helps to predict species' abundances. Moreover, we can determine the contribution of each branch of the tree to the likelihood, revealing particular clades in which interaction strengths are strongly tied to phylogeny. We find strong evidence of phylogenetic signal across a range of published experiments and a variety of models. The flexibility of our framework permits incorporation of ecological information beyond phylogeny, such as functional groups or traits, and provides a principled way to test hypotheses about which factors shape the structure and function of ecological communities.

Introduction

In 1837, Charles Darwin jotted *"I think"* and sketched a phylogenetic tree in his notebook, setting in motion a revolution in biology. Almost two centuries later, the notions of phylogeny and shared ancestry have a central place in biological research, from biochemistry to ecology. In community ecology, phylogeny was recognized early on as a powerful tool to organize our thinking about ecological communities, and the integration of phylogenetic information has promised to shed light on community assembly and patterns of species co-occurrence. Early studies relied on taxonomic information, focusing for example on species-to-genus ratios (Elton 1946; Simberloff 1970) as a proxy for competition; the idea that competition might leave a fingerprint in the taxonomic structure of ecological communities can be traced all the way back to Darwin: "As species of the same genus have usually, though by no means invariably, some similarity in habits and constitution, and always in structure, the struggle will generally be more severe between species of the same genus, when they come into competition with each other, than between species of distinct genera" (Darwin 2004).

Decades later, the availability of molecular-based phylogenies has allowed for more thorough, quantitative testing of this hypothesis. Under the umbrella of "community phylogenetics", many studies have sought to detect the signatures of community assembly processes in the phylogenetic structure of natural communities. For example, in an influential study, Webb (2000) investigated the role of shared evolutionary history in structuring rain forest tree communities. By comparing the phylogenetic diversity of local communities to regional species pools, Webb showed that closely-related species were more likely to co-occur locally than expected when sampling randomly from the regional pool. These results are consistent with the "environmental filtering" hypothesis (Webb et al. 2002; Cavender-Bares et al. 2009; Keck and Kahlert 2019): particular environmental conditions select for species possessing specific traits, and because species with similar traits are likely to share a common evolutionary history, local phylogenetic diversity is lower than expected by chance ("phylogenetic under-dispersion"). The opposite pattern, phylogenetic over-dispersion, is typically interpreted as a signature of competitive exclusion, as articulated by Darwin. In this scenario, niche differences between species, rather than the matching of species and the environment, are the driving force behind community assembly (Webb et al. 2002; Cavender-Bares et al. 2009; Graham et al. 2009). The key assumption underlying both hypotheses is that the traits determining species interactions – and consequently coexistence – have a strong phylogenetic signal. However, this simple picture has been challenged in numerous studies, which have shown that the interpretation of phylogenetic under/over-dispersion can depend on the spatial scale, the number of clades included in the study, and various statistical issues inherent to observing only the endpoints of evolution and community assembly (Cavender-Bares, Keen, and Miles 2006; Mayfield and Levine 2010; Mazel et al. 2018; Davies 2021). These issues make it difficult to assess the relevance of phylogeny for understanding ecological processes and predicting ecological patterns. To overcome these obstacles, a combination of manipulative experiments and principled statistical approaches are called for.

Under the assumption that species' interactions are tied to their phylogenetic relatedness, phylogeny can be further linked to ecosystem functioning. A basic consequence of this assumption is that phylogenetically diverse communities should harbor functionally diverse species, with less niche overlap and weaker competition, leading to increased ecosystem functioning (e.g., higher biomass, Venail et al. 2015; Davies et al. 2016). Thus, an important line of evidence for understanding the connection between phylogeny and community structure comes from studies of biodiversity and ecosystem functioning (BEF). Manipulative BEF experiments have shown not only that community biomass increases with species richness, but also that the phylogenetic diversity of experimental communities mediates their functioning (Maherali and Klironomos 2007; Connolly et al. 2011; Flynn et al. 2011; Cadotte 2013; Venail et al. 2015; Huang et al. 2020). Broadly, the results of BEF experiments are consistent with patterns found in agriculture and prairie grasslands (Jochum et al. 2020), making inferences from these experimental systems relevant for natural communities. However, careful statistical analyses of these general trends have led to conflicting conclusions regarding phylogeny and community functioning (Díaz and Cabido 2001; Petchey and Gaston 2006; Cadotte, Cardinale, and Oakley 2008; Cadotte 2013, 2015; Venail et al. 2015; Cardinale et al. 2015). Overlooked in earlier studies is the fact that species richness and phylogenetic diversity are positively correlated, and thus it is necessary to control for the richness in evaluating the effect of phylogeny (Venail et al. 2015; Davies 2021). Additionally, many analyses have relied on relating community functioning to pairwise or averaged phylogenetic dissimilarity between species, ignoring the complex relationship between evolutionary

tree topology and realized community structure (Mayfield and Levine 2010; Serván et al. 2020), as well as the processes that drive trait (co)evolution (Mazel et al. 2017; Swenson 2019). Despite considerable evidence that trait similarity, and therefore niche overlap, frequently increases with shared evolutionary history, simply correlating phylogenetic diversity with measured ecological properties fails to provide a clear-cut test of the connection between community function and phylogeny.

Previous attempts to incorporate phylogeny in the analysis of BEF data share two overaching features: a) they typically consider a single measurement for the function of each community (e.g., total biomass for each experimental plot, or the log-ratio of species abundances in multispecies communities to their abundances in monoculture), and b) they employ a statistical approach based on regressing these measurements against phylogenetic diversity or dissimilarity (Cadotte, Cardinale, and Oakley 2008; Flynn et al. 2011; Venail et al. 2015). Here we show that by abandoning these two features we can perform a strong and principled test for phylogenetic signal in these data, and we find that indeed phylogeny structures species' interactions and thus patterns of coexistence and abundance.

The basic goal of this work is to build a framework in which, first, phylogenetic information is used to parameterize a model of interactions among a pool of n species; second, species' interactions are used to predict coexistence and abundances for different assemblages that can be formed from the pool (Maynard, Miller, and Allesina 2020); and third, predictions are scored against experimental data from BEF experiments (Tilman et al. 2001; Van Ruijven and Berendse 2010; Cadotte 2013). To test whether phylogeny structures species' interactions, we repeat this procedure many times, using both the "true" (molecular) phylogeny for a given community, as well as a large number of randomized trees. By comparing the resulting likelihoods, we assess whether the true phylogeny allows superior prediction of the ecological data, which would indicate that interactions are structured by this phylogenetic topology.

Our method departs considerably from previous approaches to testing for phylogenetic effects in the same datasets (Connolly et al. 2011, 2013; Cadotte 2013; Venail et al. 2015). Specifically, we make use of the biomasses recorded for individual species, rather than the total biomass for each plot, thereby gaining statistical power and resolution; we use the full tree topology in our statistical model, rather than summary statistics (e.g., phylogenetic diversity Cadotte, Cardinale, and Oakley (2008); Flynn et al. (2011)), allowing us to distinguish between similar trees; and, finally, we relate the tree structure directly to species' interactions, and then to abundances via a model consistent with population dynamics, rather than using phylogenetic diversity as a covariate in a regression model.

Overall, our approach produces strong evidence for phylogenetic effects: the probability that a model parameterized with a random tree fits the empirical data better than a model based the actual phylogenetic tree is typically small, an outcome that is consistent across three different experimental settings, comprising a total of 19 distinct data sets. Additionally, we are able to identify which clades are most important for accurately predicting community function. Finally, we show that our framework can be extended in a number of ways and used to test other hypotheses about factors that structure ecological communities.

Results

As detailed in the Methods section, we build a statistical model to test the effect of phylogeny on species interactions. We use this model to analyze data from three BEF experiments (Tilman et al. 2001; Van Ruijven and Berendse 2010; Cadotte 2013) in which plants selected from a pool of n species are grown together in a large number of different combinations. For these data sets, we have access to the recorded biomass of each species i across different plots in which it was grown along with other species. We use $\tilde{x}_i^{(k,r)}$ to denote the biomass of species i when growing in the assemblage k (corresponding to the set of extant species in the plot) and replicate r (as the same assemblage may be found in different plots). Based on the topology of the phylogenetic tree, we build a matrix of interactions for each data set. We then use this matrix of interactions to compute the expected biomass of species i in assemblage k, $x_i^{(k)}$, and by relating the expected and observed values via a likelihood function, we can find the maximum likelihood estimates of the free parameters in the model.

These operations are accomplished in three steps. First, the topology of the phylogenetic tree is combined with tunable parameters to produce a matrix of competitive interactions; the parameterization of the matrix is guided by models for consumer-resource dynamics (Methods and Supplementary Information; see also Serván et al. (2020)), and we consider four nested model formulations (model 1, the simplest, has $2n-1$ free parameters, models 2 and 3 have $3n-2$, and model 4 has $4n-3$). In all of these models, each branch i of the tree is associated with a parameter $\lambda_i \geq 0$, and larger values of λ_i result in stronger competition between members of the clade subtended by branch i. Models 2 and 3 each consider an additional vector of parameter (α and γ) related to consumption and uptake rates, respectively, and model 4 incorporates both parameter vectors (see Supplementary Information for details). Second, we use the matrix of interactions to compute the expected abundance of each species in every possible assemblage, following the approach of Maynard *et al.* (Maynard, Miller, and Allesina 2020, summarized in the Methods section). Thus, for each species in each assemblage, we obtain a prediction, influenced by both the structure of the tree and the tunable parameters. Finally, we specify a statistical model describing how experimental data are distributed around these predicted means (requiring an extra tunable parameter to control the shape of this distribution). This allows us to compute an overall likelihood for the free parameters given a specific tree, data set, and model formulation. We perform numerical optimization to search for the maximum likelihood estimates for the free parameters (Methods), ultimately allowing us to associate a likelihood with each tree topology (and model formulation). Figure 1 represents these steps graphically, and the Methods section and Supplementary Information contain the mathematical and statistical details.

Results for the data set published by Cadotte (2013) are shown in Figure 2 a, with one panel per model formulation. For this data set, even the simplest model, with a total of 16 free parameters (15 used to parameterize the interactions, one for the distribution), is able to recapitulate the observed abundances with good fidelity (8 species, measured in 27 distinct communities, for a total of 87 observations, including replicated assemblages). We find a correlation between the logarithm of the observed biomasses and the logarithm of the predicted biomasses close to 88%. When we consider more complex models, the log-likelihood improves marginally (from -348.37 to -342.55, see Table 1 for details; the correlation also grows to about 89%) at the cost of a substantial increase in the number of free parameters (30 parameters for model 4). In this case, and in all cases we consider, almost any metric for model selection (e.g., AIC, BIC) would favor the simplest model (model 1).

While these results indicate that models structured by phylogeny can describe the observed data quite well, they do not tell us if the particular tree topology is responsible for the quality of fit, or if other models of similar complexity (i.e., with a similar number of free parameters) would fit just as well. To answer this question, we compare the quality of fit, measured by its likelihood, against alternative models. First, as a natural reference point, we consider a "star tree", in which all of the species descend directly from the root of the tree (Tucker et al. 2018). This configuration can be obtained from any tree topology by setting the parameters λ_i corresponding to the internal branches to zero; therefore, the likelihood for the star tree provides a lower bound for the distribution of possible trees (continuous line in Figure 2 b). Next, we fit the model many times, each time using a different random tree to structure interactions. We consider two possible ways to build random trees: generated via the Yule model (teal in Figure 2 b) or randomized by

Figure 1: Diagram illustrating the steps of our approach. We combine a set of parameters and a phylogenetic tree to parameterize a matrix of interactions between all the species in a pool. This parameterization is inspired by consumer-resource models (Supplementary Information). Using the matrix of interactions, we predict whether a given assemblage of species will coexist, and at what abundances. These predicted abundances are thus determined by the structure of the phylogenetic tree as well as the tunable parameters. Abundances observed in experimental communities are assumed to follow a statistical distribution determined by these expectations, allowing us to compute the likelihood of the parameters given the tree, data and model. We maximize the likelihood of the parameters given the phylogenetic tree, obtaining a unique likelihood score for the tree topology, and then constrast it with model fits using randomized trees. By repeating this procedure many times, we build a distribution of likelihoods over the set of random trees. Finally, we compute a p-value which measures the probability of obtaining a better likelihood when using a random tree in place of the true phylogenetic tree.

Figure 2: a) Predicted vs. observed biomasses for the data published by Cadotte (2013). Each point corresponds to the biomass measured for a given species in a given plot. The predicted value is necessarily the same across replicates for a given species/assemblage combination. The correlation between the logarithm of the predicted and observed biomasses is reported in each panel. b) Distribution of the likelihoods for the four models, using random trees as an input. The colors indicate whether the random trees were obtained from the Yule process (teal) or by shuffling the leaves of the original tree (grey). The vertical dashed line represents the likelihood using the original tree. The solid line represents the likelihood using a "star" tree, which provides a lower bound for the distribution (see text). Approximate p-values obtained for one thousand randomizations are reported in the panels.

shuffling species' identities but maintaining the true tree topology (grey in Figure 2 b). Both approaches maintain the same number of free parameters, while the second approach maintains additional structure, such as the overall asymmetry of the tree. We build one thousand distinct random trees for each type of randomization and maximize the likelihood for each tree and model. From this ensemble of randomizations, we compute a p-value, quantifying the fraction of random trees that yield a better likelihood than the true phylogeny. Intuitively, $1 - p$ is the probability that using the true phylogenetic relationships for the community will improve the model fit.

For the Cadotte data set, Figure 2 shows that the star tree sets a lower bound for the likelihood, and that only a few a random trees perform better than the true tree (dashed black line) when using model 1. The p-values are approximately 0.029 and 0.028 for the shuffled and Yule trees, respectively. As we increase the number of free parameters, the flexiblity of the models increases, reducing the effect of the tree topology on the quality of fit. Accordingly, p-values increase for the more complex models (both close to 0.13 for model 2, and higher for models 3 and 4).

When we repeat this procedure for systems with a larger number of species, as in the Biodiversity II experiment (Tilman et al. 2001), we find even stronger evidence for phylogenetic effects. Additionally, when we analyze another system with eight species but more observations, from the Wageningen biodiversity experiment (Van Ruijven and Berendse 2010), we also find substantial evidence for phylogenetic effects. The p-values for every data set and model-randomization combination are reported in Table 1. In Figure 3 we summarize these results by plotting the distribution of p-values. This distribution is greatly enriched in low values. Out of a total of 152 tests (19 data sets, four models, two randomizations), we find that 86 p-values are ≤ 0.05, while we would expect less than 8 by chance alone. Another 14 are below 0.1, and 52 are above 0.1 (we would expect more than 136 by chance). This pattern is evident across the three experimental settings and across model formulations, even though we expect the more complex models to dilute phylogenetic signal. If we reduce the number of tests by pooling the results of the two randomizations (having found that the likelihood distributions for the two randomizations are very similar), we find 42 p-values ≤ 0.05 out of 76 tests (19 data sets, 4 models; 11 p-values ≤ 0.05 for model 1 and 3, 10 for models 2 and 4). Many p-values remain below this threshold when they are adjusted according to the Bonferroni (22 adjusted p-values ≤ 0.05) or Hommel (33 adjusted p-values ≤ 0.05) corrections. We conclude that these data sets bear a strong signature of phylogenetic structure.

Our method also allows us to identify which features of the phylogenetic tree are the most important to capture the patterns in the empirical data. Take the simplest model, in which each branch of the tree (defining a clade) is associated with an increase in the strength of competition between the species in the corresponding clade. When species belonging to a certain clade do not experience increased competition with the other members of the clade, the maximum likelihood estimate of the corresponding branch length parameter will shrink to near zero. Figure 4 represents this feature visually: branches that do not strongly affect the pattern of interaction strengths are small, while those that help explain the data are associated with a positive value. As noted above, if the interactions had no structure, or structure not corresponding to the "input" tree toplogy, all internal branches would shrink to zero, giving rise to the star tree, in which only the branches associated with the root (encoding "mean-field" interspecific competition) and the tips (intraspecific competition) would have positive lengths. Thus, branches with positive values correspond to clades that help explain the structure of the empirical data.

For the data from Cadotte (2013), the likelihood maximization singles out the clade of grasses (Poaceae, clade *b* in the phylogenetic tree in Figure 4), and separates the only legume (*D. canadense*) from the rest of the species (clade *a* in the figure). Likewise, the analysis of the data by Van Ruijven and Berendse (2010) (year 2003 shown) highlights the competition between the four grasses (Poaceae, clade *a* in Figure 4), as well as the two members of Asteraceae (clade *b*). We can confirm this interpretation by stratifying the random trees according to the presence of these highlighted clades. For example, when fitting the data from Cadotte (2013), we find that those trees possessing clade *a* perform slightly better than other random trees, those that group the species that are not grasses or legumes (clade *b*) perform much better, and those retainng both *a* and *b* (as in the true tree) perform best than all. In fact, this last set of trees performs as well as or better than the true tree (likelihood marked by a dashed line), indicating that these two partitions are the salient components of the phylogenetic structure. Similar results are found for the Wageningen experiments:

Figure 3: Pooled p-values obtained by testing for phylogenetic effects using four models, two randomizations, and nineteen data sets (152 p-values in total). We count the p-values falling in each bin of size 5% to build the histogram; colors indicate which of the three experiments the p-value corresponds to. The distribution of p-values is greatly enriched in low values, as expected when the null hypothesis of no phylogenetic effects (no difference between the true and random trees) can be rejected.

trees clustering the grasses or Asteraceae perform better than those that do not, and trees clustering both clades correctly yield a much higher likelihood. For more speciose trees, we also see evidence that a few clades contribute strongly to the quality of fit, although in this case sampling enough random trees to find all possible combinations of these key clades would require an astronomical number of randomizations. For example, in the Biodiversity II experiment (year 2012 data) we see support for increased competition between the legumes (clade a), and all random trees clustering the legumes together yield high likelihood; the grasses are split into several distinct clades, with strong support for increased competition between two C4 grasses (clade b), and two C3 grasses (clade c). Again, randomized trees conserving these clades perform better on average than those that do not.

We have presented the results regarding interpretability of model parameters for the simplest model, but the same apply to the models with greater flexibility. All models presented here point out in the same direction of these data sets being phylogenetically structured, as evidenced by the overall results presented in Table 1 and the distribution of p-values. Even when adding more flexibility to the model through the incorporation of parameters related with consumption and uptake rates associated with each species the presence of phylogenetic signal is quite robust (see Supplementary Information for the fit of models 2-4).

Figure 4: Maximum likelihood trees using model 1 for three selected data sets. The length and color of the branches correspond to increased strength of competition between the species subtended by the branch. Some branch lengths are close to zero, indicating that the species in these clades do not experience elevated competition with one another. For example, in the data from Cadotte (2013) the model does not support an increase in the strength of competition between *R. hirta* and *S. altissima*, compared to their interactions with *M. fistulosa*, despite the additional evolutionary history shared by these two species. As a result, this clade collapses into a polytomy with three species. Box-plots show the distribution of log-likelihoods for randomizations that retain the clades highlighted in the phylogenies (indicated by letters; boxes without labels correspond to randomizations where no labeled clades were retained). The dashed line indicates the log-likelihood of the original tree.

Discussion

In this study, we leveraged the statistical framework of Maynard, Miller, and Allesina (2020) to fit species interaction models where interaction strengths are systematically tied to evolutionary history, summarized by phylogeny. By fitting models using both the true phylogenetic tree for a set of species, as well as a randomized ensemble, we have devised a simple and direct test to determine whether phylogeny structures species' interactions, and thus patterns of coexistence and abundance in empirical data. Analyzing nineteen data sets from three different biodiversity-ecosystem functioning experiments, we found substantial support for phylogenetic structure in species' interactions. In this section, we briefly discuss the advantages and limitations of this approach, as well as possible extensions.

Advantages of the approach

Our methodology uses biomasses recorded for all species in a plot, as opposed to the total biomass in the plot. While previous studies (Tilman et al. 2001; Connolly et al. 2011; Cadotte 2013; Huang et al. 2020) focused only on total biomass in order to test BEF hypotheses, examining these data at the level of species yields a larger number of measurements, giving greater statistical power to detect phylogenetic structure, while also reducing the risk of overfitting. Because BEF and other community data sets often contain species-level biomasses (or abundances), our approach makes fuller use of the painstaking empirical work that went into building these large data sets.

This approach is grounded in a simple framework to infer interactions and predict biomasses from community "endpoints" — a framework that is gaining traction in the ecological literature (Xiao et al. 2017; Fort 2018; Maynard, Miller, and Allesina 2020; Ansari et al. 2021; Skwara et al. 2023). One of the main advantages of this framework is that it makes predictions that are compatible with models of population dynamics: if we were to simulate communities using the fitted interaction strengths as parameters of the Generalized Lotka-Volterra model, we would find the same predicted abundances as equilibria of the dynamical system. This is in contrast with models based on a regression framework, which are inherently incompatible with population dynamics (see Supplementary Information). To include phylogeny in this framework, we took a principled approach, formalizing the intuitive expectation (already clear to Darwin) that shared ancestry translates into stronger competition. This translation between phylogenetic structure and interaction strengths was guided by consumer-resource models (Methods, Supplementary Information), permitting a clear ecological interpretation of the parameters.

This basic approach is quite modular and flexible; for example, here we used four related models—each structured by phylogeny but including different levels of additional complexity—and employed two different randomizations to test for phylogenetic effects. Other model formulations or randomizations could be used to test for phylogenetic structure within the same framework, and one could additionally alter the assumptions of the statistical model (e.g., we assumed that empirical values were Gamma distributed, but one could substitute any other distribution with positive support) or otherwise tailor the pipeline for a specific data set and ecological question.

The models we have employed use the full topology of the phylogenetic tree to structure interactions. This means that, in principle, we can use likelihoods to discriminate between trees with similar coarse structure (e.g. sum of branch lengths, a commonly used summary statistic). We believe this is a substantial step forward compared to approaches that rely on a single metric for trees induced by each assemblage, as the detailed structure of a phylogeny might mediate important differences in community patterns. Additionally, we depart from the simplistic assumption that interaction strength should increase linearly (or according to another simple function) with shared ancestry. Instead, we associate a free parameter with each branch, thereby allowing some branches to take on greater importance, and other to shrink to zero, guided by the data. Essentially, we assume that interaction strengths do increase with shared ancestry, but make no other assumption about the form of this relationship. This allows us to capture phylogenetic effects even if the underlying processes of trait evolution are highly variable in time and across the phylogeny (Mazel et al. 2017; Tucker et al. 2018).

This model flexibility also allows us to identify which branches play a key role in structuring interactions. Previous efforts to move beyond straightforward phylogenetic distances used network tools to identify important branches in phylogenies for experimental communities (Davies et al. 2016). Analyzing the Biodiversity II data, as we did here, Davies et al. (2016) used this approach to detect an important effect of legumes on community function. In agreement with their conclusions, we also find that the clade including the legumes plays a central role in structuring these experimental communities (clade *a* in Biodiversity II, Figure 4). For the Wageningen experiment (Ruijven and Berendse 2005; Van Ruijven and Berendse 2010), an examination of inferred branch lengths reveals temporal patterns. This experiment showed increasing complementary effects over time, suggesting differentiation of resource use or other traits, or increasing facilitation. These adaptive processes at the community level take time and were not detected in the first year of the Wageningen experiment, however, in the experiment's subsequent years there was a strong signal of complementary effects that increased in time and with species richness (Van Ruijven and Berendse 2010). Our approach recovers these temporal effects: in the first years of the experiment (2000, 2001 in Supplementary Information and 2003 in Figure 4) we find evidence for a strong phylogenetic signal in the clades containing the grass species and most of the dicots. From 2004 on (Supplementary Information), the fit of our model shows that the branch lengths of the dicot clade are reduced, with these species mostly collapsing into a polytomy and only the grasses retaining a strong phylogenetic signal. The decrease in the fitted branch lengths over time in the experimental data suggests reduced competition, which is consistent with the reported increase in complementary effects (Ruijven and Berendse 2005; Van Ruijven and Berendse 2010) and points to the phylogenetic origins of these effects.

Limitations of the approach

There are important limitations, both practical and theoretical, to this methodology. Practically, our approach is computationally very intensive, especially due to the large number of randomizations needed to obtain accurate p-values. This issue is exacerbated for data sets with many species because, for each parameterization, it is necessary to invert an interaction sub-matrix for each observed assemblage. The computational cost of this operation scales approximately with the cube of the size of the matrix (number of species). This problem can be alleviated by fitting randomizations in parallel with computer clusters, but it remains a key limitation (for example, the calculations presented here required approximately 5 years of computing time). The computational challenge is made worse by the fact that we expect the likelihood surface of the parameter space to contain many local maxima. Thus, finding the global maximum likelihood is challenging, and at a minimum requires initializing the numerical search from several initial conditions.

On the theoretical side, we have considered four nested models, capturing progressively more ecological detail, and moving from the simpler to the more complex formulations adds approximately n parameters at each step. Models with more parameters are more prone to over-fitting (i.e., fitting the noise as well as the signal in the data), and in fact we find that the contribution of the phylogenetic tree can be swamped by the large number of parameters in the most complex models (e.g., Figure 2). As we have seen in the Results section, some—and sometimes many—of these parameters shrink to zero through likelihood maximization, and could therefore be removed without affecting the goodness of fit. This suggests an approach to reduce the number of parameters, by analyzing degenerate trees in which some branch have zero length (i.e., non-binary phylogenetic trees), or coarse-grained trees (e.g., based on higher taxonomic levels).

Beyond phylogeny

Interestingly, from an ecological perspective, the collapse of many branch lengths suggests that only parts of the phylogeny—key clades, possibly corresponding to key functional traits—matter for species' interactions. Thus, we might hope to test for other, possibly simpler, kinds of structure. This is possible through a straightforward extension of our approach. In this framework, the topology of a tree is encoded by binary matrix V (see Methods), in which rows correspond to ancestral species, columns to extant species, and $V_{ij} = 1$ whenever the row is an ancestor of the column (and zero otherwise). The same approach could be extended to any binary matrix V (provided the resulting matrix of interactions and all its sub-matrices are

invertible). For example, we could postulate models in which there are k generalized "groups" of species (with or without nested or overlapping membership). Then the rows of V would be groups, and $V_{ij} = 1$ whenever species j belongs to group i. Groups might correspond to functional traits: in this case the rows of V are traits and $V_{ij} = 1$ whenever species j possesses trait i. These simple yet powerful extensions allow our basic pipeline to be used to test for community structure well beyond phylogenetic signal. While we find evidence for phylogenetic structure in the data sets considered here, the strong assumption that phylogeny serves as a proxy for ecological similarity might not always hold in natural communities, or might be subject to distortion. For example, trait similarity might be a consequence of convergent evolution, as opposed to shared evolutionary history (Cavender-Bares, Keen, and Miles 2006; Losos 2008). Further, traits determining species interactions—and hence influencing ecosystem functioning—may be under strong selection, and consequently evolutionary history, inferred from neutral markers, might not reflect functional trait similarity (J. P. Wright et al. 2006; Davies et al. 2016). In such cases, our approach makes it possible to explicitly test the role of traits known to be under selection or known to influence species interactions and coexistence (Fry, Power, and Manning 2014). In fact, setting aside the computational challenge of fitting a large number of trees, one could in principle attempt to find the maximum likelihood tree or group structure and effectively "reverse-engineer"" the traits or other factors that contribute most to community structure and functioning.

Importantly, regardless of the chosen structure, the same randomization scheme can be used to test whether the traits or groups of interest are informative about ecological outcomes. Using an appropriately chosen randomized ensemble, one can always compare a V matrix of interest against randomizations to compute a p-value, which quantifies the value of using the chosen structure to constrain interactions, as compared to models that are equally complex, but biologically "uninformed".

Beyond competition

So far, we have not relaxed the assupmtion that species' interactions are strictly competitive. Even when considering species at same trophic level, the view of competition as the only force structuring communities might be too limited (A. J. Wright et al. 2017; Frainer et al. 2018). One of the striking patterns observed in BEF experiments is "overyelding", in which a species achieves a higher biomass in mixed communities than it does in monoculture (Connolly et al. 2013). This pattern has been associated with complementary effects, but might also reflect facilitation (A. J. Wright et al. 2017). Current approaches make it challenging to disentangle these two mechanisms, but the flexibility of our framework might also make it possible to detect the role of facilitation in structuring these ecological communities.

There are at least two ways our framework could be extended to allow for other types of interactions besides competition. First, by letting the clade-specific interactions (λ_i) take any sign, positive or negative, we can model facilitation or mutualism together with competition. Second, we can let the auxilary parameters modulating phylogenetic effects (γ, θ; see Methods) take arbitrary values, thereby allowing asymmetric interactions, such as predation or paratism. None of these changes increases the number of free parameters, but should always result in a better fit to the data by removing constraints on the parameters.

Naturally, this flexibility comes with costs, including increased risk of overfitting and more complex interpretation of the model parameters. To avoid these issues and retain maximal interpretability, we have concentrated on less flexible models with competition only, although these extensions represent a natural area for future inquiry.

Conclusions

We have introduced a principled, robust statistical approach to detect phylogenetic effects on species interactions. Given the topology of a tree, we find a maximum likelihood model by optimizing over a set of free parameters. Using data from three different biodiversity-ecosystem functioning experiments, we have found that the models structured by phylogeny provide a good fit to observations, and provide strong support for the presence of phylogenetic effects. Our approach departs from existing regression-based frameworks, making it compatible with population dynamic models; moreover, it more fully exploits all of the information in the published data, both in terms of measurements (using biomasses recorded at the level of species instead of at the level of plots), and in terms of phylogeny (using the full tree structure instead of summary statistics). Even though it is computationally intensive, our approach can easily be extended to account for functional groups, explicit traits or other types of interactions beyond competition.

Methods

Modeling framework

As outlined in the main text, our statistical framework is composed of three steps. Each of the steps could be carried out in a number of ways.

a) From tree topology to interactions

We represent the topology of a binary phylogenetic tree with n leaves (i.e., the species in the pool) as a binary matrix V of size $(2n-1) \times n$. The rows of the matrix represent either the extant species or their ancestors (including the common ancestor to all species, i.e., the root of the tree), while the columns represent the species. A coefficient $V_{ij} = 1$ if i is an ancestor of j (or j itself), and is zero otherwise. This matrix is called the "basis matrix" of the tree in the literature (Bravo et al. 2009); take a vector of $2n-1$ branch lengths α: then $V^T D(\alpha) V$ is the variance-covariance matrix of the tree.

Inspired by consumer-resource models (Supplementary Information), we parameterize the $n \times n$ matrix of interactions between the species as:

$$A = D(\gamma) V^T D(\lambda) V D(\theta) \quad , \tag{1}$$

where γ and θ are positive vectors of length n with entries summing to one (thus yielding $n-1$ free parameters each), λ is a positive vector of length $2n-1$, $D(\zeta)$ is a diagonal matrix with the elements of ζ on the diagonal, and V^T denotes the transpose of matrix V. We therefore have $4n-3$ free parameters in total. We also consider simplified models in which either $D(\gamma)$, $D(\theta)$, or both are taken to be $\frac{1}{n}I$, i.e., a scaled version of the identity matrix, reducing the number of parameters to $3n-2$, $3n-2$ and $2n-1$, respectively. Details on these parameterizations and their derivation from consumer-resource models are reported in Supplementary Information.

Provided with a tree topology describing the evolutionary history of the n species in the pool (defining the associated matrix V), and empirical data in which different sets of species taken from the pool are grown together, we want to find the maximum likelihood estimates of the parameters stored in γ, λ, and θ. To achieve this goal, we first relate interactions to predicted abundances, and then predicted abundances to experimental data.

b) From interactions to predictions

We want to relate the species interactions, encoded in the phylogenetically-structured matrix A (Eq. 1) with the abundances species would attain in different assemblages. Take k to be the set of species in a given assemblage. We assume that the abundance of species i in assemblage k, denoted $x_i^{(k)}$, is linearly related to the abundance of the other species in the assemblage, with each effect mediated by the respective interaction strength:

$$x_i^{(k)} = \frac{1}{A_{ii}} \left(1 - \sum_{j \in k; j \neq i} A_{ij} x_j^{(k)} \right) \quad .$$

Thus, species i grown by itself would reach $1/A_{ii}$ (its carrying capacity), while when growing with j it would yield a lower value whenever $A_{ij} > 0$, i.e., when species i and j are competitors.

Writing $A^{(k)}$ for the sub-matrix of A obtained by retaining only the columns and rows of species in assemblage k, we can write this system of equations in more compact form as:

$$A^{(k)} x^{(k)} = 1^{(k)} \quad ,$$

where $1^{(k)}$ is a vector of ones with as many elements as there are species in k. Thus, provided with a matrix A we can compute the value of $x^{(k)}$:

$$x^{(k)} = \left(A^{(k)}\right)^{-1} 1^{(k)} \quad . \tag{2}$$

There are two possible outcomes: a) all the elements of $x^{(k)}$ are positive, which we interpret as the coexistence of the species in assemblage k, with expected abundances specified by the vector $x^{(k)}$; b) one or more of the elements are negative, which we interpret as the impossibility of coexistence for this set of species. This framework was proposed numerous times in the ecological literature (Xiao et al. 2017; Fort 2018; Maynard, Miller, and Allesina 2020; Ansari et al. 2021) and here we follow the approach of Maynard *et al.* (Maynard, Miller, and Allesina 2020; Skwara et al. 2023) quite closely.

Provided with a matrix A defined in Eq. 1, we can thus predict the expected coexistence and abundance of species in all possible assemblages using Eq. 2. Finally, we can relate these predictions to the empirical data.

c) From predictions to likelihoods

Each data set is represented by a set of measurements $\tilde{x}_i^{(k,r)}$, specifying the biomass of species i when growing in assemblage k (i.e., defined by the extant species in the plot) and replicate r. The data set contains a large number of assemblages, possibly with replicated experiments (i.e., different plots in which the same assemblage is measured). We take these measurement to be noisy estimates of the "true", expected biomass of species i in assemblage k:

$$\tilde{x}_i^{(k,r)} \stackrel{iid}{\sim} \text{Gamma}\left(\alpha, \frac{\alpha}{x_i^{(k)}}\right) \quad , \tag{3}$$

that is, we assume the measurements to have been sampled independently from a Gamma distribution with parameters α (scale) and $\alpha/x_i^{(k)}$ (rate), such that the expected value is $\mathbb{E}(\tilde{x}_i^{(k,r)}) = x_i^{(k)}$ and the variance is $\mathbb{V}(\tilde{x}_i^{(k,r)}) = \frac{1}{\alpha}\left(x_i^{(k)}\right)^2$. The parameter α is taken to be specific to the data set. Thus, for a given data set all measurements are taken from distributions with a constant coefficient of variation, $\frac{1}{\sqrt{\alpha}}$. Whenever $\alpha > 1$, the mode of the distribution is larger than zero, and the shape of the distribution displays a central tendency.

The choice of this particular Gamma distribution is convenient (e.g., it has support on the positive real line, a constant coefficient of variation, and adds a single free parameter); alternatively, one could have chosen a Log-Normal distribution, or any other appropriate alternative. We do not expect the choice of a particular distribution to affect the qualitative conclusions drawn here.

Numerical search for maximum likelihood estimates

In summary, provided with the parameters of the model (the vectors γ, θ and λ, as well as the parameter α controlling the shape of the distribution), the empirical data, represented by the empirical measurements of the biomass of the species in the different assemblages (possibly, in replicated experiments), and the phylogenetic tree for the species pool, we can: a) use the parameters and the tree to build the matrix of interactions (Eq. 1), b) compute the predicted abundances for all assemblages (Eq. 2), and c) compute a likelihood by multiplying the densities attained by the empirical samples for the distributions specified in Eq. 3.

The fact that we can go from parameters to likelihoods allows us to search numerically for the maximum likelihood estimates of the parameters. Briefly, a) we initialize the parameters such that A is the sum of the identity matrix and a random matrix with small norm (practically, ensuring coexistence in all possible assemblages); b) search for parameters increasing the likelihoods via a combination of simulated annealing,

followed by hill-climbing (slightly alter parameters and accept the move if likelihood is increasing), and numerical optimization via the Nelder-Mead and BFGS algorithms provided by the *optim* function in *R* (R Core Team 2021); c) step b) is repeated 500 times, each time improving the likelihood until (tentative) convergence to the maximum likelihood estimates. Parameterizations yielding lack of coexistence for assemblages that were observed to coexist experimentally are discarded. Parameterizations yielding a likelihood that is lower than what obtained for the corresponding star tree are also discarded. Finding optimal parameters is computationally very difficult (there can be many local optima in parameter space), and very expensive (due to Eq. 2 requiring to invert a matrix for each assemblage). We implemented the approach using a mix of *Stan* (Stan Development Team 2023) and *R* (R Core Team 2021) code, started the search five times from random parameters, and stored the best likelihood for each data set/model combination. The code, raw and organized data, and results are available here (`github.com/StefanoAllesina/lemos-costa-2023`).

Testing for phylogenetic signal

In the preceding sections, we showed that, given empirical data and a matrix V associated with a tree, we can find the maximum likelihood estimates of the parameters in Eq. 1, and thus the maximum likelihood associated with the tree, $\mathcal{L}_V$. To test whether phylogeny can help explaining species interactions, and thus the observed abundances in all assemblages, we repeat the procedure for a random tree V', yielding $\mathcal{L}_{V'}$. By repeating this procedure several times, we can estimate the probability that a random tree yields a higher likelihood than the "true" tree associated with V. As such we take $p \approx \frac{1}{m} \sum_i^m \mathcal{H}(\mathcal{L}_V \leq \mathcal{L}_{V_i'})$, where m is the number of random trees, $\mathcal{H}(\cdot)$ the Heaviside step function and V_i' is the matrix associated with the i^{th} random tree.

Finally, we need to define what we mean by a random tree. We consider two choices: a) a random tree with the same number of leaves constructed according to the Yule model, a speciation model in which every branch has the same probability of speciating at a time; b) a tree with the same topology as the original, but where we shuffled species' identities. Therefore, for each data set we present randomizations for four related models (all variations on Eq. 1) and two distinct ways of constructing the random trees.

When we randomize the tree, we might encounter a tree we have already used for a previous parameterization; because we want to compute the p-value with respect to the ensemble of unique random trees, we keep drawing random trees until we have a thousand distinct ones; in case of duplicate trees, we store the best likelihood we have found.

Data and data organization

The methods by Maynard, Miller, and Allesina (2020) were devised for the analysis of experiments carried out in laboratory conditions for a small species pool, for which a large proportion of all the possible combinations of species have been observed. Biodiversity-Ecosystem Functioning experiments, on the other hand, are often carried out in experimental plots, starting from a pool of a dozen or more species—because the number of possible assemblages one can form with n species is 2^n, and only about 100 plots are typically available, it follows that the number of sampled assemblages is necessarily small compared to the theoretical limit. We therefore attempt selecting data sets in which a) only species that have appeared in multiple plots are considered to be part of the pool; b) unidentified biomass or non-target species are not dominating the plots; c) we have sufficient replication or diversity of assemblages to be able to confidently fit the parameters of the model.

To this end, we consider data collected as part of the Biodiversity II experiment (Tilman et al. 2001, data from 2001 to 2018, inclusive, downloaded from the Cedar Creek website), the Wageningen diversity experiment (Van Ruijven and Berendse 2010, provided by Dr. Jasper van Ruijven), and the experiment carried out at the Toronto's Koffler Scientific Reserve by Marc W. Cadotte (data accompanying Cadotte 2013).

The data from Cadotte (2013) contains plots in which a subset of 14 species were planted. Some did not grow, and went extinct during the experiment. We consider a species that did not reach 2.5% of its maximum observed biomass to be extinct in the plot. We then considered species that appeared in at least 5 distinct assemblages; if a plot contained species that did not meet this requirement, it was discarded; the procedure was then repeated until all species appeared in at least 5 distinct assemblages, leaving 8 species, forming 27 distinct assemblages, for a total of 87 measurements (including replicated assemblages). Code to implement this procedure, along with the raw and elaborated data are accompanying this work.

The same general procedure was applied to the data from Van Ruijven and Berendse (2010), with the difference that we required each species to be present in six distinct assemblages. After the selection procedure, all eight species were retained, and we obtained nine data sets, one for each yearly sampling from 2001 to 2009, inclusive.

The data from Tilman et al. (2001) is the most difficult to process, because besides the original target pool of eighteen species, many other species are found in the plot, as well as substantial amount of litter or unidentified biomass. We therefore included in the "species pool" a much larger number of species, excluded plots in which the total biomass for species not in the pool surpassed 10% of the total biomass, and then applied the procedure illustrated for the other data sets. In this way, we were able to extract nine data sets containing between 13 and 20 species and a substantial number of assemblages.

Summary statistics on each data set, including number of species, number of unique assemblages and number of measurements is reported in Table 1.

We normalized species names using the World Flora Database (*http://www.worldfloraonline.org/*) and constructed the phylogenetic tree associated with each set of species using the *V.PhyloMaker* package in *R* (Jin and Qian 2019).

Ackowledgments

We thank C. Serván, P. Lechón-Alonso, M. Chakraverti-Wuerthwein, and S. Kundu for fruitful comments and discussion, and to J. Van Ruijven for sharing the data from the Wageningen experiment. This work was supported by the National Science Foundation (DEB #2022742 to SA).

Derivation of the model from consumer-resource dynamics

1)

2) $C = D(\nu) \times V$

$$= \begin{pmatrix} \nu_1 & \nu_1 & \nu_1 & \nu_1 & \nu_1 \\ \nu_2 & 0 & 0 & 0 & 0 \\ 0 & \nu_3 & \nu_3 & \nu_3 & \nu_3 \\ 0 & \nu_4 & \nu_4 & 0 & 0 \\ 0 & \nu_5 & 0 & 0 & 0 \\ 0 & 0 & 0 & \nu_6 & 0 \\ 0 & 0 & 0 & \nu_7 & \nu_7 \\ 0 & 0 & 0 & \nu_8 & 0 \\ 0 & 0 & 0 & 0 & \nu_9 \end{pmatrix}$$

3) $A = C^T C = V^T D(\nu^2) V$

$$= \begin{pmatrix} \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \\ \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \\ \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \\ \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \\ \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \end{pmatrix} +$$

$$\begin{pmatrix} \nu_2^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} +$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & \nu_3^2 & \nu_3^2 & \nu_3^2 & \nu_3^2 \\ 0 & \nu_3^2 & \nu_3^2 & \nu_3^2 & \nu_3^2 \\ 0 & \nu_3^2 & \nu_3^2 & \nu_3^2 & \nu_3^2 \\ 0 & \nu_3^2 & \nu_3^2 & \nu_3^2 & \nu_3^2 \end{pmatrix} +$$

$$\ldots +$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \nu_9^2 \end{pmatrix}$$

$$V = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

4)

$$A = \begin{pmatrix} \nu_1^2 + \nu_2^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 & \nu_1^2 \\ \nu_1^2 & \nu_1^2 + \nu_3^2 + \nu_4^2 + \nu_5^2 & \nu_1^2 + \nu_3^2 + \nu_4^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 \\ \nu_1^2 & \nu_1^2 + \nu_3^2 + \nu_4^2 & \nu_1^2 + \nu_3^2 + \nu_4^2 + \nu_6^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 \\ \nu_1^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 + \nu_7^2 + \nu_8^2 & \nu_1^2 + \nu_3^2 + \nu_7^2 \\ \nu_1^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 & \nu_1^2 + \nu_3^2 + \nu_7^2 & \nu_1^2 + \nu_3^2 + \nu_7^2 + \nu_9^2 \end{pmatrix}$$

Figure 5: Consumer-resource framework for model 1. 1) Each clade has access to its private resource, as encoded in matrix V (the rows are the resources, the columns the species, and a coefficient is one if the species can access the resource). 2) Each resource has a value, stored in vector ν; in this case the consumption matrix is simply $C = D(\nu)V$. 3) The competition matrix A is obtained by summing the competition coefficients (ν_j^2) for each clade a species belongs to. 4) The resulting matrix is block-structured, with a block per clade.

We consider n species of consumers along with their phylogenetic tree. We imagine the situation in which each clade in the tree has access to a clade-specific resource. Thus we can build a matrix V with $2n-1$ rows (the resources, one per clade, including a resource accessible to all species) and n columns (the consumers). The matrix has a coefficient of $V_{ij} = 1$ whenever consumer j has access to resource i, and zero elsewhere. Next, we model consumption, by assuming that each resource has a certain value (taken to be the same irrespective of the consumer), and that each consumer has a given attack rate (taken to be the same for each resource):

$$C = D(\nu)VD(\alpha) = \nu\alpha^T \circ V \quad ,$$

where $D(\nu)$ is a diagonal matrix with the value of each resource on the diagonal, $D(\alpha)$ a diagonal matrix of

attack rates and ∘ is the Hadamard product. We obtain a system of $2n - 1 + n$ differential equations, which follow the Generalized Lotka-Volterra model:

$$
\begin{cases}
\frac{dx_i}{dt} = x_i \left(-\delta_i + \sum_{j=1}^{2n-1} C_{ji} y_j \right) \\
\frac{dy_j}{dt} = y_j \left(\rho_j - \kappa_j y_j - \sum_{i=1}^{n} C_{ji} x_i \right)
\end{cases} ,
$$

in which $x_i = x_i(t)$ is the density of consumer i, $y_j = y_j(t)$ that of resource j, and all parameters are positive. δ_i is the intrinsic death rate of consumer i, ρ_j is the growth rate of resource j and κ_j its self-limitation. If we assume that resources quickly equilibrate for any level of consumer (practically, if $\rho_j \gg \sum_{i=1}^{n} C_{ji} x_i$), we can perform a "separation of time-scales" (as in MacArthur 1970), and solve the equation for the equilibrium of y_j:

$$
y_j^\star = \frac{1}{\kappa_j} \left(\rho_j - \sum_{i=1}^{n} C_{ji} x_i \right) \quad .
$$

Plugging these values into the equations for the consumers yields:

$$
\begin{aligned}
\frac{dx_i}{dt} &= x_i \left(-\delta_i + \sum_j C_{ij} \frac{1}{\kappa_j} \left(\rho_j - \sum_i C_{ji} x_i \right) \right) \\
&= x_i \left(-\delta_i + \sum_j C_{ij} \frac{\rho_j}{\kappa_j} + \sum_j \sum_k C_{ji} \frac{1}{\kappa_j} C_{jk} x_k \right) , \\
&= x_i \left(\zeta_i - \sum_k B_{ik} x_k \right)
\end{aligned}
$$

where we have defined $\zeta_i = \left(\sum_j C_{ij} \rho_j / \kappa_j \right) - \delta_i$ and $B = C^T D(\kappa)^{-1} C$. Finally, we move ζ_i outside the parenthesis, obtaining:

$$
\frac{dx_i}{dt} = x_i \zeta_i \left(1 - \sum_k \frac{1}{\zeta_i} B_{ik} x_k \right) \quad .
$$

The matrix of interactions is therefore defined as:

$$
\begin{aligned}
A &= D(\zeta)^{-1} B = D(\zeta)^{-1} C^T D(\kappa)^{-1} C \\
&= D(\zeta)^{-1} D(\alpha) V^T D(\nu) D(\kappa)^{-1} D(\nu) V D(\alpha) \\
&= D(\gamma) V^T D(\lambda) V D(\theta) ,
\end{aligned}
$$

where $\theta = \alpha$, $\gamma = \frac{\alpha}{\zeta}$ and $\lambda = \frac{\nu^2}{\kappa}$. This is exactly the Eq. 1 in the main text, which we use for model 4. Further assuming that $\zeta = 1/\alpha$ we obtain model 2; choosing $\alpha = 1$ yields model 3 and, finally, using both simplifications we obtain model 1. Figure 5 illustrates the parameterization for the case of the simplest model: $A = V^T D(\lambda) V$. We note that the variance-covariance matrix of a tree with branch lengths λ would have the same form; as such, an alternative parameterization is thus to take competition coefficients to be the result of the evolution via Brownian motion of many traits (Ives and Helmus 2011).

Analysis of previous approaches

Traditional approaches used to fit data produced in Biodiversity-Ecosystem Functioning experiments rely on regression models that consider the effects of biodiversity, in terms of species richness and composition, and some measurement of ecosystem function, such as biomass, carbon fixation, and litter decomposition.

These models usually consider total biomass of each plot as the response variable, and regress it against the number of species or other properties of the plots; for example, they consider a design matrix of zeros and ones reporting the presence/absence of each species in a given plot. These models can be supplemented by measures of phylogenetic distance between the species, species' functional groups, species-specific effects, pairwise interactions, etc. Because of their underlying structure, these models do not take full advantage of the data, which usually reports individual biomasess for each species in each plot—the data gets aggregated when computing the total biomass per plot. These models suffer from three main limitations: (i) the use of aggregate data to fit the models when measurements are made on an individual basis; (ii) lack of association/compatibility with population dynamical models (but see Parain et al. 2019); and (iii) underlying assumptions regarding the error structure of the statistical models, which are not in accordance with the structure of the data or the assumptions of the model. Following the formulation by Connolly et al. (2011), a possible regression model would have the following structure:

$$y_i = \beta_0 + \sum_{j=1}^{S} \beta_j P_{ij} + \delta_{AV} \sum_{j<k}^{S} P_{ij}P_{ik} + +\kappa C_{D_i} + \epsilon_i \tag{4}$$

where y_i is a measurement of ecosystem function for plot i (e.g., total biomass, respiration, oxidation), S is the total number of species considered in the experiment (total species richness), β_0 is an intercept, β_j is the effect of species j in monoculture, P_{ij} marks the presence of species j in plot i, δ_{AV} is the average interaction effect, C_{D_i} is the phylogenetic diversity in plot i, computed as $C_{D_i} = \sum_{j<k}^{S}(D_{jk} - \bar{D})P_j P_k$, where D_{jk} is the phylogenetic distance between species j and k (accounted as the sum of branch lengths between the two species, $\bar{D}$ is the average distance of all species in the pool, and ϵ_i is the error term.

The amount of data needed to fit can be very large. One way to overcome this limitation is to rely on simulations where data is generated in agreement with biological assumptions and test model performance with the simulated data. Here we simulate ecological communities of competing species following the Generalized Lotka-Volterra dynamics with the matrix of interaction given by the variance-covariance matrix describing a phylogenetic tree—what should be an ideal situation for fitting the model. Starting from a pool of $n = 10$ species, we assume all species have the same growth rates, so that dynamics are determined solely by phylogenetic relations. We draw a random tree, computed its variance-covariance matrix, and assembled all $2^n - 1 = 1023$ communities that can be formed from that species pool and recorded each species' biomass in each community. The structure of the matrix of interactions can be thought as emerging from models of trait evolution; we follow the evolution of many trait values, each changing under Brownian motion, giving rise to the variance-covariance matrix which we associated with competition coefficients; in this way, differences in trait value are proportional to evolutionary distances (Ives and Helmus 2011). One key aspect of considering the matrix of interaction as traits evolving under Brownian motion is that all sub-communities that can be formed with a subset of species are feasible and dynamically stable (see Serván et al. 2020 for details).

Having produced a large synthetic data set containing more than a thousand "plots", we can assess the quality of fit using a variety of models of varying complexity. Model $m0 : y \sim P$ regresses total biomass against presence/absence of species (P); model $m1 : y \sim P + C_D$ regresses the total biomass against presence absence of species (P) and the phylogenetic distance of species in the plot (C_D); $m2 : y \sim P + C_D + P : C_D$ is the same as $m1$ but also considers the interaction between presence and phylogenetic distance; $m3 : y \sim P + \sum P_i P_j$ considers the presence absence of each species and their pairwise interaction; $m4 : y \sim P + \sum P_i P_j + C_D$ is the same as $m3$ adding the phylogenetic distance of each plot; and finally $m5 : y \sim P + \sum P_i P_j + C_D + P : C_D$ is the same as $m4$ including the interaction between presence and phylogenetic distance. Below we show the plots for each of the models, including the predicted versus observed biomasses and diagnostic plots from linear regression.

Results show that a) none of the models can fully capture the simulated data (despite the lack of stochasticity); b) all of the models violate some of the main assumptions of linear regressions: residuals are not independent and normally distributed, variances are not constant, and thus Quantile-Quantile plots deviate substantially from the 1:1 line. We conclude that these models structures are not adequate to capture models for population dynamics—departing from competition, choosing more complex models for population

 dynamics, or introducing variation and stochasticity can only worsen the fit.

Figure 6: Fitting regression models to simulated data from a phylogenetically-structured Generalized Lotka-Volterra model. Each figure reports the structure of the model (top) as well as scatterplots showing the relationship between observed and fitted total biomasses per plot; corresponding distribution of standardized residuals (for linear regression, these should be centered at zero and have the same variance irrespective of the fitted values); Quantile-Quantile plots (points should fall on the 1 to 1 line); scale location plots, showing whether residuals change with the input (they should form a horizontal band of points).

Figure 7: Fitting regression models to simulated data from a phylogenetically-structured Generalized Lotka-Volterra model. Each figure reports the structure of the model (top) as well as scatterplots showing the relationship between observed and fitted total biomasses per plot; corresponding distribution of standardized residuals (for linear regression, these should be centered at zero and have the same variance irrespective of the fitted values); Quantile-Quantile plots (points should fall on the 1 to 1 line); scale location plots, showing whether residuals change with the input (they should form a horizontal band of points).

Figure 8: Fitting regression models to simulated data from a phylogenetically-structured Generalized Lotka-Volterra model. Each figure reports the structure of the model (top) as well as scatterplots showing the relationship between observed and fitted total biomasses per plot; corresponding distribution of standardized residuals (for linear regression, these should be centered at zero and have the same variance irrespective of the fitted values); Quantile-Quantile plots (points should fall on the 1 to 1 line); scale location plots, showing whether residuals change with the input (they should form a horizontal band of points).

Figure 9: Fitting regression models to simulated data from a phylogenetically-structured Generalized Lotka-Volterra model. Each figure reports the structure of the model (top) as well as scatterplots showing the relationship between observed and fitted total biomasses per plot; corresponding distribution of standardized residuals (for linear regression, these should be centered at zero and have the same variance irrespective of the fitted values); Quantile-Quantile plots (points should fall on the 1 to 1 line); scale location plots, showing whether residuals change with the input (they should form a horizontal band of points).

Figure 10: Fitting regression models to simulated data from a phylogenetically-structured Generalized Lotka-Volterra model. Each figure reports the structure of the model (top) as well as scatterplots showing the relationship between observed and fitted total biomasses per plot; corresponding distribution of standardized residuals (for linear regression, these should be centered at zero and have the same variance irrespective of the fitted values); Quantile-Quantile plots (points should fall on the 1 to 1 line); scale location plots, showing whether residuals change with the input (they should form a horizontal band of points).

Full results

The results for each data set are reported in Table 1. The scatterplots for the data stemming from the Biodiversity II and the Wageningen experiments are reported in Figure 11 and Figure 13, respectively. The histograms of the likelihoods for the randomized trees are reported in Figure 12 and Figure 14. The fitted trees and associated boxplots with likelihoods for randomizations keeping certain clades are reported in Figure 15 - Figure 22 for the Biodiversity II experiment and Figure 23 - Figure 30 for the Wageningen experiment. And finally the fits for models 2-4 with the associated tree and vectors are also in **?@fig-models2-4**.

Table 1: For each data set (Bio II, Tilman et al. (2001); Cadotte, Cadotte (2013); Wag, Van Ruijven and Berendse (2010)), we report the number of species (nsp); the number of unique assemblages (ncomm); the number of measurements (nmeas); the model used to fit the data; the corresponding log-likelihood (logLik); the total number of parameters (npar); the p-values for randomized tree either produced by the Yule model (prnd) or by shuffling the species' identities (pshuffle); and the correlation between the logarithms of the observed and predicted biomasses (cor).

data set	n_{sp}	n_{comm}	n_{meas}	model	logLik	n_{par}	p_{rnd}	$p_{shuffle}$	cor
Bio II 6/01	13	28	142	1	-515.125	26	**0.011**	**0.006**	0.916
				2	-506.803	38	**0.001**	**0.004**	0.924
				3	-510.384	38	0.606	0.612	0.923
				4	-496.525	50	0.153	0.137	0.936
Bio II 7/06	20	63	416	1	-1 562.862	40	**0.032**	**0.034**	0.848
				2	-1 543.540	59	**0.028**	**0.032**	0.857
				3	-1 548.530	59	**0.013**	**0.028**	0.856
				4	-1 528.100	78	**0.001**	**0.007**	0.866
Bio II 8/07	20	69	445	1	-1 829.444	40	**0.017**	**0.014**	0.829
				2	-1 810.571	59	**0.020**	**0.018**	0.842
				3	-1 816.684	59	**0.033**	0.056	0.840
				4	-1 791.473	78	**0.049**	**0.041**	0.852
Bio II 7/08	17	33	247	1	-1 019.927	34	**0.008**	**0.008**	0.833
				2	-1 005.384	50	**0.045**	**0.049**	0.848
				3	-1 014.225	50	**0.030**	**0.024**	0.842
				4	-989.244	66	**0.014**	**0.008**	0.854
Bio II 8/11	20	59	451	1	-1 872.468	40	0.147	0.188	0.865
				2	-1 851.060	59	**0.037**	**0.033**	0.876
				3	-1 845.121	59	**0.002**	**0.005**	0.880
				4	-1 822.608	78	**0.000**	**0.000**	0.888
Bio II 7/12	16	40	322	1	-1 265.636	32	0.253	0.261	0.814
				2	-1 257.352	47	0.141	0.118	0.821
				3	-1 256.901	47	0.193	0.216	0.822

Table 1: For each data set (Bio II, Tilman et al. (2001); Cadotte, Cadotte (2013); Wag, Van Ruijven and Berendse (2010)), we report the number of species (nsp); the number of unique assemblages (ncomm); the number of measurements (nmeas); the model used to fit the data; the corresponding log-likelihood (logLik); the total number of parameters (npar); the p-values for randomized tree either produced by the Yule model (prnd) or by shuffling the species' identities (pshuffle); and the correlation between the logarithms of the observed and predicted biomasses (cor).

data set	n_{sp}	n_{comm}	n_{meas}	model	logLik	n_{par}	p_{rnd}	$p_{shuffle}$	cor
				4	-1 239.699	62	**0.009**	**0.009**	0.843
Bio II 8/14	16	39	271	1	-1 100.731	32	**0.000**	**0.001**	0.832
				2	-1 089.324	47	**0.027**	**0.019**	0.844
				3	-1 093.747	47	**0.006**	**0.003**	0.838
				4	-1 077.425	62	**0.012**	**0.013**	0.856
Bio II 8/15	18	43	337	1	-1 344.643	36	**0.005**	**0.012**	0.830
				2	-1 335.333	53	**0.002**	**0.004**	0.835
				3	-1 333.390	53	**0.027**	**0.028**	0.840
				4	-1 315.121	70	**0.004**	**0.009**	0.854
Bio II 7/17	19	53	409	1	-1 599.741	38	**0.001**	**0.001**	0.841
				2	-1 587.881	56	**0.011**	**0.014**	0.850
				3	-1 580.571	56	**0.002**	**0.000**	0.851
				4	-1 562.737	74	**0.001**	**0.000**	0.863
Cadotte	8	27	87	1	-348.372	16	**0.027**	**0.029**	0.878
				2	-344.637	23	0.131	0.132	0.890
				3	-345.628	23	0.512	0.507	0.881
				4	-342.545	30	0.696	0.711	0.891
Wag 00	8	58	235	1	-856.445	16	0.419	0.378	0.948
				2	-850.295	23	0.456	0.435	0.951
				3	-823.108	23	0.246	0.248	0.960
				4	-817.592	30	0.260	0.297	0.962
Wag 01	8	58	231	1	-815.679	16	**0.001**	**0.001**	0.933
				2	-811.568	23	0.103	0.115	0.935
				3	-786.491	23	**0.000**	**0.001**	0.948
				4	-780.650	30	0.071	0.066	0.950
Wag 02	8	56	222	1	-770.436	16	0.067	0.066	0.858
				2	-756.722	23	**0.040**	**0.024**	0.879
				3	-754.152	23	0.068	0.071	0.877

Table 1: For each data set (Bio II, Tilman et al. (2001); Cadotte, Cadotte (2013); Wag, Van Ruijven and Berendse (2010)), we report the number of species (nsp); the number of unique assemblages (ncomm); the number of measurements (nmeas); the model used to fit the data; the corresponding log-likelihood (logLik); the total number of parameters (npar); the p-values for randomized tree either produced by the Yule model (prnd) or by shuffling the species' identities (pshuffle); and the correlation between the logarithms of the observed and predicted biomasses (cor).

data set	n_{sp}	n_{comm}	n_{meas}	model	logLik	n_{par}	p_{rnd}	$p_{shuffle}$	cor
				4	-726.884	30	**0.003**	**0.003**	0.908
Wag 03	8	51	196	1	-754.342	16	**0.043**	**0.043**	0.849
				2	-737.109	23	**0.036**	**0.018**	0.873
				3	-731.931	23	0.076	**0.027**	0.880
				4	-713.488	30	**0.030**	**0.028**	0.903
Wag 04	8	53	184	1	-731.749	16	0.145	0.140	0.796
				2	-721.201	23	0.115	0.088	0.823
				3	-699.034	23	**0.005**	**0.003**	0.853
				4	-690.901	30	0.214	0.215	0.871
Wag 05	8	53	197	1	-816.254	16	0.060	**0.049**	0.766
				2	-807.928	23	0.125	0.071	0.786
				3	-804.602	23	**0.028**	**0.020**	0.802
				4	-792.838	30	0.380	0.387	0.820
Wag 06	8	54	197	1	-746.003	16	0.106	0.104	0.721
				2	-740.846	23	0.239	0.243	0.741
				3	-720.442	23	**0.028**	**0.028**	0.790
				4	-714.023	30	0.166	0.166	0.803
Wag 07	8	55	206	1	-808.122	16	**0.023**	**0.032**	0.769
				2	-799.968	23	0.148	0.152	0.788
				3	-791.224	23	**0.036**	0.075	0.807
				4	-786.148	30	0.080	0.077	0.809
Wag 08	8	54	210	1	-803.785	16	0.278	0.282	0.780
				2	-792.913	23	0.986	0.961	0.797
				3	-794.166	23	0.236	0.271	0.799
				4	-779.235	30	1.000	1.000	0.822

Figure 11: Scatterplot of predicted vs. observed biomass for the Biodiversity II data. Each panel represents a specific sampling date and model. The correlation between the logarithm of the observed biomass and the logarithm of the predicted biomass is reported in each panel.

Figure 12: Histograms of the log-likelihoods for models parameterized using random tree and data from the Biodiversity II experiment. The distributions for the random trees generated by the Yule model are in teal, and those for the shuffled trees in grey. Each panel reports the p-values obtained for the Yule and shuffled trees, respectively.

Figure 13: Scatterplot of predicted vs. observed biomass for the Wageningen biodiversity experiment data. Each panel represents a specific sampling date and model. The correlation between the logarithm of the observed biomass and the logarithm of the predicted biomass is reported in each panel.

Figure 14: Histograms of the log-likelihoods for models parameterized using random tree and data from the Wageningen diversity experiment. The distributions for the random trees generated by the Yule model are in teal, and those for the shuffled trees in grey. Each panel reports the p-values obtained for the Yule and shuffled trees, respectively.

Figure 15: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 16: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 17: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 18: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 19: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 20: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 21: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 22: Resulting trees for the Biodiversity II experiments in Cedar Creek, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 23: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 24: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 25: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 26: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 27: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 28: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 29: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 30: Resulting trees for the Wageningen experiments, fit for phylogenetic tree from model 1 and boxplots of likelihoods for randomizations that kept the selected clades highlighted in the trees. Dashed line represents likelihood of the original tree

Figure 31: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 32: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 33: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 34: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 35: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 36: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 37: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 38: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 39: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 40: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 41: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 42: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 43: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 44: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 45: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 46: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 47: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 48: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

Figure 49: Fit for the phylogenetic tree and associated parameter vectors (Gamma and Theta) for models 2-4. The length and color of the branches correspond to increased strength in competition between the species in the clade and the size of the vector corresponds to the fitted value for the corresponding species in the tree. bioII refers to the Biodiversity II experiment (Tilman et al. (2001)); cadotte is from Cadotte (2013); and vr refers to the Wageningen experiment from Van Ruijven and Berendse (2010).

References

Ansari, Aamir Faisal, Yugandhar BS Reddy, Janhavi Raut, and Narendra M Dixit. 2021. "An Efficient and Scalable Top-down Method for Predicting Structures of Microbial Communities." *Nature Computational Science* 1 (9): 619–28.

Bravo, Héctor Corrada, Stephen Wright, Kevin Eng, Sunduz Keles, and Grace Wahba. 2009. "Estimating Tree-Structured Covariance Matrices via Mixed-Integer Programming." In *Artificial Intelligence and Statistics*, 41–48. PMLR.

Cadotte, Marc W. 2013. "Experimental Evidence That Evolutionarily Diverse Assemblages Result in Higher Productivity." *Proceedings of the National Academy of Sciences* 110 (22): 8996–9000.

———. 2015. "Phylogenetic Diversity and Productivity." *Functional Ecology* 29 (12): 1603–6.

Cadotte, Marc W, Bradley J Cardinale, and Todd H Oakley. 2008. "Evolutionary History and the Effect of Biodiversity on Plant Productivity." *Proceedings of the National Academy of Sciences* 105 (44): 17012–17.

Cardinale, Bradley J, Patrick Venail, Kevin Gross, Todd H Oakley, Anita Narwani, Eric Allan, Pedro Flombaum, et al. 2015. "Further Re-Analyses Looking for Effects of Phylogenetic Diversity on Community Biomass and Stability." *Functional Ecology* 29 (12): 1607–10.

Cavender-Bares, Jeannine, Adrienne Keen, and Brianna Miles. 2006. "Phylogenetic Structure of Floridian Plant Communities Depends on Taxonomic and Spatial Scale." *Ecology* 87 (sp7): S109–22.

Cavender-Bares, Jeannine, Kenneth H Kozak, Paul VA Fine, and Steven W Kembel. 2009. "The Merging of Community Ecology and Phylogenetic Biology." *Ecology Letters* 12 (7): 693–715.

Connolly, John, Thomas Bell, Thomas Bolger, Caroline Brophy, Timothee Carnus, John A Finn, Laura Kirwan, et al. 2013. "An Improved Model to Predict the Effects of Changing Biodiversity Levels on Ecosystem Function." *Journal of Ecology* 101 (2): 344–55.

Connolly, John, Marc W Cadotte, Caroline Brophy, Áine Dooley, John Finn, Laura Kirwan, Christiane Roscher, and Alexandra Weigelt. 2011. "Phylogenetically Diverse Grasslands Are Associated with Pairwise Interspecific Processes That Increase Biomass." *Ecology* 92 (7): 1385–92.

Darwin, Charles. 2004. *On the Origin of Species, 1859.* Routledge.

Davies, T Jonathan. 2021. "Ecophylogenetics Redux." *Ecology Letters* 24 (5): 1073–88.

Davies, T Jonathan, Mark C Urban, Bronwyn Rayfield, Marc W Cadotte, and Pedro R Peres-Neto. 2016. "Deconstructing the Relationships Between Phylogenetic Diversity and Ecology: A Case Study on Ecosystem Functioning." *Ecology* 97 (9): 2212–22.

Díaz, Sandra, and Marcelo Cabido. 2001. "Vive La Différence: Plant Functional Diversity Matters to Ecosystem Processes." *Trends in Ecology & Evolution* 16 (11): 646–55.

Elton, Charles. 1946. "Competition and the Structure of Ecological Communities." *The Journal of Animal Ecology*, 54–68.

Flynn, Dan FB, Nicholas Mirotchnick, Meha Jain, Matthew I Palmer, and Shahid Naeem. 2011. "Functional and Phylogenetic Diversity as Predictors of Biodiversity–Ecosystem-Function Relationships." *Ecology* 92 (8): 1573–81.

Fort, Hugo. 2018. "On Predicting Species Yields in Multispecies Communities: Quantifying the Accuracy of the Linear Lotka-Volterra Generalized Model." *Ecological Modelling* 387: 154–62.

Frainer, André, Brendan G McKie, Per-Arne Amundsen, Rune Knudsen, and Kevin D Lafferty. 2018. "Parasitism and the Biodiversity-Functioning Relationship." *Trends in Ecology & Evolution* 33 (4): 260–68.

Fry, Ellen L, Sally A Power, and Pete Manning. 2014. "Trait-Based Classification and Manipulation of Plant Functional Groups for Biodiversity–Ecosystem Function Experiments." *Journal of Vegetation Science* 25 (1): 248–61.

Graham, Catherine H, Juan L Parra, Carsten Rahbek, and Jimmy A McGuire. 2009. "Phylogenetic Structure in Tropical Hummingbird Communities." *Proceedings of the National Academy of Sciences* 106 (supplement_2): 19673–78.

Huang, Mengjiao, Xiang Liu, Marc W Cadotte, and Shurong Zhou. 2020. "Functional and Phylogenetic Diversity Explain Different Components of Diversity Effects on Biomass Production." *Oikos* 129 (8): 1185–95.

Ives, Anthony R, and Matthew R Helmus. 2011. "Generalized Linear Mixed Models for Phylogenetic Analyses of Community Structure." *Ecological Monographs* 81 (3): 511–25.

Jin, Yi, and Hong Qian. 2019. "V. PhyloMaker: An r Package That Can Generate Very Large Phylogenies for Vascular Plants." *Ecography* 42 (8): 1353–59.

Jochum, Malte, Markus Fischer, Forest Isbell, Christiane Roscher, Fons van der Plas, Steffen Boch, Gerhard Boenisch, et al. 2020. "The Results of Biodiversity–Ecosystem Functioning Experiments Are Realistic." *Nature Ecology & Evolution* 4 (11): 1485–94.

Keck, François, and Maria Kahlert. 2019. "Community Phylogenetic Structure Reveals the Imprint of Dispersal-Related Dynamics and Environmental Filtering by Nutrient Availability in Freshwater Diatoms." *Scientific Reports* 9 (1): 11590.

Losos, Jonathan B. 2008. "Phylogenetic Niche Conservatism, Phylogenetic Signal and the Relationship Between Phylogenetic Relatedness and Ecological Similarity Among Species." *Ecology Letters* 11 (10): 995–1003.

MacArthur, Robert. 1970. "Species Packing and Competitive Equilibrium for Many Species." *Theoretical Population Biology* 1 (1): 1–11.

Maherali, Hafiz, and John N Klironomos. 2007. "Influence of Phylogeny on Fungal Community Assembly and Ecosystem Functioning." *Science* 316 (5832): 1746–48.

Mayfield, Margaret M, and Jonathan M Levine. 2010. "Opposing Effects of Competitive Exclusion on the Phylogenetic Structure of Communities." *Ecology Letters* 13 (9): 1085–93.

Maynard, Daniel S, Zachary R Miller, and Stefano Allesina. 2020. "Predicting Coexistence in Experimental Ecological Communities." *Nature Ecology & Evolution* 4 (1): 91–100.

Mazel, Florent, Arne O Mooers, Giulio Valentino Dalla Riva, and Matthew W Pennell. 2017. "Conserving Phylogenetic Diversity Can Be a Poor Strategy for Conserving Functional Diversity." *Systematic Biology* 66 (6): 1019–27.

Mazel, Florent, Matthew W Pennell, Marc W Cadotte, Sandra Diaz, Giulio Valentino Dalla Riva, Richard Grenyer, Fabien Leprieur, et al. 2018. "Prioritizing Phylogenetic Diversity Captures Functional Diversity Unreliably." *Nature Communications* 9 (1): 2888.

Parain, Elodie C, Rudolf P Rohr, Sarah M Gray, and Louis-Felix Bersier. 2019. "Increased Temperature Disrupts the Biodiversity–Ecosystem Functioning Relationship." *The American Naturalist* 193 (2): 227–39.

Petchey, Owen L, and Kevin J Gaston. 2006. "Functional Diversity: Back to Basics and Looking Forward." *Ecology Letters* 9 (6): 741–58.

R Core Team. 2021. *R: A Language and Environment for Statistical Computing.* Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Ruijven, Jasper van, and Frank Berendse. 2005. "Diversity–Productivity Relationships: Initial Effects, Long-Term Patterns, and Underlying Mechanisms." *Proceedings of the National Academy of Sciences* 102 (3): 695–700.

Serván, Carlos A., José A. Capitán, Zachary R. Miller, and Stefano Allesina. 2020. "Effects of Phylogeny on Coexistence in Model Communities." *bioRxiv.* https://doi.org/10.1101/2020.09.04.283507.

Simberloff, Daniel S. 1970. "Taxonomic Diversity of Island Biotas." *Evolution,* 23–47.

Skwara, Abigail, Paula Lemos-Costa, Zachary R Miller, and Stefano Allesina. 2023. "Modelling Ecological Communities When Composition Is Manipulated Experimentally." *Methods in Ecology and Evolution* 14 (2): 696–707.

Stan Development Team. 2023. *Stan Modeling Language Users Guide and Reference Manual, 2.32.* https://mc-stan.org.

Swenson, Nathan G. 2019. *Phylogenetic Ecology: A History, Critique, and Remodeling.* University of Chicago Press.

Tilman, David, Peter B Reich, Johannes Knops, David Wedin, Troy Mielke, and Clarence Lehman. 2001. "Diversity and Productivity in a Long-Term Grassland Experiment." *Science* 294 (5543): 843–45.

Tucker, Caroline M, T Jonathan Davies, Marc W Cadotte, and William D Pearse. 2018. "On the Relationship Between Phylogenetic Diversity and Trait Diversity." *Ecology* 99 (6): 1473–79.

Van Ruijven, Jasper, and Frank Berendse. 2010. "Diversity Enhances Community Recovery, but Not Resistance, After Drought." *Journal of Ecology* 98 (1): 81–86.

Venail, Patrick, Kevin Gross, Todd H Oakley, Anita Narwani, Eric Allan, Pedro Flombaum, Forest Isbell, et al. 2015. "Species Richness, but Not Phylogenetic Diversity, Influences Community Biomass Production and Temporal Stability in a Re-Examination of 16 Grassland Biodiversity Studies." *Functional Ecology*

29 (5): 615–26.

Webb, Campbell O. 2000. "Exploring the Phylogenetic Structure of Ecological Communities: An Example for Rain Forest Trees." *The American Naturalist* 156 (2): 145–55.

Webb, Campbell O, David D Ackerly, Mark A McPeek, and Michael J Donoghue. 2002. "Phylogenies and Community Ecology." *Annual Review of Ecology and Systematics* 33 (1): 475–505.

Wright, Alexandra J, David A Wardle, Ragan Callaway, and Aurora Gaxiola. 2017. "The Overlooked Role of Facilitation in Biodiversity Experiments." *Trends in Ecology & Evolution* 32 (5): 383–90.

Wright, Justin P, Shahid Naeem, Andy Hector, Clarence Lehman, Peter B Reich, Bernhard Schmid, and David Tilman. 2006. "Conventional Functional Classification Schemes Underestimate the Relationship with Ecosystem Functioning." *Ecology Letters* 9 (2): 111–20.

Xiao, Yandong, Marco Tulio Angulo, Jonathan Friedman, Matthew K Waldor, Scott T Weiss, and Yang-Yu Liu. 2017. "Mapping the Ecological Networks of Microbial Communities." *Nature Communications* 8 (1): 2042.