Experimental nerve block study on painful withdrawal reflex responses in humans

Oumie Thorell[1,2*], David A. Mahns[1], Jan Otto[3], Jaquette Liljencrantz[4], Mats Svantesson[2], Håkan

Olausson[2], Saad S. Nagi[1,2]

[1]School of Medicine, Western Sydney University, Australia; [2]Department of Biomedical and Clinical

Sciences, Linköping University, Sweden; [3]Department of Neurology, University Hospital Schleswig-

Holstein Kiel, Germany; [4]Department of Anesthesiology and Intensive Care, Institute of Clinical

Sciences, Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden

* Corresponding author

E-mail: oumie.thorell@liu.se (OT)

# Abstract

The nociceptive withdrawal reflex (NWR) is a protective limb withdrawal response triggered by painful stimuli, used to assess spinal nociceptive excitability. Conventionally, the NWR is understood as having two reflex responses: a short-latency Aβ-mediated response, considered tactile, and a longer-latency Aδ-mediated response, considered nociceptive. However, nociceptors with conduction velocities similar to Aβ tactile afferents have been identified in human skin. In this study, we investigated the effect of a preferential conduction block of Aβ fibers on pain perception and NWR signaling evoked by intradermal electrical stimulation in healthy participants. We recorded a total of 198 NWR responses in the intact condition, and no dual reflex responses occurred within our latency bandwidth (50-150 ms). The current intensity required to evoke the NWR was magnitude higher than the perceptual pain threshold, indicating that NWR did not occur before pain was felt. In the block condition, when the Aβ-mediated tuning fork sensation was lost while Aδ-mediated nonpainful cooling was still detectable (albeit reduced), we observed that the reflex was abolished. Further, short-latency electrical pain intensity at pre-block thresholds was greatly reduced, with any residual pain sensation having a longer latency. Although electrical pain was unaffected at suprathreshold current intensities, the reflex could not be evoked despite a two-fold increase in the pre-block current intensity and a five-fold increase in the pre-block pulse duration. These observations lend support to the possible involvement of Aβ-fiber inputs in pain and reflex signaling.

## Introduction

Pain involves cognitive, genetic, and psychosocial factors [1-4]. Currently, pain assessment mainly relies on individuals' self-report which has limitations. In the early 20[th] century, reflexes were extensively studied by Sherrington who observed coordinated muscle movements, such as flexion followed by stepping movements, that correlated with the intensity of noxious stimulation [5]. This phenomenon is often termed the nociceptive withdrawal reflex (NWR) and involves a complex interplay between top-down and bottom-up influences [6-9]. It was initially suggested that the NWR may serve as a readout of pain, but several studies have since highlighted that the relationship between pain and reflex is not clearcut [10-16]. Nonetheless, the reflex is a useful tool for monitoring the excitability of nociceptive spinal systems.

The NWR is often reported as consisting of two distinct electromyographic (EMG) responses with an intervening silent period [10, 17, 18]. These responses, RII and RIII, are attributed to different peripheral afferents: RII to large, thickly myelinated afferents with $A\beta$-fiber conduction velocities (CV) and RIII to smaller, thinly myelinated afferents with $A\delta$-fiber CVs [19]. It is generally argued that the RII is non-painful, and it is the RIII that represents spinal nociceptive signaling [10, 20, 21]. Thus, it is common practice to exclude the first NWR component from the reflex analysis. However, there is no consensus on where in time the separation between RII and RIII should occur. Using a 90-ms cutoff, for instance, it was found that NWR responses shorter than 90 ms were just as painful as those that were ≥90 ms [22]. Further, the NWR may comprise of single EMG responses occurring at different latencies and stimulation intensities, questioning the involvement of distinct peripheral afferent classes [20, 22].

It was recently reported that human skin is equipped with a specific class of high-threshold mechanoreceptors with Aβ conduction velocities. These receptors encode noxious mechanical stimuli and produce painful percepts when selectively activated through low-current intraneural stimulation [23-25]. This discovery raises the question of whether Aβ inputs contribute to painful NWR signaling in humans. Part of the ambiguity in the literature around latencies is due to the paucity of direct recordings from Aδ afferent fibers in humans, thereby relying on indirect measurements to infer their conduction velocities. Indeed, in animals, the conduction velocity of D-hair afferents is used as a cutoff between Aβ and Aδ populations [26]; however, D-hair afferents have not yet been characterized in humans.

In the current study, we employed preferential Aβ-fiber conduction blocks and tested pain and reflex responses evoked by intradermal electrical stimulation before, during, and upon recovery from the block. Nerve conduction blocks are widely used in the somatosensory field to study the functions of primary afferent fibers [27-29]. In the block condition where tuning fork sensation was abolished while nonpainful cooling remained relatively preserved – readouts of Aβ- and Aδ-fiber activity, respectively – we found that reflex responses at pre-block thresholds could not be evoked. Further, short-latency pain intensity at pre-block electrical thresholds was significantly reduced during the block, with any residual pain sensation having a longer latency. Although pain could be evoked at higher stimulus intensities, the reflex could not be evoked during the block despite considerable increases to the pre-block stimulus intensities and duration, hinting at the potential involvement of Aβ fibers in driving our nocifensive behaviors.

## Methods

## Participants

Twenty-five healthy participants (17 males and 8 females), aged 18-39 years, took part in this study. The exclusion criteria included neurological and musculoskeletal disorders, skin diseases, diabetes, and the use of pain-relieving or psychoactive medications. The study was approved by the Swedish Ethical Review Authority (dnr 2020-04207). Written informed consent was obtained from all participants before the start of the experiment. The study was conducted in accordance with the Helsinki Declaration. Participants were seated comfortably in a chair with the knee flexed to ~130□.

## EMG recordings

Three self-adhesive recording electrodes (Kendall ECG electrodes, 57x34 mm, Medtronics, USA) were attached to the right tibialis anterior (TA) muscle serving as active, reference, and ground points. EMG recording settings comprised a 1 mV range, 1 kHz low pass filter, 0.3 Hz high pass filter, and 20 kHz sampling rate (LabChart v8.1.16, ADInstruments, Dunedin, New Zealand).

## Intradermal electrical stimulation

Two uninsulated tungsten microelectrodes (FHC Inc., Bowdoin, USA) were inserted just below the metatarsophalangeal joint of the right foot sole, separated by 5 mm, to deliver focal electrical stimulation. The microelectrodes had a tip diameter of 5-10 μm and a shaft of 200 μm. Each stimulus trial consisted of 5 square pulses delivered at a frequency of 200 Hz with a pulse duration of 0.2 ms. These stimuli were generated using a constant current bipolar stimulator (DS8R, Digitimer, Hydeway, UK). In cases where a reflex could not be evoked with a pulse duration of 0.2 ms, a longer pulse

110     duration of 1 ms was used. The interstimulus intervals were varied from trial to trial (at least >6 s) to

111     prevent habituation and/or cognitive suppression of the reflex response. To avoid visual and auditory

112     cues, a partition was placed between the participant and the experimenter, and a silent mouse was

113     used to trigger the stimuli.

## Perception and reflex threshold measurements

Participants were instructed to remain relaxed during the recordings, which were performed without

any muscle contraction. Current intensities were slowly increased in increments of 1-3 mA until the

first (nonpainful) sensation was reported. This was taken as the detection threshold ($DET^{th}$). The same

procedure was repeated to establish the minimum current required to evoke a painful percept (pain

threshold ($PAIN^{th}$). NWR thresholds ($NWR^{th}$) were determined based on at least two successful trials

at same current intensity. Participants were asked to rate the intensity of pain on a visual analog scale

(VAS) ranging from 0 to 10, with 0 representing "no pain" and 10 representing the "worst imaginable

pain" (Response meter, ADInstruments, Dunedin, New Zealand). Participants were instructed to move

the analog scale only if the sensation was perceived as painful, and they were free to interpret pain

according to their individual experiences. Pain qualities were captured using a short-form McGill Pain

Questionnaire [30] immediately following $PAIN^{th}$ and $NWR^{th}$ measurements (a total of six times).

## Reflex analysis

Reflex latencies, Z scores, and pain ratings were analyzed in MATLAB (r2021b, MathWorks Inc, Natick,

Massachusetts). Z scores were calculated as the difference between peak amplitude (50-150 ms post-

stimulus onset) and mean baseline amplitude (-0.15 to 0 ms relative to the stimulus onset), divided by

the standard deviation of baseline EMG activity. The minimum current intensity required to evoke a

reflex response was taken as the $NWR^{th}$. Responses with latencies exceeding 150 ms after stimulus

onset were excluded from the analysis to avoid voluntary/startle responses that can follow reflex

elicitation [31].

## Nerve block

An ischemic nerve block progressively affects large myelinated fibers that signal vibration, followed by

small myelinated fibers that signal innocuous cold, and finally, unmyelinated fibers that signal warmth

sensations [27-29, 32-34]. To induce the block, an air-filled pressure cuff (Riester Gmbh, Jungigen,

Germany) was placed over the right ankle and inflated to 300 mm Hg for up to an hour [35]. The block

was applied distal to the TA muscle EMG was recorded from.

In order to track the progression of the nerve block, vibration perception (test for Aβ-fiber function)

was tested on the foot sole in three ways: *1.* With a tuning fork (128 Hz, American Diagnostic

Corporation, NY, USA) using a two-alternative forced choice detection task (2AFC) where participants

reported whether the tuning fork was perceived as "vibration" or "no vibration"; *2.* With a punctate

Piezo electric stimulator (probe diameter: 1.3 cm, Dancer Design, UK) where participants rated the

intensity of vibration (20 or 200 Hz) on an analog scale ranging from 0 ("no vibrating sensation") to 10

("highest vibrating intensity"); *3.* Using a 3AFC detection task where participants reported whether

vibration at 200, 20, and 0 Hz was perceived as "high", "low" or "no" vibration. Participants wore

earplugs during vibration tests to prevent auditory cues. When participants could no longer

distinguish whether the tuning fork was stationary or vibrating, the blockade of Aβ fibers was

considered successful. In addition, it was expected that vibratory stimuli would be rated as less

intense during the block, and participants would be unable to discriminate between 20 and 200 Hz

frequencies.

To assess Aδ- and C-fiber functions, we conducted simple detection tasks by placing a cold and hot

metal rod, which had been immersed in ice and a water bath at 45°C, respectively, against the

metatarsophalangeal joint of the foot and contralateral (intact) foot sole every ~5 minutes.

Participants were asked to verbally report what they felt and whether the intensity between the two

sites was the same or different. This allowed for frequent testing of thermal sensibility. As soon as the

tuning fork sensation was lost, and other vibratory tests were performed, detection thresholds for

cooling and warming were measured on the foot sole using the method of limits. The thermode probe

had dimensions of 30 x 30 mm (TSA-II, Medoc Ltd., Ramat Yishai, Israel) and the rate of temperature

change was 1°C/s, starting from a neutral temperature of 32°C [36]. Each modality was tested four

times. In the condition where the vibration sense was blocked while temperature senses remained,

perceptual responses at $PAIN^{th}$ and reflex responses at $NWR^{th}$ were tested (at least 3 times).

In the pre-block condition, reaction time measurements were conducted 10 times at the $PAIN^{th}$

stimulus intensity. During this assessment, participants were asked to press a button as soon as they

felt a painful sensation. The inter-stimulus intervals were pseudorandomized (mean: 3.7 s, min: 1.1 s,

max: 7.5 s). During the block, if any pain was reported by the participants at the $PAIN^{th}$, the reaction

time measurements were tested again.

Upon release of the nerve block, vibratory and thermal sensibilities were monitored and upon

recovery to pre-block levels (typically within 20-30 min), $PAIN^{th}$ and $NWR^{th}$ were measured again.

## Control experiment

180 To confirm that any effect of the nerve block on pain and reflex responses was due to the blockade of

181 peripheral A-fiber inputs rather than central or other factors, in five participants we ran control

182 experiments with the ischemic cuff applied to the contralateral (left) leg. Pain and reflex responses

183 were measured from the standard (right) leg.

## Statistical analysis

The experiment followed a quantitative, repeated-measurement design in which participants were

always tested in three conditions: before (Baseline), during (Block), and after (Recovery) of the nerve

block. Descriptive statistics and analyses were performed in Prism (9.0.2, GraphPad Software, San

Diego, USA). Non-parametrical statistical tests were chosen because of the dataset's medium to small

sample sizes, non-normal distribution (as indicated by normality tests), skewed distribution (QQ-plots,

skewness, and kurtosis), and/or high standard deviation (SD) in relation to mean values (>50%).

Wilcoxon test was used to compare two related groups. Friedman's test was used to compare

multiple related groups and Kruskal-Wallis for multiple non-related groups. Dunn's test was used as a

post hoc for multiple comparisons. Spearman's rank was used to investigate correlations with a 95%

confidence interval (CI). A p-value of $< 0.05$ was considered statistically significant. The a-priori sample

size estimation was based on a pilot study where we observed an effect size (f) of 0.255. We then

used a 1-$\beta$ error probability (power) of 0.80, and $\alpha$ error probability of 0.05, which gave a total sample

size of 27. Post hoc power analysis, based on f(0.27), $\alpha$(0.05), and a sample size of 25, gave a power of

82.6. Sample size and power calculations were performed in G* Power (open software, v3.1.9.7). Data

are shown as median with interquartile range.

# Results

Under baseline conditions, NWR was successfully evoked in 22 out of 25 participants. The current

intensity required to reach $DET^{th}$ was the lowest, followed by $PAIN^{th}$, and finally $NWR^{th}$ (Fig 1A-B).

Consequently, all reflex thresholds occurred in response to a painful stimulus (pain intensity range:

0.3-6 on VAS of 0-10). A breakdown into individual reflex responses reveals that only 9 out of 198, or

less than 5%, were rated as nonpainful, and in no participant were two consecutive reflex responses

rated as nonpainful. The NWR responses had Z scores ranging from 1 to 61 and occurred at latencies

between 65 and 137 ms after stimulus onset (mean latency: 91 ms). Out of these, 80 (40.4%) NWR

responses had latencies under 90 ms, a cutoff for defining RII reflexes as used in prior studies [e.g. 15,

16, 37]. In another study, involving transcutaneous electrical stimulation, dual reflex responses (RII

and RIII) were observed in 12% of reflex recordings [22]. In the current study using intradermal

electrical stimulation, no instances of dual RII-RIII responses were observed (Fig 1C). In a subset of

participants (n = 7), the reflex could only be evoked using a 1-ms pulse duration. There were no

differences in pain ratings or NWR latencies at $PAIN^{th}$ and $NWR^{th}$ between the 0.2-ms and 1-ms

responses, hence the data were combined (Fig S1A-D).

**Fig 1. Characterization of pain and reflex responses evoked by intradermal electrical stimulation. A.**

The first trace shows the absence of an EMG response at the nonpainful detection threshold ($DET^{th}$).

The second trace shows a pain rating at the pain threshold ($PAIN^{th}$), although no EMG response was

detected. The third trace shows an EMG response at the NWR threshold ($NWR^{th}$). **B**. $DET^{th}$, $PAIN^{th}$, and

$NWR^{th}$ were significantly different ($DET^{th}$: 0.5 (0.2) mA, $PAIN^{th}$: 2.0 (1.5) mA, $NWR^{th}$: 13.5 (12.0) mA,

f(2) = 44.00, p < 0.0001, post hoc test: **P = 0.0027; ****P < 0.0001, n = 22, Friedman test). **C**. A total

of 198 NWR responses were recorded with no instances of dual EMG bursts within our latency

bandwidth (50-150 ms).

## Preferential block of Aβ fibers

Somatosensory tests were performed to gauge the progression of the ischemic nerve block. During

baseline and recovery conditions, participants performed with 100% accuracy in the vibration

discrimination tasks (2AFC, 3AFC). During the block for >20 min but <1 hour, participants could no

longer distinguish whether the tuning fork was stationary or vibrating. Further, the vibration intensity

ratings declined significantly (Fig 2A-B), and vibration discrimination was significantly impaired (Fig

2C).

Cold detection thresholds (CDTs) were significantly altered (median difference from baseline = 7.2°C)

during the nerve block (Fig. 2D). However, in no participant did the mean CDTs shift to the cold pain

threshold range (reported as ≤10-14°C) [38], indicating that cooling remained detectable within the

nonpainful range during the block. The change in CDT (baseline vs. block) was unrelated to the block

duration (Fig S2A). Further, considering 23°C as the lower border of normal values for innocuous cold

detection in the foot [39], we found no differences when comparing pain ratings (at $PAIN^{th}$) between

participants with CDT above or below the lower border of innocuous CDT (Fig S2B).

Warm detection thresholds (WDTs) were not significantly different between baseline and block

conditions, or between block and recovery conditions (Fig 2E). A significant correlation was found

between the change in WDT and the duration of the nerve block (Fig S2C).

## NWR abolished by preferential Aβ-fiber block

During the nerve block, all responses at pre-block $NWR^{th}$ were abolished (Fig. 2F). Despite further

increases in stimulus intensity (up to 2 times the pre-block $NWR^{th}$) and/or prolonging of the pulse

duration (extended to 1 ms), the reflex did not recover during the nerve block. This was true even for

those participants (n=5) whose block CDTs were within 1-3°C of their intact CDTs, yet no reflex

responses were evoked.

**Fig 2. Assessment of vibratory and thermal perception and NWR during nerve block. A-C**. Vibration

intensity ratings for 200 Hz (A) and 20 Hz (B) declined significantly during the nerve block (*200 Hz*:

baseline 9.9 (1.4), block 1.8 (4.4), recovery 9.9 (1.2), f(2) = 32.38, p < 0.0001; *20 Hz*: baseline 5.1 (1.6),

block 0.3 (1.6), recovery 5.1 (1.7), f(2) = 33.77, p < 0.0001, post hoc test: ****P < 0.0001, ns > 0.9999,

n = 22, Friedman test). Vibration discrimination (6 trials per condition) was also significantly impaired

during the block (*3AFC*: baseline 6.0 (0.0), block 3.0 (3.0), recovery 6.0 (0.0), f(2) = 40, p < 0.0001, post

hoc test: ****P < 0.0001, ns > 0.9999, n = 21, Friedman test). **D-E**. Cold detection thresholds (CDTs)

significantly changed during the block (baseline 28.8 (1.9)°C, block 21.6 (5.6)°C, recovery 28.0 (2.8)°C,

f(2) = 35.27, p < 0.0001, post hoc test: ****P < 0.0001, ns = 0.395, n = 22, Friedman test). The dotted

line at 23°C represents the lower border of normal values for innocuous cold detection, with cold pain

emerging ≤10-14°C. Warm detection thresholds (WDTs) remained unchanged during the block but

were elevated in the recovery condition compared to baseline (baseline 35.8 (3.6)°C, block 37.0

(3.3)°C, recovery 38.4 (5.5)°C, f(2) = 7.44, p = 0.024, post hoc test: *P = 0.031, ns (baseline vs block) >

0.9999, ns (baseline vs recovery) = 0.150, n = 22, Friedman test). **F.** The reflex responses were

completely abolished during the nerve block (NWR latencies: baseline 90.0 (14.0) ms, block 0.0 (0.0)

ms, recovery 92.5 (13.0) ms, p = 0.052, U = 4113, n = 198, Mann Whitney test).

## Reduced pain during preferential Aβ-fiber block

During the nerve block, pain ratings at the pre-block $PAIN^{th}$ current intensity dropped significantly,

resulting in the complete abolition of pain in 14 out of 22 participants (Fig 3A). Reaction time

measurements at the pre-block $PAIN^{th}$ current intensity were significantly delayed (baseline 258.8 ms,

block 426.2 ms, n=6), suggesting perception mediated via slower-conducting first-order afferents (Fig

S3A). In four participants, pain ratings at $PAIN^{th}$ increased during the nerve block, an effect unrelated

to the block duration (Fig S3B). Further, these four participants were not different from the other

participants when comparing reflex latencies, pain ratings at $PAIN^{th}$ or $NWR^{th}$, vibration sensibility, or

temperature thresholds (Fig S3C-H).

The most frequently chosen descriptors for characterizing pain quality at $PAIN^{th}$ were "sharp" and

"stabbing" (Fig 3C). At $NWR^{th}$, "shooting" and "hot-burning" were also frequently chosen (Fig 3D). The

proportion of descriptor intensity ranked as moderate and severe increased with increasing stimulus

intensity from $PAIN^{th}$ to $NWR^{th}$ (mild-moderate-severe: 74-21-5% *to* 53-39-8%, respectively; Fig 3C-D).

During the block, while the NWR was abolished (Fig 2F), pain ratings at the pre-block $NWR^{th}$ did not

differ from baseline levels. However, the overall occurrence of descriptors (and their corresponding

intensity) reduced by 71% and 66% at $PAIN^{th}$ and $NWR^{th}$, respectively (Fig 3E-F).

In the control experiment, pain ratings at $PAIN^{th}$ and $NWR^{th}$, as well as NWR latencies, remained

unchanged when the nerve block was applied to the contralateral leg (Fig S4A-C).

**Fig 3. Effect of nerve block on pain intensity and quality. A.** Reduction in pain ratings at $PAIN^{th}$ during

nerve block**.** Pain ratings at the pre-block $PAIN^{th}$ current intensity were significantly reduced during

the block (baseline 0.9 (0.7), block 0.0 (0.7), recovery 1.0 (0.9), f(2) = 11.55, p = 0.003, post hoc test:

*P = 0.013, **P = 0.008, ns > 0.999,  n = 22, Friedman test). Pain was completely abolished in 14

participants, greatly reduced in another 4, and increased in the remaining 4 (highlighted in red). **B.**

Pain ratings at the pre-block $NWR^{th}$ current intensity did not significantly change across conditions

(baseline 2.8 (3.1), block 1.8 (3.2), recovery 3.0 (2.2), f(2) = 6.181, p = 0.0331, post hoc test: ns

(baseline vs block and baseline vs recovery) = 0.071, ns (baseline vs block) > 0.9999, n = 22, Friedman

test). The 4 participants who showed an increase in $PAIN^{th}$ during nerve block in (a) are highlighted in

red. **C-D**. Pain qualities at the pre-block $PAIN^{th}$ current intensity are shown on the left, while pain

qualities at the pre-block $NWR^{th}$ current intensity are shown on the right. **E-F.** Pain qualities at block

$PAIN^{th}$ current intensity are shown on the left and pain qualities at block $NWR^{th}$ current intensity are

shown on the right.  On each occasion, participants chose any number of descriptors and ranked their

intensity as mild, moderate, or severe. Thus, the maximum number of "events" for each descriptor

equals the number of participants (n=22). The y-axis shows how many times a descriptor was chosen,

and the x-axis shows the complete list of descriptors from the McGill short-form questionnaire.

# Discussion

A preferential Aβ-fiber block significantly reduced pain and completely abolished NWR responses. The involvement of specific classes within the A-fiber population remains to be delineated. However, the abolition of $NWR^{th}$ responses during the block, despite using suprathreshold intensities in a condition in which cold perception, while reduced, was still detectable, invites speculation that Aβ nociceptors might be involved. Likewise, the reduction in $PAIN^{th}$ ratings during the block aligns with previous findings of reduced mechanical pain perception in patients with selective Aβ deafferentation and normal mechanical pain perception in patients with selective small-fiber deafferentation [23, 40].

The recently discovered Aβ nociceptors in human skin are particularly well-suited to signal percepts and responses requiring rapid transmission of nociceptive information from the periphery [23, 25, 41]. In microneurography, intraneural stimulation of Aβ nociceptors produces painful percepts at the same current intensities where intraneural stimulation of Aβ tactile afferents produces nonpainful percepts [23].

In the literature, the short-latency reflex component is considered non-nociceptive with the fast-conducting presumed "tactile" inputs thought to serve a role in posture correction or the inhibition of the late reflex response [10, 13, 20]. In the current study using intradermal electrical stimulation, we found no instances of dual NWR responses (e.g. two reflex responses within the same reflex recording time window). The NWR responses in our data had latencies ranging from 65 to 137 ms (mean latency: 91 ms), corresponding to a potential mix of RII and RIII latencies; however, they were rated as equally painful regardless of latency. Further, when examining the data from all individual reflex trials, only 9 out of 198 reflex responses (4.5%) were perceived as non-painful. The non-painful reflexes did

not have the shortest latencies and the absence of pain could perhaps be a momentary shift in the

attention of the participants, as they never rated two consecutive reflex responses as non-painful.

In the current study, we used intradermal stimulation (needle electrodes), whereas the conventional

approach is to use surface electrodes. In a previous study where surface electrical stimulation was

used, 12.4% of the NWR responses had a dual component. Further, 14.2% of the NWR responses

were rated as non-painful, and those had latencies ranging from 64 to 140 ms, with a median latency

at 83.0 ms [22]. While a study using both needle and surface electrodes noted the absence of the

early component of the reflex response (RII) during surface stimulation [11], another study found no

difference between the two methods [42]. Intradermal electrodes, as used in the current study, are

likely to stimulate the terminal branches of cutaneous afferents rather than the nerve endings

themselves, resulting in a less synchronized afferent volley, perhaps resembling a more natural

stimulus compared to surface electrodes [20]. The targeted nerves (sural or tibial) or stimulation

paradigms (duration and number of pulses) could be other factors determining if dual NWR responses

are observed or not.

We used short-duration electrical pulses (0.2 ms) to preferentially activate larger-diameter fibers,

based on the strength-duration relationship for electric excitation of myelinated axons [43-45]. In a

few participants, the reflex was elicited using a longer pulse duration (1 ms), but this did not suggest

activation of different nerve fibers, as latencies and pain ratings were not different from reflexes

elicited at shorter pulse durations (Fig. S1A-D). Consistent with earlier studies, and our pilot

observations, using a single pulse failed to evoke NWR responses, indicating that temporal summation

of repeated stimuli is required for reflex elicitation [7, 42, 46]. Typically, this need for multiple stimuli

is overlooked when calculating conduction velocity in the afferent limb, as measurements are

routinely made from the onset of the first pulse which usually fails to evoke an NWR. As NWR

responses are typically elicited by trains of pulses separated by 2-5 ms, the conduction velocity of the

afferent fibers contributing to the response could be underestimated, leading to bias towards the

slower (Aδ) conduction range. Furthermore, the observations that NWR latencies are reduced by 3-4

ms during voluntary muscle contraction [17] and that motor neuronal response is enhanced for 50-

150 ms following low-threshold electrical stimulation [47] indicate that the motor neuronal pool is a

key determinant of the timing and amplitude of the NWR.

NWR relies on temporal summation driven by high-frequency repeated stimuli. In this context, the

high impulse rates (up to 300 Hz) produced by Aβ nociceptors [23, 48] in response to noxious stimuli

suggests that this class of afferent fiber is ideally suited for detection (and rapid relay) of information

about noxious stimuli and to contribute to the generation of NWR. Indeed, the function of pain as a

warning system necessitates the rapid transmission of information from the periphery to execute

appropriate motor responses and meet behavioral requirements.

During the block, pain ratings at the pre-block $PAIN^{th}$ current intensity were significantly reduced,

along with the frequency of pain descriptors and their corresponding intensities. The most frequently

chosen pain descriptor "*sharp*" matches the percept evoked by selective activation of single Aβ

nociceptors using low-current intraneural stimulation [23]. During the block, overall pain ratings at

the pre-block $NWR^{th}$ current intensity did not change, but the frequency of pain descriptors and their

corresponding intensities were greatly reduced. It may be that the increase in stimulus intensities

from $PAIN^{th}$ to $NWR^{th}$ led to the activation of additional afferent types (Aδ and possibly C fibers),

which might explain the persistence of pain at the pre-block $NWR^{th}$ current intensity during the block.

This shift towards reliance on small-diameter inputs is reflected in the prolonged reaction times for

electrical pain at $PAIN^{th}$ during the block (baseline 258.8 ms, block 426.2 ms), which now fall within

the same range as reaction times for cold detection – a known Aδ function [49].

During the block, while pain ratings at pre-block $PAIN^{th}$ were reduced or abolished in most

participants, in four of them, the pain was intensified. However, only the pain ratings, and not the

NWR responses, differed from the other participants (Fig S3C).

Nociceptive responses can be modulated by another nociceptive stimulus in a phenomenon known as

diffuse noxious inhibitory control (DNIC), where 'pain inhibits pain'. This is caused by inhibition in the

spinal dorsal horn by nociceptive input from an adjacent part of the body [50]. The DNIC phenomenon

has also been tested in relation to the NWR, resulting in increased NWR thresholds following

exposure to noxious cold, heat, and muscular exercise [51]. To investigate potential DNIC effects on

electrical pain and reflex, we performed control experiments with the blood pressure cuff applied to

the contralateral leg. However, we found that pain and the NWR persisted ipsilaterally, indicating that

the diminution of pain and the complete abolition of NWR were the result of a preferential loss of Aβ

inputs when the cuff was applied to the ipsilateral leg rather than a DNIC effect.

During the block, CDT was significantly affected (reduced sensitivity), and detection thresholds are

known to be variable. However, no difference was found in pain ratings at pre-block $PAIN^{th}$ between

participants with reduced and normal cold sensitivity during the block. Importantly, during the block,

despite a two-fold increase in pre-block $NWR^{th}$ and a five-fold increase in pulse duration, the reflex

could not be evoked, even though all participants could still detect cooling in the nonpainful range

(>10-14°C) mediated by Aδ fibers. Warm sensibility, a function of C fibers, remained unaffected during

the block (p > 0.999). WDT values were statistically elevated in the recovery condition compared to

baseline, which could possibly be a consequence of increased blood flow to the limb upon cuff

release, masking the participants' ability to detect warm temperatures. A longer waiting time for full

recovery (>30 min) might have eliminated that difference.

We made several important observations regarding pain and the NWR in our experimental conditions.

The NWR consisted of a single response, evoked at stimulus intensities that were always painful, with

no NWR thresholds observed below the pain threshold. During a preferential block of Aβ afferents,

pain at the perception threshold was diminished, and the NWR was abolished. These results suggest

the possible involvement of very fast-conducting afferents in pain perception and NWR signaling and

may be relevant for understanding the functions of recently discovered human Aβ nociceptors.

# Acknowledgements

We would like to thank Magnus Kronander for his helpful contributions to data compilation and

analysis.

# References

1.  Khera T, Rangasamy V. Cognition and Pain: A Review. Front Psychol. 2021;12:673962. Epub 20210521.

doi: 10.3389/fpsyg.2021.673962. PubMed PMID: 34093370; PubMed Central PMCID:

PMCPMC8175647.

2.  Rischer KM, Gonzalez-Roldan AM, Montoya P, Gigl S, Anton F, van der Meulen M. Distraction from

pain: The role of selective attention and pain catastrophizing. Eur J Pain. 2020;24(10):1880-91. Epub

20200813. doi: 10.1002/ejp.1634. PubMed PMID: 32677265; PubMed Central PMCID:

PMCPMC7689692.

3.  Fillingim RB. Individual differences in pain: understanding the mosaic that makes pain personal. Pain.

2017;158 Suppl 1(Suppl 1):S11-S8. doi: 10.1097/j.pain.0000000000000775. PubMed PMID: 27902569;

PubMed Central PMCID: PMCPMC5350021.

4.  Willer JC. Anticipation of pain-produced stress: electrophysiological study in man. Physiol Behav.

1980;25(1):49-51. doi: 10.1016/0031-9384(80)90181-x. PubMed PMID: 7413819.

5.  Sherrington CS. Flexion-reflex of the limb, crossed extension-reflex, and reflex stepping and standing. J

Physiol. 1910;40(1-2):28-121. doi: 10.1113/jphysiol.1910.sp001362. PubMed PMID: 16993027;

PubMed Central PMCID: PMCPMC1533734.

6.  Carstens E, Hartung M, Stelzer B, Zimmermann M. Suppression of a hind limb flexion withdrawal reflex

by microinjection of glutamate or morphine into the periaqueductal gray in the rat. Pain.

1990;43(1):105-12. doi: 10.1016/0304-3959(90)90055-I. PubMed PMID: 1980535.

7.  Sandrini G, Serrao M, Rossi P, Romaniello A, Cruccu G, Willer JC. The lower limb flexion reflex in

humans. Prog Neurobiol. 2005;77(6):353-95. Epub 20051228. doi: 10.1016/j.pneurobio.2005.11.003.

PubMed PMID: 16386347.

8.  Rhudy JL, Williams AE, McCabe KM, Nguyen MA, Rambo P. Affective modulation of nociception at

spinal and supraspinal levels. Psychophysiology. 2005;42(5):579-87. doi: 10.1111/j.1469-

8986.2005.00313.x. PubMed PMID: 16176380.

9.    Andersen OK. Studies of the organization of the human nociceptive withdrawal reflex. Focus on

sensory convergence and stimulation site dependency. Acta Physiol (Oxf). 2007;189 Suppl 654:1-35.

doi: 10.1111/j.1748-1716.2007.01706.x. PubMed PMID: 17439638.

10.    Hugon M. Exteroceptive Reflexes to Stimulation of the Sural Nerve in Normal Man. In: Desmedt JE,

editor. Human Reflexes, Pathophysiology of Motor Systems, Methodology of Human Reflexes. 3.

Brussels: Karger; 1973. p. 713-29.

11.    de Willer JC. Comparative study of perceived pain and nociceptive flexion reflex in man. Pain.

1977;3(1):69-80. doi: 10.1016/0304-3959(77)90036-7. PubMed PMID: 876668.

12.    Dowman R. Spinal and supraspinal correlates of nociception in man. Pain. 1991;45(3):269-81. doi:

10.1016/0304-3959(91)90051-X. PubMed PMID: 1876436.

13.    Ellrich J, Treede RD. Convergence of nociceptive and non-nociceptive inputs onto spinal reflex

pathways to the tibialis anterior muscle in humans. Acta Physiol Scand. 1998;163(4):391-401. doi:

10.1046/j.1365-201X.1998.t01-1-00392.x. PubMed PMID: 9789583.

14.    Popescu A, LeResche L, Truelove EL, Drangsholt MT. Gender differences in pain modulation by diffuse

noxious inhibitory controls: a systematic review. Pain. 2010;150(2):309-18. Epub 20100616. doi:

10.1016/j.pain.2010.05.013. PubMed PMID: 20557999.

15.    Ruscheweyh R, Kreusch A, Albers C, Sommer J, Marziniak M. The effect of distraction strategies on pain

perception and the nociceptive flexor reflex (RIII reflex). Pain. 2011;152(11):2662-71. Epub 20110916.

doi: 10.1016/j.pain.2011.08.016. PubMed PMID: 21925793.

16.    Ydrefors J, Karlsson T, Wentzel Olausson U, Ghafouri B, Johansson AC, Olausson H, et al. Automated

Nociceptive Withdrawal Reflex Measurements Reveal Normal Reflex Thresholds and Augmented Pain

Ratings in Patients with Fibromyalgia. J Clin Med. 2020;9(6). Epub 20200625. doi: 10.3390/jcm9061992.

PubMed PMID: 32630430; PubMed Central PMCID: PMCPMC7356211.

17.   Hagbarth KE. Spinal withdrawal reflexes in the human lower limbs. J Neurol Neurosurg Psychiatry.

1960;23(3):222-7. doi: 10.1136/jnnp.23.3.222. PubMed PMID: 14398771; PubMed Central PMCID:

PMCPMC497412.

18.   Grimby L. Normal plantar response: integration of flexor and extensor reflex components. J Neurol

Neurosurg Psychiatry. 1963;26(1):39-50. doi: 10.1136/jnnp.26.1.39. PubMed PMID: 13950500;

PubMed Central PMCID: PMCPMC495534.

19.   Lloyd DPC. Neuron Patterns Controlling Transmisson of Ipsilateral Hind Limb Reflexes in Cat. Journal of

Neurophysiology. 1943;6(4):293-315. doi: 10.1152/jn.1943.6.4.293.

20.   Shahani BT, Young RR. Human flexor reflexes. J Neurol Neurosurg Psychiatry. 1971;34(5):616-27. doi:

10.1136/jnnp.34.5.616. PubMed PMID: 5122389; PubMed Central PMCID: PMCPMC493877.

21.   Danziger N, Fournier E, Bouhassira D, Michaud D, De Broucker T, Santarcangelo E, et al. Different

strategies of modulation can be operative during hypnotic analgesia: a neurophysiological study. Pain.

1998;75(1):85-92. doi: 10.1016/S0304-3959(97)00208-X. PubMed PMID: 9539677.

22.   Thorell O, Ydrefors J, Svantesson M, Gerdle B, Olausson H, Mahns DA, Nagi SS. Investigations into an

overlooked early component of painful nociceptive withdrawal reflex responses in humans. Frontiers in

Pain Research. 2023;3. doi: 10.3389/fpain.2022.1112614.

23.   Nagi SS, Marshall AG, Makdani A, Jarocka E, Liljencrantz J, Ridderstrom M, et al. An ultrafast system for

signaling mechanical pain in human skin. Sci Adv. 2019;5(7):eaaw1297. Epub 20190703. doi:

10.1126/sciadv.aaw1297. PubMed PMID: 31281886; PubMed Central PMCID: PMCPMC6609212.

24.   Yu H, Usoskin D, Nagi SS, Hu Y, Kupari J, Bouchatta O, et al. Single-Soma Deep RNA sequencing of

Human DRG Neurons Reveals Novel Molecular and Cellular Mechanisms Underlying Somatosensation.

bioRxiv. 2023:2023.03.17.533207. doi: 10.1101/2023.03.17.533207.

25.   Ng KKW, Lafee O, Bouchatta O, Makdani AD, Marshall AG, Olausson H, et al. Human Foot Outperforms

the Hand in Mechanical Pain Discrimination. eNeuro. 2024;11(2). Epub 20240215. doi:

10.1523/ENEURO.0412-23.2024. PubMed PMID: 38272674; PubMed Central PMCID:

PMCPMC10875634.

26.  Djouhri L, Lawson SN. Abeta-fiber nociceptive primary afferent neurons: a review of incidence and

properties in relation to other afferent A-fiber neurons in mammals. Brain Res Brain Res Rev.

2004;46(2):131-45. doi: 10.1016/j.brainresrev.2004.07.015. PubMed PMID: 15464202.

27.  Mackenzie RA, Burke D, Skuse NF, Lethlean AK. Fibre function and perception during cutaneous nerve

block. J Neurol Neurosurg Psychiatry. 1975;38(9):865-73. doi: 10.1136/jnnp.38.9.865. PubMed PMID:

1185225; PubMed Central PMCID: PMCPMC492115.

28.  Torebjork HE, Hallin RG. Perceptual changes accompanying controlled preferential blocking of A and C

fibre responses in intact human skin nerves. Exp Brain Res. 1973;16(3):321-32. doi:

10.1007/BF00233334. PubMed PMID: 4686614.

29.  Nagi SS, Rubin TK, Chelvanayagam DK, Macefield VG, Mahns DA. Allodynia mediated by C-tactile

afferents in human hairy skin. J Physiol. 2011;589(Pt 16):4065-75. Epub 20110704. doi:

10.1113/jphysiol.2011.211326. PubMed PMID: 21727219; PubMed Central PMCID: PMCPMC3180003.

30.  Melzack R. The short-form McGill Pain Questionnaire. Pain. 1987;30(2):191-7. doi: 10.1016/0304-

3959(87)91074-8. PubMed PMID: 3670870.

31.  Dowman R. Possible startle response contamination of the spinal nociceptive withdrawal reflex. Pain.

1992;49(2):187-97. doi: 10.1016/0304-3959(92)90142-X. PubMed PMID: 1608645.

32.  Laursen RJ, Graven-Nielsen T, Jensen TS, Arendt-Nielsen L. The effect of compression and regional

anaesthetic block on referred pain intensity in humans. Pain. 1999;80(1-2):257-63. doi: 10.1016/s0304-

3959(98)00214-0. PubMed PMID: 10204738.

33.  Mahns DA, Nagi SS. An investigation into the peripheral substrates involved in the tactile modulation of

cutaneous pain with emphasis on the C-tactile fibres. Exp Brain Res. 2013;227(4):457-65. Epub

20130421. doi: 10.1007/s00221-013-3521-5. PubMed PMID: 23604625.

34.    Gasser HS, Erlanger J. THE RÔLE OF FIBER SIZE IN THE ESTABLISHMENT OF A NERVE BLOCK BY

519           PRESSURE OR COCAINE. American Journal of Physiology-Legacy Content. 1929;88(4):581-91. doi:

520           10.1152/ajplegacy.1929.88.4.581.

521    35.    Sharma JP, Salhotra R. Tourniquets in orthopedic surgery. Indian J Orthop. 2012;46(4):377-83. doi:

522           10.4103/0019-5413.98824. PubMed PMID: 22912509; PubMed Central PMCID: PMCPMC3421924.

523    36.    Hilz MJ, Stemper B, Axelrod FB, Kolodny EH, Neundorfer B. Quantitative thermal perception testing in

524           adults. J Clin Neurophysiol. 1999;16(5):462-71. doi: 10.1097/00004691-199909000-00008. PubMed

525           PMID: 10576229.

526    37.    Sandrini G, Arrigo A, Bono G, Nappi G. The nociceptive flexion reflex as a tool for exploring pain control

527           systems in headache and other pain syndromes. Cephalalgia. 1993;13(1):21-7. doi: 10.1046/j.1468-

528           2982.1993.1301021.x. PubMed PMID: 8448783.

529    38.    Magerl W, Krumova EK, Baron R, Tolle T, Treede RD, Maier C. Reference data for quantitative sensory

530           testing (QST): refined stratification for age and a novel method for statistical comparison of group data.

531           Pain. 2010;151(3):598-605. Epub 20101020. doi: 10.1016/j.pain.2010.07.026. PubMed PMID:

532           20965658.

533    39.    Rolke R, Baron R, Maier C, Tolle TR, Treede DR, Beyer A, et al. Quantitative sensory testing in the

534           German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values.

535           Pain. 2006;123(3):231-43. Epub 20060511. doi: 10.1016/j.pain.2006.01.041. PubMed PMID: 16697110.

536    40.    Ridderström M, Svantesson M, Thorell O, Magounakis T, Minde J, Olausson H, Nagi SS. High prevalence

537           of carpal tunnel syndrome in individuals with rare nerve growth factor-beta mutation. Brain Commun.

538           2020;2(2):fcaa085. Epub 20200625. doi: 10.1093/braincomms/fcaa085. PubMed PMID: 32954334;

539           PubMed Central PMCID: PMCPMC7472894.

540    41.    Yu H, Usoskin D, Nagi SS, Hu Y, Kupari J, Bouchatta O, et al. Single-Soma Deep RNA sequencing of

541           Human DRG Neurons Reveals Novel Molecular and Cellular Mechanisms Underlying Somatosensation.

bioRxiv. 2023. Epub 20230928. doi: 10.1101/2023.03.17.533207. PubMed PMID: 36993480; PubMed

Central PMCID: PMCPMC10055202.

42.   Meinck HM, Piesiur-Strehlow B, Koehler W. Some principles of flexor reflex generation in human leg

muscles. Electroencephalogr Clin Neurophysiol. 1981;52(2):140-50. doi: 10.1016/0013-4694(81)90161-

9. PubMed PMID: 6167423.

43.   Hill AV. The Strength-Duration Relation for Electric Excitation of Medullated Nerve. Proceedings of the

Royal Society of London Series B, Biological Sciences. 1936;119(815):440-53.

44.   Wesselink WA, Holsheimer J, Boom HB. A model of the electrical behaviour of myelinated sensory

nerve fibres based on human data. Med Biol Eng Comput. 1999;37(2):228-35. doi:

10.1007/bf02513291. PubMed PMID: 10396827.

45.   Szlavik RB, de Bruin H. The effect of stimulus current pulse width on nerve fiber size recruitment

patterns. Med Eng Phys. 1999;21(6-7):507-15. doi: 10.1016/s1350-4533(99)00074-0. PubMed PMID:

10624746.

46.   Tøorring J, Pedersen E, Klemar B. Standardisation of the electrical elicitation of the human flexor reflex.

J Neurol Neurosurg Psychiatry. 1981;44(2):129-32. doi: 10.1136/jnnp.44.2.129. PubMed PMID:

7217968; PubMed Central PMCID: PMCPMC490843.

47.   Shahani B. Flexor reflex afferent nerve fibres in man. J Neurol Neurosurg Psychiatry. 1970;33(6):786-91.

doi: 10.1136/jnnp.33.6.786. PubMed PMID: 5531898; PubMed Central PMCID: PMCPMC493593.

48.   Bouchatta O, Brodzki M, Manouze H, Carballo GB, Kindström E, de-Faria FM, et al. PIEZO2-dependent

rapid pain system in humans and mice. bioRxiv. 2023:2023.12.01.569650. doi:

10.1101/2023.12.01.569650.

49.   Yarnitsky D, Ochoa JL. Warm and cold specific somatosensory systems. Psychophysical thresholds,

reaction times and peripheral conduction velocities. Brain. 1991;114 ( Pt 4):1819-26. doi:

10.1093/brain/114.4.1819. PubMed PMID: 1884180.

50.    Le Bars D, Villanueva L, Bouhassira D, Willer JC. Diffuse noxious inhibitory controls (DNIC) in animals

567           and in man. Patol Fiziol Eksp Ter. 1992;(4):55-65. PubMed PMID: 1303506.

568    51.    Willer JC, Roby A, Le Bars D. Psychophysical and electrophysiological approaches to the pain-relieving

569           effects of heterotopic nociceptive stimuli. Brain. 1984;107 ( Pt 4):1095-112. doi:

570           10.1093/brain/107.4.1095. PubMed PMID: 6509310.

**A.**

Stimulus onset

1. DET$^{th}$ (EMG)

2. PAIN$^{th}$ (response meter)

3. NWR$^{th}$ (EMG)

NWR response

**B.**

**C.**

Total NWR = 198

**A.**

**B.**

**C.**

**D.**

**E.**

**F.**