Deep Whole-Genome Resequencing of Marek's Disease Virus Strains Reveals Intrastrain Variation

Alejandro Ortigas-Vásquez[1], Utsav Pandey[2], Daniel Renner[1], Chris Bowen[1], Sue Baigent[3], John Dunn[4], Hans Cheng[5], Yongxiu Yao[3], Andrew F. Read[1], Venugopal Nair[3], Dave A. Kennedy[1], Moriah L. Szpara[1,2]*

[1]Departments of Biology and [2]Biochemistry and Molecular Biology, Center for Infectious Disease Dynamics and the Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, Pennsylvania 16802, USA

[3] Avian Oncogenic Virus Group, The Pirbright Institute, Ash Road, Pirbright, Woking GU24 0NF, United Kingdom

[4] United States Department of Agriculture, Agricultural Research Service, US National Poultry Research Center, Southeast Poultry Research Laboratory, Athens, Georgia 30605, USA

[5] United States Department of Agriculture, Agricultural Research Service, US National Poultry Research Center, Avian Disease and Oncology Laboratory, East Lansing, Michigan 48823, USA

*Corresponding Author
Dr. Moriah L. Szpara, email: moriah@psu.edu

CRediT (Contributor Roles Taxonomy; https://credit.niso.org/contributor-roles-defined/)
Conceptualization – AOV, MLS, DAK
Data curation – UP, DWR, AOV
Formal analysis – AOV
Funding acquisition – DAK, VN, MLS, AFR
Investigation – UP, CDB, SB, YY, JD, HC
Methodology – UP, MJJ
Project administration – MLS, DAK
Resources – JD, HC, SB, YY, VN, DWR, AFR
Software – UP, DWR, AOV
Supervision – MLS, DAK
Validation – CDB, DWR, AOV
Visualization – AOV, MLS, DAK
Writing – AOV
Writing – review & editing – AOV, MLS, DAK, HC, JD

Abstract

Current strategies to understand the molecular basis of Marek's disease virus (MDV) virulence

primarily consist of cataloguing divergent nucleotides between strains with different phenotypes.

However, each MDV strain is typically represented by a single low-coverage consensus genome

despite the confirmed existence of mixed viral populations. To assess the reliability of

consensus-only interstrain comparisons, we obtained two additional consensus genomes of

vaccine strain CVI988 (Rispens) and two additional consensus genomes of the very virulent

strain Md5 by sequencing viral stocks and cultured field isolates. In conjunction with the

published genomes of CVI988 and Md5, this allowed us to perform 3-way comparisons between

consensus genomes of the same strain. We found that consensus genomes of CVI988 can vary in

as many as 240 positions involving 12 open reading frames (ORFs). We found that Md5

genomes varied only in 12 intergenic positions, with no variation within ORFs. Phylogenomic

analyses showed all three Md5 consensus genomes clustered closely together, while also

showing that CVI988-GenBank diverged from CVI988-UK and CVI988-USDA. Comparison of

CVI988 consensus genomes with those of seven closely related isolates from China revealed 19

SNPs that appear to be unique to CVI988-GenBank. Finally, we evaluated the genomic

homogeneity of CVI988 and Md5 populations by identifying positions with >2% read support

for alternative alleles in two ultra-deeply sequenced samples. We were able to confirm that both

populations of CVI988 and Md5 were mixed, exhibiting a total of 30 and 27 minor variant

positions, respectively. We found no evidence of minor variants in the positions corresponding to

the 19 unique SNPs in CVI988-GenBank. Taken together, our findings challenge prior claims of

natural recombination between CVI988 and virulent strains and confirm the need to resequence

MDV strains in order to ensure the accuracy of interstrain comparisons.

Keywords: Intrastrain; Diversity; Population; Genome; MDV; CVI988; Md5; Variation

Introduction

Marek's disease virus (MDV) is an oncogenic, lymphoproliferative and neuropathic

alphaherpesvirus affecting chickens (*Gallid herpesvirus 2*, Genus *Mardivirus*; Family

Herpesviridae). It was first reported in 1907 by Hungarian veterinarian József Marek (1).

Successful isolation of MDV in culture was only realized six decades later, closely followed by

the development of the very first vaccine, HPRS-16/Att (2–4). HPRS-16/Att was soon replaced

by a better replicating, equally protective vaccine derived from strain FC126 of turkey

herpesvirus (5). This vaccine, known as HVT, would go on to become the first widely used

commercial vaccine for Marek's disease (MD) (6). HVT was successful in reducing MD

incidence in commercial flocks until the late 1970s, when reports of strains with increased

virulence that were able to bypass vaccine-induced immunity started to emerge across the USA

(7, 8). A bivalent vaccine consisting of HVT alongside MDV serotype 2 strain SB-1 was

introduced in response. However, a similar shift in virulence occurred in just over a decade (9,

10). Outside the USA, a vaccine known as CVI988 had been in use since the mid 1970s (11).

Developed in the Netherlands by a group of veterinary scientists under Dr. Bart Rispens, CVI988

offered the best protection against the emerging variants, and it quickly became the "gold

standard" vaccine around the world (12). CVI988 is colloquially known today as the "Rispens"

vaccine for MDV. Though CVI988 has mostly kept MD losses at bay for the past 30 years, there

are now reports of strains able to bypass CVI988-conferred immunity, and the most recent

estimate places MDV-associated losses in the poultry industry at around US $1-2 billion per

annum (13–16).

Comparative genomics has been a driving force in efforts to understand the molecular basis of

MDV-induced virulence ever since the first viral genomes were sequenced over two decades ago

(17, 18). The current strategy involves cataloging variable nucleotide positions between the

consensus genomes of different MDV strains. Comparisons are made between strains grouped on

the basis of their ability to break through vaccine protection, or "pathotype," with the hope that

this will help to identify the genes and polymorphisms associated with virulence (19, 20).

However, despite MDV strains being known to exist as mixed populations, the degree to which

the published consensus genomes represent the average for each strain has never been assessed

(21–23). In addition, all available consensus genomes for MDV are low-coverage genomes that

only account for the major variants (≥50%) present in the original viral population from which

they were sequenced. The lack of information on minor variants (≤50%) can be problematic, due

to previous reports suggesting that the diversity within a given viral population contributes to the

overall pathotype of a strain (24–27). Although the extent of this contribution is still unknown,

this nonetheless suggests that failing to account for minor variants can negatively impact our

ability to correlate sequencing data to pathotype. Finally, many of the available reference

genomes were not sequenced directly from viral stocks (i.e., a virus population grown in host

cells). This includes the only available CVI988 consensus genome, which was sequenced from a

bacterial artificial chromosome (BAC) clone, and the Md5 consensus genome, which was

obtained from a set of five overlapping cosmid clones (28). By randomly selecting one viral

haplotype from a population, these methods inherently reduce the chances of obtaining a

representative consensus genome. In addition, the use of BAC and cosmid approaches generates

an artificially homogenous population of viral genomes, further moving away from the real-

world biology of herpesvirus strains.

To address these issues, it is necessary to test the reliability of available MDV consensus

genomes and assess the genomic homogeneity of MDV populations using next-generation

sequencing (NGS) approaches. Whole-genome resequencing (WGR) is a simple yet powerful

approach that consists of comparing newly sequenced individuals against a reference (29). Here,

we have adapted this strategy for viruses, by sequencing multiple populations of the same strain

and determining whether these yield a consistent consensus genome, under the assumption that

large shifts in genomic variant frequencies will manifest as consensus-level differences. We

independently sequenced two populations of the vaccine strain CVI988 and two populations of

the very virulent prototype strain Md5, from a combination of viral stocks and cultured field

isolates. The newly-sequenced genomes were compared with previously published consensus

genomes using whole-genome pairwise alignments. Using this approach, we were able to detect

variable positions between consensus genomes of each strain. We also evaluated the individual

contribution of each consensus genome to overall viral diversity using phylogenomic

approaches. We found 19 single nucleotide polymorphisms (SNPs) that were only present in

CVI988-GenBank. Next, we assessed the genomic homogeneity of CVI988 and Md5 by

analyzing ultra-deep Illumina sequencing data (>1,000x depth) and identified positions where at

least 2% of the reads supported an alternative allele (30). This method enabled detection of

minor variants and allowed us to confirm that the ultra-deep sequenced populations of CVI988

and Md5 were mixed. These findings challenge prior claims of natural recombination between

CVI988 and virulent strains, and highlight the need to resequence MDV strains in order to

accurately link viral genotypes to virulence phenotypes.

Methods

Viral culture and DNA isolation of USDA CVI988 and Md5

Chicken embryo fibroblasts (CEFs) were used to culture viruses. Cultures were maintained in 1:1

mixture of Leibovitz's L-15 and McCoy's 5A (LM) media supplemented with fetal bovine serum

(FBS), 200 U/ ml penicillin, 20 µg/ml streptomycin, and 2 µg/ml amphotericin B in a 37 °C, 5%

CO_2 incubator. Cells were plated with 4% FBS LM media and maintained in 1% FBS LM media.

For storage as viral stocks, infected cells were suspended in freezing media composed of 10%

DMSO, 45% FBS, and 45% LM media and kept in liquid nitrogen.

DNA library preparation and sequencing of USDA CVI988 and Md5

Total DNA isolations for each MDV culture were quantified for total DNA by Qubit (Invitrogen)

and either viral or vaccine DNA present by a previously described qPCR assay (30). Samples

were then acoustically sheared using Covaris and high-throughput, deep sequencing DNA library

preparations were produced using the Kapa HyperPrep Library Kit (Roche) as per

manufacturer's protocol, with 14 PCR cycles. Then, samples were processed through a

previously described oligo-enrichment procedure, where MDV-specific probes (31) (myBaits,

Arbor Biosciences) were used with the myBaits Target Capture Kit (Arbor Biosciences) to

enrich for MDV DNA (32). An additional PCR amplification (14 cycles) yielded libraries which

were assessed for quality by Qubit, as well as a KAPA-specific qPCR, which allowed for

accurate dilutions of libraries to 4pM before sequencing. Finally, libraries were sequenced using

v3 chemistry 300bp × 300bp paired-end sequencing on an Illumina Miseq, as per manufacturer's

instruction.

DNA library preparation and sequencing of UK CVI988 and Md5

Total DNA isolations for each MDV culture were quantified for total DNA by Qubit (Invitrogen)

and either viral or vaccine DNA present by a previously described qPCR assay (31). Samples

were then acoustically sheared using Covaris and high-throughput, deep sequencing DNA library

preparations were produced using the Kapa HyperPrep Library Kit (Roche) as per

manufacturer's protocol, with 14 PCR cycles. Then, samples were processed through a

previously described oligo-enrichment procedure, where MDV-specific probes (32) (myBaits,

Arbor Biosciences) were used with the myBaits Target Capture Kit (Arbor Biosciences) to

enrich for MDV DNA (33). An additional PCR amplification (14 cycles) yielded libraries which

were assessed for quality by Qubit, as well as a KAPA-specific qPCR, which allowed for

accurate dilutions of libraries to 4 pM before sequencing. Finally, libraries were sequenced using

v3 chemistry 300 bp × 300 bp paired-end sequencing on an Illumina MiSeq, as per the

manufacturer's instruction.

Processing of sequencing reads and genome assembly

Viral reads were identified by aligning raw reads in FASTQ files against a local database of 97

MDV genomes using BLASTN with an E-value cutoff of 1×10^{-2}. The selected reads were then

subjected to the quality control and preprocessing step (Step 1) of our published viral genome

assembly (VirGA) workflow (34). These properly paired reads were then used for *de novo*

assembly using metaSPAdes v3.14.0 (35). The resulting file containing the metaSPAdes

scaffolds in FASTA format served as input for the remaining steps of VirGA (Steps 3-4), which

include genome linearization, annotation and post-assembly quality assessments. For the

reference-guided contig-ordering step, we used a trimmed version (TRL and TRS regions

removed) of the published consensus genome of each strain (**Table 1**). New consensus genomes

were also trimmed for downstream analyses. Tandem repeats were identified in trimmed

consensus genomes using Tandem Repeats Finder (TRF) v4.09 (36). Manual verification and

masking of tandem repeats was conducted using Geneious Prime v2023.0.4.

Pairwise identity and genetic distance comparisons

Trimmed consensus genomes with masked tandem repeats were aligned using MAFFT v7.505

with default settings (37). The resulting alignment file in clustal format was imported into

Geneious to visualize variable positions (SNPs and Indels) between the genomes. Published

MDV consensus genomes with PubMed Identifiers (n = 29) and all consensus genomes included

in the present study (n = 6) were aligned using MAFFT (see **Supplementary Table 5** for

details). A neighbor-joining (NJ) tree was constructed from the multi-genome alignment file in

clustal format using the Geneious Tree Builder tool with a Tamura-Nei genetic distance model

and no outgroup.

Identification of minor variant positions

191   A custom script was used to identify positions with at least 2% read support for an alternative

192   allele – i.e., a minor variant – in genomes with >1,000x coverage. Positions in tandem repeats

193   and homopolymers were excluded. All remaining minor variant positions were verified through

194   manual inspection of BAM alignment files using the Integrative Genomics Viewer (IGV)

195   v2.12.3 (38).

Results

Resequencing viral stocks and field isolates of CVI988 and Md5 using Illumina whole-genome sequencing

For this study, we obtained complete genome sequences for two populations each of the vaccine strain CVI988 and the virulent strain Md5 (vv pathotype). For Md5, both the USDA and UK viral populations were derived from standard lab stocks (i.e., a virus population initiated from a prior viral stock and expanded on host cells). The UK CVI988 was likewise derived from lab stock. In contrast, the USDA Rispens was isolated from sample collected at a Pennsylvania farm, before isolation and expansion on host cells in the lab. All viral stocks were propagated similarly for viral DNA isolation and sequencing library preparation (see Methods for details). The UK viral genomes were additionally enriched using published methods for oligonucleotide-bait based enrichment in solution (32), which can enable a greater depth of coverage by enriching viral sequences from the overall milieu of DNA. Viral consensus genomes for each stock were determined using a previously published combination of *de novo* and reference-guided assembly methods (**Table 1**; see Methods for details).

The MDV genome has a double-stranded DNA (dsDNA) structure, with a full length of approximately 175-180 kb. The genomic layout is split into two large unique regions known as Unique Long (UL) and Unique Short (US), each flanked by inverted repeat sequences (IRL, IRS) and terminal repeat sequences (TRL, TRS) (**Figure 1A**, center). The α-type packaging sequences are located at the genomic termini and at the IRL/IRS junction. None of the new genomes contained any truncated or missing proteins. The average G+C content of all genomes was 44.0%. The coverage depth of Illumina sequencing data exceeded 100X for all viral genomes,

and it reached an average of 1,000X for the two genomes subjected to oligonucleotide-bait based

enrichment (**Table 1**).

Pairwise alignment of same-strain consensus genomes reveals greater intrastrain

variation in CVI988 than Md5

For each strain, trimmed consensus genomes with masked tandem repeats were aligned and

visualized. Variable positions corresponding to SNPs, insertions and deletions (Indels) were

identified and verified through manual inspection of sequence-read alignment files (i.e., BAM

files). A total of 240 variable positions were detected in the three consensus genomes of vaccine

strain CVI988, impacting a total of 12 ORFs. These include the genes UL22 / gH (MDV034),

UL49 /VP22 (MDV062), MDV074, MDV075, Meq /RLORF7 (MDV076 / MDV005), vIL-8 /

RLORF2 (MDV078 / MDV003), MDV080, MDV085, US2 (MDV091), US3 (MDV092),

SORF4 / S4 (MDV093) and US6 /gD (MDV094) (**Figure 1A, Supplementary Table 1**). In

contrast, only 12 variable positions were detected in the three consensus genomes of very

virulent strains Md5 (**Supplementary Table 2**). No variable positions were found within ORFs

in the Md5 consensus genomes.

The *meq* oncogene contributes significantly to overall intrastrain variation in

consensus genomes of CVI988

Out of 240 variable positions detected in the CVI988 consensus genomes (**Supplementary**

Table 1), 217 were found in IRL (n = 207 bp) and IRS (n = 10 bp). A large portion of these

variable loci (n = 177 bp) corresponded to a 59 amino acid (AA) insertion in the *meq* oncogene,

which was found in the CVI988-GenBank and CVI988-USDA consensus genomes but absent in

the CVI988-UK consensus genome (**Figure 2**). For *meq*, previous studies have suggested the

existence of up to three isoforms: Meq, Long Meq (L-meq) and Short Meq (S-meq) (39–42).

Meq is the standard isoform and is 339 AA in length. L-meq is a longer version of Meq that is

characterized by a 59 AA insertion in the transactivation domain (TAD), which is often

accompanied by a 3 bp CCA deletion immediately upstream from the insertion site. S-meq is a

shorter version of Meq that exhibits a 41 AA deletion in the TAD. The CCA deletion that is

typically associated with the presence of the L-meq isoform was confirmed to exist as a

subpopulation of considerable size (~ 40%) in the UK CVI988 population, but it did not appear

to be linked to the presence of the 177 bp insertion (43). All three Md5 genomes exhibited the

standard isoform of *meq* (**Figure 2**), again indicating less diversity than found in the CVI988

genomes.

Increased intrastrain variation in structural repeat regions is associated with

greater homopolymer instability

Outside of *meq*, 37 of the 40 remaining variable positions found in structural repeat regions of

CVI988 consensus genomes were associated with differences in homopolymer lengths

(**Supplementary Table 1**). By contrast, all variable positions found in the unique regions of

CVI988 consensus genomes (n = 20) were SNPs, with an even distribution between synonymous

(n = 10) and non-synonymous (n = 10) SNPs. In Md5 consensus genomes, all of the 12 variable

positions were found in either IRL (n = 7) or IRS (n = 5) and were all associated with differences

in homopolymer lengths (**Supplementary Table 2**).

264 **Intrastrain variation in unique regions is associated with 19 unique SNPs in the**

265 **published CVI988 consensus genome**

Out of the 20 SNPs in unique regions of CVI988 consensus genomes, 19 were unique to

CVI988-GenBank, including 3 SNPs in UL and 16 SNPs in US (**Figure 1B**). Phylogenomic

analysis confirmed divergence of CVI988-GenBank from CVI988-UK and CVI988-USDA

(**Figure 3**). To evaluate whether these 19 SNPs could be found in closely related strains, we

aligned the three consensus genomes of CVI988 with the consensus genomes of seven co-

clustering strains (HNLC503, HC/0803, SY/1209, Hrb/1504, DH/1307, DH/1504, CH/10; all

from China) and compared the nucleotides present at these positions. All 19 SNPs remained

exclusive to CVI988-GenBank. CVI988-UK, CVI988-USDA and all seven Chinese isolates

shared the same nucleotides in all of these positions (**Figure 1B**). These nucleotides also

matched the sequences of these 19 positions in the three Md5 genomes included in this study

(**Figure 1B**).

278 **Ultra-deep sequencing reveals presence of minor variants in CVI988 and Md5**

279 **populations**

In addition to consensus-level comparisons, we performed minor variant analysis on the deeply

sequenced UK genomes (**Figure 4**, see **Methods** for details). This included UK CVI988

(>10,000 ×) and UK Md5 (>8,500 ×; **Table 1**). A total of 30 minor variant positions were

detected in the original UK CVI988 population across a combination of coding and regulatory

regions. These minor variants impacted a total of 9 ORFs: MDV011, UL13 (MDV025), UL26

(MDV038), MDV069, MDV072, MDV077, ICP4 (MDV084), US3 (MDV092) and SORF4 / S4

(MDV093) (**Supplementary Table 3**). Similarly, a total of 27 minor variants were detected in

the UK Md5 population. These impacted the following ORFs: UL19 (MDV031), UL41 / VHS

(MDV054), UL44 gC (MDV057), UL48 / VP16 (MDV061), UL52 MDV066, MDV069 and

MDV072 (**Supplementary Table 4**). Twelve of the 14 minor variant positions found across

MDV069 and MDV072 were common to both strains. No minor variants were identified in the

US region of the UK Md5 population. Additionally, we verified that no minor variant positions

coincided with the 19 unique SNPs in CVI988-GenBank (**Figure 1B**).

Discussion

The goal of comparative genomics is to provide a link between genotype and phenotype. The

accuracy with which these two elements are represented is therefore directly proportional to our

ability to make biologically relevant conclusions through the use of comparative genomic

methods. DNA viruses have long been considered to be inherently stable, especially when

compared to RNA viruses, which are now described as mutant swarms, or "quasispecies" (44,

45). This had led to the assumption that viral consensus genomes, which represent the majority

nucleotide at any given position, can accurately represent the genotype of MDV strains such as

vaccine strain CVI988 and the very virulent strain Md5. As demonstrated here and in prior

studies, these and other herpesvirus populations can harbor minor variants both in culture and *in*

vivo (46–49). Here, by resequencing and ultra-deep sequencing viral stocks and field isolates of

these strains, we have sought to evaluate the impact of conducting interstrain comparisons

without accounting for the underlying variation in MDV strains.

CVI988 is an attenuated and non-oncogenic vaccine strain that has long been known to exist as a

mixed population due to undergoing multiple passages *in vitro*, in different laboratories and in

multiple vaccine manufacturing companies. There have also been reports suggesting that the

effectiveness of CVI988 vaccines can vary depending on the manufacturer, and that even

different batches from the same manufacturer can show differences in effectiveness (50, 51). In

contrast, Md5 is a virulent field strain of MDV that is currently used as the prototype very

virulent (vv) strain in the USDA-ARS-Avian Disease and Oncology Laboratory (ADOL)

pathotype classification system (52). As such, the fact that we found 240 variable positions

between consensus genomes of CVI988 while only finding 12 variable positions between

consensus genomes of Md5 is consistent with their history and real-world biology. Whether the

larger number of variant loci in CVI988 reflects the strain's attenuation through serial passage,

or if it is a feature of this strain's genetic background, remains to be determined. In addition, the

fact that we were able to detect 30 minor variant positions in CVI988-UK and 27 minor variant

positions in Md5-UK confirms that both strains exist as mixed viral populations. Over time

minor variant differences may expand during passage(s) in culture or during *in vivo* infections,

leading to consensus level differences (46, 53). Minor variants may thus provide an explanation

for the differences observed between consensus genomes of each strain.

Having access to the minor variant profiles of CVI988 and Md5 has the added advantage of

providing insight into the potential effects of intrastrain variation in loci of high interest, such as

pp38. Currently available qPCR assays to distinguish between CVI988 and virulent MDV strains

are based on a single nucleotide polymorphism (SNP) located at position 320 of the pp38 gene

(54). Our results show pp38 to be a highly conserved gene between different populations of

CVI988 and Md5, with no minor variant positions coinciding with this gene. This not only

confirms the robustness of qPCR assays based on SNP #320 of pp38, but also establishes pp38

as a locus with high levels of within-strain conservation. Overall, our study provides a solid

starting point for assessing intrastrain diversity in MDV strains. However, access to additional

consensus genomes of CVI988 and Md5 will still be required in order to obtain the complete

picture of MDV intrastrain diversity, as well as resequencing of other virulent and attenuated

MDV strains.

Among our most significant findings was the discovery of 19 SNPs present in the unique regions

of CVI988-GenBank but absent in the two new CVI988 consensus genomes sequenced as part of

this study. These 19 SNPs were also not present as minor variants in the CVI988-UK viral

population, where these positions were sequenced at an average depth of >10,000 ×. CVI988-

GenBank was sequenced by Spatz *et al*. in 2007 from a BAC clone of CVI988 at 6× coverage

(**Table 1**). This BAC clone was derived by Petherbridge *et al*. in 2003 from a commercial

vaccine supplied by Fort Dodge Animal Health (55). As part of the cloning process a BAC

vector was inserted into US2, and the missing region was later reconstructed using PCR and

wild-type CVI988 DNA. While explaining the origin of these 19 unique SNPs in CVI988-

GenBank is beyond the scope of our current study, our results strongly suggest that consensus

genomes of CVI988 stocks do not typically differ from virulent strains such as Md5 in any of

these positions.

The implications of these findings are multiple. Firstly, these 19 SNPs have been reported as

differences between CVI988 and virulent strains like Md5, RB-1B, GA and Md11 by many prior

studies, starting with the original publication by Spatz *et al.* (56–58). Secondly, and perhaps most

significantly, the 16 unique SNPs in US of CVI988-GenBank impacting US2, US3, SORF4 and

US6 / gD have been used as the main evidence of naturally occurring recombination between

CVI988 and virulent strains, starting in 2018 by He *et al.* (59). Multiple studies have followed,

resulting in a total of seven isolates from China being described as natural recombinants of

CVI988 on the basis of these 16 SNPs (60–62). Our two new consensus genomes of CVI988

share the same nucleotides in all 16 positions with these seven isolates, implying that they are

likely to be field isolates of CVI988 and not natural recombinants as previously described.

Notably, the US regions of CVI988-UK and DH/1504 share 100% identity, with all other

genomes sharing over 99.8% identity in US. These findings indicate that an overreliance on the

published CVI988 consensus genome has led to a historical overestimation of genomic

divergence between CVI988 and virulent strains. At the same time, these data challenge previous

notions regarding the role of recombination in MDV evolution and highlight the urgent need to

use genomic resequencing more widely in order to verify the actual instances of sequence

divergence and/or recombination.

Finally, it is worth mentioning that current Illumina-based approaches do not provide insight into

certain regions in MDV genomes where additional intrastrain variation is likely to be found.

Specifically, tandem repeats have been suggested to be an important source of diversity in

herpesviruses yet remain mostly inaccessible to NGS strategies both at the consensus and sub-

consensus levels (48, 63). The use of long read technologies (e.g., Oxford Nanopore and Pacific

Biosciences) could potentially help to overcome these limitations and make it possible to reliably

assess intrastrain variation at tandem repeats and homopolymers (64, 65). In addition, the use of

long reads could enhance our ability to link minor variant positions into proper haplotypes and

help us to better understand the real-world biology of MDV strains.

In conclusion, by resequencing viral stocks and field isolates of the vaccine strain CVI988 and

the very virulent strain Md5, we have shown that consensus genomes of these strains can differ

in multiple positions across all genomic regions. We have also established that these differences

are most likely made possible by fluctuations in a mixed population of viral genomes, the

existence of which we have demonstrated using ultra-deep Illumina sequencing. In addition, we

have shown how not accounting for underlying diversity in MDV strains has led to a historical

overestimation of divergence between CVI988 and virulent strains and provided evidence that

contradicts prior claims of naturally occurring recombination between the two. In doing so, we

have demonstrated the value of resequencing MDV strains as part of comparative genomic

studies and challenged prevailing notions regarding the magnitude and significance of diversity

in DNA viruses.

Acknowledgements

The authors thank members of the Szpara, Kennedy, and Nair labs for helpful feedback and

discussion. We thank Matthew J. Jones for contributions to DNA quantification and Andrew S.

Bell for contributions to library preparation. This work was supported by NSF-NIH EEID award

1 R01 GM140459 (DK, MS, VN).

Tables

Table 1: Sequencing statistics for new and previously published CVI988 and Md5

consensus genomes

MDV Strain	# MDV-specific reads	# Reads used for assembly	Coverage	GenBank Accession	Genome length (bp)	Prior citation
CVI988	N/A	N/A	6x	DQ530348	178,311	(28)
CVI988-USDA	227,487	138,122	149x	TBD	176,089	-
CVI988-UK	11,997,583*	9,730,216	10,055x	TBD	177,063	-
Md5	N/A	N/A	6x	NC_002229	177,874	(18)
Md5-USDA	1,030,131	727,124	793x	TBD	174,996	-
Md5-UK	10,138,498*	8,296,132	8,530x	TBD	176,079	-

399 *The UK viral stock populations were subjected to oligo-bait enrichment to increase the yield of viral
400 DNA (32, 33).
401

Figure Legends

Figure 1: Comparative analysis of CVI988 and Md5 consensus genomes

A) A total of three trimmed genomes (TRL and TRS removed) were obtained for each strain (CVI988 = Blue, Md5 = Red) and aligned using MAFFT. Variable positions including SNPs and Indels for each individual genome are highlighted (dark lines) based on their relative locations in the genome (UL = Unique Long, IRL = Internal Repeat Long, IRS = Internal Repeat Short, US = Unique Short). The total number of variable positions for each individual genome and the total for each strain are indicated on the left; these loci are listed in detail in **Supplementary Table 1**. For each strain, ORFs with variable positions in at least one of the three consensus genomes belonging to that strain are highlighted and labeled (with alternative names when possible). The total number of divergent ORFs for each strain is 12 for CVI988 and zero for Md5. **B)** A total of 19 unique SNPs (UL = 3, US = 16) were identified in CVI988-GenBank. These SNPs were not present in CVI988-UK or CVI988-USDA. They were also not present in seven closely related isolates from China (see **Figure 3**). These SNPs impacted a total of 5 ORFs, including VP22, US2, US3, SORF4 and gD. Md5-GenBank, Md5-UK and Md5-USDA shared nucleotides at these positions with all genomes except CVI988-GenBank.

Figure 2: Comparison of the *meq* oncogene in Md5 and CVI988 consensus genomes

Schematic representation of the location and structure of *meq* isoforms present at consensus level in Md5 (red) and CVI988 (blue) populations. The *meq* oncogene is comprised of the proline/glutamine (Pro/Gln) rich domain, the basic region (BR), the leucine zipper (ZIP) and the

transactivation domain (TAD) at the C-terminal region. Differences in the AA sequence of the

two strains are highlighted and labeled (A = Alanine, K = Lysine, S = Serine, E = Glutamic

acid). Meq is 339 AA in length. The L-Meq isoform contains a 59-amino acid insertion in the

transactivation domain, resulting in a total length of 398 AA.

Figure 3: Neighbor-joining tree of previously published MDV strains and the new

consensus genomes of CVI988 and Md5

A total of 29 MDV strains with published consensus genomes in GenBank (**Supplementary**

Table 5) were used to infer a neighbor-joining tree using Geneious Prime v2023.0.4. The

genomes of each strain used for our comparative analyses are highlighted alongside their

respective branches (Md5 = Red, CVI988 = Blue).

Figure 4: Minor variant distribution in CVI988 and Md5 UK populations

For each of the ultra-deep sequenced viral populations (see **Table 1**), positions where the minor

allele was present in ≥ 2% of the reads were identified and classified depending on whether they

fell on a coding or non-coding region and whether they resulted in a synonymous or non-

synonymous substitution. ORFs associated with minor variant positions of all types are labeled

and highlighted beneath the graph using the same criteria. Minor variants corresponding to

tandem repeats and homopolymers were excluded. (**A**) Minor variant distribution in the UK

CVI988 population. (**B**) Minor variant distribution in the UK Md5 population.

Figures

Figure 1: Comparative analysis of CVI988 and Md5 consensus genomes

Figure 2: Comparison of the *meq* oncogene in Md5 and CVI988 consensus genomes

Figure 3: Neighbor-joining tree of previously published MDV strains and the new

consensus genomes of CVI988 and Md5

Figure 4: Minor variant distribution in CVI988 and Md5 UK populations

Supplementary Tables

Supplementary Table 1: Variable positions in CVI988 consensus genomes

Genome	ORF	Alternative name	Position	This CVI988 genome	Other CVI988 genomes	Description
USDA	MDV034	UL22 (gH)	52,546	T	C	Synonymous substitution
GenBank	MDV062	UL49 (VP22)	111,859	C	A	Synonymous substitution
GenBank	MDV062	UL49 (VP22)	111,866	C	G	Ser155 -> Cys155
GenBank	MDV062	UL49 (VP22)	111,876	G	C	Ala152 -> Pro152
USDA	MDV074	-	127,041	C	-	Frameshift
USDA	MDV074	-	127,042	G	-	Frameshift
USDA	MDV074	-	127,043	G	-	Frameshift
USDA	MDV074	-	127,044	C	-	Frameshift
USDA	MDV074	-	127,045	T	-	Frameshift
UK	MDV075	-	128,398	-	C	Homopolymer contraction, non-coding
GenBank	MDV075	-	130,655	C	-	Homopolymer extension, non-coding
GenBank	MDV075	-	130,656	C	-	Homopolymer extension, non-coding
USDA	MDV075	-	128,207	T	-	Homopolymer extension, non-coding
USDA	MDV075	-	128,208	T	-	Homopolymer extension, non-coding
USDA	MDV075	-	128,877	-	T	Homopolymer contraction, non-coding
GenBank	MDV075	-	131,742	A	T	Met48 -> Lys48
GenBank	MDV075	-	131,766	C	-	Homopolymer extension, frameshift
GenBank	MDV075	-	131,979	A	-	Homopolymer extension, non-coding
GenBank	MDV075	-	131,980	A	-	Homopolymer extension, non-coding
UK	MDV075	-	129,719	-	A	Homopolymer contraction, non-coding
GenBank	-	-	132,535	T	-	Homopolymer extension
GenBank	-	-	132,536	T	-	Homopolymer extension
GenBank	-	-	134,710	A	-	Homopolymer extension
UK	-	-	132,665	-	C	Homopolymer contraction
USDA	-	-	133,221	G	-	Homopolymer extension
UK	-	-	133,498	-	G	Homopolymer contraction
UK	MDV076	Meq	134,320	-	G…G	Meq isoform lacking 177 bp insertion
UK	-	-	135,147	-	G	Homopolymer contraction
GenBank	-	-	137,581	G	-	Homopolymer extension
UK	-	-	135,235	T	-	Homopolymer extension
GenBank	-	-	137,670	-	TT	Homopolymer contraction
GenBank	-	-	138,647	T	-	Homopolymer extension, frameshift
GenBank	MDV078	vIL-8	140,028	A	-	Homopolymer extension, frameshift
USDA	MDV080	-	138,538	-	T	Homopolymer contraction, frameshift
GenBank	-	-	141,642	A	G	Synonymous substitution
USDA	-	-	139,196	C	T	Synonymous substitution
GenBank	MDV085	-	152,350	C	-	Homopolymer extension, frameshift
GenBank	MDV085	-	152,351	C	-	Homopolymer extension, frameshift
USDA	-	-	150,420	A	-	Homopolymer extension
USDA	-	-	150,421	A	-	Homopolymer extension

Genome	ORF	Alternative name	Position	This CVI988 genome	Other CVI988 genomes	Description
USDA	-	-	150,422	A	-	Homopolymer extension
GenBank	MDV086	-	153,340	A	-	Homopolymer extension, frameshift
UK	-	-	151,727	-	T	Homopolymer contraction
GenBank	-	-	154,385	C	-	Homopolymer extension
USDA	-	-	151,853	C	A	Synonymous substitution
USDA	-	-	152,098	-	G	Homopolymer contraction
GenBank	MDV091	US2	159,254	G	A	Val51 -> Ala51
GenBank	MDV092	US3	159,527	C	G	Ser4 -> Thr4
GenBank	MDV092	US3	159,552	T	C	Synonymous substitution
GenBank	MDV092	US3	160,197	T	C	Synonymous substitution
GenBank	MDV093	SORF4	160,899	C	T	Leu22->Ser22
GenBank	MDV093	SORF4	161,025	G	A	Tyr64 -> Cys64
GenBank	MDV093	SORF4	161,269	G	A	Synonymous substitution
GenBank	MDV094	US6 (gD)	161,518	T	C	Synonymous substitution
GenBank	MDV094	US6 (gD)	161,531	C	T	Ile29 -> Thr29
GenBank	MDV094	US6 (gD)	161,709	T	C	Synonymous substitution
GenBank	MDV094	US6 (gD)	161,799	A	C	Synonymous substitution
GenBank	MDV094	US6 (gD)	162,018	T	A	Synonymous substitution
GenBank	MDV094	US6 (gD)	162,065	G	A	Glu207 -> Gly207
GenBank	MDV094	US6 (gD)	162,163	C	G	Gly240 -> Arg240
GenBank	MDV094	US6 (gD)	162,183	C	T	Synonymous substitution
GenBank	MDV094	US6 (gD)	162,218	A	T	Ile258 -> Asn258

461 **Supplementary Table 2: Variable positions in Md5 consensus genomes**

Genome	ORF	Position	Consensus	Call	Description
USDA	-	127,709	C	-	Homopolymer contraction
GenBank	-	129,507	-	A	Homopolymer extension
GenBank	-	129,508	-	A	Homopolymer extension
GenBank	-	129,588	-	T	Homopolymer extension
GenBank	-	130,253	-	T	Homopolymer extension
GenBank	-	134,602	-	G	Homopolymer extension
USDA	-	132,800	G	-	Homopolymer contraction
USDA	-	149,305	C	-	Homopolymer contraction
UK	-	152,067	C	-	Homopolymer contraction
UK	-	152,227	T	-	Homopolymer contraction
UK	-	152,227	G	-	Homopolymer contraction
UK	-	152,239	T	-	Homopolymer contraction

463 **Supplementary Table 3: Minor variants in UK CVI988**

Position in the genome	Major allele	Minor allele	Minor allele frequency	Major allele # forward reads	Major allele # reverse reads	Minor allele # forward reads	Minor allele # reverse reads	ORF	Type
3018	T	C	4.65%	5651	2646	275	130	MDV011	Upstream non-coding
3021	G	T	2.22%	5817	2636	126	66	MDV011	Upstream non-coding
3056	A	G	3.21%	5990	6705	199	222	MDV011	Upstream non-coding
3095	A	G	17.30%	4691	5465	919	1206	MDV011	Non-synonymous (Thr2 -> Ala2)
3204	A	G	10.06%	2353	2384	266	264	MDV011	Non-synonymous (Glu38->Gly38)
22747	A	T	2.65%	2845	3416	63	108	MDV025	Synonymous
22749	T	G	2.09%	2853	3418	46	88	MDV025	Non-synonymous (Thr265->Pro265)
45308	G	T	2.55%	4836	3741	72	153	MDV038	Non-synonymous (Ala239->Ser239)
45326	A	C	2.70%	4512	2768	61	141	MDV038	Non-synonymous (Thr245->Pro245)
108639	G	A	2.11%	6326	5563	138	119	MDV069	Upstream non-coding
108648	G	A	2.26%	6323	5494	151	123	MDV069	Upstream non-coding
108691	A	G	2.34%	5802	5540	145	128	MDV069	Upstream non-coding
108705	T	C	2.66%	3553	5549	127	122	MDV069	Upstream non-coding
108707	T	G	2.66%	3552	5519	122	126	MDV069	Upstream non-coding
109134	A	G	2.43%	4877	5946	116	154	MDV072	Non-synonymous (Leu875->Pro875)
109143	A	G	2.66%	4802	5647	123	163	MDV072	Non-synonymous (Val872->Ala872)
109186	G	A	3.76%	3676	4473	141	178	MDV072	Non-synonymous (His858->Tyr858)
109200	C	T	3.25%	3332	4093	81	169	MDV072	Non-synonymous (Cyst853->Tyr853)
109202	G	T	3.27%	3270	4023	80	167	MDV072	Synonymous
109263	T	G	4.08%	2886	2890	84	162	MDV072	Non-synonymous (His832->Pro832)
109298	C	T	3.28%	2684	2155	51	113	MDV072	Non-synonymous (Trp820->STOP)
121037	A	G	14.59%	25181	19947	5447	2316	MDV076	Non-synonymous (Thr180->Ala180)
121045	A	G	16.71%	23815	19819	6391	2431	MDV077	Non-synonymous (Phe111->Leu111)
121070	C	-	58.08%	29714	19086	14302	14301	MDV076	Deletion
121071	C	-	57.02%	30172	19826	14287	14286	MDV076	Deletion
121072	A	-	57.12%	29125	19029	14288	14287	MDV076	Deletion
124200	T	C	2.20%	3776	3761	86	84	MDV077	Upstream non-coding
131065	T	C	2.60%	11669	13033	441	219	MDV084	Synonymous
144079	A	G	2.16%	6886	5749	157	123	MDV092	Non-synonymous (Glu54->Gly54)
150529	A	G	3.79%	252	271	7	18	MDV093	Downstream non-coding

465   **Supplementary Table 4: Minor variants in UK Md5**

Position in the genome	Major allele	Minor allele	Minor allele frequency	Major allele # forward reads	Major allele # reverse reads	Minor allele # forward reads	Minor allele # reverse reads	ORF	Type
36117	A	G	7.89%	3400	3374	278	303	MDV031	Non-coding
84011	T	G	4.15%	2275	1949	107	76	MDV054	Non-synonymous (His136->Pro136)
84098	G	T	3.82%	2575	1105	106	40	MDV054	Non-synonymous (Thr107->Asn107)
84812	T	C	3.09%	1091	3673	33	119	MDV054	Non-coding
88814	A	G	12.92%	3903	2728	590	397	MDV057	Synonymous
96655	C	T	5.32%	4509	2255	259	121	MDV061	Non-synonymous (Asp35->Asn35)
100749	G	A	5.96%	2422	4690	229	222	MDV066	Non-synonymous (Val192->Ile192)
108784	G	A	2.61%	4183	3701	53	158	MDV069	Non-coding
108793	G	A	2.80%	4122	3688	65	160	MDV069	Non-coding
108836	A	G	3.17%	3769	3694	68	177	MDV069	Non-coding
108850	T	C	3.90%	2128	3659	63	172	MDV069	Non-coding
108852	T	G	4.03%	2122	3624	63	179	MDV069	Non-coding
108914	G	T	3.05%	2982	3659	51	158	MDV069	Non-coding
109246	G	A	2.01%	3501	4391	92	70	MDV072	Non-synonymous (Ala886->Val886)
109279	A	G	2.97%	3503	4076	142	90	MDV072	Non-synonymous (Leu875->Pro875)
109288	A	G	3.29%	3495	3995	158	97	MDV072	Non-synonymous (Val872->Ala872)
109331	G	A	3.57%	2986	3374	151	85	MDV072	Non-synonymous (His858->Tyr858)
109345	C	T	2.55%	2852	3217	78	81	MDV072	Non-synonymous (Cyst853->Tyr853)
109347	G	T	2.57%	2803	3185	76	82	MDV072	Synonymous
109408	T	G	3.58%	2753	2417	110	82	MDV072	Non-synonymous (His832->Pro832)
109443	C	T	2.87%	2726	2034	90	51	MDV072	Non-synonymous (Trp820->STOP)
118716	C	T	2.68%	6309	4595	161	139	MDV076	Upstream non-coding
124098	G	A	3.12%	5261	3764	168	123	MDV078	Synonymous
125178	G	C	2.55%	158	301	4	8	MDV079	Non-synonymous (Ala107->Pro107)
131736	T	G	2.25%	6893	3297	137	98	MDV084	Non-synonymous (Glu1725->Ala1725)
134220	G	A	21.44%	3490	2598	963	699	MDV084	Non-synonymous (Ala897->Val897)
136722	T	C	31.19%	7233	6650	3294	3002	MDV084	Non-synonymous (Gln63->Arg63)

467   **Supplementary Table 5: Published reference genomes for additional 29 strains used**

468   **to construct NJ tree**

Strain	Accession	Publication Date	Length	PMID	Region
Kgs-c1	LC589272	15-Oct-20	174,999	33228722	Asia
J-1	KU744555	18-Feb-17	176,118	27112385	Asia
814	JF742597	23-Feb-12	172,541	21984218	Asia
GA	AF147806	29-Jun-04	174,077	10823954	North America
MD70/13	MF431495	12-Dec-17	177,844	29151863	Europe
648a	JQ806361	4-Apr-13	176,080	22923089	North America
RB-1B	EF523390	29-Oct-07	178,246	17721813	North America
CU-2	EU499381	5-Aug-08	176,922	18351449	North America
HNGS101	MG432697	6-Mar-18	175,888	29407378	Asia
JL/1404	KU744559	18-Feb-17	176,083	27112385	Asia
LTS	KU744557	18-Feb-17	176,023	27112385	Asia
CC/1409	KU744560	18-Feb-17	175,561	27112385	Asia
GX0101	JX844666	3-Dec-12	178,101	23166235	Asia
LHC2	MW247189	19-Apr-22	177,327	35266322	Asia
WC/1203	KU744558	18-Feb-17	176,057	27112385	Asia
LCY	KX290013	12-Apr-17	175,319	28194622	Asia
LCC	KU744556	18-Feb-17	175,525	27112385	Asia
HS/1412	KU744561	18-Feb-17	175,532	27112385	Asia
Polen5	MF431496	12-Dec-17	177,821	29151863	Europe
EU-1	MF431494	12-Dec-17	177,828	29151863	Europe
ATE2539	MF431493	12-Dec-17	177,868	29151863	Europe
LMS	JQ314003	1-Aug-12	177,526	22476905	Asia
HNLC503	MG518371	29-Apr-19	178,195	31698116	Asia
HC/0803	MW531728	10-Jul-22	174,808	35266322	Asia
SY/1209	MW247208	19-Apr-22	177,240	35266322	Asia
Hrb/1504	MW247204	19-Apr-22	177,839	35266322	Asia
DH/1307	MW247201	19-Apr-22	177,235	35266322	Asia
DH/1504	MW247202	19-Apr-22	177,991	35266322	Asia
CH/10	MW247200	19-Apr-22	177,401	35266322	Asia

References

1. Marek J. 1907. Multiple Nervenentzündung (Polyneuritis) bei Hühnern. Dtsch Tierärztl Wochenschr 15:417–421.

2. Churchill AE, Biggs PM. 1967. Agent of Marek's disease in tissue culture. Nature 215:528–530.

3. Witter RL, Burgoyne GH, Solomon JJ. 1968. Preliminary studies on cell cultures infected with Marek's disease agent. Avian Dis 12:169–185.

4. Churchill AE, Payne LN, Chubb RC. 1969. Immunization against Marek's disease using a live attenuated virus. Nature 221:744–747.

5. Witter RL, Nazerian K, Purchase HG, Burgoyne GH. 1970. Isolation from turkeys of a cell-associated herpesvirus antigenically related to Marek's disease virus. Am J Vet Res 31:525–538.

6. Okazaki W, Purchase HG, Burmester BR. 1970. Protection against Marek's disease by vaccination with a herpesvirus of turkeys. Avian Dis 14:413–429.

7. Witter RL, Sharma JM, Fadly AM. 1980. Pathogenicity of Variant Marek's Disease Virus Isolants in Vaccinated and Unvaccinated Chickens. Avian Dis 24:210–232.

8. Schat KA, Calnek BW, Fabricant J. 1982. Characterisation of two highly oncogenic strains of Marek's disease virus. Avian Pathol J WVPA 11:593–605.

9. Witter RL. 1982. Protection by attenuated and polyvalent vaccines against highly virulent strains of Marek's disease virus. Avian Pathol J WVPA 11:49–62.

10. Witter RL. 1997. Increased virulence of Marek's disease virus field isolates. Avian Dis 41:149–163.

11. Rispens BH, van Vloten H, Mastenbroek N, Maas HJ, Schat KA. 1972. Control of Marek's disease in the Netherlands. I. Isolation of an avirulent Marek's disease virus (strain CVI 988) and its use in laboratory vaccination trials. Avian Dis 16:108–125.

12. Schat KA. 2016. History of the First-Generation Marek's Disease Vaccines: The Science and Little-Known Facts. Avian Dis 60:715–724.

13. Schumacher D, Tischer BK, Teifke J-P, Wink K, Osterrieder N. 2002. Generation of a permanent cell line that supports efficient growth of Marek's disease virus (MDV) by constitutive expression of MDV glycoprotein E. J Gen Virol 83:1987–1992.

14. Zhang Y, Li Z, Bao K, Lv H, Gao Y, Gao H, Qi X, Cui H, Wang Y, Ren X, Wang X, Liu C. 2015. Pathogenic characteristics of Marek's disease virus field strains prevalent in China and the effectiveness of existing vaccines against them. Vet Microbiol 177:62–68.

15. Sun G-R, Zhang Y-P, Lv H-C, Zhou L-Y, Cui H-Y, Gao Y-L, Qi X, Wang Y-Q, Li K, Gao L, Pan Q, Wang X-M, Liu C-J. 2017. A Chinese Variant Marek's Disease Virus Strain with Divergence between Virulence and Vaccine Resistance. Viruses 9:E71.

16. Davison F, Venugopal N. 2004. Marek's Disease: An Evolving Problem, 1st ed. Academic Press.

17. Lee LF, Wu P, Sui D, Ren D, Kamil J, Kung HJ, Witter RL. 2000. The complete unique long sequence and the overall genomic organization of the GA strain of Marek's disease virus. Proc Natl Acad Sci U S A 97:6091–6096.

18. Tulman ER, Afonso CL, Lu Z, Zsak L, Rock DL, Kutish GF. 2000. The genome of a very virulent Marek's disease virus. J Virol 74:7980–7988.

19. Firth C, Kitchen A, Shapiro B, Suchard MA, Holmes EC, Rambaut A. 2010. Using time-structured data to estimate evolutionary rates of double-stranded DNA viruses. Mol Biol Evol 27:2038–2051.

20. Renzette N, Gibson L, Jensen JD, Kowalik TF. 2014. Human cytomegalovirus intrahost evolution-a
new avenue for understanding and controlling herpesvirus infections. Curr Opin Virol 8:109–115.

21. Kuny CV, Szpara ML. 2021. Alphaherpesvirus Genomics: Past, Present and Future. Curr Issues Mol
Biol 42:41–80.

22. Renner DW, Szpara ML. 2018. Impacts of Genome-Wide Analyses on Our Understanding of Human
Herpesvirus Diversity and Evolution. J Virol 92:e00908-17.

23. Gimeno IM, Cortes AL, Faiz NM, Barbosa T, Villalobos T. 2015. Evaluation of Factors Influencing
Efficacy of Vaccine Strain CVI988 Against Marek's Disease in Meat-Type Chickens. Avian Dis
59:400–409.

24. Gimeno IM, Witter RL, Reed WM. 1999. Four distinct neurologic syndromes in Marek's disease:
effect of viral strain and pathotype. Avian Dis 43:721–737.

25. Spatz SJ, Volkening JD, Gimeno IM, Heidari M, Witter RL. 2012. Dynamic equilibrium of Marek's
disease genomes during in vitro serial passage. Virus Genes 45:526–536.

26. Spatz SJ. 2010. Accumulation of attenuating mutations in varying proportions within a high passage
very virulent plus strain of Gallid herpesvirus type 2. Virus Res 149:135–142.

27. Hildebrandt E, Dunn JR, Perumbakkam S, Niikura M, Cheng HH. 2014. Characterizing the
molecular basis of attenuation of Marek's disease virus via in vitro serial passage identifies de novo
mutations in the helicase-primase subunit gene UL5 and other candidates associated with reduced
virulence. J Virol 88:6232–42.

28. Spatz SJ, Petherbridge L, Zhao Y, Nair V. 2007. Comparative full-length sequence analysis of
oncogenic and vaccine (Rispens) strains of Marek's disease virus. J Gen Virol 88:1080–1096.

29. Stratton M. 2008. Genome resequencing and genetic variation. Nat Biotechnol 26:65–66.

30. Lauring AS. 2020. Within-Host Viral Diversity: A Window into Viral Evolution. Annu Rev Virol
7:63–81.

31. Kennedy DA, Cairns C, Jones MJ, Bell AS, Salathé RM, Baigent SJ, Nair VK, Dunn PA, Read AF.
2017. Industry-Wide Surveillance of Marek's Disease Virus on Commercial Poultry Farms. Avian
Dis 61:153–164.

32. Trimpert J, Groenke N, Jenckel M, He S, Kunec D, Szpara ML, Spatz SJ, Osterrieder N, McMahon
DP. 2017. A phylogenomic analysis of Marek's disease virus reveals independent paths to virulence
in Eurasia and North America. Evol Appl 10:1091–1101.

33. Shipley MM, Rathbun MM, Szpara ML. 2020. Oligonucleotide Enrichment of HSV-1 Genomic
DNA from Clinical Specimens for Use in High-Throughput Sequencing. Methods Mol Biol Clifton
NJ 2060:199–217.

34. Parsons LR, Tafuri YR, Shreve JT, Bowen CD, Shipley MM, Enquist LW, Szpara ML. 2015. Rapid
genome assembly and comparison decode intrastrain variation in human alphaherpesviruses. mBio
6:e02213-14.

35. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile
metagenomic assembler. Genome Res 27:824–834.

36. Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res
27:573–580.

37. Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

38. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011.
Integrative genomics viewer. Nat Biotechnol 29:24–26.

39. Sato J, Murata S, Yang Z, Kaufer BB, Fujisawa S, Seo H, Maekawa N, Okagawa T, Konnai S,
Osterrieder N, Parcells MS, Ohashi K. 2022. Effect of Insertion and Deletion in the Meq Protein
Encoded by Highly Oncogenic Marek's Disease Virus on Transactivation Activity and Virulence.
Viruses 14:382.

40. Murata S, Yamamoto E, Sakashita N, Maekawa N, Okagawa T, Konnai S, Ohashi K. 2021. Research
Note: Characterization of S-Meq containing the deletion in Meq protein's transactivation domain in
a Marek's disease virus strain in Japan. Poult Sci 100:101461.

41. Lee SI, Takagi M, Ohashi K, Sugimoto C, Onuma M. 2000. Difference in the meq gene between
oncogenic and attenuated strains of Marek's disease virus serotype 1. J Vet Med Sci 62:287–292.

42. Renz KG, Cooke J, Clarke N, Cheetham BF, Hussain Z, Fakhrul Islam AFM, Tannock GA,
Walkden-Brown SW. 2012. Pathotyping of Australian isolates of Marek's disease virus and
association of pathogenicity with *meq* gene polymorphism. Avian Pathol 41:161–176.

43. Bell AS, Kennedy DA, Jones MJ, Cairns CL, Pandey U, Dunn PA, Szpara ML, Read AF. 2019.
Molecular epidemiology of Marek's disease virus in central Pennsylvania, USA. Virus Evol
5:vey042.

44. Domingo E, Perales C. 2019. Viral quasispecies. PLoS Genet 15:e1008271.

45. Singh K, Mehta D, Dumka S, Chauhan AS, Kumar S. 2023. Quasispecies Nature of RNA Viruses:
Lessons from the Past. Vaccines 11:308.

46. Trimpert J, Groenke N, Kunec D, Eschke K, He S, McMahon DP, Osterrieder N. 2019. A
proofreading-impaired herpesvirus generates populations with quasispecies-like structure. Nat
Microbiol https://doi.org/10.1038/s41564-019-0547-x.

47. Shipley MM, Renner DW, Ott M, Bloom DC, Koelle DM, Johnston C, Szpara ML. 2018. Genome-
wide surveillance of genital herpes simplex virus type 1 from multiple anatomic sites over time. J
Infect Dis 218:595–605.

48. Renner DW, Szpara ML. 2018. The impacts of genome-wide analyses on our understanding of
human herpesvirus diversity and evolution. J Virol 92:e00908-17.

49. Loncoman CA, Vaz PK, Coppo MJ, Hartley CA, Morera FJ, Browning GF, Devlin JM. 2017. Natural
recombination in alphaherpesviruses: Insights into viral evolution through full genome sequencing
and sequence analysis. Infect Genet Evol J Mol Epidemiol Evol Genet Infect Dis 49:174–185.

50. López de Juan Abad BA, Cortes AL, Correa M, Gimeno IM. 2019. Evaluation of Factors That
Influence Dose Variability of Marek's Disease Vaccines. Avian Dis 63:591–598.

51. Dambrine G, Labaille J, Boissel É, Dupuy C, Rasschaert D. 2016. Marek's disease herpesvirus: a
model of vaccine-dependent adaptation. Virol Montrouge Fr 20:273–286.

52. Witter RL, Calnek BW, Buscaglia C, Gimeno IM, Schat KA. 2005. Classification of Marek's disease
viruses according to pathotype: philosophy and methodology. Avian Pathol J WVPA 34:75–90.

53. Kuny CV, Bowen CD, Renner DW, Johnston CM, Szpara ML. 2020. In vitro evolution of herpes
simplex virus 1 (HSV-1) reveals selection for syncytia and other minor variants in cell culture.
Virus Evol 6:veaa013.

54. Baigent SJ, Nair VK, Le Galludec H. 2016. Real-time PCR for differential quantification of CVI988
vaccine virus and virulent strains of Marek's disease virus. J Virol Methods 233:23–36.

55. Petherbridge L, Howes K, Baigent SJ, Sacco MA, Evans S, Osterrieder N, Nair V. 2003. Replication-
competent bacterial artificial chromosomes of Marek's disease virus: novel tools for generation of
molecularly defined herpesvirus vaccines. J Virol 77:8712–8718.

56. Spatz SJ, Silva RF. 2007. Sequence determination of variable regions within the genomes of gallid
herpesvirus-2 pathotypes. Arch Virol 152:1665–1678.

57. Spatz SJ, Rue CA. 2008. Sequence determination of a mildly virulent strain (CU-2) of Gallid
herpesvirus type 2 using 454 pyrosequencing. Virus Genes 36:479–489.

58. Spatz SJ, Zhao Y, Petherbridge L, Smith LP, Baigent SJ, Nair V. 2007. Comparative sequence
analysis of a highly oncogenic but horizontal spread-defective clone of Marek's disease virus. Virus
Genes 35:753–766.

59. He L, Li J, Zhang Y, Luo J, Cao Y, Xue C. 2018. Phylogenetic and molecular epidemiological
studies reveal evidence of recombination among Marek's disease viruses. Virology 516:202–209.

60. He L, Li J, Peng P, Nie J, Luo J, Cao Y, Xue C. 2020. Genomic analysis of a Chinese MDV strain
derived from vaccine strain CVI988 through recombination. Infect Genet Evol J Mol Epidemiol
Evol Genet Infect Dis 78:104045.

61. Xu H, Krieter AL, Ponnuraj N, Tien YY-T, Kim T, Jarosinski KW. 2022. Coinfection in the host can
result in functional complementation between live vaccines and virulent virus. Virulence 13:980–
989.

62. Zhang Y, Lan X, Wang Y, Lin Y, Yu Z, Guo R, Li K, Cui H, Qi X, Wang Y, Gao L, Pan Q, Liu A,
Gao Y, Wang X, Liu C. 2022. Emerging natural recombinant Marek's disease virus between
vaccine and virulence strains and their pathogenicity. Transbound Emerg Dis 69:e1702–e1709.

63. van Belkum A, Scherer S, van Alphen L, Verbrugh H. 1998. Short-sequence DNA repeats in
prokaryotic genomes. Microbiol Mol Biol Rev MMBR 62:275–293.

64. Rhoads A, Au KF. 2015. PacBio Sequencing and Its Applications. Genomics Proteomics
Bioinformatics 13:278–289.

65. Wang Y, Zhao Y, Bollas A, Wang Y, Au KF. 2021. Nanopore sequencing technology, bioinformatics
and applications. Nat Biotechnol 39:1348–1365.