CAR T cells recognizing CD276 and Dual-CAR T cells against CD276/FGFR4 promote rhabdomyosarcoma clearance in an orthotopic mouse model

Andrea Timpanaro[1,2,3]; Caroline Piccand[1,2,3]; Dzhangar Dzhumashev[1,2,3]; Stenija Anton-Joseph[1,2,3], Andrea Robbi[1,2]; Janine Moser[1,2]; Jochen Rössler[1,2]; Michele Bernasconi[1,2*]

[1] Department of Pediatric Hematology and Oncology, Inselspital, Bern University Hospital, 3010 Bern, Switzerland

[2] Translational Cancer Research, Department for BioMedical Research (DBMR), University of Bern, 3008 Bern, Switzerland

[3] Graduate School for Cellular and Biomedical Sciences, University of Bern, 3012 Bern, Switzerland

* Correspondence: michele.bernasconi@unibe.ch

Abstract (350)

Background: Rhabdomyosarcoma (RMS) is the most common soft tissue sarcoma in childhood. Prognosis is still poor especially for metastatic, high-grade, and relapsed RMS. New therapies are urgently needed, especially systemic therapies. Chimeric Antigen Receptor T cells (CAR Ts) are very effective against B-cell malignancies, but efficacy against solid tumors needs to be improved. CD276 is upregulated in RMS and detected at low levels in normal tissues. FGFR4 is a very specific target for RMS. Here, we optimized CAR Ts for RMS recognizing the two targets and their combination, and we performed *in vitro* and *in vivo* studies of effectiveness for tumor growth inhibition.

Methods: RMS cells killing and cytokines production by CD276-CAR Ts with combinations of CD8α or CD28 HD/TM domains and 4-1BB and/or CD28 co-stimulatory domains were tested *in vitro*. Four different single-domain antibodies were used to select the most specific FGFR4-CAR construct. The selected CD276- and FGFR4-CAR Ts were used to generated Dual-CAR Ts, and tumor killing was tested *in vivo* in three orthotopic RMS mouse models.

Results: CD276.V-CAR Ts (276.MG.CD28HD/TM.CD28CSD.3z) showed the strongest killing of RMS cells, and the highest release of IFN-γ and Granzyme B *in vitro*. FGFR4.V-CAR Ts (F8-FR4.CD28HD/TM.CD28CSD.3z) showed the most specific killing. CD276-CAR Ts successfully eradicated RD- and Rh4-derived RMS tumors *in vivo*, achieving complete remission in 3/5 and 5/5 mice, respectively; but in CD276^{low} JR-tumors, they achieved complete remission in only 1/5 mice. FGFR4 CAR Ts instead delayed growth of Rh4-tumor. CD276/FGFR4-Dual-CAR Ts promoted Rh4-tumors clearance in 5/5 mice.

Conclusions: CD276- and CD276/FGFR4-directed CAR Ts showed very strong RMS cells killing *in vitro* and eradication of CD276^{high} RMS tumors *in vivo*. CD276^{low} tumors escaped the therapy showing a correlation of antigen density and effectiveness. FGFR4-CAR Ts showed specific killing *in vitro* and only delayed RMS growth *in vivo*. Our results show that combined expression of CD276-CAR with other CAR does not reduce its activity. Introducing immunotherapy with CD276-CAR Ts in RMS seems to be feasible and promising, although CAR constructs design and target combinations have to be further improved to achieve synergism and to eradicate tumors with low target expression.

Keywords

Rhabdomyosarcoma, CAR T cells, CD276, FGFR4, immunotherapy

Background

Pediatric rhabdomyosarcoma (RMS) is the most common soft tissue sarcoma in children and young adults (1), and each year it accounts for 3% of childhood cancers (2). Based on histology, RMS can be classified in four main subtypes: embryonal RMS (eRMS; 60-70%), alveolar RMS (aRMS; 20-30%), pleomorphic RMS (pRMS) and spindle cell/sclerosing RMS (s-scRMS) accounting for 7–15% of the cases (3). RMS classification has been further refined in 'fusion positive' (FP) and 'fusion negative' (FN) RMS (4), based on two main characteristic chromosomal translocations, the t(2;13)(q35;q14) or the t(1;13)(p36;q14) resulting in the expression of a PAX3-FOXO1 or PAX7-FOXO1 fusion transcription factor, respectively (5, 6, 7). The onset of RMS has been attributed mainly to cells of myogenic lineage, but more recently also non-myogenic mesenchymal cells were indicated as possible RMS progenitors (8, 9, 10, 11). As a result, RMS can develop not only in skeletal muscles, like head and neck, trunk, and extremities, but also in distant sites, including genitourinary and biliary tract. Despite overall 5-year survival rates have improved with the combined use of surgery, radiation therapy, and chemotherapy, in pediatric patients with metastatic and high-risk disease, such as aRMS, prognosis remains poor (12). Therefore, new therapies are needed for children and young adults with high-risk and recurrent RMS.

Immunotherapies using Chimeric Antigen Receptor (CAR) T cells (13, 14) have shown extraordinary results in patients with B cell malignancies (15), leading to the FDA approval of six CAR T cell therapies so far (16). Target molecules for these CART cells are surface proteins on leukemia or lymphoma cell surface, such as CD19 and CD20. However, CAR T cell therapy remains challenging for solid tumors, mainly due to a lack of ideal tumor specific target molecules, and to the immunosuppressive tumor microenvironment (TME) of solid tumors limiting the activity of CAR T cells (17, 18).

Several targets have been investigated pre-clinically and clinically for immunotherapies of pediatric solid tumors (reviewed in (19, 20, 21)), including for RMS. More than 10 years ago, CAR T cell therapy for RMS has been explored targeting the fetal acetylcholine receptor gamma subunit gamma (fAChRγ) surface protein which is not present in humans after birth and is therefore highly restricted to RMS (22). The fAChRγ-CAR T cells exhibited specific, MHC-independent, *in vitro* lysis and cytokine secretion in response to antigen-expressing target cells and were moderately immunoprotective in murine tumor xenograft models (23, 24). Unfortunately, their activity was not sufficient to eradicated tumors, probably due to fAChRγ low expression levels. Other surface proteins explored for RMS CAR T cell therapy in RMS are IGF1R and ROR1. The corresponding CAR T cells showed effective *in vitro* RMS cell lysis, but have not been tested *in vivo* (25). Other approaches have further investigated RMS

cell surface antigens for CAR T cell therapy: PDGFRα-CAR T-cell showed good activity *in vitro* and tumor control in a subcutaneous RMS xenograft mouse model (26) as PDGFRα is aberrantly expressed in RMS (27); HER2/ErbB2-CAR-engineered cytokine-induced killer cells showed potent activity on RMS spheroids (28) based on an expression level of the receptor tyrosine kinase HER2/ErbB2 in 10-33% of RMS on the protein level (29, 30). Importantly, in a phase I clinical trial of HER2/ErbB2-CAR T cells (NCT00902044) two complete responses were observed in RMS patients without any safety concerns (31). In particular, one patient with bone marrow metastatic involvement showed complete response after HER2/ErbB2-CAR T cell therapy (31). Despite all these promising results, more targets and better CAR designs need to be investigated to cover all RMS patients and to avoid relapses mediated by antigen-escape.

In our previous work on identifying potential surface antigens by surfaceome profiling of RMS for an immunotherapy approach (32), we identified two targets that are particularly appealing for a CAR T cell therapy: CD276 and FGFR4.

The type I transmembrane protein CD276 (B7-H3) belongs to the B7 immune co-stimulatory and co-inhibitory family (33), and is expressed on antigen-presenting cells (APC) interacting with immune checkpoint inhibitors, such as CTLA-4, PD-1, and CD28 (34). CD276 has both co-stimulatory and co-inhibitory functions in the regulation of different T cell subsets. CD276 is expressed at high levels in a high proportion of human malignancies, including the pediatric solid tumors RMS (35, 36), osteosarcoma (36, 37), intrinsic pontine glioma (38), medulloblastoma, and Ewing sarcoma (36). Importantly, an immune evasive role for CD276 in the context of RMS has been recently described, further supporting the important impact its targeting might have on disease control by the immune system (39). CD276-CAR T cells showed efficacy in preclinical models of pancreatic ductal adenocarcinoma, ovarian cancer, and neuroblastoma without toxicity (40). Of note, a CD276-CAR armed with an antigen-binding domain derived from Enoblituzumab (MGA271), a humanized antibody from MacroGenics recognizing an epitope of CD276 with high tumor reactivity, was particularly effective against pediatric solid tumors osteosarcoma, Ewing sarcoma and medulloblastoma (36), but has not been tested for RMS *in vitro*, nor *in vivo*.

The other target of interest, the Fibroblast Growth Factor Receptor 4 (FGFR4) is a receptor tyrosine kinase, that shows very specific expression in RMS, and in normal tissues is only expressed at low levels in liver and in mature skeletal tissue (41, 42, 43, 44). FGFR4-CAR T cells are active *in vitro* against both aRMS and eRMS cell lines (45, 46), and show effective response against disseminated disease *in vivo*, but not in an RMS orthotopic model (47). Interestingly, the combination of FGFR4-CAR T cells with pharmacological inhibition of the

myeloid component in stroma allowed FGFR4-CAR T cells to eradicate orthotopic RMS tumors in mice (47).

Concerning the design of CAR constructs, there is a need to optimize it to the special situation of RMS. Clearly, each domain of the CAR – the extracellular antigen-binding domain (ABD), the hinge domain (HD); the transmembrane domain (TM); and the intracellular co-stimulatory domains (CSD) - can impact the activity of CAR T cells (48, 49). The ABD determines the affinity and specificity, while its efficacy is influenced by the target accessibility and density on the surface of tumor cells (50). H/TM and CSD contribute to regulate the activation threshold (51). The TM domain can affect the stability of CAR immune synapses and regulates the signaling threshold. CD28- or CD8α-derived TM are the most frequently used. CAR T cells armoring CD28 TM domain show higher CAR stability on the T cell surface, and a stronger response, but may be prone to overactivation, resulting in activation-induced cell death (AICD) (52). On the other hand, CAR T cells with a CD8α TM domain may have a lower activation threshold, increasing their cytotoxic potential, but release lower levels of TNF-α and IFN-γ (51, 53). The clinically used CSDs are derived from 4-1BB or CD28 proteins. CSDs are necessary for proper activation, expansion, and most importantly, for persistence of CAR T cells. CARs containing a CD28 CSD show high cytokine release upon antigen recognition through activation of the PI3K pathways, promoting the survival of cytotoxic CAR T cells (54, 55); whereas, 4-1BB confers higher T cell persistence by promoting central memory T cells differentiation through the activation of the NF-κB signaling pathways (55, 56).

Here, we aimed at selecting *in vitro* the most active CAR construct targeting CD276, FGFR4, or the combination of both, on RMS cells. A combinatorial panel of eight CAR constructs, comprising second- and third-generation constructs, with CD8α or CD28 HD/TM domains and intracellular CSDs from 4-1BB and/or CD28 were produced and tested. The most promising CAR T cells were tested for their activity *in vivo* in an orthotopic RMS mouse model.

Materials and Methods

Cell culture

Human RMS cell lines RD, Rh4, Rh5, Rh18, Rh28, Rh30, Rh36, JR, RMS, RUCH-3, TTC-442 were kindly provided by Prof. Beat Schäfer (University Children's Hospital of Zurich). PDXs IC-pPDX-104 from a 7-year-old female with recurrent primary aRMS with PAX3-FOXO1 translocation, IC-pPDX-35 from a 13-year-old male with recurrent metastatic aRMS with PAX3-FOXO1 translocation, were established at the Institut Curie in Paris, France (57, 58); PDX RMS-ZH003 was established from a recurrent aRMS with PAX3-FOXO1 translocation at the University Children's Hospital Zurich, as described in (57, 58). Immortalized human healthy primary myoblasts KM155C25Dist, kindly provided by the platform for immortalization of human cells MyoLine from the Institut de Myologie (Sorbonne University, Paris, France), referred to as myoblasts, and MRC-5 cell lines were used as negative controls. All the cell lines, with the exception of human myoblasts, were cultured in DMEM medium (BioConcept, #1-26F01-I), supplemented with 10% fetal bovine serum (FBS, Thermofisher, #10270106), L-glutamine 2 mM (BioConcept, #5-10K00-H) and 100 U/ml Penicillin-Streptomycin (BioConcept, #4-01F00-H) at 37°C and 5% CO_2 in a humidified incubator. Myoblasts were cultured in Skeletal Muscle Cell Growth Medium (PromoCell, #C-23060) supplemented with Skeletal Muscle Cell Growth Medium SupplementMix (PromoCell, #C-39365). Jurkat cells were cultured in RPMI-1640 medium (Gibco, #42401-018), supplemented with 10% fetal bovine serum (FBS, Thermofisher, #10270106), L-glutamine 2 mM (BioConcept, #5-10K00-H) and 100 U/ml Penicillin-Streptomycin (BioConcept, #4-01F00-H) at 37°C and 5% CO_2 in a humidified incubator. PDXs were cultured in pre-coated P10 dishes with 1:10 diluted Matrigel (Corning, #354234) in 3 ml of pre-cooled Advanced DMEM/F-12 (Gibco, #12634010), supplemented with Glutamax (Gibco, #35050), 100 U/ml Penicillin-Streptomycin (BioConcept, #4-01F00-H), 1x B-27 (Life Technologies, #17504044), 20 ng/ml bFGF (PeproTech, #AF-100-18B), 20 ng/ml EGF (PeproTech, #AF-100-15), and 1.25 mM N-acetyl-L-cysteine (Sigma, #A9165).

Plasmids

The lentiviral plasmid pLenti.PGK.chFP.W was a gift from Rusty Lansford (Addgene, #51008). It encodes mCherry reporter gene and was used to produce lentiviruses to infect RMS cell lines for live-imaging experiments. To express truncated-CD19 protein (tCD19) on RMS cell lines as target for CD19 CAR T cells, the lentiviral plasmid pLenti-fLuc-P2A-tCD19 was

generated by excising with AjiI and MluI digestion the Neomycin resistance in the plasmid Lenti-luciferase-P2A-Neo (Addgene, #105621) and replacing it by GIBSON assembly with the tCD19 sequence amplified by PCR from the vector pcDNA3.1+CD19-C-DYK (Genescript, OHU27320D) using the primers P2A-tCD19 Forward and P2A-tCD19 Reverse (Suppl. Table S1).

Production of lentiviruses

Lentiviruses were generated by transient co-transfection of Lenti-X 293T cells (Takara, #632180). Briefly, $5x10^6$ Lenti-X 293T cells were seeded onto 10 cm dish, pre-coated with collagen type I rat tail (1:10 diluted in sterile water, Sigma, #9007-34-5), and the day after, transfected with the calcium phosphate method. 10 µg of psPAX2 (Addgene, Plasmid #12260), 4 µg of pVSV-G (Addgene, Plasmid #8454), and 10 µg of lentiviral plasmid of interest were used. The plasmids were mixed in 500 µl 0.25 M CaCl2, added dropwise to 500 µl of HBS 2x (NaCl 0.28 M, HEPES 0.05 M, Na_2HPO_4 1.5 mM, pH 7.05), and incubated for 15 min at room temperature. The solution was then dropwise added to Lenti-X 293T cells. Lentiviruses-containing supernatants were harvested 48h post-transfection, filtered with 0.45 µm filters, and concentrated via overnight centrifugation at 4'000xg at 4°C. Lentiviruses were aliquoted and stored at -80°C until use.

Lentiviruses titration

Titration of Lentiviruses was carried out with Jurkat T cells. 250'000 cells were plated onto 24-well plates and were transduced with ten-fold dilutions of Lentiviruses. 48h post-transduction, fluorescent cells were quantified by Flow Cytometry. The titer was calculated with the following formula:

$$\frac{TU}{mL} = \frac{[(\text{Number of cells transduced} * \text{\% fluorescent cells} * \text{Dilution Factor})]}{(\text{Transduction Volume in mL})}$$

mCherry$^+$ RMS cell lines

The lentiviral plasmid pLenti.PGK.chFP.W (Addgene, #51008) was used to generate mCherry$^+$ lentiviruses to transduce RD and Rh4 cell lines at a multiplicity of infection (MOI) of 5. RMS cells stably expressing mCherry were sorted 48h after transduction by mCherry and amplified.

Firefly Luciferase (fLuc$^+$) RMS cell lines

pHIV-iRFP720-E2A-Luc was a gift from Antonius Plagge (Addgene plasmid # 104587) and was used to generate iRFP720$^+$fLuc$^+$ lentiviruses and to stably transduce RD and Rh4 cell lines at an MOI of 5. iRFP720$^+$fLuc$^+$ RMS cell lines were sorted 48h after transduction and amplified.

Truncated CD19 (tCD19$^+$) RMS cell lines

The pLenti-fLuc-P2A-tCD19 plasmid was used to generate tCD19$^+$fLuc$^+$ lentiviruses to stably transduce RD and Rh4 cells at an MOI of 5. 48h after transduction, tCD19$^+$fLuc$^+$ RMS cell lines were stained with BV421-conjugated anti-CD19 antibody for 30 min at room temperature, sorted by BV421, and amplified.

CRISPR/Cas9

Two in silico-designed sgRNA sequences targeting CD276 and FGFR4 genes, respectively, were selected by using the Graphical User Interface for DNA Editing Screens (GUIDES), available at guides.sanjanalab.org (59). The two sgRNA sequences (Supplementary Table S2) were cloned into the pLentiguide-PURO plasmid (Addgene, #52963). The plasmid pLentiCRISPR-mCherry (Addgene, #75161) was used to stably express Cas9 nuclease and mCherry in RD and Rh4 cell lines. 48h after lentiviral transduction at an MOI of 5, cells expressing mCherry were sorted by FACS. Cas9$^+$mCh$^+$ RMS cells were then transiently transfected with pLentiguide-PURO plasmid and selected with 1 µg/ml Puromycin (InVivoGen, #ant-pr-1) for two days, single cell cultures were then generated by limiting dilutions, and cells were expanded.

CAR plasmids generation

The lentiviral plasmids (Table S3) used for the CAR T cells production were generated with GeneArt Gibson assembly HiFi cloning kit (Thermofisher, #A46624EA). The sequences of FMC63 scFv targeting CD19, CD8αHD, and CD28TM CAR domains were amplified from the plasmids pSLCAR-CD19-hCAR-BB (Addgene, #135992) and pSLCAR-CD19-28z (Addgene, #135991) kindly provided by Dr. Scott McComb (National Research Council of Canada, Ottawa, Canada) (60). The sequence of CD276.MG scFv was kindly provided by Prof. Robbie Majzner (Stanford University, Stanford, CA). To generate the CD276 and F8-FR4 CAR constructs, the FMC63 scFv, flanked by Esp3I sites, was swapped with the PCR-amplified

sequences of CD276.MG scFv or F8-FR4 sdAb. The swapping reaction was performed by incubating the DNA samples with Esp3I (BsmBI) (Thermofisher, #ER0451) and T4 ligase (Thermofisher, #EL0011) at 37°C for 10 min, followed by 5 cycles at 16°C 10 min and 37°C 60 min and a final enzyme inactivation at 80°C for 5 min (61). The plasmids were then transformed in One Shot TOP10 chemically competent cells (Thermofisher, #C404010), amplified, and purified by NucleoBond Xtra Midi kit (Macherey-Nagel, #740410.50). Purity and concentration were evaluated with Nanodrop. Plasmids were sequenced at Microsynth (Balgach, Switzerland).

PBMCs and T cells isolation

PBMCs were isolated with Lymphoprep (STEMCELL, #07801) from different donors' buffy coats obtained from the "Rotkreuz-Organisation für die Menschen im Kanton Bern", and frozen in FBS (Thermo Scientific, #10270106) with 10% DMSO (Sigma, #D2438). T cells were purified from thawed PBMCs by using EasySep^{TM} Human CD8$^+$ T cell Isolation Kit (STEMCELL, #17953) or EasySep Human T cell Isolation Kit (STEMCELL, #17951) following the manufacturer's instructions, and used for CAR T cells production.

CAR T cells production

On day 0, $1x10^6$ T cells freshly purified from PBMCs were activated with CD3/CD28 T Cell Activators (STEMCELL, #10971). T cells were expanded for 3 days in XF Expansion medium (STEMCELL, #10981) with addition of 10 ng/ml IL-2 (STEMCELL, #78036.2) and 10 ng/ml IL-21 (STEMCELL, #78082). On day 4, T cells were transduced with Lentiviruses at an MOI of 5. On day 7, GFP$^+$ CAR T cells were sorted by FACS and expanded until day 13 for cytotoxicity assays, cytokine production, or *in vivo*.

T cells transduction

250'000 Jurkat cells or purified T cells were seeded onto a 24-well plate and transduced at an MOI of 5 via spin-infection for 1h at 800xg and 32°C in 450 µl RPMI-1640 medium (Gibco, #42401-018) containing 50 µl of 100-fold concentrated Lentiviruses. After spin-infection, the cells were diluted 2.5-fold in RPMI-1640 supplemented with 10 ng/ml IL-2 and further incubated for 4h with LV at 37°C. The next day, the cells were centrifuged and resuspended in fresh cell culture medium. After 48h, transduction efficiency was assessed by Flow Cytometry based on GFP and MycTag expression.

Bioluminescence killing assays

Bioluminescence killing assays were performed by co-incubating 5'000 iRFP720$^+$fLuc$^+$ RMS target cells, seeded 24h in advance in bioluminescence 96-well plates (Thermofisher, #136101), with GFP$^+$ CAR T cells at the Effector:Target (E:T) ratios of 0.5:1, 1:1, 5:1, and 10:1, in 200 µl total of complete RPMI-1640 medium. Proliferation of RMS cells was monitored for 72h. The supernatant was gently aspirated, and each well was filled with 75µl. The plates were then frozen at -20°C until analysis. The day of the analysis, bioluminescence released by RMS target cells was measured by using ONE-Glo^{TM} Luciferase assay system (Promega, #E6120) with GLO-Max device (Promega). The results were analyzed with Microsoft Excel software.

Western blotting (WB)

Cells were detached with a scraper, resuspended in PBS and pelleted by centrifugation at 10'000xg for 3 min, 4°C. Pellets were lysed in RIPA buffer (#89900, Thermofisher) with addition of Halt™ Protease Inhibitor Cocktail (100x) (#78430, Thermofisher) for 20 min on ice. Cells were further homogenized with 1 min sonication (10% duty cycle, Branson Sonifer 250). Protein concentration was measured by Pierce BCA Protein Assay Kit (#23225, Thermofisher). Loading buffer (Invitrogen, #NP0007) was added to 30 µg of protein for each sample. After heating at 65°C for 10 min, samples were separated by SDS-PAGE in Tris-Glycine running buffer (BioRad, #1610732, 10x: 25mM Tris base, 1.92 M glycine, 0.1 % SDS), at 100V for about 2h. Separated proteins were then transferred to a PVDF membrane in transfer buffer (BioRad, #1610732. 10x: 250 mM Tris base, 1.92 M glycine, 0.1 % SDS) with 10% methanol, at a constant voltage of 100V at 4°C for 1.5 h. PVDF membrane was blocked by 30 min incubation in TBS-T (TBS 10x: 200mM Tris (Sigma, #1185-53-1), 1.37M NaCl (Sigma, #S9888), pH 7.6; supplemented with Tween 0.1%; (Sigma, #P1379)) and 10% skimmed milk. Membranes were then incubated overnight at 4°C with the corresponding primary antibody in TBS-T. After 1h incubation at room temperature with the secondary antibody, the signal was developed with SuperSignal West Femto Maximum Sensitivity Substrate (Thermofisher, #34094) and measured by using ChemiDoc MP Imaging System (Bio-Rad).

Live-imaging cytotoxicity assays

Live-imaging cytotoxicity assays were performed by co-culturing 5'000 mCh$^+$ RMS target cells, seeded 24h in advance in transparent 96-well plates (Greiner bio-One, #650180), with GFP$^+$ CAR T cells at Effector:Target (E:T) ratios of 0.5:1, 1:1, 5:1, and 10:1, in 200 µl total of complete RPMI-1640 medium. Proliferation of RMS cells was monitored for 48h and images were acquired every hour using Incucyte S3 device (Sartorius). Analyses of the results were performed with IncuCyte 2019B Rev2 software.

Enzyme-linked immunosorbent assays

ELISAs were performed to assess the amount of cytokines released by CAR T cells during co-incubation with RMS target cells. 25'000 RMS target cells were seeded 24h in advance in transparent 24-well plates (Greiner bio-One, #662160), and CAR T cells were co-cultured at the Effector:Target (E:T) ratios of 1:1, 5:1, and 10:1, in 500µl total volume of complete RPMI-1640 medium. After 24h, the supernatant was collected, centrifuged at 600xg, transferred to a new plate, and stored at -80°C until use. ELISAs were performed to measure the release of IL-2 (Invitrogen, #88-7025-88), IFN-γ (Invitrogen, #88-7316-88), and Granzyme B (Biozol, #3485-1H-20) following the manufactures' instructions.

Antibodies

Western Blotting primary antibodies and working dilutions used were: goat anti-CD276 (1 µg/ml; R&D systems, #AF1027); mouse anti-α-tubulin (1:1'000; Cell Signaling, #2144); rabbit anti-MycTag (1:1'000; Cell Signaling, #2278S); mouse anti-human FGFR4 (Santa Cruz, #SC-136988). Secondary antibodies: anti-goat IgG, HRP-linked (1:10'000 Cell Signaling, #HAF109); anti-mouse IgG, HRP-linked (1:10'000; Cell Signaling, #7076); anti-rabbit IgG HRP-linked (1:10'000; Cell Signaling, #7074P2). **Flow Cytometry** primary antibodies and working dilutions: BV421-CD19 antibody (BioLegend, #115537); PE-CD276 (1:100; R&D, #AF1027); PE-FGFR4 (1:100; Biolegend, #324306); CARs expression on CAR T cells was assessed with PE-MycTag (Cell Signaling, #64025S). CD8/CD4 T cells ratio was assessed with BD Multitest CD3/CD8/CD45/CD4 (BD Biosciences, #342317). APC-CD69 (BD Biosciences, #555533); BV421-CD62L (BioLegend, #304828); BV510-CD3 (BioLegend, #563109); BV650-CCR7 (BioLegend, #353234); PerCP-CD4 (BioLegend, #300528); PE-CD95 (FAS) (BioLegend, #305608); PE-Cy7-CD45RA (BioLegend, #304108). UltraComp eBeads™ compensation beads (Thermofisher, #01-2222-41) were used for compensation. CD276 and FGFR4 surface copy numbers was calculated with the PE Fluorescence Quantitation Kit (BD Biosciences, #340495) according to the manufacturer's protocol. **Immunohistochemistry**

primary antibodies and working dilutions: mouse monoclonal anti-L1CAM (Sigma-Aldrich, #L4543, dilution 1:50), mouse monoclonal anti-human FGFR4 (Santa Cruz, #SC-136988), goat polyclonal anti-CD276 (1 μg/ml; R&D systems, #AF1027), rabbit polyclonal anti-JAM-3 (Sigma-Aldrich, #HPA003417, dilution 1:20), rabbit polyclonal anti-ADGRL2 (Cusabio, #PA013060ESR2HU, dilution 1:100), rabbit monoclonal anti-CD3 (Abcam, #16669, dilution 1:200). **Immunohistochemistry secondary** antibodies and working dilutions: polyclonal goat anti-mouse, (DAKO, #P0447), polyclonal rabbit anti-goat (DAKO, #P016002, dilution 1:200), polyclonal goat anti-rabbit, (DAKO, #P0448, dilution 1:100). The protein expression was detected using the liquid DAB$^+$ substrate chromogen system (DAKO, #K3468).

Immunohistochemistry

Immunohistochemistry analyses were performed on freshly cut, formalin-fixed, and paraffin-embedded tissue slides. Briefly, deparaffinized and rehydrated tissues were treated with 10mM sodium citrate buffer (pH 6) for antigen retrieval. The samples were next blocked with 10% goat or rabbit serum (depending on the host) and 1% BSA for 1h, before being incubated overnight at 4°C with the corresponding primary antibody. Endogenous peroxidase activity was blocked with peroxidase block (DAKO, #K1492). The tissues were incubated with the corresponding secondary antibody for 1h at room temperature, tissues were counterstained in hematoxylin, dehydrated and mounted with Eukitt mounting media (Sigma-Aldrich, #03989). Images were captured with Leica Microsystem DM6000 B microscope.

Flow Cytometry

Cell staining was performed in 100 μL FACS buffer (PBS + 2% BSA), by diluting the corresponding primary antibodies following the optimized concentrations described above. Flow cytometry experiments were evaluated with CytoFLEX device (Beckman Coulter). The results were analyzed by FlowJo v10.8.1 Software (BD Life Sciences).

Mice

NOD.Cg-PrkdcscidlL2rgtm1Wjl/SzJ (NSG) (4-6 weeks old) immunodeficient female mice were purchased from Charles River (Germany) and housed at the Central Animal Facility of the University of Bern, under pathogen-free conditions. The experiments were approved by Animal Welfare Office of Canton Bern and in accordance with Swiss Federal Law (Authorization BE22/2022). After 2 weeks of housing, mice were injected in the left

gastrocnemius muscle with 50µl of iRFP720$^+$fLuc$^+$ Rh4, RD, or JR cells (5x10^5 cells) in sterile PBS. Four days after the injection of iRFP720$^+$fLuc$^+$ RMS cells, the size of tumors was estimated by bioluminescence imaging with the In Vivo Imaging System (IVIS). On day 5, 5x10^6 CAR T cells per mouse were resuspended in 100 µl sterile PBS and intravenously injected into the tail vein. Tumor size was measured by using the following formula: Volume = ((short length)^2 * (long length))/2.

Protein isolation from tumors

Expression of CD276 and FGFR4 on RMS cells derived from xenografted tumors was evaluated by WB. Tumors were excised on day 60, samples were snap-frozen and stored at -80°C until use. Tumors were then thawed, chopped, shaken at 4°C in RIPA buffer (Thermofisher, #89900) with addition of Halt™ Protease Inhibitor Cocktail (Thermofisher, #78430). Samples were further homogenized with 1 min sonication (10% duty cycle, Branson Sonifer 250) and centrifuged at 10'000xg for 15min, 4°C. Protein quantification and WB were performed by following the protocol reported above.

Single cells isolation from tumors

Expression of CD276 and FGFR4 on RMS cells derived from xenografted tumors was evaluated by Flow Cytometry. Tumors were chopped and incubated in Balanced Salt Solution (HBSS; Thermofisher, #88284) with addition of collagenase III (200 U/ml or 1 mg/ml, STEMCELL, #07422) and DNAse I (200 U/ml, Thermofisher, #18047019) for 1 h, 37°C. Cells were filtered with 40 µm-strainer and red blood cells were lysed by incubation with ACK lysis buffer (10x: 1.5M NH$_4$Cl (Sigma, #213330), 0.1M KHCO$_3$ (Sigma, #237205), 0.001M Disodium EDTA (Thermofisher, #AM9260G) for 5 min, room temperature. Samples were then centrifuged at 400xg for 5 min and washed twice before following the Flow Cytometry protocols as described above.

Results

Expression of CD276 in RMS cell lines and PDXs

CD276-CAR T cells have shown excellent activity against several pediatric tumors but have not been tested against RMS (62). In order to verify CD276 expression levels in RMS, we selected six FP-RMS cell lines (JR, RMS, Rh4, Rh5, Rh28, Rh30), five FN-RMS cell lines (RD, Rh18, Rh36, TTC-442, RUCH-3), and three PDXs (IC-pPDX-104, IC-pPDX-35, RMS-

ZH003), and quantified total CD276 expression by WB (Fig. 1A) and cell surface expression by FACS (Fig. 1B-E). WB analysis revealed high total expression levels of CD276 in RMS PDXs, and in most RMS cell lines, with the exception of JR, RMS, and Rh18, where lower expression levels were observed. WB confirmed low expression in control myoblasts, PBMCs, and T cells (Fig. 1A). Flow Cytometry revealed high surface expression of CD276 in all the samples (Fig. 1B and 1D) and no detectable expression in PBMCs and T cells controls. Interestingly, lower levels of surface CD276 were observed for JR, RMS, and Rh18, in agreement with WB results. The number of CD276 on the cell surface were estimated with PE-beads. The most representative RMS cell lines RD, Rh30, and Rh4, display around 50'000 copies on their surface. Consistently with the previous analyses, JR, RMS, and Rh18 displayed the lowest number of copies of CD276 on the surface, with ca. 7'000, 10'000, and 25'000 copies, respectively (Fig. 1C and 1E). These experiments confirmed consistent expression of CD276 on RMS cell lines and PDXs, and support their targeting by CD276-CAR T cells.

Generation of CD276-CAR constructs and validation

Aiming at comparing different CAR constructs to identify the CAR conferring the highest cytotoxicity to CAR T cells targeting RMS, we generated a panel of second- and third-generation CARs with HD and TM deriving from either CD8α or CD28, intracellular CSDs from 4-1BB and/or CD28, and CD3z as SD, as schematically summarized in Fig. 2A. By Gibson assembly, we initially produced eight modular CAR plasmids armed with the CD19 targeting FMC63 scFv used in the clinic and regarded as the golden standard (CD19.I-VIII, Fig. 2B). These eight CD19-CAR plasmids were as negative controls, or as positive control with RMS cells overexpressing a truncated CD19. CD276-CARs were generated by swapping the ABD with the CD276.MG scFv sequence (62). A summary of the sixteen CAR plasmids can be found in Table S3. Next, in order to validate correct expression of the constructs, Jurkat cells were transduced with lentiviruses coding for the different constructs, and both correct size and surface expression of the CARs were assessed on CAR Jurkat cells by WB and Flow Cytometry, respectively. Anti-MycTag detection revealed high expression levels at the expected size for all the CARs (ca. 60 kDa), showing a clear shift from second- to third-generation CARs, due to the additional CSDs (Fig. S1A). Flow Cytometry on live cells confirmed high detection of MycTag on the surface of CAR Jurkat cells (Fig. S1B).

Manufacturing and characterization of CD276 CAR T cells

T cells isolated from PBMCs were seeded on day 1 in XF expansion medium, supplemented with IL-2 and CD3/CD28 activators, as schematically represented in Fig. 3A. Expansion, viability, and CD4:CD8 ratios of T cells were monitored for 14 days prior to both *in vitro* and *in vivo* experiments. Initially, activation of T cells was monitored over one week by measuring CD69 expression. The highest levels of CD69 expression were observed four days after activation (Fig. S2A). T cells were therefore transduced with lentiviruses on day 4 and the efficiency was measured after 48h by GFP and MycTag detection by FACS. Infection efficiency was over 40% for most of the CD276 constructs (mean of 55.6% ± 12.1, Fig. S2B). In order to normalize the expression of the CAR constructs, and to allow a precise comparison of their activity, CAR T cells were then sorted by GFP on day 7. After 14 days in culture, all the CAR T cells showed an up to 35-fold in vitro expansion (Fig. 3B) and a viability of more than 70%. In the moment of the co-incubation with target cells, the CD4:CD8 ratio of the amplified CAR T cells was around 1:1 (Fig. 3C). CARs size and expression on the surface of CAR T cells were assessed by WB and Flow Cytometry based on MycTag detection. This confirmed the correct expression at ca. 60kDa and different size between the second- and third-generation constructs for CD19-CARs (Fig. 4A), and CD276-CARs (Fig. 4C), and high CAR expression levels were observed in all the samples (Fig. 4B and 4D).

Phenotype of CAR T cells

In order to determine the phenotype of CAR T cells enriched during expansion, and to assess any change in T cell subsets due to culture conditions or upon co-incubation with tumor cells, T cell phenotype was investigated before and after co-incubation with tumor cells. On day 0, T cells were isolated from PBMCs and cultured in XF expansion medium, supplemented with IL-2 and CD3/CD28 activators. Activated T cells were compared to untreated T cells after one day (Fig. 5A: untreated T cells: upper panels; activated T cells: lower panels). To note is that a lower CD3 expression was observed in activated T cells than in non-activated controls (Fig. 5A, left panels), probably due to CD3 sequestering and masking by the CD3/CD28 activators. The analysis of $CD3^+$ T cells revealed in both groups more than 60% of $CD45RA^+/CD62L^+$ T cells (T_N) (Fig. 5A, central panels), and interestingly showed a transition into $CCR7^+/CD95^+$ stem cell-like memory T cells (T_{SCM}) only in activated T cells (Fig. 5A, right panels). The percentage of $CD45RA^-/CD62L^+$ central memory T cells (T_{CM}) was higher in non-activated T cells (12% vs 7%), while $CD45RA^+/CD62L^-$ effector T cells (T_{EFF}) were higher in activated T cells (9% vs 23%). After lentiviral transduction and sorting on day 7, as expected from removal of activators, CD3 levels in GFP^+ CAR T cells were similar to those of non-transduced T lymphocytes (Fig. 5B, left panel). On day 7, >90% of CAR T cells converted into T_{SCM} cells

(Fig. 5B, right panel). Lastly, FACS analysis performed on day 14 pre-coincubation with RMS cells, and on day 16 post-co-incubation with RMS cells, showed that ca. 70% of CAR T cells remained differentiated as T_{SCM} cells, but interestingly around 30% of CAR T cells transitioned towards a central memory phenotype (T_{CM}) (Fig. 5C, upper middle panel). To conclude, we observed an initial differentiation of activated T cells from T_N to T_{SCM} during the first 7 days of expansion. During the second week, just one third of CAR T cells converted into T_{CM} cells. However, measurements performed pre- and post-co-incubation experiments revealed no crucial phenotype alteration on CAR T cells, except for a 10% shift of T_N cells toward T_{CM} cells (Fig. 5C, middle lower panel).

CD276-CAR T cells cytotoxicity and cytokines release

The eight different CD276-CAR constructs were then tested to determine the CAR construct promoting the highest cytotoxicity *in vitro*. fLuc$^+$ RD and Rh4 cells, both expressing high levels of CD276 on their surface, were selected as the most representative cell lines for FP-RMS and FN-RMS, respectively (Fig. 1C). After 48h co-incubation, survival of RMS cells was determined by luciferase assays. At the effector to target (E:T) ratios of 5:1 and 10:1, all experimental groups efficiently killed almost 100% of fLuc$^+$ RMS cells (Fig. 6, upper panels). Nevertheless, at the lower E:T ratios we could discriminate CD276.V (276.28.28.28.3z, cyan) as the CAR showing the highest efficacy against both fLuc$^+$ RD and Rh4. CD19-CAR T cells, used here as a negative control, showed no significant killing capacity against RMS cells, whereas when co-incubated with RMS cells expressing truncated CD19 (tCD19$^+$ fLuc$^+$ RD and Rh4 cells), CD19-CAR T cells specifically eliminated most of RMS cells with high efficacy (Fig.6, lower panels).

Levels of IL-2, IFN-γ, and Granzyme B were measured to quantify cytokine release from the eight CD276-CAR T cell constructs after 24h co-incubation with RD or Rh4 cells. The results showed highest expression of IL-2 by CD276.V co-incubated with RD cells, and by CD276.I, CD276.II, and CD276.V co-incubated with Rh4 cells (Fig. 7A). These results were confirmed also in terms of IFN-γ release, with maximum values of around ~400 pg/ml released by CD276.V-CAR T cells, when co-incubated with both RD and Rh4 cells (Fig. 7B). Considering the absolute values, Granzyme B (~8000 pg/ml) was detected at higher levels than IL-2 and IFN-γ, especially when CD276.V-CAR T cells were co-incubated with RD cells (Fig. 7C). The constructs CD276.II and CD276.V confirmed to be among the most effective ones when co-incubated with Rh4 cells, releasing ~4000 pg/ml.

Expression of FGFR4 in RMS cell lines and PDXs

Antigen loss is a common mechanism of tumor resistance against CAR T cellsthat can be addressed by CAR T cells able to recognize multiple targets on the surface of tumor cells. Therefore, we next focused on FGFR4, another promising target for RMS, as also recently confirmed by our RMS surfaceome profiling (32). Expression levels of FGFR4 on RMS cell lines and PDXs were assessed by Western blotting and Flow Cytometry. Western blotting put in evidence high expression levels of FGFR4 in Rh4, Rh28, JR, and RMS cell lines, and in the PDXs RMS ZH003 and IC-pPDX-104; medium FGFR4 expression levels were instead observed in Rh36, RD, and TTC-442; whereas low levels of FGFR4 were detected in Rh18, RUCH-3, Rh5, and Rh30 cell lines, and in IC-pPDX-35 (Fig. 8A). Flow Cytometry analysis performed on live cells confirmed expression of FGFR4 on most of the RMS cell lines (Fig. 8B). Quantification of surface FGFR4 copy number revealed a relatively low copy number of FGFR4 on RMS cells, compared to CD276. Despite the lower density of FGFR4 on RMS samples compared to CD276, the absence of expression in most normal tissues makes FGFR4 a promising target in RMS for CAR T cell-based strategies (63, 64, 65).

Cytotoxicity of FGFR4- and CD276/FGFR4 Dual-CAR T cells against RMS cell lines

Having identified CD276.V as the most effective CAR construct in terms of cytotoxicity and cytokine release, and after validation of FGFR4 expression on several RMS cell lines, we generated FGFR4-CARs in the CAR.V backbone (28.28.28.3z). Four previously selected FGFR4 targeting single-domain antibodies (sdAb) A8-FR4, B1-FR4, B5-FR4, and F8-FR4 (63) were used as ABD to generate lentiviral CAR plasmids and to produce the corresponding CAR T cells. To determine the killing capacity of sdAb-FGFR4-CAR T cells and to screen for the most specific and effective construct, we performed cytotoxicity assays with fLuc$^+$ Rh4 cells. Rh4 was chosen among a panel of RMS cell lines due to the high surface expression of FGFR4 with 3'343 FGFR4 copies per cell (Fig. 8C). In order to estimate the specificity of the different sdAb, FGFR4-KO Rh4 cells were generated by CRISPR/Cas9 (Fig. S3) and were used as negative controls in cytotoxicity assays. All the four FGFR4-CAR constructs were able to trigger more than 80% tumor cells killing as CAR T cells at E:T ratios of 5:1 and 10:1. However, at lower ratios B5-FR4.V-CAR T cells showed the highest cytotoxic effect, eliminating ~40% and ~60% of Rh4 cells at the ratios of 0.5:1 and 1:1, respectively. A8-FR4.V and F8-FR4.V-CAR T cells showed a killing capacity of ~20% and ~30% respectively, whereas B1-FR4.V was the weakest in terms of cytotoxicity, with no killing at 0.5:1 (Fig. 9A, left panel). On the other hand, co-incubation of FGFR4-CAR T cells with fLuc$^+$ FGFR4 KO Rh4 cells surprisingly revealed an unspecific activity of CARs A8-FR4.V, B1-FR4.V, and B5-FR4.V, resulting in an

effective killing of FGFR4 KO Rh4. F8-FR4.V was the only CAR showing no activity against FGFR4 KO Rh4 cells (Fig. 9A, right panel). Therefore, F8-FR4.V-CAR T cells were selected for the next *in vitro* and *in vivo* experiments. In order to overcome possible antigen loss or antigen escape *in vivo* and to further improve cytotoxicity of the CAR T cells, we investigated a possible combinatorial effect of CD276.V/F8-FR4.V-Dual-CAR T cells, targeting both CD276 and FGFR4, in Rh4 cells, defined as CD276^{high}FGFR4^{high}; and JR, defined as CD276^{low}FGFR4^{high}. The results from cytotoxicity assays indicate that, when co-incubated with Rh4, Dual-CAR T cells killed Rh4 cells with high efficacy, *i.e.* more than 50% compared to the control at an E:T ratio of 0.5:1, slightly better than CD276.V-CAR T cells, suggesting a synergistic effect. By contrast, we observed only a moderate killing of Rh4 cells of around 20% by F8-FR4.V-CAR T cells (Fig. 9B, left panel). Experiments performed with JR cell line, instead, suggest a cytotoxic activity of Dual-CAR T cells analogous to the F8-FR4.V-CAR T cells, with no clear synergistic effect. CD276.V-CAR T cells showed a lower activity, likely due to the lower surface expression of CD276 in JR cells (Fig. 9B, right panel).

CD276-CAR T cells eradicate RMS tumors in an orthotopic mouse model

Next, *in vivo* experiments were performed in immunodeficient NSG mice carrying orthotopic RD-derived tumors (CD276^{high}FGFR4^{low}), with the aim to assess the *in vivo* activity of CD276.V-CAR T cells. RD was selected as a representative FN-RMS cell lines presenting high CD276 surface copy number (60'000, Fig.1C). On day 0, four treatment groups were defined: 1) PBS; 2) Untransduced T cells (referred to as UTD); 3) CD19.V-CAR T cells (referred to as CD19); 4) CD276.V-CAR T cells (referred to as CD276). We used PBS, UTD, and CD19 as negative control groups. 0.5×10^6 fLuc$^+$ RD cells were orthotopically injected into the left gastrocnemius muscle of five mice per group, and on day 4, the tumor size was estimated by bioluminescence to prove the uniform tumor growth in all the groups (Fig.10A). On day 5, 5×10^6 untransduced T cells (UTD), or 5×10^6 CAR T cells were injected intravenously. Tumor growth was then monitored weekly by bioluminescence (Fig. 10B). During the third week after RD injection, we observed RMS tumor eradication in 3/5 mice in the CD276 group. By contrast, in the three control groups RD tumors regularly grew. At the end of the experiment at week 6, fLuc$^+$ RD cells were still undetectable by bioluminescence in 3/5 CD276 mice, indicating a maintained complete remission, whereas in the other two CD276 mice the tumor growth was controlled until week 6. This experiment revealed a promising activity of CD276.V-CAR T cells in promoting clearance of orthotopic RMS tumor. No drop in mouse weight, nor evident side effects were observed in the CD276 treated group. Notably, alopecia-like phenotype appeared only in the CD19.V-CAR T cells group during week 5 and persisted until the end of

the experiment. All the controls showed a similar tumor growth during the whole experiment. These results revealed a good activity of CD276.V-CAR T cells against CD276[high] FN-RMS cells, leading to RMS eradication in 3/5 mice and growth reduction in 2/3.

CD276- and CD276/F8-FR4 Dual-CAR T cells induce RMS tumor clearance

Next, we tested in immunodeficient NSG mice the activity of CD276.V-CAR T cells against orthotopic tumors derived from Rh4 cells (CD276[high]FGFR4[high]) expressing high levels of CD276 (50'259, Fig. 1C) and FGFR4 (3'343, Fig. 9C). Here, we also tested F8-FR4.V-CARs against FGFR4, as well as Dual-CAR T cells targeting both CD276 and FGFR4. On day 0, five treatment groups were defined: 1) Untransduced T cells (referred to as UTD); 2) CD19.V-CAR T cells (referred to as CD19); 3) CD276.V-CAR T cells (referred to as CD276); 4) F8-FR4.V-CAR T cells (referred to as F8-FR4); 5) CD276/F8-FR4 Dual-CAR T cells (referred to as Dual). On day 0, 0.5×10^6 fLuc$^+$ Rh4 cells were engrafted into NSG mice and, on day 5, 5×10^6 untransduced (UTD) T cells, and 5×10^6 CAR T cells were intravenously injected in the mice (Fig.11A). At week 2, 2/5 mice from the CD276 group and 2/5 from the Dual group showed a significantly smaller tumor burden, compared to the controls (Fig.11B). At week 3, all the mice of both CD276 and Dual groups presented a complete remission, which lasted until the end of the experiment (day 59). The F8-FR4 group exhibited a delayed tumor growth in 4/5 mice, compared to the controls until week 6 of the treatment, but failed to completely eradicate the tumors. In the control mice, instead, rapid growth of Rh4-derived xenograft was observed from week 3, so that CD19 mice had to be sacrificed on day 39, due to high tumor burden (>800 mm3 tumor size, Fig.11C). For the same reason, after 53 days, all the control and the F8.FR4.V mice had to be sacrificed. At the end of the experiment (day 59), all the mice from CD276.V and Dual-CAR T cells groups were tumor free (Fig. 11D) and did not show any obvious side effect. No drop in mouse weight was observed in the treated groups. These results confirm the excellent activity of CD276.V-CAR T cells against CD276[high] tumors, leading to complete sustained remission in 5/5 mice, while F8-FR4.V-CAR T cells showed a limited activity, leading to growth delay in 4/5 mice. Interestingly, Dual-CAR T cells showed the same activity as CD276.V-CAR T cells, completely eliminating tumors in 5/5 mice.

Limited eradication in CD276[low] RMS tumors

In order to test CD276-CAR T cells activity *in vivo* against tumor cells expressing lower levels of CD276, and to investigate a potential synergistic effect promoted by CD276/F8-FR4.V Dual-CAR T cells, the same experimental groups of CAR T cells described above (Fig. 11)

were investigated in JR cells (CD276^{low}FGFR4^{high}) expressing low levels of CD276 (7'964, Fig. 1C) and medium-high levels of FGFR4 (2'245, Fig. 9c). On day 0, six treatment groups were defined: 1) Untreated (referred to as PBS); 2) Untransduced T cells (referred to as UTD); 3) CD19.V-CAR T cells (referred to as CD19); 4) CD276.V-CAR T cells (referred to as CD276); 5) F8-FR4.V-CAR T cells (referred to as F8-FR4); 6) CD276/F8-FR4 Dual-CAR T cells (referred to as Dual) (Fig.12A). We used CD19 as negative control group. On day 0, 0.5x10^6 fLuc$^+$ JR cells were engrafted into NSG mice and, on day 5, 5x10^6 untransduced (UTD) T cells, and 0.5x10^6 CAR T cells, were intravenously injected into the mice. During week 2, bioluminescence imaging revealed a reduction in tumor size in 1/5 mice from the CD276 group and in 1/5 mice from the F8-FR4 group, compared to the controls (Fig. 12B). However, from week 3, only one mouse (nr. 2) from the CD276 group exhibited a complete remission, which lasted until the end of the experiment (Fig. 12B and 12C). Unfortunately, none of the mice from the other experimental groups showed tumor eradication, nor a significantly delay in tumor growth. Instead, all the mice showed a rapid and homogenous tumor increase (Fig. 12C), and the mice were euthanized during week 6, having reached the termination criteria. We did not observe any drop in mouse weight, except for one mouse (nr. 3) of the PBS group, which was sacrificed once it lost 15% of the initial weight (Fig. 12D). Mice of the CD19 group exhibited fur loss on the back of the neck, but no further side effects were observed. These results revealed the limit of CD276.V-CAR T cells against tumors expressing lower levels of CD276 and confirmed the limited activity of F8-FR4-CAR T cells. Disappointingly, no synergistic effect could be achieved by Dual-CARs.

Decreased expression of CD276 and FGFR4 in tumors compared to cultured cells

To investigate if expression of the CAR targets in orthotopic tumors correspond to what observed on cultured cell lines, CD276 and FGFR4 expression was assessed by Flow Cytometry on single cells extracted from Rh4 tumors (Fig. 13), Flow cytometry results showed a lower CD276 expression on tumor xenografts-derived Rh4 cells, with a fluorescence intensity 14-fold higher than control, compared to cultured Rh4 cells (600-fold over control) (Fig. 13A), while a much lower expression of FGFR4 was observed in tumor xenografts-derived Rh4 cells (2-fold over isotype control) compared to cultured wild-type Rh4 cells (97-fold over control) (Fig. 13B). An estimation of the magnitude of the decrease of CD276 and FGFR4 based on the fluorescence intensity (geometric mean) indicates a similar decrease in tumors, 40-fold (CD276) and 50-fold (FGFR4), compared to cultured cells.

To confirm these results, IHC staining of CD276 and FGFR4 was performed on formalin-fixed and paraffin-embedded sections of RD-, Rh4-, and JR-derived tumor xenografts from

control mice (Fig. 13C). IHC staining results on tumors confirmed uniform and higher expression of CD276 in RD and Rh4 derived tumors, compared to JR derived tumors (Fig. 13C). FGFR4 was detected at high levels in Rh4-, at medium levels in JR-, and at low levels in RD-derived tumors (Fig.13C). Taken together, these results indicate that both targets are clearly detectable in Rh4 mouse tumor xenografts, but they are expressed at different levels in RD and JR. In conclusion, CAR targets seem to be expressed in tumors at lower levels compared to cultured cell lines, but their relative expression seems to be maintained.

CAR T cells persistence in mice spleen

CAR T cells persistence *in vivo* is an important aspect that determines the therapeutic impact. To investigate this, we performed CD3 staining on JR-derived tumors collected from CD276.V-CAR T cells treated mice at the end of the experiment, after 5 weeks from tumor cells implantation, and from Rh4-derived tumors collected from F8-FR4.V-CAR T cells treated mice at the end of the experiment, after 7 weeks from tumor cells implantation. IHC CD3 staining revealed the presence of CAR T cells in both tumors (Fig. 14A). In JR tumors CAR T cells were visible only in structures that appear to be vessels (Fig. 14A, black arrowheads) and CAR T cells seem not to have penetrated the tumor tissue, while in Rh4 tumors CAR T cells appeared to have infiltrated in the tumor. Since Rh4-derived tumors were cleared by CD276.V- and Dual-CAR T cells, we investigated if persistent CAR T cells could be still detected in the spleen of mice treated with CD276.V-CAR T cells. IHC analyses detected T cells in the spleen of Rh4-bearing mice injected with control untransduced T cells (UTD control) and in the spleen of CD276.V-CAR T cells treated mice, suggesting persistence of CAR T cells even after tumor eradication. IHC analyses on normal tissues from control mice and from Rh4-tumor bearing mice were performed to evaluate any sign of toxicity caused by CAR T cells. Hematoxylin and Eosin staining in liver, muscle, brain, heart, and spleen of untreated and treated mice showed similar phenotype (Supplementary Fig. S4). No evident necrosis nor inflammation were observed in the tissues analyzed, suggesting that CD276.V- and Dual-CAR T cells specifically recognize the targets of interest without damaging the investigated normal mouse tissues.

In conclusion, these results indicate that even after unsuccessful treatment, both CD276.V- and F8-FR4-CAR T cells can be detected in JR tumors and suggest that CD276-CAR T cells might be limited in reaching tumors cells. In Rh4-bearing mice, where tumors could be eradicated, CAR T cells could be detected even several weeks after tumor clearance, indicating a good persistence potential.

Discussion

Here, we investigated the impact of specific domains included in the CAR constructs and adopted a pragmatic approach, by *in vitro* screening of a combinatorial pool of eight CD276-directed CAR constructs, with CD8α or CD28 HD/TM domains and one or two intracellular CSDs from 4-1BB and/or CD28. CD276.V-CAR T cells (276.MG.28HD/TM.28CSD.3z), were the most effective in killing FN-RMS RD and FP-RMS Rh4 cells *in vitro* and released the highest levels of IFN-γ and Granzyme B. CD276.V-CAR T cells were able to eradicate CD276^{high} Rh4 tumors *in vivo* and persisted even after tumor elimination.

These results confirm recent studies demonstrating that CAR T cells with CD28 HD/TM domains show more reactivity to low-antigen density tumors, faster killing, and higher levels of IL-2 release when compared to CARs with CD8α HD/TM domains (51). Superior anti-tumor efficacy of CD28 HD/TM was also reported in HER2-positive breast cancer and CD276-positive neuroblastoma (51). On the other side, CD8α HD/TM domains are associated with a lower production of cytokines, such as IL-2, IFN-γ, and TNF-α, when compared to CD28 HD/TM domains. However, despite these reduced levels, CD8α-mediated cytokine-release syndrome (CRS) was still observed in patients (66). In this context, we chose the killing capacity of CAR T cells and the release of cytokines such as IFN-γ, as predictive readouts, due to the important role played by IFN-γ in targeting solid tumors (67).

From investigations of different co-stimulatory domains (CSDs) in mouse models, we know that CD28 CSD is associated with a higher release of inflammatory cytokines and with an exhausted phenotype (56). Moreover, an augmented CRS occurrence was observed in patients treated with CD19-CD28 CAR T cells, compared to CD19-4-1BB CAR T, highlighting a more cytotoxic effect by CD28 CSD (68, 69). Our results are in good agreement with a stronger activity conferred by CD28 CSD.

Our results on successful use of CD276.V-CAR T in CD276^{high} RMS tumors are in line with the results obtained with other CAR constructs targeting CD276 in preclinical studies against several types of tumors: 276.8αHD/TM.28CSD.BBCSD.3ζ against glioblastoma (70); 276.MG.8αHD/TM.BBCSD.3ζ against medulloblastoma (62), neuroblastoma (62, 71), osteosarcoma (62), and Ewing sarcoma (62); 276.8H9S3.3.8α.28TM.BBCSD.3ζ against gastric cancer (72); and 276.8αHD.28TM.28CSD.3ζ and 276.8αHD/TM.BBCSD.3ζ against esophagus squamous cell carcinoma (73). All these CAR constructs generally allowed T cells to increase cytokine release *in vitro*, to prolong survival, and to persist *in vivo*.

In our screening, second-generation CAR T cells with one CSD, outperformed third-generation CAR T cells with two CSD. This result was surprising, since several preclinical

studies with third-generation CD19-CAR T cells hinted at a higher intracellular signaling activity compared to the second-generation CAR constructs (74). Co-expression of both CD28 and 4-1BB CSDs was reported to compensate the possible limitations of single CSD, improving expansion and showing superior persistence (75). For these reasons, we expected that third-generation CAR T cells would show a more effective killing capacity and a higher cytokine release. On the other side, some studies revealed lower surface expression of third-generation CARs and inferior persistence, compared to second-generation (76).

We established a CAR T cell manufacturing protocol leading to a stem-like memory T cells (T_{SCM}) phenotype in the majority of CD276.V-CAR Ts. *In vitro* analyses pre- and post-co-incubation with tumor cells confirmed persistent expression of T_{SCM} markers in ̴60% of CAR T cells. T_{SCM} cells represent the least differentiated grade of memory T cells and have self-renewal ability and anti-tumor ability (77, 78). Using T_{SCM} CAR T cells can be beneficial, since it was demonstrated that they can persist in patients for up to 12 years and present a good safety profile (79). Recent studies have also observed that T_{SCM} play a central role in the early anti-leukemia response and late immune surveillance (80). Therefore, enriching CAR T cells with a high percentage of T_{SCM} cells should represent an advantage for CAR T cells *in vivo*.

In vivo, CD276.V-CAR T cells successfully eradicated Rh4-derived tumors at week 2 of treatment with 100% efficiency whereby in 5/5 mice the tumor did not relapse. The efficacy against RD-derived tumors was of around 60%, with 3/5 mice showing a clear maintained complete remission. Notably, the other two mice showed an initial significant treatment response, but not sufficient to achieve a complete remission. Since Rh4 and RD cells lines show similar CD276^{high} surface density *in vitro*, a plausible explanation of the different efficacy might be found in the different growth rate: slow growing FN-RMS RD possibly impede CD276-CAR T cells activity due to the small size of the RD-derived tumors that might not provide sufficient stimulation to activate CAR T cells and to allow their expansion and persistence. In fact, RD tumors could only be monitored by bioluminescence, and not measured by caliper. This might theoretically also reduce homing and decrease the probability by CD276-CAR T cells to encounter the target, protecting RD tumor cells during the first week of experiment. Future experiments with either implantation of more RD cells, or with CAR T cells injection at a later time point with a larger RD-tumor size, might answer this question. CD276.V-CAR T cells failed to control growth ofCD276^{low} JR-derived tumors.

CD276.V-CAR T cells were detectable in tumors after 5 weeks, indicating that a lack of persistence might not be the reason for the limited activity. This lack of activity was disappointing, since CD276.V killing of JR cells was comparable to Rh4 and RD cells *in vitro*, but not surprising, since the dependence of CARs on antigen density for effective activity has

already been recognized (51, 62, 81). On the same line, F8-FR4-CAR T cells failed to eradicate Rh4-derived tumors and the strongest effect observed was just a 1-week delay in tumor expansion. F8-FR4-CAR T target FGFR4 which is expressed at lower levels than CD276 on Rh4 cells, with ~3'000 FGFR4 copies per cell vs ~50'000 of CD276. Therefore, a possible explanation for the lack of tumor eradication might simply be an insufficient activation of F8-FR4-CAR T cells. However, recent work published by Sullivan et al. (47) showed that a combination of FGFR4-CAR T cells with pharmacologic inhibition of the myeloid component in stroma in Rh30 RMS-derived tumors allowed FGFR4-CAR T cells to eradicate orthotopic RMS in mice (47). It is interesting to note that Rh30 cells were reported to express 2'700 FGFR4 molecules per cell, while we measured 2'000 molecules/cell in the Rh30 cell line in our laboratory and 3'700 molecules per cell in Rh4 cells. These results suggest that it is possible to overcome the limitations posed by a low-density target by manipulating the myeloid compartment in RMS. Alternatively, expression of c-Jun gene in F8-FR4.V-CAR T cells, which have the the same CAR backbone used to target low GD2 expressing cells, could lead to *in vivo* eradication of tumors with low antigen density without side effects (82).

To increase CAR T cells activity, and to overcome a possible antigen loss, we investigated a Dual-CAR T cell strategy by targeting CD276 and FGFR4 at the same time. Dual-CAR T cells co-expressing on their surface equivalent numbers of CD276.V-CARs and F8-FR4.V CARs were tested *in vitro* against Rh4 cells showing a slightly better killing capacity compared to CD276.V-CAR T cells, while a lower cytotoxicity was observed for F8-FR4.V CAR T cells. In Rh4-tumor bearing mice, CD276/FGFR4 Dual-CAR T cells showed a similar complete eradication as CD276.V-CAR T cells. This is an important observation in the optic of combining CD276 with other CARs, since it implies that they are still fully active even if the number of CD276 CARs on the surface of T cells is reduced,. Unfortunately, Dual-CAR T cells failed to show a synergistic effect, and were not effective against CD276^{low}FGFR4^{high} JR cells, just like F8-FR4.V- and CD276.V-CAR T, indicating a direct correlation between target surface density and efficacy of CAR T cells. In fact, despite the high efficacy shown with RD and Rh4, CD276.V-CAR T cells exhibited reduced killing capacity against JR.

Currently, there are three ongoing CD276 clinical trials: NCT04691713 is recruiting patients with advanced CD276-positive solid tumors; NCT04432649 is a Phase 2 study investigating the activity of 4SCAR-276 CAR T cells in brain tumors and Ewing sarcoma; NCT04185038 is a Phase I clinical trial for Diffuse Intrinsic Pontine Glioma/Diffuse Midline Glioma and recurrent or refractory pediatric central nervous system tumors. Our results support the inclusion of patients with CD276-positive RMS in current clinical trials.

Conclusions

CD276.V-CAR T cells containing CD28-derived HD/TM and CSD are able to control and eradicate orthotopic RMS tumors, when their target is expressed above a critical threshold. FGFR4-targeted CAR T cells can only partially control RMS growth, possibly because of low target expression and immunosuppressive environment. CD276/FGFR4 Dual-CAR T cells showed the same activity as CD276-CAR T cells, suggesting that dilution of CD276-CAR does not detrimentally impact their activity. Targeting multiple antigens, increasing signal strength through optimized CAR design, remedying the limitations imposed by the immunosuppressive TME will be necessary to achieve consistently complete remission.

List of abbreviations

ABD: antigen binding domain; AICD: activation-induced cell death; APC: antigen presenting cells; aRMS: alveolar RMS; CAR: chimeric antigen receptor; eRMS: embryonal RMS; FN-RMS: fusion negative RMS; FP-RMS: fusion positive RMS; HD: hinge domain; IHC: immunohistochemistry; PDX: patient derived xenograft; RMS: rhabdomyosarcoma; sdAb: single-domain antibodies; TM: transmembrane domain; TME: tumor microenvironment; UTD: untreated; WB: Western blotting.

Declarations

Ethics approval and consent to participate

The mice experiments were approved by Animal Welfare Office of Canton Bern and in accordance with Swiss Federal Law (Authorization BE22/2022).

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests. JR is currently an employee of Novartis Pharma, Basel.

Funding

This research was funded by the "Bernese Foundation for Children and Young Adults with Cancer / Berner Stiftung für krebskranke Kinder und Jugendliche", the "Stiftung für klinisch-experimentelle Tumorforschung / Foundation for Clinical-Experimental Cancer Research" to M.B., and by a research award by the "Childhood Cancer Switzerland" to A.T.

Authors' contributions

AT: Design and execution of experiments, supervision, analysis, interpretation of data, writing, and funding acquisition. CP: Design and execution of experiments, analysis and interpretation of data, critical reading of the manuscript. DD: Execution of experiments, analysis and interpretation of data. AR: Execution of experiments, analysis and interpretation of data. JM: Execution of experiments, analysis and interpretation of data. JR: Supervision, funding acquisition, and critical reading of the manuscript. MB: Design of the study, interpretation of data, supervision, writing, and funding acquisition. All authors read and approved the final manuscript.

Acknowledgements

We thank the Institute of Myology, Paris France, for providing humanized myoblasts. We are also very grateful to Dr. Stephan Müller and the Flow Cytometry and Cell Sorting (FCCS) facility of the DBMR (University of Bern) for their support. We are grateful to Prof. Andreina Schöberlein and her lab for their help with immunohistochemistry.

Authors' information (optional)

References

1. Skapek SX, Ferrari A, Gupta AA, Lupo PJ, Butler E, Shipley J, et al. Rhabdomyosarcoma. Nature Reviews Disease Primers. 2019;5(1):1.
2. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer J Clin. 2022;72(1):7-33.
3. Fletcher CDM, Baldini EH, Blay JY, Gronchi A, Lazar AJ, Messiou C, et al. WHO Classification of Tumours. Soft Tissue and Bone Tumours. Board TWCoTE, editor. Lyon: IARC Press; 2020. 201-15 p.

4. Parham DM, Barr FG. Classification of rhabdomyosarcoma and its molecular basis. Adv Anat Pathol. 2013;20(6):387-97.

5. Davis RJ, D'Cruz CM, Lovell MA, Biegel JA, Barr FG. Fusion of PAX7 to FKHR by the variant t (1; 13)(p36; q14) translocation in alveolar rhabdomyosarcoma. Cancer research. 1994;54(11):2869-72.

6. Galili N, Davis RJ, Fredericks WJ, Mukhopadhyay S, Rauscher FJ, 3rd, Emanuel BS, et al. Fusion of a fork head domain gene to PAX3 in the solid tumour alveolar rhabdomyosarcoma. Nat Genet. 1993;5(3):230-5.

7. Barr FG, Galili N, Holick J, Biegel JA, Rovera G, Emanuel BS. Rearrangement of the PAX3 paired box gene in the paediatric solid tumour alveolar rhabdomyosarcoma. Nat Genet. 1993;3(2):113-7.

8. Charytonowicz E, Cordon-Cardo C, Matushansky I, Ziman M. Alveolar rhabdomyosarcoma: is the cell of origin a mesenchymal stem cell? Cancer Lett. 2009;279(2):126-36.

9. Hatley ME, Tang W, Garcia MR, Finkelstein D, Millay DP, Liu N, et al. A mouse model of rhabdomyosarcoma originating from the adipocyte lineage. Cancer Cell. 2012;22(4):536-46.

10. Rubin BP, Nishijo K, Chen HI, Yi X, Schuetze DP, Pal R, et al. Evidence for an unanticipated relationship between undifferentiated pleomorphic sarcoma and embryonal rhabdomyosarcoma. Cancer Cell. 2011;19(2):177-91.

11. Drummond CJ, Hanna JA, Garcia MR, Devine DJ, Heyrana AJ, Finkelstein D, et al. Hedgehog Pathway Drives Fusion-Negative Rhabdomyosarcoma Initiated From Non-myogenic Endothelial Progenitors. Cancer Cell. 2018;33(1):108-24 e5.

12. Haduong JH, Heske CM, Allen-Rhoades W, Xue W, Teot LA, Rodeberg DA, et al. An update on rhabdomyosarcoma risk stratification and the rationale for current and future Children's Oncology Group clinical trials. Pediatric blood & cancer. 2022;69(4):e29511.

13. June CH, Sadelain M. Chimeric Antigen Receptor Therapy. N Engl J Med. 2018;379(1):64-73.

14. June CH, O'Connor RS, Kawalekar OU, Ghassemi S, Milone MC. CAR T cell immunotherapy for human cancer. Science. 2018;359(6382):1361-5.

15. Melenhorst JJ, Chen GM, Wang M, Porter DL, Chen C, Collins MA, et al. Decade-long leukaemia remissions with persistence of CD4+ CAR T cells. Nature. 2022;602(7897):503-9.

16. Zhang X, Zhu L, Zhang H, Chen S, Xiao Y. CAR-T Cell Therapy in Hematological Malignancies: Current Opportunities and Challenges. Front Immunol. 2022;13:927153.

17. Sterner RC, Sterner RM. CAR-T cell therapy: current limitations and potential strategies. Blood Cancer J. 2021;11(4):69.

18. Hong M, Clubb JD, Chen YY. Engineering CAR-T Cells for Next-Generation Cancer Therapy. Cancer Cell. 2020;38(4):473-88.

19. Hegde M, Moll AJ, Byrd TT, Louis CU, Ahmed N. Cellular immunotherapy for pediatric solid tumors. Cytotherapy. 2015;17(1):3-17.

20. Dyson KA, Stover BD, Grippin A, Mendez-Gomez HR, Lagmay J, Mitchell DA, et al. Emerging trends in immunotherapy for pediatric sarcomas. J Hematol Oncol. 2019;12(1):78.

21. Bernasconi M, Dzhumashev D, Timpanaro A, Rössler J. Targeted therapies for rhabdomyosarcoma. Schweizer Krebsbulletin. 2019;39(June 2019):129-33.

22. Gattenloehner S, Vincent A, Leuschner I, Tzartos S, Muller-Hermelink HK, Kirchner T, et al. The fetal form of the acetylcholine receptor distinguishes rhabdomyosarcomas from other childhood tumors. Am J Pathol. 1998;152(2):437-44.

23. Rossig C, Brenner MK. Genetic modification of T lymphocytes for adoptive immunotherapy. Mol Ther. 2004;10(1):5-18.

24. Gattenlohner S, Marx A, Markfort B, Pscherer S, Landmeier S, Juergens H, et al. Rhabdomyosarcoma lysis by T cells expressing a human autoantibody-based chimeric receptor targeting the fetal acetylcholine receptor. Cancer Res. 2006;66(1):24-8.

25. Huang X, Park H, Greene J, Pao J, Mulvey E, Zhou SX, et al. IGF1R-and ROR1-specific CAR T cells as a potential therapy for high risk sarcomas. PLoS One. 2015;10(7):e0133152.
26. Xiao W, Wang J, Wen X, Xu B, Que Y, Yu K, et al. Chimeric antigen receptor-modified T-cell therapy for platelet-derived growth factor receptor alpha-positive rhabdomyosarcoma. Cancer. 2020;126 Suppl 9:2093-100.
27. Karkampouna S, La Manna F, Benjak A, Kiener M, De Menna M, Zoni E, et al. Patient-derived xenografts and organoids model therapy response in prostate cancer. Nat Commun. 2021;12(1):1117.
28. Merker M, Pfirrmann V, Oelsner S, Fulda S, Klingebiel T, Wels WS, et al. Generation and characterization of ErbB2-CAR-engineered cytokine-induced killer cells for the treatment of high-risk soft tissue sarcoma in children. Oncotarget. 2017;8(39):66137-53.
29. Ganti R, Skapek SX, Zhang J, Fuller CE, Wu J, Billups CA, et al. Expression and genomic status of EGFR and ErbB-2 in alveolar and embryonal rhabdomyosarcoma. Mod Pathol. 2006;19(9):1213-20.
30. Armistead PM, Salganick J, Roh JS, Steinert DM, Patel S, Munsell M, et al. Expression of receptor tyrosine kinases and apoptotic molecules in rhabdomyosarcoma: correlation with overall survival in 105 patients. Cancer. 2007;110(10):2293-303.
31. Navai SA, Derenzo C, Joseph S, Sanber K, Byrd T, Zhang H, et al. Abstract LB-147: Administration of HER2-CAR T cells after lymphodepletion safely improves T cell expansion and induces clinical responses in patients with advanced sarcomas. Cancer Research. 2019;79(13 Supplement):LB-147.
32. Timpanaro A, Piccand C, Uldry AC, Bode PK, Dzhumashev D, Sala R, et al. Surfaceome Profiling of Cell Lines and Patient-Derived Xenografts Confirm FGFR4, NCAM1, CD276, and Highlight AGRL2, JAM3, and L1CAM as Surface Targets for Rhabdomyosarcoma. Int J Mol Sci. 2023;24(3).
33. Chapoval AI, Ni J, Lau JS, Wilcox RA, Flies DB, Liu D, et al. B7-H3: a costimulatory molecule for T cell activation and IFN-gamma production. Nat Immunol. 2001;2(3):269-74.
34. Souweidane MM, Kramer K, Pandit-Taskar N, Zhou Z, Haque S, Zanzonico P, et al. Convection-enhanced delivery for diffuse intrinsic pontine glioma: a single-centre, dose-escalation, phase 1 trial. Lancet Oncol. 2018;19(8):1040-50.
35. Modak S, Kramer K, Gultekin SH, Guo HF, Cheung NK. Monoclonal antibody 8H9 targets a novel cell surface antigen expressed by a wide spectrum of human solid tumors. Cancer Res. 2001;61(10):4048-54.
36. Majzner RG, Theruvath JL, Nellan A, Heitzeneder S, Cui Y, Mount CW, et al. CAR T Cells Targeting B7-H3, a Pan-Cancer Antigen, Demonstrate Potent Preclinical Activity Against Pediatric Solid Tumors and Brain Tumors. Clin Cancer Res. 2019.
37. Wang L, Zhang Q, Chen W, Shan B, Ding Y, Zhang G, et al. B7-H3 is overexpressed in patients suffering osteosarcoma and associated with tumor aggressiveness and metastasis. PLoS One. 2013;8(8):e70689.
38. Zhou Z, Luther N, Ibrahim GM, Hawkins C, Vibhakar R, Handler MH, et al. B7-H3, a potential therapeutic target, is expressed in diffuse intrinsic pontine glioma. J Neurooncol. 2013;111(3):257-64.
39. Lavoie RR, Gargollo PC, Ahmed ME, Kim Y, Baer E, Phelps DA, et al. Surfaceome Profiling of Rhabdomyosarcoma Reveals B7-H3 as a Mediator of Immune Evasion. Cancers (Basel). 2021;13(18).
40. Du H, Hirabayashi K, Ahn S, Kren NP, Montgomery SA, Wang X, et al. Antitumor Responses in the Absence of Toxicity in Solid Tumors by Targeting B7-H3 via Chimeric Antigen Receptor T Cells. Cancer Cell. 2019;35(2):221-37 e8.
41. Crose LE, Etheridge KT, Chen C, Belyea B, Talbot LJ, Bentley RC, et al. FGFR4 blockade exerts distinct antitumorigenic effects in human embryonal versus alveolar rhabdomyosarcoma. Clin Cancer Res. 2012;18(14):3780-90.

42. Khan J, Wei JS, Ringner M, Saal LH, Ladanyi M, Westermann F, et al. Classification and diagnostic prediction of cancers using gene expression profiling and artificial neural networks. Nat Med. 2001;7(6):673-9.

43. Baird K, Davis S, Antonescu CR, Harper UL, Walker RL, Chen Y, et al. Gene expression profiling of human sarcomas: insights into sarcoma biology. Cancer Res. 2005;65(20):9226-35.

44. Davicioni E, Finckenstein FG, Shahbazian V, Buckley JD, Triche TJ, Anderson MJ. Identification of a PAX-FKHR gene expression signature that defines molecular classes and determines the prognosis of alveolar rhabdomyosarcomas. Cancer Res. 2006;66(14):6936-46.

45. Shivaprasad N, Xiong Y, Yohe M, Schneider D, Shern J, Baskar S, et al. 649. Developing FGFR4 Chimeric Antigen Receptor CAR T Cell Therapy Against Rhabdomyosarcoma. Molecular Therapy. 2016;24:S257-S8.

46. Alijaj N, Moutel S, Gouveia ZL, Gray M, Roveri M, Dzhumashev D, et al. Novel FGFR4-Targeting Single-Domain Antibodies for Multiple Targeted Therapies against Rhabdomyosarcoma. Cancers. 2020;12(11):3313.

47. Sullivan PM, Kumar R, Li W, Hoglund V, Wang L, Zhang Y, et al. FGFR4-Targeted Chimeric Antigen Receptors Combined with Anti-Myeloid Polypharmacy Effectively Treat Orthotopic Rhabdomyosarcoma. Mol Cancer Ther. 2022;21(10):1608-21.

48. Bailey SR, Maus MV. Gene editing for immune cell therapies. Nat Biotechnol. 2019;37(12):1425-34.

49. Rafiq S, Hackett CS, Brentjens RJ. Engineering strategies to overcome the current roadblocks in CAR T cell therapy. Nat Rev Clin Oncol. 2020;17(3):147-67.

50. Smith EL, Staehr M, Masakayan R, Tatake IJ, Purdon TJ, Wang X, et al. Development and Evaluation of an Optimal Human Single-Chain Variable Fragment-Derived BCMA-Targeted CAR T Cell Vector. Mol Ther. 2018;26(6):1447-56.

51. Majzner RG, Rietberg SP, Sotillo E, Dong R, Vachharajani VT, Labanieh L, et al. Tuning the Antigen Density Requirement for CAR T-cell Activity. Cancer Discov. 2020;10(5):702-23.

52. Hudecek M, Lupo-Stanghellini MT, Kosasih PL, Sommermeyer D, Jensen MC, Rader C, et al. Receptor affinity and extracellular domain modifications affect tumor recognition by ROR1-specific chimeric antigen receptor T cells. Clin Cancer Res. 2013;19(12):3153-64.

53. Dotti G, Gottschalk S, Savoldo B, Brenner MK. Design and development of therapies using chimeric antigen receptor-expressing T cells. Immunol Rev. 2014;257(1):107-26.

54. Cappell KM, Kochenderfer JN. A comparison of chimeric antigen receptors containing CD28 versus 4-1BB costimulatory domains. Nat Rev Clin Oncol. 2021.

55. Li R, Ma C, Cai H, Chen W. The CAR T-Cell Mechanoimmunology at a Glance. Adv Sci (Weinh). 2020;7(24):2002628.

56. Salter AI, Ivey RG, Kennedy JJ, Voillet V, Rajan A, Alderman EJ, et al. Phosphoproteomic analysis of chimeric antigen receptor signaling reveals kinetic and quantitative differences that affect cell function. Sci Signal. 2018;11(544).

57. Danielli SG, Porpiglia E, De Micheli AJ, Navarro N, Zellinger MJ, Bechtold I, et al. Single-cell profiling of alveolar rhabdomyosarcoma reveals RAS pathway inhibitors as cell-fate hijackers with therapeutic relevance. Sci Adv. 2023;9(6):eade9238.

58. Manzella G, Schreck LD, Breunis WB, Molenaar J, Merks H, Barr FG, et al. Phenotypic profiling with a living biobank of primary rhabdomyosarcoma unravels disease heterogeneity and AKT sensitivity. Nat Commun. 2020;11(1):4629.

59. Meier JA, Zhang F, Sanjana NE. GUIDES: sgRNA design for loss-of-function screens. Nat Methods. 2017;14(9):831-2.

60. Bloemberg D, Nguyen T, MacLean S, Zafer A, Gadoury C, Gurnani K, et al. A High-Throughput Method for Characterizing Novel Chimeric Antigen Receptors in Jurkat Cells. Mol Ther Methods Clin Dev. 2020;16:238-54.

61. Engler C, Kandzia R, Marillonnet S. A one pot, one step, precision cloning method with high throughput capability. PLoS One. 2008;3(11):e3647.

62. Majzner RG, Theruvath JL, Nellan A, Heitzeneder S, Cui Y, Mount CW, et al. CAR T Cells Targeting B7-H3, a Pan-Cancer Antigen, Demonstrate Potent Preclinical Activity Against Pediatric Solid Tumors and Brain Tumors. Clin Cancer Res. 2019;25(8):2560-74.

63. Alijaj N, Moutel S, Gouveia ZL, Gray M, Roveri M, Dzhumashev D, et al. Novel FGFR4-Targeting Single-Domain Antibodies for Multiple Targeted Therapies against Rhabdomyosarcoma. Cancers (Basel). 2020;12(11).

64. Crose LE, Etheridge Kt Fau - Chen C, Chen C Fau - Belyea B, Belyea B Fau - Talbot LJ, Talbot Lj Fau - Bentley RC, Bentley Rc Fau - Linardic CM, et al. FGFR4 blockade exerts distinct antitumorigenic effects in human embryonal versus alveolar rhabdomyosarcoma. (1557-3265 (Electronic)).

65. Taylor JGt, Cheuk AT, Tsang PS, Chung JY, Song YK, Desai K, et al. Identification of FGFR4-activating mutations in human rhabdomyosarcomas that promote metastasis in xenotransplanted models. J Clin Invest. 2009;119(11):3395-407.

66. Ying Z, Huang XF, Xiang X, Liu Y, Kang X, Song Y, et al. A safe and potent anti-CD19 CAR T cell therapy. Nat Med. 2019;25(6):947-53.

67. Larson RC, Kann MC, Bailey SR, Haradhvala NJ, Llopis PM, Bouffard AA, et al. CAR T cell killing requires the IFNgammaR pathway in solid but not liquid tumours. Nature. 2022;604(7906):563-70.

68. Zhao X, Yang J, Zhang X, Lu XA, Xiong M, Zhang J, et al. Efficacy and Safety of CD28- or 4-1BB-Based CD19 CAR-T Cells in B Cell Acute Lymphoblastic Leukemia. Mol Ther Oncolytics. 2020;18:272-81.

69. Ying Z, He T, Wang X, Zheng W, Lin N, Tu M, et al. Parallel Comparison of 4-1BB or CD28 Co-stimulated CD19-Targeted CAR-T Cells for B Cell Non-Hodgkin's Lymphoma. Mol Ther Oncolytics. 2019;15:60-8.

70. Tang X, Zhao S, Zhang Y, Wang Y, Zhang Z, Yang M, et al. B7-H3 as a Novel CAR-T Therapeutic Target for Glioblastoma. Mol Ther Oncolytics. 2019;14:279-87.

71. Tian M, Cheuk AT, Wei JS, Abdelmaksoud A, Chou HC, Milewski D, et al. An optimized bicistronic chimeric antigen receptor against GPC2 or CD276 overcomes heterogeneous expression in neuroblastoma. J Clin Invest. 2022;132(16).

72. Sun F, Yu X, Ju R, Wang Z, Wang Y. Antitumor responses in gastric cancer by targeting B7H3 via chimeric antigen receptor T cells. Cancer Cell Int. 2022;22(1):50.

73. Xuan Y, Sheng Y, Zhang D, Zhang K, Zhang Z, Ping Y, et al. Targeting CD276 by CAR-T cells induces regression of esophagus squamous cell carcinoma in xenograft mouse models. Transl Oncol. 2021;14(8):101138.

74. van der Stegen SJ, Hamieh M, Sadelain M. The pharmacology of second-generation chimeric antigen receptors. Nat Rev Drug Discov. 2015;14(7):499-509.

75. Ramos CA, Rouce R, Robertson CS, Reyna A, Narala N, Vyas G, et al. In Vivo Fate and Activity of Second- versus Third-Generation CD19-Specific CAR-T Cells in B Cell Non-Hodgkin's Lymphomas. Mol Ther. 2018;26(12):2727-37.

76. Kalaitsidou M, Kueberuwa G, Schutt A, Gilham DE. CAR T-cell therapy: toxicity and the relevance of preclinical models. Immunotherapy. 2015;7(5):487-97.

77. Kagoya Y, Nakatsugawa M, Yamashita Y, Ochi T, Guo T, Anczurowski M, et al. BET bromodomain inhibition enhances T cell persistence and function in adoptive immunotherapy models. J Clin Invest. 2016;126(9):3479-94.

78. Scholz G, Jandus C, Zhang L, Grandclement C, Lopez-Mejia IC, Soneson C, et al. Modulation of mTOR Signalling Triggers the Formation of Stem Cell-like Memory T Cells. Ebiomedicine. 2016;4:50-61.

79. Biasco L, Scala S, Basso Ricci L, Dionisio F, Baricordi C, Calabria A, et al. In vivo tracking of T cells in humans unveils decade-long survival and activity of genetically modified T memory stem cells. Sci Transl Med 2015;7(273):273ra13.

80. Biasco L, Izotova N, Rivat C, Ghorashian S, Richardson R, Guvenel A, et al. Clonal expansion of T memory stem cells determines early anti-leukemic responses and long-term CAR T cell persistence in patients. Nat Cancer. 2021;2(6):629-42.

81. Walker AJ, Majzner RG, Zhang L, Wanhainen K, Long AH, Nguyen SM, et al. Tumor antigen and receptor densities regulate efficacy of a chimeric antigen receptor targeting anaplastic lymphoma kinase. Mol Ther. 2017;25(9):2189-201.

82. Heitzeneder S, Bosse KR, Zhu Z, Zhelev D, Majzner RG, Radosevich MT, et al. GPC2-CAR T cells tuned for low antigen density mediate potent activity against neuroblastoma without toxicity. Cancer Cell. 2022;40(1):53-69 e9.

Figures, tables and additional files

Fig. 1. CD276 expression in RMS cell lines and PDXs.

(**A**) CD276 expression levels were evaluated in eleven RMS cell lines, three PDX, myoblasts, PBMCs, and isolated T cells by Western Blotting, revealing high expression in all RMS cell lines, except for the FP-RMS JR and RMS, medium-high levels in the three investigated PDXs, and no detection in the controls. (**B**) Surface CD276 expression on eleven RMS cell lines, and PBMCs and T cells as controls, were assessed by Flow Cytometry confirming high expression of CD276 in all the considered RMS samples, especially in RUCH-3, RD and Rh4, and no detectable expression in the controls. The correspondent isotype controls are shaded in light grey. (**C**) The number of CD276 molecules on the surface of RMS cells, PDXs, and controls was estimated by Quantibrite PE beads. (**D**) Surface CD276 expression on three RMS PDXs, and PBMCs and T cells as controls, was assessed by Flow Cytometry

confirming high expression of CD276 in all three PDXs. The correspondent isotype controls are shaded in light grey. (**E**) The number of CD276 molecules on the surface of RMS PDXs and controls was estimated by Quantibrite PE beads.

Fig. 2. Schematic representation of the investigated CAR constructs.

(**A**) The backbone used for the expression of the CARs is characterized by an EFS promoter, followed by a Kozak sequence and a CD28 leader sequence. The ORF is composed of two MycTag regions; an Antigen Binding Domain (ABD; anti-CD19 scFv FMC63 for the initial cloning) flanked by two Esp3I restriction sites for ABD swapping with anti-CD276 scFv CD276.MG or with sdAbs targeting FGFR4; a Hinge Domain (HD) and a Transmembrane domain (TM) derived from CD8α or CD28; one or two Co-Stimulatory Domains (CSDs), derived from CD28 or 4-1BB; and a CD3ζ Signaling Domain (SD). Downstream, GFP is linked to the CD3ζ domain by the self-cleaving peptide P2A as reporter. (**B**) Modular CAR structure of the different constructs generated by Gibson assembly with either CD8α HD/TM (green) or CD28 HD/TM (cyan). In each panel, the schematic representation of two second-generation CARs, characterized by a single CSD (CD28 or 4-1BB), and two third-generation CARs, characterized by the expression of both CSDs (CD28-4-1BB or 4-1BB-CD28). Created with Biorender.com.

Fig. 3. Manufacturing, expansion, viability, and CD4$^+$:CD8$^+$ ratios of the most representative CD276-CAR T cells.

(**A**) Manufacturing, selection, and test of CAR T cells. After isolation from PBMCs, T cells were incubated on day 1 with IL-2 and anti-CD3/CD28 activators. On day 4, T cells were infected with lentiviruses and CAR T cells were sorted by GFP expression on day 7 and further expanded for 7 days. Surface CARs expression and low activation before cytotoxicity assays CAR T cells were then tested by WB and FACS. On day 14, CAR T cells were co-incubated with RMS cells for killing assays and cytokine release quantification. (**B**) Expansion and viability of CAR T cells were monitored from day 4 to day 13, resulting in up to 35-fold expansion, with a viability >70%. (**C**) CD4$^+$:CD8$^+$ ratio was assessed on day 14 to rely on a balanced number of helper and cytotoxic T cells. In all the CAR T cells the CD4$^+$:CD8$^+$ ratio was around 1:1. Created with BioRender.com

Fig. 4. Validation of the CAR constructs expression in T cells.

(**A, B**) CD19-CAR expression in T cells were assessed by (**A**) Western Blotting and (**B**) by Flow Cytometry on day 14 after first activation of T cells. (**C, D**) CD276-CAR expression in T cells were assessed by (**C**) Western Blotting and (**D**) by Flow Cytometry on day 14 after first activation of T cells. Anti-MycTag antibody was used to discriminate the different CAR generations by size in Western Blotting. (**A**, **C**) The respective bands confirm the predicted CAR size around 60 kDa with a visible shift between the second- and third-generations CARs. (**B**, **D**) Surface expression of CAR constructs was detected with an anti-MycTag PE-conjugated antibody by Flow Cytometry, confirming robust expression of the CAR construct on the surface of T cells. All the experiments were performed in triplicate and with three different donors. Shown are results from one representative donor.

Fig. 5. Phenotypic characterization of CAR T cells during expansion and co-incubation experiments.

Flow Cytometry during CAR T cells manufacturing was used to quantify percentages of cell memory (T_{CM}), effector memory (T_{EM}) and effector (T_{EFF}) T cells on CD3 positive cells by CD45RA and CD62L staining. Naïve (T_N) and stem cell-like (T_{SCM}) T cells were quantified on CD3$^+$/CD45RA$^+$/CD62L$^+$ by CD95 and CCR7 staining. (**A**) On day 1, activated (upper panels) and non-activated (lower panels) T cells were compared. T_{CM} cells percentage (central panel) were higher in non-activated T cells compared to activated. On the other hand, T_{EFF} percentage was higher in activated T cells. A transition towards T_{SCM} is evident upon activation (right panel). Of note, CD3 staining was weaker in activated T cells, due to the presence of CD3/CD28 cross-linking activators. (**B**) On day 7, activated CAR T cells showed very high percentage of T_{SCM} cells and lower percentages of T_{EM} and T_{EFF} cells, compared to day 1. (**C**) On day 14, before co-incubation assays, almost ~70% of activated CAR T cells had a T_{SCM} phenotype and ~27% a T_{CM} phenotype; on day 16, after cytotoxicity experiments, ~60% were T_{SCM} cells and ~36% were T_{CM} cells.

Fig. 6. Evaluation of killing capacity by CD276-directed CAR T cells after co-incubation with RD and Rh4 cell lines.

In order to determine the CAR conferring the highest cytotoxicity, CAR T cells were co-incubated for 48h with fLuc⁺ RD and Rh4 cells at different E:T ratios. Luciferase assays showed effective killing capacity by all the generations of CAR T cells at E:T ratios of 5:1 and 10:1. Lower E:T ratios highlighted optimal efficacy by CD276.V-CAR T cells, compared to the other constructs (upper panels). CD19-CAR T cells showed no cytotoxic effect when co-incubated with fLuc⁺ RMS cells (lower panels). All the experiments were performed in triplicate and with three different donors. Shown are results from one representative donor.

Fig. 7. Cytokine release by CD276-CAR T cells after 24h co-incubation with RD and Rh4 cell lines. Concentrations of IL-2, IFN-γ, and Granzyme B in the supernatant released during co-incubation of CAR T cells with RD (left panels) and Rh4 cells (right panels) at the E:T ratio of 5:1 were measured by ELISA. (**A**) IL-2 levels detected upon co-incubation with RD and Rh4 cells revealed CD276.I (red), CD276.II (orange), and CD276.V (cyan) as the constructs with the highest IL-2 release. (**B**) CD276.V (cyan) released maximum levels of IFN-γ (400 pg/ml) with both RD and Rh4 cell lines. (**C**) The highest quantity of Granzyme B (>8000 pg/ml) measured during co-incubation with RD cells was released by CD276.V-CAR T cells. Similar values were observed among CD276.II, CD276.V, CD276.VI, and CD276.VII, when co-cultured with Rh4 cells. In all the experiments, IL-2, IFN-γ, and Granzyme B levels were at background levels when CD276.V-CAR T cells were co-incubated with CD276 KO RMS cells, similar to background levels detected in RD and Rh4 alone supernatant (brown). All the experiments were performed in triplicate and with three different donors. Shown is a representative donor.

Fig. 8. FGFR4 expression in RMS cell lines and PDXs.

(**A**) FGFR4 expression levels were evaluated in eleven RMS cell lines, three PDXs, and in the controls as Myoblasts, PBMCs, and T cells. WB results revealed high expression in Rh36, Rh4, Rh28, JR, and RMS cell lines, whereas low levels were detected in Rh18, RUCH-3, Rh5, and Rh30. No FGFR4 expression was observed in the controls. (**B**) Flow Cytometry results confirmed expression of FGFR4 on the surface of most RMS cell lines, especially high in Rh4, Rh18, and JR. The corresponding isotype controls are highlighted in grey. (**C**) Estimation of FGFR4 expression on RMS cell lines by Quantibrite PE-Beads showed a surface expression in the range of 600-4000 copies per cell.

Fig. 9. Selection of FGFR4 sdAb and killing capacity of Dual-CAR T cells.

(**A**) To select the FGFR4 sdAb that conferred the most specific and potent killing capacity to the 28H-28TM-28CSD-3z CAR construct, four different sdAb FGFR4-CAR T cells were co-incubated for 48h with fLuc⁺ Rh4 cells at different E:T ratios. RMS cells survival measured by Luciferase assays showed effective killing capacity by all the FGFR4-CAR T cells at E:T ratios of 5:1 and 10:1. Low E:T ratios put in evidence B5-FR4 sdAb-based CAR as the most effective (pink line and full circles, left). However, experiments performed with FGFR4 KO Rh4 showed unspecific killing of the CAR T cells A8-FR4.V, B1-FR4.V, and B5-FR4.V. Nevertheless, F8-FR4-V-CAR T cells did not kill FGFR4 KO Rh4 cells and showed moderate killing with fLuc⁺ Rh4 cells (right). (**B**) Evaluation of CD276/FGFR4 Dual-CAR T cells. The killing capacity of Dual-CAR T cells was evaluated by co-incubation experiments with Rh4 and JR cell lines for 48h at different E:T ratios. Luciferase assays showed effective killing capacity by CD276.V-, F8-FR4.V-, and Dual-CAR T cells at E:T ratios of 5:1 and 10:1. Low E:T ratios exhibited high efficacy by CD276.V- and Dual-CAR T cells, compared to F8-FR4.V-CAR T cells (left panel). Similar experiments performed with JR cells showed high killing capacity by F8-FR4.V- and Dual-CAR T cells, but moderate efficacy by CD276.V-CAR T cells (right). CD19-CAR T cells and untransduced T cells showed no significant killing against both RMS cells.

Fig. 10. *In vivo* experiment with CD276.V-CAR T cells in mice orthotopically engrafted with RD cells. (**A**) Schematic overview of the experimental design. On day -10, PBMCs were thawed, T cells were purified and activated. On day -6, CAR T cells were produced and expanded until day 5. In the meantime, on day 0, 0.5×10^6 fLuc$^+$ RD cells were orthotopically injected into the left gastrocnemius muscle of mice. After 5 days, CAR T cells were injected into mice by tail vein injection. (**B**) Tumor progression was monitored by bioluminescence photometry. Efficient tumor eradication by CD276.V-CAR T cells was observed in 3/5 mice after 3 weeks after RD cells injection on day 0, whereas CD276.V-CAR T cells showed significant tumor burden control in 2/5 mice. Maintained complete remission was confirmed in 3/5 mice in the group treated with CD276-CAR T cells until the end of the experiment (day 110).

Fig. 11. *In vivo* experiment with CD276-CAR T cells, FGFR4-CAR T cells, and CD276/FGFR4 Dual-CAR T cells in mice orthotopically engrafted with Rh4 cells.

(**A**) Schematic overview of the experimental design. On day -10, PBMCs were thawed, T cells were purified and activated. On day -6, CAR T cells were produced and expanded until day 5. In the meantime, on day 0, 0.5×10^6 fLuc$^+$ Rh4 cells were orthotopically injected into the left gastrocnemius muscle of mice. After 5 days, CAR T cells were injected into mice by tail vein injection. (**B**) Tumor progression was monitored by bioluminescence photometry. Efficient tumor eradication by CD276.V-CAR T cells and Dual-CAR T cells was observed in 5/5 mice 3 weeks after Rh4 cells injection on day 0. Maintained complete remission was confirmed in both groups until the end of the experiment (day 60). (**C**) Tumor growth curves for the investigated groups as measured by caliper. Humane endpoint was set to 1000 mm^3. (**D**) Kaplan-Meier plot showing survival of treated and control mice. Humane endpoint was set to 1000 mm^3.

Fig. 12. *In vivo* experiment with CD276-CAR T cells, FGFR4-CAR T cells, and CD276/FGFR4 Dual-CAR T cells in mice orthotopically engrafted with JR cells.

(**A**) Schematic overview of the experimental design. On day -10, PBMCs were thawed, T cells were purified, and activated. On day -6, CAR T cells were produced and expanded until day 5. In the meantime, on day 0, 0.5×10^6 fLuc$^+$ JR cells were orthotopically engrafted into the left gastrocnemius muscle of mice. After 5 days, CAR T cells were injected into mice by tail vein injection. (**B**) Tumor progression monitored by bioluminescence photometry. Tumor eradication was observed in 1/5 mouse in the CD276.V-CAR T cells treated group, which was maintained until the end of the experiment at day 50. (**C**) Tumor growth curves for the investigated groups as measured by caliper. CD276.V-CAR T cells and Dual-CAR T cells delayed tumor growth in 3/5 and 1/5 mice, respectively. (**D**) Kaplan-Meier plot showing survival of treated and control mice. Humane endpoint was set to 1000 mm^3.

Fig. 13. Expression of CD276 and FGFR4 on RMS-derived xenografts. (**A**) Expression of CD276 was assessed by Flow Cytometry on untreated Rh4-derived xenograft tumors and compared to cultured wild-type Rh4 cells by using a PE-conjugated anti-CD276 antibody. CD276 expression was calculated

dividing the geometric mean of antibody-stained cells by the geometric mean of the controls and showed a lower CD276 expression on Rh4 tumor-derived cells, compared to the cultured Rh4 cells. (**B**) Expression of FGFR4 was assessed by Flow Cytometry on Rh4-derived xenograft tumors from untreated mice and compared to cultured wild-type Rh4 cells by using a PE-conjugated anti-FGFR4 antibody. FGFR4 expression was calculated dividing the geometric mean of antibody-stained cells by the geometric mean of the isotype controls. The results indicated a strong decrease in FGFR4 expression in Rh4 tumor xenograft-derived cells. (**C**) Assessment of CD276 and FGFR4 expression in tumor tissue xenografts by IHC. After treatment, CD276 is still highly detectable on RD- and Rh4-derived tumor xenografts, but low staining detection was observed in JR-derived tumors. FGFR4 seems expressed at low levels on RD, at medium levels on JR, and at high levels on Rh4.

Fig. 14. T cells detection in tumors and spleen. (**A**) Detection of T cells in orthotopic tumors derived from JR or Rh4 cells. Tumors were collected at the end of the experiment, after 5 weeks from JR implantation, and after 7 weeks from Rh4 implantation. CD3 staining revealed presence of CAR T cells in JR- and Rh4-derived xenografts. However, in JR xenografts CAR T cells were only detectable in the vessels (black arrows and inset), whereas T cells seemed to have infiltrated Rh4 tumors. (**B**) Evidence of CAR T cells persistence in the spleen CD3 staining revealed presence of T cells in the spleen of mice bearing Rh4 orthotopic tumors and injected with untransduced T cells (UTD) and CD276.V-CAR T cells at the end of the experiment, after 8 weeks and 9 weeks from Rh4 implantation, respectively.