A pangenomic atlas reveals that eco-evolutionary dynamics shape plant pathogen type VI secretion systems

Authors
Nathalie Aoun*[1], Stratton J. Georgoulis*[1], Jason K. Avalos[1], Kimberly J. Grulla[1], Kasey Miqueo[1], Cloe Tom[1], and Tiffany M. Lowe-Power[1].

*These authors contributed equally to this work.

Affiliations:
[1]Department of Plant Pathology, University of California Davis, Davis, CA 95616

Corresponding author:
Tiffany M. Lowe-Power;

E-mail: tlowepower@ucdavis.edu

Abstract

Soil-borne *Ralstonia solanacearum* species complex (RSSC) bacteria disrupt rhizosphere and endophytic microbial communities as they invade roots and fatally wilt plants. RSSC pathogens secrete antimicrobial toxins using a type VI secretion system (T6SS). To investigate how evolution and ecology have shaped pathogen T6SS biology, we analyzed the T6SS gene content and architecture across the RSSC pangenome and their evolutionarily relatives. Our analysis reveals that two ecologically similar Burkholderiaceae taxa, xylem pathogenic RSSC bacteria and *Acidovorax*, have convergently evolved to wield large arsenals of T6SS toxins. To understand the mechanisms underlying genomic enrichment of T6SS toxins, we compiled an atlas of 1,069 auxiliary ("*aux*") T6SS toxin clusters across 99 high-quality RSSC genomes. We classified 25 types of *aux* clusters with toxins that predominantly target lipids, nucleic acids, or unknown cellular substrates. The *aux* clusters were in diverse genetic neighborhoods and had complex phylogenetic distributions, suggesting frequent horizontal gene flow. Phages and other mobile genetic elements account for most of the *aux* cluster acquisition on the chromosome but very little on the megaplasmid. Nevertheless, RSSC genomes were more enriched in *aux* clusters on the megaplasmid. Secondary replicons like megaplasmids often evolve more rapidly than the more evolutionarily stable chromosome. Although the single ancestral T6SS was broadly conserved in the RSSC, the T6SS was convergently lost in atypical lineages with vectored transmission. Overall, our data suggest dynamic interplay between the lifestyle of soil-transmitted RSSC lineages and the evolution of T6SSs with robust arsenals of toxins. This pangenomic atlas poises the RSSC as an emerging, tractable model to understand the role of the T6SS in shaping pathogen populations.

Introduction

To defend their habitat or colonize new niches, bacteria often attack competing microbes by secreting toxic molecules and proteins (1, 2). Many gram-negative bacteria use a molecular weapon known as the type 6 secretion system (T6SS) to deliver toxic protein effectors to target cells (3, 4). T6SS effectors bind to the tip of a multimeric, spear-like projectile (5–7). Upon contracting, the sheath forcibly ejects the projectile, puncturing nearby cells and delivering the associated effectors directly to the intracellular space of target cells within range (3, 8). T6SS effectors kill target cells by degrading cellular components including DNA, lipid membranes, and bacterial cell wall polymers. To prevent damage to kin cells, each T6SS effector has a cognate immunity protein that prevents toxicity, usually by directly binding the effector (8, 9).

Most T6SSs include a double membrane spanning complex (TssJLM) which recruits a baseplate complex (TssEFGK) and a spike complex (VgrG heterotrimer, PAAR sharp tip, effectors) (10–13). Hcp then polymerizes to form the shaft of the projectile, and repeating units of TssBC form a sheath around the shaft that contacts to fire the projectile complex. After firing, TssH disassembles the TssBC sheath, allowing the monomers to reassemble again in a new complex (14, 15). While just one copy of most *tss* genes is required for a functional T6SS, T6SS⁺ genomes usually encode multiple copies of *vgrG*, typically collocated with the cognate effector gene, immunity gene, and genes encoding the PAAR sharpening tip protein and any adapter proteins (13, 16). More copies of *vgrG* in a genome implies a greater diversity of effectors, potentially allowing the T6SS to be deadly against a greater diversity of targets. Clusters of *vgrG* and associated genes may be located with the other *tss* genes or in other locations termed "T6SS auxiliary clusters" ("*aux* clusters") (17, 18). Horizontal gene transfer has previously been implicated in the spread of *aux* clusters between bacterial genomes (19–22), contributing to convoluted evolutionary patterns of gain and loss that result in diverse T6SS effector arsenals. Even closely related strains often have non-identical T6SS effector repertoires, allowing interstrain competition within a bacterial species (23, 24). T6SS-mediated interstrain competition has been shown to contribute to the outcome of host colonization when two or more strains of the same species attempt to colonize a new host (25). Large-scale analyses suggest that plant-associated and human-associated bacteria are more likely to have a T6SS (26).

Plant pathogens in the *Ralstonia solanacearum* species complex (RSSC) are known to have a functional T6SS (27–29). RSSC pathogens cause severe wilt disease on diverse plants, including economically important crops like potatoes, tomatoes, and bananas. Like many host-associated bacteria, RSSC plant pathogens must successfully transition between disparate ecosystems to complete their disease cycle. RSSC bacteria persist in soil or water and chemotax through the microbially dense rhizosphere to infect a new host (30). *Ralstonia* enter the roots and spread throughout the xylem, growing to such a high density that the mass of cells and biofilm physically block water transport, wilting the plant (31, 32). Transcriptomic analyses indicate that the T6SS is upregulated when RSSC pathogens reach quorum, as in an infected plant (33). T6SS-inactive *Ralstonia* mutants have pleiotropic phenotypes, including altered biofilm and motility, which has made it difficult to define the precise ecological role for the RSSC T6SS.

We hypothesized that intermicrobial competition during plant colonization has shaped the evolution of RSSC pathogens and their T6SS weapons. Here, we used an evolutionary genomics approach to shed light on the eco-evolutionary dynamics between the lifestyle of RSSC strains and their T6SS arsenals. We carried out a pangenome analysis of T6SS-related genes on diverse genomes in the RSSC

compared to Burkholderiaceae family relatives. We found that that plant pathogenic *Ralstonia* genomes are evolutionarily enriched in T6SS effectors. Scrutinizing the diversity and distribution of T6SS effectors in the RSSC, we found a complex distribution of T6SS *aux* clusters suggesting that *aux* clusters are frequently gained and lost. More of the identified T6SS gene clusters were present on the megaplasmid than the main chromosome. Intriguingly, most of the chromosomally located *aux* clusters were associated with mobile genetic elements, while there was rarely evidence of mobile genetic elements associated with megaplasmid-located *aux* clusters. We infer that the T6SS is an ancestral trait in the RSSC whose genetic architecture is shaped by complex ecological and evolutionary processes. Our analyses reveal that the T6SS is a dynamic weapon intimately linked to the evolutionary success of soilborne *Ralstonia*, a plant pathogen of global concern.

Results

Bacteria in the family Burkholderiaceae vary in their T6SS gene content.

We used an evolutionary framework to explore T6SS gene content among the RSSC relative to their evolutionary neighbors in the Burkholderiaceae family. Using the genome taxonomy database (GTDB) (34–39), we identified complete genomes of 288 Burkholderiaceae species (Table S1, S2). We used BLASTp searches to estimate the abundance of T6SS genes in the Burkholderiaceae genomes and a curated set of 99 high-quality RSSC genomes. To estimate the number of secretion systems per genome, we averaged the number of BLASTp hits for the core T6SS components: TssABCEFGHJKLM and Hcp (40). The majority of the Burkholderiaceae species representatives encoded a T6SS (62.4%) (Figure S1, Table S1). The three species representatives of the RSSC wilt pathogens (*R. solanacearum* IBSBF1503, *R. pseudosolanacearum* GMI1000, and *R. syzygii* PSI07), each encoded a single copy of the core T6SS genes. In contrast, there were multiple T6SSs encoded in approximately 27% of the Burkholderiaceae genomes; between three to six sets of T6SS core genes were identified in genomes in the *Burkholderia*, *Trinickia*, *Caballeronia*, *Variovorax*, *Pseudoduganella*, *Massilia*, and *Paraburkholderia* genera (Figure S2; Table S1), which is consistent with previous reports for some of these taxa (41–45).

Genomes of RSSC plant pathogens are enriched in *vgrG* genes.

T6SS effector/immunity pairs are often encoded in gene clusters with their corresponding *vgrG,* any adaptors that mediate effector translocation, and any genes encoding the PAAR sharpening tip protein (46–48). Because VgrG proteins have conserved sequences, we estimated the abundance of effector/immunity pairs by searching genomes for *vgrG* copies. We carried out multiple low-stringency BLASTp searches for *vgrG* alleles in the 288 Burkholderiaceae species representatives and the 99 high-quality RSSC genomes. As expected, most of the T6SS[null] genomes lacked intact *vgrG* genes (Figure S2). Generally, there was a positive correlation between the copy number of T6SS core genes and *vgrG* alleles (Figure 1A, S2).

Two taxa with single copies of the core T6SS genes were enriched in *vgrG* genes: the plant-pathogenic RSSC and the genomo-genus Acidovorax_A (Figure 1A). Acidovorax_A is a genomo-genus defined by GTDB (34–39) that contains two xylem-infecting plant pathogen species, *Acidovorax avenae* and *Acidovorax citrulli*. To investigate whether the RSSC had an evolutionary enrichment of *vgrG* copies, we compared the number of *vgrG* genes in the RSSC to other species in the *Ralstonia* genus (n=70) and the closely related genera *Cupriavidus* (n=120) and *Pandoraea* (n=75). Plant-pathogenic RSSC genomes typically had over twice as many *vgrG* copies as their close relatives (median of n=12 per genome; Figure

1B). Most *Pandoraea* spp. did not encode *vgrG*. *Cupriavidus* and the non-RSSC *Ralstonia* encoded a similar number of *vgrG* copies (medians of n=4 or n=5 per genome, respectively) (Figure 1B).

A single T6SS subtype is largely conserved among plant pathogenic *RSSC*.

We used synteny analyses to investigate the organization of T6SS genes among the plant pathogenic RSSC. The genomes we investigated have the same T6SS subtype, $T6SS^{i4b}$ (49, 50) (Figure 2A), encoded on the 2.1 Mb secondary replicon known as the megaplasmid (51). The RSSC T6SS has two conserved regions interspersed between three variable regions that mainly contain *vgrG*-linked effector/immunity clusters and small transposable elements (Figure 2A-B).

Excluding the three *vgrG* paralogs, we identified 19 core genes in the T6SS locus (Figure 2A). We BLAST-searched the 19 genes against the 288 Burkholderiaceae species representatives. Although the structural genes were well-conserved, we observed that six of the core $T6SS^{i4b}$ genes were variably present in 21.4-70.9% of the other T6SS-containing genomes: RS_RS20665, *GH10* (glycosyl hydrolase 10), *ompA*, *tagF*, RS_RS20775, and the M15 peptidase (Figure S1). One gene encoding a hypothetical protein, RS_RS20790, was exclusively found in *Ralstonia* genomes (Figure S1).

Several RSSC lineages have lost T6SS.

Upon searching 398 publicly available RSSC genomes, we identified several lineages that lacked the T6SS (Figure 2C). The $T6SS^{null}$ lineages include the two insect and mechanically vectored phylotype IV lineages that cause Sumatra Disease of Clove (SDC) and Blood Disease of Banana (BDB), one of the lineages that causes Moko Disease of Banana (phylotype IIA-6), one phylotype I lineage, and all but one of the eight phylotype III strains with sequenced genomes. Of these $T6SS^{null}$ lineages, the SDC lineage strains encode the *gh10* and *ompA* genes, suggesting that the other T6SS genes were lost. We also identified multiple genomes that lacked the conserved *tssL*-to-*tssH* and the *GH10*-to-RS_RS20790 region, (n=30 and n=24, respectively; Figure 2C, Table S2). There were sporadic genomes that lacked a BLASTp hit for certain core genes, but these could be false negatives based on carrying out BLASTp based searches on draft genomes of varying quality.

To infer whether the $T6SS^{i4b}$ had been lost in the $T6SS^{null}$ lineages, we carried out BLAST searches and synteny analyses with all the T6SS gene clusters identified in RSSC and the non-RSSC *Ralstonia*. Of the $T6SS^+$ non-RSSC *Ralstonia*, 98% of the genomes encoded a $T6SS^{i4b}$ with the same gene organization except for the absence of the GH10-encoding gene (Figure S3). The simplest explanation for the phylogenetic pattern is that the $T6SS^{i4b}$ is ancestral to the genus *Ralstonia* and has independently been lost in multiple lineages of the RSSC and non-RSSC *Ralstonia*.

The RSSC pangenome encodes dozens of auxiliary T6SS effector/immunity clusters.

Because RSSC genomes contain a median of 12 *vgrG* genes and only three *vgrG* paralogs are encoded in the main $T6SS^{i4b}$ locus, we hypothesized that there were additional T6SS loci on the chromosome or megaplasmid. We performed an iterative synteny analysis of the gene neighborhoods of all *vgrG* alleles in the 99 high-quality RSSC genomes, which allowed us to infer the presence of 1069 auxiliary effector/immunity (*aux*) clusters that we classified into 25 types: *aux1-aux25* (Table S3). Furthermore, we used synteny analyses to map the *aux* clusters to their locations on the chromosome, megaplasmid, or small accessory plasmids. Figures S4-S24 show an atlas for each *aux* cluster, depicting the phylogenetic distribution, gene organization and annotations, and genomic locations.

The *aux* clusters contain between three and 14 genes and vary in size from 2.27 kb (*aux17*) to 16.6 kb (*aux16*). We used the NCBI Conserved Domain Database and PaperBlast to infer the function of each *aux* cluster gene (Figures S4-S24). Although there is a conserved standalone PAAR domain gene in the T6SS main, small PAAR domain-containing genes were also encoded in some or all structural variants of

five *aux* types (*aux1, aux2, aux5, aux7, aux9*). While some T6SS effectors bind directly to VgrG spike proteins, others must form complexes with adaptor proteins (52). We identified adaptors with DUF1795 (24) and DUF2169 (53) in three *aux* types and DUF4123 domains (54) in four *aux* types. Several of the *aux* clusters include genes containing polymorphic effectors with recombination hotspot (RHS) domains (*aux3, aux4,* and *aux12*), DUF4150 PAAR-like superfamily domains (*aux8, aux13*, and *aux16*) (55), marker-for-type-six (MIX_III) domains (*aux14* and *aux20*) (56), found-in-type-six (FIX) domains (*aux10, aux17, aux22, aux23*, and *aux24*) (17), and FIX-like domains (*aux19* and *aux21*). Finally, genes also included other domains that have been previously identified in T6SS effectors (DUF3274, DUF2235, DUF2875, and DUF6531) (57–59) and immunity proteins (DUF1910, DUF1911, DUF3304, and Sel1 repeats) (17, 18, 57) (Figures S4-S24). We identified putative arrays of immunity genes or orphan immunity genes in structural variants in 13 of the *aux* types: *aux2, aux3, aux4, aux5, aux6, aux8, aux11, aux12, aux14, aux15, aux18, aux22*, and *aux23*.

The largest two clusters, *aux8* and *aux16*, are atypical because they were the only ones that contain genes upstream of the *vgrG* gene: one DUF4124-encoding gene and 1-2 ankyrin repeat-encoding genes (Figures S11 and S19). All genes in *aux1-aux24* were arranged unidirectionally, but *aux25* had an atypical three-gene layout of *vgrG*, a 2.3 kb hypothetical gene, and then a conserved inverted gene encoding a PvdO family nonheme iron enzyme (Figure S24).

RSSC-encoded T6SS effectors target diverse substrates, including lipids and DNA.
T6SS effectors damage important cellular components in target cells. We used the NCBI Conserved Domain Database (60) and PaperBlast (61) to infer the mode-of-action of the effectors (Figure 3, Table S3). The effectors from nine *aux* clusters were lipases of previously defined families (62): Tle1 (*aux6* and *aux11*), Tle3 (*aux2, aux7,* and *aux9*), Tle4 (*aux1* and *aux15*), Tle5 (*aux6*), and a lipase with a novel domain architecture (*aux18*). The effectors of seven *aux* clusters were nucleases with HNH nuclease domains (*aux3, aux12,* and *aux16*), GHH2 nuclease domains (*aux8*), or PoNe nuclease domains (*aux10, aux19,* and *aux21*). The polymorphic RHS effectors in *aux4* varied in the C-terminal effector, containing either ATP-targeting (p)ppApp synthase or actin-ADP ribosylase. For the remaining nine clusters, we could infer which gene was the effector based on the presence of known polymorphic effector domains for eight of the clusters (*aux3, aux14, aux17, aux20, aux22, aux23, aux24,* and *aux25),* but we could not identify domains or motifs that hint at the mode-of-action (Figure 3E). For the final cluster, *aux13*, the effector might be either the MAEBL-domain protein or one of the three hypothetical proteins.

We investigated whether RSSC lineages varied in the abundance of lipase, nuclease, ATP/Actin manipulating, and unknown-targeting effectors (Figure 3). All T6SS⁺ RSSC genomes encoded at least one lipase. Phylotype I and IIC lineages encoded the most lipases (median of 5 and 8, respectively) (Figure 3A-B). Phylotype IIB-1, which contains the clonal pandemic lineage (regulated as a U.S. Select Agent under the name "*R. solanacearum* R3Bv2")(63), had the fewest lipases of the T6SS⁺ genomes with only a single lipase per genome. Nucleases were common, but 17% of T6SS⁺ strains lacked any obvious nucleases. Sporadic phylotype I strains, the IIB-1 lineage, and the IV-8 lineage had more nucleases than the other lineages (Figure 3A). Genomes with more nucleases tended to have fewer lipases. The ATP and actin-targeting effectors are rare and sporadically distributed. None of the analyzed phylotype II genomes encoded an actin-targeting effector. Most of the T6SS⁺ genomes (91%) had one or more effectors with unknown targets.

T6SS gene conservation varies among RSSC clades.
The copy number of *aux* types ranged from zero to three copies among the RSSC genomes (Figure S25). Some *aux* types were never found with more than a single copy per genome. *aux2, aux5,* and *aux7*

were sometimes found in two or three copies, and *aux1, aux4, aux6, aux9,* and *aux17* were sometimes found in two copies across the RSSC genomes. The *aux10, aux11, aux12, aux18, aux19, aux21, aux24* and *aux25* were found in only 20% of the RSSC genomes. Meanwhile, *aux1*, *aux*2, *aux*3, *aux*5, *aux*6, *aux*7, *aux*14, and *aux*15 were found in more than 50% of the RSSC genomes (Figure S25).

To compare the phylogenetic distribution of different *aux* types, we visualized the copy number of each *aux* type across an RSSC phylogenetic tree (Figure 4C and Figure S26). To organize the *aux* types based on their patterns of presence in each genome, we calculated the Bray-Curtis dissimilarity metric based on the *aux* cluster content of each strain and used the *aux* cluster content dissimilarity matrix to draw a dendrogram (Figure 4C, Table S4). The dendrogram divides *aux* clusters into two groups: one group where *aux* clusters are highly prevalent across the strains and another group where *aux* clusters are less common (Figure 4C). *aux1-3*, *aux6*, and *aux14* were present in at least one genome of every phylotype, excluding the mostly T6SS[null] phylotype III. *aux4, aux9*, *aux10*, and *aux22* were found in each of the remaining three mostly T6SS[+] phylotypes. In contrast, some *aux* types were restricted to particular phylotypes: *aux13* and *aux19* were present only in phylotype I, while *aux12* and *aux25* are only found in phylotype II. Additionally, there were species-specific patterns for some *aux* clusters. *aux15* and *aux22* are present in *R. pseudosolanacearum* (phylotype I and III) but absent in *R. solanacearum* (phylotype II) and *R. syzygii* (phylotype IV). *aux24* and *aux25* are rare and were only identified in two and one RSSC genomes, respectively (Figure 4C). As a result of these highly variable patterns of *aux* cluster distribution, closely related strains typically have overlapping but non-identical *aux* cluster repertoires (Figure 4C).

aux clusters are enriched on the megaplasmid.

In bacterial genomes, secondary replicons like the RSSC megaplasmid often contain more accessory genes which evolve more quickly than chromosomal genes (64–66). Because *aux* clusters are part of the RSSC accessory genome, we hypothesized that they would be more common on the megaplasmid. We found that *aux* clusters were enriched on the megaplasmid, accounting for 71.6% of the 1069 *aux* clusters that we classified (Figure 5, Figure S27, and Table S3). Considering that the megaplasmid is smaller than the chromosome (approximately 2.1 Mb and 3.5 Mb, respectively), *aux* cluster density is dramatically higher on the megaplasmid. Although most *aux* types are more common on the megaplasmid, certain *aux* types were more common on the chromosome: *aux2, aux8, aux21, aux19, aux4, aux17,* and *aux13* (Figure 5B and Table S3).

To determine whether there are genomic islands where *aux* clusters are located, we used Clinker synteny analyses to map the specific location of each of the 1069 *aux* clusters. Except for the T6SS main locus, there were no obvious hotspots where *aux* clusters were located (Figure 5C and panel C of Figure S4-S24). The three variable regions of the T6SS main locus each contained *aux* clusters that had the same superfamily of effectors. The variable region downstream of *tssM* only contained *aux15* or *aux18*, which had lipase effectors (Figure S18). The variable region between *tssH* and *tssA* exclusively contained *aux14* and/or *aux20*, which have a MIX_III domain effector (Figure S17). All *aux* clusters found downstream of *tssJKL* contained effectors with FIX domains, including *aux22-24*, which were exclusively found at T6SS main, and *aux10* and *aux17*, which were also found associated with the T6SS main cluster as well as additional loci (Figure S13, S20). Certain *aux* types were located in consistent genomic loci while others were highly variable. For instance, *aux3* is found in 79% of surveyed RSSC genomes and is always found in

the same locus on the megaplasmid (Figure S6). In contrast, *aux2* is found in 84% of our genome set, but it is located in nine distinct loci (Figure S5).

Some of our RSSC genomes included accessory plasmids that encoded two different aux types: *aux6* and *aux11*. However, both of these *aux* types were also found on the chromosome or megaplasmid in other RSSC genomes (Figure S9, S14).

Loss-of-function mutations are common among *aux* clusters.

To infer the role of gene loss in the evolutionary history of *aux* clusters, we used Clinker synteny analyses to identify apparently functionally "intact" *aux* clusters and "decayed" clusters with one or more putative loss-of-function mutations (Figure S28). Loss-of-function mutations included large deletions, frameshifts, premature stop codons, or disruption by insertion sequence (IS) elements (Table S3). Of the 1069 *aux* clusters classified, about 23.5% contained one or more *vgrG*, effector, or immunity genes that had one of these loss-of-function mutations (Table S3). Of the 251 decayed *aux* clusters, 70.5% had mutations in *vgrG*, 34.7% had mutated effectors, and 17.5% mutated immunity genes. Only 8.8% of *aux* clusters had mutations in the immunity gene while retaining evidently functional effectors and *vgrG* genes. Some *aux* types were more frequently decayed than others (Figure 4A, Figure S26, and Table S3). Clusters with nuclease effectors (*aux3, aux8, aux10, aux12, aux16, aux19, and aux21*) rarely decayed, as well as *aux13* and *aux23* whose effectors target unknown substrates. The most commonly decayed *aux* clusters were *aux1* (43/88 decayed), *aux2* (42/115 decayed) and *aux14* (36/75 decayed).

Most T6SS⁺ RSSC genomes have at least one decayed *aux* cluster. We investigated whether RSSC lineages varied in their proportion of decayed *aux* clusters, which could suggest that these lineages had less ecological pressure to maintain large toxin repertoires. As expected, the T6SS^null lineages (III, IV-BDB, and IV-SDC) mostly lack intact *aux* clusters although the BDB genomes had decayed *aux* clusters, and two T6SS^null phylotype III strains had intact *aux6* and *aux21* clusters (Figure S26, S29, and Table S3). Although phylotype IIB is T6SS⁺, IIB genomes encode fewer intact *aux* clusters than other T6SS⁺ clades, with a median of 5 intact *aux* clusters per genome (Figure 4B-C). A small clade within phylotype I that had five decayed *aux* clusters per genome (Figure 4, S26). Overall, there is no particularly strong phylogenetic pattern to the prevalence of decayed *aux* clusters.

Mobile genetic elements facilitate horizontal acquisition of chromosomal *aux* clusters.

The phylogenetic distribution of *aux* clusters among RSSC genomes suggests a complicated pattern of gene flow with frequent gain events in addition to the loss events documented above. We hypothesized that horizontal gene transfer (HGT) between RSSC clades may explain the convoluted phylogenetic pattern of *aux* cluster presence and absence. In bacteria, mobile genetic elements like phages and conjugative plasmids are common vehicles for the horizontal transmission of genes (67). To investigate if mobile genetic elements (MGEs) were associated with T6SS genes in RSSC genomes, we ran a combination of bioinformatic methods on the *aux* clusters, including genetic neighborhood visualization and comparison with Clinker (68), prophage prediction with Phaster (69) and domain analysis with NCBI CDD (60).

Of the 72 unique genetic neighborhoods around *aux* clusters, we classified 47.2% as MGE-associated, 44.4% as not MGE-associated, and the rest as inconclusive. We assigned the "inconclusive" status if there was minor but insufficient evidence that the cluster is associated with an MGE. For example, one cluster was encoded next to a single pseudogenized phage portal gene. We identified prophages (Myoviridae, Inoviridae, and Siphoviridae families), conjugative plasmids, and 16 other unclassified MGEs that were co-inherited with *aux* clusters (Figure 6A, Table S3). Of the prophages, the Myoviridae and

Siphoviridae phages had highly conserved gene structure, whereas the filamentous Inoviridae phages were highly diverse.

Many of the MGEs were associated with lipase *aux* types. The φRSY1-like prophages in the Myoviridae family carried certain lipase clusters: *aux5* at 3/6 locations, *aux7*, and *aux9* at 2/6 locations (70). Siphoviridae prophages carried a lipase cluster and two clusters with unknown targets: *aux5*, *aux13*, and *aux16* at 3/3 locations. An unclassified 8-gene MGE with a Toprim topoisomerase carried two lipase clusters *aux1* and *aux7* at 3/8 locations. Finally, the conjugative plasmids carried two lipase clusters *aux6* and *aux11*. Inoviridae prophages carried clusters with diverse toxins *aux4, aux6, aux10, aux17* at 2/5 locations*, aux19,* and *aux21*.

Several of the unclassified MGE contain anti-phage defense systems, such as a type 2 bacteriophage exclusion (BREX) system with the *aux21^{loc4}* MGE and as DISARM restriction-modification system with the *aux6^{loc4}* MGE (71, 72).

We investigated the genomic locations of MGE-associated, inconclusive, and non-MGE *aux* clusters. Surprisingly, we found that there was a strong linkage between chromosomal location and MGE-association (Figure 6B). Whereas 27/32 chromosomal *aux* clusters were MGE-associated, only 5/38 megaplasmid *aux* clusters were MGE-associated. All accessory plasmids with aux clusters were conjugative plasmid MGEs. We identified several hotspots on chromosome where *aux*-carrying MGEs inserted, and most of the hotspots were dominated by a single prophage family (Figure 6C). For example, hotspot B was dominated by the Siphoviridae phages, while C and D were both dominated by Inoviridae prophages. In contrast, hotspot A was occupied by four distinct unclassified MGEs.

Discussion

Bacteria in the RSSC are aggressive pathogens that cause lethal wilt disease in plants, resulting in drastic losses of economically important crops. RSSC are renowned for manipulating a broad range of host plants with massive repertoires of 60-80 type 3 secreted effectors (73). Here, we document that RSSC are also evolutionary enriched in a large arsenal of type 6 secreted toxins. The high prevalence of T6SS genes in RSSC genomes suggests that possessing a T6SS is adaptive for the lifestyle of these pathogens. RSSC pathogens pass through a variety of microbial ecosystems to complete their disease cycle. When transmitting and infecting plants, they encounter a variety of ecological interactors. We speculate that the RSSC have an ecological role similar to spotted hyenas. Similar to how hyenas attack with their teeth and claws, RSSC bacteria may wield their T6SS against competitors (other RSSC and plant-colonizing bacteria), predators (bacterivorous amoebae) and, possibly, their prey (plant hosts).

Our data support the model that there is eco-evolutionary feedback between bacterial lifestyle and enrichment of T6SS-reated genes (26). We discovered that the RSSC and the *Acidovorax citrulli* species complex, two taxa of ecologically similar plant pathogens, have convergently evolved to deploy a large array of effectors from a single T6SS. These pathogens both cause acute infections of the plant xylem (74, 75). In contrast, plant symbiotic *Rhizobia* and *Agrobacterium* cause chronic infections with a spectrum of mutualistic to pathogenic outcomes (76–78). These chronic symbionts have only a single T6SS and only a handful of effectors, which suggests a reduced need to combat diverse microbes and a greater emphasis on reducing collateral damage to the host. In contrast, other Burkholderiaceae encode up to 6 distinct T6SSs in their genomes (79, 80), which is consistent with a model that organisms with complex lifestyles

benefit from having multiple T6SSs that are independently regulated and specialized for different targets and contexts.

The relationship between bacterial lifestyle and T6SS gene content is also evident within the RSSC. Most RSSC pathogens are transmitted through soil or fresh water. However, two of the phylotype IV clades that lack a T6SS have adopted novel lifestyles: the strains causing Sumatra Disease of Clove (SDC) and Blood Disease of Banana (BDB) (81, 82). SDC strains are spread by piercing sucking insects (83) while BDB is mechanically transmitted by insects or agricultural tools (84). While the two SDC-lineage genomes lacked *aux* clusters, they encoded two T6SS core genes (*gh10* and *ompA*). Similarly, all three genomes in the BDB lineage lacked T6SS core genes but contained pseudogenized *aux* clusters. Overall, this pattern suggests that the T6SS is ancestral to the RSSC and has been recently lost in T6SS^{null} lineages. Like many bacteria that transition to a host-restricted lifestyle, these lineages have undergone a reduction in genome size. While SDC and BDB strains have chromosomes of comparable size to soil-transmitted relatives, their megaplasmids are reduced in size (85).

Although the T6SS is widely conserved in soil-transmitted RSSC, the absence of the T6SS in several soil-transmitted lineages demonstrates that the T6SS is non-essential for this lifestyle. Soil-transmitted RSSC that lack a T6SS included all but one of the phylotype III genomes and a minor clade of phylotype I (Figure 2C). Three of the T6SS^{null} phylotype III strains had intact *aux* clusters, which could suggest that the T6SS was recently lost from this lineage. However, two of the three intact *aux* clusters were associated with mobile genetic elements (MGEs), so it is alternatively possible these *aux* clusters were recently gained through HGT after the loss of the T6SS.

Within phylotype II, several RSSC lineages cause Moko disease of bananas and plantains. Pioneering studies on Moko disease by Luis Sequeira and Ivan Buddenhagan demonstrated that the causal pathogens are facultatively transmitted by either soil transmission or contact transmission of insects with sap, similar to BDB (86, 87). We and other groups have sequenced isolates from the Sequiera and Buddenhagan collection from the 1960s Moko epidemic (88–91), and we now know that multiple RSSC lineages were responsible for the epidemic, including T6SS$^+$ lineages (IIB-4 and IIB-3) and T6SS^{null} lineages (IIA-6). Further epidemiological research is needed to understand how variation in transmission routes shapes the evolution and behavior of RSSC lineages.

Our results provide new insight into the epidemiology of RSSC in regions with multiple lineages. It has long been known that RSSC strains can inhibit each other's growth and competitively exclude each other *in planta* (92, 93). A thorough epidemiological survey of Malagasy vegetable plots demonstrated that T6SS$^+$ phylotype I strains are displacing the T6SS^{null} phylotype III strains native to the island (94). Subsequent functional analysis demonstrated that the phylotype I Malagasy strains secrete bacteriocin toxins that inhibit growth of the phylotype III strains (95). Our results suggest that the T6SS may confer an additional advantage to phylotype I strains when directly competing against T6SS^{null} phylotype III strains *in planta*.

The convoluted phylogenetic distribution of *aux* clusters suggests that RSSC populations have dynamic T6SS gene flow with frequent gain and loss events. Although there is evidence that certain *aux* types have been primarily vertically inherited, most *aux* clusters are clearly horizontally inherited based on their phylogenetic distribution patterns and the context of their genetic neighborhoods. Notably, the φRSY1-like Myoviridae prophages definitively transfer *aux* clusters. φRSY1 was originally from soil from a

Ralstonia-infested field, and whole genome sequencing of the purified virion particles confirms that the genomes of φRSY1-like phages encode *aux* clusters and, thus, can transmit *aux* clusters.

Like many bacteria in the Burkholderiaceae, RSSC have a bipartite genome. When bacteria have a secondary replicon like the megaplasmid, the genes on the secondary replicon evolve more quickly than chromosomal genes (64–66). Intriguingly, we discovered that *aux* clusters are dramatically enriched on the megaplasmid. We speculate that the enrichment of *aux* clusters on the evolutionarily dynamic megaplasmid could allow RSSC lineages to rapidly diversify with respect to their intraspecies and interspecies competitive behavior. In-depth studies on the molecular evolution of *aux* cluster genes are needed to understand if they evolve at different rates than other RSSC genes and if there are differences in the evolutionary behavior of megaplasmid and chromosomal MGE-associated *aux* clusters.

In *Ralstonia*'s bipartite genome, we found clear distinctions in the mechanisms for horizontal gene transfer for *aux* clusters on the chromosome vs. megaplasmid. Chromosomal *aux* clusters were predominantly MGE-associated, which is consistent with prior reports that RSSC prophages are moderately enriched on the evolutionarily stable chromosome and that they target specific insertion sequences (96). In contrast, megaplasmid *aux* clusters were rarely associated with MGEs. Nevertheless, the patterns of phylogenetic distribution suggest that both chromosomal and megaplasmid *aux* clusters have been horizontally transmitted. This opens the question—what genetic mechanisms contributed to horizontal gene flow of megaplasmid *aux* clusters? RSSC are naturally competent (97), so they could readily acquire genes by uptake of environmental DNA and integration by homologous recombination. However, it is surprising that there would be a bias towards homologous recombination for the megaplasmid.

The presence of *aux* clusters in the genomes of phages indicates that these phages may function as mutualists of RSSC. Across the diversity of RSSC-infective phages, most do not transport *aux* clusters (96, 98–100). Nevertheless, with growing interest in employing phages for control of bacterial plant pathogens (99, 101–105), it will remain important to evaluate candidate biocontrol agents for their ability to improve the ecological fitness of the pathogens.

In closing, we propose that evolution has positioned the RSSC to be an ideal model to understand the role of the T6SS in shaping pathogen populations and host-associated ecosystems. Our systematic analysis opens a plethora of evolutionarily grounded questions for future investigation. For example, do RSSC pathogens target novel cellular targets with their nine effector families that lack known toxin domains? Are there genetic and epigenetic mechanisms that promote preferential integration of *aux* clusters on the megaplasmid? Moreover, what other physiological functions are enriched on the megaplasmid and secondary replicons in other bacteria with multipartite genomes? RSSC genomes also encode large repertoires of other genes that allow them to sense and change their environments, including root-sensing MCP chemoreceptors (106), plant-manipulating T3SS effectors (73), and anti-phage defense systems (107). Do these genes exhibit biased distribution across the replicons? We anticipate that this study will fuel many new discoveries in T6SS biology and pathogen evolution.

Materials and Methods

Identification of genes encoding the type 6 secretion system (T6SS) machinery in RSSC genomes

All publicly available genomes in the genus *Ralstonia*, including *R. syzygii* genomes deposited as "Blood Disease Bacterium" strains, were downloaded from NCBI and uploaded to KBase (108) for analysis.

Genomes were analyzed with the Genome Taxonomy Database Toolkit GTDB-Tk - *v1.7.0* to identify the genomospecies (34–39). We used CheckM - *v1.0.18* to evaluate the completeness and contamination of the assemblies. Only genomes with completeness greater than 99.82 % and contamination less than 0.96 % were retained for further analysis. We also excluded assemblies with 100 or more contigs. With the selected assemblies we generated a phylogenetic tree of RSSC using Insert Genome Into SpeciesTree – *v.2.2.0.* The KBase SpeciesTree uses 49 genes broadly conserved across bacteria to build the phylogenetic tree. The protein sequence for each GMI1000 T6SS component was queried against the *Ralstonia* genomes using BLASTp - *v2.13.0.* We considered all BLASTp results with ≥ 20% identity and ≥ 80% coverage to be hits. We analyzed phylogenetic patterns of T6SS gene presence or absence in the RSSC phylogenetic tree on iTOL (109).

We selected high quality genomes for detailed analysis of the repertoires of *vgrG*-linked effector/immunity gene clusters. Limiting the analysis to complete genomes would have excluded most phylotype II, III, and IV genomes, so we included any genome assembled into as many as 28 contigs. We carried out a series of low stringency BLASTp searches against the RSSC genomes with VgrG protein sequences from phyl. I GMI1000, phyl. II IBSBF1503, and phyl. IV PSI07 (Parameters: ≥ 1% identity, ≥ 1% coverage, and Bit Score ≥ 10). Vgr BLASTp results were exported and further analyzed in Excel.

Classifying the Vgr-linked effector/immunity gene clusters through synteny analysis
We downloaded the genetic neighborhood of each *vgrG* gene from NCBI. The gene locus was queried against the NCBI Nucleotide database. From the corresponding Graphics page, we downloaded a Genbank Flat file that encompassed a 10-200 kb region surrounding the *vgrG* gene. The .flat files were converted to .gbk, and the synteny analysis and visualization tool Clinker was iteratively used to identify gene clusters with shared genetic architecture (68). *vgrG*-linked clusters were named *aux1-aux25,* which is short for auxiliary cluster because these effector/immunity clusters are not core T6SS genes. Per genome, the set of *vgrG*-linked clusters were matched to references for each auxiliary cluster (*aux1-aux25*) using Clinker (68).

Because the VgrG BLASTp search would not identify *aux* clusters with pseudogenized or missing *vgrG* genes, we carried out further BLASTp queries with the effector and immunity protein sequences. We exported the gene neighborhoods from NCBI and used Clinker to confirm the *aux*-type of each cluster.

Identification of T6SS and *vgrG* genes in representative Burkholderiaceae genomes
The Genome Taxonomy Database (GTDB) (34–39) was used to identify complete genomes in the Burkholderiaceae family. GTDB classifies genomes into genomospecies based on a 95% ANI threshold. Per GTDB species, we selected one representative genome to import into a public KBase Narrative (https://narrative.kbase.us/narrative/142785). We used BLASTp as described above to identify T6SS_main genes and VgrG proteins. The full results are present in Table S1.

Classifying the genetic architecture of T6SS_main loci across the *Ralstonia* genus
Clinker was used to visually compare the genetic architectures of T6SS_main loci. We downloaded the genetic neighborhood of each T6SS_main locus from NCBI and compared them to reference T6SS_main loci from plant-colonizing bacteria (49, 110). The phylogenetic distributions of the T6SS subtypes were visualized with iTol (109).

Identifying effectors, immunity proteins, and adaptors encoded in each *aux* cluster
We queried protein sequences ("WP_" accessions) against the literature using PaperBlast (61) and the NCBI Conserved Domain Database (CDD) (60) using CD-Search. Effectors were identified based on homology with bona fide T6SS effectors (e.g., PaperBlast matches to Tle1, Tle3, Tle4, and Tle5

phospholipases). Immunity proteins were identified based on homology to bona fide immunity proteins (PaperBlast) and/or the presence of known immunity domains. Adapters were identified based on the presence of domains like DUF1795, DUF2169, or DUF4123. Sometimes the VgrG or effector encoded a PAAR spike motif and sometimes the cluster had a small, standalone PAAR-containing protein.

Identifying genetic decay in the clusters and orphan immunity genes

We recorded which genes were annotated as pseudogenized. Moreover, the Clinker visualizations enabled us to identify transposons and/or insertion sequences that had been inserted into and disrupted genes. The visualizations revealed if an effector or *vgrG* gene was conspicuously absent from a cluster.

Identifying the genomic location of each *vgrG*-linked *aux* cluster.

Clinker was used to identify which *aux* clusters were in the same genomic location across genomes. Per *aux*-type (i.e., all "*aux1*" clusters), Clinker was run, and unique auxiliary loci were identified and named. Each auxiliary locus was mapped to its relative position with respect to the GMI1000 genome by BLAST-searching neighboring genes against the GMI1000 genome and retrieving a Genbank Flat file of the approximate region. Auxiliary loci and their mapped location are referred to as "*aux1-loc1*" and "*aux1-loc2*". Synteny between the GMI1000 sequence and the *aux* cluster's neighborhood was visualized with Clinker. If GMI1000 encoded the exact cluster, the locus for the *vgrG* gene was recorded. If GMI1000 lacked the cluster, the nearest GMI1000 gene upstream of the *vgrG* was recorded.

When downloading Genbank Flat files from complete genomes, we recorded whether the cluster was encoded on the chromosome, megaplasmid, or a small accessory plasmid. For incomplete genomes, we inferred whether the cluster was on the chromosome or megaplasmid by matching the cluster's *aux-loc* to a syntenous cluster from a complete genome.

To visualize the relative location of each cluster across the replicons (Figure S4-S24), we used Clinker to compare a representative *aux* cluster from each of the 90 *aux-loc* subtypes to the relative region in GMI1000. The GMI1000 gene upstream of each VgrG was recorded as the "nearby location marker" in Table S3.

Identifying *aux* clusters diversity in RSCC

To assess *aux* clusters diversity in RSSC we measured Bray-Curtis dissimilarity matrix which accounts for the presence, absence, and relative abundance of *aux* clusters. We used 'vegdigest' of the 'vegan v.2.6-2' package in RStudio (version 2023.03.0+386) to compute the Bray-Curtis dissimilarity matrix. With this matrix, we constructed an UPGMA tree with 'hclust' of the 'Phangorn v.2.11.1' in RStudio (version 2023.03.0+386).

Identifying mobile genetic elements associated with *aux* clusters

To identify MGEs associated with various T6SS auxiliary clusters, we started with BLASTp searches on the *aux* clusters with ~100 Kb on each side of their genetic neighborhoods in PHASTER identification and annotation of prophage tool (69). In parallel, we checked gene annotations of the genetic neighborhoods. To check if hypothetical proteins could be associated to mobile genetic elements, we checked for the presence of any mobile genetic elements-related domains on the Conserved Domain Database of NCBI (60). Based on this analysis we recorded the presence, inconclusive, or absence of MGE on the chromosome, megaplasmid, and the conjugative plasmid (when present). The barplot showing the proportions of the presence, inconclusive, or absence of MGE (Figure 6) was built using GraphPad Prsim (version 10.0.1). We highlighted the *aux* clusters associated with MGE in orange using AFFINITY Designer

(version 1.10.6). *aux* clusters and corresponding locations were summarized on the chromosome, megaplasmid, and conjugative plasmid maps using AFFINITY Designer (version 1.10.6). The synteny

Acknowledgments

We would like to thank Dr. Jonathan Jacobs for insightful recommendations about senteny analysis. We would also like to thank Dr. Anneliek ter Horst, Dr. Joanne Emerson, and Dr. Poliane Alfenas-Zerbini for sharing expertise in phage genomics.

References

1. Raaijmakers JM, Mazzola M. 2012. Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. Annu Rev Phytopathol 50:403–424.

2. Klein TA, Ahmad S, Whitney JC. 2020. Contact-dependent interbacterial antagonism mediated by protein secretion machines. Trends Microbiol 28:387–400.

3. Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ. 2006. Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. Proc Natl Acad Sci U S A 103:1528–1533.

4. Bingle LE, Bailey CM, Pallen MJ. 2008. Type VI secretion: a beginner's guide. Curr Opin Microbiol 11:3–8.

5. Durand E, Cambillau C, Cascales E, Journet L. 2014. VgrG, Tae, Tle, and beyond: the versatile arsenal of type VI secretion effectors. Trends Microbiol 22:498–507.

6. Alcoforado Diniz J, Liu Y-C, Coulthurst SJ. 2015. Molecular weaponry: diverse effectors delivered by the Type VI secretion system. Cell Microbiol 17:1742–1751.

7. Flaugnatti N, Le TTH, Canaan S, Aschtgen MS, Nguyen VS, Blangy S, Kellenberger C, Roussel A, Cambillau C, Cascales E, Journet L. 2016. A phospholipase A_1 antibacterial Type VI secretion effector interacts directly with the C-terminal domain of the VgrG spike protein for delivery. Mol Microbiol 99:1099–1118.

8. Russell AB, Hood RD, Bui NK, LeRoux M, Vollmer W, Mougous JD. 2011. Type VI secretion delivers bacteriolytic effectors to target cells. Nature 475:343–349.

9. Hood RD, Singh P, Hsu F, Güvener T, Carl MA, Trinidad RRS, Silverman JM, Ohlson BB, Hicks KG, Plemel RL, Li M, Schwarz S, Wang WY, Merz AJ, Goodlett DR, Mougous JD. 2010. A type VI secretion system of *Pseudomonas aeruginosa* targets a toxin to bacteria. Cell Host Microbe 7:25–37.

10. Cascales E, Cambillau C. 2012. Structural biology of type VI secretion systems. Philos Trans R Soc B Biol Sci 367:1102–1111.

11. Coulthurst SJ. 2013. The Type VI secretion system - a widespread and versatile cell targeting system. Res Microbiol 164:640–654.

12. Brunet YR, Zoued A, Boyer F, Douzi B, Cascales E. 2015. The type VI secretion TssEFGK-VgrG phage-like baseplate is recruited to the TssJLM membrane complex via multiple contacts and serves as assembly platform for tail tube/sheath polymerization. PLoS Genet 11:1–21.

13. Ho BT, Dong TG, Mekalanos JJ. 2014. A view to kill: the bacterial type 6 secretion system. Cell Host

Microbe 15:9–21.

14. Bönemann G, Pietrosiuk A, Diemand A, Zentgraf H, Mogk A. 2009. Remodelling of VipA/VipB tubules by ClpV-mediated threading is crucial for type VI protein secretion. EMBO J 28:315–325.

15. Basler M, Mekalanos JJ. 2012. Type 6 secretion dyramics within and between bacterial cells. Science 337:1–7.

16. Shneider MM, Buth SA, Ho BT, Basler M, Mekalanos JJ, Leiman PG. 2013. PAAR-repeat proteins sharpen and diversify the type VI secretion system spike. Nature 500:350–353.

17. Jana B, Fridman CM, Bosis E, Salomon D. 2019. A modular effector with a DNase domain and a marker for T6SS substrates. Nat Commun 10.

18. Crisan C V., Chande AT, Williams K, Raghuram V, Rishishwar L, Steinbach G, Watve SS, Yunker P, Jordan IK, Hammer BK. 2019. Analysis of *Vibrio cholerae* genomes identifies new type VI secretion system gene clusters. Genome Biol 20:1–14.

19. Borgeaud S, Metzger LC, Scrignari T, Blokesch M. 2015. The type VI secretion system of *Vibrio cholerae* fosters horizontal gene transfer. Science 347:63–67.

20. Salomon D, Klimko JA, Trudgian DC, Kinch LN, Grishin N V, Mirzaei H, Orth K. 2015. Type VI secretion system toxins horizontally shared between marine bacteria. PLoS Pathog 11:e1005128.

21. Peñil-Celis A, Garcillán-Barcia MP. 2019. Crosstalk between type VI secretion system and mobile genetic elements. Front Mol Biosci 6:126.

22. Vassallo CN, Troselj V, Weltzer ML, Wall D. 2020. Rapid diversification of wild social groups driven by toxin-immunity loci on mobile genetic elements. ISME J 14:2474–2487.

23. Russell AB, Peterson SB, Mougous JD. 2014. Type VI secretion system effectors: poisons with a purpose. Nat Rev Microbiol 12:137–148.

24. Alcoforado Diniz J, Coulthurst SJ. 2015. Intraspecies competition in *Serratia marcescens* is mediated by type VI-secreted Rhs effectors and a conserved effector-associated accessory protein. J Bacteriol 197:2350–2360.

25. Speare L, Cecere AG, Guckes KR, Smith S, Wollenberg MS, Mandel MJ, Miyashiro T, Septer AN. 2018. Bacterial symbionts use a type VI secretion system to eliminate competitors in their natural host. Proc Natl Acad Sci U S A 115:E8528–E8537.

26. Geller AM, Zlotkin D, Levy A. 2021. Large-scale discovery of candidate type VI secretion effectors with antibacterial activity. bioRxiv 2021.10.07.463556.

27. Zhang L, Xu J, Xu J, Zhang H, He L, Feng J. 2012. TssM is essential for virulence and required for type VI secretion system in *Ralstonia solanacearum*. Microb Pathog 119:125–134.

28. Zhang L, Xu J, Xu J, Zhang H, He L, Feng J. 2014. TssB is essential for virulence and required for Type VI secretion system in *Ralstonia solanacearum*. Microb Pathog 74:1–7.

29. Bocsanczy AM, Achenbach UCM, Mangravita-Novo A, Chow M, Norman DJ. 2014. Proteomic comparison of *Ralstonia solanacearum* strains reveals temperature dependent virulence factors. BMC Genomics 15:1–14.

30. Yao J, Allen C. 2006. Chemotaxis is required for virulence and competitive fitness of the bacterial

wilt pathogen *Ralstonia solanacearum*. J Bacteriol 188:3697–3708.

31. Lowe-Power TM, Khokhani D, Allen C. 2018. How *Ralstonia solanacearum* exploits and thrives in the flowing plant xylem environment. Trends Microbiol 26:929–942.

32. Ingel B, Caldwell D, Duong F, Parkinson DY, McCulloh KA, Iyer-Pascuzzi AS, McElrone AJ, Lowe-Power TM. 2022. Revisiting the source of wilt symptoms: X-ray microcomputed tomography provides direct evidence that *Ralstonia* biomass clogs xylem vessels. PhytoFrontiers 2:41–51.

33. Khokhani D, Lowe-Power TM, Tran TM, Allen C. 2017. A single regulator mediates strategic switching between attachment/spread and growth/virulence in the plant pathogen *Ralstonia solanacearum*. MBio 8:1–20.

34. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. Nat Biotechnol 36:996.

35. Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ, Hugenholtz P. 2020. A complete domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol 38:1079–1086.

36. Rinke C, Chuvochina M, Mussig AJ, Chaumeil PA, Davín AA, Waite DW, Whitman WB, Parks DH, Hugenholtz P. 2021. A standardized archaeal taxonomy for the Genome Taxonomy Database. Nat Microbiol 6:946–959.

37. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. 2022. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res 50:D785–D794.

38. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2020. GTDB-Tk: A toolkit to classify genomes with the genome taxonomy database. Bioinformatics 36:1925–1927.

39. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2022. GTDB-Tk v2: memory friendly classification with the genome taxonomy database. Bioinformatics 38:5315–5316.

40. Gallique M, Bouteiller M, Merieau A. 2017. The type VI secretion system: a dynamic system for bacterial communication? Front Microbiol 8:1454.

41. Schell MA, Ulrich RL, Ribot WJ, Brueggemann EE, Hines HB, Chen D, Lipscomb L, Kim HS, Mrázek J, Nierman WC, DeShazer D. 2007. Type VI secretion is a major virulence determinant in *Burkholderia mallei*. Mol Microbiol 64:1466–1485.

42. Schwarz S, West TE, Boyer F, Chiang WC, Carl MA, Hood RD, Rohmer L, Tolker-Nielsen T, Skerrett SJ, Mougous JD. 2010. Burkholderia type VI secretion systems have distinct roles in eukaryotic and bacterial cell interactions. PLoS Pathog 6:77–78.

43. Spiewak HL, Eberl L, Shastri S, Zhang L, Schwager S, Thomas MS, Vergunst AC. 2018. *Burkholderia cenocepacia* utilizes a type VI secretion system for bacterial competition. Microbiologyopen 8:1–26.

44. de Campos SB, Lardi M, Gandolfi A, Eberl L, Pessi G. 2017. Mutations in two *Paraburkholderia phymatum* type VI secretion systems cause reduced fitness in interbacterial competition. Front Microbiol 8:1–11.

45. Pehl MJ, Jamieson WD, Kong K, Forbester JL, Fredendall RJ, Gregory GA, Mcfarland JE, Healy JM,

Orwin PM. 2012. Genes that influence swarming motility and biofilm formation in *Variovorax paradoxus* EPS. PLoS One 7.

46. Dong TG, Ho BT, Yoder-Himes DR, Mekalanos JJ. 2013. Identification of T6SS-dependent effector and immunity proteins by Tn-seq in *Vibrio cholerae*. Proc Natl Acad Sci U S A 110:2623–2628.

47. Hachani A, Allsopp LP, Oduko Y, Filloux A. 2014. The VgrG proteins are "à la carte" delivery systems for bacterial type VI effectors. J Biol Chem 289:17872–17884.

48. Liang X, Moore R, Wilton M, Wong MJQ, Lam L, Dong TG. 2015. Identification of divergent type VI secretion effectors using a conserved chaperone domain. Proc Natl Acad Sci 112:9106–9111.

49. Gallegos-Monterrosa R, Coulthurst SJ. 2021. The ecological impact of a bacterial weapon: Microbial interactions and the type VI secretion system. FEMS Microbiol Rev 45:1–18.

50. Boyer F, Fichant G, Berthod J, Vandenbrouck Y, Attree I. 2009. Dissecting the bacterial type VI secretion system by a genome wide *in silico* analysis: what can be learned from available microbial genomic resources? BMC Genomics 10:1–14.

51. Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, Billault A, Brottier P, Camus JC, Cattolico L, Chandler M, Choisne N, Claudel-Renard C, Cunnac S, Demange N, Gaspin C, Lavie M, Moisan A, Robert C, Saurin W, Schiex T, Siguier P, Thébault P, Whalen M, Wincker P, Levy M, Weissenbach J, Boucher CA. 2002. Genome sequence of the plant pathogen *Ralstonia solanacearum*. Nature 415:497–502.

52. Liu Y, Zhang Z, Wang F, Li D-D, Li Y-Z. 2020. Identification of type VI secretion system toxic effectors using adaptors as markers. Comput Struct Biotechnol J 18:3723–3733.

53. Kim N, Han G, Jung H, Lee H, Park J, Seo Y. 2022. T6SS accessory proteins, including DUF2169 domain-containing protein and pentapeptide repeats protein, contribute to bacterial virulence in T6SS Group_5 of *Burkholderia glumae* BGR1. Plants 11.

54. Unterweger D, Kostiuk B, Ötjengerdes R, Wilton A, Diaz-Satizabal L, Pukatzki S. 2015. Chimeric adaptor proteins translocate diverse type VI secretion system effectors in *Vibrio cholerae*. EMBO J 34:2198–2210.

55. Ma LS, Hachani A, Lin JS, Filloux A, Lai EM. 2014. Agrobacterium tumefaciens deploys a superfamily of type VI secretion DNase effectors as weapons for interbacterial competition *in planta*. Cell Host Microbe 16:94–104.

56. Salomon D, Kinch LN, Trudgian DC, Guo X, Klimko JA, Grishin N V, Mirzaei H. 2014. Marker for type VI secretion system effectors. Proc Natl Acad Sci 111:9271–9276.

57. Berni B, Soscia C, Djermoun S, Ize B, Bleves S. 2019. A type VI secretion system trans-kingdom effector is required for the delivery of a novel antibacterial toxin in *Pseudomonas aeruginosa*. Front Microbiol 10:1218.

58. Ray A, Schwartz N, Santos MDS, Zhang J, Orth K, Salomon D. 2017. Type VI secretion system MIX-effectors carry both antibacterial and anti-eukaryotic activities. EMBO Rep 18:1978–1990.

59. Zhang J, Guan J, Wang M, Li G, Djordjevic M, Tai C, Wang H, Deng Z, Chen Z, Ou H. 2023. SecReT6 update: a comprehensive resource of bacterial Type VI Secretion Systems. Sci China Life Sci

66:626–634.

60. Wang J, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu S, Marchler GH, Song JS, Thanki N, Yamashita RA, Yang M, Zhang D, Zheng C, Lanczycki CJ, Marchler-Bauer A. 2023. The conserved domain database in 2023. Nucleic Acids Res 51:D384--D388.

61. Price MN, Arkin AP. 2017. PaperBLAST: text mining papers for information about homologs. mSystems 2.

62. Russell AB, Leroux M, Hathazi K, Agnello DM, Ishikawa T, Wiggins PA, Wai SN, Mougous JD. 2013. Diverse type VI secretion phospholipases are functionally plastic antibacterial effectors. Nature 496:508–512.

63. Champoiseau PG, Jones JB, Allen C. 2009. *Ralstonia solanacearum* Race 3 Biovar 2 causes tropical losses and temperate anxieties. Plant Heal Prog https://doi.org/10.1094/PHP-2009-0313-01-RV.Introduction.

64. Cooper VS, Vohr SH, Wrocklage SC, Hatcher PJ. 2010. Why genes evolve faster on secondary chromosomes in bacteria. PLoS Comput Biol 6.

65. Harrison PW, Lower RPJ, Kim NKD, Young JPW. 2010. Introducing the bacterial "chromid": not a chromosome, not a plasmid. Trends Microbiol 18:141–148.

66. DiCenzo GC, Finan TM. 2017. The divided bacterial genome: structure function, and evolution. Microbiol Mol Biol Rev 81:1–37.

67. Haudiquet M, de Sousa JM, Touchon M, Rocha EPC. 2021. Selfish, promiscuous and sometimes useful: how mobile genetic elements drive horizontal gene transfer in microbial populations. Philos Trans R Soc Lond B Biol Sci 377.

68. Gilchrist CLM, Chooi Y-H. 2021. clinker & clustermap.js: automatic generation of gene cluster comparison figures. Bioinformatics 37:2473–2475.

69. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids Res 44:W16--21.

70. Askora A, Kawasaki T, Fujie M, Yamada T. 2017. Lysogenic conversion of the phytopathogen *Ralstonia solanacearum* by the P2virus ΦRSY1. Front Microbiol 8:1–11.

71. Goldfarb T, Sberro H, Weinstock E, Cohen O, Doron S, Charpak-Amikam Y, Afik S, Ofir G, Sorek R. 2015. BREX is a novel phage resistance system widespread in microbial genomes. EMBO J 34:169–183.

72. Ofir G, Melamed S, Sberro H, Mukamel Z, Silverman S, Yaakov G, Doron S, Sorek R. 2018. DISARM is a widespread bacterial defence system with broad anti-phage activities. Nat Microbiol 3:90–98.

73. Peeters N, Carrère S, Anisimova M, Plener L, Cazalé AC, Genin S. 2013. Repertoire, unified nomenclature and evolution of the Type III effector gene set in the *Ralstonia solanacearum* species complex. BMC Genomics 14.

74. Schaad NW, Sowell JR. G, Goth RW, Colwell RR, Webb RE. 1978. *Pseudomonas pseudoalcaligenes* subsp. *citrulli* subsp. nov. Int J Syst Bacteriol 28:117–125.

75. Burdman S, Walcott R. 2012. *Acidovorax citrulli*: generating basic and applied knowledge to tackle

a global threat to the cucurbit industry. Mol Plant Pathol 13:805–815.

76.   van Rijn P, Vanderleyden J. 1995. The *Rhizobium*-plant symbiosis. Microbiol Rev 59:124–142.

77.   Escobar MA, Dandekar AM. 2003. *Agrobacterium tumefaciens* as an agent of disease. Trends Plant Sci 8:380–386.

78.   Nishiguchi MK, Hirsch AM, Devinney R, Vedantam G, Riley MA, Mansky LM. 2008. Deciphering evolutionary mechanisms between mutualistic and pathogenic symbioses. Vie Milie 58:87–106.

79.   Shalom G, Shaw JG, Thomas MS. 2007. *In vivo* expression technology identifies a type VI secretion system locus in *Burkholderia pseudomallei* that is induced upon invasion of macrophages. Microbiology 153:2689–2699.

80.   Angus AA, Agapakis CM, Fong S, Yerrapragada S, Estrada-de Los Santos P, Yang P, Song N, Kano S, Caballero-Mellado J, De Faria SM, Dakora FD, Weinstock G, Hirsch AM. 2014. Plant-associated symbiotic *Burkholderia* species lack hallmark strategies required in mammalian pathogenesis. PLoS One 9.

81.   Roberts SJ, Eden-Green SJ, Jones P, Ambler DJ. 1990. *Pseudomonas syzygii*, sp. nov., the cause of Sumatra disease of cloves. Syst Appl Microbiol 13:34–43.

82.   Eden-Green SJ, Sastraatmadja H. 1990. Blood disease of banana confirmed in Java. Infomusa 38:13.

83.   Safni I, Subandiyah S, Fegan M. 2018. Ecology, epidemiology and disease management of *Ralstonia syzygii* in Indonesia. Front Microbiol 9:1–11.

84.   Ray JD, Subandiyah S, Prakoso AB, Rincon-Florez VA, Carvalhais LC, Drenth A. 2022. Transmission of Blood disease in banana. Plant Dis 1–38.

85.   Remenant B, de Cambiaire JC, Cellier G, Jacobs JM, Mangenot S, Barbe V, Lajus A, Vallenet D, Medigue C, Fegan M, Allen C, Prior P. 2011. *Ralstonia syzygii*, the Blood Disease bacterium and some Asian *R. solanacearum* strains form a single genomic species despite divergent lifestyles. PLoS One 6:1–10.

86.   Buddenhagen IW, Elsasser TA. 1962. An insect-spread Bacterial wilt epiphytotic of Bluggoe banana. Nature 194:164–165.

87.   French ER, Sequeira L. 1970. Strains of *Pseudomonas solanacearum* from Central and South America: a comparative study. Phytopathology 60:506–512.

88.   Ailloud F, Lowe T, Cellier G, Roche D, Allen C, Prior P. 2015. Comparative genomic analysis of *Ralstonia solanacearum* reveals candidate genes for host specificity. BMC Genomics 16:1–11.

89.   López-Alvarez D, Leiva AM, Barrantes I, Pardo JM, Dominguez V, Cuellar WJ. 2020. Complete genome sequence of the plant pathogen *Ralstonia solanacearum* strain CIAT-078, isolated in Colombia, obtained using Oxford Nanopore technology. Microbiol Resour Announc 9:1–2.

90.   Pais AKL, da Silva JR, Dos Santos LVS, Albuquerque GMR, de Farias ARG, Junior WJS, Balbino V de Q, Silva AMF, da Gama MAS, de Souza EB. 2021. Genomic sequencing of different sequevars of *Ralstonia solanacearum* belonging to the Moko ecotype. Genet Mol Biol 44:1–6.

91.   Beutler J, Holden S, Georgoulis S, Williams D, Norman DJ, Lowe-Power TM. 2023. Whole genome squencing suggests that "nonpathogenicity on banana (NPB)" is the ancestral state of the *Ralstonia*

solanacearum IIB-4 lineage. PhytoFrontiers 6–11.

92. Cuppels DA, Hanson RS, Kelman A. 1978. Isolation and characterization of a bacteriocin produced by *Pseudomonas solanacearum*. J Gen Microbiol 109:195–303.

93. Huerta AI, Milling A, Allen C. 2015. Tropical strains of *Ralstonia solanacearum* outcompete race 3 biovar 2 strains at lowland tropical temperatures. Appl Environ Microbiol 81:3542–3551.

94. Rasoamanana H, Ravelomanantsoa S, Yahiaoui N, Dianzinga N, Rébert E, Gauche MM, Pecrix Y, Costet L, Rieux A, Prior P, Robène I, Cellier G, Guérin F, Poussier S. 2020. Contrasting genetic diversity and structure among Malagasy *Ralstonia pseudosolanacearum* phylotype I populations inferred from an optimized Multilocus Variable Number of Tandem Repeat Analysis scheme. PLoS One 15:1–22.

95. Rasoamanana H, Ravelomanantsoa S, Nomenjanahary M, Gauche M, Prior P, Guérin F, Robène I, Pecrix Y, Poussier S. 2023. Bacteriocin production correlates with epidemiological prevalence of Phylotype I Sequevar 18 *Ralstonia pseudosolanacearum* in Madagascar. Appl Environ Microbiol 89.

96. Gonçalves OS, Souza F de O, Bruckner FP, Santana MF, Alfenas-Zerbini P. 2021. Widespread distribution of prophages signaling the potential for adaptability and pathogenicity evolution of *Ralstonia solanacearum* species complex. Genomics 113:992–1000.

97. Bertolla F, Frostegård Å, Brito B, Nesme X, Simonet P. 1999. During infection of its host, the plant pathogen *Ralstonia solanacearum* naturally develops a state of competence and exchanges genetic material. Mol Plant Pathol 12:467–472.

98. Greenrod STE, Stoycheva M, Elphinstone J, Friman VP. 2022. Global diversity and distribution of prophages are lineage-specific within the *Ralstonia solanacearum* species complex. BMC Genomics 23:1–19.

99. Wang X, Wei Z, Yang K, Wang J, Jousset A, Xu Y, Shen Q, Friman VP. 2019. Phage combination therapies for bacterial wilt disease in tomato. Nat Biotechnol 37:1513–1520.

100. Trotereau A, Boyer C, Bornard I, Pécheur MJB, Schouler C, Torres-Barceló C. 2021. High genomic diversity of novel phages infecting the plant pathogen *Ralstonia solanacearum*, isolated in Mauritius and Reunion islands. Sci Rep 11:1–10.

101. Ramírez M, Neuman BW, Ramírez CA. 2020. Bacteriophages as promising agents for the biological control of Moko disease (Ralstonia solanacearum) of banana. Biol Control 149:1–9.

102. Fujiwara A, Fujisawa M, Hamasaki R, Kawasaki T, Fujie M, Yamada T. 2011. Biocontrol of *Ralstonia solanacearum* by treatment with lytic bacteriophages. Appl Environ Microbiol 77:4155–4162.

103. Álvarez B, López MM, Biosca EG. 2019. Biocontrol of the major pathogen Ralstonia solanacearum in irrigation water and host plants by novel waterborne lytic bacteriophages. Front Microbiol 10:1–17.

104. Bhunchoth A, Phironrit N, Leksomboon C, Chatchawankanphanich O, Kotera S, Narulita E, Kawasaki T, Fujie M, Yamada T. 2015. Isolation of *Ralstonia solanacearum*-infecting bacteriophages from tomato fields in Chiang Mai, Thailand, and their experimental use as biocontrol agents. J Appl Microbiol 118:1023–1033.

105. Thapa Magar R, Lee SY, Kim HJ, Lee S-W. 2022. Biocontrol of bacterial wilt in tomato with a cocktail

of lytic bacteriophages. Appl Microbiol Biotechnol 106:3837–3848.

106.   Hida A, Oku S, Kawasaki T, Nakashimada Y, Tajima T, Kato J. 2015. Identification of the *mcpA* and *mcpM* genes, encoding methyl-accepting proteins involved in amino acid and L-malate chemotaxis, and involvement of McpM-mediated chemotaxis in plant infection by *Ralstonia pseudosolanacearum* (Formerly <i>. Appl Environ Microbiol 81:7420–7430.

107.   da Silva Xavier A, de Almeida JCF, de Melo AG, Rousseau GM, Tremblay DM, de Rezende RR, Moineau S, Alfenas-Zerbini P. 2019. Characterization of CRISPR-Cas systems in the *Ralstonia solanacearum* species complex. Mol Plant Pathol 20:223–239.

108.   Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia JM, Chia JM, Colasanti R, Conrad N, Davis JJ, Davison BH, Dejongh M, Devoid S, Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, Joachimiak MP, Keegan KP, Kondo S, Kumar V, Land ML, Meyer F, Mills M, Novichkov PS, Oh T, Olsen GJ, Olson R, Parrello B, Pasternak S, Pearson E, Poon SS, Price GA, Ramakrishnan S, Ranjan P, Ronald PC, Schatz MC, Seaver SMD, Shukla M, Sutormin RA, Syed MH, Thomason J, Tintle NL, Wang D, Xia F, Yoo H, Yoo S, Yu D. 2018. KBase: The United States department of energy systems biology knowledgebase. Nat Biotechnol 36:566–569.

109.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Res 49:W293--W296.

110.   Bernal P, Llamas MA, Filloux A. 2018. Type VI secretion systems in plant-associated bacteria. Environ Microbiol 20:1–15.

111.   Hulsen, T. 2021. BioVenn–an R and Python package for the comparison and visualization of biological lists using area-proportional Venn diagrams. Data Science, 4(1):51–61.

Figures and Tables:

Figure 1. RSSC genomes are enriched in *vgrG*-linked effector/immunity clusters.
(**A**) The number of *vgrG* alleles and T6SS core components (TssABCEFGHJKLM and Hcp) were compared across species within the Burkholderiaceae family. Using a custom genome database in KBase, we identified genes with BLASTp searches. The Grey "X" symbols indicate the nine genera with few-to-no T6SS genes (*Acidovorax, Alcaligenes, Bordetella, Comamonas, Hydrogenophaga, Pandoraea, Polaromonas, Polynucleobacter, Rhodoferax*). Error bars indicate SD, and dashed lines indicate the 95% confidence bands around the linear regression. (**B**) The number of *vgrG* genes were estimated in RSSC genomes (n=99) relative to closely related species/genera: other *Ralstonia* spp. (n=70), *Cupriavidus* spp. (n=120), and *Pandoraea* spp. (n=75). *vgrG* alleles were identified via BLASTp on KBase. Letters indicate p<0.0001 by Kruskal-Wallis multiple comparisons test.

A. General organization of the RSSC T6SS_main locus

B. Synteny of T6SS_main

C. Presence/Absence analysis of conserved T6SS_main genes

Figure 2. Although the T6SS is broadly conserved, multiple RSSC lineages lack the T6SS.
Most RSSC encode a type 6 secretion system (T6SS) with a conserved gene order with three variable regions and two conserved regions. (**A**) shows the conserved genes and (B) displays the synteny of the locus across phylogenetically diverse RSSC strains. Core T6SS genes encoding structural components and associated genes are shown as grey arrows, VgrG spike proteins are shown as purple arrows, and IS elements and transposases as orange arrows. Numbers superimposed over genes in the variable regions identify which auxiliary effector/immunity clusters are encoded at each spot. Linkages are drawn between homologous genes. Synteny and global amino acid identity was visualized with Clinker and aesthetics were adjusted in Affinity Designer. (**C**) Left: A phylogenetic tree of 398 RSSC genomes was created with the KBase "Species Tree" app, which compares 49 conserved bacterial genes. Right: The presence/absence of core T6SS genes based on BLASTp analysis. Grey rectangles indicate at least one homolog was identified using the KBase BLASTp v2.13.0+ app with percent identity cut-offs of > 15% and aligned-length cut-offs of > 20%. The data was visualized on iTOL. Lineages that lack the T6SS are indicated.

Figure 3. RSSC T6SS effectors are predicted to target lipids, nucleic acids, ATP, actin, and unknown targets.

To identify effectors and analyze their sequence for enzymatic domains, all Vgr-linked sequences were queried against NCBI Conserved Domain Database and PaperBlast. (A) Phylogenetic analysis of the effector profiles. (B-E) Comparison of clusters with (B) unknown effectors, (C) Lipases, (D) Nucleases, and (E) *aux4*, a polymorphic effector cluster that varies in the effector target.

Fig 4. RSSC pathogens vary in their repertoires of T6SS auxiliary effector/immunity clusters ("*aux*").

(**A**) Synteny analysis classified 1069 *vgr*-linked clusters into 25 subtypes (*aux1-aux25*) that varied in their prevalence from common to rare. Clusters were classified as "decayed" (pink) if genes were pseudogenized or were disrupted by transposons and were classified as "intact" (purple) if they lacked obvious mutations. (**B**) Relative abundance of intact *aux* clusters across major phylogenetic divisions of the RSSC. Lines indicate the median, and a cladogram shows the relative evolutionary relationships of the groups. Letters indicate significance groups by Kruskal-Wallis test with multiple comparisons (p<0.05). (**C**) Presence of each cluster (*aux1-aux25*) was determined across complete or nearly complete RSSC genomes by a combination of BLASTp and Clinker analyses. The left tree represents phylogeny of the *Ralstonia* genomes based on analysis of 49 genes. The top tree represents Bray-Curtis dissimilarity of the 25 *aux* clusters across the analyzed genomes. The phylogenetic tree was built with the Kbase SpeciesTree app, Bray-Curtis dissimilarity indices were calculated with the 'vegan v.2.6-2' package in RStudio (2023.03.0+386), and the visualization was created with iToL.

Figure 5. T6SS *aux* clusters are enriched on the megaplasmid.
(**A**) The number of *aux* clusters on each replicon (the chromosome, megaplasmid, and accessory conjugative plasmids) per RSSC genome. Letters indicate p<0.0001 by Brown-Forsythe and Welch ANOVA with Dunnett's T3 multiple comparisons test. (**B**) The occurrence of each *aux* cluster type on the replicons was identified. *aux* clusters are sorted by the substrate that the effector degrades (See Fig 3-4). Nine *aux* clusters were always (dark grey circles) or sometimes (white circles) located in the T6SS main locus. The three locations where *aux* clusters are found in the T6SS main locus are indicated above the graph. (**C**) The locations of *aux* clusters on the chromosome (green arrowheads) and megaplasmid (purple arrowheads) of RSSC species representatives were visualized. The species representatives (*R. pseudosolanacearum* GMI1000, *R. solanacearum* IBSBF1503, and *R. syzygii* PSI07) lack accessory conjugative plasmids.

Figure 6. Mobile genetic elements mediate horizontal gene transfer of *aux* clusters.
(**A**) Synteny analysis of representative *aux* loci associated with MGE. MGE genes are orange, *aux* cluster genes are brown, and all other genes are grey. (**B**) Proportion of of *aux* loci on the chromosomes, megaplasmids, and accessory plasmids that are MGE-associated, inconclusive, or not MGE-associated. (**C**) Specific locations of each *aux*-loci across the RSSC genomes. Each triangle represents one *aux* locus, and colors correspond to the legend in B. The letters identify hotspots where multiple *aux*-loci were found across the RSSC pangenome.