Cross-tolerance evolution is driven by selection on heat tolerance in

Drosophila subobscura

Luis E. Castañeda

Programa de Genética Humana, Instituto de Ciencias Biomédicas (ICBM), Facultad de

Medicina, Universidad de Chile, Santiago, Chile

Corresponding author: Luis E. Castañeda (luis.castaneda@uchile.cl)

ABSTRACT

The evolution of heat tolerance is a crucial mechanism for the adaptive response to

global warming, but it depends on the genetic variance carried by populations and on

the intensity of thermal stress in nature. Experimental selection studies have greatly

benefited research into heat tolerance, providing valuable insights into its evolutionary

process. However, the impact of varying levels of heat stress intensity on the associated

changes in resistance traits has not yet been explored. Here, the correlated evolution of

increasing knockdown temperature in *Drosophila subobscura* was evaluated on the

knockdown time at different stress temperatures (35, 36, 37, and 38 ºC), thermal death

time (TDT) curves, and desiccation and starvation resistance. The selection of heat

tolerance was performed using different ramping temperatures to compare the impact of

heat intensity selection on resistance traits. Correlated evolution was found for these

four resistance traits in *D. subobscura*, indicating that the evolutionary response to

tolerance of higher temperatures also confers the ability to tolerate other stresses such as

desiccation and starvation. However, these correlated responses depended on the

intensity of thermal selection and sex, which may limit our ability to generalize these

results to natural scenarios. Nevertheless, this study confirms the value of the

experimental evolutionary approach for exploring and understanding the adaptive

responses of natural populations to global warming.

Keywords: correlated evolution, global warming, heat stress intensity, stress resistance

evolution, thermal tolerance landscape.

INTRODUCTION

Rising environmental temperatures are a major challenge for ectotherms (i.e., organisms

whose body temperature depends on the ambient temperature) because their

morphology, physiology, behavior, and performance depend on the thermal

environment (Huey and Stevenson 1979; Cossins and Bowler 1987; Angilletta 2009).

Furthermore, rising environmental temperatures increase the risk of extinction for many

species living near their upper thermal limits (Deutsch et al. 2008; Huey et al. 2009;

Hoffmann and Sgrò 2011). However, ectotherms can avoid the negative effects of heat

through behavioral thermoregulation, evolutionary change, and/or phenotypic plasticity

of the upper thermal limits (Visser 2008).

Evolutionary adaptation depends on the genetic variation exhibited by upper

thermal limits; however, some studies have suggested that heat tolerance has a limited

evolutionary potential to respond to increasing environmental temperatures (Chown et

al. 2009; Mitchell and Hoffmann 2010; Kellermann et al. 2012). Yet, theoretical and

empirical evidence suggests that heritability estimates for heat tolerance tend to be

lower when heat tolerance is measured in longer assays (e.g., slow-ramping assays or

static assays using sublethal temperatures) than in shorter assays (e.g., fast-ramping

assays or static assays using extremely high temperatures) (Chown et al. 2009; Mitchell

and Hoffmann 2010; Rezende et al. 2011; Blackburn et al. 2014; Heerwaarden et al.

2016; Castañeda et al. 2019). Thus, the intensity of heat stress may influence our

predictions regarding the evolutionary potential of heat tolerance, but how do

populations respond to variable selection driven by heat stress? Selection under

laboratory conditions has a long history of providing information on the adaptive

evolution of specific selective agents (Lenski and Bennett 1993; Garland Jr 2003; Fuller

et al. 2005; Gibbs and Gefen 2009). In particular, the experimental evolution of heat

tolerance has been assessed in several species, including fish, corals, and insects (Baer

and Travis 2000; Kelly et al. 2012; Geerts et al. 2015; Esperk et al. 2016). Experimental

evolution of heat tolerance has also been studied in several *Drosophila* species,

including D. melanogaster (Gilchrist and Huey 1999; Folk et al. 2006), *D. subobscura*

(Quintana and Prevosti 1990; Mesas et al. 2021; Mesas and Castañeda 2023), and *D.*

buzzatti (Krebs and Loeschcke 1996). Most of these studies reported the evolution of

heat tolerance using fast ramping protocols, ranging from 0.4 ºC/min in Folk et al. (Folk

et al. 2006) to 1 °C/min in Gilchrist and Huey (Gilchrist and Huey 1999) or static high-

temperature assays (40 °C), as in Bubliy and Loeschcke (2005). Recently, Mesas et al.

(Mesas et al. 2021) reported that selected lines of *D. subobscura* evolved higher heat

tolerance, regardless of the heating rate used during the selection experiments (slow-

ramping rate: 0.08 °C/min and fast-ramping rate: 0.4 °C/min).

Interestingly, several of these selection experiments on heat tolerance in

Drosophila have found correlated responses in other traits such as starvation resistance,

desiccation resistance, and heat shock proteins (Hoffmann et al. 1997; Feder et al. 2002;

Bubliy and Loeschcke 2005). However, the intensity of thermal stress is expected to

have important effects on the correlated responses of other traits to heat tolerance

selection (Fragata and Simões 2022). For example, fast-ramping selected lines have

evolved thermal performance curves with higher optimum temperatures and narrower

thermal breadths than slow-ramping selected lines (Mesas et al. 2021). In addition,

Mesas and Castañeda (Mesas and Castañeda 2023) reported that the evolution of heat

tolerance was associated with reduced activity of the enzymes involved in the glucose-

6-phosphate branch point and increased performance of life-history traits in slow-

ramping selected lines. However, they did not observe any changes in the metabolic rate

of the selected lines, as predicted by Santos et al. (2012). In summary, there is evidence

that heat stress intensity determines the magnitude of the evolutionary responses of

performance, metabolic, and life-history traits to heat tolerance selection; however, the

correlated evolution of resistance traits has not yet been tested. This information should

explain how thermal stress intensity might determine the cross-tolerance evolution to

stressful environmental conditions. Natural populations are regularly subjected to

multiple environmental stressors, and it is well-established that enhanced tolerance to

one stressor can enhance tolerance to another (Rodgers and Gomez Isaza 2023). Cross-

tolerance induced by thermal stress has been widely studied in several arthropod

species, increasing resistance to desiccation, insecticides, and pathogens (Kalra et al.

2017; Rodgers and Gomez Isaza 2021; Singh et al. 2022). However, the cross-tolerance

patterns at the evolutionary level can be constrained or facilitated by genetic

correlations among resistance traits depending on the environmental context (Lande and

Arnold 1983; Bubliy and Loeschcke 2005; Gerken et al. 2016).

97   Previous research has examined the impact of varying levels of heat stress on the

heat knockdown temperature of *D. subobscura*, as well as its associated impacts on

thermal performance curves (Mesas et al. 2021), energy metabolism, and fitness-related

traits (Mesas and Castañeda 2023). The evolutionary response of these traits was

evaluated using two thermal selection protocols that differed in the rate of temperature

increase (hereafter, ramping rate) to measure the heat knockdown temperature: slow-

ramping selection (0.08ºC min^{-1}) and fast-ramping selection (0.4ºC min^{-1}). The present

study investigates the effects of heat intensity selection for increasing knockdown

temperature on the cross-tolerance evolution of four different resistance traits in *D.*

subobscura: knockdown time at different stress temperatures, thermal-death-time curves

(TDT), desiccation resistance, and starvation resistance. In particular, TDT curves

represent an integrative approach to assess how the probability of survival depends on

the intensity and duration of heat stress, as they allow the estimation of the critical

thermal maxima (CT_{max}) and thermal sensitivity using the thermal tolerance

measurements obtained at different stress temperatures (Rezende et al. 2014). Here, it is

expected that fast-ramping selected lines will exhibit higher knockdown time at highly

stressful temperatures and higher CT_{max} because fast-ramping protocols reduce the

confounding effects (e.g., hardening, rate of resource use) on heat tolerance associated

with the assay length (see Rezende et al. 2011; Santos et al. 2012; Mesas et al. 2021). In

contrast, slow-ramping selected lines should exhibit higher desiccation and starvation

resistance because individuals with higher starvation and desiccation resistance exhibit

higher thermal tolerance during long assays.

Materials and Methods

Sampling and maintenance

D. subobscura females were collected in the spring 2014 at the Botanical Garden of

the Universidad Austral de Chile (Valdivia, Chile; 39º 48' S, 73º 14' W) using plastic

traps containing banana/yeast baits. Two hundred females were collected and placed

individually in plastic vials containing David's killed-yeast *Drosophila* medium to

establish isofemale lines. In the next generation, 100 isofemale lines were randomly

selected, and 10 females and 10 males per line were placed in an acrylic cage to

establish a large, outbred population. In the next generation, the flies from this cage

were divided into three population cages (R1, R2, and R3), attempting to assign the

same number of flies to each cage. After three generations, the flies in each replicate

cage were divided into four population cages, trying to assign the same number of flies

to each cage. This procedure established 12 population cages assigned to each

artificial selection protocol in triplicate: fast-ramping selection, fast-ramping control,

slow-ramping selection, and slow-ramping control lines (Fig. S1). During the selection

experiments, population cages were maintained at 18 °C (12:12 light-dark cycle) in a

discrete generation, controlled larval density regime (Castañeda et al. 2015). Each

population cage had a population size of 1000-1500 breeding adults.

Heat tolerance selection

For each replicate line, 120 four-day-old virgin females were randomly mated with

two males for two days, after which the females were individually placed in a capped

5-mL glass vial, and the males were discarded. The vials were attached to a plastic

rack and immersed in a water tank with an initial temperature of 28 °C, controlled by a

heating unit (model ED, Julabo Labortechnik, Seelbach, Germany). After an

equilibration period of 10 min, the temperature was increased to 0.08 ºC min^{-1} for the

slow-ramping selection protocol or 0.4 ºC min^{-1} for the fast-ramping selection

protocol. Assays were stopped when all flies collapsed. Each assay was recorded using

a high-resolution camera (model D5100, Nikon, Tokyo, Japan) and then visualized to

score the knockdown temperature for each fly, defined as the temperature at which

each fly ceased to move. Flies were ranked by knockdown temperature, and four

151 virgin females were selected from the progeny of the 40 flies with the highest

152 knockdown temperature (top 30% of each assay) to establish the next generation. For

153 the fast and slow control lines, the knockdown temperature was measured as described

154 above, but the progeny was randomly selected to establish the next generation,

155 regardless of the knockdown temperature of their mother.

156  This artificial selection experiment was performed for 16 generations, after which

157 flies from each selection treatment were placed in separate acrylic cages and

158 maintained without selection (e.g., relaxed selection) at 18 °C and a 12:12 light-dark

159 cycle.

161 ***Knockdown time in static assays***

162 Eggs were collected from each population cage and transferred to vials at a density of

163 40 eggs/vial. At 4 days of age, ten females and ten males from each population cage

164 were tested to measure their heat knockdown time at four different static temperatures:

165 35, 36, 37, and 38°C. This experimental design allowed the measurement of 960 flies

166 (10 flies × 2 sexes × 4 static temperatures × 4 selection treatments × 3 replicated lines).

167 Static assays were performed similarly to knockdown temperature assays, but static

168 temperatures were used instead of ramping temperatures. A total of 240 flies were

169 measured for each static temperature, except for the assay at 35°C (178 flies) because

170 two flies died before the start of the assay, and a video file of one assay was corrupted

171 (data for 60 flies were lost). For the 37°C assay, four flies died before the assay began,

172 and the collapse time could not be measured for six flies. Finally, for the 38°C assay,

173 three flies died before the start of the assay and the collapse time could not be measured

174 for five flies. Heat knockdown assays were performed in generation 23 (Fig. S1).

Desiccation and starvation resistance

Eggs from each replicate cage were collected and maintained in vials at a density of 40

eggs/vial. Only fast control lines were measured as control lines. This decision was

based on logistical reasons (i.e., the high number of vials) and statistical support

because fast and slow control lines did not differ in their knockdown times and CT_{max}

values (see *the Results* section).

For desiccation resistance assays, five flies from each sex were separately placed

in a vial containing five desiccant droplets (Drierite) and sealed with parafilm (flies had

no access to food or water during the assay). For starvation resistance assays, five flies

from each sex were separately placed in a vial containing agar only (flies had access to

water but no food). For both desiccation and starvation resistance assays, the number of

live flies was counted every 3 h until all the flies were dead. Desiccation and starvation

resistance were measured in 126 vials containing 10 flies each, respectively (7 vials × 2

sexes × 3 selection treatments × 3 replicate lines). These experiments were conducted at

18 °C using flies from generation 24 (Fig. S1).

Statistical analysis

Normality and homoscedasticity were tested for all variables, and the knockdown times

were squared root transformed to meet the parametric assumptions. All analyses were

performed with R software (R Development Core Team 2011).

Heat tolerance

For the knockdown temperature, control and selection lines were compared separately

for the fast- and slow-ramping selection because it is well known that the knockdown

temperature is higher in fast-ramping than in slow-ramping assays (Chown et al. 2009;

see Mesas et al. 2021). For the knockdown time analysis, a mixed linear model with

ramping selection (fixed effect with fast-control, slow-control, fast-selection, and slow-

selection lines as levels), sex (fixed effect with females and males as levels), and

replicate lines nested within the thermal selection (random effect with replicates 1, 2

and 3 as levels) was performed using the library *lme4* package for R (Bates et al. 2015).

Fixed effects were tested by a type III ANOVA and the random effect was tested by a

likelihood ratio test comparing the model with and without the replicate lines. Both tests

were performed using the library *lmerTest* package for R (Kuznetsova et al. 2017). If

the selection effect was significant, *a posteriori* comparisons were performed using

false discovery rate adjustment implemented in the *emmeans* package for R (Lenth et al.

2018).

Knockdown times were also used to plot the survival curves based on the

Kaplan-Meier formula using the *survfit* function implemented in the *survival* package

for R (Therneau 2023).

Thermal death time curves (TDT)

Average knockdown times were calculated for each sex, replicate lines, and selection

treatment combination (Table S1). These values were regressed against the assay

temperatures according to Equation 1 (Rezende et al. 2014):

$$log_{10}t = \frac{CT_{max}-T}{z} \qquad \text{eqn. 1}$$

, where T is the assay static temperature (ºC), CT_{max} is the upper thermal limit (ºC), t is

the knockdown time (min), and z is the thermal sensitivity. These curves allowed the

estimation of CT_{max} as the extrapolated temperature that would result in a knockdown

time of $\log_{10} t = 0$ (i.e., knockdown time at 1 min) and the estimation of the thermal

sensitivity ($z = -1/$slope), where the lower z values, the higher the thermal sensitivity.

228   Using equation 1, 24 TDT curves (2 sexes × 3 replicate lines × 4 selection

protocols) were fitted, from which CT_{max} and z values were estimated as described

above. A linear model with ramping selection treatment (levels: fast-control, slow-

control, fast-selection, and slow-selection lines), sex (levels: females and males), and

their interaction was performed to evaluate their effects on CT_{max} and z values. TDT

curve analysis did not include replicate lines as a random effect because only one CT_{max}

and z value was estimated by each replicate line. Additionally, a linear mixed model

with ramping selection (fixed effect with fast-control, slow-control, fast-selection, and

slow-selection lines as levels), sex (fixed effect with females and males as levels), and

replicate lines nested within the thermal selection (random effect with replicates 1, 2

and 3 as levels), and assay temperatures (as covariate) was fitted on the knockdown

time using the *lmer* package for R.

Desiccation and starvation resistance

To determine the lethal time at which 50% of flies of each vial were dead (LT_{50}), a

generalized linear model following a binomial distribution was fitted with the

proportion of flies alive as the dependent variable and time as the predictor variable.

The generalized linear model was run using the g*lm* function of the *lme4* package for R

(Bates et al. 2015). The LT_{50} of each vial was then estimated using the function *dose.p*

from the *MASS* package for R (Venables and Ripley 2002).

248   To estimate the median LT_{50} and the 95% confidence intervals for each selection

treatment and sex, each LT_{50} was transformed into a survival object using the *Surv* and

survfit functions of the *survival* package for R (Therneau 2023). This procedure also

allowed to estimate the survival curves in each vial. Finally, to test the effect of

selection treatment (levels: control, fast-selection, and slow-selection lines) and sex

(levels: females and males) on desiccation and starvation resistance, a Cox proportional

regression model was fitted with LT_{50} as the dependent variable, and selection protocol

and sex as predictor variables. The Cox model was run using the *coxph* function of the

survival package (Therneau 2023).

RESULTS

Knockdown temperature evolution

Knockdown temperature evolved in response to artificial selection for increased heat

tolerance, regardless of the ramping assay protocol: the knockdown temperature was

significantly higher in fast-ramping selected lines than in fast-ramping control lines

(mean fast-ramping selected lines [95% CI] = 37.71 ºC [37.63 – 37.78] and mean fast-

ramping control lines [95% CI] = 37.23 ºC [37.0 – 37.38]; $F_{1,4}$ = 32.0, P = 0.005); and

the knockdown temperature in slow-ramping selected lines was significantly higher

than in slow-ramping control lines (mean slow-ramping selected lines [95% CI] =

35.48ºC [35.41 – 35.55] and mean fast-ramping control lines [95% CI] = 34.97 ºC

[34.82 – 35.12]; $F_{1,4}$ = 41.7, P = 0.003). These results were previously reported by

Mesas et al. (2021) and are reported here to show that selected lines used in this study

evolved higher thermal tolerance compared to control lines.

Knockdown time evolution

As expected, the knockdown time decreased significantly as the assay temperatures

increased ($F_{1,877}$ = 649.1, $P < 2\times10^{-16}$). The mean knockdown time and 95% CI for each

static assay are as follows: 35º C = 33.77 min [32.1 – 35.5]; 36º C = 16.98 min [16.1

– 17.9]; 37º C = 8.84 min [8.4 – 9.3]; and 38º C = 6.78 min [6.3 – 7.0].

Knockdown times differed significantly between selection treatments when flies

were assayed at 36 and 37ºC (Table 1; Table S2; Fig. 1). At these temperatures, slow

and fast selected lines showed higher heat tolerance than slow and fast control lines

(Table S1- S3; Fig. 1C, E). Also, fast-selected lines showed a higher heat tolerance than

slow-selected lines in flies assayed at 37ºC but not at 36ºC (Table S1-S3; Fig. 1C, E),

whereas fast and slow control lines did not differ (Table S3; Fig 1). On the other hand,

replicate lines had no significant effect on knockdown time, indicating consistent

evolutionary responses within each selection and control treatment (Table S2).

Concerning sex, females showed a higher thermal tolerance than males but only when

flies were assayed at 35 and 38ºC (Table 1; Fig. 1B, H). Finally, non-significant

interactions between selection and sex were found for all assay temperatures (Table 1).

TDT curves evolution

Linear regressions between $\log_{10}(LT_{50})$ and assay temperatures enabled the estimation

of 24 TDT curves (4 selection treatments × 3 replicate lines × 2 sexes) with high

coefficients of determination (mean R^2 = 0.946, range: 0.820 – 0.989; Table S4),

confirming that heat knockdown time is linearly related to stressful sublethal

temperatures. From these TDT curves, the mean CT_{max} [95% CI] was 41.21ºC [41.02

– 41.41], and the mean z [95% CI] was 4.18ºC [4.03 – 4.32]. CT_{max} were significantly

different between selection treatments ($F_{3,20}$ = 4.46, P = 0.015; Fig. 2A). A post hoc

analysis showed that fast-ramping selected and slow-ramping control lines were

significantly different in their CT_{max} values (t_{20} = 3.195, P = 0.02). In contrast, fast and

slow control lines had similar CT_{max} values (t_{20} = 0.911, P = 0.80). Thus, when control

lines are pooled, CT_{max} still differs between selection treatments ($F_{2,18} = 6.69$, $P =$

0.007), with fast-ramping (mean CT_{max} [95% CI] = 41.55 ºC [41.2 – 41.9]) and slow-

ramping selected lines (mean CT_{max} [95% CI] = 41.43 ºC [41.1 – 41.8]) had higher

CT_{max} than control lines (mean CT_{max} [95% CI] = 40.94 ºC [40.7 – 41.2]) ($t_{18} = 3.27$, P

$= 0.01$ and $t_{18} = 2.64$, $P = 0.04$, respectively). CT_{max} was not different between the

selected lines ($t_{18} = 0.54$, $P = 0.85$). On the other hand, sex and the interaction between

selection treatments and sex had no significant effect on CT_{max} ($F_{1,18} = 0.004$, $P = 0.95$

and $F_{3,18} = 2.11$, $P = 0.15$, respectively). Regarding z (i.e., thermal sensitivity), it shows

no significant effects of selection treatments ($F_{3,16} = 0.91$, $P = 0.46$; Fig. 2), sex ($F_{1,16} =$

1.30, $P = 0.27$), nor the interaction between selection treatments and sex ($F_{3,16} = 2.23$, P

$= 0.12$). In summary, the evolution of a higher CT_{max} is not associated with an

evolutionary change in thermal sensitivity (Fig. 2B). Indeed, the relationship between

CT_{max} and z did not change with the evolution of increasing thermal tolerance ($r_{control-lines}$

$= 0.979$ and $r_{selected-lines} = 0.929$; Z-test = 0.76, $P = 0.45$). Additionally, using a linear

mixed model with the assay temperature as a covariate, this result was corroborated by a

non-significant interaction between selection treatment and assay temperature ($F_{3,865} =$

0.30, $P = 0.82$).

Desiccation resistance evolution

Survival analysis showed a significant effect of sex and selection treatment on

desiccation resistance, but not for the interaction of the two effects (Table S5). Males

showed a higher risk of desiccation than female flies (hazard ratio = 7.11, $P < 2 \times 10^{-7}$;

Fig. 3). Females showed a significant difference between selected and control lines

(LTR: $\chi^2_2 = 6.72$, $P = 0.03$; Fig. 3A). Specifically, females of the slow-ramping

selection lines showed a higher desiccation resistance than females of the control lines

(hazard ratio = 0.42, P = 0.009), whereas females of the fast-ramping selection and

control lines showed similar desiccation risk (hazard ratio = 0.56, P = 0.072). On the

other hand, males showed no differences in desiccation resistance between selected and

control lines (LTR: χ^2_2 = 1.88, P = 0.4; Fig. 3B).

Starvation resistance evolution

A significant effect of sex, selection treatment, and the interaction between the two

effects on starvation resistance was found in the survival analysis (Table S6). Males had

a higher risk of starvation than female flies (hazard ratio = 22.75, $P < 1\times10^{-16}$; Fig. 4).

In female flies (Fig. 4A), fast-ramping selection and slow-ramping selection lines

showed a higher starvation risk than control lines (hazard ratio = 2.37, P = 0.009; and

hazard ratio = 2.20, P = 0.014, respectively). In contrast, male flies had an opposite

pattern (Fig. 4B): slow-ramping selection lines had a lower starvation risk than control

lines (hazard ratio = 0.50, P = 0.03), but nonsignificant differences were found between

fast-ramping selection and control lines (hazard ratio = 0.64, P = 0.16).

Discussion

Studying the evolutionary responses of thermal limits is key to understanding the

adaptive responses and evolutionary constraints to global warming. Cross-tolerance

studies can provide valuable information on the evolutionary response to multiple

environmental stressors. Cross-tolerance evolution has been reported among different

resistance traits (Hoffmann and Parsons 1993; Bubliy and Loeschcke 2005; Stazione et

al. 2020; Singh et al. 2022), but the magnitude of the evolutionary response could be

explained by the trait under direct selection or the stress intensity (Gerken et al. 2016).

Here, artificial selection for heat tolerance (i.e., knockdown temperature) resulted in

correlated responses in heat knockdown time, the thermal tolerance landscape (TDT

curves), desiccation resistance, and starvation resistance. However, these responses

depended on the intensity of thermal selection and sex, suggesting that the evolutionary

response to tolerate higher temperatures also confers partial tolerance to other stresses

such as desiccation and starvation.

Different approaches to measuring the upper thermal limit of ectotherms

produce different genetic and phenotypic estimates. Fast-ramping assays generally

estimate higher upper thermal limits and higher heritabilities than slow-ramping assays

(Chown et al. 2009; Rezende et al. 2011). For instance, the heritability of thermal

tolerance was 0.13 for fast assays and 0.08 for slow assays in *D. subobscura* (Castañeda

et al. 2019). Because heritability is commonly used as a predictor of the evolutionary

response of a trait to natural or artificial selection, the evolutionary response of heat

tolerance would be expected to depend on the ramping rate used during selection.

However, previous work did not support this prediction for *D. subobscura*, finding that

the evolution of heat tolerance was independent of the ramping rate (Mesas et al. 2021),

but the correlated responses of the thermal performance curves or the energy

metabolism depended on the intensity of the thermal selection (Mesas et al. 2021;

Mesas and Castañeda 2023). In the present study, the evolution of knockdown

temperature (e.g., heat tolerance measured in dynamic assays) induced a correlated

response on the heat knockdown time (e.g., heat tolerance measured in static assays)

when it was assayed at intermediate temperatures (36 and 37ºC), but not at less or more

extreme assay temperatures (35 and 38ºC). These findings can be explained because

stress tolerance at 35ºC should depend on the physiological state of the organism during

prolonged thermal assays (e.g., availability of energy resources; see Rezende et al.

2011, but also see Overgaard et al. 2012) and not only on heat tolerance, whereas heat

tolerance at 38ºC could be limited by physical properties of ectotherms (e.g., protein

denaturation, membrane permeability). However, a previous study found a clinal pattern

for heat tolerance in *D. subobscura* only for flies assayed in static assays (specifically at

38ºC), but this clinal pattern was not detected using ramping assays (Castañeda et al.

2015). Differences between these two studies could be explained by the number of

generations under thermal selection, which could result in a different evolutionary

response of heat tolerance. According to Begon (1976), *D. subobscura* can have

between 4 and 6 generations per year, which makes it possible to estimate about 125

generations of selection from the introduction of *D. subobcura* in Chile until the study

by Castañeda et al. (2015). On the other hand, the type of selection is completely

different between the two studies (e.g., natural versus artificial selection), which could

lead to various evolutionary outcomes. In any case, beyond these results from specific

thermal assays, these findings support the idea that (1) the use of a single static

temperature would miss genetic or phenotypic effects on heat tolerance, and (2)

unifying several knockdown time estimates into a single approach (TDT curves) should

be necessary to elucidate genetic and phenotypic patterns of heat tolerance in

ectotherms (Rezende et al. 2014; Jørgensen et al. 2021).

TDT curves evolved in response to heat tolerance selection in *D. subobscura*.

TDT curves showed that fast- and slow-ramping selected lines evolved higher CT_{max}

than control lines (ΔCT_{max} = 0.49 ºC). This differential CT_{max} value is slightly lower

than the population differences (0.9ºC) observed between the lowest and highest latitude

populations (~8 latitudinal degrees) of *D. subobscura* studied by Castañeda et al. (2015)

and even lower than the CT_{max} variation reported among *Drosophila* species (Jørgensen

et al. 2019; Alruiz et al. 2022). On the other hand, although CT_{max} and z (i.e., thermal

sensitivity) are phenotypically correlated (see Castañeda et al. 2015; Molina et al.

2023), the evolutionary increase in CT_{max} was not associated with a correlated response

in thermal sensitivity (z). This result suggests that both thermal parameters are not

genetically constrained, but further evidence from quantitative genetic studies is needed

to assess the genetic association between CT_{max} and z. A caveat for this finding could be

related to the fact that thermal selection for heat tolerance was carried out over 16

generations, followed by 7 generations of relaxed selection (i.e., no selection).

However, previous evidence suggests that differences in heat tolerance between control

and selected lines were consistent between generations 16 and 25 (Mesas et al. 2021).

Indeed, Passananti et al. (2004) also reported that phenotypic values did not change

after 35 generations of relaxed selection in desiccation-selected populations of *D.*

melanogaster.

It was expected that flies selected for higher heat tolerance using slow-ramping

rate protocols would exhibit greater desiccation and starvation resistance than flies

selected using fast-ramping selection protocols. This is because flies assayed for heat

tolerance in long assays are also exposed to desiccation and starvation stress (Santos et

al. 2012). This study provides partial support for this hypothesis. First, slow-ramping

selected lines evolved a higher desiccation resistance than control and fast-ramping

selected lines. However, this was only observed in female flies, while males of the

different selection treatments did not show any difference in desiccation resistance. On

the other hand, starvation resistance evolved in opposite directions depending on sex:

females of the fast-ramping and slow-ramping selected lines showed lower starvation

resistance than females of the control lines, whereas males of the slow-ramping selected

lines showed higher starvation resistance than males of the control and fast-ramping

selected lines. Differential evolutionary responses between the sexes could be due to

heat thermal selection only being applied to females, which could have exacerbated the

evolutionary responses of female flies. However, previous studies that artificially

selected exaggerated male traits also found fitness consequences in females (Harano et

al. 2010). Differential evolutionary responses between females and males can then be

explained by sexually antagonistic selection on genetically correlated traits (Eyer et al.

2019; Fanara et al. 2023). Kwan et al. (2008) reported that desiccation-selected females

had higher desiccation resistance than desiccation-selected males (see also Chippindale

et al. 2004), which can be explained by males using resources at a faster rate than

females (e.g., males lose weight, water, and metabolites faster than females). Sexual

dimorphism in stress resistance traits has been mainly explained by differences in

cuticular composition, resource storage, and energy conservation between the sexes

(Schwasinger-Schmidt et al. 2012; Rusuwa et al. 2022). Although energy content was

not measured here, Mesas and Castañeda (2023) found that body mass and metabolic

rate were similar between control and heat-tolerance selected lines of *D. subobscura*,

suggesting that neither resource storage nor energy conservation explains the sex-

dependent correlated response for stress resistance traits. However, the same study

found that heat-tolerance selected lines had higher fecundity than control lines, whereas

previous studies have found negative associations between fecundity and starvation

resistance in *D. melanogaster* (Bubliy and Loeschcke 2005; Kalra et al. 2017). Then,

the decrease in starvation resistance in females of the heat-selected lines could be

related to increased fecundity, which is consistent with the reported trade-off between

stress resistance traits and life-history traits (van Noordwijk and de Jong 1986; Rion and

Kawecki 2007).

In conclusion, the present study shows that heat tolerance evolution is associated

with evolutionary responses in other stress resistance traits, which could be explained

by pleiotropic effects or linkage disequilibrium among the traits evaluated. However,

further evidence (e.g., quantitative genetic or genome-wide analysis studies) is needed

to elucidate the genetic basis of the cross-tolerance evolution in *D. subobscura*. In

addition, this study provides evidence for rapid evolutionary responses in ectotherms

mediated by thermal selection, but the evolutionary outcomes depend on the intensity of

the thermal stress (Mesas and Castañeda 2023) and sex (Rogell et al. 2014; Rusuwa et

al. 2022). This study also highlights the importance of *D. subobscura* as a suitable

model to study thermal adaptation mediated by natural selection (Huey 2000; Gilchrist

et al. 2008; Castañeda et al. 2013, 2015), and laboratory selection (Santos et al. 2005,

Santos et al. 2021; Simões et al. 2017; Mesas et al. 2021; Mesas and Castañeda 2023).

In addition, this study highlights the relevance of experimental evolutionary studies for

understanding the adaptive responses to climate change (Mitchell and Whitney 2018;

Brennan et al. 2022; Kelly 2022). Finally, these results suggest that ectotherms may

evolve in response to climate warming, but evolutionary responses may differ between

sexes and/or the warming rates experienced by natural populations, which may make it

difficult to propose general trends in the fate of ectotherms in a changing world where

temperature is not the only driver of climate change, but species are also expected to be

exposed to changes in precipitation patterns and food availability.

Acknowledgments

I thank Andres Mesas, Angélica Jaramillo, Julio Figueroa, and Jaiber Solano for their

help with fly maintenance, experiments, and data management. I also thank Enrico

Rezende and Mauro Santos for their valuable suggestions and advice while designing

the artificial selection experiment. A preprint version of this article has been peer-

reviewed and recommended by PCIEvolBiol

(https://doi.org/10.24072/pci.evolbiol.100709).

Funding

This work was funded by the Fondo Nacional de Desarrollo Científico y Tecnológico

(FONDECYT) [grant number 1140066] – Chile. Currently, L.E. Castañeda is supported

by the Agencia Nacional de Invertigación y Desarrollo (ANID) through the grants

Anillo ATE230025, FOVI220194, and FOVI 230149.

Data availability

Data and scripts are available at https://doi.org/10.6084/m9.figshare.24085107.v5

Conflict of interest disclosure

The author declares having no financial conflicts in relation to the content of the article.

References

Alruiz, J. M., I. Peralta-Maraver, F. Bozinovic, M. Santos, and E. L. Rezende. 2022.
        Thermal tolerance in Drosophila: Repercussions for distribution, community
        coexistence and responses to climate change. J. Anim. Ecol. 91:655–667.

Angilletta, M. J. 2009. Thermal Adaptation: A Theoretical and Empirical Synthesis.
        Oxford University Press.

Baer, C. F., and J. Travis. 2000. Direct and correlated responses to artificial selection on
        acute thermal stress tolerance in a livebearing fish. Evolution (N. Y). 54:238–244.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects
        models using lme4. J. Stat. Softw. 67:1–48.

Begon, M. 1976. Temporal variations in the reproductive condition of Drosophila
        obscura Fallén and D. subobscura Collin. Oecologia 23:31–47.

Blackburn, S., B. Van Heerwaarden, V. Kellermann, and C. M. Sgrò. 2014.

Evolutionary capacity of upper thermal limits: beyond single trait assessments. J.

Exp. Biol. 217:1918–1924.

Brennan, R. S., J. A. deMayo, H. G. Dam, M. Finiguerra, H. Baumann, V. Buffalo, and

M. H. Pespeni. 2022. Experimental evolution reveals the synergistic genomic

mechanisms of adaptation to ocean warming and acidification in a marine copepod.

Proc. Natl. Acad. Sci. U. S. A. 119:1–10.

Bubliy, O. A., and V. Loeschcke. 2005. Correlated responses to selection for stress

resistance and longevity in a laboratory population of Drosophila melanogaster. J.

Evol. Biol. 18:789–803.

Castañeda, Luis (2023). Cross-tolerance evolution is driven by selection on heat

tolerance in Drosophila subobscura. figshare. Dataset.

https://doi.org/10.6084/m9.figshare.24085107.v5

Castañeda, L. E., J. Balanya, E. L. Rezende, and M. Santos. 2013. Vanishing

chromosomal inversion clines in Drosophila subobscura from Chile : is behavioral

thermoregulation to blame ? 182:249–259.

Castañeda, L. E., E. L. Rezende, and M. Santos. 2015. Heat tolerance in Drosophila

subobscura along a latitudinal gradient: Contrasting patterns between plastic and

genetic responses. Evolution (N. Y). 69:2721–2734.

Castañeda, L. E., V. Romero-Soriano, A. Mesas, D. A. Roff, and M. Santos. 2019.

Evolutionary potential of thermal preference and heat tolerance in Drosophila

subobscura. J. Evol. Biol. 32:818–824.

Chippindale, A., M. R. Rose, A. G. Gibbs, A. K. Chippindale, and M. R. Rose. 2004.

Methuselah Flies - A Case Study in the Evolution of Aging. , doi:

10.1142/9789812567222.

Chown, S. L., K. R. Jumbam, J. G. Sørensen, and J. S. Terblanche. 2009. Phenotypic
variance, plasticity and heritability estimates of critical thermal limits depend on
methodological context. Funct. Ecol. 23:133–140.

Cossins, A. R., and K. Bowler. 1987. Temperature Biology of Animals. Chapman and
Hall.

Deutsch, C. A., J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C.
Haak, and P. R. Martin. 2008. Impacts of climate warming on terrestrial
ectotherms across latitude. Proceeding Natl. Acad. Sci. 105:6668–6672.

Esperk, T., A. Kjærsgaard, R. J. Walters, D. Berger, and W. U. Blanckenhorn. 2016.
Plastic and evolutionary responses to heat stress in a temperate dung fly: Negative
correlation between basal and induced heat tolerance? J. Evol. Biol. 29:900–915.

Eyer, P. A., A. J. Blumenfeld, and E. L. Vargo. 2019. Sexually antagonistic selection
promotes genetic divergence between males and females in an ant. Proc. Natl.
Acad. Sci. U. S. A. 116:24157–24163.

Fanara, J. J., P. L. Sassi, J. Goenaga, and E. Hasson. 2023. Genetic basis and
repeatability for desiccation resistance in Drosophila melanogaster (Diptera:
Drosophilidae). Genetica, doi: 10.1007/s10709-023-00201-0. Springer
International Publishing.

Feder, M. E., T. B. C. Bedford, D. R. Albright, and M. Pawel. 2002. Evolvability of
Hsp70 expression under artificial selection for inducible thermotolerance in
independent populations of Drosophila melanogaster. Physiol. Biochem. Zool.
75:325–334.

Folk, D. G., P. Zwollo, D. M. Rand, and G. W. Gilchrist. 2006. Selection on
knockdown performance in Drosophila melanogaster impacts thermotolerance and
heat-shock response differently in females and males. J. Exp. Biol. 209:3964–

3973.

Fragata, I., and P. Simões. 2022. The other side of the evolution of heat tolerance:

correlated responses in metabolism and life-history traits. Peer Community Evol.

Biol. 100155.

Fuller, R. C., C. F. Baer, and J. Travis. 2005. How and when selection experiments

might actually be useful. Integr. Comp. Biol. 45:391–404.

Garland Jr, T. 2003. Selection experiments : an under-utilized tool in biomechanics and

organismal biology. Pp. 23–57 *in* V. L. Bels, J.-P. Gasc, and A. Casinos, eds.

Vertebrate Biomechanics and Evolution. BIOS Scientific Publisher Ltd, Oxford.

Geerts, A. N., J. Vanoverbeke, B. Vanschoenwinkel, W. Van Doorslaer, H. Feuchtmayr,

D. Atkinson, B. Moss, T. A. Davidson, C. D. Sayer, and L. De Meester. 2015.

Rapid evolution of thermal tolerance in the water flea Daphnia. Nat. Clim. Chang.

5:665–668.

Gerken, A. R., T. F. C. Mackay, and T. J. Morgan. 2016. Artificial selection on chill-

coma recovery time in Drosophila melanogaster: Direct and correlated responses to

selection. J. Therm. Biol. 59:77–85. Elsevier.

Gibbs, A. G., and E. Gefen. 2009. Physiological Adaptations in Laboratory

Environments. Pp. 523–550 *in* T. Garland Jr and M. R. Rose, eds. Experimental

Evolution: Concepts, Methods, and Applications of Selection Experiments.

University of California Press, Berkeley.

Gilchrist, G. W., and R. B. Huey. 1999. The direct response of Drosophila melanogaster

to selection on knockdown temperature. Heredity (Edinb). 83 (Pt 1):15–29.

Gilchrist, G. W., L. M. Jeffers, B. West, D. G. Folk, J. Suess, and R. B. Huey. 2008.

Clinal patterns of desiccation and starvation resistance in ancestral and invading

populations of   Drosophila subobscura. Evol. Appl. 1:513–523.

Harano, T., K. Okada, S. Nakayama, T. Miyatake, and D. J. Hosken. 2010. Intralocus

sexual conflict unresolved by sex-limited trait expression. Curr. Biol. 20:2036–

2039. Elsevier Ltd.

Heerwaarden, B. Van, M. Malmberg, and C. M. Sgrò. 2016. Increases in the

evolutionary potential of upper thermal limits under warmer temperatures in two

rainforest Drosophila species. Evolution (N. Y). 70:456–464.

Hoffmann, A. A., H. Dagher, M. Hercus, and D. Berrigan. 1997. Comparing different

measures of heat resistance in selected lines of Drosophila melanogaster. J. Insect

Physiol. 43:393–405.

Hoffmann, A. A., and P. A. Parsons. 1993. Direct and correlated responses to selection

for desiccation resistance: a comparison of Drosophila melanogaster and D.

simulans. J. Evol. Biol. 6:643–657.

Hoffmann, A. A., and C. M. Sgrò. 2011. Climate change and evolutionary adaptation.

Nature 470:479–85.

Huey, R. B. 2000. Rapid Evolution of a Geographic Cline in Size in an Introduced Fly.

Science (80-.). 287:308–309.

Huey, R. B., C. A. Deutsch, J. J. Tewksbury, L. J. Vitt, P. E. Hertz, J. Héctor, Á. Pérez,

T. Garland, P. R. S. B, R. B. Huey, C. A. Deutsch, J. J. Tewksbury, and L. J. Vitt.

2009. Why tropical forest lizards are vulnerable to climate warming. Proceeding R.

Soc. B 2009:1939–1948.

Huey, R. B., and R. D. Stevenson. 1979. Integrating thermal physiology and ecology of

ectotherms: a discussion of approaches. Am. Zool. 19:357–366.

Jørgensen, L. B., H. Malte, M. Ørsted, N. A. Klahn, and J. Overgaard. 2021. A unifying

model to estimate thermal tolerance limits in ectotherms across static, dynamic and

fluctuating exposures to thermal stress. Sci. Rep. 11:1–14. Nature Publishing

Group UK.

Jørgensen, L. B., H. Malte, and J. Overgaard. 2019. How to assess Drosophila heat

tolerance: Unifying static and dynamic tolerance assays to predict heat distribution

limits. Funct. Ecol. 33:629–642.

Kalra, B., A. M. Tamang, and R. Parkash. 2017. Cross-tolerance effects due to adult

heat hardening, desiccation and starvation acclimation of tropical drosophilid-

Zaprionus indianus. Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol.

209:65–73. Elsevier.

Kellermann, V., J. Overgaard, A. A. Hoffmann, C. Fljøgaard, J. C. Svenning, and V.

Loeschcke. 2012. Upper thermal limits of Drosophila are linked to species

distributions and strongly constrained phylogenetically. Proc. Natl. Acad. Sci. U.

S. A. 109:16228–16233.

Kelly, M. W. 2022. Experimental evolution reveals complex responses to

environmental change. Proc. Natl. Acad. Sci. U. S. A. 119:1–2.

Kelly, M. W., E. Sanford, and R. K. Grosberg. 2012. Limited potential for adaptation to

climate change in a broadly distributed marine crustacean. Proc. R. Soc. B Biol.

Sci. 279:349–356.

Krebs, R. A., and V. Loeschcke. 1996. Acclimation and selection for increased

resistance to thermal stress in Drosophila buzzatii. Genetics 142:471–479.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest package:

Tests in linear mixed effects models. J. Stat. Softw. 82:1–26.

Kwan, L., S. Bedhomme, N. G. Prasad, and A. K. Chippindale. 2008. Sexual conflict

and environmental change: trade-offs within and between the sexes during the

evolution of desiccation resistance. J. Genet. 87:383–394.

Lande, R., and S. J. Arnold. 1983. The measurement of selection on correlated

characters. Evolution (N. Y). 37(6):1210–1226.

Lenski, R. E., and A. F. Bennett. 1993. Evolutionary response of Escherichia coli to

thermal stress. Am. Nat. 142:47–64.

Lenth, R., H. Singmann, J. Love, P. Buerkner, and M. Herve. 2018. Package

"Emmeans". R Package Version 4.0-3.

Mesas, A., and L. E. Castañeda. 2023. Evolutionary responses of energy metabolism,

development, and reproduction to artificial selection for increasing heat tolerance

in Drosophila subobscura. Evolution 77:509–518.

Mesas, A., A. Jaramillo, and L. E. Castañeda. 2021. Experimental evolution on heat

tolerance and thermal performance curves under contrasting thermal selection in

Drosophila subobscura. J. Evol. Biol. 34:767–778.

Mitchell, K. A., and A. A. Hoffmann. 2010. Thermal ramping rate influences

evolutionary potential and species differences for upper thermal limits in

Drosophila. Funct. Ecol. 24:694–700.

Mitchell, N., and K. D. Whitney. 2018. Can plants evolve to meet a changing climate?

The potential of field experimental evolution studies. Am. J. Bot. 105:1613–1616.

Molina, A. N., J. M. Pulgar, E. L. Rezende, and M. J. Carter. 2023. Heat tolerance of

marine ectotherms in a warming Antarctica. Glob. Chang. Biol. 29:179–188.

Overgaard, J., T. N. Kristensen, and J. G. Sørensen. 2012. Validity of thermal ramping

assays used to assess thermal tolerance in arthropods. PLoS One 7:e32758.

Passananti, H. B., K. A. Beckman, and M. R. Rose. 2004. Relaxed stress selection in

Drosophila melanogaster. Pp. 323–352 in M. R. Rose, H. B. Passananti, and M.

Matos, eds. Methuselah Flies: A Case Study in the Evolution of Aging. World

Scientific, Singapore.

Quintana, A., and A. Prevosti. 1990. Genetic and environmental factors in the resistance

of Drosophila subobscura adults to high temperature shock. Theor. Appl. Genet.

80:847–851.

R Development Core Team, R. 2011. R: A Language and Environment for Statistical

Computing. R Foundation for Statistical Computing.

Rezende, E. L., L. E. Castañeda, and M. Santos. 2014. Tolerance landscapes in thermal

ecology. Funct. Ecol. 28:799–809.

Rezende, E. L., M. Tejedo, and M. Santos. 2011. Estimating the adaptive potential of

critical thermal limits: methodological problems and evolutionary implications.

Funct. Ecol. 25:111–121.

Rion, S., and T. J. Kawecki. 2007. Evolutionary biology of starvation resistance : what

we have learned from Drosophila. 20:1655–1664.

Rodgers, E. M., and D. F. Gomez Isaza. 2021. Harnessing the potential of cross-

protection stressor interactions for conservation: A review. Conserv. Physiol. 9.

Rodgers, E. M., and D. F. Gomez Isaza. 2023. The mechanistic basis and adaptive

significance of cross-tolerance: a 'pre-adaptation' to a changing world? J. Exp.

Biol. 226:jeb245644.

Rogell, B., W. Widegren, L. R. Hallsson, D. Berger, M. Björklund, and A. A.

Maklakov. 2014. Sex-dependent evolution of life-history traits following

adaptation to climate warming. Funct. Ecol. 28:469–478.

Rusuwa, B. B., H. Chung, S. L. Allen, F. D. Frentiu, and S. F. Chenoweth. 2022.

Natural variation at a single gene generates sexual antagonism across fitness

components in Drosophila. Curr. Biol. 32:3161-3169.e7. Elsevier Ltd.

Santos, M. A., A. Carromeu-Santos, A. S. Quina, M. Santos, M. Matos, and P. Simões.

2021. No evidence for short-term evolutionary response to a warming environment

in Drosophila. Evolution (N. Y). 75:2816–2829.

Santos, M., W. C,spedes, J. Balany…, V. Trotta, F. C. F. Calboli, A. Fontdevila, and L.

Serra. 2005. Temperature-related genetic changes in laboratory populations of

Drosophila subobscura : Evidence against simple climatic-based explanations for

latitudinal clines. Am. Nat. 165:258–273.

Santos, M., L. E. Castañeda, and E. L. Rezende. 2012. Keeping pace with climate

change: What is wrong with the evolutionary potential of upper thermal limits?

Ecol. Evol. 2:2866–2880.

Schwasinger-Schmidt, T. E., S. D. Kachman, and L. G. Harshman. 2012. Evolution of

starvation resistance in Drosophila melanogaster: Measurement of direct and

correlated responses to artificial selection. J. Evol. Biol. 25:378–387.

Simões, P., I. Fragata, S. G. Seabra, G. S. Faria, M. A. Santos, M. R. Rose, M. Santos,

and M. Matos. 2017. Predictable phenotypic, but not karyotypic, evolution of

populations with contrasting initial history. Sci. Rep. 7:1–12.

Singh, K., M. Arun Samant, and N. G. Prasad. 2022. Evolution of cross-tolerance in

Drosophila melanogaster as a result of increased resistance to cold stress. Sci. Rep.

12:19536. Nature Publishing Group UK.

Stazione, L., F. M. Norry, F. H. Gomez, and P. Sambucetti. 2020. Heat knockdown

resistance and chill-coma recovery as correlated responses to selection on mating

success at high temperature in Drosophila buzzatii. Ecol. Evol. 10:1998–2006.

Therneau, T. M. 2023. A package for survival analysis in R. R package version 3.2-7.

van Noordwijk, A. J., and G. de Jong. 1986. Acquisition and Allocation of Resources:

Their Influence on Variation in Life History Tactics. Am. Nat. 128:137–142.

Venables, W. N., and B. D. Ripley. 2002. Modern Applied Statistics with S. Fourth edi.

Springer US, New York.

Visser, M. E. 2008. Keeping up with a warming world; assessing the rate of adaptation

to climate change. Proc. R. Soc. B Biol. Sci. 275:649–659.

Table 1. Mixed linear effect model for the knockdown time of *Drosophila subobscura*

assayed at four static temperature assays. For simplicity, results for the random effect

(replicate lines) are not shown because they were not statistically significant (see

Materials and Methods). Significant effects (P values < 0.05) are indicated in boldface

type.

Knockdown time	Selection	Sex	Selection × Sex
Static assay at 35ºC	$F_{3,170} = 0.62$ $P = 0.60$	$F_{1,170} = 8.64$ **$P = 0.004$**	$F_{3,170} = 0.64$ $P = 0.59$
Static assay at 36ºC	$F_{3,232} = 9.86$ **$P = 3.8×10^{-6}$**	$F_{1,232} = 2.65$ $P = 0.10$	$F_{3,232} = 0.74$ $P = 0.53$
Static assay at 37ºC	$F_{3,222} = 18.39$ **$P = 1.1×10^{-10}$**	$F_{1,222} = 0.001$ $P = 0.97$	$F_{3,222} = 2.05$ $P = 0.11$
Static assay at 38ºC	$F_{3,224} = 1.93$ $P = 0.13$	$F_{1,224} = 4.63$ **$P = 0.032$**	$F_{3,224} = 2.44$ $P = 0.07$

Figure 1. Heat-induced mortality in *Drosophila subobscura* flies assayed at four static

temperatures. Left panels show the heat knockdown time of slow-ramping control (solid

black line), fast-ramping control (dashed black line), slow-ramping selection (red line),

and fast-ramping selection lines (blue lines). The right panels show the heat knockdown

time of female (purple line) and male (green line) flies. Dotted lines indicate the median

knockdown time for each selection protocol (left panels) and sex (right panels).

Figure 2. (A) Thermal death curves for control (black solid and dashed lines) and

selected (red and blue lines) lines for increasing heat tolerance in *Drosophila*

subobscura. Symbols represent the average knockdown time at the different assay

temperatures. Each symbol represents the average knockdown time for each replicate

line for each thermal regime: slow-control (black circle), fast-control (black triangle),

slow-ramping (red circle), and fast-ramping (blue triangle). (B) Relationship between

CT_{max} and z for slow-ramping control (solid black line), fast-ramping control (dashed

black line), slow-ramping selection (red line), and fast-ramping selection lines (blue

lines). Each symbol represents the CT_{max} and z estimated for each replicate line.

Figure 3. Desiccation survival curves of (A) females and (B) males from control (black

line), slow-ramping selection (red line), and fast-ramping selection lines (blue lines) of

Drosophila subobscura. Dashed lines indicate the median mortality time for each

selection protocol (pooled replicate cages).

Figure 4. Starvation survival curves of (A) females and (B) males from control (black

line), slow-ramping selection (red line), and fast-ramping selection lines (blue lines) of

Drosophila subobscura. Dashed lines indicate the median mortality time for each

selection protocol (pooled replicate cages).

740