Kisspeptin fiber and receptor distribution analysis suggests its potential role in central sensorial processing and behavioral state control

Running title: Kiss1r and KP fiber mapping in rodent brain

Limei Zhang[1,2,*c], Vito S. Hernández[1,2,*c], Mario A. Zetter[1,5], Oscar R. Hernández-Pérez[1], Rafael Hernández-Gonzáles[6], Ignacio Camacho-Arroyo[7], Lee E. Eiden[2,c], Robert P. Millar[1,3,4]

[1]Dept. Physiology, School of Medicine, National Autonomous University of Mexico, Mexico City, Mexico

[2]Section on Molecular Neuroscience, NIMH-IRP, NIH, Bethesda, MD, USA

[3]Centre for Neuroendocrinology, University of Pretoria

[4]Dept. of Integrative Biomedical Sciences, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, South Africa

[5]Department of Medicine and Health, University of La Salle, Mexico City, Mexico

[6]Vivarium, School of Medicine, National Autonomous University of Mexico, Mexico City, Mexico

[7]Research Unit in Human Reproduction, National Institute of Perinatology-Faculty of Chemistry, National Autonomous University of Mexico, Mexico City, Mexico

[*]These authors share first authorship.

[c]Correspondence: limei@unam.mx, vitohdez@unam.mx

Abstract

Kisspeptin (KP) signaling in the brain is defined by its anatomical framework, including the locations and chemotypes of KP-producing neurons, their fibers, receptors, and connections. Recent technological advances prompted us to re-evaluate these chemoanatomical aspects, initially studied during the early years following the discovery of KP and identification of its receptor. We recently characterized seven KP neuronal populations in the mouse brain using sensitive RNAscope methods, with two populations as new additions to the literature, and their short-term response to gonadectomy [1]. These results prompted us to continue in mapping KP fiber distribution in rats and mice and examined *Kiss1r* mRNA expression relative to that of the vesicular GABA transporter mRNA (*Slc32a1*), which provides a major chemoanatomical marker throughout the mouse brain. We identified KP fibers in 118 brain regions, predominantly in extra-hypothalamic areas involved in sensory processing and behavioral state control. Remarkably, KP fiber density was similar in aged long-term (one year) gonadectomized compared to intact mice, either male or female. *Kiss1r* was prominently expressed in brain sensory and state control regions, such as septal nuclei, tuberomammillary nuclei, the suprachiasmatic nucleus, the locus coeruleus, the hippocampal pyramidal and granule cell layers, the thalamic nuclei, and the cerebellar granule cell layer, as well as the interposed nucleus. *Kiss1r* was also co-expressed in *Kiss1*-expressing hypothalamic neurons and surrounding regions, suggesting autocrine and paracrine modes of KP action. These findings provide additional insight for further exploration of KP's role in brain control of complex physiological processes, independently of reproductive status.

key words: behavior state system, sensory-motor circuits, extrahypothalamic kisspeptin signaling, Kiss1r, autocrine

1. Introduction

Kisspeptin (KP) is a neuropeptide encoded by the *Kiss1* gene, initially known as metastin, identified via its role in inhibiting cancer metastasis [2]. It is now well-recognized as a critical regulator of the mammalian reproductive system, playing a pivotal role in the activation of the hypothalamo-pituitary-gonadal (HPG) axis by stimulating the release of gonadotropin-releasing hormone (GnRH) [3, 4]. Beyond its reproductive functions, KP is increasingly implicated in a variety of physiological processes, including the regulation of emotions and cognitive functions [5]. For instance, kisspeptin receptors are expressed in the amygdala, a critical region for emotional processing, and their activation has been associated with anxiolytic effects and the modulation of stress responses [6]. Additionally, KP has been implicated in cognitive functions such as learning and memory, potentially through its interactions with the hippocampus and prefrontal cortex [7]. These findings are supported by behavioral studies in animal models demonstrating that KP administration can enhance memory retention and reduce anxiety-like behavior [8] and modify fear response [9, 10]. As research continues to elucidate the molecular pathways and neural circuits involved, it has become increasingly clear that KP should play a multifaceted role in brain function, beyond reproductive function.

In a recent study in mouse, our group reported a comprehensive KP cell population description and molecular signature characterization within excitatory or inhibitory neurons throughout mouse brain, using a sensitive dual and multiple-channels in situ hybridization method. We described seven distinct KP populations, four of them within the hypothalamus and three of them as extrahypothalamic populations, two of which are newly-identified KP cell groups. Regulation of KP expression by short-term gonadectomy (GNX) was also reported quantitatively [1]. These results prompted us to extend the study by examining kisspeptin immunopositive fiber distribution in both rat and mouse, in comparison with KP receptor (Kiss1r) distribution using RNAscope methods.

There had been comprehensive anatomical studies on KP neuronal populations and fiber distributions in the brain of diverse animal species in the literature of the first decade after the discovery of KP, albeit hypothalamus and reproductive functions were the main focus of those studies, with less attention to KP expression in extrahypothalamic regions [11] [7, 12-14]. We examined KP fiber distributions in whole rat and mouse brain serial samples using a pan-specific KP antibody which recognizes the Kp decapeptide epitope within Kp-54, 14 and 10, with a less than 0.1% cross reactivity with other RF-amide peptides [15] and an immunohistochemistry procedure that maximizes staining sensitivity and reproducibility followed in our laboratories [16]. We report here 118 brain regions that contain KP fibers and terminals, with most of them located in extra-hypothalamic regions associated with central sensory processing and behavioral state control [17]. Surprisingly, the down-regulation of kisspeptin signaling from the rodent rostro-periventricular region (RP3V), observed after a short period of GNX in rodents reported previously by several groups including our own [1] was not evident in animals after long-term (12 months) GNX. Data from this study provide a new functional anatomical basis for further hypothesis generation and testing of the role of KP in a wider scope of brain function beyond reproductive function.

87 2. Results and focused discussion
88

89 2.1 Regional distribution of kisspeptin-expressing fibers and neurons in the mouse and rat brain
90

Due to the emerging evidence for kisspeptin function beyond its well-documented role in orchestration of
GnRH secretion in the hypothalamus, it is imperative that reliable data concerning the comprehensive
distribution of kisspeptin cell bodies and nerve fibers be obtained. At this time, there appear to be data
sets that are reliable but incomplete, and some that, while comprehensive, have been questioned as
potentially unreliable, due to adventitious cross-reactivity with other RF peptides. We have used a KP
antibody with now well-documented authenticity and lack of cross-reactivity, and applied state-of-the-art
tissue preparation and staining methods, in order to furnish a reliable and comprehensive description of
the distribution of KP cell bodies and fibers throughout the brain contributing to a better understanding
of the possible role(s) of KP first messenger function relevant to both reproductive and non-reproductive
behavior beyond the regulation of GnRH secretion from the hypophysiotropic hypothalamus.
101

In this study, we performed KP-immunohistochemistry (IHC) using the pan-specific KP antibody
which recognizes the KP decapeptide epitope within KP-54, 14 and 10, with a less than 0.1% cross
reactivity with other RF-amide peptides [15] and an improved immunohistochemistry procedure followed
in our laboratories [16], with picric acid added to the conventional fixative, strict time frames for fixation
(15 min) followed by thorough washing, tissue slicing, and initiation of the IHC reaction immediately
thereafter, coupled with long incubation with primary antibody (2 days at 4°C), to maximize the
visualization of KP-positive fibers and neurons throughout the rodent brain. With this method, we
observed greater intensity and a wider expression pattern of KP immunoreactivity in both mouse and rat
brain than previously reported [11] [7, 12-14]. We found only one previous comprehensive report on the
distribution of KP immunoreactivity in the rodent brain. In that study, authors used the antibody produced
by Phoenix Pharmaceuticals [18] which has been cited as potentially cross-reactive with other RFRP
peptides [12]. We have included the data reported by Brailoiu et al., [18] in rat, using the Phoenix
Pharmaceuticals antibody anti-metastin (MT), in the table 1, with blue letters to facilitate comparison with
the findings of the current investigation.
116

117 In table 1 (Tab. 1), we report 118 regions of the rodent brain in which we observed KP-ir labelling
118 with differential strengths. When making detailed comparison with a previous publication [18], we found
119 all the regions reported in this early study were replicated with our experiments. However, numerous new

Table 1. Distribution of kisspeptin immunoreactive (antibody GQ2) fiber distribution in mouse & rat brain with comparison of Brailoiu et al, JCN, 2005 (metastin, MT)

Structure	Abbr.	irMT cells (rat)	irGQ2 cells rat	irGQ2 cells mouse	irMT fibers (rat)	irGQ2 fiber rat	irGQ2 fiber mouse	Axons from RP3V viral tracers	Axons from Arc viral tracers
Telencephalon									
Cerebral cortex									
Cingular cortex	Cg		-	-		++	+		
Dorsal peduncular cortex	DP		-	-		++	+		
Dorsal tenia tecta	DTT		-	-		++	+	+	
Infralimbic cortex	IL		-	-		++	+		
Prelimbic cortex	PrL		-	-		++	+		
Ventral orbital cortex	VO		-	-		++	+		
Olfactory area									

Region	Abbr								
Lat. septal n. ventral	LSV		-	-	++	+++	+++	+++	++
Medial septal nucleus	MS		-	-	+	+++	+	+	+
Septo-fimbrial n.	SFi		+	+		+	+	++	
Septo-hippocampal n.	SHi		+	+		+	+		
N. horizontal limb of the diagonal band	HDB		-	-	+	+	+	++	+
N. vertical limb of the diagonal band	VDB		-	-	+/-	+	+	+++	+
Diencephalon									
Hypothalamus									
Anteroventral periventricular n.	AVPe		+++	+++		+++	+++	+++	+++
Rostral periventricular area	RP3V		+++	+++		+++	+++		
Anterior hypothalamic area anterior	AHA		-	-		++	++	++	+++
Anterior hypothalamic area posterior	AHp		+	+	+	++	++		+++
Arcuate nucleus	Arc	+	++	++	+++	++++	++++	+++	++++
Dorsal hypothalamic	DH		++	++	+	+	+	+++	
Dorsomedial hypothalamic area	DM	+++	++	++	++	++	++	+++	++
Lat. hypothalamic area	LH		-	-	+	+	+	+	+
Lat. preoptic area	LPO		-	-	+	++	++	+++	
Medial preoptic area	MPA		-	-	+	++	++	+++	++
Median preoptic n.	MnPO		-	-	+	+++	+++	++++	+++
Parastrial nucleus	PS		-	-	+++	+++	+++		
Paraventricular hypothalamic n.	PVN	+	-	-	+	++	+	+++	+
Paraventricular hypothalamic n. medial magnocellular div	PVNmm		-	-		+++	+++	+++	+
Pa hypothalamic n. lateral magnocellular div.	PVNlm		-	-		+	+	+++	+
Pa hypothalamic n. lateral parvocellular div.	PVNmp		-	-		+++	+++	++	
Periventricular hypothalamic n. anterior	PVa		-	-		++	++		+++
Premammillary n. ventral	PMV		+	+		+++	++	+++	+++
Posterior hypothalamic area	PH		-	-	+	+	+	++	++
Retrochiasmatic area	RCh		-	-	+	+	+	++	++
Supraoptic nucleus	SON		-	-	+	++	++	+	
Tuber cinereum area	TC		-	-	+	++	+	++	
Ventral tuberomammillar n.	VTM		-	-		+++	++		+++
Ventromedial hypothalamic nucleus	VMH	+	-	-	+	+	+		+
Thalamus									
Anteromedial thalamic nucleus	AM		-	-		++	++		
Central medial thalamic nucleus	CM		-	-	+	+	+	++	
Mediodorsal thalamic nucleus	IMD		-	-	+	+	+		
Paratenial thalamic nucleus	PT		-	-	+	+	+		
Periventricular thalamic nucleus	PVT		-	-	+++	++++	++++	++	+
Reuniens thalamic nucleus	Re		-	-	+	+	+		
Rhomboid thalamic nucleus	Rh		-	-	+	+	+		
Submedius thalamic nucleus	SubD		-	-		+	+		
Zona incerta	ZI		-	-	+	+	+		
Epithalamus									
Habenular nucleus	Hb		-	-	++	++	++	+	
Mesencephalon									
Central nucleus of the inferior colliculus	CIC		-	-	+	++	++		

Region	Abbr								
Cuneiform nucleus, dorsal part	CnFD		-	-		++	++		
Dorsal raphe nucleus	DR		-	-	+	++	++		
Deep mesenceph. n.	DpMe		-	-	+	+	+		
Interpeduncular n.	IP		-	-		+	+		
Median raphe n.	MnR		-	-	+	+	+		
Periaqueductal gray, lateral	LPAG		-	-		++	++	++	+
Periaqueductal gray dorsomedial	DMPAG	+/-	-	-	+++	+++	+++	++	+
Substantia nigra	SN		-	-	+	+	+		
Superior colliculus, intermediate and deep gray layer	DpG		-	-	+	+	+		
Supramammilary n.	SuM		-	-	+	+	++		
Ventral tegmental area	VTA		-	-		+	+	+	
Metencephalon and myelencephalon									
Ambiguus nucleus	Amb		-	-	+	+	+		
Area postrema	AP		+	+		+++	+++		
Caudoventrolateral reticular nucleus	CVL	+	-	-	+	++	++		
Dorsomedial tegmental area	DMTg		-	-		+	+		
Gigantocellular reticular nucleus	Gi		-	-	+	++	++		
Intermediate reticular nucleus	IRt		-	-	++	++	++		
Lateral parabrachial nucleus	LPB		-	-	+++	++++	++++	+	
Lateral reticular nucleus	LRt	+			++	++	++		
Lateral superior olive	LSO		-	-	++	+	+		
Locus coeruleus	LC		-	-	++	+++	+++	+	
Medial vestibular nucleus, magnocellular part	MVeMC		-	-	.	++	++		
Medioventral periolivary nucleus	MVPO		-	-		+++	+++		
Medullary reticular nucleus, dorsal part	MdD		-	-	+	++	++		
Medullary reticular nucleus, ventral part	MdV		-	-	+	+	+		
Nucleus of the solitary tract	Sol	+++	++	++	+++	++++	++++		
Parvicellular reticular nucleus	PCRt		-	-		++	++		
Parabrachial pigmented nucleus	PBP		-	-		+	+		
Pontine nuclei	Pn		-	-	+	-	-		
Pontine n, brachium pontis	bpPn		-	-		+	+		
Pontine raphe nucleus	PnR		-	-	+	-	-	+	
Pontine reticular nucleus, caudal part	PnC		-	-	+	+	+	++	
Pontine reticular nucleus, oral part	PnO		-	-	+	+	+	++	
Principal sensorial 5 n. dorsomedial	Pr5DM		-	-		++++	++++		
Raphe magnus nucleus	RMg		-	-	+	+	+		
Raphe obscurus nucleus	Rob		-	-	+	+	+		
Raphe pallidus nucleus	RPa		-	-	+	+	+		
Rostroventrolateral reticular nucleus	RVL	-	-		++	++	+		
Rubrospinal tract	rs	-	-	-	++	++	++		
Spinal trigeminal tract	Sp5	+	-	-	+++	-	-		
Spinal 3, gelatinous layer	Ge5		-	-		++++	+		
Superior cerebellar peduncle	scp		-	-		+++	+++		

In figure 1 we show KP immunohistochemical reactivity (-ir) in regions, including AVPV and VLPO where KP functions to regulate GnRH secretion, in comparison with other brain regions, with equally intense KP innervation, that are relevant in regulation of brain state, including MnPO, SHy, MS, LS and BST (panel A). Also depicted (Panel B), micrograph of a coronal section around Bregma -0.15mm, where 4 brain main conducting fiber systems can be clearly seen, i.e. the anterior commissure (ac), the stria medullaris (sm), the stria terminalis (st) and the fornix (f). KP-ir fibers running inside these conducting systems are indicated with red arrows. The dense innervation pattern of KP-ir fibers in the paraventricular anterior nucleus of the thalamus is shown. In panels C and D we show examples of KP-ir cell bodies and fibers in amygdalar complex. KP-ir fibers can be seen inside the stria terminalis (st), bed nucleus of stria terminalis, intra-amygdalar division (STIA), baso-medial amygdala (BMA), nu. basalis of Meynert (NBM) and medial amygdala (MEA). Some fibers within the medial amygdala, postero-dorsal division (MEApd), are observed coming from the stria terminalis, with main axons branching within the MEApd, of possible hypothalamic origin. In Tab 1, we summarize an analysis of experiments of two Kiss1-Cre mice injected Cre-dependent tracer in the arcuate nucleus (Arc, experiments 232311236 and 232310521) and two in the rostral periventricular region (R3PV, experiments 299247009 and 301989585) from the Allen Mouse Brain Connectivity Atlas. Fibers appear to reach amygdala via the stria terminalis (C1, red arrows). Panels Es and Fs show photomicrographs from the hypothalamic tuberal region, where the KP-ir cell bodies are seen in arcuate nucleus, and in premammillary nucleus, ventral division. KP-ir fibers are observed in other regions relevant for behavioral state control at this level, such as ventral, magnocelular and dorsal tubero mammillary nu., posterior hypothalamus, zona incerta (ZI) and subthalamic nu. (STN) Panel E2 shows the third ventricle at the level of Arc posterior. Abundant KP-ir fibers extended toward the lumen of the ventricle (also see the A3, red arrows). Panels G-K are micrographs of samples from parasagittal sections of mouse hindbrain. Panel G show the KP-ir cell bodies and dense KP-ir fibers in the solitary tract (sol) nu. (NTS) of mouse medulla. H and I show the dense innervation pattern of KP-ir fibers in pontine nuclei: lateral parabrachial nu. (LPBN) and locus coeruleus (LC) respectively. J and K show the dense KP-ir fibers in the spinal trigeminal tract (Sp5, K) and in the gelatinous layer of Sp5 (Ge5). Note that within the Ge5, the KP-ir fiber make perisomatic contacts with the cell bodies of the Ge5 (J, red arrows).

In Figure 2, we present chartings of mouse KP cell and fiber distribution. Panels A, B and C, (in register with [11] for purposes of comparison) show intense KP innervation of the prefrontal structures, such as cingular cortex (Cg), dorsal peduncular cortex (DP), dorsal tenia tecta (DTT), infralimbic cortex

(IL), prelimbic cortex (PrL), ventral orbital cortex (VO), nucleus accumbens shell (AcbSh), medial septal
nucleus (MS), nucleus of the diagonal band of Broca (NDB) and, dorsal division of the lateral septal
nucleus (LSd), which were not reported in the cited report. We also observed abundant innervation in
thalamic structures such as nucleus reuniens (Re), periventricular anterior nucleus (PVA) and the
epithalamic lateral habenula (LHb). In the amygdaloid complex, we observed KP fibers in cortico-
amygdala (CoA), nucleus of lateral olfactory tract (NLOT), the medial amygdala (MEA), mainly in the
postero-dorsal division, (MEApd), the basomedial amygdala (BLA) and the nu. of the stria terminalis,
intra-amygdala division (BSTIA) (Fig. 2, D-G). Midbrain and hindbrain sensory relay and state control
centers, such as premammillary nu. (PM), dorsal raphe nu. (DR), periaqueductal gray (PAG), subthalamic
nu. (STh) parabrachial complex (PBc), locus coeruleus (LC), Kölliker-Fuse nu. (KF) and nucleus of
solitary tract (NTS) are all seen with KP-ir fibers which were not reported previously (Fig. 2, H-L).
181

Fig. 1 KP immunohistochemical reactive (-ir) samples in mice using a polyclonal sheep anti-kisspeptin antibody [15] (GQ2, kindly supplied by Mohammad Ghatei†, Imperial College, London, UK), see Material and Methods section). This antibody recognizes kisspeptins 54, 14,13, and 10 that conserve the C-terminal decapeptide essential for biological activity and shows virtually no cross-reactivity (<0.01%) with other related human RF-amide peptides [15]. Panel A shows a low magnification micrograph at coronal plane (Bregma 0.38mm dorsal, 0.02 ventral [19]), where hypothalamic kisspeptinergic regions (regions containing KP-ir cell bodies, such as the antero-ventral periventricular nucleus (AVPV,A3), ventrolateral preoptic nucleus (VLPO) are shown. Besides, regions with high densities of KP innervation, which are relevant for brain state control, such as medial nucleus of preoptic area (MnPO), septo-hypothalamic nucleus (SHy, A1, A5, which also contains KP cell bodies, indicated with black arrows. For ISH image of SHy Kiss1 expressing neurons please see [1]), medial septal nucleus (MS, A2)), lateral Septal nu. (LS), bed nucleus of stria terminalis (BST, A4), with its subdivisions medio-anterior (STMA), medio-ventral (STMV) and latero-ventral (STLV). A3 shows the KP-ir in AVPV, where KP-ir neurons are indicated with black arrows and abundant KP-ir fibers innervating the ventricular surface are indicated with red arrows. Panel B, micrograph of a coronal section around Bregma -0.15mm, where 4 brain main conducting fiber systems can be clearly seen, i.e. the anterior commissure (ac), the stria medullaris (sm), the stria terminalis (st) and the fornix (f). KP-ir fibers running inside these conducting systems are indicated with red arrows. Also, the dense innervation pattern of KP-ir fibers in the paraventricular anterior nucleus of the thalamus is shown. Panel C shown a low magnification coronal micrograph of amygdala around the coordinate Bregma -1.58mm. KP-ir fibers can be seen in stria terminalis (st), bed nucleus of stria terminalis, intra-amygdalar division (STIA), baso-medial amygdala (BMA), nu. basalis of Meynert (NBM) and medial amygdala (MEA). Interestingly, some fibers within the medial amygdala, postero-dorsal division (MEApd), are observed coming from the st, that the main axon entered the MEApd and branched locally (C1, red arrows). Panel D shows a micrograph of MEApd at Bregma -2.18mm. Numerous KP-ir cell bodies are indicated with red arrows. D1 shows a lower magnification micrograph with the the circunscript region being the one of the panel D. Panels Es and Fs show the migrophotographs taken from the hypothalamic tuberal region, where the KP-ir cell bodies are seen in arcuate nucleus (Arc. E and F1), in Premammillary nucleus, ventral division (PMv, E and E1, F3, for ISH image of PMv Kiss1 expressing neurons please see [1]) and in the dorso-medial hypothalamic region (DMH, F, F1 and F2, black arrows). KP-ir fibers are observed in other regions relevant for behavioral state control at this level, such as ventral tuberomammillar nu. (VTM), magnocelular tuberomammillar nu. (MT) and dorsal tubero mammillary nu. (DTM), posterior hypothalamus (PH), zona incerta (ZI) and subthalamic nu. (STN) (E). E2 shows the third ventricle at the level of Arc posterior. Abundant KP-ir fibers extended toward the lumen of the ventricle (red arrows).Panel Gs- K are micrographs of samples from parasagittal sections of mouse hindbrain. Gs show the KP-ir cell bodies and dense KP-ir fibers in the solitary tract (sol) nu. (NTS) of mouse medulla. H and I show the dense innervation pattern of KP-ir fibers in pontine nuclei: lateral parabrachial nu. (LPBN) and locus coeruleus (LC) respectively. J and K show the dense KP-ir fibers in the spinal trigeminal tract (Sp5, K) and in the gelatinous layer of Sp5 (Ge5). Note that within the Ge5, the KP-ir fiber make perisomatic contacts with the cell bodies of the Ge5 (J, red arrows).

Fig. 2 Charting to represent the distribution of kisspeptin immunoreactivity in mouse brain. Sequential atlas planes show in the left side, KP immunoreactive cell bodies from the seven KP cell populations. Color-filled circles symbolizing the molecular signature previously reported [1]: KP-glutamatergic (green), KP-GABAergic (red) and expressing both VGluT2 and VGAT (blue). The distribution of KP-ir fibres are represented by red lines in the right hemisphere (the thickness of the lines has been exaggerated to ensure that the lines can be seen on the atlas planes). KP-ir fibers observed in brain areas relevant for behavioral state control are shaded in yellow: accumbens nucleus, shell (AcbSh), septohippocampal nu. (SHi), lateral septal nu. dorsal (LSD), medial septal nu. (MS), horizontal division of nucleus of dianogal band of Broca (HDB), septohypothalamic nucleus (SHy), bed nucleus of stria terminalis (BST), medial preoptic area (MPO), paraventricular anterior parvicellular division (PaAP), nucleus basalis of Meynert (NBM), lateral habenula (LHb), lateral hypothalamus (LH), tuberomammillary nu. (TM), ventral Premammillary nu. (PMv), periaqueductal grey (PAG), dorsal raphe nu. (DR), locus coeruleus (LC), lateral parabrachial nu. (LPB), nu. of solitary tract (NTS). See table 1 for rest of abbreviations.

244 *Fig. 3 Mapping of KP immunoreactive fibers and cell bodies **in rat brain** in ten septo-temporal planes based on*
245 *microscopic observations. Blue dots represent KP-ir cell bodies and red lines represent KP-ir fibers. Relevant*
246 *anatomical regions have supporting photomicrographs form IHC-DAB reactions using the GQ2 anti-kisspeptin*
247 *antibody. See table 1 for abbreviations.*

 <u>2.2 Detailed description of regional distribution of KP fibers with an emphasis on sensory processing and</u>
250 <u>behavioral state control</u>
251
252 The distribution of kisspeptin fibers in several regions involving numerous crucial central sensory relays
253 and behavioral-state control structures suggests wider participation of this peptide in other functions
254 besides its known participation in reproduction and energy balance. In the following sections, we describe
255 these regions in detail.
256
257 *2.2.1 Telencephalon*
258
In rodent telencephalon, the most prominent KP fiber expressing regions are located in the *lateral*
septal region in male and female mice and rats. The strongest expression was observed in the dorsal lateral
septal nucleus (LSD) (Fig. 1 A & A2, Fig. 2 B-D, and Fig 3 B-C*),* followed by ventral LS (LSV) (Fig. 2
C, and 3 C). LS is involved in the integration of sensory and emotional information, and can modulate the
activity of other brain regions involved in the regulation of behavioral states, such as the hypothalamus,
amygdala and hippocampus [20, 21]. Recent studies have reported that the main afferents of the above
regions are the subdivisions of the hippocampal formation and the main efferent regions of the LSD are
the ventral tegmental area and the supramammilary nucleus, whereas the LSV project mainly to the medial
hypothalamic area and medial preoptic area [21]. Activity of lateral septum can lead to changes in
behavioral state, including increased exploratory behavioral, social interactions and decreased anxiety,
particularly in response to social and emotional stimuli. These changes were reported from clinical studies
with KP infusion in human subjects [22]. LS is also a key player for arousal and wakefulness, as well as
for regulating processes related to mood and motivation through its extensive connections with numerous
regions such as the hippocampus, amygdala, hypothalamus and with the mesocorticolimbic dopamine
system [23].
274
The *medial septum* (MS) also contained mild and sparse innervation in rats and mice (Fig. 1 A and
2 B). This structure is crucially involved in regulating the hippocampal oscillations [24, 25], and recent
studies have shown that the medial septum acts as a neural hub that regulates theta oscillations and
locomotion speed, influencing goal-directed behaviors such as spatial navigation [26]. Other behaviours
influenced by the medial septum include sleep, attention, learning and memory, motivation and emotional
processing [27-29].
281
282 Sparse fibers were found in *septofimbrial nucleus* (Fig. 3 B) which is an important source of
283 cholinergic inputs to the hippocampus, the entorhinal cortex, influencing hippocampal plasticity and
284 learning and memory under emotional stimuli; and the septohippocampal (SHi, Fig. 2 B) nuclei that
285 provide important GABAergic inputs to hippocampus playing a role in the generation of hippocampal
286 theta oscillation [30]. The above two regions are described for the first time in this study.
287
Similar to Brailoiu et. al [18], we found sparse fibers in the *horizontal and vertical limbs of the*
nucleus of diagonal band of Broca (NDB) (Fig. 2 B). The diagonal band of Broca interconnects the

amygdala and the septal area and projects extensively to hippocampus CA2 subfield which is involved in
memory retrieval and social functions [31, 32]. It contains the second largest cholinergic neuronal population
in the brain (after the nucleus basalis of Meynert) [33]. It is particularly important in the aging of the brain,
since the cholinergic neuronal loss in dementias has been commonly observed in human. The influence of
KP to brain cholinergic system is under-investigated.
295
296
The presence of kisspeptin fibers in the *nucleus accumbens,* in both core and shell divisions (AcbC
and AcbSh), with a mild density in mice and a moderate density in rats (Table 1, Fig. 2 A and 3 B-D) was
a remarkable finding of this study. AcbC receives inputs from the prefrontal cortex, the amygdala, and the
ventral tegmental area (VTA), which is a key component of the brain reward system. The AcbC integrates
these inputs to evaluate the salience and value of rewards and thus helps to regulate motivated behavior.
The nucleus accumbens shell is an important brain region involved in the processing of reward-related
information and the regulation of motivated behavior, including food intake, drug seeking, and social
behavior [34].
305
306   Sparse fibers were observed in the *caudate putamen* (CPu Fig. 2 D & 3 C), ventral pallidus (VP,
307 Fig. 3 D), the *sustantia innominata* (SI), as well as main fiber pathways such as *the internal capsule* (ic)
308 and *external capsule* (ec)
309
In telencephalic *neocortical regions*, we found sparse fibers distributed through the *dorsal tenia*
tecta (DTT) and the *dorsal peduncular* (DP), *prelimbic* (PrL) , *infralimbic* (IL), and *cingular* (Cg) cortices,
regions that have not been reported in the Brailou study. The observed fibers were mainly located in layers
2/3, and no differences between species were observed (Table 1 and Fig. 2 A & 3 B). The *prelimbic* (PrL),
infralimbic (IL) and *cingulate cortices* (Cg) have been classically considered to be involved in mediating
the role of context in the expression of fear or reward induced behaviors [35] [36, 37]. The DTT and the DP
have been classically involved in olfactory information processing, and recently these two structures have
been shown to send inputs with information related to psychosocial stress to the dorsomedial
hypothalamus, where a cardiovascular and thermogenic sympathetic stress response can be integrated [38].
In other areas classically related to the processing of olfactory information, such as the *ventral and*
ventroposterior of the *anterior olfactory nucleus*, (AOV and AOVP) and the *olfactory tubercle* (Tu), we
found sparse fibers only in the rat (Tab. 1, Fig. 3 B-D).
322
In the *amygdalar complex*, we confirmed the KP-ir+ cell population mainly confined within the
medial amygdala's postero-dorsal division (MePD) which has been reported already elsewhere [39, 40] (Tab.
1 & Figs. 1 C-D, 2 F-G and 3 F-G). Kisspeptin immunolabelled neurons in the MePD and an axon terminal
process were observed to innervate the adjacent nucleus of the stria terminalis intra-amygdaloid division
and the stria terminalis. Also, we observed KP expressing somas in the septo-hippocampal (SHi) and
septo-fimbrial (SFi) region surrounding the anterior commissure, and extending into the septum, these
neurons were characterized neurochemically in mice in our previous report [1].

In relation to density of fibers in the septo-bed nucleus of stria terminalis complex, regions with a moderate/intense density of fibers were found within the septal part of the *bed nucleus of the stria terminalis* (BSTMA, BSTMP, BSTLV and BSTLP, Fig. 1 A; Fig. 2 B-D and Fig. 3 B-C). The *temporal intra-amygdala division of the BST* (BSTIA, Fig. 1 C; Fig. 2 F and Fig. 3 H) showed a selective expression of KP fibers. Other regions in the extended amygdala that also showed mild expression of KP fibers were the central amygdalar nucleus (CeC and CeM Fig. 3 G-H) and the medial amygdalar nucleus (MePD and MePV Fig. 2, F-G and 3. G-H). Sparse fibers were found in the *oval nucleus of the stria terminalis* (BSTLD), the *basolateral amygdala* (BLA), the *basomedial amygdala* (BMA), the *lateral part of the central amygdalar nucleus* (CeL), the *cortico-amygdala* (CoA) the *nucleus of the lateral olfactory tract* (NLOT) and the *stria terminalis* (*st*, Fig. 1 C). Compared with the study of Brailou et al, here we report the presence of fibers in the anterior, medial, and cortical amygdala and the intra-amygdaloid division of the stria terminalis (Tab. 2).

2.2.2. Diencephalon

The *hypothalamus* is the region where four of the seven KP neuronal populations have been reported recently by our group, i.e. the rostral periventricular (RP3V) population mainly located in the *medial preoptic area* (MPO), and in the *anteroventral periventricular area* (AVPe) , the arcuate (Arc) population, the dorsomedial hypothalamus (DMH) and the ventral premammillary nu. (PMV) (Fig. 1A; Fig. 2 C,D,F,G,H and Fig. 3 B). After a careful analysis of immunoreacted serial sections, following the pathways of the labelled fibers, we identified some extrahypothalamic regions targets of these regions. Ascending fibers originated in the RP3V population, travel lateral-dorsal-caudally toward the *stria medularis* (*sm*), and the *stria terminalis* (*st*) conduction systems (Fig. 1, B). In their path, they extensively innervate the *bed nucleus of the stria terminalis* (Tab. 2). It is worth noting the presence of KP fibers innervating the border between the *medial (MHb) and lateral (LHb) habenula* (Fig. 2, F-G and Fig. 3 B), it is known that the main input to the habenula is the stria medularis (*sm*). Other fibers originating in the MPO/AVPV region travel to innervate the *organum vasculosum of the lamina terminalis* (OVLT) and then ascend dorsally and laterally to innervate the *accumbens core* (AcbC) and the *lateral septum* regions.

We observed moderately high densities of KP-ir fibers (see Tab. 2) in the caudal periventricular area and ventral periventricular areas of the *arcuate nucleus* (Arc, Fig. 1 E, and Fig. 2 F-H). Some targets of the KP neural population in the *posterior hypothalamus* were identified by following the labelled axon in serially collected sections in coronal and parasagittal planes Some fibers originating in the Arc travelled in the base of the brain reaching the *internal capsule* (*ic*) and *optic tract* through which they travel in a path to the *subthalamic nucleus* (STh), the lateral part of the *supramammillary nucleus* (SuML) and continuing to the *central nucleus of the amygdala* (CeA). Some efferent fibers traveled caudally to innervate the ventral premammillary nucleus (VPM), the *ventral tuberomammillary nucleus* (VTu) the *supramammillary nucleus* (SuM). Some additional fibers surrounded the ventromedial hypothalamus (VMH) and travelled dorsally to innervate the anterior hypothalamic area and continued to innervate medial nuclei of the thalamus including the paraventricular thalamic (PVT) that showed the highest

370   innervation, but also the Intermedio dorsal (IMD) the centro medial (CM), medio dorsal (MD), paratenial
371   (PT), rheuniens (Re) and rhomboid (Rh) nuclei of the thalamus.
372
373   The population of weakly KP immunopositive cells scattered in the *medio-lateral dorsal*
374   *hypothalamic area* (DH) spanning from the posterior part of the *anterior hypothalamus* (AHp) to the
375   posterior hypothalamus (PH), centered in the *tuber-cinerum area* (TC) (Fig.   1 F; Fig. 2 F-H, and Fig. 3
376   B) were observed to project to immediately located *zona incerta* (ZI). Finally, the last population of KP
377   neurons in the PMV was very weakly labelled and it was not possible to identify the axonal projections.
378
Other relevant regions in the hypothalamus with KP fibers presence are the *median preoptic*
nucleus (MPO), the *parastriatal nucleus* (PS), the *paraventricular nucleus medial magnocellular division*
(PVNmm) and the *parvocellular division* (PVNmp), and in rat also in the ventral *premammillary nucleus*
(PMV) and *ventral tuberomammillary nucleus* (VTu) and *suprammamilary nucleus* (SuM). A mild density
of fibers was observed in *dorsomedial hypothalamic nucleus*, MPO , lateral preoptic area, the
periventricular hypothalamic nucleus anterior (PVa), and the *supraoptic nucleus* (SO). Sparse fibers were
found in the *dorsal hypothalamus* (DH), *zona incerta* (ZI), *lateral hypothalamic area* (LH), *posterior*
hypothalamic area (PH), *retrochiasmatic area* (RCh), and *ventromedial hypothalamic nucleus* (VMH).
The KP system in the hypothalamus has been extensively discussed in the literature [12].
388
In the *thalamus*, the highest density of fibers was observed in the *paraventricular thalamic nucleus*
(PVT) in rats and mice (Fig. 1 B; Fig. 2 E-F and 3 B). Mild fibers densities were also observed in other
nuclei of the thalamus, such as in the *centromedial* (CM), *mediodorsal* (MD), *paratenial* (PT), *reuniens*
(Re), and *rhomboid* (Rh). The epithalamic *habenular nucleus* (Hb) showed a moderate density of fibers
mainly located in the border between its medial and lateral subdivisions (Fig. 2 F-G and 3 B), some of
these fibers were observed arriving trough the stria medullaris.
395
396   *2.2.3. Mesencephalon*
397
Within the mesencephalon, Brailoiu et al [18] reported a sparse number of kisspeptin-
immunoreactive fibers only in the *dorsomedial periaqueductal gray substance* (DMPAG). In our study
that region had the highest density of kisspeptin fibers within the mesencephalon (Fig. 2 J and Fig. 3 B).
However, we also visualized a small number of fibers in the *central nucleus of the inferior colliculus* (CIC,
Fig. 2 I and Fig. 3 D), *dorsal raphe nucleus* (DR, Fig. 2 J and Fig. 3, A), and sparse fibers in the *deep*
mesencephalic nucleus (DpMe), *interpeduncular nucleus* (IP), *median raphe nucleus* (MnR), *substantia*
nigra (SN), *intermediate and deep gray layer of the superior colliculus* (DpG), *supramammillary nucleus*
(Sum), *substantia nigra pars reticulata* (SNpr) and *ventral tegmental area* (VTA). The funtional
implications for those Kp projections are unknown.
407
408   *2.2.4. Metencephalon and myelencephalon*
409

In the metencephalon and myelencephalon, we observed kisspeptin immunopositive somas in the *nucleus tractus solitarius* (NTS or Sol), and *area postrema* (AP) Fig. 1 G; Fig. 2 L and Fig. 3 A-D). We did not observed immunopositive neurons in the caudoventrolateral or the lateral reticular nuclei (CVL and LRt), nor the spinal trigeminal tract, as reported in Brailoiu et al [18]. With respect to the density of the fibers, intense (++++) labelling was found in the *lateral parabrachial nucleus* (LPB, Fig. 1H, Fig. 2 K), the *Kolliker-Fuse nucleus* (KF Fig. 2 K), *NTS*, (Fig. 1 G; Fig. 2 L and Fig. 3 A-D) in the dorsomedial part of the *principal sensorial component of the trigeminal nucleus* (Pr5DM, Fig. 1 K and Fig 3 D-G) and locus coeruleus (Fig. 1 I; Fig 2 K and Fig. 3C). The rat brain also showed intense Kp labeling in the gelatinous layer and the spinal nerve 5 (Fig. 1 J-K and Fig 3 E-F). Moderate (+++) density was observed in the superior cerebral peduncle. Low (++) intensity staining was observed in several nuclei of the reticular formation, including the caudoventral reticular nucleus (CVL); the gigantocellular reticular nucleus (Gi); the intermediate reticular nucleus, the lateral reticular nucleus (LRt); the medullary dorsal reticular nucleus (MdD), the parvicellular reticular nucleus (PCRt) and the rostroventrolateral reticular nucleus (RVL). Sparse (+) fibre density was observed in the ambiguus nucleus (Amb); lateral superior olive (LSO); ventral medullary reticular nucleus (MdV); in subregions of the pontine nucleus such as the brachium pontis (bpPn) the reticular oral part (PnO) and the reticular caudal part (PnC); in the raphe magnus (RMg), obscurus (Rob), pallidus (RPa).

2.3 Gonadectomy's long term effect on kisspeptin-expressing systems in old rodents: abolition of the GNX's effect after 12 months in most brain areas

Numerous studies have reported that gonadal status exerts key feedback to KP expression with the neurons of ARC being inhibited and those in the rostral hypothalamus being potentiated by gonadal steroids [1, 41-43]. We analyzed the brains of male and female mice that underwent gonadectomy in their young adulthood (10-12 weeks) and were perfused/fixed 12 months later (15 months of age). We examined KP-ir fibers densities aiming to assess if castration early in reproductive life could modify the density of fibers in aged brain, compared with intact subjects of the same age.

Table 2 is a report of a semiquantitative analysis of 21 regions within the scope of this study. In female subjects, 19 of the 21 regions were found similar in term of density of KP-ir fibers between old intact subjects ("FC" stands for female control) and subjects undewent ovariectomy in young adulthood ("OVX" stands for ovariectomy, 12 months before the perfusion/fixation). The septohypothalamic nucleus showed increased KP-ir fiber density in OVX subjects. This region, beside of hosting *Kiss1/Slc32a1/Ar* expressing neurons [1] (Fig. 1, A and A1), received direct RP3V[KP] and Arc[KP] projections showed by Allen Mouse Brain Atlas, in two experiments using Cre-dependent AAV tracing of axonal projections from RP3V and Arc as injection sites in Kiss1-cre mice (mouse.brain-map.org/experiment/show/232311236 and mouse.brain-map.org/experiment/show/232310521). Hence, the origin of this increased fiber density is still not clear. In the RP3V and MPO region, the samples from OVX subjects also showed an increased KP immunoreactivity (Fig. 4). This region, beside of hosting a main KP neuronal populations, also receives Arc[KP] axonal projections demonstrated by the data base of Allen Institute (mouse.brain-map.org/experiment/show/232310521). In male subjects, 18 of the 21 regions found no apparent

differences between intact male subjects ("MC" stands for male control) and orchiectomized subject ("ORX" stands for orchiectomy). Three regions were found having higher densities of KP-ir fibers the ORX group than MC: dorsal hypothalamus (DH, Fig. 4), paraventricular anterior of thalamus (PVA) and tuberomammillary nuclues. Also, in these 3 regions, MC group showed higher densities than the FC group (Fig. 4).

Table 2 KP immunopositive fiber density analysis in regions related to behavioral states in old female and male mice (14-15 mo.) vs. mice 12 mo. post gonadectomy

	FC	OVX	FC vs OVX	MC	ORX	MC vs ORX	FC vs MC
AcbSh	+++	+++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC
BSTam	+++	+++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC
DB	+++	+++	FC ≈ OVX	+	++	MC ≈ ORX	FC > MC
DH	+	+	FC ≈ OVX	++	++	MC < ORX	**FC < MC**
LC	++++	++++	FC ≈ OVX	++++	++++	MC ≈ ORX	FC ≈ MC
LSD	+	+	FC ≈ OVX	+	+	MC ≈ ORX	FC ≈MC
MEA	+	+	FC ≈ OVX	+	+	MC ≈ ORX	FC ≈ MC
MS	++	++	FC ≈ OVX	+	+	MC ≈ ORX	FC > MC
NTS	++++	++++	FC ≈ OVX	++++	++++	MC ≈ ORX	FC ≈ MC
PaAP	+++	+++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC
PBN	++++	++++	FC ≈ OVX	++++	++++	MC ≈ ORX	FC ≈ MC
TM	+	+	FC ≈ OVX	++	++	MC < ORX	**FC < MC**
PMv	+	+	FC ≈ OVX	++	++	MC ≈ ORX	**FC < MC**
PVA	++	++	FC ≈ OVX	+++	++++	MC < ORX	**FC < MC**
PVN	+++	+++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC
RP3V	+++	++++	FC < OVX	++	++	MC ≈ ORX	FC > MC
AH (adj SCN)	+++	+++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC
SHy	+++	++++	FC < OVX	++	++	MC ≈ ORX	FC > MC
STIA	+	+	FC ≈ OVX	+	+	MC ≈ ORX	FC ≈ MC
Wall 3V	+++	+++	FC ≈ OVX	+	+	MC ≈ ORX	FC ≈ MC
ROIs adj. (f/st/sm/ac)	++	++	FC ≈ OVX	++	++	MC ≈ ORX	FC > MC

Table 2. Semiquantitative comparison of kisspeptin immunoreactive cell and fiber distribution in KP expressing regions of control vs gonadectomized mice. Since no differences in the direction of changes were observed in male vs female mice the data reflected here is derived from pooled observations form both sexes.   n.a.: does not apply. FC: female control (intact 15 mo. old); OVX: 12 mo. post-ovariectomy (15 mo. old); MC: male control (intact 15 mo. old); ORX: 12 mo. post-orchiectomy (15 mo. old); AcbSh, nucleus accumbens share; BSTam, bed nucleus of stria terminalis, anteromedial division; DB, nucleus diagonal band de Broca; DH, dorsal hypothalamus; LC, locus coeruleus; LSD, lateral septal nucleus, dorsal division; MEA, medial amygdala; MS, medial septal nucleus; NTS, nucleus of tract solitary; PaAP, paraventricular nucleus anterior parvocellular division of the hypothalamus; PBN, parabrachial nucleus and pons; TM, tuberomammillary nucleus of the hypothalamus; PMv, premammillary nucleus, ventral subdivision of the hypothalamus; PVA, paraventricular nucleus anterior division of the thalamus; PVN, paraventricular nucleus of the hypothalamus; RP3V, rostro-periventricular complex of third ventricle of the hypothalamus; AH anterior hypothalamic area (adjacent to suprachiasmatic nucleus, SCN); SHy, septo-hypothalamic nucleus; STIA, nucleus of stria terminalis, intra-amygdalar division; Wall 3V, ependymal layer of the 3rd ventricle; ROIs adj. (f/st/sm/ac), regions of interest adjacent to brain conducting system (fornix, f, stria terminalis, st, stria medullaris, sm, anterior commissure, ac).

Fig. 4 Examples of KP-ir fiber density in old mice: comparation of 4 groups. FC: samples from female
control, intact mice of 15-month-old; OVX: female mice underwent ovariectomy in young adulthood (3-
month-old) and perfused/fixed after 12 months after OVX); MC: male intact mice; ORX: male mice
underwent orchiectomy in young adulthood (3 month old) and perfused/fixed after 12 months after
ORX).

482
483   2.3 Mapping of *Kiss1r and its co-expression with Slc32a1, Slc17a6 and Kiss1* with RNAscope methods
484   suggests the *Kiss1r* is much more widely distributed in the CNS and Kiss1-Kiss1r signaling system can
485   function in *autocrine* and *paracrine* modes
486

The kisspeptin receptor (Kiss1R), was first identified in rat in 1999 as an orphan receptor, previously
termed Gpr54 [44]. Kiss1R is a metabotropic receptor coupled to Gq/11, and its inactivation has been
associated with hypogonadism [45]. After kisspeptin binds its receptor, phospholipase C (PLC) is activated
and hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP2) to produce diacylglicerol (DAG) and
inositol 1,4,5-trisphosphate (IP3) and the ensuing rise in Ca++ mobilized from intracellular stores and the
activation of PKC [46, 47].
493

There have been few reports on Kiss1R expression in the CNS [46, 48, 49], however, a comprehensive
description, especially the Kiss1R co-expression within glutamate-GABA context is incomplete. Using
RNAscope dual and multiplex ISH methods, we report here semi-quantitative expression levels of *Kiss1r*
in mouse brain sagittal serial sections, in figure 5, based on microscopic observation as different intensities
of blue shading. Contrasting with the discrete expression of *Kiss1 in seven neuronal populations* [1], *Kiss1r*
expression is widespread and not noticeably restricted to defined nuclei in rodent brain. *Kiss1r* was
preferentially expressed in non-GABAergic neurons (Table 3), such as hippocampal projection neurons,
main glutamatergic thalamic nuclei, mammillary complex neurons, cerebellar granule cells, and the
interposed nucleus. *Kiss1r* was also co-expressed in *Kiss1*-expressing hypothalamic neurons, suggesting
an autocrine mechanism for KP signaling. Besides, most neurons neighboring *Kiss1 expressing* cells also
expressed *Kiss1r*. These observations suggest that the KP/Kiss1R pathway may
use *autocrine* and *paracrine* mechanisms in addition to classical neurotransmission through axon
innervation and transmitter co-release, as well as through cerebrospinal fluid (Fig. 1, Panels A3 and E2).
507
508

Fig. 4 *Kiss1r* expression assessment in relation to Slc32a1, Slc17a6, Kiss1 expression suggests that the KP-Kiss1R system uses autocrine, paracrine, and neuroendocrine modes for signal transduction.
(**A–C**) Mapping of *Kiss1*(RNA encoding KP) expression (symbolized by the intensity of blue shading) in three septo-temporal planes in relation to main *Kiss1* containing brain nuclei and subfields, based on microscopic observations. Shaded regions with different blue intensity symbolize the strength of *Kiss1r*. For abbreviations see the corresponding table for abbreviations in table 2.

Kiss1r *Slc32a1*

rostral ⟷ caudal

(AOB, gl)
(AOB, mi & gr)
(CLA)
(EP)
(CP)
(ACB)
(NDB)
(LPO)
(BST)
(Rt & AM, thalamus)
(hippocampus)
(DG)
(CA1)
(CA2)
(CA3)
(SCs)
(PAG)
(PB)
(LC)

Table 2. Distribution, cell types and strength of main *Kiss1r expressing cell groups* in mouse brain with comparison of co-expression of *Slc32a1* (VGAT)

Cell group / sub-field	Co-expressing Slc17a6 (VGAT)	NO Slc32a1 (VGAT)
Cerebrum: cortical plate		
Olfactory area		
Main Olfactory bulb (MOB)		
Granular cell layer (gr)	+++	+
Inner plexiform layer (ipl)	-	-
Mitral cell layer (mi)	+	+++
Outer plexiform layer (opl)	-	+
Glomerular layer (gl)	+++	+
Accessory olfactory bulb (AOB)	n/r	+
Mitral cell layer (mi)	+	+++
Glomerular layer (gl)	++++	+
Granular layer (gr)	++++	+
Ant olfactory n. (AON)	+	++
Taenia Tecta (TT)	+	++
Piriform area: Pir2	-	+++
Piriform area: Pir3	-	++
N. lat. olfactory tract (NLOT)	-	+++
Cortical amygdalar area (CoA)	-	+++
Hippocampal formation		
dorsal dentate gyrus, granule cells layer (gcl)	+	++++
dorsal hippocampus CA1, pyramidal cell layer (pcl))	+	++++
dorsal hippocampus CA2, pcl	+	++++
dorsal hippocampus CA3, pcl	+	++++
hilus	++	++
Subiculum	++	++
Isocortex		
Layer I	+	++
Layer II-II	+	++
Layer IV	+	++
Layer V	+	++
Layer VI	+	+++
Agranular insular cortex	+	++++
Somatomotor areas		
2ry motor area, layer 2-3:	+	++
2ry motor area layer 5:	+	+++
1ry motor area, layer 2-3	+	++
1ry motor area, layer 5	+	++
Agranular insular cortex	+	+++
Orbital frontal cortex (OFC)		
OFC 1	+	++
OFC 2/3	+	+++
OFC 5	+	+++
Prefrontal Cortex (PFC)		
Ant cingulate cortex (ACC):		
ACC 2/3:	+	+++
ACC 5:	++	+++
Prelimbic (PL)		
PL 2/3	+	+++
PL 5	++	+++
Infralimbic (IL)		
IL 2/3	++	+++
IL 5	++	+++
Prim somatosensory a. SSp,		
SSp 1	+	++
SSp 2/3	+	++
SSp 4 (mouth)	+	++
SSp 5	+	++
SSp 6a	+	+++
Gustatory areas (GU)	+	++
Auditory area	+	++
Visual area	+	++
Visceral area (VISC)	+	++
Temporal association area	+	+++
Ectorhinal area	+	+++
Perirhinal area	+	++
Retrosplenial area	+	+++
Post parietal association area	+	+++
Cortical subplate		
Claustrum	+	++
Endopiriform nucleus	+	++
Lateral amygdalar nucleus (LA)	+	+++
Post amygdalar nucleus (PA)	+	+++
Basomedial amygdala (BMA)	+	+++
Basolateral amygdala (BLA)	+	+++
Cerebral nuclei		
Striatum		
Caudado putamen	+++	+
Lateral septal nucleus (LS)	++	++
Septalhypothalamic nucleus (SHy)	+++	+++
Anterior amygdala area (AAA)	++	+
Central amygdalar nucleus (CEA)	++	+
Medial amygdalar nucleus (MEA)	++	+
Pallidum (basal forebrain)		
Medial Septal nucleus (MS)	++	++
Bed nucleus of stria terminalis (BNST)	++	++
Nucleus basalis of Meyner	+	+++
Globus pallidus	+	++
Substantia innominata (SI)	+	++
Nucleus of diagonal band of Broca (NDB)	+	++
Nucleus Accumbens	++	+
Brain stem, inter-brain		
Thalamus		
Somato-motor related		
Subparafacicular nucleus, magnocellular part	-	+++
Subparafacicular area	-	+++
Peripeduncular nucleus	-	+++
Medial geniculate complex	-	+++
Polymodal association cortex related		
Lat. posterior n. thal	-	+++
Post. limiting nucleus	-	+++
Suprageniculate n.	-	+++
Anterodorsal n.	-	+++
Anteromedial n.	-	++
Parataenial n.	-	++
Intermedial n.	-	+++
Laterodorsal n.	-	+++
Centrolateral n	-	+++
Intermediodorsal n.	-	+++
Mediodorsal n.	-	+++
Pariventricular n.	-	+++
Parateanial n.	-	+++
N. of reuniens[n2]	-	+++
Posterior pretectal n.	-	+++
Precommissural n.	-	+++
Reticular nucleus of the thalamus (Rt)	+++	-
Epithalamus		
Medial habenula	-	++++
Lateral habenula	+	++++
Hypothalamus		
Paraventricular n (PVN)	+	++
Arcuate Nucleus	+++	-
Suprachiasmatic nucleus	+++	+
Rostro-periventricular of the 3V (RP3V)	+++	++
Anterodorsal preoptic n.	++	++
Anteroventral n.	+++	++
Dorsomedial n.	+++	++
Median preoptic n.	++	++
Medial preoptic area	++	++
Vascular organ of lamina terminalis	-	-
Lateral preoptic area (LPO)	++	+
Anterior hyp. area (AH)	+++	+
Premammilary n. (PM)	-	+++++
Lateral mammillary n.	-	+++++
Medial mammillary n. (MM)	-	+++++
Supramammilary n.(SUM)	++	++
Median preoptic n. (MnPO)	+	+++
Lateral hyp. area (LH)	++	++
Subthalamic nucleus (STN)	-	+++++
Retrochiasmatic area	+++	++

Region		
Tuberomammillary nucleus (TM, magnocellular subnucleus MTu)	+++	+++
Zona incerta (ZI)	++	+
Ventromedial hyp. n (VMH)	-	+++++
Post. hypothalamic n.	+	+
Midbrain		
Superior colliculus, sensorial (SCS)	+	+++
Superior colliculus, motor (SCM)	+	++
Inf. Colliculus (IC)	++	++
N. of the brachium of IC (BIC)	+	+
Ventral tegmental area (VTA)	+	+++
Sustantia nigra reticulata	+	+++
Midbrain reticular n.	+	++
Periaqueductal gray	++	+++
Midbrain raphe nuclei	+	+++
Pedunculopontine n.	+	++
Dorsal n. raphe	++	++
Pons		
N. lateral lemniscus	+	+
Principal sensorial nucleus of trigeminal nerve	-	++
Koelliker-Fuse subnucleus	-	++
Parabrachial n. lateral div.	-	+++
Superior olivary comp (lat)	-	++
Tegmental reticular n.	++	+++
Dorsal tegmental n.	++	+++
Pontine gray	-	++++
Pontine central gray	++	+++
Supratrigeminal nucleus	++	++
Locus Coeruleus (state)	-	++++
Laterodorsal tegmental n.	++	+++
Pontine reticular n.	++	+++
Superior central n. raphe	++	++
Medulla		
N. tractus solitarii medial	+	+++
N. tractus solitarii lateral	+	+++
Hypoglossal (XII) n.	+	++
Dorsal motor n. of the Vagus nerve (X)	+	+++
Dorsal cochlear n.	-	++
Ventral cochlear n.	-	++
Spinal n. trigeminal	+	++
N. prepositus	-	+++
Facial motor n. (VII)	++	++
N. ambiguus	++	+++
Magnocellular reticular n.	++	+++
Spinal vestibular n.	+++	++
N. X	+	++
N. raphe magnus (state related)	+	+++
N. raphe pallidus (state related)	+	+++
N. raphe obscurus (state rel.)	+	+++
Cuneate n.	+	+
Inferior olivary	+	+++
Cerebellar cortex		
Purkinje's cells	-	-
Golgi's cells	+	-
Granule cells[n4]	-	++++
Cerebellar nuclei		
Interposed n.	+	+
Dentate n.	-	++
Fastigial n.	+	++

* Semiquantitative annotations are used here (the percentage of expressing cell/total Nissl stained nuclei: "-", not detectable; "+", weak (<20%); "++", low (40%-20%); "+++", moderate (60%-40%); "++++", intense (80%-60%); "+++++", very intense (>80%).

** Functional neuroanatomy order and annotations are based on Allen Institute Mouse Reference Atlas.

3. General discussion and conclusions

In this study, we provide a comprehensive description of the distribution of kisspeptin (KP) fibers and cell bodies throughout the brain, as well as the distribution of the Kiss1 receptor (Kiss1R) using GABAergic neuronal groups as markers for regional chemoarchitecture. Our findings were obtained using immunohistochemistry with a highly specific antibody and an improved method that maximized the visualization of KP immunoreactivity. Additionally, we employed a sensitive dual in situ hybridization technique (RNAscope dual and multiplex) to further delineate the expression patterns. Our immunohistochemical analysis revealed a stronger and more extensive distribution of KP immunoreactivity throughout the mouse and rat brains than previously reported.

The distribution patterns suggest that many hypothalamic and extrahypothalamic regions are as strongly innervated as the classical hypothalamic regions known for regulating the hypothalamic-pituitary-gonadal (HPG) axis, but likely serve non-reproductive functions. Indeed, many of these regions are critical for regulating brain states, including telencephalic cortical areas, septal regions (SHy, MS, LS), diencephalic structures (MnPO, SCh, BST, NBM), midbrain regions (PAG, DR, SN/VTA), and metencephalic areas such as the raphe, reticular formation, and LC.

Another prominent finding was the profuse innervation of KP-immunoreactive fibers in regions involved in sensory processing, including the main and accessory olfactory bulbs, olfactory cortical regions, superior colliculus, solitary tract, trigeminal sensory regions, and several thalamic nuclei. A highlight of this study is the analysis of brains from male and female mice that underwent gonadectomy

at 3 months of age, with tissue collection occurring one year post-surgery. Notably, the KP fiber density in 21 regions involved in non-reproductive functions was similar or even increased in gonadectomized animals compared to controls, suggesting that the KP system remains functional in other processes beyond reproductive senescence.

Regarding the distribution of Kiss1r, we observed its preferential expression in non-GABAergic cells, including hippocampal projection neurons, major glutamatergic thalamic nuclei, mammillary complex neurons, cerebellar granule cells, and the interposed nucleus. Additionally, Kiss1r was expressed in Kiss1-expressing hypothalamic cells, indicating a potential autocrine/paracrine role for the peptide.

The widespread expression of Kiss1r in regions associated with sensory and behavioral state functions, along with the presence of kisspeptin-immunopositive fibers beyond the hypothalamus, suggests a broader and less understood role for kisspeptin signaling in the brain. Recent technological advancements in molecular mapping and imaging have enabled a more systematic examination of Kiss1r distribution and its interaction with kisspeptin fibers throughout the brain. This study aims to bridge this knowledge gap by providing a comprehensive map of Kiss1r expression and kisspeptin fiber distribution in the mouse brain, offering insights into its potential functions within non-reproductive neural circuits.

We have reported in this study that a high density of KP fibers is present in areas related to pain sensory processing, such as the principal sensorial trigeminal nucleus, the gelatinous layer of the trigeminus and in the periaqueductal gray. In the literature, only a few studies have investigated Kp's role in the modulating of the perception of exteroceptive or interoceptive non-reproductive functions. It has been shown that KP-10 activates calcium signaling pathways in cultured dorsal root ganglion cells, suggesting a possible participation in pain modulation [50]. Other regions classically involved in sensory processing have moderate/dense innervation of KP fibers, *i.e.,* superior colliculus for visual processing; the inferior colliculus and superior olive for audition; the parabrachial nucleus and nucleus of the solitary tract (NTS or Sol) for gustatory processing; and the olfactory tubercle, nucleus of the olfactory tract and medial amygdala for olfaction. To our knowledge, the effect of KP on the activity of these regions and its effect on sensory integration has not been studied.

3. Material and methods

3.1. Animals

Wistar rats (N=12, male = 6, female = 6) and C57BL/6 mice (N=20, male = 10, female = 10) were obtained from the local animal vivarium and housed three per cage under controlled temperature and illumination (12h/12h) with water and food *ad-libitum*. Mice subjected to gonadectomy were singly housed for recovery for one week after surgery and then returned with their cage mates. All animal procedures were approved by the local research ethics supervision commission (license number CIEFM-079-2020). We analysed two brains sectioned in the parasagittal plane and two brains sectioned in the coronal plane for each sex and species. To evaluate the effect of gonadectomy on the expression of

607 kisspeptin through the brain of male and female mice, we compared the immunoactivity of 4 control vs.
608 4 gonadectomised (one-year post-surgery) mice.
609

610 3.2. Gonadectomy
611

612 *3.2.1 Mouse ovariectomy*
613

A dorsal approach to remove the ovaries was used [51]. Briefly, animals were anaesthetised with
ketamine 90mg/kg + xylazine 20mg/kg i.p. under anaesthesia animals were placed in ventral recumbency
and a 3cm/side squared region centred over the third lumbar vertebra shaved. The shaved area was
disinfected with Povidone and 70% alcohol, and a sterile fenestrate drape was placed over it. A single
midline dorsal incision approx. 1 cm long was made in the skin at the level of the bottom of the rib cage.
The incision was moved laterally to cut the right and left abdominal walls. After cutting the wall (obliques
and transverse abdominal muscles), blunt dissection through the body wall with mosquito haemostats was
performed. Forceps were gently and carefully introduced to retrieve and exteriorise the corresponding
ovary. The ovary, ovarian vessels, and ligament were ligated with 4-0 synthetic absorbable sutures and
cut. The wall muscle layers incision was closed with a 4-0 synthetic, monofilament, absorbable suture.
The skin incision was closed with a subcuticular suture.
625

2.2.2 Mouse orchiectomy

For the excision of the testicles, we followed a midline incision technique [52]. Briefly, animals
were anaesthetised with ketamine 75mg/kg + xylazine 16mg/Kg i.p. and placed on the operating table on
their back. The central ventral side of the scrotum area was disinfected with Povidone and 70% alcohol.
A single incision on the ventral side of the scrotum (1cm long) penetrating the skin was performed. Each
cremaster muscle was cut, and each testis was exposed by gently pulling it through the incision using blunt
forceps. Cauda of the epididymis, caput epididymis, vas deferens, and testicular blood vessels were
exposed while holding the testicular sack with sterile tooth forceps. A single ligature 4-0 synthetic
absorbable suture around the vas deferens and blood vessels was made, prior to excision and gentle
removal of the testis. The remaining content was replaced in each one of the testicular sacks. The skin
incision was closed with a subcuticular suture.
637

638 3.3 Immunohistochemistry
639

Animals were deeply anaesthetised with *i.p.* injected sodium pentobarbital (100mg/kg). They were
perfused transaortically with 0.9% saline followed by a cold fixative containing 4% paraformaldehyde in
0.1 M sodium phosphate buffer (PB, pH 7.4) plus 15% v/v saturated picric acid for 15 min. Brain tissues
were rinsed immediately after the 15 min fixation lapse until the PB yellowish colour was fully cleaned.
Brains were then sectioned the same day, using a Leica VT 1000S vibratome at 70μm thickness in the
coronal and parasagittal planes and the immunohistochemical (IHC) reaction was performed the same day
of the perfusion-fixation. This procedure yielded a more comprehensive IHC labelling than longer fixation
and delayed IHC as previously reported and discussed [16]. One of every three sections was selected, and

sections from animals that would be compared (*i.e.*, female control *vs.* female gonadectomy) were incubated in the same vial. To block the unspecific antibody binding, sections were incubated for one hour in a solution consisting of 20% normal donkey serum (NDS) diluted in tris-buffered saline (TBS, pH 7.4) plus 0.3% Triton-X (TBST). After the blocking step, sections were incubated for 48h at 4°C in a polyclonal sheep anti-kisspeptin antibody [15] (GQ2, kindly supplied by Mohammad Ghatei†, Imperial College, London, UK), diluted at a 1:2000 concentration in TBST + 1% NDS. This antibody recognises kisspeptins 54, 14,13, and 10 that conserve the C-terminal decapeptide essential for biological activity and shows virtually no cross-reactivity (<0.01%) with other related human RF-amide peptides [15]. After three wash steps, sections were incubated for 24 hours with the secondary donkey anti-sheep biotinylated antibody (Cat: 713-065-147, Jackson ImmunoResearch Inc, PA, USA) diluted at a concentration of 1:500 in TBST + 1% NDS. To detect the biotinylated antibody, VECTASTAIN Elite ABC HRP Kit (Cat: PK-6100, Vector Laboratories, Inc. CA, USA) was used, with H_2O_2 and 3,3'-Diaminobenzidine as substrates. Finally, sections were mounted on glass slides, dehydrated, dipped in xylene, and cover-slipped using Permount^{TM} mounting medium.

3.4 Anatomical assessment of kisspeptin innervation density and comparison between control and gonadectomised animals

To assess the projection patterns of kisspeptin immunopositive fibers through the brain three researchers independently examined the slides under a light microscope, after identifying the regions based on the Paxinos Rat & Mouse brain atlases. Scores of "+" (sparse, main axons passing, without branching nor axon terminals); "++", mild, some branching; "+++", moderate, abundant branching and axon terminals; "++++", intense, extensive branching and intense axon terminals, with confluent fibers that it was difficult to see a separation between them, were reported.

3.5 Anatomical assessment of mRNA expression in the context of Slc32a1 expressing neurons through the brain.

The detailed methods have been previously described (Zhang, Hernandez et al., 2018; 2020; 2021). In summary, mice were deeply anesthetized with sodium pentobarbital (Sedalpharma, Mexico, 100 mg/kg b.w., i.p.) and decapitated using a small animal guillotine (Kent Scientific Corp.). The brains were then extracted and rapidly frozen in dry ice. Fresh-frozen brains were sectioned sagittally into 12 μm thick slices using a Leica CM-1520 cryostat and mounted onto positively charged Superfrost Plus microscope slides (Cat. 22-037-246, Fisher Scientific, Pittsburgh, PA).

Probes used for the detection of mRNAs in the double in-situ hybridization (DISH) were custom-designed by Advanced Cell Diagnostics (https://acdbio.com/rnascope-25-hd-duplex-assay) for kisspeptin receptor (Mm-Kiss1r, channel 2 probe), kisspeptin (Mm-Kiss1, channel 1 probe) and VGAT (Mm-Slc32a, channel 2 probe), All experimental procedures were carried out in accordance with the manufacturer's protocol.

After sectioning, the slides were fixed in chilled 4% paraformaldehyde (Cat. 30525-89-4, Sigma-Aldrich)
prepared in phosphate-buffered saline (Cat. M32631, Sigma-Aldrich), dehydrated in ethanol, and treated
with hydrogen peroxide for 15 minutes. They were then treated with protease III for 20 minutes at room
temperature, incubated for two hours with the channel 1 and 2 probe mixtures, and underwent
amplification and detection steps following the manufacturer's guidelines. Finally, the slides were
counterstained with hematoxylin and cover slipped.
694
695   Conceptualization: LZ, VSH, MAZ, RPM, LEE; data curation: LZ, VSH, MAZ, ORHP; investigation: LZ, VSH,
696   MAZ, ORHP, RHG; methodology: LZ, VSH, RHG; project administration: LZ, VH, LEE; resources: LZ, LEE,
697   RPM, ICA, RHG; visualization LZ, VSH; writing – original draft: LZ, VH, LEE; review & editing: all the co-
698   authors.
699

700   Funding and fellowship acknowledgement
701
The research leading to these results received funding from the National Autonomous University of Mexico, under
Grant Agreement UNAM-PAPIIT-IG200121, Mexican Nacional Council for Humanity, Science and Technology
(CONAHCYT), under Grant Agreement CF-2023-243 (LZ), NIMH-IR, NIH, under Grant Agreement MH002386
(LEE). The investigators of this study have been supported by the following fellowships: sabbatical year fellowship
from the PASPA program of the Dirección General de Personal Académico (DGAPA) of the Universidad Nacional
Autónoma de México (UNAM) (LZ, VSH); Mexican CONAHCYT sabbatical fellowship (LZ) and Fulbright-
García Robles Fellowship (VSH), for sabbatical research year in LEE's laboratory of NIMH, NIH; DGAPA-UNAM
PREI program for a sabbatical research stay (RPM) in LZ's lab in UNAM, Mexico.
710

711
References
713

1. Hernández VS, Zetter MA, Hernández-Pérez O, Hernández-González R, Camacho-Arroyo I, Millar
RP, Eiden LE, Zhang L. A comprehensive chemotyping and gonadal regulation of seven kisspeptinergic
neuronal populations in the mouse brain. *bioRxiv*. 20242024.07.23.604881.
2. Lee JH, Miele ME, Hicks DJ, Phillips KK, Trent JM, Weissman BE, Welch DR. KiSS-1, a novel
human malignant melanoma metastasis-suppressor gene. *J Natl Cancer Inst*. 1996; **88**(23): 1731-7.
3. Pinilla L, Aguilar E, Dieguez C, Millar RP, Tena-Sempere M. Kisspeptins and reproduction:
physiological roles and regulatory mechanisms. *Physiol Rev*. 2012; **92**(3): 1235-316.
4. Clarke H, Dhillo WS, Jayasena CN. Comprehensive Review on Kisspeptin and Its Role in
Reproductive Disorders. *Endocrinol Metab (Seoul)*. 2015; **30**(2): 124-41.
5. Mills EGA, O'Byrne KT, Comninos AN. Kisspeptin as a Behavioral Hormone. *Semin Reprod Med*.
2019; **37**(2): 56-63.
6. Stephens SBZ, Kauffman AS. Regulation and Possible Functions of Kisspeptin in the Medial
Amygdala. *Front Endocrinol (Lausanne)*. 2017; **8**191.
7. Liu X, Herbison AE. Kisspeptin Regulation of Neuronal Activity throughout the Central Nervous
System. *Endocrinol Metab (Seoul)*. 2016; **31**(2): 193-205.
8. Ebrahimi Khonacha S, Janahmadi M, Motamedi F. Kisspeptin-13 Improves Spatial Memory
Consolidation and Retrieval against Amyloid-beta Pathology. *Iran J Pharm Res*. 2019; **18**(Suppl1): 169-
81.
9. Abdul Satar NM, Ogawa S, Parhar IS. Kisspeptin-1 regulates forebrain dopaminergic neurons in
the zebrafish. *Sci Rep*. 2020; **10**(1): 19361.
10. Sivalingam M, Ogawa S, Parhar IS. Habenula kisspeptin retrieves morphine impaired fear
memory in zebrafish. *Sci Rep*. 2020; **10**(1): 19569.
11. Clarkson J, d'Anglemont de Tassigny X, Colledge WH, Caraty A, Herbison AE. Distribution of
kisspeptin neurones in the adult female mouse brain. *J Neuroendocrinol*. 2009; **21**(8): 673-82.
12. Lehman MN, Hileman SM, Goodman RL. Neuroanatomy of the kisspeptin signaling system in
mammals: comparative and developmental aspects. *Advances in experimental medicine and biology*.
2013; **784**27-62.
13. Mikkelsen JD, Simonneaux V. The neuroanatomy of the kisspeptin system in the mammalian
brain. *Peptides*. 2009; **30**(1): 26-33.
14. Lehman MN, Coolen LM, Steiner RA, Neal-Perry G, Wang L, Moenter SM, Moore AM, Goodman
RL, Hwa-Yeo S, Padilla SL, Kauffman AS, Garcia J, Kelly MJ, Clarkson J, Radovick S, Babwah AV, Leon S,
Tena-Sempere M, Comninos A, Seminara S, Dhillo WS, Levine J, Terasawa E, Negron A, Herbison AE.
The 3(rd) World Conference on Kisspeptin, "Kisspeptin 2017: Brain and Beyond":Unresolved questions,
challenges and future directions for the field. *J Neuroendocrinol*. 2018e12600.
15. Dhillo WS, Chaudhri OB, Patterson M, Thompson EL, Murphy KG, Badman MK, McGowan BM,
Amber V, Patel S, Ghatei MA, Bloom SR. Kisspeptin-54 stimulates the hypothalamic-pituitary gonadal
axis in human males. *J Clin Endocrinol Metab*. 2005; **90**(12): 6609-15.
16. Zhang L, Hernandez VS. Synaptic innervation to rat hippocampus by vasopressin-immuno-
positive fibres from the hypothalamic supraoptic and paraventricular nuclei. *Neuroscience*. 2013;
228139-62.

17.   Swanson LW. *Brain Architecture: Understanding the Basic Plan* Oxford, UK: Oxford University
Press, 2011.
18.   Brailoiu GC, Dun SL, Ohsawa M, Yin D, Yang J, Chang JK, Brailoiu E, Dun NJ. KiSS-1 expression
and metastin-like immunoreactivity in the rat brain. *J Comp Neurol*. 2005; **481**(3): 314-29.
19.   Paxinos G, Franklin KBJ. *Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates*
Elsevier Science, 2012.
20.   Risold PY, Swanson LW. Connections of the rat lateral septal complex. *Brain research Brain*
research reviews. 1997; **24**(2-3): 115-95.
21.   Rizzi-Wise CA, Wang DV. Putting Together Pieces of the Lateral Septum: Multifaceted Functions
and Its Neural Pathways. *eNeuro*. 2021; **8**(6).
22.   Comninos AN, Wall MB, Demetriou L, Shah AJ, Clarke SA, Narayanaswamy S, Nesbitt A, Izzi-
Engbeaya C, Prague JK, Abbara A, Ratnasabapathy R, Salem V, Nijher GM, Jayasena CN, Tanner M,
Bassett P, Mehta A, Rabiner EA, Honigsperger C, Silva MR, Brandtzaeg OK, Lundanes E, Wilson SR,
Brown RC, Thomas SA, Bloom SR, Dhillo WS. Kisspeptin modulates sexual and emotional brain
processing in humans. *The Journal of clinical investigation*. 2017; **127**(2): 709-19.
23.   Sheehan TP, Chambers RA, Russell DS. Regulation of affect by the lateral septum: implications
for neuropsychiatry. *Brain research Brain research reviews*. 2004; **46**(1): 71-117.
24.   Salib M, Joshi A, Katona L, Howarth M, Micklem BR, Somogyi P, Viney TJ. GABAergic Medial
Septal Neurons with Low-Rhythmic Firing Innervating the Dentate Gyrus and Hippocampal Area CA3. *J*
Neurosci. 2019; **39**(23): 4527-49.
25.   Unal G, Crump MG, Viney TJ, Eltes T, Katona L, Klausberger T, Somogyi P. Spatio-temporal
specialization of GABAergic septo-hippocampal neurons for rhythmic network activity. *Brain Struct*
Funct. 2018; **223**(5): 2409-32.
26.   Espinosa N, Caneo M, Alonso A, Moran C, Fuentealba P. Optogenetic stimulation of septal
somatostatin neurons disrupts locomotory behavior and regulates hippocampus cholinergic theta
oscillations. *bioRxiv*. 20222022.09.30.510330.
27.   Shen L, Zhang GW, Tao C, Seo MB, Zhang NK, Huang JJ, Zhang LI, Tao HW. A bottom-up reward
pathway mediated by somatostatin neurons in the medial septum complex underlying appetitive
learning. *Nat Commun*. 2022; **13**(1): 1194.
28.   Srividya R, Mallick HN, Kumar VM. The changes in thermal preference, sleep-wakefulness, body
temperature and locomotor activity in the rats with medial septal lesion. *Behav Brain Res*. 2005;
164(2): 147-55.
29.   Tsanov M. Speed and Oscillations: Medial Septum Integration of Attention and Navigation.
Frontiers in systems neuroscience. 2017; **11**67.
30.   Joshi A, Salib M, Viney TJ, Dupret D, Somogyi P. Behavior-Dependent Activity and Synaptic
Organization of Septo-hippocampal GABAergic Neurons Selectively Targeting the Hippocampal CA3
Area. *Neuron*. 2017; **96**(6): 1342-57 e5.
31.   Liu AKL, Lim EJ, Ahmed I, Chang RC, Pearce RKB, Gentleman SM. Review: Revisiting the human
cholinergic nucleus of the diagonal band of Broca. *Neuropathol Appl Neurobiol*. 2018; **44**(7): 647-62.
32.   Dudek SM, Alexander GM, Farris S. Rediscovering area CA2: unique properties and functions.
Nature reviews Neuroscience. 2016; **17**(2): 89-102.
33.   Semba K, Fibiger HC. Organization of central cholinergic systems. *Prog Brain Res*. 1989; **79**37-
63.

34.   Floresco SB. The nucleus accumbens: an interface between cognition, emotion, and action.
Annu Rev Psychol. 2015; **66**25-52.
35.   Riaz S, Puveendrakumaran P, Khan D, Yoon S, Hamel L, Ito R. Prelimbic and infralimbic cortical
inactivations attenuate contextually driven discriminative responding for reward. *Sci Rep*. 2019; **9**(1):
3982.
36.   Capuzzo G, Floresco SB. Prelimbic and Infralimbic Prefrontal Regulation of Active and Inhibitory
Avoidance and Reward-Seeking. *J Neurosci*. 2020; **40**(24): 4773-87.
37.   de Lima MAX, Baldo MVC, Oliveira FA, Canteras NS. The anterior cingulate cortex and its role in
controlling contextual fear memory to predatory threats. *Elife*. 2022; **11**.
38.   Kataoka N, Shima Y, Nakajima K, Nakamura K. A central master driver of psychosocial stress
responses in the rat. *Science*. 2020; **367**(6482): 1105-12.
39.   Gresham R, Li S, Adekunbi DA, Hu M, Li XF, O'Byrne KT. Kisspeptin in the medial amygdala and
sexual behavior in male rats. *Neurosci Lett*. 2016; **627**13-7.
40.   Pineda R, Plaisier F, Millar RP, Ludwig M. Amygdala Kisspeptin Neurons: Putative Mediators of
Olfactory Control of the Gonadotropic Axis. *Neuroendocrinology*. 2017; **104**(3): 223-38.
41.   Starrett JR, Moenter SM. Hypothalamic kisspeptin neurons as potential mediators of estradiol
negative and positive feedback. *Peptides*. 2023; **163**170963.
42.   Rance NE. Menopause and the human hypothalamus: evidence for the role of
kisspeptin/neurokinin B neurons in the regulation of estrogen negative feedback. *Peptides*. 2009;
30(1): 111-22.
43.   Dubois SL, Acosta-Martinez M, DeJoseph MR, Wolfe A, Radovick S, Boehm U, Urban JH, Levine
JE. Positive, but not negative feedback actions of estradiol in adult female mice require estrogen
receptor alpha in kisspeptin neurons. *Endocrinology*. 2015; **156**(3): 1111-20.
44.   Lee DK, Nguyen T, O'Neill GP, Cheng R, Liu Y, Howard AD, Coulombe N, Tan CP, Tang-Nguyen
AT, George SR, O'Dowd BF. Discovery of a receptor related to the galanin receptors. *FEBS Lett*. 1999;
446(1): 103-7.
45.   Franssen D, Tena-Sempere M. The kisspeptin receptor: A key G-protein-coupled receptor in the
control of the reproductive axis. *Best Pract Res Clin Endocrinol Metab*. 2018; **32**(2): 107-23.
46.   Irwig MS, Fraley GS, Smith JT, Acohido BV, Popa SM, Cunningham MJ, Gottsch ML, Clifton DK,
Steiner RA. Kisspeptin activation of gonadotropin releasing hormone neurons and regulation of KiSS-1
mRNA in the male rat. *Neuroendocrinology*. 2004; **80**(4): 264-72.
47.   Ringel MD, Hardy E, Bernet VJ, Burch HB, Schuppert F, Burman KD, Saji M. Metastin receptor is
overexpressed in papillary thyroid cancer and activates MAP kinase in thyroid cancer cells. *J Clin
Endocrinol Metab*. 2002; **87**(5): 2399.
48.   Herbison AE, de Tassigny X, Doran J, Colledge WH. Distribution and postnatal development of
Gpr54 gene expression in mouse brain and gonadotropin-releasing hormone neurons. *Endocrinology*.
2010; **151**(1): 312-21.
49.   Higo S, Honda S, Iijima N, Ozawa H. Mapping of Kisspeptin Receptor mRNA in the Whole Rat
Brain and its Co-Localisation with Oxytocin in the Paraventricular Nucleus. *J Neuroendocrinol*. 2016;
28(4).
50.   Kelestimur H, Bulut F, Canpolat S, Ozcan M, Ayar A. Kisspeptin leads to calcium signaling in
cultured rat dorsal root ganglion neurons. *Gen Physiol Biophys*. 2021; **40**(2): 155-60.
51.   Stout Steele M, Bennett RA. Clinical Technique: Dorsal Ovariectomy in Rodents. *Journal of
Exotic Pet Medicine*. 2011; **20**(3): 222-6.

52.   Sophocleous A, Idris AI. Ovariectomy/Orchiectomy in Rodents. In: Idris AI, ed. *Bone Research*
Protocols. New York, NY: Springer New York 2019: 261-7.
1.