Bacteria Are a Major Determinant of Orsay Virus Transmission and Infection in *Caenorhabditis elegans*

Brian G. Vassallo[1,2], Noémie Scheidel[1], Sylvia E. J. Fischer[1], and Dennis H. Kim[1,*]

1. Division of Infectious Diseases, Department of Pediatrics, Boston Children's Hospital and Harvard Medical School; Boston, 02115, USA
2. Department of Biology, Massachusetts Institute of Technology; Cambridge, 02139, USA

*Corresponding author. Email: dennis.kim@childrens.harvard.edu

Abstract

The microbiota is a key determinant of the physiology and antiviral immunity of animal hosts.

The factors governing the transmissibility of viruses between susceptible hosts are incompletely

understood. Bacteria serve as food for *Caenorhabditis elegans* and represent an integral part of

the natural environment of *C. elegans.* We determined the effects of bacteria isolated with *C.*

elegans from its natural environment on the transmission of Orsay virus in *C. elegans* using

quantitative virus transmission and host susceptibility assays. We observed that *Ochrobactrum*

species promoted Orsay virus transmission, whereas *Pseudomonas lurida* MYb11 attenuated

virus transmission relative to the standard laboratory bacterial food *Escherichia coli* OP50. We

found that pathogenic *Pseudomonas aeruginosa* strains PA01 and PA14 further attenuated virus

transmission. We determined that the amount of Orsay virus required to infect 50% of a *C.*

elegans population on *P. lurida* MYb11 compared with *Ochrobactrum vermis* MYb71 was

dramatically increased, over three orders of magnitude. Host susceptibility was attenuated even

further in presence of *P. aeruginosa* PA14. Genetic analysis of the determinants of *P. aeruginosa*

required for attenuation of *C. elegans* susceptibility to Orsay virus infection revealed a role for

regulators of quorum sensing. Our data suggest that distinct constituents of the *C. elegans*

microbiota and potential pathogens can have widely divergent effects on Orsay virus

transmission, such that associated bacteria can effectively determine host susceptibility versus

resistance to viral infection. Our study provides quantitative evidence for a critical role for

tripartite host-virus-bacteria interactions in determining the transmissibility of viruses among

susceptible hosts.

Introduction

Viruses are ubiquitous and abundant[1,2]. Infection can have profound consequences for the health of an individual host. The ability of viruses to transmit from one individual to another can scale these consequences causing morbidity and mortality throughout whole populations. Many factors influence virus transmission[3–5]. Abiotic factors such as temperature[6] and humidity[7] and biotic factors such as viral load[8,9] and host immune status[10,11] all interact to determine transmission rates. Despite these findings, the determinants of virus transmissibility remain incompletely understood.

The microbiota has emerged as a host-associated factor that modulates multiple aspects of virus infection and thereby alters transmission rates among host organisms[12–14]. In general, the microbiota is critical for the proper development of the immune system and for the effective activation of antimicrobial immune responses even at sites distal to microbiota colonization[15]. More specifically, bacteria and bacterial surface structures such as lipopolysaccharide and peptidoglycan have been shown to stabilize poliovirus and reovirus *in vitro*[13,16]. These observations likely explain why microbiota depletion by antibiotic treatment was sufficient to provide protection against the same viruses in mice[13]. The bacterial symbiont *Wolbachia* protects numerous insect species from multiple viruses either by upregulating antiviral defenses or competing for intracellular nutrients[17–21]. Individual bacteria have also been found to enhance viral infection; *Serratia marcescens* promoted infection of the mosquito *Aedes aegypti* by Dengue, Zika, and Sindbis viruses by secreting a protein, enhancin, that degrades the mucus layer covering epithelial cells[14].

Caenorhabditis elegans is a nematode often found in microbially rich environments such

as rotting vegetation[22]. The first naturally occurring virus capable of infecting the *C. elegans* was

isolated from Orsay, France[23]. Orsay virus is a part of a group of nematode infecting viruses

closely related to the *Nodaviridae* family of viruses which infect arthropods and fish[23,24]. Like

other Nodaviruses, Orsay virus is a positive-sense, single-stranded RNA virus with a bipartite

genome[23]. Transmission of Orsay virus occurs horizontally through the fecal-oral route.

Fluorescence *in situ* hybridization and immunofluorescence imaging has revealed that Orsay virus

solely infects *C. elegans* intestinal cells[23,25]. Viral infection activates host defense mechanisms

including the RNA-interference response and a transcriptional program known as the

intracellular pathogen response[23,26–29].

C. elegans is a bacterivore and is propagated in the laboratory on lawns of *Escherichia coli*

OP50. *Pseudomonas aeruginosa,* an opportunistic pathogen of humans, is found in the soil and

water and can also infect *C. elegans*[30–32]. Infection of *C. elegans* with *P. aeruginosa* activates

innate immunity and stress responses, as well as behavioral avoidance responses[33–35]. Recently,

the bacteria resident in the natural environment of *C. elegans* in the wild have been of increasing

interest[22,36–38]. The intestinal lumen of free-dwelling *C. elegans* is occupied by taxonomically and

functionally diverse bacteria that can affect *C. elegans* fitness and physiology[36,37,39].

In this study, we sought to understand how bacteria that are constituents of the *C.*

elegans microbiota quantitatively affect the transmission of Orsay virus in *C. elegans.* We

observed that monoaxenic cultures of different bacterial species had widely divergent effects on

the transmission of Orsay virus and conducted genetic analysis of the bacterial determinants

involved in modulating virus transmission. Our data point to a key species-specific role for

bacteria as critical determinant of virus transmission.

Results
Wide variation in the effects of bacteria on the transmission of Orsay virus.

Orsay virus spreads horizontally through the fecal-oral route and transmission can be

observed spreading from a single animal to a population of animals on a plate[40]. We sought to

assess the impact of bacteria on Orsay virus transmission rates and utilized a collection of

bacteria isolated from the environment with wild *C. elegans* to assemble a panel of Gram-

negative bacteria for comparison with the standard laboratory bacterial food, *Escherichia coli*

OP50[36,37,39,41]. We set up a transmission assay by placing infected animals ("spreaders") together

with uninfected animals on a monoaxenic lawn of each bacteria and examining the incidence

proportion, or, the number of new infections produced after 24 h as determined by detection of

induction of the *pals-5p::GFP* reporter, which is induced by infection with Orsay virus (Fig. 1A-

C)[27]. We observed a wide range in the measured incidence proportion (Fig. 1B and 1C). Exposure

to many of the naturally-associated bacterial strains resulted in transmission that was

comparable to the incidence proportion observed with *E. coli* OP50 with some prominent

exceptions (Fig. 1B and 1C). Two *Ochrobactrum* species promoted virus infection in nearly all

individuals in the transmission assay and increased the incidence proportion 2.7-fold and 2.9-fold

compared to *E. coli* OP50 (Fig. 1B and 1C). We confirmed that transmission from the initial

spreader animals in the assay, and not multiple rounds of infection, was responsible for the

increased incidence proportion observed in the presence of *Ochrobactrum vermis* MYb71 (Supp.

Fig. 1A and 1B). On the other hand, the presence of *P. lurida* MYb11 reduced the incidence

proportion 4.1-fold compared to *E. coli* OP50 and 11-fold compared to *O. vermis* MYb71 (Fig. 1B

and C).   These results present a striking divergence in the effects of individual bacterial

constituents of the *C. elegans* microbiota on the level of transmission of Orsay virus from infected

animals.

Of further note, we observed a substantial amount of variability in the incidence

proportion in the presence of most bacteria (Fig. 1C). Considering both the spreaders and

uninfected animals are isogenic populations, the variability we observe between technical and

experimental replicates may be due to the inherently stochastic elements of virus transmission.

It is striking that the two *Ochrobactrum* species and *P. lurida* MYb11 were exceptions to this rule,

as it suggests these bacteria somehow mitigate the semi-random elements of virus transmission

under these conditions.

We reasoned that increased transmission of Orsay virus could be due to bacterial effects

on the shedding of virus by infected spreader animals or bacterial modulation of host

susceptibility to viral infection. To determine quantitatively how different bacterial species

modulate host susceptibility to viral infection independently of the potential differential effects

of bacteria on host shedding of viruses, we added a fixed amount of Orsay virus to plates of each

bacterial species and monitored infection using the *pals-5p::GFP* reporter. Specifically, we added

Orsay virus in doses that ranged across four orders of magnitude and then quantified the fraction

of individuals that became infected after 24 h (Fig. 1A and 1D). At lower doses of Orsay virus, the

presence of *O. vermis* MYb71 resulted in a higher fraction of infected animals than those

observed in the presence of *E. coli* OP50 or *P. lurida* MYb11 (Fig. 1D). At higher doses of Orsay

virus, the presence of *P. lurida* MYb11 resulted in a reduced fraction of infected animals

compared to *O. vermis* MYb71 and *E. coli* OP50 (Fig. 1D). We further quantified these impacts on

host susceptibility by calculating the dose of Orsay virus required to infect 50% of the population

after 24 h of exposure, which we defined as the ID_{50} (Fig. 1D and 1E). On average we observed

that the ID_{50} in the presence of *E. coli OP50* was 17-fold higher than the ID_{50} in the presence of

O. vermis MYb71 (Fig. 1E). The ID_{50} observed in the presence of *P. lurida* was 120-fold and 2300-

fold higher than the ID_{50} in the presence of *E. coli* OP50 or *O. vermis* MYb71 respectively (Fig. 1E).

We corroborated our observations from the *pals-5p::GFP* reporter by scoring a

susceptibility assay using fluorescence *in situ* hybridization to detect the RNA1 segment of the

Orsay virus genome in the intestinal cells of infected animals (Fig. 1F). As expected, we observed

that at lower doses the presence of *O. vermis* MYb71 resulted in a greater fraction of infected

animals compared to *E. coli* OP50 or *P. lurida* MYb11, while at higher doses the presence of *P.*

lurida MYb11 resulted in a reduced fraction of infected animals (Fig. 1F). Together these data

establish that individual members of the *C. elegans* microbiota can modulate host susceptibility

to Orsay virus over three orders of magnitude with dramatic consequences for the

transmissibility of Orsay virus.

Incidence Proportion = # of New Infections / Total Population

$$\text{Incidence Proportion} = \frac{\text{\# of New Infections}}{\text{Total Population}}$$

Figure 1: *Ochrobactrum* species and *P. lurida* MYb11 divergently modulate Orsay virus transmission and infection rates.
(A) Schematic representation of the transmission and susceptibility assays: Transmission can be assessed by combining infected spreader individuals (green nematode), uninfected reporter individuals (white nematodes), and various bacteria. Susceptibility can be assessed by combining uninfected reporter individuals with exogenous Orsay virus, and various bacteria. 24 h later, infection is assessed. This schematic was made using Biorender. (B) Representative images of GFP expression among individuals exposed to spreaders or no spreaders in the presence of *O. vermis* MYb71, *E. coli* OP50, or *P. lurida* MYb11. (C) Incidence proportion, calculated as indicated, of Orsay virus transmission quantified on different bacteria from the environment of *C. elegans*. Data shown are from three experiments combined, each dot represents the incidence proportion from a single plate. (D) Dose response curves of *C. elegans* to exogenous Orsay virus. The dashed line indicates the calculated dose at which 50% of the population was infected, the ID_{50}, the exact value which is given above the x-axis for each bacterium. The solid curves represent the 95% confidence interval of the modeled log-logistic function while in the presence of each bacterium. Data are from a single representative experiment, Dots represent individual plates. ((a.u.), arbitrary units). (E) Table shows the mean +/- the standard deviation of the ID_{50} ratio of the indicated bacteria measured in three experiments as in (D). (F) The fraction of individuals with staining as assessed by Fluorescence *in situ* hybridization targeting the RNA1 segment of Orsay virus. Data shown are from three experiments combined, each dot represents three pooled technical replicates from a susceptibility assay as in (D). For all plots the black bar is the mean and error bars are the 95% confidence interval (C.I.). p-values determined using one-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

***P. aeruginosa* attenuates Orsay virus transmission.**

In view of the effect of *P. lurida* MYb11 on attenuating Orsay virus infection of *C. elegans,*

we examined the effect of the distantly related bacterium, *Pseudomonas aeruginosa,* an

opportunistic pathogen of humans that has been characterized extensively[30,42]. We observed

that in the presence of either *P. aeruginosa* strains PA01 or PA14, transmission from spreader

animals to uninfected individuals was nearly completely blocked and the incidence proportion

was 11-fold and 32-fold lower than that observed in the presence of *E. coli* OP50, respectively

(Fig. 2A). The attenuating effect of *P. aeruginosa* PA01 and PA14 on virus transmission was

further reduced 3.1-fold and 9.5-fold, respectively, compared to the incidence proportion

observed in the presence of *P. lurida* MYb11 (Fig. 2A). Once more we observed substantial

variation in the incidence proportion in the presence of *E. coli* OP50, while the incidence

proportions in the presence of either *O. vermis* MYb71, *P. aeruginosa* PA01, or *P. aeruginosa*

PA14 had minimal variation, suggesting a potent overall effect on transmission under these

conditions (Fig 2A).

We confirmed that host susceptibility to Orsay virus was reduced in the presence of *P.*

aeruginosa by performing susceptibility assays at two doses of exogenous Orsay virus. At each

dose, fewer animals were infected following exposure to equivalent doses of Orsay virus in the

presence of *P. aeruginosa* PA01, PA14, or *P. lurida* MYb11 as compared to the fraction of animals

infected in the presence of *E. coli* OP50 (Fig. 2B). At the highest dose of virus used, we still

observed robust attenuation of infection in the presence of *P. aeruginosa* PA14 as 0.5% of the

animals were infected as compared to 54%, 70%, and 97% in the presence of *P. aeruginosa* PA01,

P. lurida MYb11, or *E. coli* OP50, respectively (Fig. 2B). We confirmed that no individuals were

detectably infected with Orsay virus while in the presence of *P. aeruginosa* PA14 using FISH

staining (Fig. 2C). FISH additionally confirmed that the presence of *P. aeruginosa* PA01 and *P.*

lurida MYb11 attenuated average infection to 33% and 62% of the population respectively

compared to *E. coli* OP50 which supported infection of 85% of the population. Further, *P.*

aeruginosa PA01, *P. lurida* MYb11, and *E. coli* OP50 all supported higher levels of infection than

P. aeruginosa PA14. (Fig. 2C). Together these results demonstrate a striking capacity of the

presence of *P. lurida and P. aeruginosa* species to sharply reduce and even effectively block, host

susceptibility to infection with Orsay virus.

Figure 2: *P. aeruginosa* attenuates Orsay virus transmission and infection rates.
(A) Incidence proportion of Orsay virus transmission quantified on different bacteria from *C. elegans*' environment and *P. aeruginosa* PA01 and *P. aeruginosa* PA14. Data shown are from three experiments combined, each dot represents the incidence proportion from a single plate. (B) The fraction of individuals GFP positive following exposure to two doses of exogenous Orsay virus. Data are from a single representative experiment and dots represent individual plates. (C) The fraction of individuals with staining following exposure to 100a.u. Orsay Virus as assessed by fluorescence *in situ* hybridization targeting the RNA1 segment of the Orsay Virus genome. Data shown are from three experiments combined, each dot represents three pooled technical replicates from an experiment. For all plots the black bar is the mean and error bars are the 95% confidence interval (C.I.). p-values determined using one-way ANOVA followed by Tukey's Honest Significant Difference (HSD) test (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

172 **Attenuation of Orsay virus transmission by *P. lurida* and *P. aeruginosa* is not due to effects on**

173 **Orsay virus replication.**

174   We considered the possibility that the attenuation of Orsay virus transmission in the

175 presence of *P. lurida* and *P. aeruginosa* strains might be due to inhibitory effects of the bacteria

176 on the replication of Orsay virus once transmitted to a susceptible animal host. To evaluate this

177 possibility, we made use of a plasmid-based system in which viral RNA1 is expressed through a

178 transgene introduced into *C. elegans,* so that replication of RNA1 can be assessed independent

179 of the entry of exogenous virus into the host[43]. In this system, the Orsay virus RNA1 segment,

180 which encodes the RNA-dependent RNA polymerase (RdRP), is expressed following heat-shock.

181 The expressed RNA1 may then be translated to produce the RdRP which can then replicate RNA1

182 through a negative-strand intermediate. The expression of RNA1 via heat-shock bypasses any

183 differences in viral entry or pre-replication steps, allowing for a direct test of RNA1 replication

184 efficiency under different conditions[43]. A plasmid expressing a mutated RdRP (RNA1[D601A])

185 incapable of supporting further RNA1 replication after the initial heat-shock serves as a control

186 for heat shock efficiency[43]. Using this system, we observed that Orsay virus RNA1 replication

187 efficiency was unaffected by the presence of *Pseudomonas* species relative to RNA1 replication

188 observed in the presence of *E. coli* OP50 (Fig. 3A). Additionally, there were no differences

189 observed in heat-shock efficiency between the different bacteria (Fig. 3A). These data suggest

190 that bacteria-induced differences in RNA1 replication in the host do not explain the substantial

191 attenuation of Orsay virus transmission and infection rates caused by *P. aeruginosa* PA01 or PA14

192 and *P. lurida* MYb11.

193   We sought to confirm the successful replication of the plasmid expressed RNA1 of Orsay

194 virus by assessing whether we could detect the replication of Orsay virus RNA1 following rare

events of natural infection of *C. elegans* in the presence of *P. aeruginosa* PA14. We exposed

young adult wild-type (N2) or RNAi defective animals (*rde-1(ne219)*) to exogenous Orsay virus

while in the presence of *P. aeruginosa* PA14 and quantified the amount of virus present at 2 h

and 24 h post-exposure, with the difference in RNA levels at these points reflecting viral genome

replication. In the N2 background viral RNA1 levels increased by 3.5-fold at 24 h relative to levels

at 2 h post-exposure (Fig. 3B). Orsay virus is more effective at replicating in the *rde-1* background

compared to N2 and in the *rde-1* background viral RNA1 levels increased by 9000-fold at 24 h

relative to levels at 2 h post-exposure (Fig. 3B)[23]. Together, these results suggest that Orsay virus

RNA1 can also replicate once natural infection is achieved in both the wild-type and *rde-1*-

defective background while animals are in the presence of *P. aeruginosa* PA14. These data

confirm our findings that bacteria induced differences in viral replication do not explain the

substantial attenuation of Orsay virus transmission in the presence of *P. aeruginosa* PA14.

Figure 3: *P. aeruginosa* and *P. lurida* MYb11 do not eliminate Orsay virus replication.
(A) Animals carrying either a wild-type (RNA1(WT)) or replication defective (RNA1(D601A)) transgenic viral RNA replicon system were used to assess Orsay virus replication efficiency independently of virus entry. P-values determined using one-way ANOVA followed by Dunnett's test (B) Orsay virus RNA1 levels of N2 and *rde-1(ne219)* animals exposed to *P. aeruginosa* PA14 and exogenous Orsay virus 2- and 24- h post infection (hpi). P-values were determined using Welch's t-Test For each plot, each dot represents five pooled technical replicates, the black bar is the mean, error bars are the standard deviation, RNA1 levels were quantified by qPCR, and the data shown are for three independent experiments. (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

Attenuation of Orsay virus transmission by *Pseudomonas* species is dependent on regulators

of bacterial quorum sensing.

Extensive studies on *Pseudomonas* species have demonstrated the importance of quorum

sensing for regulating many community-level behavior in response to growing population

density[44]. *P. aeruginosa* relies upon three quorum sensing systems: *las*, *rhl*, and *pqs*. These

systems are arranged hierarchically, however crosstalk between them is extensive (Fig. 4A)[44]. In

P. aeruginosa, an additional layer of regulation stems from two-component regulatory systems

such as *gacA/gacS* which regulates numerous genes that together influence quorum sensing,

virulence, and biofilm development (Fig. 4A)[45–47]. *P. aeruginosa* possesses three

exopolysaccharide biosynthetic clusters that each can contribute to biofilm formation: *pel*, *psl*,

and *alg*. However, *P. aeruginosa* PA01 preferentially produces Psl while *P. aeruginosa* PA14 is

unable to synthesis Psl and produces Pel[48–52]. We hypothesized that quorum sensing might

mediate the effect of *Pseudomonas* strains to attenuate virus transmission, while differences in

exopolysaccharide production might mediate the enhanced attenuation of virus transmission

observed in the presence of *P. aeruginosa* PA14 compared with what was observed in the

presence of *P. aeruginosa* PA01. Therefore, we tested a panel of *P. aeruginosa* PA01 and *P.*

aeruginosa PA14 quorum sensing and biofilm mutants to determine whether quorum sensing or

biofilm formation was involved in the attenuation of Orsay virus infection mediated by *P.*

aeruginosa.

Mutations of any the regulators of the *las*, *rhl*, or *pqs* quorum sensing systems suppressed

the attenuation of Orsay virus infection caused by the presence of wild-type *P. aeruginosa* PA01

(Fig. 4B). Knockout of *gacA* in *P. aeruginosa* PA01 also suppressed the attenuation of Orsay virus

infection (Fig. 4B). On the other hand, mutation of any of the three exopolysaccharide production

pathways had no effect on infection (Fig. 4B). These data support a role for quorum sensing

regulated processes in reducing Orsay virus infection rates but do not implicate a role for

exopolysaccharide biosynthesis.

For *P. aeruginosa* PA14, mutation of *gacA* or *rhlR* suppressed the attenuation of Orsay

Virus infection observed in the presence of wild-type *P. aeruginosa* PA14 (Fig. 4C). Loss of the

rhlR regulators *rhlI* or *pqsE* alone had no effect on the attenuation of Orsay virus infection, but

simultaneous mutation of both *rhlI* and *pqsE* did suppress the attenuation of Orsay virus infection

observed in the presence of wild-type *P. aeruginosa* PA14, similar to that observed for the *rhlR*

mutant, suggesting *rhlI* and *pqsE* function redundantly to regulate *rhlR* in this context (Fig. 4C)[53].

Mutation of *lasI* or *lasR* suppressed the attenuation of Orsay virus infection to a lesser extent

(Fig. 4C). Independent mutation of two genes responsible for *pel* or *alg* exopolysaccharide

production had no effect on infection rates (Fig. 4C). These data suggest a role for quorum

sensing in mediating *P. aeruginosa* PA14 suppression of Orsay virus infection, as we observed for

P. aeruginosa PA01. However, our results obtained in the presence of *P. aeruginosa* PA14 suggest

that there may be some differential regulation of the bacterial effectors responsible in

comparison to *P. aeruginosa* PA01, or additional non-quorum sensing related factors that also

mediate suppression.

As *P. aeruginosa* mutants could suppress the effects of wild-type *P. aeruginosa* on *C.*

elegans infection in the presence of exogenous virus, we next confirmed that these *P. aeruginosa*

mutants similarly affected not just host susceptibility, but also transmission from infected

spreader animals. All *P. aeruginosa* PA01 quorum sensing mutants suppressed the attenuation

of transmission by increasing the incidence proportion >6-fold compared to the wild-type *P.*

aeruginosa PA01 (Fig. 4D). *P. aeruginosa* PA14 *rhlR* and *gacA* mutants suppressed the attenuation

of transmission by increasing the incidence proportion 19-fold and 17-fold respectively compared

to wild-type *P. aeruginosa* PA14, but a *lasI* mutant had minimal effect consistent with the pattern

we observed in the susceptibility assay using exogenous Orsay virus (Fig. 4D). Likewise, the

magnitude of suppression of the attenuation of Orsay virus transmission observed in the *P.*

aeruginosa rhlR or *gacA* mutants was greater in the *P. aeruginosa* PA01 background compared

to the *P. aeruginosa* PA14 background, potentially suggesting the existence of additional strain

specific factors that act specifically in *P. aeruginosa* PA14 to attenuate Orsay virus transmission.

It is additionally interesting to note that similar to the incidence proportion in the

presence of *E. coli* OP50, the variation in the incidence proportion in the presence of all *P.*

aeruginosa mutants increased (Fig 4D). This was true even for the incidence proportion observed

in the presence of *P. aeruginosa* PA14 *lasI* mutant, which while not statistically different from

wild-type *P. aeruginosa* PA14, had a qualitatively different distribution nonetheless (Fig 4D).

Figure 4: Mutation of *P. aeruginosa* quorum sensing regulators and the two-component response regulator *gacA* suppresses the attenuation Orsay virus infection.

(A) Diagram demonstrating the three quorum sensing systems in *P. aeruginosa*. Each system encodes the enzyme(s) (LasI, RhlI, PqsA(BCDH)) to produce an autoinducer (3OC12HSL – 3-oxo-C12-homoserine lactone, C4HSL –butanoyl homoserine lactone , PQS – Pseudomonas quinolone signal) that is recognized by its cognate receptor (LasR, RhlR, PqsR) that influences gene transcription and the activity of the other quorum sensing systems. An additional level of regulation stems from the two-component *gacS/gacA* system. External signals are recognized by the histidine kinase GacS, which phosphorylates the response regulator GacA that indirectly influences quorum sensing processes. IM is inner membrane, OM is outer membrane. Arrows represent crosstalk between the various system. Adapted from Rutherford and Bassler 2012 and Song, Li, and Wang 2023. (B-C) The fraction of individuals GFP positive following exposure to exogenous Orsay virus. Data are from a single representative experiment, bars represent mean, dots represent individual plates. (B) Wild-type *P. aeruginosa* PA01 compared to mutant *P. aeruginosa* PA01 strains using 10a.u. of Orsay virus. (C) Wild-type *P. aeruginosa* PA14 compared to mutant *P. aeruginosa* PA14 strains using 100a.u. of Orsay virus. (D) Incidence proportion of Orsay virus transmission in the presence of *P. aeruginosa* wild-type versus select *P. aeruginosa* mutants. Data are from three experiments, bars represent mean, dots represent individual plates. For all plots error bars represent 95% C.I. For B-E, p-values were determined using one-way ANOVA followed by Dunnett's test. (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

***P. lurida gacA* is required for attenuation of Orsay virus transmission.**

We next sought to determine whether our findings from the interaction of *C. elegans* and

Orsay virus in the presence of *P. aeruginosa*, a well-characterized pathogen of *C. elegans*, could

inform us further regarding the mechanisms underlying the attenuation of Orsay virus

transmission in the presence of *P. lurida* MYb11, a non-pathogenic constituent of the *C. elegans*

microbiota[39]. In particular, we identified a *gacA* ortholog using Orthovenn2 and generated a

putative knockout allele by removing the central 194 out of 214 amino acids (Fig. 5A)[54]. Following

exposure to exogenous Orsay virus, knockout of *gacA* in *P. lurida* MYb11 led to infection of 80%

of the population compared to infection of only 14% of the population in the presence of wild-

type *P. lurida* MYb11 from exogenous Orsay virus (Fig. 5B). The *gacA* mutation also suppressed

the attenuation of Orsay virus transmission, increasing the incidence proportion 2.9-fold

compared to wild-type *P. lurida* MYb11 (Fig. 5C). These data suggest that *gacA* has a conserved

role across distant *Pseudomonas* species in the attenuation of Orsay virus transmission and

infection of *C. elegans*.

P. aeruginosa genes linked to virulence reduce Orsay virus transmission and infection.

We designed a candidate-based screen using a non-redundant transposon insertion

library to gain further insight into the genetic regulation of *P. aeruginosa* mediated reduction of

Orsay Virus transmission and infection (Fig. 6A)[55]. We identified candidate genes to include in our

screen from three sources: *gacA*-regulated genes, *rhlR*-regulated genes (but *rhlI*- or *pqsE*-

independent), and the set of genes required for full virulence in *C. elegans* previously identified

by Feinbaum et al. (Fig 6A)[47,56,57]. Of the 201 genes tested, 15 putative hits were identified with

the corresponding *P. aeruginosa* PA14 mutants exhibiting suppression of the attenuation of

Orsay virus infection by wild-type *P. aeruginosa* PA14 (Fig. 6A, Supp. Table 3). Using these 15

candidate genes, we performed an additional set of susceptibility assays which confirmed six of

the hits (Fig. 6B, Supp. Table 3). The identified hits grouped into two clusters based on the

strength of their suppression. Transposon insertion in the genes *ptsP*, *prpC*, and *kinB* led to

marked suppression of the attenuation of Orsay Virus leading to infection of 97%, 89%, and 82%

of the population respectively compared to only 4.2% of the population in the presence of wild-

type *P. aeruginosa* PA14. Transposon insertion in three additional genes, *clpA*, *glnK*, and *fabF1*

resulted in weaker, but robust suppression of the attenuation of Orsay virus leading to infection

of 34%, 28%, and 17% of the population respectively. We additionally tested whether transposon

insertion into these genes suppressed attenuation of Orsay virus transmission. While we

observed a trend towards mutations affecting susceptibility also affecting virus transmission, we

again observed a high degree of variation in the transmission assay, such that only mutation of

ptsP led to statistically significant suppression of Orsay virus attenuation (Fig. 6C). However,

similar to the effect observed with *lasI* mutation, the distributions of the incidence proportion

observed in the presence of the *prpC*, *kinB*, or *glnK* mutants were qualitatively different from the

nearly uniform incidence proportion observed in the presence of wild-type *P. aeruginosa* PA14

suggesting that mutation of these genes was impacting the attenuation of Orsay virus

transmission to some extent (Fig. 6C).

All six of the hits originated from the set of genes required for full *P. aeruginosa* PA14

virulence in *C. elegans*. We noted that these hits represented only six out of the 41 tested genes

that are known to influence virulence, indicative of some degree of specificity in the

consequences of these mutations beyond their general effect on *P. aeruginosa* PA14 virulence[57].

While these six genes have orthologs within *P. lurida* MYb11, knockout of the *P. lurida* MYb11

orthologs of *ptsP*, *prpC*, or *kinB* failed to suppress the attenuation of Orsay virus infection or

transmission by *P. lurida* MYb11(Supp. Fig. 2A and 2B)[57,58]. One explanation of these results is

that these genes play different roles within *P. lurida* MYb11 and therefore their mutation did not

have the same consequences for Orsay virus attenuation. Alternatively, these results might

further suggest that *ptsP*, *prpC*, and *kinB* act to attenuate Orsay virus infection via some specific

effect on *P. aeruginosa* PA14 virulence.

Figure 6: Mutation of *P. aeruginosa* PA14 genes related to virulence suppresses the attenuation of Orsay virus infection.
(A) Diagram describing the design and results of the screen to identify suppressors of *P. aeruginosa* PA14 Orsay virus attenuation. (B) The fraction of individuals GFP positive following exposure to 100 a.u. of exogenous Orsay virus. Data are from a single representative experiment. (C) Incidence proportion of Orsay virus transmission while individuals are present on *P. aeruginosa* PA14 wild-type compared to *P. aeruginosa* PA14 mutants. Data are from three experiments and dots represent individual plates. For all plots the black bar is the mean and error bars are the 95% confidence interval (C.I.). p-values determined using one-way ANOVA followed by Dunnett's test (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

323
Discussion

In our study, we quantitatively define the effect of bacteria on the transmission of Orsay

virus in the *C. elegans* host. While the presence of *O. vermis* MYb71 enhanced the transmission

of Orsay virus, the presence of *P. lurida* MYb11 or *P. aeruginosa* strains PA01 and PA14

attenuated Orsay virus transmission. The enhancement of Orsay virus transmission by *O. vermis*

MYb71 and reduction of transmission by *P. lurida* MYb11 and *P. aeruginosa* PA01 and PA14 was

mirrored in assays assessing infection rates by exogenous virus. Our results using exogenous virus

demonstrate that host susceptibility to Orsay virus infection may vary by over three orders of

magnitude in the presence of *O. vermis* MYb71 versus *P. lurida* MYb11 which are found in the

natural environment in association with *C. elegans*. Moreover, we observe that pathogenic *P.*

aeruginosa, which may be found in natural environments, can further attenuate Orsay virus

transmission[31]. Our work is also consistent with a recent study by González and Félix that also

reported that monoaxenic cultures of other bacteria from the environment of *C. elegans* impact

host susceptibility[59].

Our observations that two *Ochrobactrum* species promoted transmission of Orsay virus

are intriguing given that other *Ochrobactrum* species have also been linked to viral infections.

Ochrobactrum intermedium promoted poliovirus infection in mice and enhanced poliovirus

stability in vitro[13]. *Ochrobactrum anthropi*, an opportunistic human pathogen, was identified

using a random forest analysis as one of the most predictive features differentiating the upper

respiratory tract of human patients recovering from influenza infection versus healthy controls[60].

Our observations in the *C. elegans* host raise the speculative possibility that *Ochrobactrum*

colonization may have roles in evolutionarily diverse hosts in modulating viral infection.

P. lurida MYb11 and *P. aeruginosa* PA01 and PA14 shared the ability to attenuate Orsay

virus infection and transmission, even as *P. lurida* and *P. aeruginosa* have markedly distinct

effects on the *C. elegans* host. *P. lurida* MYb11 promotes developmental rate and fitness,

although may do so at a minor cost to overall lifespan[39,61]. In contrast, *P. aeruginosa* is a pathogen

of *C. elegans* and numerous other organisms[30,62]. We identified that *gacA* regulates the

attenuation of Orsay virus by *P. aeruginosa* PA01 and PA14 and *P. lurida* MYb11. The *gacS/gacA*

system regulates many phenotypes in a variety of γ-proteobacteria[63]. In Pseudomonads,

gacA/gacS regulate processes related to quorum sensing, virulence, and biocontrol in a

population density dependent manner[63]. These data raise the possibility that *Pseudomonas*

quorum sensing pathways are required for the effects on Orsay virus transmission and infection.

Quorum sensing also influences additional aspects of *P. aeruginosa* physiology, including

swarming behaviors, biofilm formation, secondary metabolism rates, and overall transcription

patterns[44].To gain additional insights into the *P. aeruginosa* effects on Orsay virus transmission

we conducted a candidate-based screen including genes influenced by quorum sensing and genes

that regulate virulence towards *C. elegans*[47,56,57]. We identified six genes, *ptsP*, *prpC*, *kinB*, *clpA*,

glnK, and *fabF1*, that when mutated suppressed *P. aeruginosa* PA14 attenuation of Orsay virus

infection. Each of these genes has been shown to be required for full *P. aeruginosa* PA14

virulence in *C. elegans*[57]. However, we observed that only these six out of the total 41 virulence-

related genes identified in Feinbaum *et al*. influenced Orsay virus infection, arguing against a role

for general virulence in the attenuation of Orsay virus transmission and infection caused by *P.*

aeruginosa PA14 and *P. aeruginosa* PA01[57].

C. elegans is unlikely to associate with a single bacterium in its natural environment.

However, *C. elegans* shows clear behavioral preferences for grazing on certain bacteria from its

environment and may eat monoxenic lawns in the wild[64]. Our data demonstrate that individual

bacteria can have a profound impact on Orsay virus transmission rates in a species-specific

manner. Additionally, the tractability of the *C. elegans*-Orsay virus experimental system allowed

us to identify molecular determinants of viral transmission and will be useful for identifying

additional biotic factors that influence viral transmission. Our work builds on the expanding body

of knowledge showing that the microbiota can influence the interactions between viruses and

their animal hosts.

Supplementary Figure 1: Transmission does not occur from individuals infected during the course of a transmission or susceptibility assay.
(A-B) Individuals were exposed to 0 a.u. or 5 a.u. exogenous Orsay virus in the presence of *O. vermis* MYb71. 8 h post infection five infected individuals from the 5 a.u. plate were transferred to the 0 a.u. plate to assess whether transmission from these individuals would occur. Both plates were scored 16 h after the transfer. Data are from a single representative experiment, bars represent mean, dots represent individual plates. Error bars are the 95% confidence interval (C.I.). p-value determined by Student's t-test (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

Supplementary Figure 2: Knockout of the *P. lurida* MYb11 orthologs of *ptsP*, *prpC*, or *kinB* does not suppress Orsay virus attenuation in susceptibility or transmission assays.
(B) The fraction of individuals GFP positive following exposure to 10 a.u. of exogenous Orsay virus in the presence of the indicated bacteria. Data are from a single representative experiment. (C) Incidence proportion of Orsay virus transmission in the presence of the indicated bacteria. The *E. coli* OP50 reference is shared with Figure 5C. Data are from three experiments and dots represent individual plates. For all plots the black bar is the mean and error bars are the 95% confidence interval (C.I.). p-values determined using one-way ANOVA followed by Dunnett's test (NS non-significant, *p<0.05, **p<0.01, ***p<0.001).

Materials and methods

C. elegans strains and growth conditions

C. elegans were maintained on NGM agar plates (17 g agar, 2.5 g peptone, 3 g NaCl per 1 L water) containing *E. coli* OP50[41]. Strains bearing the *glp-4(bn2)* temperature sensitive mutation were maintained at 16°C, while all other were maintained at 20°C. All assays were performed on SKA assay plates at 20°C (17 g agar, 3.5 g peptone, 3 g NaCl per 1 L water)[30]. A full list of *C. elegans* strains used in this study is contained within the Supplement (Supplement Table 1). For all assays relying upon *pals-5p::GFP*, induction was manually assessed using a Nikon SMZ18 stereofluorescent microscope.

Bacterial strains and growth conditions

All bacteria were grown in Luria broth (10 g tryptone, 5 g yeast, 10 g NaCl per 1 L water). *E. coli* OP50 and *P. aeruginosa* strains were grown at 37°C with shaking, while all other strains were grown at 27°C with shaking. A full list of bacterial strains used in this study is contained within the Supplemental Information (Supplement Table 2).

Orsay virus isolation and dose testing

Orsay virus was isolated from infected WUM31(*rde-1(ne219);jyIs9[pals-5p::gfp;myo-2p::mCherry]*) individuals. Plates with gravid WUM31 were bleached to obtain eggs. Eggs were hatched overnight in M9 solution rotating at 20°C and the L1 larvae were arrested to synchronize the population. L1 larvae were combined with 100 μL of 6x concentrated *E. coli* OP50 and 50 μL

of Orsay virus filtrate, plated on SKA plates, and once dried, placed at 20°C for 48 h. Four GFP

positive individuals were then transferred to a new 6cm NGM plate with *E. coli* OP50 and

maintained until just starved. Twenty such 6cm plates were washed with 10 mL of M9 and the

resulting suspension was Dounce homogenized. Alternatively, 3.5 cm SKA plates containing

infected WUM31 animals was allowed to starve and then equally chunked onto four 10 cm NGM

plates with *E coli* OP50. Once these plates were just starved, eight 10 cm plates were washed and

homogenized as above. The homogenized suspension was then centrifuged at 13.2xg for five

min. The supernatant was passed through a 0.22 μm filter and aliquoted. Each batch of virus was

tested for potency and an ID_{50} calculated for *E. coli* OP50. Doses referenced throughout the

manuscript refer to actual microliters applied to each plate. However, no method was used to

normalize doses between batches of virus and so we have chosen to refer to doses in terms of

arbitrary units (a.u.). The average ID_{50} while on *E. coli* OP50 for all virus batches used in this study

was 3.6 suggesting an average dose of 1 a.u. roughly corresponds to 0.28x the ID_{50} on *Escherichia*

coli.

<u>Preparation of Uninfected Individuals for transmission and susceptibility assays</u>

Prior to all assays, plates with fecund ZD2611(*glp-4(bn-2);jyIs8[pals-5p::gfp;myo-2p::mCherry]*)

were bleached to obtain eggs. Eggs were hatched overnight in M9 solution rotating at 20°C and

the L1 larvae were arrested to synchronize the population. ZD2611 L1s were dropped onto plates

containing *E. coli* OP50 and placed at 20°C for 64-72 h. At this temperature reproduction is

delayed, but not eliminated. The resulting young adults were washed off the plates in M9 and

centrifuged at 1000xg for one minute. After removing the supernatant, the young adults were

once again washed in M9, then centrifuged at 1000 x g for one minute. Young adults were directly

dropped onto prepared plates described below.

<u>Preparation of Bacteria for transmission and susceptibility assays</u>

Unless otherwise indicated, bacteria were cultured overnight and added to cover the assay

plates. Plates containing *E. coli* OP50 or *P. aeruginosa* strains were placed at 37°C for 24 h. Plates

containing *O. vermis* MYb71 or *P. lurida* MYb11 strains were placed at 25°C for 24 h. All plates

were then moved to room temperature for an additional 24 h before the addition of virus.

For the natural isolate transmission screen and early transmission assay cultures were grown to

late stationary phase. Cultures were then spun down for six minutes at 4000xg and reconstituted

to 25 mg/mL in their own supernatant. 150 µL of this suspension was combined with 50 µL of a

mix of M9 for transmission assays or 50 µL of a combination of Orsay virus filtrate and M9 for the

early transmission assay. This mixture was combined with young adult ZD2611 individuals and

directly added to assay plates before drying.

<u>Transmission assays</u>

To obtain infected spreader animals, fecund ZD2610 (*glp-4(bn-20);rde-1(ne219); jyIs8[pals-*

5p::gfp;myo-2p::mCherry]) were bleached to obtain eggs. Eggs were hatched overnight in M9

solution rotating at 20°C and the L1 larvae were arrested to synchronize the population. Arrested

ZD2610 L1s were dropped onto NGM plates containing *E. coli* OP50 and placed at 20°C for 48 h.

Animals were then washed off the plates in M9 and centrifuged at 1000xg for one min. After

removing the supernatant, the young adults were washed with M9 again then centrifuged at

1000xg for one min. Supernatant was once again removed, and individuals were immediately

mixed with 100 μL of 6x overnight *E. coli* OP50 culture and 50 μL of Orsay virus for a minimum of

18-20 h. GFP-positive ZD2610 young adults were identified and picked onto transfer plates

containing the appropriate bacteria prepared as indicated above for 4-6 h. After this period, five

spreaders were transferred to assay plates containing the same bacteria and approximately 100

uninfected ZD2611 individuals to start the assay. Transmission assays were scored 24 h after

adding spreader individuals.

For the natural isolate transmission screen, arrested ZD2610 L1s were combined with 100 μL of

6x concentrated *E. coli* OP50 and 50 μL of Orsay virus filtrate and once dried, placed at 20°C for

64-72 h. On the day of the assay, GFP-positive ZD2610 young adults were identified and picked

onto transfer plates containing the appropriate bacteria prepared as indicated above for 4-6 h.

After this period, five spreaders were transferred to assay plates containing the same bacteria

and approximately 100 uninfected ZD2611 individuals to start the assay.

Incidence proportion

Incidence proportion was calculated by dividing the number of newly infected animals by the

total non-spreader population per plate. Individual plates for which the total number of GFP

positive individuals after 24 h was less than the initial number of spreaders placed on the plate

were excluded.

<u>Susceptibility assays</u>

Unless otherwise indicated, Orsay virus filtrate at the indicated doses was diluted in filtered M9

solution. 200 uL total was applied to each assay plate and swirled to cover the entire bacterial

lawn. Plates were then dried before the addition of approximately 100 uninfected ZD2611

individuals prepared as indicated above. Assays were scored 24 h later using a Nikon SMZ18

stereo fluorescent microscope and the fraction of GFP-positive individuals was calculated.

<u>Early Transmission Assay</u>

For early transmission assay, young ZD2611 adults were exposed to 0a.u. or 5a.u. exogenous

Orsay virus in the presence of *O. vermis* MYb71. 8 hours post infection five infected individuals

from the 5 a.u. plate were transferred to the 0 a.u. plate to assess whether transmission from

these individuals would occur. Both plates were scored 16 hours after the transfer.

<u>Fluorescence *in situ* hybridization</u>

Previously published probes were obtained to target the RNA1 segment of Orsay virus[25]. Animals

were infected according to the methods described above for susceptibility assays. Animals were

processed according to the Stellaris RNA FISH Protocol for *C. elegans* (LGC Biosearch

Technologies) with minor modifications. Briefly, young adults were washed off the plate and

rinsed twice in M9. Animals were then fixed for 30 min rotating in a microcentrifuge tube at 20°C.

After washing twice with 1 mL of phosphate buffered saline animals were permeabilized in 70%

of ethanol and stored at 4°C for 1-7 d. Animals were washed with Wash Buffer A before addition

of 100 μL of hybridization buffer containing 3 μL of RNA1 probe mix. Probe was hybridized

overnight at 46°C. Animals were then washed with Wash Buffer A alone once, and then again

with Wash Buffer A containing 5 ng/mL DAPI. Lastly 100 μL of Wash Buffer B was added before

mounting animals on slides with 25 μL of Vectashield Mounting Medium. The fraction of

individuals with RNA1 staining was then quantified using a Nikon SMZ18 stereofluorescent

microscope or a Zeiss AxioImager Z1 compound fluorescent microscope.

RNA Extraction

Animals were washed 5 times in M9 and collected in TRIzol reagent (Invitrogen) and stored at -

80C before extraction. RNA extraction was performed using Direct-zol RNA microprep kits

(Zymo Research) following the manufacturer's instructions.

qPCR

cDNA was made using 500ng of RNA as template (Promega GoScript Reverse

Transcriptase/Random Primers). cDNA was diluted at 1/40 and qPCR were performed using 1ul

of diluted cDNA (GoTaq Promega) and run on QuantStudio 3 Real Time PCR system. RNA1 levels

were then quantified via qPCR using previously published primers[23].

Plasmid-based replication experiments

Adult animals carrying a transgene containing the wild-type RNA1 or RNA1D601A Orsay virus

genome segment under the control of a heat-inducible promoter were placed on the indicated

bacteria for 4 h before heat-shock at 33°C for 2 h[43]. Animals then recovered at 20°C for 20 h

before harvesting for RNA extraction and qPCR as detailed above. RNA1 levels were normalized

to the values obtained from animals bearing the wild-type RNA1 and exposed to *E. coli* OP50.

<u>Orsay virus replication in the presence of *P. aeruginosa* PA14</u>

Adult ERT54 (*jyIs8[pals-5p::gfp;myo-2p::mCherry])* or WUM31 (*rde-1(ne219); jyIs8[pals-*

5p::gfp;myo-2p::mCherry)] were exposed to exogenous Orsay virus in the presence of *P.*

aeruginosa PA14. After 2 h or 24 h, animals were harvested for RNA extraction and qPCR as

detailed above. RNA1 levels were quantified via qPCR. Within each genotype, the data for each

experiment were normalized to the 2 h timepoint.

<u>*Pseudomonas lurida* MYb11 mutant construction</u>

A protocol developed for allelic exchange in *P. aeruginosa* was modified for use in *P. lurida*

MYb11[65]. Briefly, homology arms flanking the region to be deleted were obtained using

polymerase chain reaction (PCR) and cloned into the pExG2-KanR suicide vector using Hi-Fi

Assembly (New England Biolabs)[66]. DH5α *E. coli* were transformed using a standard heat shock

protocol. Successful transformants were selected for on LB+Kanamycin (50 μg/mL) plates and

colony PCR was performed to check for proper insert size in the transformants. *E. coli* bearing

the desired plasmid were grown overnight in LB+Kanamycin (50 μg/mL). Plasmids were then

obtained using a Qiagen MiniPrep Kit (Qiagen). Plasmids were assessed for the desired sequence

by Sanger sequencing and transformed into *P. lurida* MYb11 using the following electroporation

procedure. *P. lurida* MYb11 was grown overnight then placed on ice for 30 min. The culture was

spun down at 4°C and washed twice with ice-cold water. After reconstitution in 100 μL of ice-

cold water the suspension was transferred to a pre-chilled cuvette and transformed at 630 kV

using an Eporator (Eppendorf). 900 μL of LB was added, and the suspension was transferred to a

microcentrifuge tube and incubated at 27°C for 2 h with shaking. The suspension was then plated

on LB+Kanamycin (50 μg/mL) plates and grown for 48 h at 25 °C. Colonies were picked and grown

overnight in LB. Cultures were streaked onto sucrose plates (15 g agar, 10 g tryptone, 5 g yeast,

60 g sucrose per 1 L water) to perform sucrose-based counter selection[65]. Colonies that survived

were genotyped for the expected deletion.

<u>Data visualization and statistics</u>

All experiments were performed three times. For transmission, qPCR, and FISH-based

susceptibility assays data from each experiment are combined. For susceptibility assays a single

representative experiment is shown. Data were analyzed in R Studio[67]. When comparing all the

means of more than two groups p-values were calculated using one-way ANOVA followed by the

Tukey HSD test. When comparing multiple experimental groups to a control group p-values were

calculated using one-way ANOVA followed by Dunnett's test. P-values for assays comparing only

two groups were calculated using Student's t-test or Welch's t-test as indicated. *E. coli* OP50 is

included in all experiments as a reference but was not included for statistical comparison unless

explicitely noted. Susceptibility assay curves were modeled using the drc[68] package in R. A two-

parameter log-logistic function was used to model the curve and ID_{50} values were calculated

using the ED function. Plots were made using the gdata[69], scales[70], drc[68], Rmisc[71], multcomp[72],

ggplot2[73], ggsignif[74], and cowplot[75] packages.

<u>Acknowledgements</u>

We would like to thank members of the Kim/Fischer laboratory for helpful comments during

the preparation of this work. We would like to thank David Wang, Emily Troemel, Marie-Anne

Félix, Eliana Drenkard, Simon Dove, E. Peter Greenberg, Matthew Parsek, Jon Paczkowski, and

Read Pukkila-Worley for kindly providing strains. Some strains were provided by the CGC, which

is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We would like to

further thank Brendan O'Hara, Michael Gebhardt, and Simon Dove for assistance developing

the *P. lurida* MYb11 transformation protocol. B.G.V. was partially supported by the MIT

Department of Biology Graduate Program. B.G.V., N.S., S.E.J.F., and D.H.K were supported by

NIH Grant R35GM141794.

References

1.   Edwards, R. A. & Rohwer, F. Viral metagenomics. *Nat Rev Microbiol* **3**, 504–510 (2005).

2.   Srinivasiah, S. *et al.* Phages across the biosphere: contrasts of viruses in soil and aquatic
environments. *Res Microbiol* **159**, 349–357 (2008).

3.   Pica, N. & Bouvier, N. M. Environmental factors affecting the transmission of respiratory
viruses. *Curr Opin Virol* **2**, 90–95 (2012).

4.   Thangavel, R. R. & Bouvier, N. M. Animal models for influenza virus pathogenesis,
transmission, and immunology. *J Immunol Methods* **410**, 60–79 (2014).

5.   de Vries, R. D. *et al.* Animal models of SARS-CoV-2 transmission. *Curr Opin Virol* **50**, 8–16
(2021).

6.   Lowen, A. C., Mubareka, S., Steel, J. & Palese, P. Influenza Virus Transmission Is
Dependent on Relative Humidity and Temperature. *PLoS Pathog* **3**, e151 (2007).

7.   SCHULMAN, J. L. & KILBOURNE, E. D. Airborne Transmission of Influenza Virus Infection
in Mice. *Nature* **195**, 1129–1130 (1962).

8.   Quinn, T. C. *et al.* Viral Load and Heterosexual Transmission of Human Immunodeficiency
Virus Type 1. *New England Journal of Medicine* **342**, 921–929 (2000).

9.   Edenborough, K. M., Gilbertson, B. P. & Brown, L. E. A Mouse Model for the Study of
Contact-Dependent Transmission of Influenza A Virus and the Factors That Govern
Transmissibility. *J Virol* **86**, 12544–12551 (2012).

10. Price, G. E., Lo, C.-Y., Misplon, J. A. & Epstein, S. L. Mucosal Immunization with a
Candidate Universal Influenza Vaccine Reduces Virus Transmission in a Mouse Model. *J
Virol* **88**, 6019–6030 (2014).

11. Chua, B. Y. *et al.* Inactivated Influenza Vaccine That Provides Rapid, Innate-Immune-
System-Mediated Protection and Subsequent Long-Term Adaptive Immunity. *mBio* **6**,
(2015).

12. Kane, M. *et al.* Successful Transmission of a Retrovirus Depends on the Commensal
Microbiota. *Science (1979)* **334**, 245–249 (2011).

13. Kuss, S. K. *et al.* Intestinal Microbiota Promote Enteric Virus Replication and Systemic
Pathogenesis. *Science (1979)* **334**, 249–252 (2011).

14. Wu, P. *et al.* A Gut Commensal Bacterium Promotes Mosquito Permissiveness to
Arboviruses. *Cell Host Microbe* **25**, 101-112.e5 (2019).

15. Zheng, D., Liwinski, T. & Elinav, E. Interaction between microbiota and immunity in health
and disease. *Cell Res* **30**, 492–506 (2020).

16. K Berger, A., Yi, H., B Kearns, D. & A Mainou, B. Bacteria and bacterial envelope
components enhance mammalian reovirus thermostability.
doi:10.1371/journal.ppat.1006768.

17. Hedges, L. M., Brownlie, J. C., O'Neill, S. L. & Johnson, K. N. *Wolbachia* and Virus
Protection in Insects. *Science (1979)* **322**, 702–702 (2008).

18. Teixeira, L., Ferreira, Á. & Ashburner, M. The Bacterial Symbiont Wolbachia Induces
Resistance to RNA Viral Infections in Drosophila melanogaster. *PLoS Biol* **6**, e1000002
(2008).

19. Moreira, L. A. *et al.* A Wolbachia Symbiont in Aedes aegypti Limits Infection with Dengue,
Chikungunya, and Plasmodium. *Cell* **139**, 1268–1278 (2009).

20.   Bian, G., Xu, Y., Lu, P., Xie, Y. & Xi, Z. The Endosymbiotic Bacterium Wolbachia Induces
Resistance to Dengue Virus in Aedes aegypti. *PLoS Pathog* **6**, e1000833 (2010).

21.   Pimentel, A. C., Cesar, C. S., Martins, M. & Cogni, R. The Antiviral Effects of the Symbiont
Bacteria Wolbachia in Insects. *Front Immunol* **11**, (2021).

22.   Schulenburg, H. & Félix, M.-A. The Natural Biotic Environment of *Caenorhabditis elegans*.
Genetics **206**, 55–86 (2017).

23.   Félix, M.-A. *et al.* Natural and Experimental Infection of Caenorhabditis Nematodes by
Novel Viruses Related to Nodaviruses. *PLoS Biol* **9**, e1000586 (2011).

24.   Félix, M.-A. & Wang, D. Natural Viruses of *Caenorhabditis* Nematodes. *Annu Rev Genet*
53, 313–326 (2019).

25.   Franz, C. J. *et al.* Orsay, Santeuil and Le Blanc viruses primarily infect intestinal cells in
Caenorhabditis nematodes. *Virology* **448**, 255–264 (2014).

26.   Sarkies, P., Ashe, A., Le Pen, J., McKie, M. A. & Miska, E. A. Competition between virus-
derived and endogenous small RNAs regulates gene expression in *Caenorhabditis*
elegans. *Genome Res* **23**, 1258–1270 (2013).

27.   Bakowski, M. A. *et al.* Ubiquitin-Mediated Response to Microsporidia and Virus Infection
in C. elegans. *PLoS Pathog* **10**, e1004200 (2014).

28.   Chen, K., Franz, C. J., Jiang, H., Jiang, Y. & Wang, D. An evolutionarily conserved
transcriptional response to viral infection in Caenorhabditis nematodes. *BMC Genomics*
18, 303 (2017).

29.   Le Pen, J. *et al.* Terminal uridylyltransferases target RNA viruses as part of the innate
immune system. *Nat Struct Mol Biol* **25**, 778–786 (2018).

30.   Tan, M.-W., Mahajan-Miklos, S. & Ausubel, F. M. Killing of *Caenorhabditis elegans* by
Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proceedings*
of the National Academy of Sciences **96**, 715–720 (1999).

31.   Crone, S. *et al.* The environmental occurrence of *Pseudomonas aeruginosa*. *APMIS* **128**,
220–231 (2020).

32.   Pelegrin, A. C. *et al. Pseudomonas aeruginosa* : a clinical and genomics update. *FEMS*
Microbiol Rev **45**, (2021).

33.   Kim, D. H. *et al.* A Conserved p38 MAP Kinase Pathway in *Caenorhabditis elegans* Innate
Immunity. *Science (1979)* **297**, 623–626 (2002).

34.   Kim, D. H. & Ewbank, J. J. Signaling in the innate immune response. *WormBook* 1–35
(2018) doi:10.1895/wormbook.1.83.2.

35.   Zhang, Y., Lu, H. & Bargmann, C. I. Pathogenic bacteria induce aversive olfactory learning
in Caenorhabditis elegans. *Nature* **438**, 179–184 (2005).

36.   Dirksen, P. *et al.* The native microbiome of the nematode Caenorhabditis elegans:
gateway to a new host-microbiome model. *BMC Biol* **14**, 38 (2016).

37.   Samuel, B. S., Rowedder, H., Braendle, C., Félix, M.-A. & Ruvkun, G. *Caenorhabditis*
elegans responses to bacteria from its natural habitats. *Proceedings of the National*
Academy of Sciences **113**, (2016).

38.   Berg, M. *et al.* Assembly of the Caenorhabditis elegans gut microbiota from diverse soil
microbial environments. *ISME J* **10**, 1998–2009 (2016).

39.   Dirksen, P. *et al.* CeMbio - The *Caenorhabditis elegans* Microbiome Resource. *G3*
Genes|Genomes|Genetics **10**, 3025–3039 (2020).

40. Yuan, W., Zhou, Y., Fan, Y., Tao, Y. J. & Zhong, W. Orsay δ Protein Is Required for Nonlytic
Viral Egress. *J Virol* **92**, (2018).

41. Brenner, S. THE GENETICS OF *CAENORHABDITIS ELEGANS*. *Genetics* **77**, 71–94 (1974).

42. Mahajan-Miklos, S., Tan, M.-W., Rahme, L. G. & Ausubel, F. M. Molecular Mechanisms of
Bacterial Virulence Elucidated Using a Pseudomonas aeruginosa– Caenorhabditis elegans
Pathogenesis Model. *Cell* **96**, 47–56 (1999).

43. Jiang, H., Chen, K., Sandoval, L. E., Leung, C. & Wang, D. An Evolutionarily Conserved
Pathway Essential for Orsay Virus Infection of *Caenorhabditis elegans*. *mBio* **8**, (2017).

44. Lee, J. & Zhang, L. The hierarchy quorum sensing network in Pseudomonas aeruginosa.
Protein Cell **6**, 26–41 (2015).

45. Reimmann, C. *et al.* The global activator GacA of *Pseudomonas aeruginosa* PAO positively
controls the production of the autoinducer *N* -butyryl-homoserine lactone and the
formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol Microbiol* **24**, 309–
319 (1997).

46. Parkins, M. D., Ceri, H. & Storey, D. G. Pseudomonas aeruginosa GacA, a factor in
multihost virulence, is also essential for biofilm formation. *Mol Microbiol* **40**, 1215–1226
(2001).

47. Brencic, A. *et al.* The GacS/GacA signal transduction system of *Pseudomonas aeruginosa*
acts exclusively through its control over the transcription of the RsmY and RsmZ
regulatory small RNAs. *Mol Microbiol* **73**, 434–445 (2009).

48. Friedman, L. & Kolter, R. Two Genetic Loci Produce Distinct Carbohydrate-Rich Structural
Components of the *Pseudomonas aeruginosa* Biofilm Matrix. *J Bacteriol* **186**, 4457–4465
(2004).

49. Friedman, L. & Kolter, R. Genes involved in matrix formation in Pseudomonas aeruginosa
PA14 biofilms. *Mol Microbiol* **51**, 675–690 (2003).

50. Franklin, M. J., Nivens, D. E., Weadge, J. T. & Howell, P. L. Biosynthesis of the
Pseudomonas aeruginosa Extracellular Polysaccharides, Alginate, Pel, and Psl. *Front*
Microbiol **2**, (2011).

51. Jackson, K. D., Starkey, M., Kremer, S., Parsek, M. R. & Wozniak, D. J. Identification of *psl* ,
a Locus Encoding a Potential Exopolysaccharide That Is Essential for *Pseudomonas*
aeruginosa PAO1 Biofilm Formation. *J Bacteriol* **186**, 4466–4475 (2004).

52. Matsukawa, M. & Greenberg, E. P. Putative Exopolysaccharide Synthesis Genes Influence
Pseudomonas aeruginosa Biofilm Development. *J Bacteriol* **186**, 4449–4456 (2004).

53. Mukherjee, S. *et al.* The PqsE and RhlR proteins are an autoinducer synthase–receptor
pair that control virulence and biofilm development in *Pseudomonas aeruginosa*.
Proceedings of the National Academy of Sciences **115**, (2018).

54. Xu, L. *et al.* OrthoVenn2: a web server for whole-genome comparison and annotation of
orthologous clusters across multiple species. *Nucleic Acids Res* **47**, W52–W58 (2019).

55. Liberati, N. T. *et al.* An ordered, nonredundant library of *Pseudomonas aeruginosa* strain
PA14 transposon insertion mutants. *Proceedings of the National Academy of Sciences*
103, 2833–2838 (2006).

56. Simanek, K. A. *et al.* The PqsE-RhlR Interaction Regulates RhlR DNA Binding to Control
Virulence Factor Production in *Pseudomonas aeruginosa*. *Microbiol Spectr* **10**, (2022).

57. Feinbaum, R. L. *et al.* Genome-Wide Identification of Pseudomonas aeruginosa Virulence-
Related Genes Using a Caenorhabditis elegans Infection Model. *PLoS Pathog* **8**, e1002813
(2012).
58. Sun, J. *et al.* OrthoVenn3: an integrated platform for exploring and visualizing
orthologous data across genomes. *Nucleic Acids Res* **51**, W397–W403 (2023).
59. González, R. & Félix, M.-A. Natural monobacterial environments modulate viral infection
in Caenorhabditis elegans. *bioRxiv* (2023).
60. Kaul, D. *et al.* Microbiome disturbance and resilience dynamics of the upper respiratory
tract during influenza A virus infection. *Nat Commun* **11**, 2537 (2020).
61. Kissoyan, K. A. B. *et al.* Exploring Effects of C. elegans Protective Natural Microbiota on
Host Physiology. *Front Cell Infect Microbiol* **12**, (2022).
62. Rahme, L. G. *et al.* Common Virulence Factors for Bacterial Pathogenicity in Plants and
Animals. *Science (1979)* **268**, 1899–1902 (1995).
63. Lapouge, K., Schubert, M., Allain, F. H.-T. & Haas, D. Gac/Rsm signal transduction
pathway of γ-proteobacteria: from RNA recognition to regulation of social behaviour.
Mol Microbiol **67**, 241–253 (2007).
64. Shtonda, B. B. & Avery, L. Dietary choice behavior in *Caenorhabditis elegans*. *Journal of
Experimental Biology* **209**, 89–102 (2006).
65. Hmelo, L. R. *et al.* Precision-engineering the Pseudomonas aeruginosa genome with two-
step allelic exchange. *Nat Protoc* **10**, 1820–1841 (2015).
66. Rietsch, A., Vallet-Gely, I., Dove, S. L. & Mekalanos, J. J. ExsE, a secreted regulator of type
III secretion genes in *Pseudomonas aeruginosa*. *Proceedings of the National Academy of
Sciences* **102**, 8006–8011 (2005).
67. R Core Team (2022). R: A language and environment for statistical computing. Preprint at
(2022).
68. Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-Response Analysis Using R. *PLOS ONE*
10, (2015).
69. Warnes, G. et al. gdata: Various R Programming Tools for Data Manipulation. Preprint at
(2022).
70. Wickham H & Seidel D. scales: Scale Functions for Visualization. Preprint at (2022).
71. Hope RM. Rmisc: Ryan Miscellaneous. Preprint at (2022).
72. Hothorn, T., Bretz, F. & Westfall, P. Simultaneous Inference in General Parametric
Models. *Biometrical Journal* **50**, 346–363 (2008).
73. Wickham H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, 2016).
74. Ahlmann-Eltze, C. & Patil, I. ggsignif: R Package for Displaying Significance Brackets for
'ggplot2'. *PsyArxiv* (2021).
75. Wilke C. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. Preprint at
(2020).
737