TGF-β blockade drives a transitional effector phenotype in T cells reversing SIV latency

and decreasing SIV reservoirs in vivo

Jinhee Kim[1], Deepanwita Bose[2]†, Mariluz Araínga[2]†, Muhammad R. Haque[3], Christine M

Fennessey[4], Rachel A Caddell[5], Yanique Thomas[3], Douglas E Ferrell[2], Syed Ali[2], Emanuelle

Grody[3,9], Yogesh Goyal[3,9,10], Claudia Cicala[6], James Arthos[6], Brandon F Keele[4], Monica

Vaccari[5,7], Ramon Lorenzo-Redondo[1,8], Thomas J Hope[3], Francois Villinger[2] and Elena

Martinelli[1]

[1]Department of Medicine, Division of Infectious Diseases, Feinberg School of Medicine,
Northwestern University, Chicago, IL, USA
[2]New Iberia Research Center, University of Louisiana at Lafayette, New Iberia, LA, USA
[3]Cell and Developmental Biology, Feinberg School of Medicine, Northwestern University,
Chicago, IL, USA
[4]AIDS and Cancer Virus Program, Frederick National Laboratory for Cancer Research,
Frederick MD
[5]Division of Immunology, Tulane National Primate Research Center, Covington LA, USA
[6]Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National
Institutes of Health, Bethesda, MD, USA
[7]Department of Microbiology and Immunology, Tulane University School of Medicine, New
Orleans, LA, USA
[8]Center for Pathogen Genomics and Microbial Evolution, Northwestern University Havey
Institute for Global Health, Chicago, IL, USA.
[9]Center for Synthetic Biology, Northwestern University, Chicago, IL, USA
[10]Robert H. Lurie Comprehensive Cancer Center, Feinberg School of Medicine, Northwestern
University, Chicago, IL, USA
27
28
*Corresponding author:

Elena Martinelli

elena.martinelli@northwestern.edu

One-sentence summary:

TGF-β blockade drives an effector phenotype in immune cells leading to SIV latency reversal

and enhanced immune responses in vivo.

Abstract

HIV-1 persistence during ART is due to the establishment of long-lived viral reservoirs in resting immune cells. Using an NHP model of barcoded SIVmac239 intravenous infection and therapeutic dosing of the anti-TGFBR1 inhibitor galunisertib (LY2157299), we confirmed the latency reversal properties of *in vivo* TGF-β blockade, decreased viral reservoirs and stimulated immune responses. Eight SIV-infected macaques on suppressive ART were treated with 4 2-week cycles of galunisertib. ART was discontinued 3 weeks after the last dose, and macaques euthanized 6 weeks after ART-interruption(ATI). One macaque did not rebound, while the remaining rebounded between week 2 and 6 post-ATI. Galunisertib led to viral reactivation as indicated by plasma viral load and immunoPET/CT with the ^{64}Cu-DOTA-F(ab')$_2$-p7D3-probe. Half to 1 Log decrease in cell-associated (CA-)SIV DNA was detected in lymph nodes, gut and PBMC, while intact pro-virus in PBMC decreased by 3-fold. No systemic increase in inflammatory cytokines was observed. High-dimensions cytometry, bulk and single-cell RNAseq revealed a shift toward an effector phenotype in T and NK cells.

In summary, we demonstrated that galunisertib, a clinical stage TGF-β inhibitor, reverses SIV latency and decreases SIV reservoirs by driving T cells toward an effector phenotype, enhancing immune responses *in vivo* in absence of toxicity.

Introduction

Interruption of antiretroviral therapy (ART) leads to rapid rebound of viremia in the vast majority of people living with HIV-1 (PLWH) due to the establishment of a persistent HIV-1 reservoir early after infection(*1, 2*). A key mechanism of this persistence is the ability of HIV-1 to enter a state of virological latency characterized by the silencing of viral gene expression and/or lack of viral proteins translation(*3, 4*). This allows the virus to remain invisible to the immune system and latently infected cells to survive and proliferate by homeostatic or antigen-driven proliferation(*5, 6*). Of note, the viral reservoir was initially thought to be stable. However, recent evidence suggests that stochastic HIV reactivation under ART occurs, and selective killing is favored in cells bearing replication competent virus integrated in transcriptionally active sites within the genome(*7-9*). Hence, integration site, but especially the activation status of the infected cells profoundly influences HIV-1 persistence.

Ongoing efforts to achieve a functional cure for HIV-1 are directed towards supplementing ART with immunotherapies targeting the viral reservoir. Such strategies, under the umbrella of "shock and kill", aim to reactivate the replication competent reservoir and eliminate latently infected cells by viral cytopathic effects or immune-mediated killing (*10, 11*). However, these strategies have thus far failed to achieve a reduction of the viral reservoir or post-ART virologic control in either the clinic or preclinical models. This is mainly due to the low inducibility of latent proviruses and the heterogeneity of the mechanisms of persistence(*12, 13*) (*14, 15*). In contrast, therapeutic vaccination strategies focused on enhancing HIV/SIV-specific responses have had some discreet measure of success(*16*). However, currently, no single strategy leads simultaneously to latency reversal and stimulation of effective immune responses.

The HIV life cycle and HIV's ability to replicate efficiently are especially dependent on the activation status of the infected cells. In this context, significant advances have been made to directly activate immune cells through non-canonical pathways in order to promote HIV latency reversal(*17*). However, the activation and differentiation of immune cells is intrinsically linked to

cellular metabolism(*18*). Indeed, signaling pathways that govern immune cell differentiation and

activation such as mTOR and Wnt/β-catenin, not only have been implicated in regulating HIV

latency(*19-21*), but are also critical regulators of cell metabolism(*22*). The metabolic status of an

infected cell, in turn, plays a critical role in its ability to support latency reactivation and viral

production(*19, 23*).

In this context, recent evidence suggests that TGF-β plays a critical role in the regulation of

immune cell activation and metabolic reprogramming(*24-26*). Specifically, in the context of CD8[+]

T cells, TGF-β has been shown to suppress mTOR signaling preserving the metabolic fitness of

memory CD8[+] T cells(*25*) and stem-like antigen specific CD8[+] T cells through its modulation of

Wnt/β-catenin factor TCF1(*26, 27*). TCF1 and the mTOR pathway are also critical to T cell

differentiation and responsible for the transition from activated effector cells to resting memory

cells during LCMV infection(*22, 26*). The regulation of quiescence in CD8[+] T cells that follows

continuous TGF-β stimulation is critical to the transition to a memory phenotype and it is driven

by specific metabolic changes that are linked to decreased glycolytic activity, more efficient

mitochondrial respiration, and long-term survival(*25, 26*).

Similarly, in the context of NK cells, TGF-β has been implicated in decreasing their baseline

metabolism driving lower expression of markers of NK cytotoxic activity(*28*).

While TGF-β-mediated suppression of TCR and IL-2 signaling were shown to lead to lower

CD4[+] T cell activation following cognate antigen recognition in older studies(*24, 29*), more

recent work using CD4[+] T cell-specific deletion of the TGF-β receptor demonstrated an even

more profound effect of TGF-β on all stages of CD4[+] T cell activation, proliferation and cytotoxic

response to LCMV than in CD8[+] T cells(*30*). However, little is known on the specific role of

TGF-β in regulating the transition to and from memory and effector phenotypes in CD4[+] T cells

and how this may be associated with TGF-β-driven changes in CD4[+] T cells metabolism.

Moreover, TGF-β regulates the expression of CD103 and other surface and intracellular factors

essential of T cell residency in mucosal tissues(*31-33*). Hence, TGF-β is considered the master

regulator of mucosal immunity(*34*).

We and others have recently demonstrated that TGF-β regulates HIV-1 latency in primary CD4$^+$

T cells ex vivo and in vivo(*35-37*). Latency reversal was detected in a non-human primate (NHP)

model of HIV infection following a short treatment with a clinical stage TGF-β inhibitor,

galunisertib (LY2157299)(*38*). In that previous study, we documented latency reversal

particularly at the level of the gut mucosal tissue using the ^{64}Cu-anti-gp120 Fab$_2$(7D3) probe

and immuno-PET/CT(*35*). In that study, we further validated the ability of immune-PET/CT to

identify sites of viral reactivation and replication in the gut by performing tissue resection in hot

areas of the gut identified by PET followed by confirmatory PCR for vDNA/RNA and

vRNAscope(*35*).

Herein, we demonstrate how treatment with galunisertib with a 2-week on, 2-week off regiment

that mimics the therapeutic regimen employed in the clinic in phase 1 and 2 trials of solid

tumors(*39, 40*), leads to profound transcriptional and functional changes in immune cells in the

absence of overt toxicity or increased systemic inflammation. Importantly, we observed a shift

toward a transitional effector phenotype in CD4$^+$ T cells and other immune cells both

systemically and in the lymph nodes. This shift was accompanied by, and likely responsible for,

increased viral reactivation in SIV-infected, ART treated macaques documented by molecular

techniques and PET/CT images. At the end of the treatment with galunisertib, we detected lower

viral reservoir levels, including total and intact proviral DNA in both PBMC, gut and lymph nodes

and significantly higher immune responses.

RESULTS

2-weeks on-off therapeutic regiment with Galunisertib leads to viral reactivation SIV

infected, ART-treated macaques.

To confirm galunisertib-driven HIV/SIV latency reversal and investigate the underlying

mechanisms, 8 Indian origin rhesus macaques (*Macaca Mulatta*, Mamu-A01-, -B08, -B17-, all

females) were infected intravenously with 300 TCID50 of the barcoded SIVmac239M2. We

initiated ART treatment (daily co-formulated Tenofovir [PMPA], Emtricitabine [FTC] and

Dolutegravir [DTG]) on week 6 post-infection (pi). A 2-week on, 2-week off therapeutic cycle with

galunisertib (20mg/Kg twice/daily orally) started at week ~35pi and continued for a total of 4

cycles (Fig 1A and Table S1). ART was discontinued 3 weeks after the last galunisertib dose,

and the macaques were followed for 6 weeks after ART discontinuation. Median peak plasma

viral load (pVL) was 10^8 copies/ml at week 2pi. A rhesus anti-PD1 antibody was administered at

5mg/kg before the 3rd and 4th cycle to 2 macaques (08M156 and A6X003). However, no

differences were noted for these 2 macaques in any of the parameters we measured, and the

data were pooled.

Full suppression to undetectable levels (pVL LOD 15 copies/ml) was achieved in 3 out of the 8

macaques at week 10pi. In the other 5 macaques, pVL fell below 65 copies/ml by week 22pi

with a single blip of 400 copies/ml in A8T010 at week 29pi (Fig 1B). Following the start of the

galunisertib treatment, pVLs increased in 7 out of 8 macaques from a single peak over

undetectable in 08M171 and A8L057 to several peaks and up to 10^3 copies/ml in the other

macaques. Of note pVLs in A8T010 and A8R095 was undetectable for over 5 weeks before,

respectively, blips of up to 10^2 copies/ml were detected following galunisertib treatment initiation

(Fig 1B). More frequent blips were noted during the first 2 cycles with galunisertib compared to

cycles 3 and 4 (Fig 1C). However, 08M171 and A8L057 did not experience a pVL increase until

the 4th cycle.

Importantly, we documented viral reactivation also using immunoPET/CT. The ^{64}Cu-anti-gp120

Fab$_2$(7D3) probe was injected 24hrs before and after, respectively, the beginning and the end of

the first 3 galunisertib cycles and scans performed before and after the first and the last dose.

As shown in Fig 2, Fig S1 and Movies S1-S8, the PET signal visibly increased in different areas

after cycle 2 (in A8R095, 08M156, A8L014 and A8T010) or at the beginning of cycle 3 (in A6X003, 08M134 and A8L057). In 08M171 we observed an increase in the gut area only at the beginning of cycle 2. A corresponding increase in standard uptake values (SUV) was detected in the gastrointestinal area and axillary lymph nodes (Fig 2B) and was significant in cycle 3 compared to before cycle 1 (BC1). In these anatomical areas, SUV increases correspond to increases in viral replication as shown in previous studies(*41, 42*). A PET signal increase was also noted in the area of the vertebral column (spine). Spine SUV already increased significantly with the first cycle (Fig 2B). However, neither cerebrospinal fluid (CSF) nor bone marrow (BM) was collected during the study, and we have no prior validation of the specificity of the signal in the spine. Hence, whether this signal corresponds to increased viral replication and whether this occurs in the vertebral bones or cerebrospinal fluids remains to be determined. No SUV increase was present at the level of the spleen or kidney, where probe accumulation and background signal likely masked any specific signal. However, a significant increase in SUV was detected in the liver (Fig 2B) and whole-body (Fig S2). Of note, the SUV increase in the gut area in most cases followed an increase in cell-associated vRNA detected in colorectal biopsies at the same time points during treatment (Fig S3).

A 4 cycle treatment with galunisertib decreases viral reservoir in absence of systemic inflammation.

To determine the impact of galunsiertib on SIV reservoir, we measured CA-vDNA in PBMC, colorectal biopsies and lymph nodes (LN). A significant decrease in CA-vDNA was detected in all tissues between week 35pi (beginning of cycle 1, BC1) and week 49pi (after/end of cycle 4, AC4) for gut and LN, and between week 35pi (BC1) and end of cycle 3 (AC3) for PBMC (Fig 3A, AC4 not measured for PBMC and Fig S4A). In the gut and LN (right axillary), decreases ranged from a Log to 1/3 of a Log (gut: median 0.77; range:0.33-0.94 folds LN: median: 0.93; range:0.33-0.98 folds decrease). In the PBMC the decrease was slightly less pronounced with a

median half Log decrease (median: 0.58; range 0.28-0.88 folds decrease). However, the

comparison for the PBMCs was between week ~35 and week ~45 (end of cycle 3, AC3) instead

of the end of all 4 cycles, because a snap frozen pellet was not available at the end of cycle 4

for PBMCs. The SIV reservoir in PBMCs was also monitored by SIV-IPDA comparing before

cycle 1 (BC1) to the end of cycle 4 (AC4). Of note, we observed significant decreases of both

total and intact provirus by SIV-IPDA. Intact provirus declined similarly to the CA-vDNA with a

median of half Log (median: 0.53; range:0-0.71 folds decrease).

In contrast, no decline in CA-vDNA was detected in the PBMCs of a group of 4 macaques

infected intravenously with the same stock of SIVmac239M2 for a separate study. These 4

macaques were placed on ART on week 6pi as in our study. However, no decline in CA-vDNA

was detected under ART, between weeks 28 and 52pi (untreated group, Fig S4B). Thus, we

conclude that the decline in CA-vDNA in PBMC was not due to ART-alone.

Importantly, we found no significant changes in any of the clinical variables (chemistry and

hematology, Tables S2 and S3) measured before, during and after the galunisertib therapy.

Moreover, we observed no changes in the concentrations of inflammatory chemokines and

cytokines measured in plasma before and after the first 2 treatment cycles and at the end of the

last cycle (Fig 3C). The only difference in cytokine and chemokine levels after galunisertib

treatment was a small increase in IL-10 detected at the end of the last cycle compared to before

treatment (Fig S4C).

Galunisertib treatment drives an effector phenotype in T and NK cells.

The phenotype of PBMCs before and after Galunisertib treatment was monitored by high-

parameter flow cytometry of T and NK cell subsets and phenotype (Table S4). Classical subsets

and single-color analysis of MFI revealed a substantial increase in the expression of CD95 and

a profound consistent decrease in TCF1 expression in CD4$^+$ T cells that continued throughout

the treatment (Fig 4A). In contrast, a small decrease in CD62L after the first cycle, reverted to

baseline during the following cycles. The frequency of naïve cells, defined as $CD95^-$ within $CD4^+$

T cells, decreased in parallel with the increase in CD95 (gating strategy in Fig S5). Interestingly,

there was no change in the expression of CCR7 or CD28 within $CD95^+$ $CD4^+$ T cells (Fig 4B).

Hence, the frequency of central memory and effector memory as defined by CD95 and CD28 or

CCR7 did not change (Table S5). The levels of T-bet did not change significantly (Fig S6A).

However, we detected a downregulation of the gut homing receptor integrin $\alpha4\beta7$ and a

decrease in the levels of granzyme B (GRZB, Fig S6A). Of note, the expression of activation

markers CD69, HLA-DR and Ki67 remained mostly unchanged, with the exception of an

increase in HLA-DR at the beginning of cycle 2 compared to before galunisertib (Fig 4C).

Interestingly, the effect of Galunisertib on $CD8^+$ T cells was not as pronounced as it was on

$CD4^+$ T cells. No significant increase was detected in CD95 expression and TCF1

downregulation reached significance only at the end of the treatment (Fig 4D). Markers of cell

activation like HLA-DR, Ki67 and CD69 did not change (Fig 4D and Fig S6B). However, we

detected a significant decrease in T-Bet at the beginning of cycle 2 (BC2) compared to before

treatment (BC1). Finally, in contrast to $CD4^+$ T cells, the decrease in GRZB was more

pronounced at several time points during treatment and there was a sustained significant

increase in CCR7 expression in memory $CD8^+$ T cells (Fig 4E). Interestingly, the expression of

PD1 on both $CD4^+$ T cells and $CD8^+$ T cells either did not change or was slightly downregulated.

Importantly, the frequency of $PD1^+$ $CD101^+$ ($TCF1^{low}$) exhausted memory $CD8^+$ T cells(*43*) did

not change (Fig S6B). Finally, within $NKG2A^+$ $CD8^+$ NK cells, there was a pronounced increase

in CD16 expression and an initial increase in the proliferation marker Ki67 (AC1 vs BC1; Fig

4F).

High-dimensional data visualization with tSNE and clustering analysis with FlowSOM(*44*)

confirmed the results of the classical analysis. We performed tSNE and FlowSOM after data

clean up with FlowClean and normalization with the SwiftReg algorithm(*45*). The analysis was

performed on before (BC1) and after the last cycle (AC4) only (Fig 4G and Fig S7A) and on all

time points analyzed (Fig S7B and S7C). When we compared only BC1 and AC4, the

PhenoGraph clustering algorithm identified 36 populations. FlowSOM with 36 populations

identified 6 populations of CD4$^+$ T cells, 6 of CD8$^+$ T cells and 4 of NK cells (NKG2A^{high} CD8$^+$;

Fig S7A). The remaining populations were likely monocytes and other minor subsets. Direct

comparison of each population revealed a decrease in Pop 31 (naïve CD8 T cells) and Pop 30

(central memory Ki67$^+$ CD8$^+$ T cells) and an increase in Pop 2 and 3 (effector and central

memory CD4$^+$ T cells). Finally, there was a decrease in HLA-DR high Pop0 and an increase in

CD16^{high} NKG2A$^-$ Pop14 (Fig 4H and S7A). Visual inspection of the tSNE plots (Fig 4G)

revealed 3 areas mostly occupied by cells in the post-galunisertib AC4 group (New1, 2 and 3)

which were characterized by high levels of CD16 and GRZB (New 3 is likely NK cells). In

contrast, 3 areas mostly occupied by cells in the pre-galunisertib group BC1 (Old1, 2 and 3)

were characterized by high levels of TCF1 and low CD95, confirming the finding of the classical

analysis. Analysis of all time points with phenograph-derived 38 populations in FlowSOM

recapitulated findings obtained with the classical analysis and the BC1 AC4 comparison with no

additional insights (Fig 4I).

Galunisertib treatment in vivo increases pTreg while decreasing pTfh

The frequencies of circulating Tregs and Tfh cells were monitored with an established flow

cytometry panel(*46, 47*) after the first and third cycle of galunisertib. The frequency of all CD4$^+$

Tregs (CD25^{high} FoxP3$^+$; gating in Fig S8) and CD8$^+$ Tregs increased after the first cycle and

remained higher until the end of the 3rd cycle, while CCR4$^+$ Treg were proportionally higher at

the end of the treatment compared to before (Fig 4J). In contrast, circulating Tfh (CXCR5$^+$ PD1$^+$;

gating in Fig S8) were lower at the end of the Galunisertib treatment both within total and central

memory CD4$^+$ T cells (Fig 4J). Finally, the expression of CD39 (ecto-nucleotide triphosphate

diphosphohydrolase 1), which tracks within extracellular adenosine and immunosuppressive

effects, was lower on total and central memory CD4$^+$ T cells (Table S5 and Fig 4J, respectively).

Bulk RNAseq of PBMC and single-cell (sc)RNAseq analysis of LN confirm a profound

shift toward effector phenotype.

In our previous studies, we determined that 6hrs after galunisertib treatment in naïve macaques

there was an upregulation of the AP1 complex (JUN and FOS) and several genes encoding

ribosomal proteins in CD4$^+$ T cells(*42*). To understand early and later effect of galunisertib in the

context of SIV infection, in the current study we performed bulk RNAseq of PBMC isolated 1hr

after the first administration of galunisertib in cycle 1 and at the end of the 1st 2-weeks cycle. We

found 640 genes significantly modulated (FDR q<0.05; log$_2$FC=2) in PBMC just 1hr after the first

dose of galunisertib. The majority (457 genes) were downregulated (Fig 5A). Gene set

enrichment analysis (GSEA) revealed an upregulation of the oxidative phosphorylation

(OXPHOS; ES 0.58 FDR q=0.048)) and the reactive oxygen (ES 0.55 FDR q=0.075) pathways

among the Hallmark genet sets (Fig 5A and S9). Among the Biocarta sets, there was an

enrichment in the electron transport chain (ETC; ES 0.84 FDR q=0.132) and a downregulation

of the circadian (ES -0.81 FDR q=0.163) pathways (Fig 5A and S9). Finally, among the top

modulated genes there were several genes encoding for soluble transporters, while classical

activation markers like CD69 and CD38 were downregulated (Fig S9B). An early engagement of

metabolic pathways was confirmed by enrichment analysis of significant DEG with

Metascape(*48*) with an upregulation of adipogenesis, OXPHOS and fatty acid metabolism (Fig

S9C). Of note, 2 weeks after the beginning of galunisertib, metabolic pathways were still among

the most enriched upregulated pathways in PBMCs (Fig 5B and Fig S10A). Among the top

upregulated genes there were CD44, CCR5, several integrins and GRZA and GRZB (Fig S10B).

Finally, we performed bulk RNAseq of rectal biopsy tissue before and after the first cycle with

galunisertib (Fig 5C). There were only 51 differentially expressed genes (DEGs; Adjp<0.05;

log$_2$FC=2). GSEA analysis of all DEGs (Adjp<0.05) revealed the G2_M_DNA replication

pathway highly enriched (ES 0.97 FDR q=0.012) within the Hallmark set. Among the most

interesting changes, we observed a pronounced downregulation of integrin β7, in contrast to an

increase in integrin αE (CD103; Fig 6C), suggesting that TGF-β-driven increase in αE(*49*) may

be driven by non-canonical TGF-β pathway signaling not blocked by galunisertib.

To clarify the impact of galunisertib at the single cell level and in lymphoid tissues, we also

performed scRNAseq analysis of cells isolated from LNs before (right axillar) and after (right or

left inguinal) the first cycle (Fig 6). Dimensionality reduction and clustering analysis with PCA

and uniform manifold approximation and projection (UMAP)(*50*) was performed to visualize the

data (Fig 6A). However, cells were classified based only on gene expression (Fig 6B and C).

We first classified major subsets: T cells (38,705 cells), B cells (31,385 cells), NK cells (1,628

cells), macrophages (285) and cells expressing both CD19 and CD3 (942 T/B cells) (Fig 6A

shows this classification over UMAP). Then we classified only CD4$^+$ and CD8$^+$ T cell subsets

(Fig 6C and S11A) and B/T subsets including naïve and germinal center (GC) B cells and Tfh

cells (Fig S11B and S11C). Cell number for all these subsets did not change with treatment (Fig

S12). However, differential gene expression analysis of T cells revealed an upregulation of

members of the AP1 complex, CD69, β2 macroglobulin and RPL13 among the most

upregulated genes (Fig 6D). Moreover, it confirmed a downregulation of TCF1 at the

transcriptional level (TCF7 gene; Fig 6D). Gene enrichment analysis revealed again

Myc_targets_V1, OXPHOS and mTORC1 as the most enriched hallmark pathways (Fig 6E). Of

note, among gene ontology cellular processes, RNA processing was the most highly enriched

pathway with intracellular transport and catabolic processes following right after (Fig S13A)

confirming an increase in translation and metabolism within these cells.

Since we noticed substantial differences in the impact of galunisertib on CD4$^+$ T cells compared

to CD8$^+$ T cells by flow, we focused the analysis on these subsets. In CD4$^+$ T cells we found

only 25 DEGs with a log$_2$FC=0.15, while 34 DEGs were in CD8$^+$ T cells with more

downregulated genes in the CD8$^+$ T cells compared to the CD4$^+$ T cells. The AP1 complex and

TCF7 were again upregulated and downregulated respectively in both CD4$^+$ (Fig 6F) and CD8$^+$

T cells (Fig 6G). However, STAT1 was more strongly downregulated in CD4$^+$ T cells. Enrichment

analysis showed once again upregulation of Myc_targets_V1, OXPHOS and mTORC1

pathways in both CD4$^+$ and CD8$^+$ T cells (Fig 6H and Fig S13B). Interestingly, the most enriched

set among in Biocarta was the HIV-Nef pathway (Fig 6J) demonstrating the relevance of these

galunisertib-driven changes to HIV cell cycle and transcription (highly enriched KREB pathway

as well). The 2nd most enriched Biocarta pathway was TCR signaling, linking galunisertib to cell

activation. Next, we analyzed changes in gene expression in Tfh cells. In this subset we

obtained a similar number of DEG as in other T cells and myc_targets_V1 was still the most

enriched hallmark pathway (Fig S13C and D). In B cells, the AP1 complex was again

prominently upregulated, together with CD83, CD69 and MAMU-DR. Of note, more genes were

modulated in B cells (44 genes with a log$_2$FC=0.15 and 18 with logFC0.2) than in T cells with

some differences in enriched pathways (Fig S13F). Similar genes were modulated in GC B cells

and naïve B cells with several more DEGs in naïve B cells than in GC cells (Fig S14A and B).

Finally, 144 genes were modulated in NK cells and 39 genes in macrophages (Fig S14C and D;

log$_2$FC=0.15). GRZB was prominently upregulated, but CD44 downregulated. In macrophages

genes were mostly downregulated including TCF7L2 and KLF4 suggesting an increase in

inflammatory phenotype and decrease in M2 polarization(*51*) (Fig S14D).

Galunisertib increases SIV-specific responses and changes barcode distribution

In order to understand how galunisertib affected immune cell function and SIV-specific

responses, we stimulated PBMC with 15-mer SIV peptides from SIVmac239 gag, pol and env

for 24hrs on antibody coated Elispot plates. Because of sample availability, we probed before

and after the first cycle and after the 3rd galunisertib cycle only. Interestingly, by the end of the

3rd cycle there was a significant increase in IFN-γ secretion both SIV-specific (particularly

against env) and non-specific (DMSO control). TNF-α release increased particularly in response

to Gag and non-specifically (Fig 7). However, IL-2 release appeared to increase slightly after the

first cycle (non-significant), but remained unchanged with a slight decrease by the end of the 3rd

cycle (Fig 7).

In order to understand if these changes in immune responses in combination with latency

reversal and switch toward an effector phenotype may have impacted viral population dynamics,

we analyzed changes in numbers and distribution of the viral barcodes before and after the first

3 treatment cycles. There was no barcode amplification at several time points particularly in the

lymph nodes. However, although there were no significant changes in the number or diversity

(measured as Shannon Entropy; Sh) of barcodes before and after each galunisertib cycle (Fig

S15A and Fig 8A), we found significant changes in barcode frequency distribution in most

tissues and cycles after galunisertib treatment (Fig 8B and S15B). Specifically, the relative

proportion of each barcode changed in all monkeys in all cycles in the rectal biopsies (probably

a consequence of different sampling area), but also for all LN analyzed (except for 08M134 in

cycle 1). Of note, the same LN were sampled at the beginning and after cycle 3 (Table S6).

Hence, sampling location does not explain the changes barcode distribution. Changes in the

proportion of barcodes were also detected in at least half of the PBMCs after cycle 1 and 2

(cycle 3 not analyzed). Finally, barcode diversity decreased in plasma post-ATI compared to the

time point right before ART (Fig 8C) and barcode distribution significantly changed post-ATI

compared to pre-ART in 2 of the 7 macaques that rebounded (Fig S15C).

TCF1 downregulation in CD4$^+$ T cells correlates with virological and immunological endpoints.

In order to explore a possible association between the various virological and immunological parameters and their changes, we built a correlation matrix with a curated set of variables of interest or their fold changes. This analysis revealed an association between the decrease in TCF1 expression in the CD4$^+$ T cells and several virological and immunological variables (Fig 8E). Specifically, both the levels of CA-vDNA in the colorectal tissue and the fold increase in gut-SUV strongly inversely correlated with fold changes in TCF1. Since TCF1 decreased, a larger decrease in TCF1 was directly proportional to residual vDNA in the gut at the end of cycle 4 and to the increase in PET signal in the gut (Fig 8D and 8E). A weaker, but still significant association was also present with CA-vDNA in PBMC at the end of cycle 3 and with the levels of vRNA in the lymph nodes at the end of the 4th cycle (Fig 8D and S16A). Finally, the decrease in TCF1 correlated with the increase in IFN-γ and poly-functional IFN-γ/TNF-α releasing cells (cumulative increase of SIV-specific responses to gag, pol and env; Fig 8D and 8E). Of note, the increase in PET signal in the gut also correlated with the levels of vDNA in PBMC at the end of cycle 3, vDNA in gut biopsies and residual CA-vRNA in LN at the end of cycle 4 (Fig S16B). Interestingly, the levels of residual CA-vRNA in LN also correlated with the increase in IFN-γ and poly-functional IFN-γ/TNF-α releasing cells (Pearson r=0.85 and 0.76, respectively). Finally, the residual intact pro-virus (IPDA) directly correlated with the levels of IFN-γ and IFN-γ/TNF-α produced in response to SIV peptides at the end of cycle 3 (Fig 8D; r= 0.76 and 0.88, respectively). In contrast, the change in intact provirus trented to correlate inversely with the levels of IFN-γ, so that a larger decrease directly correlated with more IFN–γ responses. However, this did not reach significance (p=0.117, Fig S16C).

Discussion

HIV-1 latency in T cells is maintained through diverse mechanisms that include blocks in

transcriptional elongation, completion, and splicing(*13*). A common characteristic shared by HIV-

1 latently infected cells of both T and myeloid cell lineages is their "resting" phenotype(*14, 52-

54*) In these cells, an inability to transcribe proviral DNA is linked to a generalized decrease in

transcriptional activity which, in turn, is linked to their metabolic status(*23*). Cellular metabolism

is in turn influenced by tissue location and environmental cues(*55*).

TGF-β is released at high level in PLWH and its levels remain high during ART(*56*). The

immunosuppressive activity of TGF-β is well-known. However, the effect of TGF-β signaling in

immune cells is highly context dependent(*57*). Hence, TGF-β plays different roles according to a

cell differentiation and activation status(*57*). In CD8$^+$ T cells and NK cells, TGF-β was shown to

decrease mTOR activity and preserve cellular metabolism (high mitochondrial activity and spare

respiratory capacity, but reduced mTOR activity) preventing metabolic exhaustion(*25, 26, 58,*

59). This effect was linked to survival of antigen-specific CD8$^+$ T cells, preservation of their

stemness and it was linked to higher expression of the TCF1 factor(*25*).

In contrast, in CD4$^+$ T cells TGF-β is known to decrease TCR activation(*29, 60*), restrict

proliferation and inhibit cytotoxicity (including granzyme and perforin release) at different stage

of infection in vivo(*30, 61*). However, the role of TGF-β in the formation and preservation of

CD4$^+$ T cell memory is still unclear(*57*). Moreover, the link between TGF-β signaling and TCF1

expression in CD4$^+$ T cells is unexplored.

Here, we used a clinical stage small drug, galunisertib, developed by Eli Lilly and used in

several phase 1 and 1/2 clinical studies against solid cancer(*39, 62, 63*) to investigate the

impact of TGF-β blockade on SIV latency, SIV reservoir and immune responses. Of note, Eli

Lilly did not terminate galunisertib development program because of toxicity(*64, 65*). Indeed, in

our studies in macaques, we observed no adverse events nor changes in chemical or

hematological variables. Moreover, there were no detectable changes in the levels of the 24

inflammatory factors that we probed in plasma during the treatment. This suggests that this

therapeutic approach may be safe in people living with HIV (PLWH).

The first important finding of our study was the confirmation of our previous report of the latency

reversal properties of TGF-β blockade in vivo(*35*). Indeed, we found increase in pVL in 7 out of

the 8 macaques upon initiation of galunisertib therapy. Although not all macaques were fully

suppressed before treatment, substantial increases in pVL ($>10^2$ copies/ml) were noted also in

fully suppressed macaques (08M171, A8R095 and A8L057). Moreover, extensive viral

reactivation was documented in tissues by immunoPET/CT. The increase detected post-

galunisertib in gut SUV correlated with CA-vRNA as in our previous studies(*35*) and,

interestingly, it was associated with a decrease in TCF1. Galunisertib led to increased probe

uptake in the gut and lymphoid tissues, as well as a generalized increase in whole-body uptake.

Yet, there was no change in the PET background signal in kidneys and spleen. This, in

conjunction with our previous studies demonstrating the specificity of the PET signal for areas of

enhanced SIV replication in gut and lymph nodes(*35, 41, 66*), demonstrates that the

galunisertib-driven increases in SUV were mostly specific, and highlighted areas of SIV latency

reversal at least in gut and lymphoid tissues. The extent to which the increased signal in the

spine and bones recapitulates an increase in SIV replication at these sites remains to be

determined.

Importantly, we observed a decrease in CA-vDNA in all the tissues that cannot be attributed to

ART alone. Indeed, although we did not have a concurrent control group, this decrease was not

present in similar studies conducted in SIVmac239M2 infected macaques on the same ART

regimen, but not treated with galunisertib. Considering studies by other groups with different

models (SIVmac251), they report 2^{nd} phase decay of SIV intact provirus (weeks ~32 to ~100pi,

Fig 2B in(*67*)) with a $t_{1/2}$ of >8 months(*67*). In contrast, in our study, the intact provirus decreased

by 3 folds (median) in a little over 3 months (from week 35pi, BC1 to week 49pi, AC4).

Interestingly, this decrease in intact pro-virus trended toward a direct correlation with IFN-γ

responses. However, IFN–γ levels also inversely correlated with the absolute value of residual

intact provirus, suggesting that IFN-γ responses were driven by residual viral reservoir while, at

the same time, were involved in clearing intact virus. Indeed, the increase in IFN-γ and TNF-α

also correlated with residual CA-vRNA in the LN at the end of the treatment. The latter, in turn,

was directly proportional to the increase in gut SUV. This is in line with increased latency

reversal explaining residual viral RNA in lymphoid tissues.

Finally, one of our most intriguing results was the profound downregulation of TCF1 in CD4$^+$ T

cells at both the transcriptional and protein levels. Although TCF1 is conventionally viewed as

an effector of the canonical Wnt pathway(*68*) and recently reached notoriety for its role in

maintaining stemness of antigen-specific memory CD8$^+$ T cells(*26, 69*), TCF1 has a plethora of

functions in T cell development and differentiation largely independent of Wnt signaling(*22*). In

CD4$^+$ T cells, TCF1 has been implicated in orchestrating all the major Th subsets, including

Th1, Th2, Th17 and Tfh(*70*). TCF1 is known to control the bifurcation between Th1 and Tfh in

favor of Tfh cells(*70*), while it negatively regulates Treg development(*22*). This is in line with our

findings of increased Treg and decreased Tfh, in the midst of a profound downregulation of

TCF1. Importantly, TCF1 is downregulated with increased cellular differentiation and

progression toward effector functions in T cells. T cell activation leads to reduced levels of TCF1

and higher levels of TCF1 are present in T cells with higher stemness and low anabolic

metabolism(*68*). These findings suggest that TCF1 has a critical role in maintaining quiescence

in immune cells likely in concert with TGF-β(*18, 61*). Our data reveal that this link may be even

more prominent and important in CD4$^+$ T cells than in CD8$^+$ T cells. Importantly, the decrease in

TCF1 was accompanied with enhancement in other effector markers such as CD95, CD16 and

GRZB (at the transcriptional level) and an increase in the transcription of AP1 complex.

However, there was no clear upregulation of other classical markers of immune activation such

as CD69 and no increase in T cell proliferation (Ki67 expression). Hence, galunisertib treatment

does not appear to lead to classical T cell activation. Instead, in vivo TGF-β blockade seems to

primarily change the metabolic state of T cells (and likely other immune cells) increasing

OXPHOS and mitochondrial function. Of note, although glycolysis is essential during cell

activation, mitochondrial pathways are engaged and remodeled early after activation and

OXPHOS upregulation has a pivotal role in the earliest stages of cell activation(*18*).

Hence, we propose a model in which TGF-β inhibition forces cells (particularly CD4$^+$ T cells) out

of quiescence to a transitional state where they reinitiate their transcriptional program and are

metabolically ready to be activated and effector. In absence of direct TCR engagement or other

activation stimuli, they do not undergo full/classical activation and proliferation, yet they can

reinitiate viral transcription and respond to SIV antigenic stimulation. Indeed, functionally, we

demonstrated that the PBMC after galunisertib treatment secrete higher levels of IFN-γ and

TNF-α. Interestingly, there was no increase in IL-2 secretion. The link between TGF-β and IL-

2(*71*) and the critical role of IL-2 in T cell proliferation again suggest that galunisertib enhances

an effector phenotype uncoupled from cellular proliferation. scRNAseq analysis demonstrated a

trend toward an effector phenotype also in other immune cells, such as B cells, NKs and

macrophages. Future studies will need to uncover in depth the effect of TGF-β blockade on

these other immune subsets.

This study has several limitations. The most important limitations are the relatively small number

of macaques and the lack of a concurrent untreated control group. We also could not investigate

in depth the viral kinetics after ART interruption because of the short follow up after ATI.

Moreover, we did not explore changes in the phenotype or turnover of cells isolated from gut

and lymph nodes and relied solely on transcriptional data for these tissues. Finally, because of

sample availability, we could not dissect the cellular origin of increased IFN-γ and TNF-α.

In conclusion, we report that in vivo treatment with a clinical stage small molecule TGF-β

inhibitor drives an effector phenotype in T cells that is likely responsible for increasing the

frequency of spontaneous latency reversal events, stimulating SIV-specific immune responses,

and decreasing the viral reservoir. Future work will determine whether the galunisertib-driven

enhanced antiviral responses and decreased viral reservoirs can significantly contribute to post-

ART virological control.

MATERIALS AND METHODS

Study design and Ethics Statement

A total of 8 adult female Indian origin *Rhesus* macaques (*Macaca mulatta*; Mamu A*01, B*08

and B*17 negative) were used for the study described in the manuscript (Table S1). All the

macaques were selected form the colonies bred and raised at the New Iberia Research Center

(NIRC), University of Louisiana at Lafayette. Treatments were in accordance with NIRC

standards which comply with the Guide for the Use and Care of Laboratory Animals. Any

abnormal observations and/or signs of illness or distress were reported per NIRC SOPs.

Appropriate procedures were performed to ensure that potential distress, pain, discomfort were

limited to that unavoidable in the conduct of the research plan. Ketamine (10mg/kg) and/or

telazol (4mg/kg) sedation were used for sample administration, collection of samples and

conducting image collections. Analgesics were used when determined appropriate by veterinary

medical staff. Monkeys were fed Teklad monkey chow supplemented with fresh fruit or

vegetables and water ad libitum.

Rhesus macaques (n=8 main study + 4 separate non-concomitant study) were infected with 300

$TCID_{50}$ of the barcoded SIVmac239M2 stock intravenously and ART (Tenofovir [PMPA],

Emtricitabine [FTC] and Dolutegravir [DTG]) was initiated on week 6 pi. Galunisertib treatment

was initiated on week 35 p.i. Powder (MedChemExpress – MCE, NJ, USA) was dissolved in

water and given orally in a treat twice daily at 20mg/kg. 4 cycles of 2 weeks daily treatment with

2 weeks wash out period were performed. Macaques 08M156 and A6X003 were given the

rhesus recombinant antibody (rhesus/human chimeric) anti-PD1 antibody [NIVOR4LALA;

comprising silenced rhesus IgG4k constant regions and variable regions from anti-human PD-1,

nivolumab; non-human primates reagents resource, NHPRR.

Blood viral load was monitored biweekly before and during ART and every 3-4 days during

Galunisertib treatment. Colorectal biopsies and LN FNA were collected before and after

galunisertib treatment. ART was terminated 3 weeks after the last galunisertib dose and

euthanasia and necropsy to harvest tissues were performed at week 58 post infection, tissues

samples were flash frozen, fixed in OCT or Z-fix.

Plasma and Tissue SIV Viral loads (VL)

Blood was collected in EDTA tubes and plasma was separated by density gradient

centrifugation and used for the determination of plasma VL by SIVgag qRT-PCR at NIRC or at

Leidos (Quantitative Molecular Diagnostics Core, AIDS and Cancer Virus Program Frederick

National Laboratory). Tissue VL from snap frozen PBMC pellets, colorectal biopsies and LN

FNA were performed as described in (*72*). SIV-TILDA was performed on freshly stored PBMC

before cycle 1 and at the end of cycle 4 of galunisertib by Accelevir, Baltimore, MD.

ImmunoPET/CT

ImmunoPET/CT for mapping SIV signals in total body scans were conducted in part as reported

previously(*42*). The probe consisted of primatized p7D3 anti-env F(ab)'2 coupled with the

chelator DOTA and labeled with Cu^{64} just prior to administration to the animals. For probe

administration, the animals were sedated, and a venous catheter was placed into an arm of leg

vein to minimize bleeding of the probe into the tissue surrounding the site of injection. The probe

for each animal consisted of ~1mg of the p7D3 F(ab)'2 labeled with 2-3 mCi of ^{64}Cu. After

injection, the animals were allowed to recuperate in their cage until the next day. At 24 hours,

the animals were again anesthetized with Telazol and the macaque's body was immobilized in

dorsal recumbency on the scanner table. Scans were conducted in a Phillips Gemini TF64

scanner. The final CT image was compiled from 200 to 300 slices, depending on macaque size.

PET Image analysis was performed using the MIM software. PET/CT fusions were generated

scaled according to calculated Standardized Uptake Values (SUV). The SUV scale for the PET

scans was selected based on the overall signal intensity of the PET scans (whole body), and

the CT scale was selected for optimal visibility of the tissues. All images and maximum image

projections were set to the same scale for visual comparisons. Additional details on MIM

analysis are described in Supplemental Methods.

Cell isolation, flow cytometry staining, classical and high-dimensional analysis

Colorectal biopsy tissues were isolated by enzymatic digestion while LN biopsies were passed

through a 70μm cell strainer as described in(*42*). Isolated cells were phenotyped with panels

listed in Supplemental Table S4. FlowJo vs 10.8 was use for both classical and high-dimesional

analysis. PeacoQC, FlowtSNE and FlowSOM plug-ins were used with default settings. SwiftReg

was used for normalization(*45*). More details on the staining procedures and analysis pipeline in

Supplemental Methods.

Bulk and scRNAseq analysis

For bulk RNAseq, snap frozen PBMC pellets from BC1, 1hr after first Gal dose and AC1 were

used for RNA extraction with the RNeasy kit with on column DNA digestion (Qiagen). Library

preparation was performed using TruSeq Stranded Total RNA with Ribo-Zero Globin and

sequencing was done with an Illumina HiSeq4000 with >20M reads/samples. Sequencing data

was demultiplexed and trimmed using Trimmomatic v0.36 to remove adapters and low-quality

reads. Trimmed reads were aligned to the Mmul10 reference genome and transcripts quantified

using the Hisat2-StringTie pipeline(*73*). Differential gene expression analysis using the

quantified gene transcripts was performed with DESeq2 R package(*74*) comparing the samples

attained before and after galunisertib treatment and controlling for intra-animal autocorrelation.

Differentially expressed genes (DEGs) were analyzed by functional enrichment analysis and

gene set enrichment analysis (GSEA) to identify specific pathways and molecular processes

altered by galunisertib.

The Parse pipeline and Partek software were. For bulk RNAseq features with <100 counts were

removed, data normalized and DESeq2 was used to obtain a DEG list. Genes with a false

discovery rate (FDR)-adjusted p-value ≤0.05 and absolute $\log_2$ fold-change (FC) (compared to

BC1) above 2 were defined as significantly differentially expressed (DEG). For scRNAseq

analysis cells isolated from lymph nodes before (BC1) and after (AC1) galuniertib were fixes

with the Parse fixation kit, barcoded and sequenced at the NUseq Core. The Partek software

was used for scRNAseq analysis. Cells with 400-8000 features, excluding features with 0 reads

in >99.99 cells, were included. Scran deconvolution was used for normalization and cell

classification was based on gene expression. Hurdle models were used to compare DEGs in

each cell subset before and after galunisertib. See supplemental methods for a detailed

description of scRNAseq analysis and additional control analysis using Seurat R package with

more stringent QC cut-offs and SCTransform normalization.

Plasma cytokines and T cell responses

Cytokines in plasma (at 1:2 dilution) were measured using the NHP Cytokine 24-Plex kit by

Meso Scale Diagnostics (MSD) according to manufacturer instructions. Frozen PBMC collected

at week 6 post-infection (pre-ART), right before the first galunisertib administration (BC1), at the

end of cycle 1 (AC1) and at the end of cycle 3 (AC3) were thawed in AIM V medium (Thermo

Fisher) with benzonase (Sigma) and plated on a FluoroSpot (CTL) plate pre-activated with 70%

Ethanol and IFN-γ, TNF-α and IL-2 capture solution. Gag, pol, and env 15-mer peptides (NIH

AIDS Reagents program) were prepared at two times the final concentration of 2.5μg/ml with

co-stimulatory reagents anti-CD28 10μg/ml and anti-CD49d 10μg/ml and added to the cells in

CTL-Test™ Medium. Parallel positive control of PMA (20ng/mL) and ionomycin (200ng/mL) or

mock DMSO solution was also plated with the stimulatory reagents. PBMCs were added at

300,000 cells per well. After 24hrs, the plate was washed, and incubated with detection and

tertiary solutions and shipped to CTL for scanning and QC.

Statistics

GraphPad Prism v10, R and Python were used for statistical analysis and data visualization.

Wilcoxon matched-pairs test and repeated measures ANOVA or mixed effect analysis (when the

data set had missing data) were used to compare the different virological and immunological

variables between baseline (BC1) and a single or multiple post-galunisertib time points. In Prism

the mixed model uses a compound symmetry covariance matrix, and is fit using Restricted

Maximum Likelihood (REML). Cytokine data from MSD assay were first Log transformed, then

normalized by subcolumn/factor in percentage with 0% as smallest value and 100% larger value

in each dataset. Each factor was analyzed separately with ANOVA for repeated measures and

Dunn's multiple comparison's post-hoc test and together by principal component analysis

(PCA). Holm-Sidak test was used for multiple comparison correction in all cases, but FlowSOM

cluster comparison where the FDR method was used. Pearson r coefficient was calculated for

pairwise tests of association in a correlation matrix with selected variables. For RNAseq

analysis see above and supplemental methods. For viral population analysis, Shannon Entropy

of the viral barcodes present in each sample was used to measure the diversity of the viral

populations (see supplemental Methods). Chi-squared tests were used in R to compare barcode

composition between before and after cycle time points within a tissue/macaque using paired

barcode relative frequencies. Unless otherwise specified $p\text{-value}<0.05$ was considered

statistically significant.

LIST OF SUPPLEMENTARY MATERIALS

Supplementary Methods

Supplementary Figure Fig. S1 to S16

Supplementary Table S1 to S6

Movies S1 to S8

REFERENCES

1. P. Gantner, S. Buranapraditkun, A. Pagliuzza, C. Dufour, M. Pardons, J. L. Mitchell, E. Kroon, C. Sacdalan, N. Tulmethakaan, S. Pinyakorn, M. L. Robb, N. Phanuphak, J. Ananworanich, D. Hsu, S. Vasan, L. Trautmann, R. Fromentin, N. Chomont, HIV rapidly targets a diverse pool of CD4(+) T cells to establish productive and latent infections. *Immunity* **56**, 653-668 e655 (2023).

2. J. B. Whitney, A. L. Hill, S. Sanisetty, P. Penaloza-MacMaster, J. Liu, M. Shetty, L. Parenteau, C. Cabral, J. Shields, S. Blackmore, J. Y. Smith, A. L. Brinkman, L. E. Peter, S. I. Mathew, K. M. Smith, E. N. Borducchi, D. I. Rosenbloom, M. G. Lewis, J. Hattersley, B. Li, J. Hesselgesser, R. Geleziunas, M. L. Robb, J. H. Kim, N. L. Michael, D. H. Barouch, Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys. *Nature* **512**, 74-77 (2014).

3. D. Persaud, Y. Zhou, J. M. Siliciano, R. F. Siliciano, Latency in human immunodeficiency virus type 1 infection: no easy answers. *J Virol* **77**, 1659-1665 (2003).

4. D. M. Margolis, N. M. Archin, Proviral Latency, Persistent Human Immunodeficiency Virus Infection, and the Development of Latency Reversing Agents. *The Journal of infectious diseases* **215**, S111-S118 (2017).

5. T. A. Wagner, S. McLaughlin, K. Garg, C. Y. Cheung, B. B. Larsen, S. Styrchak, H. C. Huang, P. T. Edlefsen, J. I. Mullins, L. M. Frenkel, HIV latency. Proliferation of cells with HIV integrated into cancer genes contributes to persistent infection. *Science* **345**, 570-573 (2014).

6. F. Maldarelli, X. Wu, L. Su, F. R. Simonetti, W. Shao, S. Hill, J. Spindler, A. L. Ferris, J. W. Mellors, M. F. Kearney, J. M. Coffin, S. H. Hughes, HIV latency. Specific HIV integration sites are linked to clonal expansion and persistence of infected cells. *Science* **345**, 179-183 (2014).

7. L. B. Cohn, I. T. Silva, T. Y. Oliveira, R. A. Rosales, E. H. Parrish, G. H. Learn, B. H. Hahn, J. L. Czartoski, M. J. McElrath, C. Lehmann, F. Klein, M. Caskey, B. D. Walker, J. D. Siliciano, R. F. Siliciano, M. Jankovic, M. C. Nussenzweig, HIV-1 integration landscape during latent and active infection. *Cell* **160**, 420-432 (2015).

8. K. B. Einkauf, M. R. Osborn, C. Gao, W. Sun, X. Sun, X. Lian, E. M. Parsons, G. T. Gladkov, K. W. Seiger, J. E. Blackmer, C. Jiang, S. A. Yukl, E. S. Rosenberg, X. G. Yu, M. Lichterfeld,

Parallel analysis of transcription, integration, and sequence of single HIV-1 proviruses.
Cell **185**, 266-282 e215 (2022).

9. X. Lian, K. W. Seiger, E. M. Parsons, C. Gao, W. Sun, G. T. Gladkov, I. C. Roseto, K. B.
Einkauf, M. R. Osborn, J. M. Chevalier, C. Jiang, J. Blackmer, M. Carrington, E. S.
Rosenberg, M. M. Lederman, D. K. McMahon, R. J. Bosch, J. M. Jacobson, R. T. Gandhi,
M. J. Peluso, T. W. Chun, S. G. Deeks, X. G. Yu, M. Lichterfeld, Progressive transformation
of the HIV-1 reservoir cell profile over two decades of antiviral therapy. *Cell Host*
Microbe **31**, 83-96 e85 (2023).

10. V. Singh, A. Dashti, M. Mavigner, A. Chahroudi, Latency Reversal 2.0: Giving the Immune
System a Seat at the Table. *Curr HIV/AIDS Rep* **18**, 117-127 (2021).

11. E. Abner, A. Jordan, HIV "shock and kill" therapy: In need of revision. *Antiviral Res* **166**,
19-34 (2019).

12. A. Ait-Ammar, A. Kula, G. Darcis, R. Verdikt, S. De Wit, V. Gautier, P. W. G. Mallon, A.
Marcello, O. Rohr, C. Van Lint, Current Status of Latency Reversing Agents Facing the
Heterogeneity of HIV-1 Cellular and Tissue Reservoirs. *Frontiers in microbiology* **10**, 3060
(2019).

13. S. A. Yukl, P. Kaiser, P. Kim, S. Telwatte, S. K. Joshi, M. Vu, H. Lampiris, J. K. Wong, HIV
latency in isolated patient CD4(+) T cells may be due to blocks in HIV transcriptional
elongation, completion, and splicing. *Sci Transl Med* **10**, (2018).

14. J. D. Siliciano, R. F. Siliciano, Low Inducibility of Latent Human Immunodeficiency Virus
Type 1 Proviruses as a Major Barrier to Cure. *J Infect Dis* **223**, 13-21 (2021).

15. Y. C. Ho, L. Shan, N. N. Hosmane, J. Wang, S. B. Laskey, D. I. Rosenbloom, J. Lai, J. N.
Blankson, J. D. Siliciano, R. F. Siliciano, Replication-competent noninduced proviruses in
the latent reservoir increase barrier to HIV-1 cure. *Cell* **155**, 540-551 (2013).

16. E. N. Borducchi, C. Cabral, K. E. Stephenson, J. Liu, P. Abbink, D. Ng'ang'a, J. P. Nkolola, A.
L. Brinkman, L. Peter, B. C. Lee, J. Jimenez, D. Jetton, J. Mondesir, S. Mojta, A.
Chandrashekar, K. Molloy, G. Alter, J. M. Gerold, A. L. Hill, M. G. Lewis, M. G. Pau, H.
Schuitemaker, J. Hesselgesser, R. Geleziunas, J. H. Kim, M. L. Robb, N. L. Michael, D. H.
Barouch, Ad26/MVA therapeutic vaccination with TLR7 stimulation in SIV-infected
rhesus monkeys. *Nature* **540**, 284-287 (2016).

17. C. C. Nixon, M. Mavigner, G. C. Sampey, A. D. Brooks, R. A. Spagnuolo, D. M. Irlbeck, C.
Mattingly, P. T. Ho, N. Schoof, C. G. Cammon, G. K. Tharp, M. Kanke, Z. Wang, R. A. Cleary,
A. A. Upadhyay, C. De, S. R. Wills, S. D. Falcinelli, C. Galardi, H. Walum, N. J. Schramm, J.
Deutsch, J. D. Lifson, C. M. Fennessey, B. F. Keele, S. Jean, S. Maguire, B. Liao, E. P.
Browne, R. G. Ferris, J. H. Brehm, D. Favre, T. H. Vanderford, S. E. Bosinger, C. D. Jones, J.
P. Routy, N. M. Archin, D. M. Margolis, A. Wahl, R. M. Dunham, G. Silvestri, A. Chahroudi,
J. V. Garcia, Systemic HIV and SIV latency reversal via non-canonical NF-kappaB signalling
in vivo. *Nature* **578**, 160-165 (2020).

18. D. O'Sullivan, The metabolic spectrum of memory T cells. *Immunol Cell Biol* **97**, 636-646
(2019).

19. J. M. Crater, D. F. Nixon, R. L. Furler O'Brien, HIV-1 replication and latency are balanced
by mTOR-driven cell metabolism. *Front Cell Infect Microbiol* **12**, 1068436 (2022).

20. E. Besnard, S. Hakre, M. Kampmann, H. W. Lim, N. N. Hosmane, A. Martin, M. C. Bassik,
E. Verschueren, E. Battivelli, J. Chan, J. P. Svensson, A. Gramatica, R. J. Conrad, M. Ott, W.

C. Greene, N. J. Krogan, R. F. Siliciano, J. S. Weissman, E. Verdin, The mTOR Complex
Controls HIV Latency. *Cell Host Microbe* **20**, 785-797 (2016).

21.   H. J. Barbian, M. S. Seaton, S. D. Narasipura, J. Wallace, R. Rajan, B. E. Sha, L. Al-Harthi,
beta-catenin regulates HIV latency and modulates HIV reactivation. *PLoS Pathog* **18**,
e1010354 (2022).

22.   F. Gounari, K. Khazaie, TCF-1: a maverick in T cell development and function. *Nat*
Immunol **23**, 671-678 (2022).

23.   J. C. Valle-Casuso, M. Angin, S. Volant, C. Passaes, V. Monceaux, A. Mikhailova, K.
Bourdic, V. Avettand-Fenoel, F. Boufassa, M. Sitbon, O. Lambotte, M. I. Thoulouze, M.
Muller-Trutwin, N. Chomont, A. Saez-Cirion, Cellular Metabolism Is a Major Determinant
of HIV-1 Reservoir Seeding in CD4(+) T Cells and Offers an Opportunity to Tackle
Infection. *Cell Metab* **29**, 611-626 e615 (2019).

24.   S. A. Oh, M. O. Li, TGF-beta: guardian of T cell function. *Journal of immunology* **191**,
3973-3979 (2013).

25.   S. S. Gabriel, C. Tsui, D. Chisanga, F. Weber, M. Llano-Leon, P. M. Gubser, L. Bartholin, F.
Souza-Fonseca-Guimaraes, N. D. Huntington, W. Shi, D. T. Utzschneider, A. Kallies,
Transforming growth factor-beta-regulated mTOR activity preserves cellular metabolism
to maintain long-term T cell responses in chronic infection. *Immunity* **54**, 1698-1714
e1695 (2021).

26.   Y. Hu, W. H. Hudson, H. T. Kissick, C. B. Medina, A. P. Baptista, C. Ma, W. Liao, R. N.
Germain, S. J. Turley, N. Zhang, R. Ahmed, TGF-beta regulates the stem-like state of PD-
1+ TCF-1+ virus-specific CD8 T cells during chronic infection. *J Exp Med* **219**, (2022).

27.   C. Ma, N. Zhang, Transforming growth factor-beta signaling is constantly shaping
memory T-cell population. *Proc Natl Acad Sci U S A* **112**, 11013-11017 (2015).

28.   S. Viel, A. Marcais, F. S. Guimaraes, R. Loftus, J. Rabilloud, M. Grau, S. Degouve, S.
Djebali, A. Sanlaville, E. Charrier, J. Bienvenu, J. C. Marie, C. Caux, J. Marvel, L. Town, N.
D. Huntington, L. Bartholin, D. Finlay, M. J. Smyth, T. Walzer, TGF-beta inhibits the
activation and functions of NK cells by repressing the mTOR pathway. *Sci Signal* **9**, ra19
(2016).

29.   J. S. Delisle, M. Giroux, G. Boucher, J. R. Landry, M. P. Hardy, S. Lemieux, R. G. Jones, B. T.
Wilhelm, C. Perreault, The TGF-beta-Smad3 pathway inhibits CD28-dependent cell
growth and proliferation of CD4 T cells. *Genes Immun* **14**, 115-126 (2013).

30.   G. M. Lewis, E. J. Wehrens, L. Labarta-Bajo, H. Streeck, E. I. Zuniga, TGF-beta receptor
maintains CD4 T helper cell identity during chronic viral infections. *J Clin Invest* **126**,
3799-3813 (2016).

31.   A. P. Nath, A. Braun, S. C. Ritchie, F. R. Carbone, L. K. Mackay, T. Gebhardt, M. Inouye,
Comparative analysis reveals a role for TGF-beta in shaping the residency-related
transcriptional signature in tissue-resident memory CD8+ T cells. *PLoS One* **14**, e0210495
(2019).

32.   N. Zhang, M. J. Bevan, Transforming growth factor-beta signaling controls the formation
and maintenance of gut-resident memory T cells by regulating migration and retention.
Immunity **39**, 687-696 (2013).

33.   T. Hirai, Y. Yang, Y. Zenke, H. Li, V. K. Chaudhri, J. S. De La Cruz Diaz, P. Y. Zhou, B. A.
Nguyen, L. Bartholin, C. J. Workman, D. W. Griggs, D. A. A. Vignali, H. Singh, D. Masopust,

D. H. Kaplan, Competition for Active TGFbeta Cytokine Allows for Selective Retention of
Antigen-Specific Tissue- Resident Memory T Cells in the Epidermal Niche. *Immunity* **54**,
84-98 e85 (2021).

34.   C. Larson, B. Oronsky, C. A. Carter, A. Oronsky, S. J. Knox, D. Sher, T. R. Reid, TGF-beta: a
master immune regulator. *Expert Opin Ther Targets* **24**, 427-438 (2020).

35.   S. Samer, Y. Thomas, M. Arainga, C. Carter, L. M. Shirreff, M. S. Arif, J. M. Avita, I. Frank,
M. D. McRaven, C. T. Thuruthiyil, V. B. Heybeli, M. R. Anderson, B. Owen, A. Gaisin, D.
Bose, L. M. Simons, J. F. Hultquist, J. Arthos, C. Cicala, I. Sereti, P. J. Santangelo, R.
Lorenzo-Redondo, T. J. Hope, F. J. Villinger, E. Martinelli, Blockade of TGF-beta signaling
reactivates HIV-1/SIV reservoirs and immune responses in vivo. *JCI Insight* **7**, (2022).

36.   S. Bergstresser, D. A. Kulpa, TGF-beta Signaling Supports HIV Latency in a Memory CD4+
T Cell Based In Vitro Model. *Methods Mol Biol* **2407**, 69-79 (2022).

37.   S. Chinnapaiyan, R. K. Dutta, M. Nair, H. S. Chand, I. Rahman, H. J. Unwalla, TGF-beta1
increases viral burden and promotes HIV-1 latency in primary differentiated human
bronchial epithelial cells. *Scientific reports* **9**, 12552 (2019).

38.   R. B. Holmgaard, D. A. Schaer, Y. Li, S. P. Castaneda, M. Y. Murphy, X. Xu, I. Inigo, J.
Dobkin, J. R. Manro, P. W. Iversen, D. Surguladze, G. E. Hall, R. D. Novosiadly, K. A.
Benhadji, G. D. Plowman, M. Kalos, K. E. Driscoll, Targeting the TGFbeta pathway with
galunisertib, a TGFbetaRI small molecule inhibitor, promotes anti-tumor immunity
leading to durable, complete responses, as monotherapy and in combination with
checkpoint blockade. *J Immunother Cancer* **6**, 47 (2018).

39.   D. Melisi, D. Y. Oh, A. Hollebecque, E. Calvo, A. Varghese, E. Borazanci, T. Macarulla, V.
Merz, C. Zecchetto, Y. Zhao, I. Gueorguieva, M. Man, L. Gandhi, S. T. Estrem, K. A.
Benhadji, M. C. Lanasa, E. Avsar, S. C. Guba, R. Garcia-Carbonero, Safety and activity of
the TGFbeta receptor I kinase inhibitor galunisertib plus the anti-PD-L1 antibody
durvalumab in metastatic pancreatic cancer. *J Immunother Cancer* **9**, (2021).

40.   A. Wick, A. Desjardins, C. Suarez, P. Forsyth, I. Gueorguieva, T. Burkholder, A. L. Cleverly,
S. T. Estrem, S. Wang, M. M. Lahn, S. C. Guba, D. Capper, J. Rodon, Phase 1b/2a study of
galunisertib, a small molecule inhibitor of transforming growth factor-beta receptor I, in
combination with standard temozolomide-based radiochemotherapy in patients with
newly diagnosed malignant glioma. *Invest New Drugs* **38**, 1570-1579 (2020).

41.   P. J. Santangelo, K. A. Rogers, C. Zurla, E. L. Blanchard, S. Gumber, K. Strait, F. Connor-
Stroud, D. M. Schuster, P. K. Amancha, J. J. Hong, S. N. Byrareddy, J. A. Hoxie, B.
Vidakovic, A. A. Ansari, E. Hunter, F. Villinger, Whole-body immunoPET reveals active SIV
dynamics in viremic and antiretroviral therapy-treated macaques. *Nat Methods* **12**, 427-
432 (2015).

42.   S. Samer, Y. Thomas, M. Araínga, C. Carter, L. Shirreff, M. Arif, J. Avita, I. Frank, M.
McRaven, C. Thuruthiyil, V. Heybeli, M. Anderson, B. Owen, A. Gaisin, D. Bose, L. Simons,
J. Hultquist, J. Arthos, C. Cicala, I. Sereti, P. Santangelo, R. Lorenzo-Redondo, T. Hope, F.
Villinger, E. Martinelli, Blockade of TGF-β signaling reactivates HIV-1/SIV reservoirs and
immune responses in vivo. *bioRxiv*, 2022.2005.2013.489595 (2022).

43.   S. J. Im, B. T. Konieczny, W. H. Hudson, D. Masopust, R. Ahmed, PD-1+ stemlike CD8 T
cells are resident in lymphoid tissues during persistent LCMV infection. *Proc Natl Acad*
Sci U S A **117**, 4292-4299 (2020).

44. S. Van Gassen, B. Callebaut, M. J. Van Helden, B. N. Lambrecht, P. Demeester, T. Dhaene,
Y. Saeys, FlowSOM: Using self-organizing maps for visualization and interpretation of
cytometry data. *Cytometry A* **87**, 636-645 (2015).

45. J. A. Rebhahn, S. A. Quataert, G. Sharma, T. R. Mosmann, SwiftReg cluster registration
automatically reduces flow cytometry data variability including batch effects. *Commun*
Biol **3**, 218 (2020).

46. M. J. Blackburn, M. Zhong-Min, F. Caccuri, K. McKinnon, L. Schifanella, Y. Guan, G. Gorini,
D. Venzon, C. Fenizia, N. Binello, S. N. Gordon, C. J. Miller, G. Franchini, M. Vaccari,
Regulatory and Helper Follicular T Cells and Antibody Avidity to Simian
Immunodeficiency Virus Glycoprotein 120. *J Immunol* **195**, 3227-3236 (2015).

47. S. Helmold Hait, D. A. Vargas-Inchaustegui, T. Musich, V. Mohanram, I. Tuero, D. J.
Venzon, J. Bear, M. Rosati, M. Vaccari, G. Franchini, B. K. Felber, G. N. Pavlakis, M.
Robert-Guroff, Early T Follicular Helper Cell Responses and Germinal Center Reactions
Are Associated with Viremia Control in Immunized Rhesus Macaques. *J Virol* **93**, (2019).

48. Y. Zhou, B. Zhou, L. Pache, M. Chang, A. H. Khodabakhshi, O. Tanaseichuk, C. Benner, S.
K. Chanda, Metascape provides a biologist-oriented resource for the analysis of systems-
level datasets. *Nat Commun* **10**, 1523 (2019).

49. Z. Qiu, T. H. Chu, B. S. Sheridan, TGF-beta: Many Paths to CD103(+) CD8 T Cell Residency.
Cells **10**, (2021).

50. H. S. Lim, P. Qiu, Quantifying Cell-Type-Specific Differences of Single-Cell Datasets Using
Uniform Manifold Approximation and Projection for Dimension Reduction and Shapley
Additive exPlanations. *J Comput Biol* **30**, 738-750 (2023).

51. X. Liao, N. Sharma, F. Kapadia, G. Zhou, Y. Lu, H. Hong, K. Paruchuri, G. H.
Mahabeleshwar, E. Dalmas, N. Venteclef, C. A. Flask, J. Kim, B. W. Doreian, K. Q. Lu, K. H.
Kaestner, A. Hamik, K. Clement, M. K. Jain, Kruppel-like factor 4 regulates macrophage
polarization. *J Clin Invest* **121**, 2736-2749 (2011).

52. M. A. Moso, J. L. Anderson, S. Adikari, L. R. Gray, G. Khoury, J. J. Chang, J. C. Jacobson, A.
M. Ellett, W. J. Cheng, S. Saleh, J. J. Zaunders, D. F. J. Purcell, P. U. Cameron, M. J.
Churchill, S. R. Lewin, H. K. Lu, HIV latency can be established in proliferating and
nonproliferating resting CD4+ T cells in vitro: implications for latency reversal. *AIDS* **33**,
199-209 (2019).

53. J. Neidleman, X. Luo, J. Frouard, G. Xie, F. Hsiao, T. Ma, V. Morcilla, A. Lee, S. Telwatte, R.
Thomas, W. Tamaki, B. Wheeler, R. Hoh, M. Somsouk, P. Vohra, J. Milush, K. S. James, N.
M. Archin, P. W. Hunt, S. G. Deeks, S. A. Yukl, S. Palmer, W. C. Greene, N. R. Roan,
Phenotypic analysis of the unstimulated in vivo HIV CD4 T cell reservoir. *Elife* **9**, (2020).

54. A. Chitrakar, M. Sanz, S. B. Maggirwar, N. Soriano-Sarabia, HIV Latency in Myeloid Cells:
Challenges for a Cure. *Pathogens* **11**, (2022).

55. E. L. Pearce, Metabolism in T cell activation and differentiation. *Curr Opin Immunol* **22**,
314-320 (2010).

56. H. Valdez, M. M. Lederman, Cytokines and cytokine therapies in HIV infection. *AIDS Clin*
Rev, 187-228 (1997).

57. R. J. Salmond, Regulation of T Cell Activation and Metabolism by Transforming Growth
Factor-Beta. *Biology (Basel)* **12**, (2023).

58. S. Regis, A. Dondero, F. Caliendo, C. Bottino, R. Castriconi, NK Cell Function Regulation by
TGF-beta-Induced Epigenetic Mechanisms. *Front Immunol* **11**, 311 (2020).

59. V. Zaiatz-Bittencourt, D. K. Finlay, C. M. Gardiner, Canonical TGF-beta Signaling Pathway
Represses Human NK Cell Metabolism. *J Immunol* **200**, 3934-3941 (2018).

60. L. Das, A. D. Levine, TGF-beta inhibits IL-2 production and promotes cell cycle arrest in
TCR-activated effector/memory T cells in the presence of sustained TCR signal
transduction. *J Immunol* **180**, 1490-1498 (2008).

61. W. Chen, TGF-beta Regulation of T Cells. *Annu Rev Immunol* **41**, 483-512 (2023).

62. T. Yamazaki, A. J. Gunderson, M. Gilchrist, M. Whiteford, M. X. Kiely, A. Hayman, D.
O'Brien, R. Ahmad, J. V. Manchio, N. Fox, K. McCarty, M. Phillips, E. Brosnan, G. Vaccaro,
R. Li, M. Simon, E. Bernstein, M. McCormick, L. Yamasaki, Y. Wu, A. Drokin, T. Carnahan,
Y. To, W. L. Redmond, B. Lee, J. Louie, E. Hansen, M. C. Solhjem, J. Cramer, W. J. Urba, M.
J. Gough, M. R. Crittenden, K. H. Young, Galunisertib plus neoadjuvant
chemoradiotherapy in patients with locally advanced rectal cancer: a single-arm, phase 2
trial. *Lancet Oncol* **23**, 1189-1200 (2022).

63. J. J. Harding, R. K. Do, A. Yaqubie, A. Cleverly, Y. Zhao, I. Gueorguieva, M. Lahn, K. A.
Benhadji, R. K. Kelley, G. K. Abou-Alfa, Phase 1b study of galunisertib and ramucirumab
in patients with advanced hepatocellular carcinoma. *Cancer Med* **10**, 3059-3067 (2021).

64. R. K. Kelley, E. Gane, E. Assenat, J. Siebler, P. R. Galle, P. Merle, I. O. Hourmand, A.
Cleverly, Y. Zhao, I. Gueorguieva, M. Lahn, S. Faivre, K. A. Benhadji, G. Giannelli, A Phase
2 Study of Galunisertib (TGF-beta1 Receptor Type I Inhibitor) and Sorafenib in Patients
With Advanced Hepatocellular Carcinoma. *Clin Transl Gastroenterol* **10**, e00056 (2019).

65. D. Melisi, R. Garcia-Carbonero, T. Macarulla, D. Pezet, G. Deplanque, M. Fuchs, J. Trojan,
H. Oettle, M. Kozloff, A. Cleverly, C. Smith, S. T. Estrem, I. Gueorguieva, M. M. F. Lahn, A.
Blunt, K. A. Benhadji, J. Tabernero, Galunisertib plus gemcitabine vs. gemcitabine for
first-line treatment of patients with unresectable pancreatic cancer. *Br J Cancer* **119**,
1208-1214 (2018).

66. P. J. Santangelo, C. Cicala, S. N. Byrareddy, K. T. Ortiz, D. Little, K. E. Lindsay, S. Gumber, J.
J. Hong, K. Jelicic, K. A. Rogers, C. Zurla, F. Villinger, A. A. Ansari, A. S. Fauci, J. Arthos,
Early treatment of SIV+ macaques with an alpha(4)beta(7) mAb alters virus distribution
and preserves CD4(+) T cells in later stages of infection. *Mucosal Immunol* **11**, 932-946
(2018).

67. E. J. Fray, F. Wu, F. R. Simonetti, C. Zitzmann, N. Sambaturu, C. Molina-Paris, A. M.
Bender, P. T. Liu, J. D. Ventura, R. W. Wiseman, D. H. O'Connor, R. Geleziunas, T. Leitner,
R. M. Ribeiro, A. S. Perelson, D. H. Barouch, J. D. Siliciano, R. F. Siliciano, Antiretroviral
therapy reveals triphasic decay of intact SIV genomes and persistence of ancestral
variants. *Cell Host Microbe* **31**, 356-372 e355 (2023).

68. G. Escobar, D. Mangani, A. C. Anderson, T cell factor 1: A master regulator of the T cell
response in disease. *Sci Immunol* **5**, (2020).

69. R. L. Rutishauser, C. D. T. Deguit, J. Hiatt, F. Blaeschke, T. L. Roth, L. Wang, K. A. Raymond,
C. E. Starke, J. C. Mudd, W. Chen, C. Smullin, R. Matus-Nicodemos, R. Hoh, M. Krone, F.
M. Hecht, C. D. Pilcher, J. N. Martin, R. A. Koup, D. C. Douek, J. M. Brenchley, R. P. Sekaly,
S. K. Pillai, A. Marson, S. G. Deeks, J. M. McCune, P. W. Hunt, TCF-1 regulates HIV-specific
CD8+ T cell expansion capacity. *JCI Insight* **6**, (2021).

70. Y. S. Choi, J. A. Gullicksrud, S. Xing, Z. Zeng, Q. Shan, F. Li, P. E. Love, W. Peng, H. H. Xue, S.
Crotty, LEF-1 and TCF-1 orchestrate T(FH) differentiation by regulating differentiation
circuits upstream of the transcriptional repressor Bcl6. *Nat Immunol* **16**, 980-990 (2015).
71. S. G. Zheng, J. Wang, P. Wang, J. D. Gray, D. A. Horwitz, IL-2 is essential for TGF-beta to
convert naive CD4+CD25- cells to CD25+Foxp3+ regulatory T cells and for expansion of
these cells. *J Immunol* **178**, 2018-2027 (2007).
72. I. Frank, M. Cigoli, M. S. Arif, M. D. Fahlberg, S. Maldonado, G. Calenda, A. Pegu, E. S.
Yang, R. Rawi, G. Y. Chuang, H. Geng, C. Liu, T. Zhou, P. D. Kwong, J. Arthos, C. Cicala, B. F.
Grasperge, J. L. Blanchard, A. Gettie, C. M. Fennessey, B. F. Keele, M. Vaccari, T. J. Hope,
A. S. Fauci, J. R. Mascola, E. Martinelli, Blocking alpha4beta7 integrin delays viral
rebound in SHIVSF162P3-infected macaques treated with anti-HIV broadly neutralizing
antibodies. *Sci Transl Med* **13**, (2021).
73. M. Pertea, D. Kim, G. M. Pertea, J. T. Leek, S. L. Salzberg, Transcript-level expression
analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. *Nat Protoc* **11**,
1650-1667 (2016).
74. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
886

Acknowledgements

General: We acknowledge the helpful staff at the New Iberia Research Center as well as the help of Dr. Suchitra Swaminathan and the rest of the staff of the Flow Cytometry Core Facility at the Robert H. Lurie Comprehensive Cancer Center of Northwestern University in Chicago. Moreover, we thank the staff of the NUseq Core at Northwestern, and in particular Dr. Ching Man Wai and Matthew Schipma for the help with RNAseq data. Finally, we thank Dr. Lifson and the Leidos staff for viral load measurements as well as Dr. Laird at Accelevir for providing SIV-TILDA results in a timely manner. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

Funding: This project has been funded by National Institutes of Health grant R56AI157822 to Dr. Martinelli, the resource for NHP immune reagents to Dr. Villinger (R24 OD010947) for probe productions and in part with federal funds from the National Cancer Institute, National Institutes of Health, under Contract No. 75N91019D00024/HHSN261201500003I. The Lurie Cancer Center is supported in part by an NCI Cancer Center Support Grant #P30 CA060553.

Authors contributions: EM conceptualized the studies; JK performed and analyzed flow cytometry and immunePET/CT data; DB, MA, DF and SA collected and processed macaque samples; MRH generated vRNA/DNA and Elispot data; RC and MV analyzed Treg/Tfh; YT and TJH contributed to immunePET/CT analysis; EG, YG and EM analyzed RNAseq data; CMF, BK and RLR analyzed barcode data; JA, CC, FV and EM interpreted the data and wrote the manuscript.

Competing interests: The authors declare no competing interests.

Data and material availability: All data needed to evaluate the conclusions in the paper are included in the manuscript and/or the supplemental material. Transcriptional profiling data (raw RNAseq data) will be uploaded to GEO.

Figure Legends

Figure 1. Four 2-weeks cycles with galunisertib lead to viral reactivation in blood. A)

Schematic representation of the study and sampling schedule B) Plasma VL in blood for each

macaque throughout the study. The longer black line indicates the period on ART, while the 4

small black lines indicate the start and end of each galunisertib cycle. C) Enlarged plasma VL

for all macaques during galunisertib therapy. Green bars indicate galunisertib cycles.

Figure 2. Galunisertib leads to viral reactivation in tissues. A) The ^{64}Cu-DOTA-Fab$_2$(7D3)

probe was injected ~24hrs before PET/CT scan before and at the end of each of the first 3

galunisertib cycles. Representative images from the PET/CT scans are show for a macaque

with a major increase at the end of cycle 2 and one showing increase at the beginning of cycle 3

B) Total SUV were calculated for each anatomical area and values analyzed with mixed-effect

analysis. Thicker black line represents the mean. P–values were calculated for comparison of

each time point with the before cycle 1 time point (BC1; AC1= after cycle 1, BC2= before cycle

2; AC2= after cycle 2; BC3= before cycle 3; AC3= after cycle 3; Holm-Sidak multiple comparison

correction; *p≤0.05 **p≤0.01 ***p≤0.01)

Figure 3. Galunisertib decreases viral reservoir in absence of systemic inflammation. A)

Levels of cell-associated (CA)-vDNA per cell equivalent are shown for the time point before

cycle 1 (BC1) and at the end of cycle 3 (AC3) or 4 (AC4) for the respective tissues. B) IPDA

data are shown for intact and total provirus for BC1 and AC4 in PBMC. pvalues are shown for

Wilcoxon matched pair signed-rank non-parametric test (*p≤0.05 **p≤0.01 ***p≤0.01) C) Heat

map of cytokine concentration in plasma at the indicated time point are shown after Log

transformation and normalization. Statistical analysis was run on each factor separately and

together (no significant differences after multiple comparison adjustment).

Figure 4. Galunisertib leads toward effector in T and NK cells, increasing Treg and
decreasing Tfh frequencies. A-F) Geometric mean fluorescent intensities (MFI) of each
marker and frequency of indicated subset within live, singlets CD3$^+$ CD4$^+$ T cells (A and C) or
CD3$^+$ CD4$^+$ CD95$^+$ T cells (B) or CD8$^+$ or CD8$^+$ CD95$^+$ T cells or NK cells (NKG2A$^+$ CD8$^+$ CD3$^-$
cells) are shown. Thick black line represents the mean. Changes from baseline (beginning of
cycle 1, BC1) are shown for graphs with at least 1 significant difference (Repeated measures
ANOVA with Holm-Sidak correction; *p≤0.05 **p≤0.01 ***p≤0.01). G) tSNE of lymphocyte, live,
singlets events after normalization for BC1 and AC4 (all 8 macaques) with FlowSOM 36 clusters
overlaid on tSNE (top left) or heatmap of each markers MFI (right) or heatmap of time point
(blue is BC1 and red is AC4; bottom left) is shown. 6 populations were manually gated on red or
blue areas (red, New1-3 and blue Old1-3). H) Bubble chart displaying changes from BC1 in
AC4. Color is proportional to the effect size and size to p-value (Wilcoxon sum rank non-
parameter test). I) Bubble chart displaying changes from BC1 in all time point (ANOVA repeated
measures with Holm-Sidak correction; *p≤0.05 **p≤0.01 ***p≤0.01). J) Frequency of indicated
subset within live, singlets CD3$^+$ CD4$^+$ T or CD3$^+$ CD4$^+$ CD95$^+$ CD28$^+$ T cells (CM=central
memory) or within CD3$^+$ CD8$^+$ T cells. Changes from baseline (BC1) are shown (ANOVA
repeated measures with Holm-Sidak correction for multiple comparisons; *p≤0.05 **p≤0.01
***p≤0.01).

Figure 5. OXPHOS and other metabolic pathways increase rapidly with TGF-β blockade.
A-C) Bulk RNAseq was performed with PBMC from before cycle 1 (24hrs) and 1hrs after the first dose of galunisertib (A) or after the last dose of cycle 1 (B) and with rectal biopsies collected before cycle 1 (24hrs) and after the last dose of cycle 1 (C). The number of differentially expressed genes (DEG) with an Adjpvalue <0.05 and $log_2FC>2$ and <-2 are shown in each respective volcano plot. Enrichment plots are shown after GSEA (with all adjp<0.05 DEGs) for significantly enriched pathways (top 1 or 2 pathway by ES). C) Lollipop graph of selected DEG of interest among significantly different genes (adjp<0.05).

Figure 6. scRNAseq of lymph node before and after cycle 1 confirms a switch toward effector and increased metabolism in all immune subsets with galunisertib. A) UMAP projection of 93234 cells from lymph nodes collected right before and at the end of cycle 1 from all 8 macaques (16 samples). Gene-based classification of major immune subset is overlaid on UMAP. In gray are unclassified cells. B-C) Bubble plots showing expression (mean normalized counts proportional to the color; size proportional to the percentage of cells) of each marker

listed in each cell subset. Marker listed are those used for classification of major immune

subsets (B) or CD4[+] and CD8[+] T cells (C). D) Significantly different genes (Adjp<0.05;

log2FC=0.15) in the T cell subset are shown with color proportional to normalized counts. E)

Significantly enriched pathways (adjp<0.01) in T cells DEGs within the hallmark collection. F-G)

Significantly different genes (Adjp<0.05; log2FC=0.15) in the CD4[+] (F) and CD8[+] (G) T cell

subset. H-I) Significantly enriched pathways (adjp<0.01) in CD4[+] T cells DEGs within the

hallmark (H) and biocarta (I) collections.

Figure 7. Galunisertib increases SIV-specific responses. Average spots (from triplicates) per

10^6 PBMC at the time of ART initiation (pre-ART), before cycle 1 (BC1), after cycle 1 (AC1) and

at the end of cycle 3 (AC3) with galunisertib after 24hrs ex vivo stimulation with 15-mer peptides

(gag, env, pol) or mock (DMSO). Each post-galunisertib time point was compared to BC1

(Mixed effect analysis with Dunnet post-hoc p-values are shown; *p≤0.05 **p≤0.01 ***p≤0.001)

986
Figure 8. TCF1 decrease associates with virological and immunological endpoints. A)

Barcode diversity measure as Shannon Entropy is shown before and after each of the first 3

galunisertib cycles for LN, PBMC and colorectal biopsies. Box-and-whisker plot represents the

median +/− the interquartile range (no significant differences using linear mixed effects models).

Blue= before; Red= end of each cycle. B) Bubble plot shows the results of statistical testing

(Chi-squared) for differences in frequency distribution of barcodes before compared to after, for

each of the first 3 cycles of galunisertib for each macaque in the indicated tissues. Blue

indicates significant differences p≤0.05. C) Barcode entropy of virus isolated at the time of ART

initiation compared to week 6 post-ATI in plasma (Wilcoxon matched pairs test; *p≤0.05). D)

Correlation matrix of several key variables of virological or immunological effect of galunisertib.

Color is proportional to Pearson r coefficient. *p≤0.05 **p≤0.01 ***p≤0.001 indicate significant

correlations. F) Association between fold increase in TCF1 (MFI) from BC1 to AC4 with CA-

vDNA levels at AC4, change in gut SUV at AC3 compared to BC1 and fold increase in IFN-γ

(AC3 vs BC1). Person r is shown. All correlations have *p≤0.05.

1001