Screening of the Pandemic Response Box identifies anti-microsporidia compounds

Qingyuan Huang[1,2], Jie Chen[1], Guoqing Pan[1#], and Aaron W. Reinke[2#]

[1]State Key Laboratory of Resource Insects, Chongqing Key Laboratory of Microsporidia Infection

and Control, Southwest University, Chongqing, China,

[2]Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

#Corresponding author

gqpan@swu.edu.cn

aaron.reinke@utoronto.ca

Abstract

Microsporidia are fungi-like obligate intracellular pathogens, which infect most animals and cause

microsporidiosis. Despite the serious threat that microsporidia pose to humans and agricultural

animals, few drugs are available for the treatment and control of microsporidia. To identify novel

inhibitors, we took advantage of the model organism *Caenorhabditis elegans* infected with its

natural microsporidian *Nematocida parisii*. We used this system to screen the Pandemic

Response Box, a collection of 400 diverse compounds with known antimicrobial activity. After

testing these compounds in a 96-well format at high (100 μM) and low (40 μM) concentrations,

we identified four inhibitors that restored the ability of *C. elegans* to produce progeny in the

presence of *N. parisii*. All four compounds reduced the pathogen load of both *N. parisii* and

Pancytospora epiphaga, a *C. elegans*-infecting microsporidia related to human-infecting species.

One of these compounds, a known inhibitor of a viral protease, MMV1006203, inhibited invasion

and prevented the firing of spores. An albendazole analog, MMV1782387, inhibited proliferation

of *N. parisii*. We tested albendazole as well as 5 other analogs and observed that MMV1782387

was amongst the strongest inhibitors of *N. parisii* and displayed the least host toxicity. Our study

further demonstrates the effectiveness of the *C. elegans*-*N. parisii* system for discovering

microsporidia inhibitors and the compounds we identified provide potential scaffolds for anti-

microsporidia drug development.

Introduction

Microsporidia are obligate intracellular pathogens phylogenetically related to the fungi (Keeling,

2009 , Wadi and Reinke, 2020). Approximately 1700 species of microsporidia have been identified

to infect invertebrates, vertebrates, and protists, with at least 17 species infecting human

(Chaijarasphong et al., 2021 , Han et al., 2020 , Murareanu et al., 2021). In addition to posing a

serious threat to human health, microsporidia are responsible for substantial economic losses in

agriculture. Species of farmed penaeid shrimp are susceptible to hepatopancreatic

microsporidiosis, which is caused by *Enterocytozoon hepatopenaei* (Chaijarasphong et al., 2021).

Nosema ceranae and *Nosema apis* are a threat to the global beekeeping industry (Higes et al.,

2013 , Forsgren and Fries, 2010). A lethal pathogen, *Nosema bombycis*, causes heavy losses or

even total culture failure in the silkworm industry (Bhat et al., 2017). Moreover, some species,

such as *Ameson portunus* and *Enterospora nucleophila*, have been identified to infect farmed

crabs and fish, respectively (Wang et al., 2017 , Picard-Sánchez et al., 2020). Among

immunocompromised individuals, *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis* are

the most common microsporidia infections in humans and are associated with diarrhea and

systemic illness (Didier, 2005 , Ruan et al., 2021). In underdeveloped countries, up to 51% of

HIV-infected individuals with diarrhea have microsporidia infection (Gumbo et al., 1999).

Infections with zoonotic microsporidia have been discovered in livestock, companion animals, and

wildlife, posing a risk to public health (Widmer et al., 2010). In recent decades, the NIAID and

CDC have recognized that many of these species of microsporidia pose a threat to human health.

Thus, microsporidia have been classified as Category B priority pathogens for biodefense

research (Didier and Khan, 2014).

There are few effective therapeutic interventions available to treat microsporidia infections (Wei

et al., 2022). Albendazole is known to inhibit microtubule polymerization by binding beta-tubulin

and is effective against a variety of parasites with little adverse effects in humans (Katiyar et al.,

1994 , Horton, 2000). In infections of *Encephalitozoon* spp., albendazole has been demonstrated

to control microsporidiosis (Li et al., 1996). However, several studies have shown that

albendazole has a limited effect on microsporidiosis caused by *E. bieneusi* and *Vittaforma cornea*

and that these species encode beta-tubulin with substitutions associated with resistance (Leder

et al., 1998 , Franzen et al., 2008 , Akiyoshi et al., 2007). Analogs of albendazole, benomyl and

carbendazim, have been used to control microsporidia in insects such as *Nosema heliothidis* in

Heliothis zea and *Nosema kingi* in *Drosophila willistoni* (Wei et al., 2022 , Armstrong, 1976 ,

Brooks et al., 1978). However, microsporidiosis in mammals cannot be treated with benomyl and

carbendazim because of their hepatotoxicity and toxic effects on reproduction (Sakr et al., 2004 ,

Barnes et al., 1983) (Kotková et al., 2017). Since benzimidazoles have been widely used for

decades, resistance fears have arisen. Fumagillin binds specifically and covalently to methionine

aminopeptidase type 2 (MetAP2) and can inhibit many species of microsporidia (Lefkove et al.,

2007 , Sin et al., 1997). However, due to concerns with host toxicity, this drug is not approved for

use in humans (Han and Weiss, 2018). Furthermore, fumagillin has been associated with

concerns regarding its toxicity in agriculture applications (Union, 2010). Therefore, there is a need

to identify additional anti-microsporidia inhibiting agents.

The model organism *Caenorhabditis elegans* has become a useful system in which to study

microsporidia infections and to identify inhibitors. The first microsporidia reported to infect *C.*

elegans was *Nematocida parisii* (Troemel et al., 2008)*.* Infection of *C. elegans* by *N. parisii* begins

when spores are ingested into the worm's intestinal lumen, where they expel their unique invasion

apparatus called the polar tube (Tamim El Jarkass and Reinke, 2020). This causes the

sporoplasm to be deposited inside intestinal cells. The parasite then proliferates intracellularly as

meronts and finally differentiates into spores, which then exit into the intestinal lumen. Infection

of *C. elegans* with *N. parisii* results in smaller body size, reduced reproductive fitness, and

shortened life span (Balla et al., 2015 , Balla et al., 2016 , Willis et al., 2021). This host-parasite

system has become a model in which to study mechanisms of microsporidia invasion, proliferation,

and spore exit (El Jarkass et al., 2022 , Reinke et al., 2017 , Balla et al., 2016 , Szumowski et al.,

2014). Other species of microsporidia have also been found to infect *C. elegans* including

Pancytospora epiphaga, which is related to human-infecting species (Bojko et al., 2022 , Wadi et

al., 2023 , Zhang et al., 2016). We recently described the use of *C. elegans* to screen compounds

for activity towards *N. parisii* using a high-throughput 96-well based assay, resulting in the

identification of inhibitors of microsporidia invasion and proliferation (Han et al., 2020 , El Jarkass

et al., 2022 , Murareanu et al., 2022).

The Pandemic Response Box (PRB) is a promising source of anti-microsporidia compounds.

Developed by the Medicines for Malaria Venture with the support of the Drugs for Neglected

Disease Initiative, this is an open-access compound library consisting of 400 small molecule

compounds that have antifungal, antibacterial and antiviral activity (Samby et al., 2022).

Compounds from this collection include those that can inhibit other types of parasites such as

nematodes, amoebas, and the causative agent of malaria (Shanley et al., 2022 , Reader et al.,

2021 , Rice et al., 2020).

To identify novel microsporidia inhibitors, we screened the PRB using a modified version of our

previously described *C. elegans*-*N. parisii* assay (Murareanu et al., 2022). We screened this

compound collection at two concentrations and identified four chemical inhibitors of microsporidia

infection. We validated these compounds and additionally show that all four compounds reduced the pathogen load of *N. parisii*. We then studied at what stage in the microsporidia life cycle each compound is active. We found that MMV1782387, an analog of albendazole, inhibits microsporidian proliferation. This compound and other albendazole analogs were identified inhibit both *N. parisii* and *P. epiphaga.* We also show that MMV1006203 inhibits spore firing and related flavone analogs also prevent *N. parisii* infection. Together our study identifies additional microsporidia inhibitors that can block microsporidia proliferation and invasion.

Results

Screen of Pandemic Response Box identifies 4 microsporidia inhibitors

To identify compounds from the Pandemic Response Box collection which inhibit microsporidia infection, we adapted our previously described 96-well infection assays (Murareanu et al., 2022). Earliest larval (L1) stage of *C. elegans* were incubated with *N. parisii* spores and compound in liquid for six days at 21°C. Host animals infected with *N. parisii* produce a reduced number of offspring and compounds which inhibit microsporidia can restore the ability of these animals to produce progeny. To quantify the number of offspring produced when incubated with each compound, animals in each well were stained with rose bengal, imaged with a flatbed scanner, and counted using WorMachine (Hakim et al., 2018)(see methods). Each compound was screened in triplicate at a concentration of 100 µM. We identified six compounds that increased progeny production in infected worms by 35-90% relative to the uninfected controls (Fig. 1A). These compounds also increased progeny production by 1.7-8.5-fold compared to infected controls. Four of these compounds significantly improved the production of *C. elegans* progeny in the presence of *N. parisii* (Fig. 1B-D). To determine the effect of the compound collection at a lower concentration, we screened each compound at 40 µM. We observed three compounds which increased progeny production over 3-5.3-fold more than infected controls and to more than 13% of the uninfected controls (Fig. 1E). Two of these compounds resulted in significantly more progeny production at 40 µM and these two compounds were also observed to be significant at 100 µM (1F and 1G). In total we identified four compounds with significant activity which we validated and characterized in subsequent experiments (Fig. 1H).

We next sought to determine whether the four compounds which significantly restored *C. elegans* progeny production also limited *N. parisii* infection. We set up assays similar to our initial screen by culturing L1 worms continuously with *N. parisii* spores in the presence of compounds in 24-

well plates. After four days, worms were fixed and then stained with direct yellow 96 (DY96), which

binds to chitin, a critical component of *N. parisii* spore walls and *C. elegans* embryos (Fig. 2A)

(Zhang et al., 2016 , Vávra and Ronny Larsson, 2014 , Willis et al., 2022). In the presence of *N.*

parisii spores, all four compounds significantly increased the proportion of adult worms containing

embryos (Fig. 2B), which is consistent with the results from our initial screen. Treatment with the

newly discovered compounds and the known microsporidia inhibitor dexrazoxane significantly

reduced animals displaying newly formed *N. parisii* spores, except for MMV1593539 (Fig. 2C)

(Murareanu et al., 2022). To determine if MMV1593539 had any impact on *N. parisii* infection, we

quantified the proportion of worms displaying a low, moderate, or high infection level (See

methods) (Fig. 2D-F). Under these infection conditions, ~90 percent of control worms displayed

high infection levels and all four compounds significantly lowered infection (Fig. 2F).

MMV1782387 inhibits the proliferation of *N. parisii*

The identified compounds could inhibit microsporidia infection through two mechanisms: by

preventing microsporidia from invading cells or by reducing proliferation after invasion. To test

whether compounds limited proliferation, we set up pulse-chase experiments where we infected

worms for three hours, washed away excess spores, and then incubated the worms with one of

the four MMV compounds or with dexrazoxane which was previously shown to limit *N. parisii*

proliferation (Fig. 3A) (Murareanu et al., 2022). Treatment with MMV1782387 and dexrazoxane

increased the gravidity of worms (Fig. 3B). Treatment with MMV1782387 also inhibited *N. parisii*

proliferation with 26.67% of worms having newly formed spores, which is less inhibition than was

observed with dexrazoxane (0.89%) (Fig. 3C).

To determine whether MMV1782387 inhibits *N. parisii* by slowing proliferation or enhancing

parasite clearance, we examined pulse-chase infected animals at either 2 day (before spore

formation) or 4 days (after spore formation) post infection with probes specific for *N. parisii* 18S

RNA (Troemel et al., 2008). Dexrazoxane was previously shown to significantly reduce the

pathogen burden within animals, without influencing the proportion of infected animals

(Murareanu et al., 2022). We observed that MMV1782387, similar to dexrazoxane, did not cause

a reduction in infected animals, but significantly reduced the amount of meronts in the worms (Fig.

3D-3F). None of the other MMV compounds we tested reduced pathogen load of the worms at

either 2- or 4-days post-treatment. Dexrazoxane also caused significantly more reduction in

pathogen levels compared to MMV1782387 (Fig. 3E and 3F). Together our results show

MMV1782387 inhibits microsporidia proliferation but does not enhance parasite clearance.

Spore firing is inhibited by MMV1006203

The initial step of microsporidia invasion is spore germination. Many microsporidia, including *N.*

parisii, germinate (also called spore firing) in the intestinal lumen to initiate infection (Fries et al.,

1996 , Troemel et al., 2008 , Lv et al., 2022). In order to determine whether any compounds

reduce germination, we conducted spore firing assays using the four MMV compounds as well as

ZPCK, which has previously been shown to inhibit spore germination (Murareanu et al., 2022). *N.*

parisii spores were incubated with compounds for 24 hours and the spores were then washed to

remove the compounds. These spores were then cultured with *C. elegans* at the L1 stage for

three hours and stained with FISH to visualize the sporoplasms and DY96 to visualize the spores.

Spore firing was determined by counting the number of spores that did not contain a sporoplasm

divided by the total number of spores. ZPCK and one of the MMV compound, MMV1006203,

significantly reduced the spore firing rate (Fig. 4A). All of the compounds, as well as ZPCK, also

significantly reduced the number of sporoplasms that had invaded each worm (Fig. 4B).

Benzimidazole and flavone analogs inhibit *N. parisii* infection

The benzimidazole molecule albendazole is one of the main treatment options currently used for

microsporidia infection. The structure of MMV1782387 is similar to albendazole and the PRB

contains several other benzimidazoles including carbendazim, fenbendazole, and oxfendazole.

However, these other benzimidazole compounds were not identified in our initial screen. We first

tested whether MMV1782387 and six benzimidazole analogs reduced the reproductive fitness of

uninfected *C. elegans* (Fig. 5G). At 40 μM, there is no difference in the percentage of worms

forming embryos when treated with any of the compounds (Fig. 5A). In contrast, menbendazole

and oxfenfazole significantly reduced progeny production in uninfected animals at 100 μM,

indicating moderate toxicity to the host (Fig. 5D). We then tested whether these benzimidazole

compounds could restore the formation of embryos in animals infected with *N. parisii* using our

continuous infection assays. Except thiabendazole, all of the compounds increased the

percentage of gravid worms in the presence of *N. parisii* at 40 μM (Fig. 5B). At 100 μM,

MMV1782387, albendazole, and carbendazim increased the percentage of gravid worms with *N.*

parisii (Fig. E). MMV1782387-treated worms displayed the largest increase in the percentage of gravid worms. We also examined the percentage of worms with newly formed spores when treated with these compounds. Except thiabendazole, all compounds displayed a reduction in infected animals at 40 µM (Fig. 5C). All compounds displayed a reduction in infected animals at 100 µM, with MMV1782387 displaying amongst the strongest inhibition of infection (Fig. 5F). These results suggest that benzimidazoles can inhibit *N. parisii* and that MMV1782387 shows both strong inhibition of *N. parisii* as well as low host toxicity in *C. elegans*.

The compound we identified which inhibits spore firing, MMV1006203, has a structure similar to flavone. To test whether flavone and an analog of MMV1006203, displurigen, could inhibit *N. parisii*, we tested these compounds in continuous infection assays at a concentration of 40 µM or 100 µM (Fig. 5L). All three of these compounds significantly increased the proportion of gravid worms and reduced infection rates at a concentration of 40 µM (Fig. 5H and 5I). At a concentration of 100 µM, all three molecules were effective at inhibiting microsporidia infection, though only MMV1006203 could significantly increase the gravidity of worms (Fig. 5J and 5K). These results show that molecules based on a flavone structure can inhibit *N. parisii* infection.

Identified MMV compounds inhibit *P. epiphaga*

To test whether the four MMV compounds we identified were effective against other microsporidia species, we tested them against *P. epiphaga*. This species of microsporidia infects the hypodermis and muscle of *C. elegans* and belongs to the *Enterocytozoonida* clade, along with the human pathogens *V. cornea* and *E. bieneusi* (Zhang et al., 2016 , Bojko et al., 2022). In order to examine whether the four inhibitors we identified from the PRB could inhibit *P. epiphaga* infection of *C. elegans*, we used FISH staining to quantify the pathogen load. All of the compounds significantly reduced *P. epiphaga* infection levels (Fig. 6A). *V. cornea* and *E. bieneusi* contain a glutamine at position 198 in beta-tubulin that is associated with albendazole resistance, but *P. epiphaga* beta-tubulin encodes for glutamate at this position which is associated with albendazole sensitivity (Franzen et al., 2008 , Akiyoshi et al., 2007 , Wadi et al., 2023). We tested albendazole against *P. epiphaga* and observed that this compound also inhibited infection (Fig. 6B).

Discussion

We screened the open-access PRB compound library, identifying four compounds with anti-
microsporidia activity. We quantified the ability of compounds to reproducibly alleviate the
reduction in *C. elegans* progeny caused by *N. parisii* infection. We validated all four compounds
identified from the initial screen, which is an improvement over our previous screen where we
qualitatively determined the effect of compounds from a single replicate and only about half of the
initially identified compounds were validated (Murareanu et al., 2022). All four compounds we
identified in this study limited both *N. parisii* and *P. epiphaga* infection. These results further
demonstrate that *C. elegans* can be used to efficiently identify compounds with activity against
multiple species of microsporidia. One limitation of our approach is that inhibitors that reduce the
reproductive fitness of *C. elegans* on their own will not be observed. However, this is also
potentially beneficial as host toxicity, at least within the *C. elegans* context, is evaluated at the
same time as inhibition of microsporidia infection.

There is a range of bioactivity associated with benzimidazoles, including anti-inflammatory,
antihypertensive, anti-bacterial, anti-parasitic, and anti-fungal properties (Labanauskas et al.,
2004 , Navarrete-Vázquez et al., 2010 , Hosamani and Shingalapur, 2011 , Wei et al., 2022). The
benzimidazole albendazole is one of the most common treatments for microsporidia. Here we
show that MMV1782387, a benzimidazole carbamate, has amongst the strongest inhibition of *N.*
parisii and relatively low host toxicity in *C. elegans*. Carbendazim, fenbendazole and oxfendazole
are present in the PRB, however, in our preliminary screening, we did not find that these
compounds improved progeny production in the presence of *N. parisii*. Beta-tubulin is the likely
target of albendazole in microsporidia, and given the similarity of the compounds, MMV1782387
may also inhibit beta-tubulin. In compound library screens, multistage activity is one of the most
preferred attributes for antipathogens (Poonam et al., 2018). We show that MMV1782387, which
inhibits proliferation of *N. parisii* can also reduce invasion by about 50%. MMV1782387 has been
shown to be effective against several fungal pathogens which cause eumycetoma and may have
potential for further development as an inhibitor of fungal pathogens (Lim et al., 2022).

Microsporidia infect host cells through spore germination, and this process may be regulated by
receptor proteins on cell membranes and external signals (Undeen, 1990 , Willis and Reinke,
2022). Furthermore, the changes in calcium ion concentrations, osmotic pressure of the external
medium, or *in vivo* host environments can induce microsporidia spore firing (Weidner and Byrd,
1982 , Pleshinger and Weidner, 1985 , UNDEEN and FRIXIONE, 1990). However, many of the
microsporidia proteins involved in spore firing are unknown. The subtilisin-like protease NbSLP1
has been implicated in germination as the active version localizes to the site of the spore where

polar tube firing occurs in *N. bombycis* spores (Dang et al., 2013) (Wang et al., 2022). We

previously found that protease inhibitors and quinones can inhibit spore firing (Murareanu et al.,

2022). Here we show that MMV1006203, a flavone, can prevent microsporidia invasion and spore

firing. This inhibition occurs after incubation of the compound with spores, followed by washing of

the spores, suggesting that this inhibitor acts directly on the spores to prevent firing. The molecular

structures of displurigen and flavone are similar to that of MMV1006203, and these compounds

also display the ability to prevent *N. parisii* infection. Other flavone compounds such as quercetin

were shown to inhibit *Encephalitozoon intestinalis* infection, but it was not determined if these

compounds block microsporidia invasion (Mead and McNair, 2006). MMV1006203 was first

shown to inhibit the human cytomegalovirus (HCMV) protease and more recently to have activity

against *Plasmodium falciparum* (Reader et al., 2021 , Dhanak et al., 1998). Further work will be

necessary to determine the molecular target of MMV1006203 in preventing microsporidia spore

firing.

Acknowledgements

We are grateful to Winnie Zhao, Yin Chen Wan, Meng Xiao, Jonathan Tersigni, Hala Tamim El

Jarkass, and Edward James for providing helpful comments on the manuscript. We thank the

Medicines for Malaria Venture for providing the PRB compound library and the individual

compounds to retest. **Funding:** This work was supported by Canadian Institutes of Health

Research grant (no. 461807 to A. W. R.) and Q. H was supported by an award from the China

Scholarship Council. **Competing interests:** The authors declare that they have no competing

interests. **Data availability:** All data is presented in Data S1.

Material and Methods

C. elegans maintenance

The food source for *C. elegans*, *Escherichia coli* OP50-1, was grown to saturation in lysogeny

broth (LB) for 18 hours at 37°C. To generate animals for infection assays, the wild-type *C. elegans*

strain N2 was grown as a mixed population and L4 stage worms were picked onto a 10 cm

nematode growth media (NGM) plates seeded with 10x OP50-1 *E. coli* and kept at 21°C for 4

days (Lewis and Fleming, 1995). In order to synchronize the worms, M9 solution was used to

remove the worms from the NGM plates, sodium hypochlorite and sodium hydroxide were used

to bleach them. Embryos from gravid adults were released into the solution and after washing
were incubated at 21°C for 18 to 24 hours until the embryos hatched.

N. parisii spore preparation

N. parisii (ERTm1) spores were prepared as described previously (Willis et al., 2021). *C. elegans*
N2 worms were infected with *N. parisii* spores on NGM plates. Worms were incubated for several
days to generate a large population of infected worms, which were harvested and frozen at -80°C.
Zirconia beads (2 mm diameter) were used for mechanical disruption of the infected worms,
followed by the removal of embryos, larvae, and debris using a 5 µm filter (Millipore). *N. parisii*
spore preparations were confirmed to be free of contaminated bacteria and stored at -80°C. Spore
concentration was measured by counting DY96-stained spores using a sperm counting slide
(Cell-VU).

Source of chemicals

Medicines for Malaria Venture (MMV, Geneva, Switzerland) provided 400 compounds dissolved
in 10 µL of DMSO at a concentration of 10 mM for inclusion in the Pandemic Response Box (PRB).
2 µL of the stock compounds were transferred to new plates containing 3 µL DMSO to generate
4 mM stocks. For retesting, the individual compounds of MMV1006203, MMV1593539,
MMV1634497 and MMV1782387 were a gift from MMV. The analog compounds of albendazole,
thiabendazole, carbendazim, oxfendazole, mebendazole, fenbendazole, flavone and displurigen
were purchased from MilliporeSigma. Stocks of all compounds were stored at -80 °C.

Phenotypic assays in 96-well plates to identify microsporidia inhibitors

Previously described methods were adapted to quantify the ability of compounds to restore the
ability of *C. elegans* to produce progeny in the presence of *N. parisii* (Murareanu et al., 2022).
Each well of a 96-well plate was filled with 25 µL of K-medium (51 mM NaCl, 32 mM KCl, 3 mM
CaCl$_2$, 3 mM MgSO$_4$, 3.25 µM cholesterol) containing 5x OP-50 and *N. parisii* spores, with the
exception of column 12 to which spores were not added. 25 µL of K-medium containing L1 worms
was then added to each well. 500 nL of compounds from the PRB were pinned into columns 2–
11 using a 96-well pinning tool manufactured by V&P Scientific. Additionally, 500 nL of DMSO
was added to columns 1 and 12 for infected and uninfected controls, respectively. Each well
contained 100 bleach-synchronized L1 worms, 1% DMSO, 15000 spores/µL *N. parisii* and 100
µM or 40 µM of compounds. A breathable adhesive porous film was used to cover the 96-well
plates, which were placed inside humidity boxes wrapped in parafilm and incubated for 6 days at

21°C with shaking at 180 rpm. Each compound was tested three times at both concentrations,

with the exception of 17 compounds for which there was not sufficient amounts at which to test at

100 µM and 9 compounds there was not sufficient amounts at which to test at 40 µM (See Data

S1).

Quantification of progeny production

Following incubation, 10 µL of 0.3125 mg/mL Rose Bengal solution was added to each well using

an Integra VIAFLO 96 Electronic pipette. Plates were then wrapped in parafilm and incubated for

16–24 hours at 37 °C, resulting in magenta staining of the worms. 240 µL M9/0.1%Tween 20

was added to each well and the plate and centrifuged for 1 minute at 2200 x g. 200 µL supernatant

was removed from each well and 150 µL of M9/0.1%Tween 20 was added to each well. Upon

mixing up the worms in the plate, 25 µL of the worms were transferred to 96-well white clear

bottom plates containing 300 µL M9/0.1%Tween 20. After 30 minutes, plates were scanned using

an Epson Perfection V850 Pro flat-bed scanner with the following settings: positive film-holder,

4800 dpi, and 24-bit color. In order to highlight stained worms, images were also modified using

GIMP version 2.8.18, with horizontal and vertical gridlines positioned such that each well is

separated by a grid and removing HTML color codes #000000 and #FFC9AF. Images were also

modified by applying unsharp masking with the following parameters (radius = 10, effect = 10,

threshold = 10). Hue saturation was adjusted by changing the lightness to 100 and the saturation

to -100 for yellow, blue, cyan, and green. For red and magenta, the lightness was changed to -

100 and the saturation to 100. Each well was exported as a single .tiff image using LZW

compression. MATLAB was used to run WorMachine (Hakim et al., 2018) with the pixel

binarization threshold set to 30, the neighboring threshold to set to 1, and the maximum object

area set toto 0.003%.

Continuous infection assays

24-well assay plate containing a total volume of 400 µL including800 L1 worms and 15,000 *N.*

parisii spores/µL were used for continuous infection assays. Assays were performed for three

biological replicates using 100 µM of each compound except for dexrazoxane (60 µM). During the

incubation period, test plates were covered with breathable adhesive porous film, the boxes were

enclosed in parafilm, and the plates were incubated at 21 °C, with shaking at 180 rpm for four

days. Incubated samples were washed with M9/0.1%Tween 20, acetone-fixed, DY96-stained,

and analyzed by fluorescence microscopy.

Pulse infection assays

To generate infected worms, ~8000 bleach-synchronized L1 worms, 30 million *N. parisii* spores,

and 5 µL 10x OP50-1 were added to 6 cm NGM plates and incubated for three hours at 21 °C

after drying. To remove excess spores, the worms were washed twice with 5 mL M9/0.1%Tween

20. Worms were then added to 24-well plates and set up as described in the continuous infection

assays, with the exception that no spores were added. For each of the biological replicates, three

wells were assayed for each compound. After incubation for 2 or 4 days as described above,

samples were fixed in acetone and stained with DY96 and a FISH probe as described below.

Spore firing assays

Spores at a concentration of 30000 spores/µL were incubated for 24 h at 21 °C with compounds

at a concentration of 200 µM, except for ZPCK was at 120 µM, and 2% DMSO. After being washed

three times with 1 mL K-medium, the spores were used in the 24-well assay plates as described

above. The final concentrations in these assays were 15000 spores/ µL, 100 µM compounds

except ZPCK (60 µM), and 1% DMSO. Each compound was tested in three biological replicates

in all assays. Incubation was performed as described above and after 3 h, samples were fixed in

acetone, stained with FISH and DY96, and examined by fluorescence microscopy.

DY96 staining, fluorescence in situ hybridization (FISH), and fluorescence microscopy

To remove excess OP50, samples were washed twice in 1 mL M9/0.1%Tween 20. They were

fixed in 700 µL acetone for 15 minutes or 500 µL PFA solution (4% PFA, 1x PBS, 0.1% Tween-

20) for 30 minutes. Then, samples were washed twice in PBS/0.1%Tween 20. For DY96 staining,

500 µL DY96 staining solution (10 µg/µL DY96, 0.1% SDS in 1xPBS + 0.1% Tween-20) was

added and samples were rotated for 30 minutes. EverBrite™ Mounting Medium with DAPI was

then added to the samples. FISH was performed using the microB FISH probe for *N. parisii* 18S

rRNA (ctctcggcactccttcctg) conjugated to Cal Fluor Red 610 (LGC Biosearch Technologies)

(Troemel et al., 2008). After washing with PBS/0.1%Tween 20, samples were incubated in

hybridization buffer (900 mM NaCl, 20 mM pH=8 Tris HCl, 0.01% SDS) containing 5 ng/µL FISH

probe at 46 °C for 1-6 hours. Samples were washed once with 1 mL wash buffer (50 mL

hybridization buffer + 5 mM EDTA). These samples were also stained with DY96 as described

above, except with DY96 at 20 µg/µL. Samples were imaged using a ZEISS Axio Imager 2 at 5x–

63x magnification and images were captured using Zeiss Zen 2.3. Gravid worms were defined as

the proportion of animals containing any number of embryos. Infected worms were defined as the

proportion of animals displaying any newly formed spores. Low infection was defined as spores

present in less than one half of an animal, moderate infection was defined as spore present in

half of an animal, and high infection was define as spores present throughout both halves of an

animal. L1 progeny of the parents could be distinguished in size and were not included in these

measurements.

P. epiphaga infection assays

P. epiphaga strain JUm1396 spores were prepared similar to as described above for *N. parisii*.

For infection experiments with *P. epiphaga*, 24-well assay plates were set up to contain a final

volume of 400 µL in K-medium, including 800 L1 worms and 80,000 P. epiphaga spores /µL. All

compounds were at a concentration of 100 µM, except for dexrazoxane which was used at 60

µM. The final concentration of DMSO was 1%. Plates were incubated as described above for four

days. The samples were fixed in 4% PFA and stained by the FISH probe specific to *P. epiphaga*

18S RNA (CAL Fluor Red 610CTCTATACTGTGCGCACGG). Fluorescence microscopy and

quantification of FISH fluorescence were performed as described above.

Statistical analyses

The data were collected from three independent experiments with 3 biological repetition and

analyzed by GraphPad Prism. The two groups were compared using the student's t-tests.

Statistical significance was defined as $p<0.05$, *$p<0.01$, ***$p<0.001$, and ****$p<0.0001$.

Figures and legends

Fig. 1. Four compounds from the Pandemic Response Box restored *C. elegans* progeny production in the presence of *N. parisii*. A, E Compounds at concentration of 100 µM (A) or 40 µM (E) were incubated with *C. elegans* and *N. parisii* for 6 days. Each point represents the mean progeny production of a compound expressed as the percentage of the DMSO uninfected control. The compound-ID is shown for compounds that had an activity of at least 35% (A) or ≥ 13% (E). Compounds are divided into their disease area as classified by the PRB and colored according to the legend at the right. 17 compounds in the collection were not screened at 100 µM and 9 compounds not screened at 40 µM due to lack of material (Data S1). **B-D** Compounds that had an activity of at least 35% at 100 µM. (B) Antifungals, (C) Antibacterials, (D) Antivirals. **F,G** Compounds that had an activity ≥ 13% at 40 µM. (F) Antifungals, (G) Antibacterials. **H** The chemical structures of the four PRB compounds with significant activity. Statistical significance was determined by Student's t-test with comparisons to the DMSO infected control Means ± SD (horizontal bars) are shown. (*p < 0.05, **p < 0.01, ***p < 0.001, ns means not significant).

Fig. 2. Identified MMV compounds inhibit *N. parisii*. A-F L1 stage animals with incubated in the presence of *N. parisii* spores and compounds for 4 days. Animals were then fixed and stained with DY96 and DAPI. Microsporidian spores and nematode embryos are stained green by DY96. **A** Representative images of continuous infection assays taken at a magnification of 50x; scale bars are 500 μm. (First and second columns) Worms incubated without (-spore) or with spores (+spore). As a result of microsporidia infection, fewer worms become gravid, and new spores are formed. Microsporidia spores and nematode embryos are stained green by DY96. (Third to Seventh column) Dexrazoxane and the identified PRB compounds inhibit spore formation and restore embryo production. **B** The percentage of worms with embryos (n = 3, N = ≥ 100 worms counted per biological replicate). **C** The percentage of worms with newly formed spores (n = 3, N = ≥ 100 worms counted per biological replicate). **D-F** The percentage of worms with (D) low-level infection, (E) moderate infection, and (F) severe infection. The P values were determined by one-way ANOVA with post hoc test. Means ± SD (horizontal bars) are shown. (***p < 0.001, ****p < 0.0001, ns means not significant).

Fig.3. MMV1782387 inhibits microsporidia proliferation. A-F L1 stage animals were incubated in the presence of *N. parisii* spores for 3 hours and then washed to remove excess spores. Compounds were then added and animals were incubated for 2 (E) or 4 (A-D, F) days, fixed, and stained with DY96, DAPI and a FISH probe specific to the *N. parisii* 18S rRNA. **A** Representative images of pulse infection assays taken at a magnification of 50x; scale bars are 500 μm. (First and second columns) Worms uninfected or infected with *N. parisii* spores. Sporoplasms and meronts are stained in red with FISH probes. (Third and fourth column) Dexrazoxane or MMV1782387 treatment reduces *N. parisii* meronts. **B** The percentage of worms with embryos (n = 3, N = ≥ 100 worms counted per biological replicate). **C** The percentage of worms with newly formed spores (n = 3, N = ≥ 100 worms counted per biological replicate). **D** The percentage of worms with FISH signal (n = 3, N = ≥ 100 worms counted per biological replicate). **E, F** Quantitation of pathogen load (FISH fluorescence area %) per worm for (E) 2 or (F) 4 days post infection (n = 3, N = 10 animals quantified per biological replicate). The P values were determined by one-way ANOVA with post hoc test. Means ± SD (horizontal bars) are shown. (*p <0.05, ***p < 0.001, ****p < 0.0001, ns means not significant).

Fig.4. MMV1006203 prevent microsporidia infection by inhibiting spore firing. A, B *N. parisii*
spores were incubated with compounds for 24 hours and then washed to remove compounds.
Spores were then incubated with L1 stage worms for 3 hours, fixed, and stained with DY96 and
N. parisii 18S rRNA FISH probe. **A** The percentage of fired spores in the intestinal lumen (n = 3,
N = ≥ 50 spores counted per biological replicate). **B** The mean number of sporoplasms per worm
(n = 3, N = ≥ 50 worms counted per biological replicate The P values were determined by one-
way ANOVA with post hoc test. Means ± SD (horizontal bars) are shown. (***p < 0.001, ****p <
0.0001, ns means not significant).

Fig.5. Benzimidazole and flavone analogs limit *N. parisii* infection. A-F, H-K L1 stage

animals were continuously incubated with *N. parisii* spores and indicated compounds for 4 days,

fixed, and stained with DY96 and DAPI. **A-F** Effect of benzimidazoles at 40 µM (A-C) or 100 µM

(D-F) on the percentage of worms with embryos in (A, D) uninfected, (B, E) infected worms and

(C, F) the percentage of worms with newly formed spores (n = 3, N = ≥ 100 worms counted per

biological replicate). **G** Chemical structures of benzimidazole analogs. **H-K** Effect of flavones at

40 µM (H-I) or 100 µM (J-K) on the percentage of worms with embryos (H, J) and the

percentage of worms with newly formed spores (I, K) (n = 3, N = ≥ 100 worms counted per

biological replicate). **L** Chemical structures of flavone analogs. The P values were determined

by one-way ANOVA with post hoc test. Means ± SD (horizontal bars) are shown. (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, ns means not significant).

Fig. 6. *P. epiphaga* **infection is impeded by the identified inhibitors. A,B** L1 stage animals

were continuously incubated with *P. epiphaga* spores for four days, fixed, and stained with a *P.*

epiphaga 18S rRNA FISH probe. Quantification of pathogen load (%FISH fluorescence area) per

worm 4 days post infection (n = 3, N = 10 animals quantified per biological replicate). The P values

were determined by one-way ANOVA with post hoc test. Means ± SD (horizontal bars) are shown.

(*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

References

AKIYOSHI D E, WEISS L M, FENG X, et al. 2007. Analysis of the β-Tubulin Genes from Enterocytozoon bieneusi Isolates from a Human and Rhesus Macaque. 54: 38-41.

ARMSTRONG E J J O I P 1976. Fumidil B and benomyl: chemical control of Nosema kingi in Drosophila willistoni. 27: 363-366.

BALLA K M, ANDERSEN E C, KRUGLYAK L, et al. 2015. A wild C. elegans strain has enhanced epithelial immunity to a natural microsporidian parasite. 11: e1004583.

BALLA K M, LUALLEN R J, BAKOWSKI M A, et al. 2016. Cell-to-cell spread of microsporidia causes Caenorhabditis elegans organs to form syncytia. 1: 1-6.

BARNES T B, VERLANGIERI A J, WILSON M C J T 1983. Reproductive toxicity of methyl-1-(butylcarbamoyl)-2-benzimidazole carbamate (benomyl) in male Wistar rats. 28: 103-115.

BHAT I, BUHROO Z, BHAT M J I J E R 2017. Microsporidiosis in silkworms with particular reference to mulberry silkworm (Bombyx Mori L.). 2: 1-9.

BOJKO J, REINKE A W, STENTIFORD G D, et al. 2022. Microsporidia: a new taxonomic, evolutionary, and ecological synthesis.

BROOKS W, CRANFORD J, PEARCE L J J O I P 1978. Benomyl: effectiveness against the microsporidian Nosema heliothidis in the corn earworm, Heliothis zea. 31: 239-245.

CHAIJARASPHONG T, MUNKONGWONGSIRI N, STENTIFORD G D, et al. 2021. The shrimp microsporidian Enterocytozoon hepatopenaei (EHP): Biology, pathology, diagnostics and control. 186: 107458.

DANG X, PAN G, LI T, et al. 2013. Characterization of a subtilisin-like protease with apical localization from microsporidian Nosema bombycis. 112: 166-174.

DHANAK D, KEENAN R M, BURTON G, et al. 1998. Benzothiopyran-4-one based reversible inhibitors of the human cytomegalovirus (HCMV) protease. 8: 3677-3682.

DIDIER E S, KHAN I A J M P O O 2014. The immunology of microsporidiosis in mammals. 307-325.

DIDIER E S J A T 2005. Microsporidiosis: an emerging and opportunistic infection in humans and animals. 94: 61-76.

EL JARKASS H T, MOK C, SCHERTZBERG M R, et al. 2022. An intestinally secreted host factor promotes microsporidia invasion of C. elegans. 11: e72458.

FORSGREN E, FRIES I J V P 2010. Comparative virulence of Nosema ceranae and Nosema apis in individual European honey bees. 170: 212-217.

FRANZEN C, SALZBERGER B J A A, CHEMOTHERAPY 2008. Analysis of the β-tubulin gene from Vittaforma corneae suggests benzimidazole resistance. 52: 790-793.

FRIES I, FENG F, DA SILVA A, et al. 1996. Nosema ceranae n. sp.(Microspora, Nosematidae), morphological and molecular characterization of a microsporidian parasite of the Asian honey bee Apis cerana (Hymenoptera, Apidae). 32: 356-365.

GUMBO T, SARBAH S, GANGAIDZO I T, et al. 1999. Intestinal parasites in patients with diarrhea and human immunodeficiency virus infection in Zimbabwe. 13: 819-821.

HAKIM A, MOR Y, TOKER I A, et al. 2018. WorMachine: machine learning-based phenotypic analysis tool for worms. 16: 1-11.

HAN B, TAKVORIAN P M, WEISS L M J F I M 2020. Invasion of host cells by microsporidia. 11: 172.

HAN B, WEISS L M J E O O T T 2018. Therapeutic targets for the treatment of microsporidiosis in humans. 22: 903-915.

HIGES M, MEANA A, BARTOLOMé C, et al. 2013. N osema ceranae (Microsporidia), a controversial 21st century honey bee pathogen. 5: 17-29.

HORTON J J P 2000. Albendazole: a review of anthelmintic efficacy and safety in humans. 121: S113-S132.

521   HOSAMANI K M, SHINGALAPUR R V J A D P 2011. Synthesis of 2‑mercaptobenzimidazole derivatives as
522   potential anti‑microbial and cytotoxic agents. 344: 311-319.
523   KATIYAR S, GORDON V, MCLAUGHLIN G, et al. 1994. Antiprotozoal activities of benzimidazoles and
524   correlations with beta-tubulin sequence. 38: 2086-2090.
525   KEELING P J P P 2009. Five questions about microsporidia. 5: e1000489.
526   KOTKOVá M, SAK B, HLáSKOVá L, et al. 2017. The course of infection caused by Encephalitozoon cuniculi
527   genotype III in immunocompetent and immunodeficient mice. 182: 16-21.
528   LABANAUSKAS L, BRUKŠTUS A, GAIDELIS P, et al. 2004. Synthesis and antiinflammatory activity of some
529   new 1-acyl derivatives of 2-methylthio-5, 6-diethoxybenzimidazole. 34: 353-355.
530   LEDER K, RYAN N, SPELMAN D, et al. 1998. Microsporidial disease in HIV-infected patients: a report of 42
531   patients and review of the literature. 30: 331-338.
532   LEFKOVE B, GOVINDARAJAN B, ARBISER J L J E R O A-I T 2007. Fumagillin: an anti-infective as a parent
533   molecule for novel angiogenesis inhibitors. 5: 573-579.
534   LEWIS J A, FLEMING J T J M I C B 1995. Basic culture methods. 48: 3-29.
535   LI J, KATIYAR S K, HAMELIN A, et al. 1996. Tubulin genes from AIDS-associated microsporidia and
536   implications for phylogeny and benzimidazole sensitivity. 78: 289-295.
537   LIM W, NYUYKONGE B, EADIE K, et al. 2022. Screening the pandemic response box identified
538   benzimidazole carbamates, Olorofim and ravuconazole as promising drug candidates for the
539   treatment of eumycetoma. 16: e0010159.
540   LV Q, ZHOU B, LIAO H, et al. 2022. Proteomic profile of polar filament and polar tube from fungal
541   pathogen microsporidium Nosema bombycis provides new insights into its unique invasion
542   organelle. 263: 104617.
543   MEAD J R, MCNAIR N J F M L 2006. Antiparasitic activity of flavonoids and isoflavones against
544   Cryptosporidium parvum and Encephalitozoon intestinalis. 259: 153-157.
545   MURAREANU B M, ANTAO N V, ZHAO W, et al. 2022. High-throughput small molecule screen identifies
546   inhibitors of microsporidia invasion and proliferation in C. elegans. 13: 5653.
547   MURAREANU B M, SUKHDEO R, QU R, et al. 2021. Generation of a microsporidia species attribute
548   database and analysis of the extensive ecological and phenotypic diversity of microsporidia. 12:
549   10.1128/mbio. 01490-01421.
550   NAVARRETE-VáZQUEZ G, HIDALGO-FIGUEROA S, TORRES-PIEDRA M, et al. 2010. Synthesis, vasorelaxant
551   activity and antihypertensive effect of benzo [d] imidazole derivatives. 18: 3985-3991.
552   PICARD-SáNCHEZ A, PIAZZON M C, AHMED N H, et al. 2020. Enterospora nucleophila (Microsporidia) in
553   gilthead sea bream (Sparus aurata): Pathological effects and cellular immune response in natural
554   infections. 57: 565-576.
555   PLESHINGER J, WEIDNER E J T J O C B 1985. The microsporidian spore invasion tube. IV. Discharge
556   activation begins with pH-triggered Ca2+ influx. 100: 1834-1838.
557   POONAM, GUPTA Y, GUPTA N, et al. 2018. Multistage inhibitors of the malaria parasite: Emerging hope
558   for chemoprotection and malaria eradication. 38: 1511-1535.
559   READER J, VAN DER WATT M E, TAYLOR D, et al. 2021. Multistage and transmission-blocking targeted
560   antimalarials discovered from the open-source MMV Pandemic Response Box. 12: 269.
561   REINKE A W, BALLA K M, BENNETT E J, et al. 2017. Identification of microsporidia host-exposed proteins
562   reveals a repertoire of rapidly evolving proteins. 8: 14023.
563   RICE C A, TROTH E V, RUSSELL A C, et al. 2020. Discovery of anti-amoebic inhibitors from screening the
564   MMV pandemic response box on Balamuthia mandrillaris, Naegleria fowleri, and Acanthamoeba
565   castellanii. 9: 476.
566   RUAN Y, XU X, HE Q, et al. 2021. The largest meta-analysis on the global prevalence of microsporidia in
567   mammals, avian and water provides insights into the epidemic features of these ubiquitous
568   pathogens. 14: 1-14.

SAKR S A, SAMEI H A, SOLIMAN M J J M S 2004. Exploring hepatotoxicity of benomyl: histological and
histochemical study on albino rats. 4: 77-83.
SAMBY K, BESSON D, DUTTA A, et al. 2022. The pandemic response Box— Accelerating drug discovery
efforts after disease outbreaks. 8: 713-720.
SHANLEY H T, TAKI A C, BYRNE J J, et al. 2022. A high-throughput phenotypic screen of the 'Pandemic
Response Box'identifies a quinoline derivative with significant anthelmintic activity. 15: 257.
SIN N, MENG L, WANG M Q, et al. 1997. The anti-angiogenic agent fumagillin covalently binds and
inhibits the methionine aminopeptidase, MetAP-2. 94: 6099-6103.
SZUMOWSKI S C, BOTTS M R, POPOVICH J J, et al. 2014. The small GTPase RAB-11 directs polarized
exocytosis of the intracellular pathogen *N. parisii* for fecal-oral transmission from *C. elegans*.
111: 8215-8220.
TAMIM EL JARKASS H, REINKE A W J C M 2020. The ins and outs of host–microsporidia interactions
during invasion, proliferation and exit. 22: e13247.
TROEMEL E R, FéLIX M-A, WHITEMAN N K, et al. 2008. Microsporidia are natural intracellular parasites
of the nematode Caenorhabditis elegans. 6: e309.
UNDEEN A H, FRIXIONE E J T J O P 1990. The role of osmotic pressure in the germination of nosema
algerae spores 1. 37: 561-567.
UNDEEN A H J J O T B 1990. A proposed mechanism for the germination of microsporidian (Protozoa:
Microspora) spores. 142: 223-235.
UNION E C J O J E 2010. Commission Regulation (EU) No 37/2010 of 22 December 2009 on
pharmacologically active substances and their classification regarding maximum residue limits in
foodstuffs of animal origin. 15: 1-72.
VáVRA J, RONNY LARSSON J J M P O O 2014. Structure of microsporidia. 1-70.
WADI L, EL JARKASS H T, TRAN T D, et al. 2023. Genomic and phenotypic evolution of nematode-
infecting microsporidia. 19: e1011510.
WADI L, REINKE A W J P P 2020. Evolution of microsporidia: An extremely successful group of eukaryotic
intracellular parasites. 16: e1008276.
WANG R, LI Q, LIU F, et al. 2022. Maturation of subtilisin-like protease NbSLP1 from microsporidia
Nosema bombycis. 1126.
WANG Y, LI X-C, FU G, et al. 2017. Morphology and phylogeny of Ameson portunus n. sp.(Microsporidia)
infecting the swimming crab Portunus trituberculatus from China. 61: 122-136.
WEI J, FEI Z, PAN G, et al. 2022. Current therapy and therapeutic targets for microsporidiosis. 13.
WEIDNER E, BYRD W J T J O C B 1982. The microsporidian spore invasion tube. II. Role of calcium in the
activation of invasion tube discharge. 93: 970-975.
WIDMER G, AKIYOSHI D E J I, GENETICS, EVOLUTION 2010. Host-specific segregation of ribosomal
nucleotide sequence diversity in the microsporidian Enterocytozoon bieneusi. 10: 122-128.
WILLIS A R, EL JARKASS H T, REINKE A W J J 2022. Studying Inherited Immunity in a Caenorhabditis
elegans Model of Microsporidia Infection. e63636.
WILLIS A R, REINKE A W. Factors That determine Microsporidia infection and host
specificity[C]//Microsporidia: Current Advances in Biology.Springer,2022:91-114.
WILLIS A R, ZHAO W, SUKHDEO R, et al. 2021. A parental transcriptional response to microsporidia
infection induces inherited immunity in offspring. 7: eabf3114.
ZHANG G, SACHSE M, PREVOST M-C, et al. 2016. A large collection of novel nematode-infecting
microsporidia and their diverse interactions with Caenorhabditis elegans and other related
nematodes. 12: e1006093.