Characterization of Turkish Pine Honey and Differentiation from other floral honey types

from Oceania, Aegean, West Coast by NMR spectroscopy and Chemometric Analysis

Kerem Kahraman[1,#], Oktay Göcenler[1,#], Çağdaş Dağ[1, 2,*]

[1]Nanofabrication and Nanocharacterization Center for Scientific and Technological Advanced

Research (n[2]STAR), Koc University, İstanbul, Turkey

[2]Koc University Isbank Center for Infectious Diseases (KUISCID), Koc University, Istanbul,

Turkey

*Corresponding author: cdag@ku.edu.tr

#equal contribution

Abstract

Honey is a viscous, supersaturated sugar solution produced by bees through the enzymatic transformation of nectar from flowers, containing a complex mixture of carbohydrates, organic acids, enzymes, and other minor constituents. Although honey has been used for thousands of years for its nutritional and medicinal properties, it has been the subject of increasing attention in recent years due to increasing adulterated honey production. Consequently, assessment of honey quality and authenticity has become essential to ensure consumer confidence of regional honey and to perverse the practice of authentic honey production. In this study, we employed nuclear magnetic resonance (NMR) spectroscopy and chemometric analysis to characterize Turkish Pine honey and compare it to flower honey originating from the Oceania, Aegean, and West Coast of North America regions. Utilizing ^{1}H NMR spectroscopy, the chemical profile of Turkish Pine honey was characterized, and unique peaks were found. Additionally, PLS-DA statistical analysis was employed to further investigate the distinction of Turkish Pine honey among other floral and regional honeys. Upon completion of our statistical analysis, we were able to effectively distinguish Turkish Pine honey from other regional honey types, allowing us to formulate a universal test for authenticity of Turkish Pine Honey.

Introduction

Honey, produced by honeybees (Apis mellifera), is a multifaceted product that has captivated human interest for millennia, celebrated for its nutritional, therapeutic, and cultural properties. Revered since ancient times, it has served humanity as both nourishment and remedy. The transformation of nectar or honeydew into honey constitutes a complex biochemical orchestration, mediated by enzymes, diastase (amylase) and invertase (α-glucosidase) (Rossano et al., 2012). Together with the intrinsic ingredients of nectar or honeydew, these enzymes give rise to a composite rich in sugars, lipids, amino acids, vitamins, minerals, and phenolic compounds (Valverde et al., 2022). It is this unique constellation of constituents that gives honey with its reputed anti-inflammatory, antioxidant, antiproliferative, and antimicrobial properties (Cianciosi et al., 2018). The chemical composition of honey displays considerable variability, influenced by factors such as geographical and botanical origin, bee species, as well as yearly and seasonal conditions. Even within the same region, distinctive seasonal periods may lead to fluctuations in both the quantity and quality of honey (Tsuruda, Chakrabarti and Sagili, 2021; Walker et al., 2022). On a global scale, Turkey ranks second only to China, boasting an annual honey production capacity of 100,000 tons, encompassing diverse types such as mono-floral, multi-floral, honeydew, and pine honey (Güngör and Sen, 2018; Erdal and Tipi, 2022; Özkök, Yüksel and Sorkun, 2022; Parin et al., 2021).

Among these honey types, pine honey presents a particularly intriguing case. This unique honeydew honey, predominantly found in the East Mediterranean region (Turkey and Greece), is produced from the ripening of secretions within the honeycomb (Tananaki, Thrasyvoulou, Giraudel, and Montury, 2007). The source of these secretions is the pine beetle, Marchalina

hellenica (Gennadius), which deposits them on pine trees. Bees subsequently collect these

secretions, infuse them with their own enzymes, and store them in beehives generating pine honey.

Notably, the distinctiveness of pine honey is not limited to its origin and production. It has an

appealing resistance to heat, making it more sought after in certain markets, especially in Turkey

and Greece, when compared to flower honey. The medical properties and composition of pine

honey have been the focus of intensive study by several research groups, uncovering a rich array

of bioactive compounds (Tsavea et al., 2022; Özkök et al., 2010). Particularly, Pine honey is

recognized for its antiradical effects, attributed to its phenolic compounds and amino acids.

Demonstrated by Eraslan et al. (2010), the antioxidant and antiradical attributes of pine honey are

well-established. Further investigations have revealed its potential antiproliferative impact, with

studies highlighting the effectiveness of pine honey's n-butanol and ethyl acetate extracts on

various cancerous cell lines, including cervical, brain tumor, prostate, colon, and breast (Sıcak,

Şahin-Yağlıoğlu, Öztürk, 2021).

Beyond these properties, pine honey also exhibits antioxidant, anti-inflammatory, and

antimicrobial effects. A recent study employed RNA-seq analysis to unravel how pine honey

influences the transcriptomic profile of Pseudomonas aeruginosa, an important soil bacterium.

This analysis illuminated pine honey's inhibitory action on the expression of the P. aeruginosa

genome (Kafantaris et al., 2021). Further extending the understanding of pine honey's composition,

Duru et al. (2021) conducted a deep investigation into its volatile components. Their research led

to the identification of 32 distinct volatile compounds.

The authenticity of honey is intrinsically linked to its geographical and botanical origin, two vital

factors that significantly influence its quality, production, and market price. As with all honey

varieties, pine honey must adhere to rigorously defined quality standards before it reaches the

consumer. Ensuring authenticity necessitates accurate and precise analytical methods to determine

honey's origin, thereby enhancing food quality and minimizing adulteration. Various techniques

are employed in assessing honey quality, ranging from instrumental analyses for sugars and

Hydroxymethylfurfural (HMF) to classical methodologies like reflectometry and titrations, and

extending to more advanced techniques such as chromatographic analyses, physicochemical and

sensory evaluations, vibrational and fluorescence methods, elemental and isotopic assessments,

and molecular and direct MS techniques (Tsagkaris et al., 2021). Among the critical

physicochemical parameters serving as authenticity markers for geographical and botanical

discrimination are phenolic compounds, color, viscosity, moisture, pH, electrical conductivity,

sugar content, water-insoluble matter, acidity, and HMF (Živkov Baloš et al., 2023). The

interpretation of these parameters is further supported by supervised and unsupervised multivariate

data analysis methods. Notably, principal component analysis (PCA) and hierarchical cluster

analysis (HCA) serve as unsupervised methods, while partial least squares discriminant analysis

(PLS-DA) and partial least squares regression (PLSR) function as supervised techniques

(Tsagkaris et al., 2021; Walker et al., 2022).

Nuclear Magnetic Resonance (NMR) spectroscopy has emerged as a particularly advantageous

tool in this context. Its value lies in its comprehensive approach, allowing for relatively easy and

rapid sample preparation and data collection. Through the utilization of ^{1}H NMR spectroscopy,

specific product fingerprints can be determined, revealing detailed insights into the botanical origin

of honey by identifying distinct organic compounds (Bertram et al., 2005). Consequently, NMR

has become pivotal in creating efficient and suitable models for honey adulteration detection (Lolli

et al., 2008).

Though Turkish pine honey has been examined in various studies, a detailed analysis focusing on

its specific biochemical metabolite composition and genuine characteristics is still lacking. In this

study, the metabolite profiles of pine honey samples collected from Turkey were qualitatively and

quantitatively characterized by NMR spectrometry. Additionally, by comparing these results with

honeys from other regions, including Oceania, the Aegean, and the West Coast of North America,

we aim to uncover distinctive properties of Turkish pine honey that have not been previously

identified.

2. Material and Method

2.1 Sample collection

A total of thirty-three honey samples, sourced from two distinct botanical origins, including flower

(n=24) and pine (n=9), were collected directly from Fijian (flower honey n=3), Hawaiian (flower

honey n=6), Turkish (flower honey n=7) (Turkish pine honey n=9) and American (flower honey

n=8) producers (Fig 1). Additionally, Manuka honey with an MGO level of 83 and maple syrup

were bought from a local market in California. All obtained samples were subsequently stored in

dark glass containers at a temperature of 4 °C prior to analysis.

2.2 Extraction and preparation of Honey samples

Honey samples were weighed in 50mL tubes and diluted 1:4 by weight using a phosphate buffer

(150mM KH_2PO_4, pH=4.5, 100% D_2O). The samples were homogenized by pipetting and

centrifuged at 1500 x g for 15 minutes at room temperature to remove insoluble particles. 600µL

NMR samples were prepared by adding DSS solution (1mM final concentration) and placed in

5mm NMR tubes.

2.3 Qualitative and quantitative data collection by [1]H NMR Spectroscopy

NMR data acquisition process was completed using 500 MHz Bruker Ascend magnet equipped

with Avance NEO console and BBO double resonance room temperature probe. Water suppression

experiments are commonly used for suppressing the water peak rather than routine 1D proton pulse

in NMR-based metabolomics studies (Dağ et al., 2022). In this study, we used 1D NOESY-presat

(noesygppr1d) pulse program for all NMR data acquisitions, and each spectrum consisted of 8K

scans of 32K complex data points with a spectral width of 9615.4 Hz. Chenomx NMR Suite 9.02

software (Chenomx inc., Canada) software was used for calculation of the target metabolites (Dağ

et al., 2022). Target metabolite concentrations and peak shape of other molecules calculated by

Chenomx based on the concentration and peak height of the DSS. NMR data were collected with

8K scans to maximize the signal to noise ratio.

2.4 Statistical analysis

Partial least squares discriminant analysis (PLS-DA), a supervised method, was conducted for a

comprehensive analysis of NMR-based metabolomics data. Dataset was subjected to cube root

transformation and Pareto scaling prior to analysis.

3. Results and Discussion

3.1 Analysis of Metabolite Composition in Turkish Pine Honey

Among identified water-soluble compounds in Pine Honey, 44 of them are listed with their average

concentration with standard deviation in Table 1. In the analysis of Turkish Pine honey, a diverse

range of chemical compounds were identified, and classified according to their chemical

classifications. Carbohydrates included Fructose, Glucose, Maltose, Sucrose, Arabinose,

Mannose, and Fucose. The Amino Acids and related compounds included Isoleucine, Leucine,

Proline, Valine, Threonine, Lysine, Alanine, Phenylalanine, Aspartate, Glutamine, Glutamate,

Methylguanidine, and Kynurenate. Keto Acids and Ketones included Pyruvate, Acetoacetate,

Acetone acetate, 2-hydroxybutyrate, lactate, Acetoacetate and Acetone. Ethanol was the only

identified Alcohol, while Tartrate, Fumarate, Succinate, Malate, Citrate, 3-phenyllactate,

Phenylacetate, and Formate were categorized as Organic Acids. Other classifications included,

Nucleotide derivatives (Uridine), Amides (Acetamide), Aromatic Compounds (Benzoate), Other

Organic Compounds (Isocaproate, 2-hydroxyisobutyrate), Quaternary Ammonium Cations

(Choline), and Other Nitrogen Compounds (Trigonelline, Pi-Methylhistidine).

Expectedly, majority of compounds detected in Turkish Pine honey are readily found in numerous

honeys (White, 1978). Concentrations of these may differ between each honey type however their

overall presence is discussed later in comparison to other honey samples. Otherwise, presence of

compounds that are not commonly found in honey acetoacetate, and Pi-Methylhistidine, are worth

discussion regarding Turkish Pine Honey. Acetoacetate, also known as acetoacetic acid or

oxobutyrate, is a short-chain keto acid that can be found in various food items, particularly various

fruits and as an artificial flavouring. Soluble in water and weakly acidic in nature, it is a ketone

body that also serves as an energy substrate during specific metabolic conditions such as fasting

or intense exercise (Johnston et al., 1954). On the other hand, Pi-Methylhistidine is a post-

translationally modified amino acid derived from the contractile proteins: actin and myosin. It is

often observed in conditions associated with nitrogen loss and is excreted through urine (Elia et

al., 1981). It is plausible that the appearance of these compounds in Turkish Pine honey could be

a direct consequence of local organisms' involvement in its production. Nevertheless, they hold

potential as distinctive biomarkers for the authentication and adulteration detection of Turkish Pine

honey.

3.2 Development of a universal test of authenticity for Turkish Pine honey

We constructed Pairwise score plots of 5 chosen PLS-DA components to distinguish between

Turkish Pine Honey and the remaining samples (Figure 2a). Following that, we selected the plot

that demonstrated the two principal components responsible for the greatest degree of separation

between the two groups, as shown in Figure 2b. Incorporation of a third component clearly

revealed clustering of Turkish Pine honey from the rest of samples (Figure 2c). As depicted in

Figure 2d, the VIP scores were classified into three distinct ranges, signifying the most important

metabolites contributing to the separation of metabolic profiles of Turkish Pine honey from the

rest of the floral honey samples. The ranges are as follows: 1.8-2.0, encompassing Acetone,

Acetamide, and Lysine; 1.5-1.8, including Citrate, Acetate, Aspartate, and Isocaproate; and 1.0-

1.5, comprising Tyrosine, Proline, Malate, 2-Hydroxyisobutyrate, Phenylacetate, Methyl

guanidine, Succinate, and Pyruvate. All these compounds were in higher prevalence in Pine honey,

except for Tyrosine and Phenylacetate, which were found to be lower. Furthermore, the detailed

analysis of the compounds with the most significant changes is represented by the box plots of

acetamide, acetate, and acetone in Figure 2e. Among these, acetate and acetone are well-known

volatile compounds found in several types of honey. Acetamide, on the other hand, deviates from

this commonality, as it is not typically detected in honey.

NMR spectrum of honey samples from different origins and maple syrup shown at figure 3a-c.

The conspicuous presence of acetamide in Turkish Pine honey compared to other samples could

signify a distinct chemical characteristic or metabolic pathway unique to this honey type. In

addition to assigned compounds, NMR spectra were compared following peak identification

(Figure 3d) to pick out peaks that are unique to pine honey samples. Peaks at 1.85, 1.87, and 1.89

ppm were observed to be unique to pine honey. These peaks can be assigned to their respective

compounds and potentially be used as biomarkers to verify the authenticity of pine honey or

distinguish it from other honey types.

In addition to this, Figure 3 presents the general proton NMR spectra of honeys from different

origins as well as maple syrup. While maple syrup can be readily distinguished at first glance due

to its unique sugar content, differentiating between honey samples is not as straightforward.

Notably, the peaks appearing in the 8-10 ppm range for Manuka honey are particularly striking.

We extended the analysis by constructing Pairwise score plots for the selected PLS-DA

components, specifically to discern the differences between Turkish Pine Honey, Turkish Floral

Honey, and Floral Honey from outside Turkey. This analysis builds upon our previous findings to

further investigate the unique characteristics of Turkish Pine Honey. We selected the plot that

exhibited the two principal components accounting for the most pronounced separation between

these three groups, as shown in Figure 4a. This plot illustrates Turkish Flower Honey as an

intermediary, bridging, characteristics of Turkish Pine Honey with Flower Honeys from outside of

Turkey. As anticipated, addition of a third dimension further accentuated the clustering of Turkish

Pine honey, clearly distinguishing all three groups particularly Turkish Flower Honey from the rest

of the groups (Figure 4b). VIP scores specify the identified compounds responsible for separation

mirroring the classification found in the previous analysis, however now in the context of an

extended comparison (Figure 4c). In our detailed examination of the VIP scores, we identified

three distinct phenomena that allowed us to discern the differential prevalence of compounds and

their role in the separation. Compounds with high VIP scores over 2, such as acetone and

acetamide, were found in higher prevalence in Pine Honey and, to a lesser extent, in Turkish

Flower Honey and least in Non-Turkish Flower honeys. Compounds with VIP scores less than 2

such as isocaproate, malate, sucrose, and acetate also exhibited a similar pattern. Overall, these

compounds can be considered as intermediatory compounds which are responsible for the bridging

separation of Turkish Pine Honey from Turkish Flower Honey and Non-Turkish Flower Honeys

as shown in Figure 4a. Compounds, formate, lysine, glucose, trigonelline Phenylalanine, 3-

phenyllactat, aspartate and citrate was found least in Turkish Flower honey on contrary to Non-

Turkish Flower honey and Turkish Pine Honey. Finally, phenylacetate was also an intermediatory

compound between Turkish Honey and Non-Turkish Honeys with notable exception that it was

found in greater extend in Non-Turkish Flower Honeys and less in Turkish Flower Honeys and

least in Turkish Pine Honey.

In a further effort to refine our understanding of the distinctions of Turkish Pine honey, we

performed a comparative analysis between Turkish Pine Honey and Oceania Flower Honey, as

well as between Turkish Pine Honey, Hawaiian Floral Honey, and Fijian Floral Honey. First, the

scores plot of PLS-DA was generated to compare Turkish Pine Honey and Oceania Floral Honey,

as illustrated in Figure 4d. There was clear separation between the two groups. The VIP scores

obtained from this PLS-DA model are displayed in Figure 4e. Compounds found in higher

prevalence in Turkish Pine Honey and associated with VIP scores higher than 1.5 include

acetamide, acetone, and isocaproate, while those with VIP scores lower than 1.5 are lysine, acetate,

and mannose. Conversely, compounds that are found to a lesser extent in Turkish Pine Honey and

exhibit VIP scores higher than 1.5 are tyrosine, glucose, and phenylacetate; those with VIP scores

lower than 1.5 include lactate, phenylalanine, valine, and isoleucine. Next, a score plot of PLS-DA

was constructed to compare Turkish Pine Honey, Hawaiian Floral Honey, and Fijian Floral Honey,

as depicted in Figure 4f. Though a complete separation between all three groups was observed,

Hawaiian Floral Honey appeared to be more closely related to Turkish Pine Honey. The VIP scores,

derived from the PLS-DA model, are presented in Figure 4g. Supporting our previous findings,

compounds such as tyrosine, phenylacetate, glucose, and lactate with VIP scores higher than 1.8

were found to the greatest extent in Fijian Floral Honey, less in Hawaiian Floral Honey, and least

in Turkish Pine Honey. Furthermore, all compounds except isocaproate and acetamide, which also

have the lowest two VIP scores, were found to the greatest extent in Fijian Floral Honey. On the

other hand, compounds such as 3-phenyllactate, valine, phenylalanine, isoleucine, and ethanol

were more prevalent in Hawaiian Floral Honey compared to Turkish Pine Honey, whereas fructose,

threonine, alanine, and pi-methylhistidine were less prevalent.

3.3 Statistical Discrimination between Turkish Pine Honey and Turkish Floral Honey

In order to display the intrinsic metabolite differences in Turkish honey we generated pairwise

score plots of the selected components as shown in Figure 5a. As formerly shown in our previous

analysis there is a noticeable separation between Turkish Flower Honey and Turkish Pine Honey.

VIP scores obtained from the PLS-DA analysis is depicted in Figure 5b showing most of the

compounds responsible for separation except tyrosine was found in greater extent in Turkish Pine

Honey. Among these compounds, Fructose and proline were the compounds with the highest VIP

scores higher than 2.0. The remaining compounds with smaller VIP scores are with respect to their

VIP scores are Citrate, Arabinose, Lysine, Aspartate, Glutamine, Malate, methylguanidin,

acetamide, formate, Sucrose, Acetone, and Leucine.

The distinctive compositional variations between Turkish Pine Honey and Turkish Flower Honey, revealed through our analysis, point toward the underlying biological and ecological factors that might be at play. Different floral sources and the unique foraging behaviors of bees may contribute to the variations in compounds such as Fructose, Proline, Citrate, and others. The presence or absence of certain compounds in specific honey types could reflect the particular plants' nectar and pollen content, soil chemistry, or the microclimate of the regions where the bees collect their materials. These metabolic variations between Turkish Pine honey and Turkish Flower Honey offer an intricate view of the relationships between honey composition and environmental factors and may prompt further targeted exploration into the ecology and biochemistry of honey production. Such understanding may also have practical implications in honey authentication, quality control of commercial Honey sold as Turkish Pine Honey, and potential health-related applications of Turkish Honey.

3.4 Chemical composition analysis of Turkish pine honey from Mugla Province

To delve into the regional variations within Turkish Pine Honey, we conducted a comparison between samples collected from the western (Datça) and eastern (Fethiye) parts of Muğla Province. The scores plot of PLS-DA, as presented in Figure 5c, illustrates a distinguishable separation between these two regions, suggesting location-specific compositional differences. The VIP scores obtained from the PLS-DA model, depicted in Figure 5d specifying the key differences. Among compounds with VIP scores greater than 2, Ethanol, which was not previously found to contribute to distinction of Turkish Pine honey, was found more prevelantly in Pine honey from Fethiye whereas Sucrose, Fructose and Isocaproate was found more in Pine honey from Datca. For

compounds with lower VIP scores than 2, Phenylalanine, Arabinose glucose, Alanine was found

more prevalently in Pine Honey from Datca and Citrate, Choline, formate, Acetone, acetate,

Leucine, and Aspartate was found more in Pine honey from Fethiye. Additonally we explored the

quantitative variations of specific metabolites though box plots namely acetate, ethanol, fructose,

and sucrose. (Figure 5e, 5f, 5g, 5h).

The distinct separation between Turkish Pine Honey from Datça and Fethiye, as evidenced by our

PLS-DA analysis, raises questions about the underlying factors influencing these compositional

differences. While specific reasons for these variations cannot be definitively pinpointed without

further investigation, several potential explanations warrant consideration considering these two

regions are not too far apart from each other. The presence of unique floral sources in each region

could contribute to differences in sugar content, such as sucrose and fructose. Ethanol, found more

prevalently in Pine honey from Fethiye, might suggest variations in fermentation processes or

other biogenic activities in the honey itself. Environmental factors, including climate and soil

composition, might impact the availability and type of nectar collected by the bees, thus affecting

the chemical profile of the honey considering Datca's geography as a peninsula deeper into the

Aegean Sea. Additionally, variations in bee foraging behaviors and hive management practices

between the two locations may play a role. By deepening our understanding of regional variations

in Turkish Pine Honey, we may open avenues for more targeted studies that can elucidate the

factors contributing to these compositional differences. Such insights could have broader

implications for the honey industry, from enhancing authenticity and traceability to optimizing

production methods tailored to specific regional characteristics.

Karabagias et al. (2018) have identified and discriminated Greek Pine honey of different geographic origins using metabolite-based nuclear magnetic resonance spectroscopy ([1]H NMR) and high-performance liquid chromatography (HPLC). The obtained physicochemical data were modeled using multivariate analyses and supervised statistical methodologies.

To characterize the origin of honey, comparative volatile analysis of Greek and Turkish pine honey has been done by a purge & trap–gas chromatograph–mass (GS-MS) spectrometer system and Kohonen self-organizing maps (KSOM) algorithm (Tananaki, Thrasyvoulou, Giraudel, and Montury, 2007). Zhang et al. (2020) have made a classification based on 1H NMR profile to separate 8 monofloral Chinese honeys, namely acacia, jujube, linden, longan, orange, rape, sunflower, vitex, into flower types and geographical origins. In the experiment, while PCA-X unsupervised modeling was carried out to determine its botanical origin, OPLS-DA supervised modeling was used to determine its geographical origin (Zhang et al., 2020). Alternating current (AC) impedance is a resistance of alternative current that can be used for evaluation of food quality and adulteration. This method is based on detecting change in impedance of a sample during stimulation which is called electrochemical impedance analysis. Hao et al. (2022) identified acacia honey adulteration by measuring the differences in AC impedance spectroscopy and metabolite analysis with 1H NMR spectroscopy. The combination of AC impedance and 1H NMR spectroscopy is a beneficial quality-control tool for identification of authenticity in honey (Hao et al., 2022).

Conclusion

In this study, 33 honey samples were examined using 1D H-NMR spectroscopy for water soluble metabolites and subjected to PLS-DA statistical analysis to reveal differences in metabolite profiles of Turkish Pine Honey, Turkish Flower Honey, and Non-Turkish Flower Honey. Statistical analyses revealed that Pi-Methylhistidine and acetoacetate are scarcely found in most honey types and can serve as potential biomarkers for Turkish Pine Honey, as well as 1D H-NMR peaks found at 1.84, 1.87, and 1.89 ppm following assignment to their respective compounds. Acetone, Acetamide, Lysine, Citrate, Acetate, Aspartate, Isocaproate, Proline, Malate, 2-Hydroxyisobutyrate, Methyl guanidine, Succinate, and Pyruvate are metabolites that were found in higher concentrations in Turkish Pine Honey samples, while Tyrosine and Phenylacetate concentrations were lower compared to floral honey types. Comparative analysis of Turkish Pine Honey, Turkish Flower Honey, and Non-Turkish Flower Honey revealed that acetone, acetamide, isocaproate, malate, sucrose, and acetate were detected in decreasing amounts for the honey types, respectively, also indicating the potential contribution of geographical factors alongside differences due to botanical origin.

PLS-DA analysis was also sufficient to observe separation between Turkish Pine Honey and Non-Turkish honey. Additionally, fructose and proline were components with the highest VIP scores when Turkish pine and floral honey samples were compared, with both being found at higher concentrations in pine honey, alongside 10 other compounds with VIP scores larger than 1, accounting for separation between the groups. Comparison between Turkish pine honey samples sourced from different locations also revealed regional differences in composition, which might be attributed to factors such as local flora, climate, and foraging behaviors of bees, although additional studies need to be conducted to address local differences.

Author Contributions

Çağdaş Dağ: Conceptualization, Methodology, Supervision, Funding acquisition, Writing-Reviewing and Editing, Writing-Original draft preparation **Kerem Kahraman:** Investigation, Formal analysis, Visualization, Writing-Reviewing and Editing **Oktay Göcenler:** Investigation, Formal analysis, Writing-Original Draft, Visualization

ACKNOWLEDGMENT

The authors acknowledge the use of the services and facilities of n^{2}STAR-Koç University Nanofabrication and Nanocharacterization Center for Scientific and Technological Advanced Research. The authors gratefully acknowledge use of the services and facilities of the Koç University Is Bank Infectious Disease Center (KUIS-CID).

REFERENCES

Boffo, E.F., Tavares, L.A, Tobias, A.C.T., Ferreira, M.M.C., Ferreira, A.G. (2012). Identification of components of Brazilian honey by 1H NMR and classification of its botanical origin by chemometric methods. *LWT - Food Science and Technology, 49*(1), 55-63. https://doi.org/10.1016/j.lwt.2012.04.024

Cianciosi, D., Forbes-Hernández, T., Afrin, S., Gasparrini, M., Reboredo-Rodriguez, P., Manna, P., Zhang, J., Bravo Lamas, L., Martínez Flórez, S., Agudo Toyos, P., Quiles, J., Giampieri, F., &

Battino, M. (2018). Phenolic compounds in honey and their associated health benefits: A Review. *Molecules*, *23*(9), 2322. https://doi.org/10.3390/molecules23092322

Consonni, R., Bernareggi, F., Cagliani, L.R. (2019). NMR-based metabolomic approach to differentiate organic and conventional Italian honey, *Food Control*, *98*, 133-140. https://doi.org/10.1016/j.foodcont.2018.11.007

Dağ, Ç., Göçenler, O., Tozkoparan, C.D. (2022). The analysis of metabolic content of traditional milk collected from three regions in turkey by NMR spectroscopy. *GIDA, 47*(5), 765-775. https://doi.org/10.15237/gida.GD22042

Elia, M., Carter, A., Bacon, S., Winearls, C. G., & Smith, R. (1981). Clinical usefulness of urinary 3-methylhistidine excretion in indicating muscle protein breakdown. *BMJ, 282*(6261), 351–354. https://doi.org/10.1136/bmj.282.6261.351

Eraslan, G., Kanbur, M., Silici, S., & Karabacak, M. (2010). Beneficial effect of pine honey on trichlorfon induced some biochemical alterations in mice. *Ecotoxicology and environmental safety*, *73*(5), 1084–1091. https://doi.org/10.1016/j.ecoenv.2010.02.017

Erdal, B., Tipi, T. (2022). Time Series Forecasting of Honey Production in Turkey, *European Journal of Science and Technology, 35*, 417-423. https://doi.org/10.31590/ejosat.1066665

Hao, S., Yuan, J., Cui, J., Yuan, W., Zhang, H. & Xuan, H. (2022). The rapid detection of acacia honey adulteration by alternating current impedance spectroscopy combined with 1H NMR profile. *Lebensmittel-Wissenschaft + Technologie, 161*, 113377. https://doi.org/10.1016/j.lwt.2022.113377

Johnston, J. A., Racusen, D. W., & Bonner, J. (1954). The metabolism of isoprenoid precursors in a plant system. *Proceedings of the National Academy of Sciences, 40*(11), 1031–1037. https://doi.org/10.1073/pnas.40.11.1031

Kafantaris, I., Tsadila, C., Nikolaidis, M., Tsavea, E., Dimitriou, T. G., Iliopoulos, I., Amoutzias,

G. D., & Mossialos, D. (2021). Transcriptomic Analysis of Pseudomonas aeruginosa Response to

Pine Honey via RNA Sequencing Indicates Multiple Mechanisms of Antibacterial Activity. *Foods*,

10(5), 936. https://doi.org/10.3390/foods10050936

Karabagias, I. K., Vlasiou, M., Kontakos, S., Drouza, C., Kontominas, M.G., Keramidas, A.D.

(2018). Geographical discrimination of pine and fir honeys using multivariate analyses of major

and minor honey components identified by 1H NMR and HPLC along with physicochemical data.

European Food Research and Technology, 244, 1249-1259. https://doi.org/10.1007/s00217-018-

3040-5

Kazalaki, A., Misiak, M., Spyros, A., Dais, P. (2015). Identification and quantitative determination

of carbohydrate molecules in Greek honey by employing 13 C NMR spectroscopy. *Analytical*

Methods, 7(14), 5962–5972. https://doi.org/10.1039/C5AY01243K

Koç, A.U., Karacaoğlu, M., Nadem, H.Ş., & Doğan, M. (2017). Determination of Shelf Life of

Chaste Tree (Vitex agnus castus) Honey and Pine Honey. *GIDA The Journal of Food, 42*(5), 577-

587. 10.15237/gida.GD17020

Luong, D.V., Tam, N.Q., Xuan, D.T.T. and Tai, N.T. (2019), NMR based metabolomic approach

for evaluation of Vietnamese honey. *VJCH, 57*, 712-716.

https://doi.org/10.1002/vjch.2019000101

Schievano, E., Stocchero, M., Morelato, E., Facchin, C., Mammi, S. (2012). An NMR-based

metabolomic approach to identify the botanical origin of honey. *Metabolomics, 8*(4), 679–

690. https://doi.org/10.1007/s11306-011-0362-8

Olawode, E. O., Tandlich, R., & Cambray, G. (2018). ^{1}H-NMR Profiling and Chemometric

Analysis of Selected Honeys from South Africa, Zambia, and Slovakia. *Molecules (Basel,*

Switzerland), *23*(3), 578. https://doi.org/10.3390/molecules23030578

Özkök, A., D'arcy, B., & Sorkun, K. (2010). Total phenolic acid and total flavonoid content of

Turkish Pine Honeydew Honey. *Journal of ApiProduct & ApiMedical Science*, *2*(2), 65–71.

10.3896/IBRA.4.02.2.01

Özkök, A., Yüksel, D., Sorkun, K. (2018). Chemometric Evaluation of the Geographical Origin of

Turkish Pine Honey, *Food and Health, 4*(4), 274-282. https://doi.org/10.3153/FH18027

Parin, F.N., Terzioğlu, P., Sıcak, Y., Yıldırım, K. & Öztürk, M. (2021). Pine honey–loaded

electrospun poly (vinyl alcohol)/gelatin nanofibers with antioxidant properties. *The Journal of The*

Textile Institute, 112(4), 628-635. https://doi.org/10.1080/00405000.2020.1773199

Rossano, R., Larocca, M., Polito, T., Perna, A. M., Padula, M. C., Martelli, G., & Riccio, P. (2012).

What are the proteolytic enzymes of honey and what they do tell us? A fingerprint analysis by 2-

D zymography of Unifloral Honeys. *PLoS ONE, 7*(11).

https://doi.org/10.1371/journal.pone.0049164

Sen, G., Güngör, E. (2018). Selecting Suitable Forest Areas For Honey Production Honey

Production Using The AHP: A Case Study in Turkey, *Cerne*, *24*(1).

Sıcak, Y., Şahin-Yağlıoğlu, A. & Öztürk, M. (2021). Bioactivities and phenolic constituents

relationship of Muğla thyme and pine honey of Turkey with the chemometric approach. *Food*

Measure, 15, 3694–3707. https://doi.org/10.1007/s11694-021-00940-8

Walker, M. J., Cowen, S., Gray, K., Hancock, P., & Burns, D. T. (2022). Honey authenticity: the

opacity of analytical reports - part 1 defining the problem. *NPJ science of food*, *6*(1), 11.

https://doi.org/10.1038/s41538-022-00126-6

Tananaki, Ch., Thrasyvoulou, A., Giraudel, J.L., Montury, M. (2007). Determination of volatile

characteristics of Greek and Turkish pine honey samples and their classification by using Kohonen

self organizing maps. *Food Chemistry,* 101(4), 1687–1693.

https://doi.org/10.1016/j.foodchem.2006.04.042

Thrasyvoulou, A. (1986). The use of HMF and diastase as criteria of quality of greek honey.

Journal of Apicultural Research, 25(3), 186–195.

https://doi.org/10.1080/00218839.1986.11100715

Tsagkaris, A. S., Koulis, G. A., Danezis, G. P., Martakos, I., Dasenaki, M., Georgiou, C. A., &

Thomaidis, N. S. (2021). Honey authenticity: analytical techniques, state of the art and challenges.

RSC advances, *11*(19), 11273–11294. https://doi.org/10.1039/D1RA00069A

Tsavea, E., Vardaka, F.-P., Savvidaki, E., Kellil, A., Kanelis, D., Bucekova, M., Grigorakis, S.,

Godocikova, J, Gotsiou, P., Dimou, M., Loupassaki, S., Remoundou, I., Tsadila, C., Dimitriou,

T.G., Majtan, J., Tananaki, C., Alissandrakis, E., Mossialos, D. (2022). Physicochemical

Characterization and Biological Properties of Pine Honey Produced across Greece. *Foods*, *11*, 943.

10.3390/foods11070943

Tsuruda, J. M., Chakrabarti, P., Sagili, R.R. (2021). Honey Bee Nutrition. *Veterinary Clinics of*

North America: Food Animal Practice, *37*(3), 505-519. 10.1016/j.cvfa.2021.06.006

Valverde, S., Ares, A. M., Stephen Elmore, J., & Bernal, J. (2022). Recent trends in the analysis of

Honey Constituents. *Food Chemistry*, *387*, 132920.

https://doi.org/10.1016/j.foodchem.2022.132920

White, J. W. (1978). Honey. *Advances in Food Research, 24*, 287–374.

https://doi.org/10.1016/S0065-2628(08)60160-3

Zhang, J., Chen, H., Fan, C., Gao, S., Zhang, Z., Bo, L. (2020). Classification of the botanical and

geographical origins of Chinese honey based on 1H NMR profile with chemometrics. *Food*

Research International, 137, 109714. https://doi.org/10.1016/j.foodres.2020.109714

Živkov Baloš, M., Popov, N., Jakšić, S., Mihaljev, Ž., Pelić, M., Ratajac, R., & Ljubojević Pelić,

D. (2023). Sunflower honey—evaluation of quality and stability during storage. *Foods, 12*(13),

2585. 10.3390/foods12132585

Figure 1. Honey Sampling Map of the study. The number of pine honey is labeled with orange

and the number of flower honey is labeled with green.

Figure 2. Multivariate Statistical Approach for the Discrimination of Turkish Pine Honey

and Floral honey a. PLS-DA pairwise score plot for top 5 components b. Scores plot of PLS-DA

(PC1 vs. PC2) carried out on 1H NMR data of Turkish pine honeys (green) and floral honeys (red).

c. 3D Scores plot of PLS-DA (PC1 vs. PC2 vs. PC3) carried out on 1H NMR data of Turkish pine

honeys (green) and floral honeys (red). d. The variable importance in projection (VIP) scores

obtained from the PLS-DA model. The colored boxes on the right demonstrate the relative

concentrations of the corresponding metabolite. e. quantitative variation of acetamide f.

quantitative variation of acetate g. quantitative variation of acetone

Figure 3. NMR spectrum of honey samples from different origins and maple syrup a.
Complete [1]H NMR spectrum of pine honey (green), flower honey (brown), Manuka honey (Blue)
and Maple syrup (red) b. 0-3.5 ppm region of [1]H NMR spectrum of pine honey (green), flower
honey (brown), Manuka honey (Blue) and Maple syrup (red) c. 5.5-10.5 ppm region of
[1]H NMR spectrum of pine honey (green), flower honey (brown), Manuka honey (Blue) and Maple
syrup (red) d. Pine honey samples unique peaks at 1.89, 1.87 and 1.85 ppm.

Figure 4. Multivariate Statistical Approach for the Discrimination of Turkish Pine Honey, Turkish Floral honey, and other Floral honey samples and specifically Oceania Floral honey samples a. Scores plot of PLS-DA (PC1 vs. PC2) carried out on ^{1}H NMR data of Turkish pine honey (blue), Turkish floral honey (green) and floral honey from out of Turkey (red). b. 3D Scores plot of PLS-DA (PC1 vs. PC2 vs. PC3) carried out on 1H NMR data of Turkish pine honey (blue), Turkish floral honey (green) and floral honey from out of Turkey (red). c. The variable importance in projection (VIP) scores obtained from the PLS-DA model. The colored boxes on the right

demonstrate the relative concentrations of the corresponding metabolite. d. Scores plot of PLS-

DA (PC1 vs. PC2) carried out on 1H NMR data of Turkish pine honey (green) and Oceania floral

honey (red). e. The variable importance in projection (VIP) scores obtained from the PLS-DA

model. The colored boxes on the right demonstrate the relative concentrations of the corresponding

metabolite. f. Scores plot of PLS-DA (PC1 vs. PC2) carried out on 1H NMR data of Turkish pine

honey (blue), Hawaiian floral honey (green) and Fijian floral honey (red). g. The variable

importance in projection (VIP) scores obtained from the PLS-DA model. The colored boxes on the

right demonstrate the relative concentrations of the corresponding metabolite

Figure 5. Multivariate Statistical Approach for the Discrimination of Turkish Pine Honey,

and Turkish Floral honey samples and Turkish Pine Honey samples from different regions

a. Scores plot of PLS-DA (PC1 vs. PC2) carried out on 1H NMR data of Turkish floral honey (red)

and Turkish pine honey (green). b. The variable importance in projection (VIP) scores obtained

from the PLS-DA model. The colored boxes on the right demonstrate the relative concentrations

of the corresponding metabolite. c. Scores plot of PLS-DA (PC1 vs. PC2) carried out on 1H NMR

data of Turkish pine honey from western Muğla (Datça) (red) and Turkish pine honey from eastern

Muğla (Fethiye) (green). d. The variable importance in projection (VIP) scores obtained from the

PLS-DA model. The colored boxes on the right demonstrate the relative concentrations of the

corresponding metabolite. e. quantitative variation of acetate f. quantitative variation of ethanol g.

quantitative variation of fructose h. quantitative variation of sucrose

Table 1. Identified metabolite concentrations in Turkish Pine Honey

Metabolite	Average (mM)	STD (mM)	Metabolite	Average (mM)	STD (mM)
Fructose	3875.375	618.867	Pyruvate	0.762	0.435
Glucose	2836.334	494.415	Isoleucine	0.425	0.128
Maltose	115.083	65.138	tartrate	0.946	0.525
Sucrose	34.762	28.534	Aspartate	4.078	2.478
Arabinose	44.803	25.899	Acetoacetate	0.355	0.262
Mannose	7.671	5.438	tyrosine	0.216	0.124
Ethanol	1.789	1.328	Leucine	1.627	3.890
Glutamine	2.537	1.230	3-phenyllactate	0.337	0.220
acetamide	1.346	0.973	2-hydroxybutyrate	0.416	0.305
lactate	1.122	0.453	Benzoate	0.230	0.244
glutamate	4.938	3.751	Uridine	0.247	0.133
proline	14.049	6.508	Acetone	0.324	0.203
acetate	1.462	0.347	Valine	0.338	0.171
threonine	0.889	0.205	Fumarate	0.149	0.141
Lysine	3.839	3.193	2-hydroxyisobutyrate	0.168	0.068
Alanine	0.816	0.300	Isocaproate	1.046	1.274
Phenylalanine	1.018	1.826	Kynurenate	0.141	0.115
Fucose	0.875	0.271	succinate	0.409	0.615
Malate	5.646	4.722	methylguanidine	0.702	1.403
Choline	1.145	0.354	formate	0.638	0.826
phenylacetate	0.329	0.249	trigonelline	0.158	0.051
Citrate	2.205	1.858	Pi-Methylhistidine	0.302	0.203

516