# Macroevolutionary inference of complex modes of chromosomal speciation in a cosmopolitan plant lineage

Carrie M. Tribble[1,*,†], José Ignacio Márquez-Corro[2,3,*], Michael R. May[4], Andrew L. Hipp[5], Marcial Escudero[6**], and Rosana Zenil-Ferguson[7**]

[1]*School of Life Sciences, University of Hawai'i at Mānoa, Honolulu, HI, 96822, USA*
[2]*Royal Botanic Gardens, Kew. Richmond, Surrey TW9 3AE, United Kingdom*
[3]*Department of Molecular Biology and Biochemistry Engineering, Universidad Pablo de Olavide. Sevilla 41013, Spain*
[4]*Department of Evolution and Ecology, University of California Davis, Davis, CA USA*
[5]*Herbarium and Center for Tree Science, The Morton Arboretum, Lisle, IL 60532 USA*
[6]*Department of Plant Biology and Ecology, University of Sevilla, Sevilla, Spain*
[7]*Department of Biology, University of Kentucky, Lexington KY 40506 USA*
*Co-first authors
**Co-senior authors
†Corresponding author: ctribble09@gmail.com

**Summary**

- The effects of single chromosome number change—dysploidy—mediating diversification remain poorly understood. Dysploidy modifies recombination rates, linkage, or reproductive isolation, especially for one-fifth of all eukaryote lineages with holocentric chromosomes. Dysploidy effects on diversification have not been estimated because modeling chromosome numbers linked to diversification with heterogeneity along phylogenies is quantitatively challenging.

- We propose a new state-dependent diversification model of chromosome evolution that links diversification rates to dysploidy rates considering heterogeneity and differentiates between anagenetic and cladogenetic changes. We apply this model to *Carex* (Cyperaceae), a cosmopolitan flowering plant clade with holocentric chromosomes.

- We recover two distinct modes of chromosomal speciation in *Carex*. In one diversification mode, dysploidy occurs frequently and drives faster diversification rates. In the other mode, dysploidy is rare and diversification is driven by heterogeneity. When we use a model that excludes process heterogeneity, we mistakenly infer a strong, uniform positive effect of dysploidy on diversification, showing that standard models may lead to confident but incorrect conclusions about diversification.

- This study demonstrates that dysploidy can have a significant role in large plant clade speciation despite the presence of other unmeasured factors that simultaneously affect diversification.

[**keywords**—*Carex*; Chromosome number; diversification; dysploidy; macroevolution process variation; macroevolution; RevBayes; TensorPhylo]

**Total word count** (excluding summary, references, and legends): 4888

Summary: 189

Introduction: 1067

Methods: 1368

Results: 619

Discussion: 1398

Acknowledgements: 107

No. of Figures: 3 (3 in color)

No. of Tables: 1

No. of Supporting Information files: 1

**Author information**

Carrie M. Tribble (0000-0001-7263-7885)

José Ignacio Márquez-Corro (0000-0003-4277-2933)

Michael R. May (0000-0002-5031-4820)

Andrew L. Hipp (0000-0002-1241-9904)

Marcial Escudero (0000-0002-2541-5427)

Rosana Zenil-Ferguson (0000-0002-9083-2972)

# Introduction

Unveiling the primary drivers of diversification remains one of the most important goals in evolutionary biology

(Sauquet & Magallón 2018). Hundreds of studies have focused on estimating changes in plant diversification processes

through time (Magallón & Castillo 2009), across clades (Magallón et al. 2019), or in association with trait evolution

(Helmstetter et al. 2023). Chromosome number changes and rearrangements are particularly likely to influence lineage

diversification (Freyman & Höhna 2018). Yet, most plant diversification studies that address chromosome evolution

focus only on the role of polyploidy, rather than on dysploidy—gains and losses of single chromosomes—which does

not involve large DNA content changes (*i.e.*, dysploidy, gains or losses of single chromosomes; Escudero et al. 2014,

Mandáková & Lysak 2018). A recent review of trait-dependent diversification in angiosperms, for example, cites seven

studies linking speciation and polyploidy (Helmstetter et al. 2023), only one which considered dysploidy linked to di-

versification (Freyman & Höhna 2018). While dysploidy has the potential to influence lineage diversification through

effects on recombination or reproductive isolation, the macroevolutionary effects of dysploidy on speciation and extinc-

tion remain unknown, and far fewer macroevolutionary studies have focused on speciation and extinction as a result

of dysploidy.

## Dysploidy and holocentric chromosomes

There are two classic models of chromosomal speciation in a macroevolutionary context. In the hybrid-dysfunction

model, dysploidy is linked to speciation events. Under this model, dysploidy causes an immediate reproductive barrier,

as reproduction between individuals with different chromosome numbers would cause problems during meiosis. Thus,

most—if not all—dysploidy events across a phylogeny would occur cladogenetically. Alternatively, the recombination-

suppression model posits that chromosomal rearrangements may become fixed in lineages via either drift or selection

(as some rearrangements may physically link adaptive loci or locally reduce recombination). Under this model, dys-

ploidy would evolve primarily anagenetically since fission and fusions do not necessarily restrict gene flow; rather,

they evolve within populations and lineages (Baker & Bickham 1986). The two above-described models are not mutu-

ally exclusive. In a recent review, Lucek et al. (2022) introduce a third option—the hybrid-dysfunction/recombination-

suppression model, under which dysploidy evolves both anagenetically and cladogenetically (see Fig. 4 in Lucek

et al. 2022). While populations may be able to continue interbreeding despite some dysploidy events (which then may

become fixed in the lineage), other dysploidy events may cause speciation, either because of an accumulation of dif-

ferences that eventually leads to incompatibility or because of the genomic signature of the dysploidy event itself. We

describe additional theory on dysploidy and macroevolution in Supplemental Section S1.

Holocentricity is distributed broadly across the Tree of Life, including 18 different lineages in animals, plants, and

rhizaria (Escudero, Márquez-Corro & Hipp 2016, Márquez-Corro, Martín-Bravo, Pedrosa-Harand, Hipp, Luceño &

Escudero 2019), approximately 15–20% of eukaryotic species (Márquez-Corro et al. 2018). Two particularly diverse

holocentric clades show extraordinary chromosome number variation: the insect order Lepidoptera (de Vos et al. 2020)

and the angiosperm sedge family Cyperaceae ($2n = 4-224$; Márquez-Corro et al. 2021, Márquez-Corro, Martín-Bravo,

Spalink, Luceño & Escudero 2019). Holocentric chromosomes, instead of having kinetochore activity concentrated in a

single point (*i.e.*, at a single centromere in monocentric chromosomes), have centromeric regions distributed along the

whole chromosome where the kinetochores assemble in most of the organisms with such chromosome type (Márquez-

Corro, Martín-Bravo, Pedrosa-Harand, Hipp, Luceño & Escudero 2019, Marques et al. 2015). In monocentric chromo-

somes, many chromosome fissions are expected to result in a loss of genetic material during meiosis and inviable

gametes, as chromosome fragments without centromeres are unable to segregate normally, and fusions are generally

inherited by combining two telocentric chromosomes (Robertsonian translocations; Robertson 1916). Holocentric chro-

mosomes, by contrast, allow chromosome fragments to segregate normally during meiosis (Faulkner 1972). Holocen-

tricity may promote chromosome number variation via fission and fusion, as these changes are expected to be neutral or

nearly so in holocentric organisms (Márquez-Corro, Martín-Bravo, Pedrosa-Harand, Hipp, Luceño & Escudero 2019).

Thus, holocentric organisms provide a unique system in which to study chromosomal speciation (Lucek et al. 2022).

Nonetheless, of the few applied macroevolutionary studies that address the role of dysploidy in lineage diversifica-

tion, most have focused on monocentric chromosomes, which have a single centromere (*e.g.*, Ayala & Coluzzi (2005),

Freyman & Höhna (2018), but see relevant work on Lepidoptera de Vos et al. (2020), among others).

*Carex*—the largest genus in Cyperaceae—is particularly well suited to studying the effect of dysploidy on plant

diversification because all *Carex* have holocentric chromosomes, the genus represents 40% of the third most species-rich

monocot family (among the tenth in angiosperms, POWO 2023) and well-developed phylogenetic and chromosome

number datasets for the genus to facilitate macroevolutionary studies (Martín-Bravo et al. 2019, Márquez-Corro et al.

2021). In *Carex*, karyotype evolves mainly through fusion, fission, and translocations, in contrast to in other sedge

lineages where karyotype evolves through both dysploidy and polyploidy (Márquez-Corro, Martín-Bravo, Spalink,

Luceño & Escudero 2019, Elliott et al. 2022, Shafir et al. 2023). *Carex* also has exceptional variability in chromosome

number, ranging from $2n = 10$ to $2n = 132$ (Márquez-Corro et al. 2021). *Carex* has experienced several rapid radiations

(Martín-Bravo et al. 2019), and shifts in optimal chromosome number are thought to have played a role in some of these

radiations (*e.g.*, *Carex* sect. *Cyperoideae*; Márquez-Corro et al. 2021, Hipp 2007).

Previous studies that have tested for a correlation between dysploidy and diversification (*e.g.*, Márquez-Corro et al.

2021) have relied on models that fail to account for alternative sources of variation in diversification rates—*e.g.*, morpho-

logical traits, climatic niche, or biotic interactions—that are not the study's focal trait (in our case, chromosome number;

Beaulieu & O'Meara 2016). Models that fail to account for alternative sources of variation in diversification rates have

high type-I error rates (Rabosky & Goldberg 2015) because they misattribute underlying diversification-rate variation

caused by unmeasured factors to variation due to the states included in the model. In other words, the null hypothesis

of those models assumes that there is no underlying diversification-rate variation. This high error rate has motivated

the development of hidden-state diversification models that account for underlying diversification-rate variation that

is not caused by the focal trait (Beaulieu & O'Meara 2016, Caetano et al. 2018). These models not only more accurately

test for associations between the focal trait and diversification, but also provide an opportunity to model how focal evolutionary processes vary across the phylogeny. Here, we design a new model of joint chromosome evolution and lineage diversification that incorporates process variation in chromosome number evolution and disentangles the effects of dysploidy from other unobserved factors that may also affect diversification rates (described in detail below). We apply our model to the most recent *Carex* time-calibrated phylogeny with chromosome number information that contains over 700 taxa and more than 50 states (Martín-Bravo et al. 2019, Márquez-Corro et al. 2021).

We test for a detectable effect of chromosome gains and losses on the rate of species formation in a large, diverse group of plants with high variation in chromosome number: *Carex* (Cyperaceae, Poales). We demonstrate decisively that dysploidy drives the formation of some new lineages, but diversification and dysploidy play out through two evolutionary modes. In one mode, chromosome rearrangements evolve rapidly and are linked to new species formation. In the other mode, rearrangements evolve less frequently and are not associated with new species formation; instead, other factors—hidden states unmeasured in our analysis—likely drive the formation of new species. Our results demonstrate the complexity of the diversification process and the important role of genomic rearrangements in determining biodiversity patterns at geological timescales. Furthermore, we illustrate how computational and statistical advances in modeling permit increasingly nuanced models that better represent the underlying complexity of biological systems.

# Materials and Methods

## Modeling chromosome number evolution

We developed the Chromosome number and Hidden State-dependent Speciation and Extinction model (ChromoHiSSE) to test the interaction between single chromosome number change, speciation, extinction, and diversification heterogeneity. Our model can be considered a natural extension of the ChromoSSE model (Freyman & Höhna 2018). Under the ChromoSSE model, lineages evolve independently under a continuous-time Markov chain (CTMC) that describes changes in chromosome numbers, speciation events, and extinction events; each of these events occurs at a particular rate (interpreted as the expected number of events per lineage per unit time). The ChromoHiSSE model includes an additional hidden trait with $m > 1$ states. The states of this hidden trait correspond to different sets of ChromoSSE parameters, and lineages evolve among hidden states as a Markov process with an estimated rate.

Under the ChromoHiSSE model, the state of a lineage is both the chromosome number, $n$, and the hidden state, $i$. For numerical stability (Mayrose et al. 2010, Zenil-Ferguson et al. 2017, Freyman & Höhna 2018), we place an upper bound on the possible number of chromosomes ($k$; transitions to $n > k$ are prohibited, *i.e.*, have rate 0). We denote the two hidden states as $i$ and $ii$. For example, the state for a lineage could be $10i$, indicating that the lineage has 10 haploid chromosomes and is in the hidden state $i$.

The ChromoHiSSE model is a stochastic process that begins with two lineages at the root, which evolve indepen-

dently forward in time. As the process evolves, a lineage can experience anagenetic events (changes in chromosome number or hidden state that happen along the branches of the phylogeny, denoted with superscript [a]), cladogenetic events (speciation events that involve changes in the number of chromosomes or hidden state for one of the daughter lineages, hence associated with speciation and denoted with superscript [c]), and extinction events. As in all continuous-time Markov chain models, each event happens at an instantaneous point in time and multiple events cannot occur concurrently. The anagenetic events are:

(1) $n$ increases by one ($n+1$ increasing dysploidy) but the hidden state stays the same, which occurs at rate $\gamma_i^a$;

(2) $n$ decreases by one ($n-1$ decreasing dysploidy) but the hidden state stays the same, which occurs at rate $\delta_i^a$, and;

(3) $n$ stays the same, but the hidden state changes (*e.g.*, $i$ to $ii$), which occurs at rate $\chi^a$.

Cladogenetic events produce two daughter lineages that then evolve independently. The states of the daughters depend on the type of event:

(4) both daughters inherit the state of the ancestor, which occurs at rate $\phi_i^c$;

(5) one daughter inherits $n$, the other inherits $n+1$ (increasing dysploidy), and both inherit the same hidden state $i$, which occurs with rate $\gamma_i^c$;

(6) one daughter inherits $n$, the other inherits $n-1$ (decreasing dysploidy), and both inherit the same hidden state $i$, which occurs with rate $\delta_i^c$;

(7) both daughters inherit $n$ from the ancestor, but one daughter changes hidden state ($i$ to $ii$), which occurs at rate $\chi^c$.

These events are depicted in Fig. 1. Additionally, all lineages go extinct at rate $\mu$, independent of $n$ or the hidden state. The lineages evolve forward in time until the present, at which point they are sampled independently with probability $f$. Extinct and unsampled lineages are pruned from the tree and the hidden state for sampled lineages is ignored; this produces a realization comprising a reconstructed phylogeny relating the sampled lineages and a chromosome number for each sampled lineage. This stochastic process allows us to compute the probability of an observed dataset (*i.e.*, the probability that a realization under this process corresponds to our observed data) given a set of parameter values.

Note that we assume that there are only two hidden states and that rates of change between hidden states are symmetric; *i.e.*, the anagenetic rate of change from $i$ to $ii$ is the same as the rate of change from $ii$ to $i$, and likewise for cladogenetic changes. In addition, we assume that the rate of extinction is constant among all lineages, regardless of the number of chromosomes or hidden state. In the supplementary material (S2), we show how to relax some of these assumptions by calculating a full ChromoHiSSE model that also includes polyploidization (and demipolyploidization).

**Figure 1:** The ChromoHiSSE model. Panel (A) describes the event rates allowed in the model for both cladogenetic (top) and anagenetic (bottom) events. Panels (B) and (C) demonstrate those rates on a tree simulated under ChromoHiSSE. Panel (B) shows the anagenetic and cladogenetic changes in the hidden states, indicated by blue (*i*) vs. orange (*ii*). Panel (C) shows the anagenetic and cladogenetic gains and losses of chromosomes, as well as speciation with no corresponding change in chromosome numbers. More red colors in (C) correspond to more chromosomes. The vertical blue and orange bars in (C) indicate clades in the blue (*i*) vs. orange (*ii*) hidden states, displaying that chromosome number changes are less frequent in the blue hidden state than in the orange.

## Testing differences between transition rates and not states

Most hidden-state SSE (HiSSE) models include two (or more) sets of diversification rates for each state hypothesized to affect diversification (*e.g.*, Helmstetter et al. 2023). For example, if a researcher wants to know if herbivorous beetles diversify faster than carnivorous ones, the hidden state approach would include two diversification parameters for the carnivorous state and two diversification parameters for the herbivorous state. If both herbivory-specific rates are higher than the carnivory rates, then the analysis suggests that herbivores diversify faster diversification than carnivores. However, our new model differs from previous HiSSE models in that we parameterize diversification rates associated with *types* of state transitions rather than with the states themselves. This means that our goal is not to test whether $n = 15$ or $n = 16$ chromosomes have different modes of diversification, but rather if changes in the karyotype (*i.e.*, increase: $n = 15$ to $n = 16$ or decrease: $n = 15$ to $n = 14$) are associated with faster or slower diversification rates compared to heterogeneity (or noise) in the diversification process. To do this, we compare speciation rates *not* associated with dysploidy ($\phi_i^c$ and $\chi^c$) to speciation rates *associated* with dysploidy ($\gamma_i^c$ and $\delta_i^c$) by defining the test statistic $T_i = \gamma_i^c + \delta_i^c - (\phi_i^c + \chi^c)$ (Fig. 2D). A posterior probability of $T_i > 0$ greater than 0.95 ($P(T_i > 0) \geq 0.95$) represents high probability of speciation by single chromosome number. In contrast, $P(T_i < 0) \geq 0.95$ suggests that speciation rates unrelated to chromosome number are significantly faster than speciation by single chromosome number. Test statistic $T_{ii}$ is defined analogously for the second hidden state.

## Chromosomal and phylogenetic data for *Carex*

We implemented our proposed ChromoHiSSE model on a large dataset that includes a phylogeny of *Carex* with 755 taxa ($\sim$ 40% of extant diversity), representing all *Carex* subgenera and most of the sections for which chromosome counts have been reported. Our dataset also includes haploid chromosome number (*n*) for all tips in the tree; both the tree and chromosome number data come from (Márquez-Corro et al. 2021). The original tree, based on a HybSeq backbone and three DNA regions (ITS, ETS and *mat*K) from ca. 1400 species out of 2000 *Carex* species, was published by (Martín-Bravo et al. 2019). Tips without chromosome number information were pruned for the current study. Chromosome number in *Carex* evolves through dysploid events, except in the small subgenus *Siderostictae* (13 species in our tree)—sister to the rest of the genus—which includes species with different reported ploidy levels (Márquez-Corro, Martín-Bravo, Spalink, Luceño & Escudero 2019, Márquez-Corro et al. 2021). To avoid modeling rare polyploidy events that occur only in a small part of the tree, we removed *Siderostictae* from the primary analysis (though see Supplemental Section S3 for results with this clade included). For the remaining taxa, we coded chromosome numbers as the most frequent haploid number or the lowest haploid cytotype in polyploid lineages from the Márquez-Corro et al. (2021) dataset for those tips. The final data matrix included *n*-values ranging from 5 to 66 chromosomes for 742 taxa.

## Computational implementation of ChromoHiSSE

We implemented ChromoHiSSE in `RevBayes` (Höhna et al. 2016), a software for specifying Bayesian probabilistic graphical models primarily for phylogenetics and phylogenetic comparative methods. Due to the large state space and cladogenesis in our model, we used the newly-developed `TensorPhylo` plugin (May & Meyer 2022) to accelerate likelihood calculations and thus achieve convergence in a more reasonable time frame. We ran two chains of the analysis and assessed convergence in the R programming language (R Core Team 2013). We processed the traces and removed 10% of the generations per chain as burnin using `RevGadgets`. We then calculated the effective sample size (ESS) value for each parameter in each chain using `coda` (Plummer et al. 2006) and verified that the harmonic mean of the ESS values of each chain was greater than 200. We additionally visually inspected all model parameters across both chains in Tracer (Rambaut et al. 2018). For any parameters that appeared to have strikingly non-normal posterior distributions, we also estimated a transformed ESS following (Vehtari et al. 2021). We subsequently combined both runs for all downstream analyses, discarding 10% of the total generations per chain as burnin prior to combining.

## Analysis and postprocessing

We summarized posteriors and plotted results in R (R Core Team 2013) using the R package `RevGadgets` (Tribble et al. 2022). We additionally transformed model parameters to produce two types of useful summary statistics. First, we calculated the total speciation rate in hidden state *i* vs. *ii* as $\lambda = \chi^c + \gamma^c + \delta^c + \phi^c$. Second, we wanted to know if the rate of speciation concurrent with changes in chromosome number ($\gamma^c + \delta^c$) is greater or less than the rate of speciation with

no change in chromosome number ($\chi^c + \phi^c$). We thus estimated an additional metric: $T = (\gamma^c + \delta^c) - (\chi^c + \phi^c)$, positive

values of *TD* indicate more speciation with chromosome number change and negative values indicate less speciation

with chromosome number change. We estimate *T* for each hidden state to estimate how chromosome number changes

associated with speciation vary between modes of the model.

All code for implementing and running the model and processing and plotting the results is available at Zenodo

DOI: 10.5281/zenodo.8320249.

# Results

## Two modes of chromosomal anagenesis

We recover two modes of anagenetic dysploidy. Branches in hidden state *i* have fast and recurrent single chromosome

number increases and decreases ($\gamma_i^a = 14.63$ and $\delta_i^a = 14.19$ chromosome number events per lineage per million

years (E/L/Myr) Table 1, Fig. 2B). For the other branches—in hidden state *ii*—chromosome number rarely changes via

dysploidy (increases are $\gamma_{ii}^a = 0.39$ and decreases are $\delta_{ii}^a = 0.14$ E/L/Myr, Table 1, Fig. 2B). Rarely, lineages transition

between hidden states ($\chi^a = 0.02$). In Table 1 we show the credible intervals for these estimates, and in figure 2B we

show the posterior distribution of all the anagenetic rate estimates.

**Table 1:** Summary statistics of posterior distributions for parameter estimates for the ChromoHiSSE model. Light blue rows correspond to parameter estimates for hidden state *i*, and light orange rows correspond to parameter estimates for hidden state *ii*. All rate estimates are given in units of events per lineage per million years (E/L/Myr)

| | Model parameter | Parameter Type | Hidden State | Median | 2.5% Quantile | 97.5% Quantile |
|---|---|---|---|---|---|---|
| $\chi^c$ | Hidden state change | Cladogenetic | $i, ii$ | 0.113 | 0.051 | 0.215 |
| $\gamma_i^c$ | Increasing dysploidy | Cladogenetic | $i$ | 0.330 | 0.021 | 0.772 |
| $\gamma_{ii}^c$ | Increasing dysploidy | Cladogenetic | $ii$ | 0.077 | 0.003 | 0.261 |
| $\delta_i^c$ | Decreasing dysploidy | Cladogenetic | $i$ | 0.539 | 0.120 | 0.916 |
| $\delta_{ii}^c$ | Decreasing dysploidy | Cladogenetic | $ii$ | 0.099 | 0.007 | 0.250 |
| $\phi_i^c$ | No character Change | Cladogenetic | $i$ | 0.072 | 0.003 | 0.363 |
| $\phi_{ii}^c$ | No character Change | Cladogenetic | $ii$ | 0.849 | 0.642 | 1.058 |
| $\chi^a$ | Hidden state change | Anagenetic | $i, ii$ | 0.021 | 0.001 | 0.093 |
| $\gamma_i^a$ | Increasing dysploidy | Anagenetic | $i$ | 14.638 | 12.282 | 17.626 |
| $\gamma_{ii}^a$ | Increasing dysploidy | Anagenetic | $ii$ | 0.397 | 0.121 | 0.769 |
| $\delta_i^a$ | Decreasing dysploidy | Anagenetic | $i$ | 14.191 | 11.903 | 17.246 |
| $\delta_{ii}^a$ | Decreasing dysploidy | Anagenetic | $ii$ | 0.142 | 0.005 | 0.505 |

**Figure 2:** Posterior distributions of rate estimates. Solid lines indicate rates corresponding to hidden state *i* and dashed lines indicate hidden state *ii*. Panel (A) shows the posterior distributions of cladogenetic rates, panel (B) shows the anagenetic rates, panel (C) shows the total speciation rate per hidden state ($\lambda$), and panel (D) shows the test statistic ($T_i$ and $T_{ii}$) of the difference in speciation rate associated with chromosome number change and speciation rate without chromosome number change. $T_i$ is significantly greater than 0 (faster speciation when associated with chromosome number change; $P(T_i > 0) = 0.986$) and $T_{ii}$ is significantly less than 0 (slower speciation when associated with chromosome number change; $P(T_{ii} < 0) = 1.000$)

## Two modes of chromosomal speciation

We identify variation in modes of chromosomal evolution across the tree via two distinct modes, corresponding to the two hidden states *i*, *ii*. While the overall total speciation rate ($\lambda$) between the two modes is very similar (Fig. 2C), rates of dysploidy and dysploidy-driven speciation vary significantly between the modes.

Hidden state *i* is characterized by a positive association between speciation and dysploidy and generally high rates of cladogenetic dysploidy (Fig. 2, solid lines, and Table 1). In this mode, the difference between the rate of speciation with chromosome number change and speciation without chromosome number change is larger than zero with 98.6% probability ($P(T_{ii} > 0) = 0.986$). Thus in hidden state *i*, there is a highly probable and positive correlation between chromosome number change and speciation rates (Fig. 2D, blue). Every estimated dysploidy-related cladogenetic rate is higher than non-dysploidy rates (*e.g.*, speciation with no state change, Fig. 2A), which suggests that speciation happens more frequently when associated with changes to karyotype.

Cladogenetic increasing and decreasing dysploidy ($\gamma_i^c$ and $\delta_i^c$) are higher than 0 with 99% probability(Table 1) and

faster than the cladogenetic rate of no character change (speciation without change in chromosome number, $\phi_i^c$). While

$\delta_i^c$ is slightly faster than $\gamma_i^c$, these rates overlap (Table 1).

In contrast, hidden state *ii* is characterized by a negative association between speciation and dysploidy and slower

rates of anagenetic and cladogenetic dysploidy (Fig. 2, dashed lines, and Table 1). The difference between speciation as-

sociated with chromosome number change and speciation without chromosome number change is less than 0 E/L/Myr

($P(T_{ii} < 0) = 1.000$), suggesting that speciation is slower when associated with a chromosome number change in hid-

den state *ii* (Fig. 2D, orange). Anagenetic rates ($\gamma_{ii}^a$ and $\delta_{ii}^a$) are quite low, approximating 0 (Table 1, Fig. 2B), suggesting

that little to no anagenetic change in chromosome number happens in hidden state *ii*.

Cladogenetic dysploidy rates are also quite low, and while the rate of decreasing dysploidy ($\delta_{ii}^c$) is slightly higher

than increasing dysploidy ($\gamma_{ii}^c$), this difference is minimal (Table 1). In summary, we identify variation in modes of

chromosomal evolution across the tree and via two distinct modes, corresponding to the two hidden states *i*, *ii*. In

hidden state *i* dysploidy is linked to speciation, whereas in hidden state *ii*, with the same total speciation rate, dysploidy

is not an important contributor to this speciation rate.

## Reconstruction of chromosome number evolution

Just over half (51.85%) of all branches in the phylogeny showed a net change in chromosome number (Fig. 3). Overall,

the dominant pattern of chromosome number in *Carex* is one of frequent but gradual change; 75.05% of per-branch

net chromosome number changes are three or fewer gains or losses. Most clades vary in chromosome number despite

some shallow evolutionary divergences (Fig. 3A). Yet there are some remarkable exceptions. For example, clade 2—

which includes sect. *Pictae*, the Hirta Clade, and sect. *Praelongae*— shows substantial variation in chromosome number,

from $n = 13$ to $n = 66$. However, a clade of hooked sedges (*Carex* sect. *Uncinia*; Fig. 3, clade 1 in grey), has a constant

chromosome number ($n = 44$) across the 33 species included in this study, with one exception: *Carex perplexa* has a count

of $n = 66$, suggesting that this species may be a demipolyploid. While both clades highlighted in Fig. 3 have similar

crown ages, number of species, and average chromosome number ($n = 35.7$ vs. $n = 44.6$), they differ dramatically in

karyotype variability and, consequently, in the inferred hidden state (clade 1 = *ii*, clade 2 = *i*, Fig. 3C).

# Discussion

227 Although the impact of dysploidy on *Carex* diversification has been addressed previously (*e.g.*, Faulkner 1972, Márquez-

228 Corro, Martín-Bravo, Pedrosa-Harand, Hipp, Luceño & Escudero 2019, Márquez-Corro et al. 2021, Hipp 2007), this is

229 the first study to jointly model chromosome number change and diversification, and to demonstrate an association

230 between higher speciation rates and dysploidy in parts of the phylogeny despite heterogeneity in the process of diver-

231 sification. While gains and losses in chromosome number spur diversification along parts of the phylogeny (hidden

state *i*), they have the opposite effect elsewhere in the tree (hidden state *ii*, see Fig. 3). Furthermore, while dysploidy does not lead to higher rates of speciation across the tree on average, speciation in some clades is driven strongly by dysploidy. We propose that these discrepancies between clades may be due to the nature of holocentric chromosomes, where a single dysploidy event in isolation may not be enough to trigger reproductive isolation (Hipp et al. 2010, Lucek et al. 2022, Márquez-Corro, Martín-Bravo, Pedrosa-Harand, Hipp, Luceño & Escudero 2019, Escudero, Hahn, Brown, Lueders & Hipp 2016, Whitkus 1988), but the accumulation of sufficient chromosomal rearrangements in a lineage may form a reproductive barrier and thus trigger speciation (Escudero, Hahn, Brown, Lueders & Hipp 2016, Whitkus 1988, Baker & Bickham 1986). This is a central hypothesis discussed in Fig. 4 of Lucek et al. (Lucek et al. 2022) known as the recombination suppression/hybrid dysfunction chromosomal speciation model. Below, we elaborate on the differences between speciation modes, discuss the evidence supporting the recombination suppression/hybrid disfunction model, and provide suggestions for future research.

Some *Carex* species are known to have striking chromosome-number polymorphism, even within populations (Luceño & Castroviejo 1991, Escudero, Arroyo, González-Ramírez & Jordano 2023, Escudero et al. 2013, Whitkus 1988). For example, *Carex scoparia* varies from $n = 28$ to $n = 35$ (Escudero et al. 2013), and individuals with different chromosome numbers are able to reproduce and exchange alleles, maintaining gene flow despite chromosome number differences. This polymorphism suggests that some chromosome number differences are not sufficient to create reproductive barriers (Hipp et al. 2009), which is supported by our results; often, inferred chromosome number changes do not result in an immediate speciation event on the phylogeny. The question remains, why does dysploidy result in speciation in some cases and not in others? One possibility is that an accumulation of changes may eventually lead to reproductive isolation, where one "last straw" dysploidy event triggers speciation (referred to as the last-straw hypothesis). We discuss potential model developments that could test this theory below in the *Modeling chromosomal speciation* section. Another possibility is that rearrangements in some parts of the genome are more stable than others, and the genomic architecture—where the fragmentation or fusion occurs in the genome—determines the evolutionary effects of dysploidy.

Patterns of repeat DNA—*e.g.*, LINEs, LTRs, and Helitrons—differ significantly between *Carex* lineages whose chromosome arrangements evolve at different rates (Cornet et al. 2023). Moreover, repeat regions in *Carex* genomes investigated correlate with chromosome breakpoints across deep phylogenetic breaks as well as within species (Escudero, Marques, Lucek & Hipp 2023, Höök et al. 2023), pointing to a mechanism by which natural selection could operate on the genome architecture of chromosome rearrangements in the genus. In fact, synteny in holocentric sedge chromosomes is more conserved than we would expect if chromosome evolution were unconstrained, even in comparisons between species that span deep nodes in the phylogeny (Escudero, Marques, Lucek & Hipp 2023). This suggests that selection may maintain large blocks of the genome, likely comprising numerous, tightly linked genes (Escudero, Marques, Lucek & Hipp 2023). The massive diversity of sedges is thus likely an outcome of two processes: recombination suppression in rearranged regions of the genome, shaping ecological divergence in during species divergence; and the typically-gradual—or, occasionally, immediate—evolution of reproductive isolation as chromosomes split and fuse.

**Figure 3:** Reconstruction of chromosome numbers (A and B) and hidden states (C) on the *Carex* phylogeny. Panel (A) shows the distribution of haploid chromosome numbers for all extant taxa included in the analysis. Panel (B) shows the reconstructed evolution of chromosome number along branches of the phylogeny, where lighter colors indicate more chromosomes. Panel (C) shows the reconstructed evolution of the hidden state along branches of the phylogeny, where blue indicates strong statistical support for state A, orange indicates strong support for state B, and intermediary colors indicate uncertainty in the estimates. Grey bars highlight two clades (labeled 1 and 2) that are discussed in the main text.

267     An important caveat to our study is the presence of missing data—particularly for tropical lineages—both in terms

268     of sequenced taxa represented in the phylogeny and for available chromosome counts. While the phylogeny used in

269     this study was assembled with a HybSeq backbone and three DNA regions for ca. 1400 out of 2000 species, phylogenetic

270     relationships in some areas of the genus are still tenuous (Martín-Bravo et al. 2019, Roalson et al. 2021, Jiménez-Mejías

et al. 2016). Additionally, there are only chromosome counts for $\sim 700$ species in the phylogeny, and data availability is more sparse in tropical lineages than temperate ones. For example, one of the most species-diverse lineages within *Carex*—the Decora Clade/*Carex* sect. *Indicae* (Roalson et al. 2021)—only includes five species with reported chromosome numbers (Márquez-Corro et al. 2021). This pantropical lineage alone would constitute ca. 10% of the genus (Roalson et al. 2021). Also, we include only a few New Zealand species from *Carex* sect. *Uncinia* (clade 1 in Fig. 3), as there are few known karyotypes for the South American representatives of the lineage. Dispersal traits such as hooked utricles and epizoochory might be as or more important for diversification as karyotype in this clade, but we cannot tease these effects apart without additional chromosome count data. *Carex* biodiversity is greatest in temperate regions including the Global North, and this bias in data availability may suggest that our findings are most robust for northern temperate *Carex*. Filling these data gaps would facilitate testing if evolutionary modes of dysploidy play out differently based on geography and/or dispersal traits.

## Chromosome number evolution

The contrasting clades 1 and 2 from Fig. 3 reflect the two modes of macroevolution described above in "Results: Reconstruction of chromosome number evolution" and in "Discussion: Multiple modes of chromosomal speciation" exemplify the importance of including hidden states in our model. *Carex* sect. *Uncinia* (hook sedges, clade 1 in Fig. 3) is characterized by hooked utricles that allow for long-distance dispersal through epizoochory (García-Moro et al. 2022). The hook sedges exemplify a *Carex* clade that is characterized by low-to-zero dysploidy but high rates of speciation (Martín-Bravo et al. 2019)—though this was not formally tested in our study—and thus may be a particularly appealing candidate for future studies focusing on potential adaptive traits (currently "hidden" and not related to chromosome number) that drive diversification. This clade is characterized by similar karyotypes of many ($n = 44$) chromosomes (Márquez-Corro et al. 2021), mostly reported from New Zealand, where ca. half of the section diversified. However, this only gives partial information on their evolutionary history, as the karyotypes of the Andean relatives have not been studied in detail yet. Most of the New Zealand chromosome counts correspond to a single, old report and there is some variation in the few existing counts from South America, so these results may be taken with caution. If the South American species show wider karyotypic variation than currently detected in comparison to the New Zealand lineage, there may be different drivers enhancing speciation on each side of the South Pacific.

## Modeling chromosomal speciation

The methodological innovations presented here allow us to integrate process noise and variation in chromosome number evolution, despite the computational challenges associated with the huge number of states defined by chromosome numbers. Process variation is fundamental to our conclusions; only by incorporating multiple diversification modes do we discover that while dysploidy is strongly associated with speciation in some clades, in other parts of the phylogeny, dysploidy is rare and does not lead to faster speciation. Leaving the hidden states out leads to a false inference about the

significance of chromosome evolution to diversification: when we implement our model without process variation, our results suggest a strong, uniform boost in speciation rate associated with dysploidy (results presented in Supplemental Section S3). Future work that builds off of our ChromoHiSSE approach will allow us to pursue promising avenues for innovative research; we highlight two examples below.

First, like all birth-death models, our ChromoHiSSE model operates with species as the fundamental unit of analysis (the tips in the tree) and thus does not formally model chromosome number polymorphism that is present in some *Carex* lineages. Second, while ChromoHiSSE tests for the effect of single changes in chromosome number on diversification rates, it cannot test for the effect of an accumulation of changes (the last-straw hypothesis). However, ChromoHiSSE could be modified to include tip-state polymorphism (*e.g.*, a dysploid series) as additional hidden states (*e.g.*, a particular tip either has 12, 13, or 14 chromosomes). Additionally, (Goldberg & Foo 2020) described a mechanism for modeling memory (thus the accumulation of chromosome number changes) in a macroevolutionary framework using hidden states, which could be applied to test the last-straw hypothesis.

## Conclusions

Our work demonstrates the important role of dysploidy on diversification in *Carex*, a model lineage for understanding how karyotype rearrangements via dysploidy affect speciation and macroevolutionary dynamics. Our results—using the new ChromoHiSSE model—paint a complex picture of how dysploidy affects speciation in a clade characterized by high species diversity, high morphological disparity, and holocentric chromosomes (Martín-Bravo et al. 2019), and our results support the recombination suppression/hybrid dysfunction chromosomal speciation model, in which only some karyotype rearrangements trigger reproductive isolation and thus speciation. Future work on the underlying genomic mechanisms of chromosomal speciation via comparative genomics will be particularly powerful for linking across scales, from molecules to lineages. Ultimately, our novel modeling approach also serves as a critical step towards even more complex and powerful macroevolutionary analyses that incorporate intraspecific chromosome number variation and track the accumulation of change through time.

## Author Contributions

RZF and ME conceived the study. CMT, JIM-C, ME, and RZF wrote the article. JIM-C, ME, ALH gathered all *Carex* chromosome number and phylogenetic data. CMT, MRM, and RZF designed models and methods, and verified mathematical/ statistical notation. CMT and MRM wrote computational code and created figures. All authors, revised and edited the manuscript.

# Data availability

RevBayes model code, and R code for processing and plotting the results are available at Zenodo DOI: 10.5281/zenodo.8320249

# Acknowledgements

CMT was supported by the School of Life Sciences at the University of Hawaiʻi at Mānoa funded through the Office of the Vice President of Research. This material is based upon work supported by the NSF Postdoctoral Research Fellowships in Biology Program under Grant No. 2109835 to CMT. JIM-C was granted by the 'Next Generation EU' funding, the Recovery Plan, Transformation and Resilience and the Ministry of Universities, under the grants 'Margarita Salas' for the requalification of the Spanish university system 2021–2023 called by the Universidad Pablo de Olavide, Seville. ME was supported by the MICINN-FEDER, Project DiversiChrom (PID2021-122715NB-I00). RZF was supported by NSF-DEB 2323170.

# Supporting Information

**Supporting Information 1.** Detailed description of the Chromosome number and Hidden State-dependent Speciation and Extinction model (ChromoHiSSE).

# References

Ayala, F. J. & Coluzzi, M. (2005), 'Chromosome speciation: humans, *Drosophila*, and mosquitoes', *Proceedings of the National Academy of Sciences* **102**, 6535–6542.

Baker, R. J. & Bickham, J. W. (1986), 'Speciation by monobrachial centric fusions', *Proceedings of the National Academy of Sciences* **83**(21), 8245–8248.

Beaulieu, J. M. & O'Meara, B. C. (2016), 'Detecting hidden diversification shifts in models of trait-dependent speciation and extinction', *Systematic biology* **65**(4), 583–601.

Caetano, D. S., O'Meara, B. C. & Beaulieu, J. M. (2018), 'Hidden state models improve state-dependent diversification approaches, including biogeographical models', *Evolution* **72**(11), 2308–2324.

Cornet, C., Pablo, M., Augustijnen, H., Nguyen, P., Escudero, M. & Lucek, K. (2023), 'Holocentric repeat landscapes: from microevolutionary patterns to macroevolutionary associations with karyotype evolution', *Molecular Ecology* **000**(000), 000–000.

de Vos, J. M., Augustijnen, H., Bätscher, L. & Lucek, K. (2020), 'Speciation through chromosomal fusion and fission in Lepidoptera', *Philosophical Transactions of the Royal Society B: Biological Sciences* **375**(1806), 20190539.

Elliott, T. L., Zedek, F., Barrett, R. L., Bruhl, J. J., Escudero, M., Hroudová, Z., Joly, S., Larridon, I., Luceño, M., Márquez-Corro, J. I., Martín-Bravo, S., Muasya, A. M., Šmarda, P., Thomas, W. W., Wilson, K. L. & Bureš, P. (2022), 'Chromosome size matters: genome evolution in the cyperid clade', *Annals of Botany* **130**(7), 999–1014.

Escudero, M., Arroyo, J. M., González-Ramírez, S. & Jordano, P. (2023), 'Founder events and subsequent genetic bottlenecks underlie karyotype evolution in the ibero - north african endemic *Carex helodes*', *Annals of Botany* **000**, 000–000.

Escudero, M., Hahn, M., Brown, B. H., Lueders, K. & Hipp, A. L. (2016), 'Chromosomal rearrangements in holocentric organisms lead to reproductive isolation by hybrid dysfunction: the correlation between karyotype rearrangements and germination rates in sedges', *American Journal of Botany* **103**(8), 1529–1536.

Escudero, M., Marques, A., Lucek, K. & Hipp, A. L. (2023), 'Genomic hotspots of chromosome rearrangements explain conserved synteny despite high rates of chromosome evolution in a holocentric lineage', *Molecular Ecology* **000**(000), 000–000.

Escudero, M., Márquez-Corro, J. I. & Hipp, A. L. (2016), 'The phylogenetic origins and evolutionary history of holocentric chromosomes', *Systematic Botany* **41**(3), 580–585.

Escudero, M., Martín-Bravo, S., Itay, M., Fernández-Mazuecos, M., Fiz-Palacios, O., Hipp, A. L., Pimentel, M., Valcárcel, V., Vargas, P. & Luceño, M. (2014), 'Karyotypic changes through dysploidy persist longer over evolutionary time than polyploid changes', *PloS One* **9**(1), e85266.

Escudero, M., Weber, J. A. & Hipp, A. L. (2013), 'Species coherence in the face of karyotype diversification in holocentric organisms: the case of a cytogenetically variable sedge (*Carex scoparia*, Cyperaceae)', *Annals of Botany* **112**(3), 515–526.

Faulkner, J. (1972), 'Chromosome studies on *Carex* section *Acutae* in north-west Europe', *Botanical Journal of the Linnean Society* **65**(3), 271–301.

Freyman, W. A. & Höhna, S. (2018), 'Cladogenetic and anagenetic models of chromosome number evolution: a Bayesian model aver-
aging approach', *Systematic Biology* **67**(2), 195–215.

García-Moro, P., Otero, A., Benítez-Benítez, C., Costa, L., Martín-Bravo, S., Naczi, R. F., Reznicek, A., Roalson, E. H., Starr, J. R. &
Jiménez-Mejías, P. (2022), 'Biogeography and systematics of *Carex* subgenus *Uncinia* (Cyperaceae): A unique radiation for the genus
*Carex* in the Southern Hemisphere', *Taxon* **71**(3), 587–607.

Goldberg, E. E. & Foo, J. (2020), 'Memory in trait macroevolution', *The American Naturalist* **195**(2), 300–314.

Helmstetter, A. J., Zenil-Ferguson, R., Sauquet, H., Otto, S. P., Méndez, M., Vallejo-Marin, M., Schönenberger, J., Burgarella, C., An-
derson, B., de Boer, H., Glémin, S. & Käfer, J. (2023), 'Trait-dependent diversification in angiosperms: Patterns, models and data',
*Ecology Letters* **26**(4), 640–657.

Hipp, A. L. (2007), 'Nonuniform processes of chromosome evolution in sedges (*Carex*: Cyperaceae)', *Evolution* **61**(9), 2175–2194.

Hipp, A. L., Rothrock, P. E., Whitkus, R. & Weber, J. A. (2010), 'Chromosomes tell half of the story: the correlation between karyotype
rearrangements and genetic diversity in sedges, a group with holocentric chromosomes', *Molecular Ecology* **19**, 3124–3138.

Hipp, A., Rothrock, P. & Roalson, E. (2009), 'The evolution of chromosome arrangements in *Carex* (Cyperaceae)', *The Botanical Review*
**75**, 96–109.

Höhna, S., Landis, M. J., Heath, T. A., Boussau, B., Lartillot, N., Moore, B. R., Huelsenbeck, J. P. & Ronquist, F. (2016), 'RevBayes:
Bayesian phylogenetic inference using graphical models and an interactive model-specification language', *Systematic Biology*
**65**(4), 726–736.

Höök, L., Näsvall, K., Vila, R., Wiklund, C. & Backström, N. (2023), 'High-density linkage maps and chromosome level genome assem-
blies unveil direction and frequency of extensive structural rearrangements in wood white butterflies (*Leptidea* spp.)', *Chromosome*
*Research* **31**(1), 2.

Jiménez-Mejías, P., Hahn, M., Lueders, K., Starr, J. R., Brown, B. H., Chouinard, B. N., Chung, K.-S., Escudero, M., Ford, B. A., Ford,
K. A., Gebauer, S., Gehrke, B., Hoffmann, M. H., Jin, X.-F., Jung, J., Kim, S., Luceño, M., Maguilla, E., Martín-Bravo, S., Míguez, M.,
Molina, A., Naczi, R. F. C., Pender, J. E., Reznicek, A. A., Villaverde, T., Waterway, M. J., Wilson, K. L., Yang, J.-C., Zhang, S., Hipp,
A. L. & Roalson, E. H. (2016), 'Megaphylogenetic Specimen-Level Approaches to the *Carex* (Cyperaceae) Phylogeny Using ITS, ETS,
and *mat*K Sequences: Implications for Classification The Global *Carex* Group)', *Systematic Botany* **41**(3), 500–518.

Lucek, K., Augustijnen, H. & Escudero, M. (2022), 'A holocentric twist to chromosomal speciation?', *Trends in Ecology & Evolution*
**37**(8), 655–662.

Luceño, M. & Castroviejo, S. (1991), 'Agmatoploidy in *Carex laevigata* (Cyperaceae). Fusion and fission of chromosomes as the mecha-
nism of cytogenetic evolution in Iberian populations', *Plant Systematics and Evolution* **117**, 149–159.

Magallón, S. & Castillo, A. (2009), 'Angiosperm diversification through time', *American journal of botany* **96**(1), 349–365.

Magallón, S., Sánchez-Reyes, L. L. & Gómez-Acevedo, S. L. (2019), 'Thirty clues to the exceptional diversification of flowering plants',
*Annals of Botany* **123**(3), 491–503.

Mandáková, T. & Lysak, M. A. (2018), 'Constraints on polyploid evolution: a test of the minority cytotype exclusion principle', *Current*
*Opinion in Plant Biology* **42**, 55–65.

Marques, A., Ribeiro, T., Neumann, P., Macas, J., Novák, P., Schubert, V., Pellino, M., Fuchs, J., Ma, W., Kuhlmann, M., Brandt, R., Vanzela, A. L. L., Beseda, T., Šimková, H., Pedrosa-Harand, A. & Houben, A. (2015), 'Holocentromeres in *Rhynchospora* are associated with genome-wide centromere-specific repeat arrays interspersed among euchromatin', *Proceedings of the National Academy of Sciences* **112**(44), 13633–13638.

Márquez-Corro, J. I., Escudero, M. & Luceño, M. (2018), 'Do holocentric chromosomes represent an evolutionary advantage? a study of paired analyses of diversification rates of lineages with holocentric chromosomes and their monocentric closest relatives', *Chromosome Research* **26**, 139–152.

Márquez-Corro, J. I., Martín-Bravo, S., Jiménez-Mejías, P., Hipp, A. L., Spalink, D., Naczi, R. F., Roalson, E. H., Luceño, M. & Escudero, M. (2021), 'Macroevolutionary insights into sedges (*Carex*: Cyperaceae): The effects of rapid chromosome number evolution on lineage diversification', *Journal of Systematics and Evolution* **59**(4), 776–790.

Márquez-Corro, J. I., Martín-Bravo, S., Spalink, D., Luceño, M. & Escudero, M. (2019), 'Inferring hypothesis-based transitions in clade-specific models of chromosome number evolution in sedges (Cyperaceae)', *Molecular Phylogenetics and Evolution* **135**, 203–209.

Márquez-Corro, J. I., Martín-Bravo, S., Pedrosa-Harand, A., Hipp, A. L., Luceño, M. & Escudero, M. (2019), *Karyotype evolution in holocentric organisms*, John Wiley & Sons, Ltd, pp. 1–7.

Martín-Bravo, S., Jiménez-Mejías, P., Villaverde, T., Escudero, M., Hahn, M., Spalink, D., Roalson, E. H., Hipp, A. L., Group, G. C., Benítez-Benítez, C. et al. (2019), 'A tale of worldwide success: Behind the scenes of *Carex* (Cyperaceae) biogeography and diversification', *Journal of Systematics and Evolution* **57**(6), 695–718.

May, M. R. & Meyer, X. (2022), 'TensorPhylo RevBayes plugin'. https://bitbucket.org/mrmay/tensorphylo/src/master.

Mayrose, I., Barker, M. S. & Otto, S. P. (2010), 'Probabilistic models of chromosome number evolution and the inference of polyploidy', *Systematic Biology* **59**(2), 132–144.

Plummer, M., Best, N., Cowles, K. & Vines, K. (2006), 'CODA: convergence diagnosis and output analysis for MCMC', *R news* **6**(1), 7–11.

POWO (2023), 'Plants of the World Online, Facilitated by the Royal Botanic Gardens, Kew', http://www.plantsoftheworldonline.org/. Retrieved: 2023-08-06.

R Core Team (2013), *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria.

Rabosky, D. L. & Goldberg, E. E. (2015), 'Model inadequacy and mistaken inferences of trait-dependent speciation', *Systematic Biology* **64**(2), 340–355.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. (2018), 'Posterior summarization in Bayesian phylogenetics using Tracer 1.7', *Systematic biology* **67**(5), 901–904.

Roalson, E. H., Jiménez-Mejías, P., Hipp, A. L., Benítez-Benítez, C., Bruederle, L. P., Chung, K.-S., Escudero, M., Ford, B. A., Ford, K., Gebauer, S., Gehrke, B., Hahn, M., Hayat, M. Q., Hoffmann, M. H., Jin, X.-F., Kim, S., Larridon, I., Léveillé-Bourret, E., Lu, Y.-F., Luceño, M., Maguilla, E., Márquez-Corro, J. I., Martín-Bravo, S., Masaki, T., Míguez, M., Naczi, R. F. C., Reznicek, A. A., Spalink, D., Starr, J. R., Uzma, Villaverde, T., Waterway, M. J., Wilson, K. L. & Zhang, S.-R. (2021), 'A framework infrageneric classification of *Carex* (Cyperaceae) and its organizing principles', *Journal of Systematics and Evolution* **59**(4), 726–762.

Robertson, W. (1916), 'Chromosome studies. I. Taxonomic relationships shown in the chromosomes of Tettigidae and Acrididae: V-shaped chromosomes and their significance in Acrididae, Locustidae, and Gryllidae: chromosomes and variation', *Journal of Morphology* **27**, 179—331.

Sauquet, H. & Magallón, S. (2018), 'Key questions and challenges in angiosperm macroevolution', *New Phytologist* **219**(4), 1170–1187.

Shafir, A., Halabi, K., Escudero, M. & Itay, M. (2023), 'A non-homogeneous model of chromosome-number evolution to reveal shifts in the transition patterns across the phylogeny', *New Phytologist* **238**(4), 1733–1744.

Tribble, C. M., Freyman, W. A., Landis, M. J., Lim, J. Y., Barido-Sottani, J., Kopperud, B. T., Höhna, S. & May, M. R. (2022), 'RevGadgets: an R Package for visualizing Bayesian phylogenetic analyses from RevBayes', *Methods in Ecology and Evolution* **13**(2), 314–323.

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. (2021), 'Rank-normalization, folding, and localization: An improved $\widehat{R}$ for assessing convergence of MCMC (with discussion)', *Bayesian Analysis* **16**(2), 667–718.

Whitkus, R. (1988), 'Experimental hybridizations among chromosome races of *Carex pachystachya* and the related species *C. macloviana* and *C. preslii* (Cyperaceae)', *Systematic Botany* **13**(1), 146–153.

Zenil-Ferguson, R., Ponciano, J. M. & Burleigh, J. G. (2017), 'Testing the association of phenotypes with polyploidy: An example using herbaceous and woody eudicots', *Evolution* **71**(5), 1138–1148.