PKA catalytic subunits dissociate from regulatory subunits to regulate

neuronal functions

Wei-Hong Xiong[a], Maozhen Qin[a], and Haining Zhong[a,1]

a. Vollum Institute, Oregon Health & Science University, Portland, OR 97239, USA

1. Correspondence should be addressed to H.Z. (email: zhong@ohsu.edu).

Haining Zhong, Ph.D.

Vollum Institute, Oregon Health & Science University

3181 SW Sam Jackson Park Road, L474

Portland, Oregon 97239

U.S.A.

(503) 494-5089 / zhong@ohsu.edu

Abstract

Protein kinase A (PKA) plays essential roles in diverse cellular functions. However, the

spatiotemporal dynamics of endogenous PKA upon activation remain debated. The classical

model predicts that PKA catalytic subunits dissociate from regulatory subunits in the

presence of cAMP, whereas a second model proposes that catalytic subunits remain

associated with regulatory subunits following physiological activation. Here we report that

different PKA subtypes, as defined by the regulatory subunit, exhibit distinct subcellular

localization at rest in CA1 neurons of cultured hippocampal slices. Nevertheless, when all

tested PKA subtypes are activated via the β-adrenergic receptor by norepinephrine,

catalytic subunits translocate to dendritic spines but regulatory subunits remain unmoved.

These differential spatial dynamics between the subunits indicate that at least a significant

fraction of PKA dissociates. Furthermore, PKA-dependent regulation of synaptic plasticity

and transmission can be supported only by wildtype, dissociable PKA, but not by non-

dissociable PKA. These results argue for the classical model in which endogenous PKA

regulatory and catalytic subunits dissociate to achieve PKA function.

Significance

Protein kinase A (PKA) is the major effector of the second messenger cAMP and plays a critical role in numerous cell biology processes. The function and specificity of PKA rely on spatial compartmentalization. However, the precise subcellular distribution and dynamics of activated PKA remain debated. Textbooks illustrate that upon activation PKA catalytic subunits dissociate from regulatory subunits. However, some recent work has proposed that PKA catalytic subunits remain associated with regulatory subunits following physiological activation. The present study tests these two models in the context of neuronal function. The results support the classical model of PKA activation. A clarified model of PKA action will contribute to the understanding of how the specificity of PKA phosphorylation may be achieved.

INTRODUCTION

Cyclic adenosine monophosphate (cAMP)-dependent kinase, or protein kinase A (PKA),

regulates diverse critical functions in nearly all mammalian cells. PKA is a tetrameric

protein consisting of two regulatory subunits (PKA-Rs) and two catalytic subunits (PKA-Cs)

(1, 2). In the inactive state, each PKA-R binds to and inhibits a PKA-C. Binding of cAMP to

PKA-R activates PKA-C. However, there are different proposals on the molecular events

that follow activation.

For decades, PKA-C is thought to dissociate from PKA-R upon cAMP binding. Freed

PKA-C molecules then move to phosphorylate their substrates. However, two recent

studies propose an alternative model, in which physiological concentrations of cAMP can

activate PKA-C but do not result in its dissociation from PKA-R (3, 4). Testing these two

models will not only elucidate the biophysical mechanism of PKA activation, but also have

distinct implications in how PKA may achieve its specificity, which is thought to rely on

spatial compartmentalization (5).

We have previously found that the majority of type IIβ PKA, as defined by PKA-R, is

anchored to microtubules in the dendritic shaft of hippocampal CA1 pyramidal neurons

where PKA-RIIβ is bound to the abundant microtubule associated protein MAP-2 (6). Upon

activation of the β-adrenergic receptor with norepinephrine, a fraction of PKA-C

dissociated from PKA-RIIβ (7). The freed PKA-C redistributed into dendritic spines,

whereas PKA-RIIβ remained anchored at the dendritic shaft (7, 8). These results are

consistent with the classical PKA activation model. However, recent studies suggest that

PKA-Rs other than PKA-RIIβ may be the more abundant isoforms in CA1 neurons (9, 10). It

remains unknown whether these PKA isoforms dissociate upon physiologically relevant

stimulations.

Here, we examined whether PKA-C dissociates from the major PKA-R isoforms,

PKA-RIβ and PKA-RIIα, in CA1 neurons. The rescue of function following knockdown of

PKA-C was examined using wildtype dissociable PKA versus a non-dissociable PKA variant

in which PKA-C is covalently linked to PKA-R. The result supports the classical model of

PKA activation via dissociation.

Results

Recent studies have suggested that PKA-RIIα and PKA-RIβ may be the prevalent

isoforms in hippocampal CA1 neurons. Therefore, we co-expressed EGFP tagged PKA-C

(PKA-C-EGFP) and a cytosolic marker (mCherry) with either PKA-RIIα or PKA-RIβ in CA1

neurons of organotypic hippocampal slice cultures (Figs. 1A and 1B, upper left panels).

PKA-C-EGFP exhibited a resting distribution that was dependent on the co-expressed PKA-

R: when co-expressed with PKA-RIIα, PKA-C was enriched in dendritic shafts; when co-

expressed with PKA-Riβ, PKA-C was more evenly distributed (quantified using the spine

enrichment indexes, or SEI, see Methods) (Fig. 1C). This distribution largely mimicked that

of the corresponding PKA-R (Figs. 1A-1C).

Notably, upon application of norepinephrine (10 μM), PKA-C of both subtypes

translocated to dendritic spines, but the subcellular localization of PKA-Rs remained

unchanged (Figs. 1D & 1E). This differential re-distribution between PKA-C and PKA-R was

more prominent following activation with forskolin (25 μM) and IBMX (50 μM). These

results can only be explained if at least a fraction of PKA-C dissociated from both PKA-RIIα

and PKA-RIβ when a physiological stimulant was used. Together with our earlier results

regarding PKA-C/PKA-RIIβ, we conclude that PKA-C dissociates from PKA-R with

physiologically relevant stimuli.

Next, we asked whether PKA regulation of neuronal function is dependent on the

dissociation of PKA-C from PKA-R. A key experiment supporting the non-dissociating PKA

activation model was that PKA regulation of cell growth could be sustained by a construct

in which PKA-C was fused to PKA-RIIα in one polypeptide chain via a flexible linker (named

R-C; Fig. 1F). We therefore asked whether this R-C construct could support PKA regulation

of neuronal function. When R-C-EGFP was expressed in CA1 neurons, this construct

exhibited a distribution highly similar to that of RIIα at the resting state (Fig. 1G and 1H).

The tendency of this construct to translocate to the spine was largely diminished compared

to PKA-C-EGFP co-expressed with wildtype, unlinked PKA-RIIα (Fig. 1I).

To determine the function of R-C, a previously established shRNA construct was

used to selectively knock down PKA-Cα in CA1 neurons in cultured hippocampal slices (7).

Given that PKA activation is required for the late phase of long-term potentiation (L-LTP)

(11), we examined the structural LTP of individual dendritic spines of CA1 neurons elicited

by focal two-photon glutamate uncaging (Figs. 2A–2C) (12). The shRNA knockdown of

PKA-C resulted in attenuated LTP at around 90 min after induction (Figs. 2A–2C). This

attenuation was not observed when a control shRNA against LacZ was expressed (Fig. 2C).

The attenuated structural LTP was rescued by co-expression of shRNA-resistant wild-type

PKA-C-EGFP together with PKA-RIIα (Figs. 2A–2C). However, the R-C construct in which

PKA-C was also resistant to shRNA knockdown but could not leave PKA-RIIα failed to

rescue the phenotype.

PKA activity has also been shown to regulate synaptic transmission. We therefore

examined evoked AMPA and NMDA receptor (AMPAR and NMDAR, respectively) currents

in paired, transfected and adjacent untransfected CA1 neurons in cultured hippocampal

slices. As shown previously (7), neurons expressing the shRNA construct against PKA-C

exhibited significantly lower AMPAR currents (Fig. 2D), but not NMDAR currents (Fig. 2E).

As a result, the AMPAR/NMDAR current ratio was also reduced (Fig. 2F). The reduced

AMPAR currents and reduced AMAPR/NMDAR current ratios were rescued by co-

expression of shRNA-resistant, wild-type dissociable PKA-C-EGFP and PKA-RIIα (Figs. 2D–

2F). However, the non-dissociable R-C construct failed to rescue the phenotype. Taken

together, the R-C construct did not support normal PKA-dependent synaptic function.

Discussion

The results indicate that at least a fraction of PKA-C molecules dissociate from all

tested PKA-R isoforms when activated by physiological stimuli. Given that PKA activity

increases by two orders of magnitude when dissociated from PKA-R (13), even a small

fraction of PKA-C dissociation will result in a marked increase of PKA kinase activity.

Notably, our results corroborate the recent results in HEK cells that used biochemical

measurements to show that PKA-C and PKA-R indeed dissociate in intact cells (14).

Furthermore, the PKA-C that is covalently linked to PKA-RIIα cannot functionally replace

wildtype PKA for normal neuronal transmission or plasticity. Although the R-C construct

has been shown to functionally replace endogenous PKA in terms of supporting the growth

of a heterologous cell line (4), our results indicate that it cannot support all necessary PKA

functions. Overall, we conclude that PKA-C dissociation from PKA-R is essential for PKA

regulation of neuronal function. Additionally, PKA specificity is mediated by spatial

compartmentalization (5). This is likely mediated by mechanism downstream of PKA

dissociation, such as membrane tethering of freed PKA-C or its buffering by extra PKA-Rs

(7, 14).

This study also demonstrates that PKA is essential for long-term structural LTP of

individual spines. PKA has been shown to facilitate the induction of LTP (i.e.,

metaplasticity) (15), and is required for the maintenance of L-LTP (11), as assayed using

electrophysiological recording of postsynaptic currents. However, it has been suggested

that the structural and synaptic current changes may not be causal (16). Our results fill the

gap to show that PKA is also required for the late phase of structural LTP.

Materials and Methods

Plasmid constructs

Constructs were made using standard mutagenesis and subcloning methods. All previously

unpublished constructs and their sequences will be submitted to Addgene. In the R-C

construct, mouse PKA-RIIα and PKA-Cα were fused via a linker with residues

WDPGSGSLEAGCKNFFPRSFTSCGSLEGGSAAA that were previously used (4).

Organotypic hippocampal slice cultures and transfections

Cultured rat hippocampal slices were prepared from P6 – P8 (typically P7) pups, as

described previously (6, 17). Animal experiments were performed in accordance with the

Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, and

were approved by the Institutional Animal Care and Use Committee (IACUC) of the Oregon

Health & Science University (#IP00002274). cDNA constructs were transfected after 1.5–3

weeks *in vitro* via the biolistic gene transfer method using the Helios gene gun and 1.6 μm

gold beads (Fig. 1) or, where long-term expression (∼ 1 week) was required, with single-

cell electroporation (Fig. 2) (18).

Two-photon imaging and two-photon glutamate uncaging

A custom built two-photon microscope based on an Olympus BW51WI microscope body

was used. Laser beams from two different Ti:Sapphire lasers (Maitai, Newport) were

aligned to allow for simultaneous two-photon excitation and photoactivation. Laser

intensities were controlled by Pockels cells (Conoptics). Imaging and photoactivation were

controlled by ScanImage (Vidrio Tech) (19). Slices were perfused in gassed artificial

cerebral spinal fluid (ACSF) containing 4 mM Ca, 4 mM Mg, and 0.5 μM tetrodotoxin (TTX)

during imaging. mEGFP fluorescence (green) was unmixed from that of the cytosolic

marker (mCherry or DsRed Express) using a dichroic (Chroma 565DCXR) and band-pass

filters (Chroma HQ510/70 for green and Semrock FF01-630/92 for red).

For single-spine structural LTP experiments, 2.25 mM MNI-caged-L-glutamate

(Tocris) was added to ACSF containing 4 mM calcium, 0.05 mM magnesium, 1 µM TTX and

4 µM 2-chloroadenosine, as previously described (20). To trigger structural plasticity, 30

pulses of 4-ms 16-mW (at back focal plane) 720-nm laser light were delivered to the spine

head at 0.5 Hz.

Image analysis was performed using custom software written in MATLAB called

SI_View (https://github.com/HZhongLab/SI_View) (21). Using the software, regions of

interest (ROIs) were manually drawn to isolate spines or their immediately adjacent

dendritic shaft. Only the spines well isolated from the dendrite laterally throughout the

entire experiments were included. Spine enrichment index was calculated as:

$$SEI = \log_2[(F_{green}/F_{red})_{spine}/(F_{green}/F_{red})_{shaft}]$$

in which F is the average fluorescence intensity in an ROI.

Electrophysiology

Whole-cell voltage-clamp recordings were performed using a MultiClamp 700B amplifier

(Molecular Devices). Electrophysiological signals were filtered at 2 kHz and digitized and

acquired at 20 kHz using custom software written in MATLAB. Slices were perfused with

artificial cerebrospinal fluid containing 4 mM Ca and 4 mM Mg. The internal solution

contained (in mM) 132 Cs-gluconate, 10 HEPES, 10 Na-phosphocreatine, 4 MgCl2, 4 Na2-

ATP, 0.4 Na-GTP, 3 Na-ascorbate, 3 QX314, and 0.2 EGTA with an osmolarity of 295

mOsmol/kg. The junction potential was calculated to be -17 mV using a built-in function in

the Clampfit software (Molecular Devices). Several less abundant anions (phosphocreatine,

ATP, GTP and ascorbate) were omitted in the calculation due to lack of data in the program.

The Cl reversal potential was -75 mV.

To reduce recurrent activities, cultured hippocampal slices were cut on both sides of

CA1 and 4 µM 2-chloroadenosine (Sigma) was present in all recording experiments. 10 µM

GABAzine (SR 95531, Tocris) was also included to suppress GABA currents. For electrical

stimulation, a bipolar, θ-glass stimulating electrode (Warner Instruments) was positioned

in the stratum radiatum 100–150 µm lateral to the recorded neuron. For all recordings, a

transfected neuron and an untransfected neuron located within 50 µm of each other were

sequentially recorded without repositioning the stimulation electrode. Measurements were

carried out on averaged traces from approximately 20 trials under each condition. For

AMPAR currents, the cells were held at -60 mV (before correcting for the junction

potential) and the current was measured as the baseline-subtracted peak current within a

window of 2–50 ms after electric stimulation. For NMDAR currents, the average currents at

140 to 160 ms after stimulation were used when the cells were held at +55 mV (before

correcting for the junction potential)

Data analysis, presentation, and statistics

Quantification and statistical tests were performed using custom software written in

MATLAB. Averaged data are presented as mean ± s.e.m., unless noted otherwise. p values

were obtained from one-way ANOVA tests, unless noted otherwise. In all figures, *: $p \leq 0.05$

and is considered statistically significant after Bonferroni correction for multiple tests, **: p

≤ 0.01, and ***: p ≤ 0.001.

Acknowledgements

We thank all members of the Mao and Zhong laboratories at the Vollum Institute for

constructive discussions. We thank Drs John Williams and Michael Muniak for critical

comments and edits on the manuscript. This work was supported by two NIH BRAIN

Initiative awards to H.Z. (U01NS094247 and R01NS127013).

Author contributions

H.Z. conceived the project. W.X. and H.Z. designed the experiments. W.X. and H.Z.

performed the experiments and data analyses. M.Q. produced critical reagents used in the

experiments. H.Z. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1.   S. H. Francis, J. D. Corbin, Structure and Function of Cyclic Nucleotide-Dependent

Protein Kinases. *Annu. Rev. Physiol.* (1994)

https://doi.org/10.1146/annurev.ph.56.030194.001321.

2.   D. A. Johnson, P. Akamine, E. Radzio-Andzelm, Madhusudan, S. S. Taylor, Dynamics of

cAMP-dependent protein kinase. *Chem. Rev.* (2001)

https://doi.org/10.1021/cr000226k.

3.   F. D. Smith, *et al.*, Intrinsic disorder within an AKAP-protein kinase A complex guides

local substrate phosphorylation. *Elife* **2013** (2013).

4.   F. D. Smith, *et al.*, Local protein kinase A action proceeds through intact holoenzymes.

Science (80-.). **356** (2017).

5.   W. Wong, J. D. Scott, AKAP signalling complexes: Focal points in space and time. *Nat.*

Rev. Mol. Cell Biol. (2004) https://doi.org/10.1038/nrm1527.

6.   H. Zhong, *et al.*, Subcellular Dynamics of Type II PKA in Neurons. *Neuron* **62**, 363–374

(2009).

7.   S. E. Tillo, *et al.*, Liberated PKA Catalytic Subunits Associate with the Membrane via

Myristoylation to Preferentially Phosphorylate Membrane Substrates. *Cell Rep.* **19**,

617–629 (2017).

8.   W. H. Xiong, M. Qin, H. Zhong, Myristoylation alone is sufficient for PKA catalytic

subunits to associate with the plasma membrane to regulate neuronal functions.

Proc. Natl. Acad. Sci. U. S. A. **118** (2021).

9.   M. Weisenhaus, *et al.*, Mutations in AKAP5 disrupt dendritic signaling complexes and

lead to electrophysiological and behavioral phenotypes in mice. *PLoS One* **5** (2010).

10.   R. Ilouz, *et al.*, Isoform-specific subcellular localization and function of protein kinase

a identified by mosaic imaging of mouse brain. *Elife* **6** (2017).

11.   T. Abel, *et al.*, Genetic demonstration of a role for PKA in the late phase of LTP and in

hippocampus-based long-term memory. *Cell* **88**, 615–626 (1997).

12.   M. Matsuzaki, N. Honkura, G. C. R. Ellis-Davies, H. Kasai, Structural basis of long-term

potentiation in single dendritic spines. *Nature* **429**, 761–766 (2004).

13.   E. Miyamoto, J. F. Kuo, P. Greengard, Cyclic nucleotide-dependent protein kinases. 3.

Purification and properties of adenosine 3',5'-monophosphate-dependent protein

kinase from bovine brain. *J. Biol. Chem.* **244** (1969).

14.   R. Walker-Gray, F. Stengel, M. G. Gold, Mechanisms for restraining cAMP-dependent

protein kinase revealed by subunit quantitation and cross-linking approaches. *Proc.*

Natl. Acad. Sci. U. S. A. **114** (2017).

15.   M. J. Thomas, T. D. Moody, M. Makhinson, T. J. O'Dell, Activity-dependent β-

adrenergic modulation of low frequency stimulation induced LTP in the hippocampal

CA1 region. *Neuron* **17**, 475–482 (1996).

16.   C. D. Kopec, B. Li, W. Wei, J. Boehm, R. Malinow, Glutamate receptor exocytosis and

spine enlargement during chemically induced long-term potentiation. *J. Neurosci.* **26**

263 (2006).

264 17. L. Stoppini, P. A. Buchs, D. Muller, A simple method for organotypic cultures of

265 nervous tissue. *J. Neurosci. Methods* (1991) https:/doi.org/10.1016/0165-

266 0270(91)90128-M.

267 18. N. Otmakhov, J. Lisman, Measuring CaMKII concentration in dendritic spines. *J.*

268 *Neurosci. Methods* (2012) https:/doi.org/10.1016/j.jneumeth.2011.09.022.

269 19. T. A. Pologruto, B. L. Sabatini, K. Svoboda, ScanImage: Flexible software for operating

270 laser scanning microscopes. *Biomed. Eng. Online* **2**, 13 (2003).

271 20. C. D. Harvey, R. Yasuda, H. Zhong, K. Svoboda, The spread of Ras activity triggered by

272 activation of a single dendritic spine. *Science (80-.).* **321**, 136–140 (2008).

273 21. L. Ma, *et al.*, Locomotion activates PKA through dopamine and adenosine in striatal

274 neurons. *Nature* **611**, 762–768 (2022).

 Figure legends

Fig. 1. PKA-C but not PKA-R or a non-dissociable R-C redistribute to spines upon

activation. (*A*, *B*) Representative two-photon images of PKA-C-EGFP co-expressed with

PKA-RIIα or PKA-RIβ at rest, or in the presence of norepinephrine (NE) or forskolin and

IBMX (F + I). mCherry (magenta) was co-expressed to reveal the neuronal morphology. (*C*–

E) Quantification and comparison of the spine enrichment index at the resting state (C) and

upon activation (*D* & *E*). As in panel *C* from left to right, n (spines/neurons) = 53/11, 34/7,

33/6, and 36/7. (*F*) Schematic of wildtype PKA versus R-C. In both cases PKA-C was tagged

by mEGFP. (*G*–*I*) Representative images (*G*), quantifications of resting distribution (*H*), and

the distribution change upon stimulation by forskolin and IBMX (*I*) of R-C compared to

PKA-RIIα-EGFP and co-expressed PKA-C-EGFP/PKA-RIIα. n (spines/neurons) = 48/10.

Fig. 2. PKA regulation of synaptic plasticity and synaptic transmission cannot be sustained

using a non-dissociable PKA variant. (*A*–*C*) Representative image (*A*), time course (*B*), and

the degree of potentiation (*C*) at the indicated timepoints in panel *B* of single-spine LTP

experiments as triggered by focal glutamate uncaging at the marked spines (gray dot). In

panel *B*, both stimulated spines (solid circles) and non-stimulated control spines (open

circles) are shown. As in panel *C* from left to right, n (spines, each from a different neuron)

= 8, 7, 17, 11, 9. (*D*–*F*) Representative traces (red) normalized to the paired control (blue)

(insets) and scatter plots of paired AMPA (*D*) and NMDA (*E*) receptor currents and

AMPA/NMDA receptor current ratios (*F*) from neighboring untransfected CA1 neurons

paired with those transfected with shRNA against PKA-C and the indicated shRNA-resistant

rescue constructs. Statistical p values were obtained using a sign test (MATLAB). From left

to right, n (neuron pairs) = 13, 11, 11, and 15.

Xiong et al._Figure 1

Xiong et al._Figure 2