Multi-Generation Ecosystem Selection Of Rhizosphere Microbial Communities Associated With

Plant Genotype and Biomass In *Arabidopsis thaliana*

Nachiket Shankar[1*], Prateek Shetty[2], Tatiana C Melo[1], Rick Kesseli[1]

[1]Department of Biology, University of Massachusetts, Boston, MA 02125,

[2]Institute of Plant Biology, Hungarian Academy of Sciences, Biological Research Centre, 6726

Szeged, Hungary

7   **Abstract**

8   The role of the microbiome in shaping the host phenotype has emerged as a critical area of

investigation, with implications in ecology, evolution, and host health. The complex and dynamic

interactions involving plants and their diverse rhizosphere microbial communities are influenced

by a multitude of factors, including but not limited to soil type, environment, and plant genotype.

Understanding the impact of these factors on microbial community assembly is key to yielding

host-specific and robust benefits for plants, yet remains challenging. Here we ran an artificial

ecosystem selection experiment, over eight generations, in *Arabidopsis thaliana* L*er* and Cvi to

select soil microbiomes associated with higher or lower biomass of the host. This resulted in

divergent microbial communities, shaped by a complex interplay between random environmental

variations, plant genotypes, and biomass selection pressures. In the initial phases of the

experiment, the genotype and the biomass selection treatment have modest but significant impacts.

Over time, the plant genotype and biomass treatments gain more influence, explaining ~40% of

the variation in the microbial community composition. Furthermore, a genotype-specific

association of a plant growth-promoting rhizobacterial taxa, *Labraceae* with L*er* and *Rhizobiaceae*

with Cvi, is observed under selection for high biomass.

Keywords: plant-associated microbiome; microbial diversity; soil health; sustainable agriculture

Introduction

The conventional understanding of the host phenotype involves genetics and environment shaping observable traits. Yet, the last two decades have underscored the microbiome's significance in shaping the host phenotype, driven by extensive research on its role in ecology, evolution, and host health. The plant microbiome represents a rich source of functional diversity that is not encoded within the host genome. The interactions between the plant host and its microbiome are dynamic and reciprocal, with the plant shaping its immediate environment by exuding specific metabolites, thereby promoting the growth of specific microbial taxa, while the microbiome in turn influences plant health and growth [1]. A multitude of biotic and abiotic factors influence the dynamic nature of the microbiome, including but not limited to plant root exudates [2–4], soil type [5–9], environment [10], and various aspects of the plant, including species [11], genotype [4,12–15] and developmental stage [16,17].

Previous studies in *Arabidopsis thaliana* have shown that the microbial diversity of the soil reduced with proximity to the endophytic compartment, and microbes found in the bulk soil were not enriched in the endophytic compartment and vice-versa [18–20]. Furthermore, the microbiome of greenhouse-grown plants was similar to that of the field-grown plants, in the different plant compartments – rhizosphere, woody stem, and endophytic compartment, indicating an active role of the plant host in creating and maintaining an environment where certain microbes have improved fitness [21]. The soil environment was a stronger predictor of rhizosphere microbiome structure, though plant genotype had a weak influence in some studies [20,22,23]. A follow-up study confirmed that variation in microbiome communities depends more on the environment than on different *Arabidopsis* varieties and sister species. Though there were some differences between

47   different host species, the core microbiome interacting with the plant host at the root endophytic

48   zone was largely consistent and reproducible [24].

49   Unfortunately, most prior studies were conducted in a single plant growth cycle, precluding

an analysis of the temporal stability of the plant-microbiome association. Observations from plant-

soil feedback studies highlight that plant-microbiome interactions can be altered by successive

growth cycles of a given plant in the same soil [25,26]. Prior studies have employed plant-mediated

selection on the soil ecosystem over multiple generations, and resulted in consistent effects on

plant characteristics such as biomass, flowering time, and germination [27,28]. By selecting for

soils where hosts show the desired phenotype, such as high biomass or altered flowering time, it

is possible to enrich microbes that modulate host traits. Additionally, studies showed the plant's

response to abiotic stress, such as drought and salt stress, could be influenced in two ways. First,

through generational selection, where plants with desirable stress responses were chosen over

multiple generations [29–31]. Second, by introducing beneficial microbes that were associated

with plants and had previously lived in similar drought or salt conditions [32]. Mueller et al., [29]

used generational selection to create microbial communities that could improve plant seed

production by up to 205%. Soil inoculum obtained from drought-exposed soils improved wheat

biomass under drought conditions [32]. Older studies, such as Swenson et al. [28], also have shown

similar results under optimal growing conditions without necessarily exploring the microbiome. A

recent study conducted on the rhizosphere microbiome of wild and domesticated tomato plants

over multiple generations demonstrated an escalating influence of host genotype on the

microbiome community [33]. A separate study on the phyllosphere microbiome of various tomato

genotypes indicated a declining impact of host genotype across successive generations [34].

Despite several earlier efforts, the development of a robust and host-specific microbial

community with long-lasting beneficial effects on the plant host remains a challenge. Using an

artificial ecosystem selection experiment, we sought to advance our understanding of how plant

genotype, environment, and biomass selection treatment (henceforth referred to as biomass

treatment) impact the assembly of host microbial communities. Treatments selecting for the

microbial communities of high and low biomass plants did affect the growth of plants in

subsequent generations, though unknown environmental factors had substantial effects throughout

the experiment. In the first few generations of the experiment, the genotype and biomass treatments

played a modest role in shaping the microbial community. Over time, the plant genotype and

biomass treatments had increasing influence, explaining ~40% of the difference in the microbial

community composition. Moreover, we observe an enrichment of known plant growth-promoting

rhizobacteria (PGPR) in the high biomass treatment, indicating potentially beneficial host-specific

interactions.

82   **Material and Methods**

Multi-generation selection of soil ecosystem

Arabidopsis thaliana Cvi and L*er* accessions were cultivated using custom-made

"rhizotubes," (Stuart Morey, Univ. of Massachusetts, Boston, unpublished). The tubes were

equipped with a black polyethylene sleeve, which effectively blocked light penetration. This

design also facilitated the convenient removal of the plant from the pot, granting full access to

the root system. (Supplementary Figure S1).

The potting soil used was PRO-MIX PGX, a commercial mixture comprising 80-90%

sphagnum peat moss and small quantities of perlite. It was autoclaved twice for 40 minutes with

a 48-hour interval between each sterilization. The sterile potting soil was then sifted through a 3

mm sieve and combined with field soil in a 6:1 ratio for the first generation. The field soil

obtained from the Center for Agricultural Research in Waltham, Massachusetts, was composed

of 44% sand, 49% silt, and 7% clay representative of agricultural and grassland ecosystems.

The mixture was homogenized using a custom-made cement mixer and attached with

sterile bins to avoid cross-contamination. Distilled water was gradually added to achieve a final

ratio of 1:2 (water to soil). The soil was incubated at room temperature for two days following

inoculation before planting the seeds. All seeds used in the experiment were obtained from a

single-parent plant. Before planting, the seeds were treated with a solution of 50% bleach vol/vol

and a drop of tween 20 for 10 minutes and rinsed ten times with sterile distilled water. We placed

3-4 seeds in the center of each pot and kept only one seedling per pot after emergence. All plants

were grown at 22 °C day/ 18 °C night with a 12/12 hour day-night cycle in a controlled growth

chamber. Relative humidity ranged from 35-60% and the light intensity was 96 µE. Fertilizer

was not applied to plants.

In the first generation of the experiment, 100 plants of each accession (Cvi and Ler) were

grown separately in individual pots. The plants were arranged in a randomized block design to

avoid batch effects. All plants were harvested 35 days after germination. The above-ground

portion of the plant, the rhizosphere, and the bulk soil from each rhizotube were separated,

ensuring there was no cross-contamination (Supplementary Figure S1). The above-ground part of

the plant, stem, and leaves were dried at 70 °C for 4 days. All the plants were weighed

individually on a closed weighing scale accurate to 1 mg. The root-soil complex (comprising the

rhizosphere and the endosphere, henceforth referred to as the rhizosphere) was obtained by

shaking the excess soil off the root and placing the root and remaining attached soil in a sterile 5

ml tube. Bulk soil samples were moved to a sterile Ziplock bag. Tubes and bags were

immediately transferred to dry ice and then stored at -80 °C for DNA analysis.

All subsequent generations were inoculated using the previously collected bulk soil from

the top five and bottom five plants by above-ground biomass. This resulted in two different

treatments for each genotype: high biomass L*er*, low biomass L*er*, high biomass Cvi, and low

biomass Cvi. This process is repeated for the remaining seven generations. Starting with the third

generation, plants were grown in uninoculated sterile potting soil, as a control for random

environmental variation (REV). In generation 6 all plants died for unknown reasons. The

experiment was then restarted using soil from generation 5. Soil sourced from the top 6-10 plants

and the bottom 6-10 plants based on above-ground biomass was used as inoculum to replant

generation 6.

DNA extraction and 16S rRNA amplicon library prep

Microbial DNA was isolated from the frozen rhizosphere samples using the Machery-

Nagel Nucleospin Soil DNA extraction kit (MACHEREY-NAGEL Inc., Allentown, PA, USA).

Approximately 0.1g of rhizosphere soil sample was used for DNA isolation. All samples were

diluted to 5 ng ul-1 with PCR-grade water. 16S rRNA gene was amplified from the isolated

DNA samples in triplicate in 96-well PCR plates. The PCR primers used were for the 16S rRNA

V4 region, 515F (5'-GTGYCAGCMGCCGCGGTAA-3') (ref. 35) and 806R (5'-

GGACTACNVGGGTWTCTAAT-3') (ref. 36) for downstream paired-end Illumina (Illumina,

Inc., San Diego, CA, USA) barcoded sequencing [37]. The PCR cycling conditions were as

follows: 94 $^{\circ}$C for 3 min; 25 cycles of 94 $^{\circ}$C for 45 s, 50 $^{\circ}$C for 60 s, 72 $^{\circ}$C for 90 s, and final

elongation at 72 $^{\circ}$C for 10 min. The triplicate amplified samples were pooled and then purified

and normalized using the SequalPrep™ Normalization Plate Kit (Invitrogen Corporation,

Carlsbad, Canada). Finally, multiplexed paired-end sequencing was carried out in the Illumina

MiSeq (SY-410-1003) platform using earth microbiome project (EMP) primers [38].

Sequence Data Analysis & Statistics

The paired-end sequences obtained from the Illumina MiSeq were demultiplexed with

QIIME 2 and converted into individual sequence fastq files for each sample [37,39]. The rest of

the sequence processing was carried out in R (Version 4.2.2) using the DADA2 package

(Version 1.24.0) (ref. 40). The reads were processed in R using the following command in

DADA2; `filterAndTrim(fnFs, filtFs, fnRs, filtRs, truncLen=c(145,145), minLen=50, maxN=0,

maxEE=c(2,2), truncQ=2, rm.phix=TRUE, compress=TRUE, multithread=TRUE). `

De novo OTU (Operational Taxonomic Units) picking was performed using DADA2 (37) which

resolves amplicon sequence variants (ASVs) to single nucleotide differences. Chimeras were

removed. The taxonomy was assigned using a 100% cut-off for species-level identification with

the (2022) GTDB 16S rRNA reference database [41]. The phylogenetic tree was constructed

using IQTree [42], with 1000 ultrafast bootstraps and Modelfinder to identify the best model. In

this study, the best-fit model was SYM+R10 chosen according to BIC. The data comprising the

OTU table, phylogenetic tree, taxonomy table, and sample meta-data were then parsed through R

studio using the phyloseq package [43]. Here, it was processed to remove: OTUs identified as

mitochondrial or chloroplast, which had less than five reads across all samples, and samples that

had fewer than 8000 reads. The packages phytools, phyloseq, microbiome, and ggplot2 were

used for further data analysis. All sequence data and metadata can be downloaded from

PRJNA1010286.

Diversity Analyses

Alpha diversity was computed using four metrics, Observed, Shannon, Chao1, and Simpson diversity indices. Beta diversity was calculated using weighted and unweighted UniFrac, and subsequently Principle Coordinate Analysis (PCoA) plots were constructed using the first two axes that explain the most variance. Linear mixed effect models using, *lmerTest* package in R were used to determine changes in alpha and beta diversity over time (lmer (DiveristyMetric, Genotype*Generation*Biomass Treatment, (1|Generation)), with generation as a fixed and random effect. Pairwise comparisons were determined with *emmeans* and adjusted p-value obtained with Tukey.

Adonis

PERMANOVA was carried out for each generation using the adonis2 test. Both weighted and unweighted UniFrac distance matrices were used to account for the abundance, presence/absence, and phylogeny of the microbiome. The following model was used `adonis2(distance.matrix~Biomass Treatment+Genotype, data = meta, permutations=999)`.

Neutral Model

To determine the impact of neutral processes like drift and dispersal or by deterministic selective forces such as plant genotype and biomass treatments on microbial community assembly, we carried out the analysis as described in Burns et al. [44], which fits the data to the neutral model for Prokaryotes from Sloan et al. [45]. The following design was used, sncm.fit (spp=generation(n)_Biomass Treatment, pool = generation(n), stats=T), where n is the nth generation and 'Biomass Treatment' is high or low biomass samples.

Differential Abundance

There are several challenges in estimating differentially abundant (DA) taxa in

microbiome data, these include high variability in abundance, zero-inflated data, and the

compositional nature of the data. Nearing et al. [46] compared several methods across 38

datasets and found that different methods often identified different sets of DA taxa. The DA taxa

are estimated using ANCOMBC2 [47], a conservative and robust approach. ANCOMBC2

incorporates bias correction, effectively addressing sampling-specific and sequencing biases

present in the data. This feature ensures that the analysis is not skewed by any systematic errors

introduced during the sampling or sequencing process. It also conducts a sensitivity analysis for

the pseudo-count addition to assess the impact of different pseudo-count values on zero counts

for each taxon. The analysis was run on L*er* and Cvi samples, from OTU to Family taxonomic

levels with (fixed effect = (Biomass Treatment + Genotype + Generation), group = Biomass

Treatment).

198   **Results**

Plant Biomass

The ecosystem selection experiment was carried out for the trait plant biomass on two

different accessions of *Arabidopsis thaliana,* L*er* and Cvi for eight generations. The two selected

treatments were high and low above-ground biomass. The absolute values for the mean biomass

of plants (n=50) from each genotype and treatment changed substantially through the course of

eight generations of the experiment. Despite the stochastic fluctuations in biomass caused by

random environmental variation (REV), the high biomass selected lines were always the same or

greater than those of the low selected biomass for both genotypes (Figures 1a and 1b). Due to the

significant drop in biomass between generations 1 and 2, a sterile potting soil uninoculated

control was planted in all subsequent generations to serve as a reference for REV. The

uninoculated control shows that the drastic drop in biomass after generations 1 and 2, was not a

result of the inoculated soil but other subtle but crucial environmental factors that could not be

held constant. Above-ground biomass for each generation in terms of deviations from the mean

was plotted to give a clearer representation of the difference in phenotype seen in every

generation (Figures 2a and 2b). Significant differences between the low and high biomass

treatments become apparent from generation 4 onwards. Plants of all treatments in generation 6

died resulting in a sharp dip, which necessitated repeating that generation (described in the

Materials and Methods).

Microbial community Composition

After preprocessing the 16S rRNA metagenome amplicon reads in DADA 2, a total of

1,897,732 reads, with an average of 12822 reads per sample were obtained. Singletons and

chimeras were removed during pre-processing. The most well-represented phyla were

Proteobacteria (36.2%), *Bacteroidetes* (10.3%), *Planctomycetes* (7.8%), and *Actinobacteria*

(7.6%) (Supplementary Figure S2). The most well-represented Classes identified in the dataset

were *Alphaproteobacteria* (24.7%), *Gammaproteobacteria* (11.4%), *Bacteroidia (10.1%)*, and

Verrucomicrobiae (6.2%) (Figure 3). The relative abundance is typical of microbiome data,

heavily weighted to a few abundant groups with a long tail of rare taxa.

Alpha Diversity

We used two metrics to assess alpha diversity (Figure 4). Observed ($F_{7,160}$ =333.054,

P<0.001) estimates species richness, and showed a statistically significant decrease in the

number of OTUs (1477 to ~580) and richness across generations. The Shannon diversity index

($F_{7,160}$ =82.23, P<0.001), which is more sensitive to the difference in abundance, also showed

this steady decrease across generations. A negative exponential function was fitted to the

Observed metric ($R2$=0.81, P<0.01) and the Shannon metric ($R2$=0.58, P<0.01). The Observed

metric displayed a better fit, indicating increased stability of the microbiome in later generations.

In addition, a linear mixed-effect model shows significant interactions between

genotype:biomass treatment (F = 8.71, P < 0.01), genotype:generation (F =3, P < 0.01), and

genotype:biomass treatment:generation (F = 3, P < 0.01), for Shannon and (Supplementary

Table S5). No significant interaction terms were found for Observed, which does not consider

the abundance of OTUs in its calculations.

Beta Diversity

Principal coordinate analyses (PCoA) of the weighted UniFrac distances were plotted

with the first two axes that captured the most variance in the data, from 73% in generation 1 to

greater than 90% in generation 8 (Figure 5). UniFrac distance considers the evolutionary

relationships between taxa, which makes it more biologically meaningful than other distance

metrics that do not consider the evolutionary history. Weighted UniFrac is more sensitive to

differences in the abundance of taxa. The microbiomes associated with the treatments and

genotypes are very similar in generation 1, but begin to diverge by generation 2. By generation 3,

the microbiomes of the L*er* and Cvi genotypes have diverged and by generation 5 the high and

low biomass treatments appear to have further distinguished the microbial communities. The

divergence of the microbial community in generations 4 and 5 caused by the high/low biomass

treatments also aligns with the onset of notable differences in the above-ground plant biomass

within these generations.

The complex interplay between genotype and biomass treatments was modeled using a

PERMANOVA test with both weighted and unweighted UniFrac distances to account for

differences in abundance and presence-absence of OTUs respectively. The resulting R^2 values

for genotype and biomass treatment and residuals as a proxy for REVs were plotted (Figure 6a

and 6b). For both weighted and unweighted UniFrac, the variance in microbial community

explained by both genotype and biomass treatment increased significantly over the eight

generations. Despite observing a decrease in the variance accounted for by the residuals, which

could potentially signify a decline in the influence of REVs, they accounted for ~50 percent of

the variability within the microbial community in generation 8. For all metrics, weighted

UniFrac exhibits more fluctuations compared to the unweighted UniFrac. Weighted UniFrac

analyses found that genotype ($F_{1,39}$=1.55, P<0.05) and biomass treatment ($F_{1,39}$=1.33, P<0.001)

explained 5.7% and 13.7%, respectively, of the dissimilarity between microbiomes in generation

1. By generation 8, genotype ($F_{1,14}$=5.14, P<0.05) and biomass treatment ($F_{1,14}$=5.48, P<0.001)

explain 24% and 22%, respectively, of the dissimilarity between microbiomes (Supplementary

Tables S6 and S7). For unweighted UniFrac, the influence of genotype on the microbial

community increased from 3.3% in generation 1 ($F_{1,39}$=1.33, P<0.01) to 26% in generation 8

($F_{1,39}$=5.13, P<0.001), and for biomass increased from 3.8% in generation 1 ($F_{1,14}$=1.55, P<.001)

to 13% in generation 8 ($F_{1,14}$=2.58, P<0.05) (Supplementary Tables S8 and S9).

Sloan Neutral Model

Many forces can alter microbial community structure and dynamics and in our study

these can be divided into selective forces such as the plant genotype being colonized and the

biomass selection process, or neutral, stochastic forces innate in any experiment and

environment. The Sloan neutral model was fit to the data to assess the importance of selective

versus neutral drivers of change in the experiment. If neutral processes were the driving force in

the microbial community assembly, the null hypothesis would be that all generations of the

experiment would fit the model equally well. The model fit to neutrality represented by the R^2

value decreases from 0.795 in generation 1 to 0.59 in generation 8 demonstrating the increasing

importance of selective drivers through the course of the experiment (Figure 7).

Differential Abundance

Differentially abundant (DA) taxa that distinguished between the high and low biomass

treatments were determined using ANCOMBC2 at the Family level. Taxa lacking classification

at the Family level were denoted by the subsequent identifiable taxonomic tier. The OTUs that

were present in higher (red) and lower (blue) abundance in the high biomass treatment are

identified (Figure 8a and 8b). Among the DA taxa in this study, several are known to benefit

plants. The Class *Bacilli* promote plant growth [48]. The Class *Gemmatimonadetes* are

positively associated with vegetation restoration, plant richness, and soil nutrients [49].

Cytophagacea are chemoorganotrophs and important for remineralizing organic materials into

micronutrients. They could support both mycelial growth and plant nutrition [50].

Enterobacteriaceae, *Paenibacillaceae*, and *JACDCH01* were all present in higher

abundance in the high biomass treatments of both L*er* and Cvi. Most *Paenibacillaceae* members

predominantly inhabited soil, frequently in close association with plant roots [51]. These

rhizobacteria were known to play a significant role in enhancing plant growth and possessing

potential applications in agriculture and *Enterobacteriaceae* had been reported to enhance plant

growth [52–54].

Additionally, *Order_NS11-12g*, *BJHT01*, *JACDCH01*, and *UBA6156* were more

abundant in Cvi under high biomass conditions and less abundant in L*er* under high biomass

conditions. The majority of these were uncultivated or candidate taxa. Two alternative families

in the Order *Rhizobiales* appeared differentially associated with the two plant genotypes:

Labraceae with L*er* and *Rhizobiaceae* with Cvi [55]. These could indicate genotype-specific

interactions with different members of the microbial community.

The relative abundance of all OTUs that exhibited higher abundance (enriched) in

samples from the high biomass treatment in the L*er* and Cvi is depicted in Figures 8 (c) and 8 (d)

respectively. These OTUs consistently exhibit an increasing trend in relative abundance across

successive generations. A linear model was employed to test this trend, resulting in R-squared

values of 0.57 (P < 0.01) for L*er* and 0.73 (P < 0.01) for Cvi. Furthermore, OTUs that were

lower in abundance (depleted) in samples from the high biomass treatment in L*er* and Cvi are

depicted in Figures 8 (e) and 8 (f) respectively. These OTUs persist in low abundance across all

generations, suggesting a lack of selective pressure on this group of microbes in the samples

from the high biomass treatment.

Discussion

We applied artificial ecosystem selection for eight generations, in *Arabidopsis thaliana*

L*er* and Cvi to select soil microbiomes associated with higher or lower biomass of the host. In

contrast to some previous studies [27,33], we did not fertilize plants thus maintaining nutrient

limitation, and thereby promoting the plant's interaction with the microbiome [56]. Over the

course of eight generations, a response to the selection, most apparent after generations 4 and 5,

was observed both in the microbiome characteristics that shifted gradually and in the plant

biomass. Microbiome selection noticeably influenced plant biomass despite large phenotypic

variation from one generation to the next. Stochasticity in growth through generations may be

attributed to random environmental variations (REVs) as indicated by Swenson et al. [28]. This

was observed in the uninoculated sterile potting soil reference that showed similar patterns of

variability as the inoculated treatments across generations (Figure 1). Ecological variability

among generations even under controlled growth chamber conditions is a common characteristic

of studies [57] and may be due to minute fluctuations in growth chamber conditions or batch and

age of potting soil.

We observed a gradual decline in microbial species richness during the initial generations

of the experiment. This was evident from the observed OTU counts, as well as the Chao1 and

Shannon diversity indices. This pattern is consistent with findings from other studies [32–34],

highlighting the impact of selection pressures as the microbial communities adapted to the host

plant environment. However, in our study which continued for about twice the number of

generations of these earlier studies, we observed a stabilization of richness and alpha diversity in

the later generations. This was further substantiated by the stronger fit to the negative exponential

function when compared to a linear model, indicating constant and proportional changes in alpha

diversity and the rhizosphere microbiome's trend towards reduced diversity loss over time,

accompanied by heightened stability over eight generations.

To further understand the complex interplay between the forces of directed selection (plant

genotype and biomass treatment), we did a PCoA. Over successive generations, the results

demonstrate a strengthening effect of plant genotypes and biomass treatment on the microbial

community (Figure 5). This result was modeled using PERMANOVA, for both weighted and

unweighted UniFrac distances. Both weighted and unweighted UniFrac show marked increases in

the proportion of variance explained by genotype and biomass treatment, with unweighted UniFrac

exhibiting larger increases from generation to generation (Figure 6). Both metrics consider

phylogenetic relatedness. Weighted UniFrac considers the absolute abundance of OTUs and is

generally used to study changes in microbial community structure. Unweighted UniFrac considers

only the presence/absence of OTUs and is generally used to study changes in microbial community

composition. The results suggest that variance in microbiome composition as explained by

genotype consistently increases, whereas variance in microbiome structure as explained by

genotype, fluctuates more with changes in abundant taxa.

Here we would like to highlight that pronounced shifts in biomass between the high/low

biomass treatments, alpha diversity, and beta diversity all occur around generations 4 and 5 (Figure

1 and 2). This marks the point at which the changes in alpha diversity stabilize (Figure 4) and the

divergence between microbial communities by high/low biomass treatments in the PCoA plots

becomes more pronounced (Figure 5). It is possible that over the first four to five generations, the

starting microbial community underwent a period of restructuring before it stabilized and formed

four distinct communities under selection by high/low biomass treatments and by plant genotype.

We propose a complex interaction between plants and their associated microbiome ensued, where

differences in root exudation patterns between the two genotypes presumably established

associations with microbes in the early generations. Simultaneously, during this period, the

biomass treatment likely promoted host-specific microbe-mediated interactions that modulate

plant biomass. This restructuring of the microbial community is driven by both structural

(abundance of OTUs) and compositional (presence/absence of OTUs) effects in the microbiome

(Figure 6a and 6b). Differences in the abundance of taxa driven by the biomass and genotype

selection pressures, was a larger contributor to the generation-to-generation variation during the

experiment.

Changes in the assembly of microbial communities can arise from either selective pressure,

like the plant genotype or biomass treatment observed in this experiment, or stochastic processes,

like minute changes in growth chamber humidity or microenvironment. To gain deeper insights

into the influence of selection on microbial community assembly, a neutral model was fit to the

data [44,45]. Interestingly, this revealed a progressive decline in fit to neutrality, indicated by

decreasing R^2 values and increasing AIC values (Figure 7). This suggests an increasing influence

of selective forces such as biomass treatment and genotype over multiple generations of ecosystem

selection. This was also previously observed by Morella et al. [34].

Previous studies in *Arabidopsis thaliana* have often detected only a weak genotypic effect

on the rhizosphere microbiome [23,24,58]. However, a majority of these studies are focused on a

single growth cycle. Insights from plant-soil feedback research emphasize that the interactions

between plants and their microbiomes can be modified by consecutive growth cycles of the same

plant species in its respective soil [25,26]. Recent research has presented conflicting findings on

the influence of host genotype [33,34,59]. The different experimental designs, communities being

assessed and the length of these experiments, alongside high stochastic environmental variation

that is clearly associated with these microbial studies likely account for the conflicting findings.

In our investigation, we find that during the initial stages, genotype and biomass treatments have

modest but significant impacts. Over time, the plant genotype and biomass treatments have an

increasing influence. Together these selective forces explained ~40% of the variation in the

microbial community composition in these later generations.

A key aim of microbiome engineering has been to develop host-specific microbial

communities that impart lasting beneficial effects on the plant host [27,29,32,60]. Here we show

the enrichment of some common taxa in high biomass treatments but also some genotype-specific

changes. Despite starting with the same soil, only three common families were enriched in the

high biomass treatment samples of both the L*er* and Cvi genotypes. In addition, well-known plant

growth-promoting rhizobacteria were enriched in a genotype-specific manner for both L*er* and Cvi

high biomass treatment samples. In the Order *Rhizobiales*, *Labraceae* was enriched in L*er,* while

Rhizobiaceae was enriched in Cvi [55]. Moreover, in high biomass conditions, Cvi had increased

levels of the taxa Order_NS11-12g, BJHT01, JACDCH01, and UBA6156, while L*er* had

decreased levels of these. These data suggest genotype-specific plant genetic control of the

rhizosphere microbiome.

To further explore how the abundance of these OTUs changed across successive

generations, we grouped all the DA OTUs that were enriched or depleted in the samples of high

biomass treatment and calculated the sum of their relative abundance. The OTUs enriched in the

samples of high biomass treatment consistently displayed an increasing trend in relative

abundance over subsequent generations (Figures 8c and 8d). A linear model was fitted to this

data with high R^2 values, reinforcing the notion that the genotype and biomass treatments exerted

selective pressure on entire microbial groups, rather than individual species. Conversely, the

OTUs that were depleted in the samples of high biomass treatment consistently remained in a

state of low relative abundance throughout the entirety of the experiment (Figures 8e and 8f).

This observation may indicate that the selective pressures exerted by the genotype and biomass

treatments do not influence this particular group of microbes.

Taken collectively, these findings suggest the presence of diverse selection pressures

influences the rhizosphere microbiome. Initially, it seems that neutral processes seem to play a

major role in determining the structure, and composition of the microbiome (Figure 7). As

generations progress, we observe an increase in the strength of directional selection, possibly

driven by genotype and biomass treatment, resulting in the enrichment of specific groups of

microbes, including some PGPRs (Figures 8c and 8d). The persistence of depleted microbes in

high biomass treatment samples suggests the possibility for negative frequency-dependent

selection, preserving them at a low abundance. On the other hand, it is possible that this group is

experiencing no observable selection pressure.

Multiple studies have shown variations in the structure of the rhizosphere microbiome,

even within closely related plant genotypes, highlighting the importance of genotype-specific root

exudates in forming associations with the corresponding microbiome [61–65]. Further exploration

must prioritize highlighting the mechanisms that uphold these interactions through the utilization

of comparative metagenome, metatranscriptome and metaproteome analyses. This will play a part

in decoding chemical crosstalk, which could eventually foster interactions between plants and

microbes, leading to an overall enhancement in plant fitness.

The relationship between plants and microorganisms in the rhizosphere involves complex

and diverse interactions, which influence crucial ecological and physiological processes. These

interactions can be mutualistic, competitive, or antagonistic in nature. Many biotic and abiotic

factors act in concert to influence the dynamic rhizosphere microbiome. Gaining insights into how

these factors influence the assembly of microbial communities is crucial for obtaining targeted and

long-lasting advantages for plants. Nevertheless, establishing a strong and host-specific microbial

community that consistently provides beneficial effects continues to be a persistent challenge. We

have shown that, despite stochastic fluctuation due to REVs, it is possible to select for microbial

communities that impact biomass in a host-specific manner within four generations, addressing

this key challenge in microbial community engineering [32,60]. The rhizosphere microbiome that

evolves under plant-mediated selection offers better chances of survivability and efficacy when

applied as inoculum to the plant [66,67]. Multi-species ecosystem selection provides a perspective

on microbiome assembly that indicates the holobiont, and the evolving relationship between host

and microbiome contributes to emerging properties beyond those predicted by host genotype or

initial microbial community composition. This study enhances our understanding of the temporal

dynamics involved in microbial assembly in the rhizosphere and has implications for sustainable

agriculture, evolution, and ecology.

Figure 1. Ecosystem selection leads to changes in above-ground plant biomass. Above ground biomass for genotypes Ler (a) and Cvi (b) of *Arabidopsis thaliana*. Dark lines represent the high biomass treatment, and light lines represent the low biomass treatment. Each point represents the mean of n=50 microcosms. The mean biomass of the selection lines differed for several generations of both genotypes (t.test p-value * < 0.05; **<0.01;*** <0.001).

Figure 2. Above ground biomass represented as deviations from means for each generation. By plotting deviations from the mean, the influence of REVs on above-ground biomass is mitigated for genotypes L*er* (a) and Cvi (b). The dark lines denote high biomass treatments, while the light lines represent low biomass treatments. Each data point is the average of n=50 microcosms. (t.test p-value $* < 0.05$; $** < 0.01$; $*** < 0.001$).

Figure 3. Taxonomic distribution of the microbial community is represented in terms of the relative abundance at the Class level. The plot shows eight generations (1-8) for the high and low biomass treatments of both the Cvi (left) and L*er* (right) genotypes of *Arabidopsis thaliana*. Ten samples were assayed for generation 1 experiments while five were done for all subsequent generations. The most represented classes across all samples were *Alphaproteobacteria* (24.7%), *Gammaproteobacteria* (11.4%), *Bacteroidia* (10.1%), and *Verrucomicrobiae* (6.2%).

Figure 4. Changes in alpha diversity. Over the course of eight generations, two alpha diversity indices were computed, including the Observed and Shannon metrics. Results indicate a significant decline in the Observed number of OTUs, which estimates species richness. The Shannon index, which considers species abundance also exhibited a noteworthy decrease from generation 1 to 8. Additionally, a negative exponential function was fit to the Observed (R^2=0.81, P<0.01) and Shannon (R^2= 0.58, P<0.01) metrics, with the Observed showing a better fit indicating increased stability of the microbiome in later generations. (wilcox test *<0.05, **<0.01, *** <0.001).

Figure 5. Host genotype and biomass treatment shape the composition of the microbiome. Principle Coordinate Analysis (PCoA) using unweighted UniFrac for generations 1 through 8. Each point represents an individual sample with microbial communities defined of Cvi represented by red/orange circles and L*er* by dark/light blue triangles. The first two coordinate axes plotted accounted highest variation in the data, ranging from 13% in generation 1 to approximately 52% in generation 8. While there was no discernible clustering in generation 1, both genotypes and high/low biomass treatments exhibited greater clustering over the course of the experiment.

Figure 6. Variance in microbiome structure and composition explained by genotype and biomass treatment increases over generations. The R^2 values were computed using the PERMANOVA test with the adonis2 function for generations 1 through 8 for (a) weighted and (b) unweighted-UniFrac distances. The resulting R^2 and variance explained by residuals values for the biomass treatment (brown), genotype (blue) and residuals (black) are plotted. Residuals are plotted as a proxy for random environmental variations (REVs). There is a marked increase in the influence of both genotype and biomass treatment on differences in the microbial community from generations 1 through 8. Despite observing a decrease in the variance accounted for by the residuals, which could potentially signify a decline in the influence of REVs, the model elucidates more than 50 percent of the variability within the microbial community in the 8th generation. These differences are more pronounced in the weighted vs unweighted UniFrac. Model (distance.matrix~biomass treatment +genotype).

506
507
508
509

Figure 7. Fit of the Neutral model diminished as generations progressed. The occupancy (prevalence of OTU in samples) plotted against log(10) abundance is depicted using the Burns model. OTUs that are neutral, i.e., not selected for or against, are indicated in black. Green and blue-colored OTUs signify positive and negative selection, respectively. This plot highlights the presence of numerous microbes that undergo directed selection throughout the experiment.

511
512
513
514

Figure 8. Differentially abundant OTUs. ANCOMBC2 results are illustrated as log₂ fold changes at the family level for high biomass vs low biomass samples. Each taxon listed in the figure represents a single OTU in either the L*er* (a) or Cvi (b) microbial community identified to the family level if possible. The scale indicates log₂ fold change, red and blue for positive and negative fold change in the high biomass samples respectively. Well-known plant growth-promoting bacteria, such as *Paenibacillaceae, Bacilli, Labraceae, Rhizobiaceae* and *Bdellovibrio* are present in higher abundance. A vast majority of the other taxa identified are uncultivated and have little to no information on them. Figures (c) and (d) show box plots of relative abundance for all OTUs that were enriched in samples of the high biomass treatment in

genotype L*er* and Cvi, respectively, with colors indicating different generations. These OTUs demonstrate a consistent trend of increasing relative abundance across successive generations. To test this trend (red line), a linear model was fitted to the data, resulting in an R-squared value of 0.57 ($P < 0.01$) for L*er* and 0.73 ($P < 0.01$) for Cvi. Figures (e) and (f) are box plots of relative abundance for all OTUs that were depleted in samples from the high biomass treatment in genotypes L*er* and Cvi, respectively, and colored by generation.

517
518

Supplementary Materials: Figure S1: Rhizotubes schematic; Figure S2: Taxonomic

distribution of the microbial community represented in terms of the relative abundance at the

Phylum level. Table S1: Statistical comparisons of biomass for L*er*; Table S2: Statistical

comparisons of biomass for Cvi; Table S3: ANOVA for Observed alpha diversity index; Table

S4: ANOVA for Shannon alpha diversity index; Table S5: Linear mixed effect model for

Shannon diversity index; Table S6: PERMANOVA weighted UniFrac distances for generation 1;

Supplementary Table S7: PERMANOVA weighted UniFrac distances for generation 8;

Supplementary Table S8: PERMANOVA unweighted UniFrac distances for generation 1; Table

S9. PERMANOVA unweighted UniFrac distances for generation 8.

Author Contributions: N.S. Conceptualization, methodology, data analysis and visualization,

data collection, writing-original draft, writing-reviewing and editing. P.S. Methodology, data

analysis, writing-original draft, writing-reviewing, and editing. T.M. Data collection. R.K.

Resources, funding, writing-reviewing, and editing.

Funding: "This research received no external funding"

Data Availability Statement: All sequence data and metadata can be downloaded from

PRJNA1010286.

Acknowledgments: We would like to express our sincere gratitude to the following individuals

for their advice at various stages of this project Professor David Sloan Wilson, Dr. Jenny Kao-

541   Kniffin, Dr. Deepa Agashe, Dr. Gergely Maróti, Professor Douglas Woodhams, Professor

542   Emeritus Michael Shiaris, Dr. Karen Ricciardi, Dr. Patrick Kearns, Dr. Kourosh Zarringhalam,

543   Dr. David Degras-Valabregue, Professor Liam Revell and Professor Jay T. Lennon, Dr. Molly

544   Bletz, Dr. Brandon LaBumbard, and Cassandra Lyons.

References

550
1. Gouda S, Kerry RG, Das G, Paramithiotis S, Shin HS, Patra JK. Revitalization of plant growth promoting
rhizobacteria for sustainable development in agriculture. Microbiol Res. 2018;206.
2. Jones DL, Nguyen C, Finlay RD. Carbon flow in the rhizosphere: Carbon trading at the soil-root interface.
Plant Soil. 2009;321(1–2):5–33.
3. Dennis PG, Miller AJ, Hirsch PR. Are root exudates more important than other sources of rhizodeposits in
structuring rhizosphere bacterial communities? FEMS Microbiol Ecol. 2010;72(3):313–27.
4. Hartmann A, Schmid M, van Tuinen D, Berg G. Plant-driven selection of microbes. Plant Soil. 2009;321(1–
2):235–57.
5. Chaparro JM, Sheflin AM, Manter DK, Vivanco JM. Manipulating the soil microbiome to increase soil health
and plant fertility. Biol Fertil Soils. 2012;48(5):489–99.
6. Marschner P, Marschner P, Crowley D, Crowley D, Yang CH, Yang CH, et al. Development of specific
rhizosphere bacterial communities in relation to plant species, nutrition and soil type. Plant Soil. 2004;261(1–
2):199–208.
7. Garbeva P, van Veen J a, van Elsas JD. Microbial diversity in soil: selection microbial populations by plant and
soil type and implications for disease suppressiveness. Annu Rev Phytopathol. 2004;42(29):243–70.
8. Tkacz A, Cheema J, Chandra G, Grant A, Poole PS. Stability and succession of the rhizosphere microbiota
depends upon plant type and soil composition. ISME J. 2015;1–11.
9. Berg G, Smalla K. Plant species and soil type cooperatively shape the structure and function of microbial
communities in the rhizosphere. FEMS Microbiol Ecol. 2009;68(1):1–13.
10. Jin H, Yang X, Liu R, Yan Z, Li X, Li X, et al. Bacterial community structure associated with the rhizosphere
soils and roots of Stellera chamaejasme L. along a Tibetan elevation gradient. Ann Microbiol. 2018 May
1;68(5):273–86.
11. Barbosa Lima A, Cannavan FS, Navarrete AA, Teixeira WG, Kuramae EE, Tsai SM. Amazonian Dark Earth
and Plant Species from the Amazon Region Contribute to Shape Rhizosphere Bacterial Communities. Microb
Ecol. 2015 May 1;69(4):855–66.
12. Marques JM, da Silva TF, Vollu RE, Blank AF, Ding GC, Seldin L, et al. Plant age and genotype affect the
bacterial community composition in the tuber rhizosphere of field-grown sweet potato plants. FEMS
Microbiol Ecol. 2014;88(2):424–35.
13. Bodenhausen N, Bortfeld-Miller M, Ackermann M, Vorholt JA. A Synthetic Community Approach Reveals
Plant Genotypes Affecting the Phyllosphere Microbiota. PLoS Genet. 2014;10(4).
14. Zancarini A, Mougel C, Voisin AS, Prudent M, Salon C, Munier-Jolain N. Soil Nitrogen Availability and Plant
Genotype Modify the Nutrition Strategies of M. truncatula and the Associated Rhizosphere Microbial
Communities. PLoS ONE. 2012;7(10).
15. Wei S, Liu B, Ni K, Ma L, Shi Y, Leng Y, et al. Rhizosphere Microbial Community Shows a Greater Response
Than Soil Properties to Tea (Camellia sinensis L.) Cultivars. Agronomy. 2023 Jan 1;13(1).
16. Yuan J, Chaparro JM, Manter DK, Zhang R, Vivanco JM, Shen Q. Roots from distinct plant developmental
stages are capable of rapidly selecting their own microbiome without the influence of environmental and soil
edaphic factors. Soil Biol Biochem. 2015;89(AUGUST):206–9.

17. Houlden A, Timms-Wilson TM, Day MJ, Bailey MJ. Influence of plant developmental stage on microbial
community structure and activity in the rhizosphere of three field crops. FEMS Microbiol Ecol.
2008;65(2):193–201.
18. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, et al. Defining the core Arabidopsis
thaliana root microbiome. Nature. 2013 Aug 1;488(7409):86–90.
19. Bulgarelli D, Rott M, Schlaeppi K, Ver E, Themaat LV, Ahmadinejad N, et al. Revealing structure and
assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature. 2012;488(7409):91–5.
20. Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, et al. Plant
compartment and biogeography affect microbiome composition in cultivated and native *Agave* species. New
Phytol. 2016 Jan;209(2):798–811.
21. Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, et al. Structure, variation, and
assembly of the root-associated microbiomes of rice. Proc Natl Acad Sci U S A. 2015 Feb 24;112(8):E911–
20.
22. Wagner MR, Lundberg DS, Del Rio TG, Tringe SG, Dangl JL, Mitchell-Olds T. Host genotype and age shape
the leaf and root microbiomes of a wild perennial plant. Nat Commun. 2016 Jul 12;7.
23. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, et al. Defining the core Arabidopsis
thaliana root microbiome. Nature. 2012;488(7409):86–90.
24. Schlaeppi K, Dombrowski N, Oter RG, Ver Loren Van Themaat E, Schulze-Lefert P. Quantitative divergence
of the bacterial root microbiota in Arabidopsis thaliana relatives. Proc Natl Acad Sci U S A. 2014;111(2):585–
92.
25. Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, et al. Root exudate metabolites drive plant-soil feedbacks
on growth and defense by shaping the rhizosphere microbiota. Nat Commun. 2018 Dec 1;9(1).
26. Van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB, Fukami T, et al. Plant-soil feedbacks:
The past, the present and future challenges. J Ecol. 2013 Mar;101(2):265–76.
27. Panke-buisse K, Poole AC, Goodrich JK, Ley RE, Kao-kniffin J. Selection on soil microbiomes reveals
reproducible impacts on plant function. ISME J. 2014;1–10.
28. Swenson W, Wilson DS, Elias R. Artificial ecosystem selection. Proc Natl Acad Sci U S A. 2000 Aug
1;97(16):9110–4.
29. Mueller UG, Juenger TE, Kardish MR, Carlson AL, Burns KM, Edwards JA, et al. Artificial Selection on
Microbiomes To Breed Microbiomes That Confer Salt Tolerance to Plants METHODS AND PROTOCOLS
[Internet]. 2021. Available from: https://journals.asm.org/journal/msystems
30. Lau JA, Lennon JT. Rapid responses of soil microorganisms improve plant fitness in novel environments. Proc
Natl Acad Sci. 2012;109(35):14058–62.
31. terHorst CP, Lennon JT, Lau JA. The relative importance of rapid evolution for plant-microbe interactions
depends on ecological context. Proc R Soc B Biol Sci. 2014 Jun 22;281(1785):20140028.
32. Jochum MD, McWilliams KL, Pierson EA, Jo YK. Host-mediated microbiome engineering (HMME) of
drought tolerance in the wheat rhizosphere. PLoS ONE. 2019 Dec 1;14(12).
33. Cordovez V, Rotoni C, Dini-Andreote F, Oyserman B, Carrión VJ, Raaijmakers JM. Successive plant growth
amplifies genotype-specific assembly of the tomato rhizosphere microbiome. Sci Total Environ. 2021 Jun
10;772.
34. Norma M. Morella, Francis Cheng-Hsuan Weng, Pierre M. Joubert, C. Jessica E. Metcalf, Steven Lindow, Britt
Koskella. Successive passaging of a plant-associated microbiome reveals robust habitat and host genotype-
dependent selection. PNAS. 2020 Jan 14;117(2).
35. Parada AE, Needham DM, Fuhrman JA. Every base matters: Assessing small subunit rRNA primers for marine
microbiomes with mock communities, time series and global field samples. Environ Microbiol. 2016 May
1;18(5):1403–14.
36. Apprill A, Mcnally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly
increases detection of SAR11 bacterioplankton. Aquat Microb Ecol. 2015 Jun 4;75(2):129–37.
37. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-throughput
microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J. 2012 Aug;6(8):1621–4.
38. Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, et al. A communal catalogue reveals
Earth's multiscale microbial diversity. Nature. 2017 Nov 1;551(7681):457.
39. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive,
scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol. 2019 Aug 1;37(8):852–7.
40. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample
inference from Illumina amplicon data. Nat Methods. 2016 Jun 29;13(7):581–3.

41. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk v2: memory friendly classification with the genome taxonomy database. Bioinforma Oxf Engl. 2022 Nov 30;38(23):5315–6.

42. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol. 2020 May 1;37(5):1530–4.

43. Bunn AG. A dendrochronology program library in R (dplR). Dendrochronologia. 2008 Oct 1;26(2):115–24.

44. Burns AR, Stephens WZ, Stagaman K, Wong S, Rawls JF, Guillemin K, et al. Contribution of neutral processes to the assembly of gut microbial communities in the zebrafish over host development. ISME J. 2016;10(3):655–64.

45. Sloan WT, Woodcock S, Lunn M, Head IM, Curtis TP. Modeling taxa-abundance distributions in microbial communities using environmental sequence data. In: Microbial Ecology. 2007. p. 443–55.

46. Nearing JT, Douglas GM, Hayes MG, MacDonald J, Desai DK, Allward N, et al. Microbiome differential abundance methods produce different results across 38 datasets. Nat Commun. 2022 Dec 1;13(1).

47. Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction. Nat Commun. 2020 Dec 1;11(1).

48. Radhakrishnan R, Hashem A, Abd Allah EF. Bacillus: A biological tool for crop improvement through bio-molecular changes in adverse environments. Front Physiol. 2017 Sep 6;8(SEP).

49. Mujakić I, Piwosz K, Koblížek M. Phylum Gemmatimonadota and Its Role in the Environment. Microorganisms. 2022 Jan 1;10(1).

50. Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E. The Prokaryotes: A Handbook on the Biology of Bacteria: Archaea. Bacteria: Firmicutes, Actinomycetes. The Prokaryotes: A Handbook on the Biology of Bacteria. 2006.

51. Grady EN, MacDonald J, Liu L, Richman A, Yuan ZC. Current knowledge and perspectives of Paenibacillus: A review. Microb Cell Factories. 2016 Dec 1;15(1).

52. Ramesh A, Sharma SK, Sharma MP, Yadav N, Joshi OP. Plant Growth-Promoting Traits in Enterobacter cloacae subsp. dissolvens MDSR9 Isolated from Soybean Rhizosphere and its Impact on Growth and Nutrition of Soybean and Wheat Upon Inoculation. Agric Res. 2014;3(1):53–66.

53. Kämpfer P, Ruppel S, Remus R. Enterobacter radicincitans sp. nov., a plant growth promoting species of the family Enterobacteriaceae. Syst Appl Microbiol. 2005 Apr 22;28(3):213–21.

54. Khalifa AYZ, Alsyeeh AM, Almalki MA, Saleh FA. Characterization of the plant growth promoting bacterium, Enterobacter cloacae MSR1, isolated from roots of non-nodulating Medicago sativa. Saudi J Biol Sci. 2016;23(1):79–86.

55. Harkes P, van Steenbrugge JJM, van den Elsen SJJ, Suleiman AKA, de Haan JJ, Holterman MHM, et al. Shifts in the Active Rhizobiome Paralleling Low Meloidogyne chitwoodi Densities in Fields Under Prolonged Organic Soil Management. Front Plant Sci. 2020 Jan 10;10.

56. Meier MA, Xu G, Lopez-Guerrero MG, Li G, Smith C, Sigmon B, et al. Association analyses of host genetics, root-colonizing microbes, and plant phenotypes under different nitrogen conditions in maize. eLife. 2022;11.

57. Porter AS, Evans-Fitz.Gerald C, McElwain JC, Yiotis C, Elliott-Kingston C. How well do you know your growth chambers? Testing for chamber effect using plant traits. Plant Methods. 2015;11(1).

58. Bulgarelli D, Garrido-Oter R, Münch PC, Weiman A, Dröge J, Pan Y, et al. Structure and function of the bacterial root microbiota in wild and domesticated barley. Cell Host Microbe. 2015;17(3):392–403.

59. Brown SP, Grillo MA, Podowski JC, Heath KD. Soil origin and plant genotype structure distinct microbiome compartments in the model legume Medicago truncatula. Microbiome. 2020 Sep 28;8(1).

60. Basu A, Prasad P, Das SN, Kalam S, Sayyed RZ, Reddy MS, et al. Plant Growth Promoting Rhizobacteria (PGPR) as Green Bioinoculants: Recent Developments, Constraints, and Prospects. Sustainability. 2021 Jan 22;13(3):1140.

61. Micallef S, Colón-Carmona A. Genetic and Developmental Control of Rhizosphere Bacterial Communities. In: de Bruijn FJ, editor. Molecular Microbial Ecology of the Rhizosphere [Internet]. 1st ed. Wiley; 2013 [cited 2023 Aug 20]. p. 257–63. Available from: https://onlinelibrary.wiley.com/doi/10.1002/9781118297674.ch24

62. Micallef SA, Channer S, Shiaris MP, Colón-Carmona A. Plant age and genotype impact the progression of bacterial community succession in the Arabidopsis rhizosphere. Plant Signal Behav. 2009 Aug;4(8):777–80.

63. Lee S, I. Sergeeva L, Vreugdenhil D. Natural variation of hormone levels in *Arabidopsis* roots and correlations with complex root architecture: Natural variation of phytohormones in *Arabidopsis* roots. J Integr Plant Biol. 2018 Apr;60(4):292–309.

64. Matsushima C, Shenton M, Kitahara A, Wasaki J, Oikawa A, Cheng W, et al. Multiple analysis of root
exudates and microbiome in rice (Oryza sativa) under low P conditions. Arch Microbiol. 2021 Nov
1;203(9):5599–611.
65. Huang XF, Chaparro JM, Reardon KF, Zhang R, Shen Q, Vivanco JM. Rhizosphere interactions: Root
exudates, microbes, and microbial communities1. Botany. 2014;92(4):267–75.
66. Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. Plant–microbiome interactions: from community assembly to
plant health. Nat Rev Microbiol. 2020;18(11).
67. Gu S, Wei Z, Shao Z, Friman VP, Cao K, Yang T, et al. Competition for iron drives phytopathogen control by
natural rhizosphere microbiomes. Nat Microbiol. 2020 Aug 1;5(8):1002–10.

708