# The molecular reach of antibodies
# determines their SARS-CoV-2 neutralisation potency

Anna Huhn[1], Daniel Nissley[2], Daniel B. Wilson[1,3,‖], Mikhail Kutuzov[1], Robert Donat[4],
Tiong Kit Tan[4], Ying Zhang[3,†], Michael I. Barton[1], Chang Liu[5,6], Wanwisa Dejnirattisai[5,7],
Piyada Supasa[5], Juthathip Mongkolsapaya[5], Alain Townsend[4,6] William James[8], Gavin Screaton[5],
P. Anton van der Merwe[1], Charlotte M. Deane[2,¶], Samuel A. Isaacson[3,†,¶], Omer Dushek[1†,¶]

[1]Sir William Dunn School of Pathology, University of Oxford, Oxford OX1 3RE, UK
[2]Oxford Protein Informatics Group, Department of Statistics, University of Oxford, Oxford OX1 3LB, UK
[3]Department of Mathematics and Statistics, Boston University, Boston, Massachusetts, USA
[4]MRC Human Immunology Unit, MRC Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK
[5]Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford OX3 7BN, UK; Oxford University Hospitals NHS Foundation Trust, Oxford, UK.
[6]Chinese Academy of Medical Science Oxford Institute, University of Oxford, Oxford OX3 7BN, UK
[7]Division of Emerging Infectious Disease, Research Department, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkoknoi, Bangkok 10700, Thailand
[8]James & Lillian Martin Centre, Sir William Dunn School of Pathology, University of Oxford, Oxford OX1 3RE, UK
‖Present Address: Kirby Institute, University of New South Wales, Sydney, New South Wales, Australia
†Present Address: Department of Mathematics and Department of Biology, Northeastern University, Boston, Massachusetts, USA
†Equal contributor
¶Corresponding authors

**Key words:** Antibody; bivalent binding; surface plasmon resonance; antibody neutralisation potency;particle-based models; SARS-CoV-2

**Pre-print server:** biorxiv

## Abstract:

Antibodies play crucial roles in health and disease and are invaluable tools for diagnostics, research, and therapy. Although antibodies bind bivalently, we lack methods to analyse bivalent binding. Here, we introduce a particle-based model and use it to analyse bivalent binding of SARS-CoV-2 RBD-specific antibodies in surface plasmon resonance assays. The method reproduces the monovalent on/off-rates and enables measurements of new parameters, including the molecular reach, which is the maximum antigen separation that supports bivalent binding. We show that the molecular reach (22-46 nm) exceeds the physical size of an antibody (15 nm) and that the variation in reach across 45 patient-isolated antibodies is the best correlate of viral neutralisation. Using the complete set of fitted parameters, the model predicts an emergent antibody binding potency that equals the neutralisation potency. This novel analytical method should improve our understanding and exploitation of antibodies and other bivalent molecules.

## Graphical abstract:

# Introduction

Antibodies are multivalent molecules that contribute to immune responses by binding their antigens on the surfaces of pathogens. IgG antibodies have two identical antigen binding fragments (Fabs) fused to a constant fragment (Fc). These Fabs enable antibodies to achieve high-affinity bivalent binding by simultaneously engaging two antigens. This is important because the monovalent Fab/antigen interaction is often too weak to be effective and indeed, bivalent binding can dramatically increase the ability of antibodies to neutralise pathogens (1–4).

Although bivalent binding is critical to antibody function, there are no accepted methods to accurately analyse it. This has limited our understanding of the factors affecting bivalent antibody binding, such as the surface density of target antigens and the molecular reach, which is the maximum distance between antigens that allows for bivalent binding. A better understanding of these parameters would enable us to predict and modulate the potency of bivalent antibodies, and should help us understand how pathogens can evade antibody responses.

A prominent method for quantifying antibody/antigen binding is to use surface plasmon resonance (SPR). Typically, monomeric and monovalent soluble antigen is injected over immobilised antibodies. A simple ordinary differential equation (ODE)-based binding model is fit to the binding traces to determine the binding kinetics ($k_{on}$, $k_{off}$) and dissociation constant ($K_D$). However, these monovalent parameters correlate poorly with antibody function (5, 6) and do not allow prediction of the concentration of antibody required for functional effects, such as blocking viral infections (4).

While more physiological SPR experiments can readily be performed by injecting soluble antibodies over immobilised antigens, the spatial and stochastic effects of bivalent binding result in complicated binding traces that cannot be analysed using existing methods (7). Here, we show that a spatial and stochastic particle-based model can fit bivalent SPR data yielding accurate estimates of the monovalent binding parameters ($k_{on}$, $k_{off}$) and two new parameters: the bivalent on-rate ($k_{on,b}$) and the molecular reach. We use this approach to analyse a panel of 45 antibodies specific for the receptor-binding-domain (RBD) of SARS-CoV-2, and find that the strongest correlate of neutralisation potency is the molecular reach. Beyond correlations, we show that the particle-based model can directly determine the antibody concentration for viral neutralisation.

# Results

**A particle-based model accurately fits bivalent SPR data highlighting the impact of molecular reach on antibody binding**

We first used SPR to study the monovalent interaction between the IgG1 FD-11A antibody and RBD (8) (Fig. 1A, left). We injected monovalent RBD over a surface immobilised with FD-11A, followed by a buffer injection, and repeated this for different RBD concentrations. We were able to fit a monovalent ODE-based model simultaneously to the entire data set, providing estimates of the $k_{on}$, $k_{off}$, and $K_D$ (Fig. 1B-left,C).

We next reversed the orientation to study the bivalent interaction by injecting FD-11A at different concentrations over a surface presenting RBD (Fig. 1A, Bivalent SPR). We then attempted to fit a widely used

Figure 1: **A particle-based model quantifies the impact of molecular reach on bivalent antibody binding and accurately analyses bivalent SPR data.** (A) Schematic of the chemical reactions in monovalent and bivalent SPR experiments described by the indicated mathematical model. The ODE-based and particle-based bivalent models assume that any free antigen on the surface or only free antigen within reach, respectively, can be bound by an antibody already bound to the surface with one arm (grey shaded regions). (B) Representative monovalent (left) or bivalent (middle, right) SPR traces using the antibody FD-11A interacting with RBD of SARs-CoV-2 (black) overlaid by the model fit (red). The family of SPR traces are generated by 2-fold dilution of RBD starting at 2000 nM (left) or by a 2-fold dilution of FD-11A starting at 300 nM (middle, right - same data but different model fit). (C) The fitted binding parameters for monovalent ($N = 4$) and bivalent ($N = 6$) SPR experiments. (D) Simulated SPR traces (bottom) for antibodies injected over a surface with a random distribution of antigen (top) but with different values of the molecular reach (columns). The antigens (circles) are coloured red if they are within reach of another antigen and blue otherwise. Parameter values: $k_{on} = 0.05 \mu M^{-1} s^{-1}$, $k_{off} = 0.05 s^{-1}$, $k_{on,b} = 1.0 s^{-1}$, and $[RBD] = 0.0025$ nm$^{-2}$. (E) The fitted bivalent binding parameters from the $N = 6$ bivalent SPR experiments. (F,G) Comparison of five antibodies analysed using monovalent and bivalent SPR from $N \geq 3$ independent experiments. (F) Comparison of the indicated parameter using both methods with the dashed line displaying perfect agreement (y=x). (G) The bivalent binding parameters. An F-test was used to determine a p-value for the null hypothesis that the dashed line and the fitted line to log-transformed binding parameters were equal (F) and a t-test with Dunnett's multiple comparison correction on log-transformed values was used to determine p-values (C). Abbreviations: ns = p> 0.05, * = p≤0.05, ** = p≤0.01, *** = p≤0.001, **** = p≤0.0001.

ODE-based bivalent model, which is included with the SPR fitting software, to these data (Fig. 1B, middle). This model adds a second binding step to the first binding step with a bivalent on-rate ($k_{on,b}$) and the same $k_{off}$. While the fit was not unreasonable, this model provided inaccurate values for $k_{off}$ and K$_D$ (Fig. 1C). A

key assumption of such ODE models is that the molecules involved are 'well-mixed'. While this assumption is reasonable for the first step, it becomes unreasonable for the second step because, once the antibody is bound to a surface immobilised antigen, the monovalent antibody/antigen complex can only bind a second antigen within reach whereas the well-mixed condition of the model assumes it can bind any free antigen on the surface. Moreover, the number of free antigens within reach is expected to be low and to decrease over time as more antibody binds. As a result, deterministic ODE models also fail to capture the local stochasticity of bivalent binding.

To address these limitations we developed a more realistic stochastic and spatially resolved particle-based model using the Gillespie method (Fig. 1A, right). In this model, the monovalent antibody/antigen complex can only bind a second antigen if it is within reach (with rate $k_{on,b}$). We used the model to simulate bivalent SPR traces with three different reach distances (Fig. 1D). With a very short reach, antibodies could not bind a second antigen, resulting in monovalent binding with fast dissociation. Increasing the reach allowed a larger fraction of antibodies to engage in bivalent binding, leading to higher levels of binding at a given concentration and much slower dissociation. This highlights the crucial role of molecular reach in determining antibody potency and binding stability.

We developed a workflow to fit the particle-based model directly to bivalent SPR data (Fig. S1). This produced an excellent fit (Fig. 1B - right) and, unlike the ODE-based bivalent model, yielded $k_{on}$, $k_{off}$ and $K_D$ values in agreement with those obtained by monovalent SPR (Fig. 1C). Importantly, the model fit provided estimates of the bivalent on-rate and reach (Fig. 1E). We found the same binding parameters when analysing bivalent SPR with different levels of RBD on the chip surface, which confirms that the particle-model is correctly capturing how the antigen density impacts bivalent antibody binding (Fig. S2).

To further validate the model, we performed monovalent and bivalent SPR on four additional RBD antibodies (Fig. S3). We confirmed that the particle-model correctly estimated the values of $k_{on}$, $k_{off}$, and $K_D$ across a 100-fold variation in affinity (Fig. 1F). As before, the particle model also provided estimates of the bivalent on-rate and the reach (Fig. 1G).

Finally, we repeated the analysis with an antibody that recognises a different antigen, namely CD19 (Fig. S4A). Once again we found agreement between the ODE-based model analysing standard monovalent SPR and the particle-based model analysing bivalent SPR (Fig. S4B).

In conclusion, a particle-based model accurately fits bivalent SPR data and allows measurement of binding parameters crucial for understanding bivalent antibody binding.

## The molecular reach is determined by both the antibody and the antigen

The molecular reach distances that our analysis produced for the five RBD antibodies (∼38 nm, Fig. 1G) and the CD19 antibody (53 nm, Fig. 4C) were notably larger than previous studies reporting that IgG1 antibodies can only bind bivalently when antigens were within ∼16 nm (9–11). They were also larger than estimates of the distance between the antigen-binding sites based on structural studies of whole antibodies (∼15 nm) (1). Furthermore, molecular dynamic (MD) simulations of the FD-11A antibody produced reaches of 3.46 to 17.58 nm with a mean of 13.05 nm (Fig. S7).

One possible explanation for these discrepancies is antigen size. Previous studies of molecular reach had used small antigens (e.g. 4-hydroxy-3-iodo-5-nitrophenylacetate (NIP, 320 Da) (9), digoxin (780 Da) (10),

6x His-Tag (1100 Da) (11)) while we used much larger protein antigens (e.g. RBD is 51,100 Da). To test this hypothesis, we extended our analysis to small antigens. We injected an anti-phosphotyrosine antibody over a small phosphorylated peptide antigen coupled to polyethylene glycol (PEG) linkers comprised of 3 or 28 PEG repeats (Fig. 2A,B, Fig. S5). The size of these peptide/linker complexes was 2234 Da (3 PEG repeats) and 3336 Da (28 PEG repeats). Consistent with previous measurements using small antigens (9–11), our analysis produced a reach of 10.3 nm using the PEG3 linker and 13.4 nm using the PEG28 linker (Fig. 2B). Similar results were obtained at different PEG3 concentrations (Fig. S6). Given that the flexibility of PEG has previously been determined (12), we used the worm-like-chain (WLC) polymer model to estimate the size of the antigen when coupled to PEG3 (0.7 nm) or PEG28 (2.1 nm). This allowed us to calculate the theoretical increase in molecular reach (see Methods), which agreed well with the experimentally measured increase (Fig. 2C).

Figure 2: **The molecular reach is similar to the predicted antibody reach when the antigen is small. (A)** Representative SPR traces of the anti-phosphotyrosine antibody (PY20) injected over surfaces coupled to a small phosphorylated peptide linked to either 3 or 28 repeats of PEG. The antibody was injected at 8 concentrations (25 nM with 2-fold dilutions). **(B)** The fitted molecular reach ($N = 9$ for PEG3, $N = 5$ for PEG28; see Fig S5 for other fitted parameters). **(C)** The difference in reach between PEG3 and PEG28 estimated by the worm-like-chain model (see Methods) or by the difference in reach from panel B.

To determine if a ∼38 nm reach was plausible for RBD binding antibodies, we used coarse-grained steered MD simulations to determine the maximum distance between RBDs that could support bivalent antibody binding. We therefore constructed a coarse-grained representation of the FD-11A antibody bound to RBD, modified to include the Lys15 biotinylation site used to anchor RBD to the SPR chip surface (Fig. 3A). One Lys15 position was held fixed while the other was pulled away at constant velocity. We computed the fraction of native contacts at both paratope/epitope interfaces (Fig. 3B) and the exerted pulling force (Fig. 3C). We identified unbinding events based on a decrease in the fraction of native contacts and this could be confirmed by visual inspection (Fig. 3D). We recorded the maximum distance between Lys15 on the two RBDs before the antibody unbound. This steered MD procedure allowed us to much more rapidly estimate the maximum possible reach for FD-11A in comparison to unrestrained simulations that would only rarely access extended conformations, which we expect are readily accessed in experiments that take place on the timescale of minutes. We ran sets of simulations using different paratope/epitope interface strengths and as expected, increasing the strength allowed the simulation to explore bivalent binding conformations with a larger maximum RBD distance (Fig. 3E). Defining the molecular reach as the maximum distance achieved over different interface strengths, we obtained a value of 34.4 nm for FD-11A, which agrees well with the 34.9 nm reach estimated by bivalent SPR (Fig. 1E). We next repeated this procedure for five additional antibodies (Fig. S8) finding agreement with the SPR-determined reach (Fig. 3F).

Taken together, these findings demonstrate that our particle-based model can produce accurate estimates of the molecular reach from bivalent binding data. These results also show the molecular reach is the maxi-

mum separation distance between antigen anchoring points that support bivalent binding, which underlines the importance of antigen size.

Figure 3: **Molecular dynamic simulations reproduce the molecular reach obtained by bivalent SPR highlighting the contribution of the antigen to reach.** (A) Coarse-grained structure of IgG1 FD-11A bound to two RBD antigens indicating the Lys15 biotinylation site (within an N-terminus AviTag), which anchors RBD to the SPR surface. The anchor point of RBD forming interface 1 was fixed while the one forming interface 2 was moved at a constant velocity. **B** The fraction of native contacts at the indicated interface (normalised to the number of contacts in the native structure) and (C) the force required to maintain the constant velocity over time. (D) Snapshots from the simulations at the indicated time points indicating the distance between Lys15 on each RBD. The maximum distance in this trajectory was 34.4 nm. (E) The maximum Lys15-Lys15 distance from $N = 50$ independent trajectories over the interface binding strength ($\eta$). The molecular reach is defined as the largest distance (horizontal dashed line). (F) The molecular reach for the indicated antibody estimated by simulation or experiment (left) and the difference between these estimates (right). A one-sample t-test is used to determine the p-value for the null hypothesis that the mean is 0.

## A systematic analysis of RBD-specific antibodies reveals a correlation between molecular reach and SARS-CoV-2 neutralisation potency

We next investigated the functional implications of molecular reach using a panel of 80 RBD-specific monoclonal antibodies previously isolated from SARS-CoV-2 infected individuals with known epitopes and viral neutralisation potencies (4) (Fig. 4A). The neutralisation potency is the concentration producing 50% inhibition ($IC_{50}$). We injected 3 concentrations of each antibody sequentially, analysing up to 32 antibodies in a single 48 hour experiment (Fig. 4B). Since we needed to re-use the same surfaces repeatedly, we used high salt (3M MgCl) to regenerate the surface between antibody injections without denaturing the antigen. In a pre-screen, we found 7 out of 80 antibodies remained bound after regeneration, and so these were excluded from subsequent experiments. We injected the FD-11A antibody at the start and end of each of the 16 ex-

periments that we performed to confirm that the antibody binding capacity of the surface remained largely intact over 48 hours despite multiple regeneration steps (Fig. S9A,B).

We next fitted the particle-based bivalent model and the ODE-based monovalent model to the bivalent SPR data. We excluded 12 antibodies because they produced a poor fit to the particle-based model and 16 antibodies because the ODE-based model produced an accurate fit (Fig. S9C-G). We reasoned that an accurate fit by the ODE-based model meant that the bivalent SPR data contained no useful information about bivalent binding parameters. Possible explanations include an antibody that only binds monovalently because of insufficient reach or an antibody that binds with very high affinity so that it does not unbind during the experiment. In these cases, it is not possible to quantify the increase in binding that bivalency provides.

The binding parameters for the remaining 45 antibodies displayed a 1000-fold variation in affinity (Fig. 4C), which was primarily the result of variations in the off-rate (Fig. 4D). The molecular reach varied from 22 to 46 nm and, interestingly, displayed some correlation with affinity and off-rate (Fig. 4C,D). This is consistent with previous work suggesting that higher-affinity antibodies can tolerate larger antigen distances when binding bivalently (9).

We found only modest correlations between neutralisation potency and the monovalent binding parameters (Fig. 5A-C). We considered that the binding parameters may independently contribute to predicting potency and therefore tried multiple linear regression, but the correlation was similar to the affinity alone (Fig. 5C vs D). We next investigated whether bivalent binding parameters could predict potency. Although the bivalent on-rate did not correlate (Fig. 5E), the molecular reach displayed the best correlation of all single binding parameters (Fig. 5F), which increased further using a multiple linear regression model that included all the parameters (Fig. 5G). This indicates that antibodies with longer reaches are better able to neutralise virus.

We reasoned that the ability of the molecular reach to predict neutralisation potency may be a result of it's ability to predict bivalent binding and/or it's ability to act as a proxy for the relevant blocking epitope. To test the latter hypothesis, we computed the distance of each antibody epitope from a reference epitope in the ACE2 binding site (Fig. 5H). As expected, we found that neutralisation potency was gradually reduced for antibodies that bound further from the blocking epitope interface (Fig. 5I). However, we found that the molecular reach did not correlate with this blocking epitope distance (Fig. 5J). We also used the previously reported epitope taxonomy for RBD-specific antibodies but again, found no difference in molecular reach depending on epitope location (Fig. S10). This suggested that molecular reach predicts bivalent binding rather than the blocking epitope.

The observation that both epitope and reach correlated with neutralisation potency but not with each other suggested that each contained independent information relevant to viral neutralisation potency. We first confirmed that including the blocking epitope distance with the monovalent binding parameters improved the correlation (Fig. 5D vs K). In support of our hypothesis, we found that the correlation with all the binding parameters improved further when including the blocking epitope distance (Fig. 5G vs L).

Taken together, these results show that the best single parameter predictor of neutralisation potency is the molecular reach, and that increasing the molecular reach increases the neutralisation potency of RBD antibodies by enhancing bivalent binding.

Figure 4: **High throughput analysis of antibody binding using bivalent SPR. (A)** Schematic of antibody identification and characterisation. **(B)** Workflow for high throughput bivalent SPR indicating that each antibody was injected at 3 concentrations with association (A, 150 s), dissociation (D, 450 s), and surface regeneration (R) steps. **(C)** Complete set of binding parameters for 45 antibodies ordered by affinity (left) or displayed as distributions (right - mean with min/max shown). **(D)** Parameter correlations.

Figure 5: **The molecular reach of RBD-specific antibodies correlates with neutralisation potency but not with the ACE2 blocking epitope. (A-G)** Single and multiple linear correlations between the indicated parameters of 45 antibodies (individual data points) and their neutralisation potency ($IC_{50}$), the concentration of antibody required to reduce *in vitro* SARS-CoV-2 infection by 50%. The 13 antibodies whose potency was not determined (n.d) were excluded from the fit (top dashed line). **(H)** Structure of RBD showing the ACE2 binding site (blocking epitope, green) and the epitope location of the RBD antibodies on the structure (middle) or in free space and rotated 180° (right). The blocking epitope distance is calculated using the indicated reference point. Images (left, middle) are taken from (4). **(I,J)** Correlation between blocking epitope distance and (I) neutralisation potency or (J) reach. **(K,L)** Multiple regression with the blocking epitope distance using (K) only the monovalent binding parameters or (L) all the binding parameters. The neutralisation potency and epitope locations were previously determined (4). Multiple linear regression included only main effects and all correlations were performed on log-transformed parameters except for molecular reach and epitope distance.

**The particle-based model calculates the viral neutralisation potency of antibodies**

The neutralisation potency of an antibody is an emergent property that depends on multiple factors including its binding parameters ($k_{on}$, $k_{off}$, $k_{on,b}$, reach), the epitope, and the antigen density on the pathogen surface. With the exception of antigen density, these factors can be combined to predict neutralisation potency as we have demonstrated (Fig. 5L) but this requires a panel of antibodies with a wide variation of measured neutralisation potencies in order to fit a regression model. We reasoned that the particle-model could incorporate all of these factors to directly calculate an emergent measure of antibody binding potency without a priori data fitting. To do this, we developed a workflow that used the particle-model to simulate the amount of free antigen on a two-dimensional surface for different antibody concentrations from which the predicted binding potency (antibody concentration required to bind 50% of antigen) can be calculated (Fig. 6A).

We first used the FD-11A antibody to validate the workflow. We simulated the amount of free antigen at 60 minutes using the FD-11A binding parameters (Fig. 1C,E) for different FD-11A concentrations and for surfaces with 7 antigen densities (Fig. 6B). To compare these simulations with data, we mixed different concentrations of FD-11A IgG or Fab with live SARS-CoV-2 for 60 minutes before adding it to Vero cells and determined infection at 20 hours (8) (Fig. 6C). We compared the predicted binding potency with the experimental neutralisation potency (Fig. 6D).

At low antigen densities (e.g. 0.00000256 nm$^{-2}$), the model predicted a poor binding potency of 600 nM that matched the expected affinity of a bivalent antibody that can only bind a single antigen ($K_D/2 = 614$ nM, where the factor of 2 accounts for the two antibody Fabs and $K_D = 1228$ nM is the monovalent affinity). This is presumably because the average distance between antigens at this density ($\sim$352 nm) is much larger than the molecular reach of FD-11A (34.9 nm) impairing bivalent binding. The predicted binding potency improved by $>$ 100-fold as the antigen density increased enabling FD-11A to bind bivalently and matched the experimental potency at an antigen density of $\approx$0.008 nm$^{-2}$. The Fab failed to neutralise virus even at concentrations above 3000 nM, which likely reflects the fact that the monovalent antibody/RBD affinity is much lower than the ACE2/RBD affinity ($K_D \sim$75 nM (13)).

We next used the workflow to predict binding potency at different antigen concentrations for all 45 antibodies. We found that the predicted and experimental potencies were similar at antigen densities of 0.0005-0.001 nm$^{-2}$ and this could be further improved if only a subset of 24 antibodies were included that bound near the blocking epitope (Fig. S11, Fig. 6E). As above for FD-11A, we found that at very low and very high antigen densities, the predicted binding potency was very low or very high for all antibodies, respectively, and it differed significantly from the measured neutralisation potency (Fig. 6E). This difference can be explained because at very low and very high antigen densities all the antibodies were unable to reach two antigens or all antibodies could reach two antigens, respectively, independent of their molecular reach. In contrast, there was no significant difference with neutralisation potency at intermediate antigen densities (0.0005-0.001 nm$^{-2}$) where the mean distance between antigens is $\approx$18-25 nm, which is on the same scale as the measured molecular reach of RBD antibodies (Fig. 4C). Importantly, these intermediate densities are similar to the estimated density of the Spike protein on the surface of SARS-CoV-2 (Fig. 6F).

These results show that, if we know the antigen density on the virion surface, it is possible to directly calculate the concentration of antibody required to neutralise SARs-CoV-2 from bivalent SPR-determined antibody binding parameters without prior data fitting. Importantly, they demonstrate that the molecular reach of antibodies is crucial in determining their neutralisation potency at antigen densities found on SARS-CoV-2 virions.

Figure 6: **The antibody binding potency calculated by the particle-model equals the antibody neutralisation potency determined by live virus experiments.** **(A)** Schematic of workflow to determine the predicted antibody binding potency defined as the concentration of antibody required to bind 50% of antigen randomly distributed on a two-dimensional surface. The model input includes the antibody/antigen concentrations and binding parameters. **(B)** The fraction of free antigen over antibody concentration after 60 minutes for surfaces with different concentrations of antigen. The binding parameters are those determined for FD-11A. **(C)** The ability of live SARS-CoV-2 virions to infect target cells when pre-mixed with different concentrations of FD-11A IgG (red) or Fab (black) determined at 20 hours. **(D)** Fitted potency values ($IC_{50}$) from the (B) simulations and (C) experiments. **(E)** Comparisons of the predicted binding potency over the experimental neutralisation potency for 25 antibodies that bind within 2.37 nm of the blocking epitope for the indicated antigen concentrations (see Fig. S11 for all antibodies and all antigen densities tested). A linear fit on log-transformed $IC_{50}$ values (solid line) is compared to the absolute line (y=x) using an F-test. The shading of each point indicates the molecular reach of the respective antibody. **(F)** Comparison of the Spike density on SARS-CoV-2 virion (14) with the intermediate antigen density producing absolute agreement using the particle-model (0.00075 nm$^{-2}$ [0.0005, 0.001])

# Discussion

Modelling bivalent antibody binding to a random distribution of antigen is challenging because the process is spatial (antibodies only bind antigens within reach) and stochastic (antibodies typically only reach a handful of antigens), which precludes the use of standard ODE-based models. We used a particle-based model in order to capture the spatial and stochastic nature of bivalent binding. This model allowed us to estimate key bivalent binding parameters from bivalent SPR data, and predict neutralisation potency of a range of antibodies from these parameters.

The molecular reach that we report for RBD-specific antibodies (up to 46 nm) is much larger than previous reports (up to ∼16 nm) (9–11). We have shown this to be the result of differences in the sizes of the antigens used, with previous reports focused on low molecular weight model antigens (<1100 Da) compared to the the RBD antigen (51 kDa). This suggests that antibodies can simultaneously bind two antigens anchored much further apart than the physical size of an antibody provided that the antigen is large. As a result, the molecular reach for bivalent antibody binding depends on both the antibody and the antigen.

The molecular reach was the best single-parameter predictor of viral neutralisation. Indeed, the distribution of distances between Spike antigens on the SARS-CoV-2 virion (15) overlaps with the distribution of molecular reaches that we report enabling antibodies with a longer reach to more efficiently bind and neutralise virus. Consistent with this, we have shown that reach is not simply a proxy for the blocking epitope but rather a predictor of bivalent binding that is likely dependent on both epitope and binding orientation.

Using the complete set of bivalent parameters, including reach, we have shown that the particle-model can be used to calculate an emergent antibody binding potency that directly predicted neutralisation potency without prior data fitting. This relied on using an antigen density in the model that is similar to the density of Spike on the surface of SARS-CoV-2. Although this calculation is based on simulations using monomeric antigen, it is noteworthy that the Spike protein is trimeric and contains three RBDs that are closer together than the molecular reaches that we report. This apparent discrepancy can be accounted for by the fact that only one RBD domain on a Spike protein is typically accessible (14) and/or that the spacing and relative orientation of the RBDs may preclude bivalent antibody binding to two RBDs on a single Spike protein.

While we have focused on analysing antibody binding by SPR, the model can readily be used to study any bivalent molecule using any instrument that measures binding. Moreover, in addition to providing the complete set of bivalent binding parameters, bivalent SPR accelerates antibody screens by enabling sequential injection of many antibodies over the same surface. We have also demonstrated that these bivalent binding parameters can be used to predict antibody function. In summary, the ability to easily acquire and now analyse bivalent binding should improve the ability to study and engineer antibodies and other native and synthetic bivalent molecules.

## Acknowledgements

We thank Citlali Solis Salas, Jack Paget, and Philip K. Maini for their effort to formulate and analyse a master equation representation of the particle-model. We thank Lisa Schimanski and Pramila Rijal for assistance in producing monoclonal antibodies. Surrogate construction was carried out using the Shared Computing Cluster administered by Boston University's Research Computing Services.

## Funding

The work received funding from a Wellcome Trust Senior Fellowship in Basic Biomedical Sciences (207537 /Z/17/Z to OD) and the National Science Foundation (NSF-DMS 1902854 to SAI, supporting SAI, DBW, and YZ). We acknowledge NIH Research Biomedical Research Centre Funding Scheme (to GRS) and the Chinese Academy of Medical Sciences Innovation Fund for Medical Science, China (2018-I2M-2-002 to DIS/GRS). We are also grateful for a Fast Grant (Fast Grants, Mercatus Center), Schmidt Futures, and the Red Avenue Foundation. The Wellcome Centre for Human Genetics is supported by the Wellcome Trust (090532/Z/09/Z).

## Competing interest statement

GRS is on the GSK Vaccines Scientific Advisory Board, a founder shareholder of RQ biotechnology, and a Jenner investigator. Oxford University holds intellectual property related to the Oxford-AstraZeneca vaccine.

## Open access

# Materials & Methods

## Proteins

**Production of monovalent Streptavidin.** Monovalent Streptavidin was produced using a method previously described to generate streptavidin tetramers of defined valency fused to Spycatcher ([16]). Two different streptavidin subunits were used: Streptavidin-SpyCatcher, which contains a functional streptavidin monomer fused to SpyCatcher at its C terminus and a 'dead' streptavidin, which contains a mutation in the streptavidin monomer that has negligible biotin-binding activity. Individual subunits were expressed in *E. coli* BL21-CodonPlus (DE3)-RIPL cells and refolded from inclusion bodies. Inclusion bodies were washed in BugBuster (Merck Millipore 70921) supplemented with lysozyme, protease inhibitors, DNase I, and magnesium sulphate as per the manufacturers' instructions. To obtain monovalent streptavidin-SpyCatcher, the subunits were mixed at a 3:1 molar ratio of dead streptavidin to streptavidin-SpyCatcher. Tetramers were refolded by rapid dilution and precipitated using ammonium sulphate precipitation. Precipitated protein was resuspended in 20 mM Tris (pH 8.0), filtered (0.22-μm filter), and loaded onto a Mono Q HR 5/5 column (GE Healthcare Life Sciences). Desired tetramers were eluted using a linear gradient of 0–0.5 M NaCl in 20 mM Tris (pH 8.0), concentrated, and buffer exchanged into 20 mM MES, 140 mM NaCl (pH 6.0).

**Production of Biotinylated-RBD.** Affinity purified SARS-CoV-2 (Wuhan) RBD was biotinylated using EZ-Link Sulfo-NHS-LC-biotin (Life Technologies, USA, A39257) accordingly to manufacturing protocol. Biotinylated RBD was subjected to Zeba™ Spin Desalting Columns (7k MWCO) (Thermo Scientific, USA, catalogue 89889) to remove excess biotin.

**Production of FD-11A, FD-5D, EY-6A, FI-3A, CR3022, REGN10987 antibodies.** Monoclonal antibodies were produced as previously described ([8], [17]). Expression plasmids were transfected into the ExpiCHO cell lines according to the manufacturer's protocol (Thermo Fisher). Supernatant containing monoclonal antibodies were clarified by centrifugation (1,400g, 5min) and 0.45 μM filtered before purification. Monoclonal antibodies were affinity purified using a MabSelect SuRe (Cytiva) pre-packed column. Purified mAbs were then desalted using Zeba Spin Desalting Column (ThermoFisher) or diafiltered using a Amicon Untracentrifugation Column (50k MWCO).

**Production of 80 mAbs specific for RBD.** Antibodies for high-throughput screen used in the present study were produced previously ([4]).

**Production of Spytag-CD19 antigen and antibody.** Spytag-CD19 fused to Spytag was produced as previously described ([18]). The extracellular domain of CD19 is expressed as a fusion protein with Spytag and Histag fused to its C-terminus, and the Sumo protein via a HRV cleavage site fused to its N-terminus (Sumo-HRV-CD19-Spytag-Histag). The protein SUMO was used to stabilise Spytag-CD19 during production. A CMV expression plasmid encoding Sumo-HRV-CD19-Spytag-Histag was transfected into Expi293F Cells (ThermoFisher) using the ExpiFectamine 293 Transfection Kit (ThermoFisher Scientific, A14524). Cells were incubated for 4-5 days for protein expression. Following, the supernatant was harvested, and in the first step the fusion protein was purified using Ni-NTA Agarose column. Next, the protein was concentrated and loaded onto a Superdex 200 10/300 GL (Cytiva, 17-5175-01) size exclusion chromatography column. Next, HRV 3C Protease Solution Kit (Pierce™, 88946) was used to cleave SUMO from the CD19 fusion protein. The protease was removed via Glutathione Agarose (Pierce™, 16100), followed by Ni-NTA Agarose to remove SUMO. Finally, the protein was stored in size exclusion buffer (25 mM NaH2PO4 and 150 mM NaCl at pH 7.5) and frozen in suitable aliquots at –80°C.

The anti-CD19 antibody (clone SJ25C1) was purchased from BioLegend (cat no. 363001).

**Production of PEG coupled phosphorylated peptide antigens and antibody.** Phosphorylated peptide antigens were a custom commercial order (Protein Peptide Research Ltd, UK) with the following sequences: Bio-(PEG)$_3$-SVPEQTEY*ATIVFPSG (PEG3) and Bio-(PEG)$_{28}$-SVPEQTEY*ATIVFPSG (PEG28), where Bio indicates biotin, PEG$_X$ is X repeats of polyethylene glycol, and * indicates phosphorylation.

The anti-phosphotyrosine antibody (clone PY20) was purchased from Absolute Antibody (Ab00294-1.1).

## Surface plasmon resonance

**Monovalent SPR for RBD antibodies.** A BIAcore 8K (Cytiva) was used to measure the affinity and the kinetics of soluble RBD to immobilised antibody. Approximately 200 RU of antibody was captured onto a Protein A Series S Sensor Chip (Cytiva) along with 200 RU of an influenza mAb AG7C in a reference flow cell. Multi cycle kinetic analysis of binding was undertaken, using a two-fold serial dilution of RBD-H in HBS-P+ buffer (Cytiva) along with a reference sample containing only HBS-P+ buffer. Measurements were made with injection times of 90 s (30 μl/min) and dissociation times of 180 s or 600 s (30 μl/ml) at 37°C. Regeneration of the sensor chip was performed with 10 mM Glycine-HCl, pH 1.7 for 30s (30 ul/min) between RBD-H concentrations. For analysis, the sensograms were double reference subtracted and fitted with a 1:1 binding model using the BIAcore Insight Evaluation Software version 2.0.15.12933 (Cytiva).

**Monovalent SPR for CD19 antibodies.** Monavalent SPR of CD19 antibodies was conducted on BIAcore T200 instrument (GE Healthcare Life Sciences). The experiment was run at 37 °C, HBS-EP was used as running buffer. An Fc capture chip was produced by amine coupling of anti-mouse IgG antibody to a CM5 chip using a commercial Fc Capture Kit (GE Healthcare). First, the chip was conditioned with 7 conditioning cycles using HBS-EP buffer. Next, the anti-CD19 antibody was injected for either 120 s or 900 s, resulting in immobilisation levels of 500 RU and 1000 RU respectively. For the control flow cell, the BBM1 antibody was immobilised to matching levels. CD19-SpyCatcher was injected for 150 s, followed by a dissociation phase of 450 s at a flow rate of 50 μl/min. After each cycle, the chip was regenerated with 10 mM Glycine-HCl, pH1.7 for 90 s, which removed both CD19 and anti-CD19 antibody from the chip. This was followed by the re-immobilisation of anti-CD19 antibody. Buffer was injected after every second cycle. For data analysis, the SPR sensogram was double-referenced against an empty flow cell and buffer injections. Subsequently, we fitted the dissociation phase with a 1:1 binding model to obtain the mean $k_{off}$ value. Next, the mean $k_{on}$ value was determined by fitting the association phase with a 1:1 binding model, while constraining the $k_{off}$ parameter to the mean $k_{off}$ value determined in the first fitting step.

**Bivalent SPR.** A BIAcore T200 instrument (GE Healthcare Life Sciences) at 37°C and and a flow rate of 100 $\mu$l/min. Running buffer was HBS-EP. Monovalent Streptavidin-SpyCatcher was coupled to CM5 sensor chips using an amino coupling kit (GE Healthcare Life Sciences) to near saturation, typically around 7000 - 8000 response units (RU). Antigens, either biotinylated (RBD) or containing a SpyTag (CD19), were injected into the experimental flow cells (FCs) for different lengths of time to produce desired immobilisation levels (typically 20 - 70 RU). The concentration of immobilised antigen was calculated using an empirical factor to convert the immobilisation level to a molar concentration. We used the formula: molar conc. = immobilisation level / (conversion factor $\times$ molecular weight). The conversion factor was previously determined to be 149 RU per g/liter (19). Usually, FC1 was kept blank as a reference for FC2, FC3, and FC4. Excess streptavidin was blocked with two 40 s injections of 250 μM biotin (Avidity). Before

antibody injections, the chip surface was conditioned with 2 injections of the running buffer. Dilution series of antibodies were injected simultaneously in all FCs, starting with the lowest concentration. Antibodies were injected for 150s followed by a buffer injection of 450s at a flow rate of 100 µl/min. After each cycle, the chip surface is regenerated with 3M MgCl (Cytiva) for 90 s at 30 µl/min to remove all remaining bound antibodies. A buffer injection was included after every 2 or 3 antibody injections; all binding data were double-referenced by subtracting the response of the control flow cell and the closest buffer injection. Before running the high-throughput SPR experiment, we conducted an initial screen to determine whether the surface can be regenerated after each antibody injection. For this, each antibody was injected at a concentration of 50 nM with an association phase of 150 sec followed by a dissociation phase of 475 sec before injecting 3M MgCl (Cytiva) for 90 sec at a flow rate of 30 µl/min followed by a buffer injection. For the high-throughput SPR experiments, up to 32 antibodies at 3 concentrations (typically between 30 and 100 nM) each were injected in sequence over the SPR chip surface. Buffer was injected after every third cycle.

## ODE-based monovalent model

As illustrated in Fig. 1A, the monovalent ODE model assumes soluble antigen can reversibly bind to immobilised antibodies. Let [Ag] denote the concentration of antigen, [Ab] the concentration of antibodies immobilised on the SPR chip, $A$ the number of unbound antibody arms, and $B$ the number of bound antibody arms. The monovalent reaction model is then

$$A \underset{k_{\mathrm{off}}}{\overset{k_{\mathrm{on}}[\mathrm{Ag}]}{\rightleftharpoons}} B$$

with the corresponding mass action ODE model of

$$\frac{dA}{dt} = -\frac{dB}{dt} = -k_{\mathrm{on}}[\mathrm{Ag}]A + k_{\mathrm{off}}B.$$

All antibody arms are initially unbound, giving the initial condition that $A(0) = 2[\mathrm{Ab}]$ and $B(0) = 0$.

The monovalent SPR experiment is modelled by an association phase ($t = 0$ to $t = t_{\mathrm{s}}$) followed by a dissociation phase after the instantaneous removal of antigen in solution at time $t_{\mathrm{s}}$ (i.e. setting [Ag] = 0 at $t_{\mathrm{s}}$). The measured monovalent SPR response trace, $R(t)$, is proportional to $B(t)$. Analytically solving the ODE model we then have that

$$R(t) = \begin{cases} C_{\mathrm{p}} \frac{2k_{\mathrm{on}}[\mathrm{Ag}][\mathrm{Ab}]}{k_{\mathrm{on}}[\mathrm{Ag}]+k_{\mathrm{off}}} \left(1 - e^{-(k_{\mathrm{on}}[\mathrm{Ag}]+k_{\mathrm{off}})t}\right), & t \leq t_{\mathrm{s}}, \\ R(t_{\mathrm{s}}^-)e^{-k_{\mathrm{off}}(t-t_{\mathrm{s}})}, & t > t_{\mathrm{s}}, \end{cases}$$

where $C_{\mathrm{p}}$ denotes the constant of proportionality between $R(t)$ (in units of RU) and $B(t)$ (in units of concentration). The monovalent model for $R(t)$ then has three unknown parameters to fit, $C_{\mathrm{p}}$, $k_{\mathrm{on}}$, and $k_{\mathrm{off}}$.

## ODE-based bivalent model

As illustrated in Fig. 1A, the bivalent ODE model assumes soluble antibodies can reversibly bind to immobilised antigens. We let [Ab] denote the concentration of antibodies in the solution, [Ag] the concentration of antigen immobilised on the SPR chip, $A$ the concentration of these antigen that are not bound to an antibody,

$B$ the concentration of antibody-antigen complexes in which the antibody has one arm bound (i.e. is "singly-bound"), and $C$ the concentration of antibodies with both arms bound to antigen (i.e. "doubly-bound"). The well-mixed bivalent reaction model is then

$$A \underset{k_{\text{off}}}{\overset{2k_{\text{on}}[\text{Ab}]}{\rightleftharpoons}} B$$

$$A + B \underset{2k_{\text{off}}}{\overset{k_{\text{on,b}}}{\rightleftharpoons}} C$$

with the corresponding mass action ODE model of

$$\frac{dA}{dt} = -2k_{\text{on}}[\text{Ab}]A + k_{\text{off}}B - k_{\text{on,b}}AB + 2k_{\text{off}}C$$

$$\frac{dB}{dt} = 2k_{\text{on}}[\text{Ab}]A - k_{\text{off}}B - k_{\text{on,b}}AB + 2k_{\text{off}}C$$

$$\frac{dC}{dt} = k_{\text{on,b}}AB - 2k_{\text{off}}C,$$

and the initial conditions that $A(0) = [\text{Ag}]$, $B(0) = 0$, $C(0) = 0$. Note that here $k_{\text{on}}$ and $k_{\text{off}}$ represent the same physical rates used in the monovalent ODE model.

As with the monovalent SPR experiments, the bivalent SPR experiment is modelled by an association phase ($t = 0$ to $t = t_{\text{s}}$) followed by a dissociation phase after the instantaneous removal of antibodies in solution at time $t_{\text{s}}$ (i.e. setting $[\text{Ab}] = 0$ at $t_{\text{s}}$). The measured SPR response, $R(t)$, is proportional to the amount of antibodies bound to immobilised antigen, and we assume $R(t) = C_{\text{p}}(B(t) + C(t))$. We then obtain a final model with four parameters to fit, $k_{\text{on}}$, $k_{\text{off}}$, $k_{\text{on,b}}$, and a constant of proportionality $C_{\text{p}}$. Note, this model assumes that a singly-bound antibody can bind any free antigen on the surface (i.e. is 'well-mixed') and therefore, does not contain a molecular reach parameter. The model was fit to bivalent SPR data for FD-11A binding RBD (Fig. 1A-C) using *lsqcurvefit* in Matlab (Mathworks, MA).

## Particle-based bivalent model

The particle-based model modifies the bivalent ODE model by explicitly resolving the position and chemical state (i.e. free or antibody bound) of each individual antigen that is immobilised on the SPR chip. We model a small portion of the SPR chip by a cube with side lengths $L$ containing $N_{\text{Ag}}$ antigens that are uniformly (randomly) distributed.

Our model is given in terms of stochastic jump processes for the states of each individual antigen or antigen-antibody complex. Let $x_i$ denote the position of the $i$th antigen within the domain, $i = 1, \ldots, N_{\text{Ag}}$. We denote by $A_i(t) \in \{0, 1\}$ the stochastic process that is one if the $i$th antigen is not bound to any antibody, and zero otherwise. Similarly, $B_i(t) \in \{0, 1\}$ will denote the stochastic process that is one if the $i$th antigen is bound to an arm of an antibody for which the other arm is unbound, and zero otherwise. Finally, $C_{ij}(t) \in \{0, 1\}$ will denote the stochastic process that is one if the antigens at $x_i$ and $x_j$ are both bound to the same antibody, and zero otherwise.

We let $\varepsilon$ label the reach of the reaction for the free arm of a singly-bound antibody-antigen complex to bind a nearby free antigen. We assume the reaction can occur with rate $k_{\text{on,b}}$ when the two antigen involved in the reaction are separated by less than $\varepsilon$. For antigens at $x_i$ and $x_j$, the rate of the reaction is given by a

Doi interaction model (20–22) as $k_{\text{on,b}}\mathbb{1}_{[0,\varepsilon]}(x_i - x_j)$, where

$$\mathbb{1}_{[0,\varepsilon]}(x_i - x_j) = \begin{cases} 1, & |x_i - x_j|_p \leq \varepsilon, \\ 0, & |x_i - x_j|_p > \varepsilon \end{cases}$$

represents the indicator function of the interval $[0, \varepsilon]$, and $|x_i - x_j|_p$ represents the periodic distance between $x_i$ and $x_j$.

Our overall reaction model is then

$$A_i \underset{k_{\text{off}}}{\overset{2k_{\text{on}}[\text{Ab}]}{\rightleftharpoons}} B_i, \qquad\qquad i \in \{1, \ldots, N_{\text{Ag}}\}$$

$$A_i + B_j \xrightarrow{k_{\text{on,b}}\mathbb{1}_{[0,\varepsilon]}(x_i - x_j)} C_{\min\{i,j\},\max\{i,j\}}, \quad i \in \{1, \ldots, N_{\text{Ag}}\}, j \in \{1, \ldots, i-1, i+1, \ldots, N_{\text{Ag}}\},$$

$$C_{ij} \xrightarrow{k_{\text{off}}} A_i + B_j, \qquad\qquad i \in \{1, \ldots, N_{\text{Ag}}\}, j \in \{i+1, \ldots, N_{\text{Ag}}\},$$

$$C_{ij} \xrightarrow{k_{\text{off}}} A_j + B_i, \qquad\qquad i \in \{1, \ldots, N_{\text{Ag}}\}, j \in \{i+1, \ldots, N_{\text{Ag}}\},$$

where $k_{\text{on}}$ and $k_{\text{off}}$ should represent the same rates as in the monovalent and bivalent ODE models. The initial condition for each stochastic process is then $A_i(0) = 1$ for all $i$, $B_i(0) = 0$ for all $i$, and $C_{ij}(0) = 0$ for all $i$ and $j$.

The corresponding mathematical model for the evolution of the stochastic jump processes is given by Kurtz's time-change representation (23, 24). Equivalently, the probability the processes are in a given state can be described by the Chemical Master Equation (CME). Let $\mathcal{I} = \{(i,j) \mid i \neq j, i = 1, \ldots, N_{\text{Ag}}, j = 1, \ldots, N_{\text{Ag}}\}$ denote the indices of all possible $A_i$ and $B_j$ pairs, and $\hat{\mathcal{I}} = \{(i,j) \mid i = 1, \ldots, N_{\text{Ag}}, j = i + 1, \ldots, N_{\text{Ag}}\}$ denote the indices of all distinct antigen pairs. For the time-change representation each possible reaction is associated with a unit rate Poisson counting process, labelled by $\{Y_{1,i}(t)\}_{i=1}^{N_{\text{Ag}}}$, $\{Y_{2,i}(t)\}_{i=1}^{N_{\text{Ag}}}$, $\{Y_{3,i,j}(t)\}_{(i,j)\in\mathcal{I}}$, and $\{Y_{4,i,j}(t)\}_{(i,j)\in\mathcal{I}}$. We can then represent the stochastic processes for the total number of occurrences of the $A_i \to B_i$, $B_i \to A_i$, $A_i + B_j \to C_{\min\{i,j\},\max\{i,j\}}$, and $C_{\min\{i,j\},\max\{i,j\}} \to A_i + B_j$ reactions respectively as

$$\mathcal{N}_{1,i}(t) = Y_{1,i}\left(2k_{\text{on}}[\text{Ab}]\int_0^t A_i(s^-)\, ds\right), \qquad\qquad i \in \{1, \ldots, N_{\text{Ag}}\}$$

$$\mathcal{N}_{2,i}(t) = Y_{2,i}\left(k_{\text{off}}\int_0^t B_i(s^-)\, ds\right), \qquad\qquad i \in \{1, \ldots, N_{\text{Ag}}\}$$

$$\mathcal{N}_{3,i,j}(t) = Y_{3,i,j}\left(k_{\text{on,b}}\mathbb{1}_{[0,\varepsilon]}(x_i - x_j)\int_0^t A_i(s^-)B_j(s^-)\, ds\right), \qquad (i,j) \in \mathcal{I}$$

$$\mathcal{N}_{4,i,j}(t) = Y_{4,i,j}\left(k_{\text{off}}\int_0^t C_{\min\{i,j\},\max\{i,j\}}(s^-)\, ds\right), \qquad (i,j) \in \mathcal{I}.$$

Our particle model is then given by

$$A_i(t) = 1 - \mathcal{N}_{1,i}(t) + \mathcal{N}_{2,i}(t) - \sum_{j \neq i} \mathcal{N}_{3,i,j}(t) + \sum_{j \neq i} \mathcal{N}_{4,i,j}(t), \qquad i \in \{1, \ldots, N_{\text{Ag}}\}$$

$$B_i(t) = \mathcal{N}_{1,i}(t) - \mathcal{N}_{2,i}(t) - \sum_{j \neq i} \mathcal{N}_{3,j,i}(t) + \sum_{j \neq i} \mathcal{N}_{4,j,i}(t), \qquad i \in \{1, \ldots, N_{\text{Ag}}\}$$

$$C_{ij}(t) = \mathcal{N}_{3,i,j}(t) + \mathcal{N}_{3,j,i}(t) - \mathcal{N}_{4,i,j}(t) - \mathcal{N}_{4,j,i}(t), \qquad (i,j) \in \hat{\mathcal{I}}.$$

As with the ODE models, the particle-model includes an association phase from $t = 0$ to $t = t_s$, followed by a dissociation phase after the assumed instantaneous removal of all antibodies in solution at the switching time $t_s$ (i.e. setting $[\text{Ab}] = 0$ at $t_s$). The measured SPR response, $R(t)$, is proportional to the average number of antibodies bound to immobilised antigen. That is, let

$$B(t) = \sum_{j=1}^{N_{\text{Ag}}} B_j(t), \qquad\qquad C(t) = \sum_{(i,j)\in\hat{\mathcal{I}}} C_{i,j}(t)$$

be the stochastic processes for the number of singly- and doubly-bound antibodies in the system at time $t$. Denoting averages by $\mathbb{E}[\cdot]$, we assume that

$$R(t) = \frac{C_{\text{p}}}{N_{\text{Ag}}} \left( \mathbb{E}[B(t)] + \mathbb{E}[C(t)] \right),$$

where $C_{\text{p}}$ denotes the constant of proportionality converting model concentrations to experimental response units. Note, as only the antibody concentration is varied during a single bivalent SPR experiment, $N_{\text{Ag}}$ remains constant and we are simply defining a re-scaled constant of proportionality compared to the ODE cases. The resulting model then has five unknown parameters to fit, $k_{\text{on}}$, $k_{\text{off}}$, $k_{\text{on,b}}$, $\varepsilon$, and $C_{\text{p}}$.

Exact realizations of the system state, i.e. $\left( \{A_i(t)\}_{i=1}^{N_{\text{Ag}}}, \{B_i(t)\}_{i=1}^{N_{\text{Ag}}}, \{C_{ij}(t)\}_{(i,j)\in\hat{I}} \right)$, can be generated by any of the many Stochastic Simulation Algorithms (SSAs) (25, 26) (also know as Gillespie methods, Kinetic Monte Carlo methods, or Doob's method). For all forward simulations we use an optimized implementation of the Gibson-Bruck Next Reaction Method (NRM) SSA (27).

**Particle-model surrogate**

While SSA simulations of the particle model (i.e. $R(t)$) can be directly fit to bivalent SPR data, we found their computational expense to be the bottleneck in our experimental workflows. To avoid parameter estimation becoming a rate-limiting step, we developed a surrogate model that approximated the particle-model but allowed for rapid data fitting and hence, parameter estimation.

We first note that from the perspective of the particle model, the antibody solution concentration, $[\text{Ab}]$, only enters in setting an effective transition rate, i.e. probability per time, of $\hat{k}_{\text{on}} \equiv 2k_{\text{on}}[\text{Ab}]$ for an individual $A_i$ to become a $B_i$. As such, in the surrogate model we treat this as one effective parameter. We then denote by

$$\boldsymbol{\theta} = (\log_{10}(\hat{k}_{\text{on}}), \log_{10}(k_{\text{off}}), \log_{10}(k_{\text{on,b}}), \varepsilon, \log_{10}(C_{\text{p}})) \tag{1}$$

the vector of particle model parameters to estimate from SPR traces, where we have log transformed the transition rates for each possible reaction.

Let

$$R_{\text{p}}(t; \boldsymbol{\theta}) = \frac{C_{\text{p}}}{N_{\text{Ag}}} \left( \mathbb{E}[B(t)] + \mathbb{E}[C(t)] \right)$$

label the particle model's predicted SPR trace for a given set of parameters. The surrogate model is a function of $t$ and $\boldsymbol{\theta}$ defined by

$$R_{\text{s}}(t; \boldsymbol{\theta}) = C_{\text{p}} S(t, \theta_1, \theta_2, \theta_3, \theta_4), \tag{2}$$

with $S(t, \theta_1, \theta_2, \theta_3, \theta_4)$ representing the surrogate's predicated SPR response for $C_{\text{p}} = 1$. $S$ is given by a five-dimensional linear interpolation table over $(t, \theta_1, \theta_2, \theta_3, \theta_4)$. For all surrogates in this work, we fixed

$N_{\text{Ag}} = 1000$ and $[\text{Ag}_{\text{sur}}] = 125\,\mu\text{M}$. The domain length was then chosen to be consistent with this density of antigen, i.e. for $N_{\text{A}}$ denoting Avogadro's number, $L$ satisfied

$$\frac{N_{\text{Ag}}}{L^3} = [\text{Ag}_{\text{sur}}] \times N_{\text{A}} \times 10^{-30}\,\frac{\text{mol}}{(\text{nm})^3 \mu\text{M}}.$$

To construct the surrogate, $(\theta_1, \theta_2, \theta_3, \theta_4)$ were each varied over a range of values $[\underline{\theta}_i, \bar{\theta}_i]$, with samples of each $\theta_i$ uniformly spaced. That is, if parameter $\theta_i$ was sampled at $M_i$ values, they were chosen as

$$\theta_i^j = \underline{\theta}_i + j\Delta\theta_i, \quad j = 0, \ldots, M_i - 1,$$

with $\Delta\theta_i = (\bar{\theta}_i - \underline{\theta}_i)/(M_i - 1)$. The sampled reaction transition rates were then uniformly spaced in log space. We similarly partitioned time such that $t \in [0, T]$ where $T$ determined the time interval over which to fit the SPR data. We again used uniform spacing in time, $t_j = j\Delta t$, for $\Delta t = T/M_t$. For the surrogate used in this work we chose $(M_1, M_2, M_3, M_4) = (42, 40, 30, 30)$, $\Delta t = 1\text{s}$, $T = 600\text{s}$, and $M_t = 600$. $t_s$ was chosen to be 150s, consistent with the SPR protocol.

With $C_{\text{p}} = 1$, for each possible combination of parameters, $\boldsymbol{\theta} = (\theta_1^{j_1}, \theta_2^{j_2}, \theta_3^{j_3}, \theta_4^{j_4}, 1)$, we then saved the values
$$S(t_k, \theta_1^{j_1}, \theta_2^{j_2}, \theta_3^{j_3}, \theta_4^{j_4}) = R_{\text{p}}(t_k; \boldsymbol{\theta}), \quad k = 0, \ldots, M_t$$

in a five-dimensional lookup table. Estimating $R_{\text{p}}(t_k; \boldsymbol{\theta})$ requires averaging SSA simulations of the particle model. Let $B^n(t)$ and $C^n(t)$ denote the sampled values of $B(t)$ and $C(t)$ in the $n$th simulation, $n = 1, \ldots, N_{\text{sims}}$, with

$$R_{\text{p}}^n(t_k; \boldsymbol{\theta}) = \frac{1}{N_{\text{Ag}}}(B^n(t_k) + C^n(t_k))$$

the corresponding response. We then approximated

$$R_{\text{p}}(t_k; \boldsymbol{\theta}) \approx \frac{1}{N_{\text{sims}}} \sum_{n=1}^{N_{\text{sims}}} R_{\text{p}}^n(t_k; \boldsymbol{\theta}).$$

Our general protocol was to average over at least $N_{\text{sims}} = 15$ SSA samples, continuing to add SSA samples until either the estimated standard error of the samples, $\{R_{\text{p}}^n(t_k; \boldsymbol{\theta})\}_{n=1}^{N_{\text{sims}}}$, was below .01 of the sample mean at all $t_k$, or $N_{\text{sims}} = 250$ SSA samples were reached.

Given the table $\{S(t_k, \theta_1^{j_1}, \theta_2^{j_2}, \theta_3^{j_3}, \theta_4^{j_4})\}_{(k,j_1,j_2,j_3,j_4)=(1,1,1,1,1)}^{(M_t+1,M_1,M_2,M_3,M_4)}$, $S(t, \theta_1, \theta_2, \theta_3, \theta_4)$ for $t \in (0, T)$ and all $\theta_j \in (\underline{\theta}_j, \bar{\theta}_j)$ could then be evaluated by linear interpolation of the bracketing tabulated values. In practice, we evaluated the surrogate at general time and parameter values using the `BSpline(Linear())` option from the Interpolations.jl library [28]. For the sizes we used, $(M_t+1, M_1, M_2, M_3, M_4) = (601, 42, 40, 30, 30)$, the table contained model responses for 1,512,000 parameter combinations, and required approximately 7.3GB of memory to store. Surrogates were generally constructed in a few hours using 500-2000 cores on the Boston University Shared Computing Cluster.

**Particle-model data fitting**

Estimates for the log transformed parameters $\boldsymbol{\theta}$, defined in Eq. [1], were generated via minimization of the squared error between surrogate predictions and experimental SPR responses for varying levels of [Ab]. Suppose $I$ SPR traces are being simultaneously fit (e.g. traces with different antibody concentrations injected over the same surface), with $R^{(i)}(t)$ labelling the $i$th SPR response trace for an experiment with

antibody concentration $[\text{Ab}^{(i)}]$. We assume a fixed and known antigen concentration, $[\text{Ag}]$, when simultaneously fitting multiple SPR traces generated using different antibody concentrations injected over the same surface. The experimental traces are ordered such that

$$[\text{Ab}^{(1)}] \leq [\text{Ab}^{(2)}] \leq \cdots \leq [\text{Ab}^{(I)}].$$

Finally, given a current estimate for $\boldsymbol{\theta}$ arising during optimization, we define

$$\boldsymbol{\theta}^{(i)} \equiv \left( \theta_1 + \log_{10}\left( \frac{[\text{Ab}^{(i)}]}{[\text{Ab}^{(1)}]} \right), \theta_2, \theta_3, \theta_4, \theta_5 \right)$$

$$= \left( \log_{10}\left( \hat{k}_{\text{on}} \frac{[\text{Ab}^{(i)}]}{[\text{Ab}^{(1)}]} \right), \log_{10}(k_{\text{off}}), \log_{10}(k_{\text{on,b}}), \varepsilon, \log_{10}(C_{\text{p}}) \right).$$

The overall loss function we then minimized was

$$L(\boldsymbol{\theta}) = \sum_{i=1}^{I} \sum_{k=0}^{M_t} \left( R^{(i)}(t_k) - R_{\text{s}}(t_k; \boldsymbol{\theta}^{(i)}) \right)^2,$$

where $R_{\text{s}}(t; \boldsymbol{\theta})$ is the surrogate response defined by Eq. [2]. In practice we minimized this loss using the XNES natural evolution optimizer from BlackBoxOptim.jl via the Optimization.jl meta-package (29, 30). All optimization related parameters were left at their default values except the maximum number of iterations, which was increased to 5000. As XNES is a stochastic optimizer, we generally applied it several times and selected the estimated parameter set across all runs having the minimal loss as the consensus estimate. For more details, see the section on "Data analysis for high-throughput bivalent SPR".

As described in the previous section, the surrogate was constructed for a fixed concentration of antigen. To avoid producing a new surrogate for each experiment, we conjectured that once our particle model system was of sufficient size (i.e. sufficiently large number of antigens for a fixed antigen concentration), the antigen concentration effectively set the average number of antigens that were within reach. In this way, we could fit SPR data generated with any antigen concentration using a single surrogate (produced with a single antigen concentration) but would need to transform the fitted reach based on the experimental antigen concentration. In other words, the biophysical reach ($\varepsilon_{\text{phys}}$) can be calculated from the fitted reach ($\varepsilon_{\text{sur}} = \theta_4$) by enforcing that the average number of antigens within reach in the surrogate model and in the SPR experiment are the same,

$$\frac{4}{3}\pi \varepsilon_{\text{sur}}^3 [\text{Ag}_{\text{sur}}] = \frac{4}{3}\pi \varepsilon_{\text{phys}}^3 [\text{Ag}].$$

We empirically confirmed that the model produced the same predicted SPR traces at different antigen densities ($[\text{Ag}]=1$, 10, and 100 $\mu$M) provided that the physical molecular reach ($\varepsilon_{\text{phys}}$) was decreased according to the above equation (Fig. S1). Therefore, the relationship between the fitted parameters $\boldsymbol{\theta}$ and the biophysical parameters are as follows,

$$k_{\text{on}} = \frac{10^{\theta_1}}{[\text{Ab}^{(1)}]} = \frac{\hat{k}_{\text{on}}}{[\text{Ab}^{(1)}]}, \quad k_{\text{off}} = 10^{\theta_2}, \quad k_{\text{on,b}} = 10^{\theta_3},$$

$$\varepsilon_{\text{phys}} = \theta_4 \left( \frac{[\text{Ag}_{\text{sur}}]}{[\text{Ag}]} \right)^{1/3}, \quad C_{\text{p}} = 10^{\theta_5}.$$

[3]

These represent the final biophysical parameter estimates reported in this work.

**Particle-model antibody binding potency predictions**

To predict the concentration of antibody required to bind 50% of antigen (antibody binding potency), the particle model is simulated using the previously mentioned Next Reaction Method-based approach. The only modification to the model is that we used a 2D square with sides of length $L$ to represent the 2D viral surface. As before, antigens are assumed to be uniformly (randomly) distributed in a 2D region with $N_{\mathrm{Ag}}$ set to 1000 and $L$ chosen to enforce a specified antigen density.

For each antibody, the model was simulated with its fitted binding parameters ($k_{\mathrm{on}}$, $k_{\mathrm{off}}$, $k_{\mathrm{on,b}}$, and the molecular reach $\varepsilon$) for different antigen and antibody concentrations to a time of 60 minutes. For each antibody concentration we averaged 20 independent simulations to estimate the average number of unbound antigens at 60 minutes. The resulting dose-response curves for the fraction of unbound antigen ($\mathrm{Ag_{unbound}}$) versus antibody concentration ($[\mathrm{Ab}]$) were fit with an inhibitory Hill model ($n < 0$) to determine $\mathrm{IC_{50}}$,

$$\mathrm{Ag_{unbound}} = \frac{1}{1 + \left(\dfrac{IC_{50}}{[\mathrm{Ab}]}\right)^n}. \qquad [4]$$

**Particle-model software**

Codes for particle model forward simulation, surrogate construction, and fitting are available in our MIT-licensed Julia library (31).

**Worm-like-chain (WLC) model to estimate size of PEG-coupled antigen**

The worm-like-chain (WLC) is a widely used polymer model that has previously been applied to PEG polymers (12). The model provides an estimate for the mean end-to-end distance of the polymer as follows: $2\sqrt{N_{PEG}l_c l_p}$ where $N_{PEG}$ is the number of PEG repeats, $l_c$ is the contour length of each repeat (0.4 nm), and $l_p$ is the persistence length of PEG (0.4 nm (12)). The WLC model predicts a mean length of 0.69 nm for PEG3 and 2.1 nm for PEG28 or a difference in length of 1.41 nm. The predicted increase in molecular reach when an antibody binds PEG28 instead of PEG3 would then be twice this difference to account for the two bound antigens involved in antibody binding (2.82 nm, Fig 2C).

**Molecular dynamics**

**Construction of full-length all-atom antibody/RBD complex structures.** The PDB structure (1HZH) of a full-length IgG1 antibody was rebuilt and minimized in CHARMM35 (32) based on SEQRES records to serve as a template for rebuilding of full-length antibodies specific for RBD. The crystal structures of the Fab/RBD complexes used are listed in Table S1. Each RBD structure was rebuilt to contain all residues of the N-terminal AviTag and signal sequence (24 residues total after cleavage) used in SPR experiments fused to RBD residues 331-526 (196 residues) resulting in a final construct of 220 residues in all cases. Any missing residues in the light or heavy chains of the Fabs were also rebuilt and all rebuilt sections of Fabs and RBD minimized in CHARMM35. Two copies of each Fab/RBD complex were then aligned to the 1HZH all-atom model in Chimera v1.16 (33) using the MatchMaker tool based on the heavy and light

chain only, with the relative orientations of the RBD and Fab structures maintained. Heavy chains from these aligned structures were then merged with those of 1HZH based on MUSCLE (34) alignments of their sequences and inspection of the aligned structures. The three residues on either side of the new bond were minimized in CHARMM35 with all other atomic positions held fixed. The final structure after 1,000 steps of unconstrained minimization *in vacuo* in Amber FF14SB (35) with OpenMM v7.5 (36) was accepted as the final all-atom structure. We note that 1HZH contains one disulfide bond between the two heavy chains in the hinge region rather than the usual two. We chose not rebuild the second disulfide bond so as not to distort the overall structure and because the disulfide that limits the extension of the hinge is in place. The lengths of all chains within each antibody in the rebuilt full-length models are listed in Table S2.

**Construction of topology-based coarse-grain models.** Coarse-grain models were parameterized following a previously published protocol that represents each amino acid as a single interaction site centered at the $C_\alpha$ coordinates of each atom. The potential energy of a conformation within this model is given by

$$
E = \sum_i k_{\mathrm{b}} (r_i - r_0)^2 + \sum_i \sum_j^4 k_{\varphi_{ij}} (1 + \cos[j\varphi_i - \delta_{ij}]) + \sum_i -\frac{1}{\gamma} \ln\left\{\exp\left[-\gamma\left(k_\alpha(\theta_i - \theta_\alpha)^2 + \varepsilon_\alpha\right)\right]\right.
$$

$$
+ \exp\left[-\gamma k_\beta(\theta_i - \theta_\beta)^2\right]\bigg\} + \sum_{ij} \frac{q_i q_j e^2}{4\pi\varepsilon_0 \varepsilon_r r_{ij}} \exp\left[-\frac{r_{ij}}{l_{\mathrm{D}}}\right] + \sum_{ij\in\{\mathrm{NC}\}} \epsilon_{ij}^{\mathrm{NC}} \left[13\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - 18\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{10}\right.
$$

$$
\left. + 4\left(\frac{\sigma_{ij}}{r_{ij}}\right)^6\right] + \sum_{ij\notin\{\mathrm{NC}\}} \epsilon_{ij}^{\mathrm{NN}} \left[13\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{12} - 18\left(\frac{\sigma_{ij}}{r_{ij}}\right)^{10} + 4\left(\frac{\sigma_{ij}}{r_{ij}}\right)^6\right]. \quad [5]
$$

These forcefield terms, which represent contributions from bonds, dihedrals, angles, electrostatics, native contacts, and non-native contacts, have been described in detail previously (37). The strength of native contacts within this coarse-grain model is determined by the Lennard-Jones-like well depths for the native contact term. The well depth, $\epsilon_{ij}^{NC}$, is computed as $\epsilon_{ij}^{NC} = \eta_{ij}\epsilon_{HB} + \eta\epsilon_{ij}$, in which $\eta_{ij}$ is the number of hydrogen bonds between residues $i$ and $j$, $\epsilon_{HB} = 0.75$ kcal/mol, $\epsilon_{ij}$ is the well-depth taken from the Betancourt-Thirumalai (38) pairwise potential for a contact between residues of types $i$ and $j$, and $\eta$ is a scaling factor that increases the effective well depths. Increasing the value of $\eta$ linearly increases the stability of all native contacts to which it is applied. Values of $\eta$ were selected individually for each domain and interface within the antibody-RBD complexes. We note that disulphide bonds are treated as harmonic bonds with $k_b = 20$ kcal/(mol×Å$^2$) rather than as native contacts.

**Selection of $\eta$ values for antibody domains and interfaces.** All coarse-grain simulations were performed in OpenMM v7.5 using the LangevinMiddleIntegrator (39) with an integration time step of 15 fs, a friction coefficient of 0.050 ps$^{-1}$, and an absolute temperature of 298 K. Values of $\eta$ were selected based on a previously published training set (40). To limit the number of domains and interfaces that required simultaneous parameter tuning, we used the same set of $\eta$ values for all domains and interfaces within each antibody and for each RBD monomer (Table S3). The intra-domain values of $\eta$ were taken in each case as the largest value from a training set (see (41) Table S5 ). Likewise, the largest training set value for interfaces was used for the HC1—LC1, HC2—LC2, and HC1:HC2 interfaces. The HC1—LC2 interface, which likely represents a crystal packing interface, was assigned the training set average value of 1.507. Simulations of FD-11A in isolation were performed using the values of $\eta$ in Table S3 but with the RBD representations deleted.

**Coarse-grain steered molecular dynamics simulations of antibody molecular reach.** Initial structures for steered molecular dynamics were generated by aligning the coarse-grained RBD-antibody-RBD

complexes such that one RBD Lys15 (i.e., the residue that is biotinylated and bound to streptavidin in the experiment) was at the coordinate system origin and the other on the positive x-axis. A spherical harmonic restraint with force constant 50 kcal/(mol×Å$^2$) was applied to the Lys15 at the origin to hold it in place during pulling. A second spherical harmonic restraint with force constant 1 kcal/(mol×Å$^2$) was applied to the second Lys15 residues to serve as the trap for steered molecular dynamics. Flat-bottom Root Mean Square Deviation (RMSD) restraints with force constant $k_{\text{RMSD}} = 50$ kcal/(mol×Å$^2$) and RMSD threshold of 5 Å were applied to residues in Fab1, Fab2, the Fc, as well as the globular portions of each RBD monomer to prevent overall unfolding while allowing structural fluctuations. Only the RBD linkers, antibody hinge, and epitope—paratope interfaces were left unrestrained. Each trajectory was then equilibrated for 1.5 ns to allow the coordinates to randomize. After equilibration, the harmonic restraint on the second Lys15 residue was pulled with a constant velocity of 1 nm/ns along the positive x-axis to a total displacement of +30 nm from its original position over the course of 30 ns. A total of 50 statistically independent trajectories were run for each of the six antibodies and each of seven different epitope—paratope interface $\eta$ values, $\eta = 1.000, 1.250, 1.500, 1.750, 2.000, 2.250, 2.500$ for a total of 2,100 statistically independent trajectories (total simulation time 66 μs). The molecular reach was computed from each trajectory as longest distance at which both epitope—paratope interfaces have a non-zero fraction of native contacts, Q. Larger values of $\eta$ sometimes result in unfolding of RBD or IgG domains during pulling due to the increased interaction strength at the epitope—paratope interfaces. We therefore applied a filter to discard trajectories that ever have a Fab or RBD RMSD > 10 Å. Varying this threshold value to 8 or 15 Å was not found to strongly influence results. The mean maximum Lys15-Lys15 distance was then computed by averaging the value of those trajectories that remain reasonably well folded. The molecular reach estimate from the simulations was taken to be the maximum mean value across all $\eta$ values.

## SARS-CoV-2 neutralisation

Neutralisation IC$_{50}$ for the FD-11A antibody was measured with a microneutralization assay as previously described (42). Two doses of SARS-CoV-2 vaccination induce robust immune responses to emerging SARS-CoV-2 variants of concern. FD-11A was preincubated with SARS-CoV-2 for 60 min at room temperature before being added to Vero CCL-81 cells. Level of infection was measured by counting the number of infectious foci.

For anti-RBD antibodies used in the high-throughput SPR experiment, neutralisation IC$_{50}$ was measured using a Focus Reduction Neutralization Test as described in (4). Briefly, serially diluted antibody was incubated with SARS- CoV-2 for 1 hr at 37ºC, afterwards transfered to Vero cell monolayers. The level of infection was measured using a focus forming assay.

## Data analysis for high-throughput bivalent SPR

In this section we explain the detailed workflow for analysis of the high-throughput bivalent SPR experiments (Fig. 4).

After double referencing the SPR curves for each antibody concentration, we aligned them to the start of the dissociation phase (setting it to $t_s = 150$s) to improve curve alignment. To eliminate artefacts arising from the start and end of the association phase (generally large spikes in RU arising from needle motion), the first 5 seconds of the association phase, the last 4 seconds of the association phase, and the first 5

seconds of the dissociation phase were excluded from the data. We also excluded all SPR curves where the maximum response across the entire injection was smaller than 6 RU because this minimal binding is within the systematic error for SPR experiments (typically 1 RU per 100 seconds of injection or 6 RU for our 600 second experiments). Finally, to produce more manageable file sizes we reduced the temporal resolution from 10 Hz to 1 Hz and this did not impact our results because the kinetics of the SPR traces were much slower than 1 Hz.

We then fitted the processed data for each antibody using the bivalent Particle-based model (see Particle-model data fitting section). The antigen and antibody concentrations for each SPR curve were provided as input parameters for the fitting process. The surrogate model used in fitting contained the following parameter ranges: $\log_{10}(k_{on}) \in [-5.0, 2.0]$ with $M_1 = 42$, $\log_{10}(k_{off}) \in [-4.0, 0.0]$ with $M_2 = 40$, $\log_{10}(k_{on,b}) \in [-3.0, 1.5]$ with $M_3 = 30$, and $\varepsilon \in [2, 35]$ with $M_4 = 30$. During fitting of the surrogate to SPR data, a box constraint that $\log_{10}(C_p) \in [1.0, 4.0]$ was used. The fitting process was repeated 100 times, and the parameters yielding the lowest fitness was recorded. The estimated bivalent model parameters from the fitting process were converted into the corresponding biophysical parameters according to Eq. [3] using the experimental antigen concentration. The bivalent SPR data was also fit with the ODE-based monovalent model (see ODE-based monovalent model section).

Finally, a quality control procedure was introduced to ensure the accuracy of the bivalent binding parameters. First, we checked that the antigen surface could be regenerated after each antibody injection (7 out of 80 antibodies could not be removed). Second, we only included data where the particle-model produces a close fit to the data (12 out of 80 antibodies could not be fit). Third, we only included data where the particle-model produced a fit that was better than the ODE-based monovalent model. We reasoned that SPR data that could accurately be fit by the ODE-based monovalent model did not contain information that could accurately determine bivalent binding (bivalent binding parameters of 16 out of 80 antibodies could not be determined).

Bivalent SPR data that passed all quality control measures for each antibody were averaged across SPR experiments. We report the geometric mean for parameters that varied on a logarithmic range in the model ($k_{on}$, $k_{off}$, $K_D$, $k_{on,b}$) and the mean for the molecular reach that varied on a linear range in the model.

## Statistical analysis

**Multiple linear regression** We fit a one way multiple linear model in Prism (v 9.5.1) with the following formula:

$$y = \beta_0 + \beta_1 x_1 + ... + \beta_n x_n$$

where $y$ is the predicted neutralisation $IC_{50}$ for each antibody, $\beta_0$ is the y-intercept to fit, $(x_1, \ldots, x_n)$ are the values for $\log_{10}(k_{on})$, $\log_{10}(k_{off})$, $\log_{10}(K_D)$, $\log_{10}(k_{on,b})$, $\varepsilon$, and the blocking epitope distance respectively, and $(\beta_1, \ldots, \beta_n)$ are the corresponding regression coefficients for each variable. Models contain either all variables, or a subset. A Least Squares regression type was used and models were compared using an Extra-Sum-of-Squares F test.

# References

1. Klein JS, Bjorkman PJ (2010) Few and Far Between: How HIV May Be Evading Antibody Avidity. *PLOS Pathogens* 6:e1000908.

2. Edeling MA, et al. (2014) Potent Dengue Virus Neutralization by a Therapeutic Antibody with Low Monovalent Affinity Requires Bivalent Engagement. *PLOS Pathogens* 10:e1004072.

3. Yan R, et al. (2021) Structural basis for bivalent binding and inhibition of SARS-CoV-2 infection by human potent neutralizing antibodies. *Cell Research 2021 31:5* 31:517–525.

4. Dejnirattisai W, et al. (2021) The antigenic anatomy of SARS-CoV-2 receptor binding domain. *Cell* 184:2183–2200.

5. Alanine DG, et al. (2019) Human Antibodies that Slow Erythrocyte Invasion Potentiate Malaria-Neutralizing Antibodies. *Cell* 178:216–228.

6. Autheman D, et al. (2021) An invariant Trypanosoma vivax vaccine antigen induces protective immunity. *Nature 2021 595:7865* 595:96–100.

7. Myszka DG (1999) Improving biosensor analysis. *Journal of Molecular Recognition* 12:279–284.

8. Huang KYA, et al. (2021) Breadth and function of antibody response to acute SARS-CoV-2 infection in humans. *PLOS Pathogens* 17:e1009352.

9. Shaw A, et al. (2019) Binding to nanopatterned antigens is dominated by the spatial tolerance of antibodies. *Nature Nanotechnology* 14:184–190.

10. Zhang P, et al. (2020) Capturing transient antibody conformations with DNA origami epitopes. *Nature Communications* 11:1–9.

11. Jendroszek A, Kjaergaard M (2021) Nanoscale spatial dependence of avidity in an IgG1 antibody. *Scientific Reports 2021 11:1* 11:1–10.

12. Kienberger F, et al. (2000) Static and Dynamical Properties of Single Poly(Ethylene Glycol) Molecules Investigated by Force Spectroscopy. *Single Mol* 1:123–128.

13. Barton MI, et al. (2021) Effects of common mutations in the sars-cov-2 spike rbd and its ligand the human ace2 receptor on binding affinity and kinetics. *eLife* 10.

14. Ke Z, et al. (2020) Structures and distributions of SARS-CoV-2 spike proteins on intact virions. *Nature 2020 588:7838* 588:498–502.

15. Klein S, et al. (2020) SARS-CoV-2 structure and replication characterized by in situ cryo-electron tomography. *Nature Communications 2020 11:1* 11:1–10.

16. Denham EM, et al. (2019) A generic cell surface ligand system for studying cell–cell recognition. *PLoS Biology* 17:1–30.

17. Huang KYA, et al. (2023) Structural basis for a conserved neutralization epitope on the receptor-binding domain of SARS-CoV-2. *Nature Communications 2023 14:1* 14:1–13.

18. Patel A, et al. (2023) Using CombiCells, a platform enabling titration and combinatorial display of cell surface ligands, to investigate the sensitivity and costimulatory requirements of TCRs and CARs. *bioRxiv* p 2023.06.15.545075.

19. Goyette J, et al. (2017) Biophysical assay for tethered signaling reactions reveals tether-controlled activity for the phosphatase SHP-1. *Science Advances* 3:e1601692.

20. Doi M (1976) Second quantization representation for classical many-particle system. *J. Phys. A: Math. Gen.* 9:1465–1477.

21. Doi M (1976) Stochastic theory of diffusion-controlled reaction. *J. Phys. A: Math. Gen.* 9:1479–1495.

22. Erban R, Chapman SJ (2009) Stochastic modelling of reaction-diffusion processes: algorithms for bimolecular reactions. *Phys. Biol.* 6:046001.

23. Kurtz TG (1978) Strong approximation theorems for density dependent markov chains. *Stochastic Processes and their Applications* 6:223–240.

24. Anderson DF, Kurtz TG (2015) *Stochastic Analysis of Biochemical Systems*, Mathematical Biosciences Institute Lecture Series (Springer).

25. Gillespie DT (1976) A general method for numerically simulating the stochastic time evolution of coupled chemical reactions. *Journal of Computational Physics* 22:403–434.

26. Marchetti L, Priami C, Thanh VH (2017) *Simulation Algorithms for Computational Systems Biology*, Texts in Theoretical Computer Science (Springer).

27. Gibson MA, Bruck J (2000) Efficient exact stochastic simulation of chemical systems with many species and many channels. *J. Phys. Chem. A* 104:1876–1899.

28. Lycken T, Holy T, Kittisopikul M, Contributors (2021) Interpolations.jl (version 0.13.5). (https://github.com/JuliaMath/Interpolations.jl).

29. Feldt R, Contributors (2021) BlackBoxOptim.jl (version 0.6.1). (https://github.com/robertfeldt/BlackBoxOptim.jl).

30. Dixit VK, Rackauckas CV, Contributors (2022) Optimization.jl (version 3.7.1). (https://github.com/SciML/Optimization.jl).

31. (2023) SPRFittingPaper2023.jl (version 0.1.0). (https://github.com/isaacsas/SPRFittingPaper2023).

32. Brooks BR, et al. (2009) CHARMM: The Biomolecular Simulation Program. *Journal of computational chemistry* 30:1545.

33. Pettersen EF, et al. (2004) UCSF Chimera–a visualization system for exploratory research and analysis. *Journal of computational chemistry* 25:1605–1612.

34. Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* 32:1792–1797.

35. Maier JA, et al. (2015) ff14SB: Improving the accuracy of protein side chain and backbone parameters from ff99SB TOC figure HHS Public Access. *J Chem Theory Comput* 11:3696–3713.

36. Eastman P, et al. (2017) OpenMM 7: Rapid development of high performance algorithms for molecular dynamics. *PLoS computational biology* 13.

37. Nissley DA, et al. (2022) Universal protein misfolding intermediates can bypass the proteostasis network and remain soluble and less functional. *Nature Communications 2022 13:1* 13:1–16.

38. Betancourt MR, Thirumalai D  (1999) Pair potentials for protein folding: choice of reference states and sensitivity of predicted native states to variations in the interaction schemes. *Protein Science : A Publication of the Protein Society* 8:361.

39. Zhang Z, Liu X, Yan K, Tuckerman ME, Liu J  (2019) Unified Efficient Thermostat Scheme for the Canonical Ensemble with Holonomic or Isokinetic Constraints via Molecular Dynamics. *The journal of physical chemistry. A* 123:6056–6079.

40. Leininger SE, Trovato F, Nissley DA, O'Brien EP  (2019) Domain topology, stability, and translation speed determine mechanical force generation on the ribosome. *Proceedings of the National Academy of Sciences of the United States of America* 116:5523–5532.

41. Nissley DA, et al.  (2020) Electrostatic Interactions Govern Extreme Nascent Protein Ejection Times from Ribosomes and Can Delay Ribosome Recycling.  *Journal of the American Chemical Society* 142:6103–6110.

42. Skelly DT, et al.  (2021) Two doses of SARS-CoV-2 vaccination induce robust immune responses to emerging SARS-CoV-2 variants of concern. *Nature Communications 2021 12:1* 12:1–12.

# Supplementary Information
# The molecular reach of antibodies
# determines their SARS-CoV-2 neutralisation potency

Anna Huhn[1], Daniel Nissley[2], Daniel B. Wilson[1,3,||], Mikhail Kutuzov[1], Robert Donat[4],
Tiong Kit Tan[4], Ying Zhang[3,ɟ], Michael I. Barton[1], Chang Liu[5,6], Wanwisa Dejnirattisai[5,7],
Piyada Supasa[5], Juthathip Mongkolsapaya[5], Alain Townsend[4,6] William James[8], Gavin Screaton[5],
P. Anton van der Merwe[1], Charlotte M. Deane[2,¶], Samuel A. Isaacson[3,†,¶], Omer Dushek[1†,¶]

Figure S1: **Workflow for fitting the particle-model to bivalent SPR data. (A)** A schematic of the particle-model chemical reactions (left) used to perform simulations of the association and dissociation phase of antibody binding in bivalent SPR (middle, red traces), which are averaged to produce the predicted SPR trace for a given parameter set (middle, black traces). These averaged SPR traces are pre-tabulated for different parameter values ($k_{on}$, $k_{off}$, $k_{on,b}$, reach - $\varepsilon$) using a computing cluster to produce a surrogate for the particle-model that is used in data fitting (right). **(B)** The average number of antigens within reach of an individual antigen depends on the molecular reach and the antigen density. **(C)** The relationship between antigen density and molecular reach when the average number of antigens within reach is 0.02 (equation plotted is from panel b). **(D)** The spatial distribution of antigen (top) and the corresponding predicted bivalent SPR traces (bottom) for different antigen concentrations and molecular reach calculated using the relationship in panel B. The SPR traces are effectively identical as shown by the small residuals (difference between the indicated SPR curves) confirming that antibody binding depends on the average number of antigens within reach, which can be achieved by a short reach at high antigen density or a long reach at low antigen density. Simulations are performed using an antibody concentration of 1 nM with $k_{on} = 0.05\ \mu$M$^{-1}$s$^{-1}$, $k_{off} = 0.02$s$^{-1}$, and $k_{on,b} = 1.0^{-1}$.

Figure S2: **The fitted binding parameters for the IgG FD-11A antibody binding RBD determined by fitting the particle-model to bivalent SPR data are independent of the RBD antigen density.** The indicated parameter is plotted over the RBD antigen density ($N = 42$). The coefficient of determination ($R^2$) and the p-value for the null hypothesis that the fitted line and a horizontal line (i.e. no relationship between the binding parameter and RBD density) produce an equal fit to the data.

Figure S3: **Representative monovalent and bivalent SPR for the indicated IgG1 antibodies binding RBD. (A)** Representative monovalent SPR traces produced by injecting RBD (2000 nM with 2-fold dilutions) over surfaces immobilised with the indicated antibody. **(B)** Representative bivalent SPR traces produced by injecting the indicated antibodies over surfaces immobilised with RBD. Antibodies were injected using a 2-fold dilution series, with a top concentration of 300 nM, 5 nM, 5 nM, 5 nM, 5 nM for FD-5D, FI-3A, CR3022, EY-6A and REGN10987 respectively. RBD was immobilised at 7 - 15 $\mu$M.

Figure S4: **The particle-model accurately analyses bivalent SPR generated using the CD19 antigen reproducing the monovalent binding parameters.** Data is generated using purified Spytag-CD19 (antigen) and an anti-CD19 antibody (SJ25C1). **(A)** Monovalent SPR (top) is produced by injecting Spytag-CD19 over a surface immobilised with SJ25C1 and bivalent SPR (bottom) is produced by injecting SJ25C1 over a surface immobilised with Spytag-CD19 (through coupling to amine-coupled Spycatcher). The SPR traces are generated by 2-fold dilution of CD19 starting at 175 nM (top) or by a 2-fold dilution of SJ25C1 starting at 33 nM (bottom). Representative SPR traces (black) and model fit (red) are shown for 1 out of 2 representative experiments. **(B)** The indicated parameter produced by monovalent or bivalent SPR. **(C)** The indicated parameter produced by bivalent SPR.

Figure S5: **The fitted binding parameters of an anti-phosphotorysine antibody (PY20) interacting with a small phosphorylated peptide antigen coupled to PEG3 or PEG28 linker.** The indicated parameter from N=9 or N=5 independent experiments for PEG3 and PEG28 coupled phosphorylated peptide antigen, respectively. A t-test with a Holm-Šídák correction for multiple comparison was used to determine p-values.

Figure S6: **The fitted binding parameters for an anti-phosphotorysine antibody (PY20) interacting with different densities of a small phosphorylated peptide antigen coupled to PEG3.** **(A)** Representative SPR traces (black) and particle-based model fits (red) for the PY20 antibody injected over surfaces with the indicated concentration of PEG3 coupled to a small phosphorylated peptide antigen. The antibody was injected at 8 different concentrations using a 2-fold dilution from a top concentration of 25 nM. **(B)** The fitted binding parameters plotted over the density of the small phosphorylated peptide antigen coupled to the SPR chip surface.

Figure S7: **The mean distance between the antigen-binding domains of the IgG1 FD-11A antibody is 13 nm.** The FD-11A Fab structure is used to generate a coarse-grain model of the complete IgG1 FD-11A antibody (see methods) and is used in coarse-grained MD simulations. **(A)** Ensemble average time series (dark purple) computed over five independent 1-$\mu$s trajectories of the distance between the centres-of-mass of the coarse-grain interaction sites that form the paratopes on each Fab binding arm (i.e. the distance between the antigen binding domains). The light purple shaded region indicates the standard error over the five trajectories at each simulation frame. The dotted black line indicates the distance between the antigen-binding sites in the structure. **(B)** Histogram of the inter-paratope or inter-antigen binding domain distances over each of the five independent trajectories. **(C)** Block averaging analysis of the merged trajectory indicates a block size of 1,000 frames is suitable. The mean inter-paratope distance is 13.05 ± 0.05 nm (error bar is estimated as the block standard error with a block size of 1000 frames). Block averaging was performed on a merged trajectory consisting of the final 975 ns of the five independent runs. **(D)** Coarse-grain structures of the reference structure (left), minimum reach structure from simulations (middle), and maximum reach structure from simulations (right). The two heavy chains are colored dark and light blue, the two light chains are colored red and pink, and the interactions sites constituting the paratopes are colored yellow. The dotted black line in the leftmost structure indicates the distance between the centers-of-mass of the paratopes in the reference state (14.94 nm).

Figure S8: **Estimating the molecular reach of RBD-specific antibodies using coarse-grained molecular dynamic simulations.** **(A)** A representative coarse-grained MD simulation trajectory for the indiciated antibody (columns) showing the calculated force along the pulling direction ($F_x$), the fraction of native contacts for each antigen interface (Q, interface 1 in pink and 2 in mauve), the RMSD as a function of time during the simulation for Fabs 1 and 2 (blue and green, respectively), and the RMSD of RBD 1 and 2 (yellow and orange, respectively). **(B)** The structure from the trajectories in panel A that produced the maximum distance between the Lys15 residues on RBD (shown below the structure) whilst the antibody was bound bivalently. This structure was achieved at the time point indicated by the vertical magenta lines in panel A. **(C)** The maximum Lys15-Lys15 distance over the interface binding strength ($\eta$) calculated over the set of trajectories where unfolding does not take place during the MD simulations. Error bars are 95% confidence intervals computed from bootstrapping with $10^6$ independent samples. Missing data points for CR3022 and FI3A indicate that all trajectories were unfolded. The molecular reach is defined as the maximum Lys15-Lys15 distance over all $\eta$ and indicated by the dashed horizontal line.

Figure S9: **Quality controls for high throughput analysis of 80 RBD antibodies using bivalent SPR. (A-B)** Surface stability of RBD was assessed by injecting FD-11A at the beginning and end of each experiment. Representative SPR traces of the first and last injection of FD-11A (A) and fraction of FD-11A binding at end of the experiment (B). The fraction is calculated as the RU at 150 s after injection of highest concentration in the first divided by the last injection. **(C)** A summary of the antibodies that passed or failed quality control and hence included or excluded from the analysis, respectively. **(D-G)** Examples of antibodies representing the four possible quality control outcomes. (D) The antibody 170 was included because the particle-model produced a good fit and the monovalent model produced a poor fit. (E) The antibody 170 displays complete regeneration (included) whereas antibody 280 shows only partial regeneration (excluded). Partial regeneration can be observed by residual RU after the injection of 3 M MgCl₂ for 90 s at the end of each SPR cycle (see arrow). (F) The antibody 264 was excluded because the particle-model produced a poor fit. (G) The antibody 150 was excluded because the particle-based model (left) and the ODE-based monovalent model (right) produced an equally good fit.

Figure S 10: **The previously introduced epitope taxonomy for RBD-binding antibodies does not stratify the molecular reach.** **(A)** The previously introduced epitope taxonomy for RBD antibodies based on five regions. Figure adapted from (4). **(B-D)** Antibody neutralisation $IC_{50}$ (B), epitope blocking distance (C), and molecular reach of each antibody (D) organised by their taxonomic class.

**Including all measured antibodies**

**Including antibodies with a epitope blocking distance < 2.37 nm**

Figure S11: **The predicted antibody binding potency produces absolute agreement with the experimental neutralsiation potency at intermediate antigen densities.** The particle-based model was used to predict the concentration of antibody required to bind 50% of antigen (Predicted $IC_{50}$) to surfaces randomly distributed with the indicated density of antigen for **(A-I)** all antibodies or **(K-S)** the subset of antibodies that bound within 2.37 nm of the blocking epitope. A linear fit to the log-transformed $IC_{50}$ values (solid line) is compared to a line of absolute agreement (dashed black line - slope 1, y-intercept 0) using an F-test for the null hypothesis that the two lines have the same slope and intercept. A Bonferroni multiple-comparison correction is applied by multiplying each p-value by 11 (number of antigen densities test). **(J,T)** Display the slope (top) and intercept (bottom) of the fitted line for the 11 different antigen densities tested.

Table S1: All-atom model building details

| Antibody | FAB structure source | Chains used (heavy, light, RBD) |
|----------|---------------------|---------------------------------|
| FD-11A | 7PQZ | A, B, E |
| FD-5D | 7PR0 | H, L, E |
| REGN10987 | 6XDG | C, A, E |
| CR3022 | 6W41 | H, L, C |
| EY6A | 6ZDH | H, L, A |
| FI3A | 7PQY | H, L, E |

Table S2: RBD-antibody-RBD chain lengths

| Antibody | Chain | Residues |
|----------|-------|----------|
| FD-5D | HC1 | 1-459 |
| | LC1 | 1-221 |
| | HC2 | 1-459 |
| | LC2 | 1-221 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |
| FD-11A | HC1 | 1-457 |
| | LC1 | 1-218 |
| | HC2 | 1-457 |
| | LC2 | 1-218 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |
| REGN10987 | HC1 | 1-450 |
| | LC1 | 1-216 |
| | HC2 | 1-450 |
| | LC2 | 1-216 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |
| CR3022 | HC1 | 1-449 |
| | LC1 | 1-221 |
| | HC2 | 1-449 |
| | LC2 | 1-221 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |
| EY6A | HC1 | 1-451 |
| | LC1 | 1-215 |
| | HC2 | 1-451 |
| | LC2 | 1-215 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |
| FI3A | HC1 | 1-447 |
| | LC1 | 1-214 |
| | HC2 | 1-447 |
| | LC2 | 1-214 |
| | RBD1 | 1-220 |
| | RBD2 | 1-220 |

Table S3: $\eta$ values for all antibody and RBD domains and interfaces except the epitope—paratope interfaces. *HC1 and LC2 share a small interface in the 1HZH crystal structure

| Identity | Structural Class | $\eta$ |
|---|---|---|
| HC1 | $\beta$ | 2.480 |
| LC1 | $\beta$ | 2.480 |
| HC2 | $\beta$ | 2.480 |
| LC2 | $\beta$ | 2.480 |
| RBD 1 | $\alpha/\beta$ | 1.916 |
| RBD 2 | $\alpha/\beta$ | 1.916 |
| HC1—LC1 interface | - | 2.124 |
| HC2—LC2 interface | - | 2.124 |
| HC1—HC2 interface | - | 2.124 |
| HC1—LC2 interface* | - | 1.507 |