Varying the position of phospholipid acyl chain unsaturation modulates hopanoid and sterol ordering

Ha-Ngoc-Anh Nguyen[a], Liam Sharp[b,c], Edward Lyman[b,d] and James P Saenz[a,e*]

[a] Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307 Dresden, Germany

[b] Department of Physics and Astronomy, University of Delaware, Newark DE 19716

[c] College of Arts and Sciences, Fairfield University, Fairfield, CT 06824

[d] Department of Chemistry and Biochemistry, University of Delaware, Newark DE 19716

[e] Medical Faculty, Technische Universität Dresden, 01062 Dresden, Germany

* corresponding author: James P Saenz

Email: james.saenz@tu-dresden.de

Abstract

The cell membrane must balance mechanical stability with fluidity to function as both a barrier and an organizational platform. Key to this balance is the thermodynamic ordering of lipids. Most Eukaryotes employ sterols, which are uniquely capable of modulating lipid order to decouple membrane stability from fluidity. Ancient sterol analogues known as hopanoids are found in many bacteria and are proposed as ancestral ordering lipids. The juxtaposition of sterols and hopanoids in extant organisms prompts us to ask why both pathways persist, especially in light of their convergent ability to order lipids. We reveal that both hopanoids and sterols order unsaturated phospholipids differently based on the position of double bonds in the phospholipid's acyl chain. We find that cholesterol and diplopterol's methyl group distributions lead to distinct effects on unsaturated lipids. In *Mesoplasma florum*, diplopterol's constrained ordering capacity reduces membrane resistance to osmotic stress, unlike cholesterol. These findings suggest cholesterol's broader lipid ordering ability may have facilitated the exploration of a more diverse lipidomic landscape in eukaryotic membranes.

Main Text

Introduction

To support life, cell membranes must balance mechanical stability (sufficient to perform robustly as a barrier) against fluidity and deformability (sufficient to support its role as an organizational platform for bioactivity). However, for bilayer forming mixtures of lipids, stability is often gained at the expense of fluidity. Different solutions to this dilemma have evolved on different branches of the tree of life. For example, thermophilic archaea synthesize double headed bolalipids that maintain membrane integrity even at extreme temperatures (1). In many eukaryotes, sterols solve this problem, their twin faces simultaneously ordering hydrocarbon chains while promoting lateral diffusivity (2). By decoupling the local motion of acyl chains and lipid translational freedom of motion, sterols allow cells to build membranes that are mechanically stable enough to withstand environmental perturbations, but fluid enough to support diffusion-dependent biochemistry.

Most non-eukaryotic organisms cannot synthesize sterols and must rely on other mechanisms to modulate their membrane properties. Some bacteria utilize a family of compounds called hopanoids (3). Hopanoids are tri-terpenoids, whose sedimentary record as early as 1.64 billion years ago (4). Since both families are synthesized from squalene, with homologous enzymes (squalene-hopene cyclase and oxidosqualene cyclase) (5), they share certain chemical similarities. Like sterols, hopanoids also reside within the membrane and modulate membrane robustness, fluidity, and resistance against abiotic stresses (3, 6). For these reasons, hopanoids are considered both bacterial and ancient sterol analogs. But why does life need two analogous classes of ordering lipids?

Despite their similarities, hopanoids and sterols possess distinct properties. Both diplopterol (Dpop), a common hopanoid in bacteria, and Chol (Chol), a mammalian sterol, interact favorably with and condense saturated lipids, a diagnostic feature of lipid ordering (7). However, the condensing effect of Dpop is impaired when an acyl chain unsaturation is introduced, and in a manner that depends on the location of the double bond. In this brief report, we examined how double bond position influences sterol and hopanoid ordering. We find that methyl group distribution on hopanoid backbones restricts its ordering ability, resulting in pronounced effects on cellular robustness in the model organism *Mesoplasma florum*. Conversely, Chol ordering is comparatively

50 insensitive to double bond position, allowing eukaryotic membranes more degrees of freedom in
51 regulating their lipidomes.
52
53 **Results**
54
55 Since unsaturation changes the interaction between Chol/Dpop and phospholipids, we hypothesize
56 that moving the position of double bonds along the phospholipid acyl chain can help us probe
57 Chol/Dpop ordering, thus revealing key structural features needed for ordering. We also aim to
58 draw a comparison between Chol and Dpop's ordering to showcase a constraint in the evolution of
59 sterol- and hopanoid-containing lipidomes.
60
We first determined the ordering of phospholipid chains by measuring monolayer surface pressure
vs area isotherms. Similar isotherms were obtained for PC phospholipids with a double bond at
one of three positions: $\Delta6$, $\Delta9$, or $\Delta11$ (chemical structures are shown in Fig. 1A), suggesting similar
packing in the pure membranes, regardless of the isomer (Fig 1B). The isotherms differ, however,
for binary mixtures of either Chol or Dpop, in a manner dependent on both terpenoid choice and
double bond position. To better quantify this difference, we calculated the condensing effect of the
terpenoid on each lipid, as well as the free energy of mixing (ΔG_{mix}) the phospholipid and terpenoid
(see Methods for definitions and details). A positive condensing effect is obtained when the
terpenoid orders lipid chains and increases lipid packing, which Chol consistently exhibited
regardless of the double bond position. In contrast, Dpop only condenses the $\Delta11$ isomer, does not
noticeably affect the $\Delta6$ isomer, and has a negative condensing effect on $\Delta9$ isomer. These
observations are mirrored by ΔG_{mix}, which reflects the thermodynamic balance between lipid-lipid
interactions and mixing entropy. Chol has favorable interaction with all 3 lipid isomers ($\Delta G_{mix} < 0$),
which progressively increases as the double bond moves away from the headgroup. On the other
hand, Dpop interaction with PC varies based on double bond positions. While Dpop was ideally
mixed with $\Delta6$-PC ($\Delta G_{mix} = 0$), the mixing of Dpop and $\Delta9$-PC was unfavorable ($\Delta G_{mix} > 0$), and only
with $\Delta11$-PC was the $\Delta G_{mix} < 0$. The distinction between Dpop interaction with $\Delta9$-PC and $\Delta11$-PC
is intriguing, as the double bonds are just 2 carbons apart.
79
To gain molecular insight into differential ordering of $\Delta9$ and $\Delta11$-PC by Dpop we performed
molecular dynamics simulations of bilayers of these binary mixtures. (Details in Methods) Figure 3
reports the 2H NMR chain order parameters (S_{CD}) for several different binary mixtures of Dpop with
phospholipids. S_{CD} is defined in Eq. 2 in methods; larger values indicate more ordered chains. A
model for Dpop compatible with the CHARMM36 family of force fields was built as described in
Methods, and first tested by simulating binary mixtures with DPPC (Fig. 3A), obtaining an ordering
effect similar to that observed in experiments on giant unilamellar vesicles (7).
87
We next considered binary mixtures with lipids with saturated chains at the sn-1 position and
monounsaturated chains ($\Delta19$ or $\Delta11$ isomers) at the sn-2 position (Fig 3B). As expected, the
saturated chains were more ordered by Dpop than the unsaturated chains, and were ordered
similarly regardless of which isomer was present in the other chain. Comparing the unsaturated
chains, the $\Delta11$ isomer was significantly more ordered than the $\Delta9$ isomer. This trend was
reproduced in PC with unsaturation in both acyl chains (Fig. 3C), where we again observed that
the $\Delta11$ isomer was significantly more ordered. In summary, the position of the double bond was
critical in determining the acyl chain's order when interacting with Dpop, mirroring the observations
in the monolayer experiments and supporting a model in which unsaturation position has a
significant effect on condensation in fluid lipid bilayers.
98
In prior work, Martinez-Seara et al. found that the positioning of the double bond relative to the
methyls protruding from cholesterol controls its condensing effect (8). We therefore performed a
similar analysis, comparing the location of the PC lipid's double bond to the positions of methyl
groups extending from the Dpop ring structure (annotated in Fig 1A) in our simulations. Figure 4
reports the distribution of double bonds of the hydrocarbon chains and Dpop methyl groups along

the membrane normal, with z = 0 being the center of the bilayer. The distribution of Δ9-PC's double
bond overlaps almost exactly with that of methyl group M2, while the Δ11-PC double bond falls in
between the M3 and M4 methyl groups. The alignment between the π bond and the methyl group
might explain the reduced condensing effect observed for Δ9-PC, while Dpop and Δ11-PC
interacted more favorably (8). This interplay between lipid structure and membrane biophysics
provides insights that could be extended to other sterols and hopanoids, possibly laying a path for
predicting how hopanoid/sterol and phospholipid structure collectively influence membrane
properties.
112
To investigate how this variation in lipid–lipid interaction might influence biomembrane function, we
employed *Mesoplasma florum* as a living model system. *M. florum* is a Mollicute with no cell wall
and a minimal genome (9). With limited machinery, Mesoplasma cannot synthesize its own lipids
and relies on supplemented lipids from the media, offering a straightforward way to manipulate its
lipidomes. By introducing either Δ9- or Δ11-PC to its lipid diet, we can create two identical biological
membrane systems differing only in their unsaturation site. We then investigated this system to
explore the effect of lipid–lipid interactions on a cellular scale.
120
Traditionally cultured with an undefined lipid diet in serums, we first test Mesoplasma's ability to
grow in a defined lipid diet. Figures 5A and 5B report growth rate and cellular lipid content,
respectively, suggesting Mesoplasma can both grow on and incorporate defined lipids into their
membrane. Since sterol ordering has previously been associated with membrane robustness (2),
we tested Mesoplasma membrane robustness with hypoosmotic shock. Live cells are subjected to
hypoosmotic conditions, forcing cells to rapidly expand. As the membrane is stressed and ruptures,
exposed cellular DNA is stained by propidium iodide. By quantifying the fluorescence intensity, we
estimated the fraction of cells lysed due to osmotic shock, inferring membrane robustness. Figure
5C shows the cell's susceptibility to osmotic shock when supplied with different lipid diets. When
cells were supplied with Chol, the addition of Δ9- or Δ11-PC did not produce significant changes in
cellular robustness. However, cells fed with Δ9-PC and Dpop exhibited higher susceptibility to lysis
than cells fed with Δ11-PC and Dpop. This data suggested that Dpop and Δ9-PC unfavorable
interaction counteracts Dpop's ability to bolster membrane robustness.
134
135 **Discussion**
136
Chol and Dpop are both prevalent lipids, accounting for approximately 40% of their respective
membrane lipidomes (10, 11). We show that both of these tri-terpenoids interact with unsaturated
lipids in a double bond position-dependent manner. While Chol condenses all unsaturated isomers
comparably, Dpop shows the strongest condensing effect on Δ11-PC, albeit less potently than
Chol. Simulations indicate that Dpop's interaction with unsaturated lipids is significantly hindered
by having multiple methyl groups extending from both sides of the cyclic ring structure, similar to
the biosynthetic precursor to Chol, lanosterol (12–14). Notably, when the double bond resides at
the Δ9 position, and overlaps with Dpop's methyl group M2, it prevents effective Dpop-induced lipid
packing. In combination with earlier simulation results for cholesterol from Martinez-Seara et al. (8),
our observations highlight the power of using simulations to explore lipid-lipid interactions,
especially in the case of less commercially available lipids like hopanoids. With careful
consideration in model development, one can explore the chemical landscape of lipids and the
consequences of their collective interactions.
150
From these observations, we hypothesized that the differentiation between Δ9 and Δ11 should be
the most significant in hopanoid-bearing membranes. In *Mesoplasma florum*, the favorable
interaction of Δ11-PC and Dpop enhanced membrane resilience to osmotic shock compared to Δ9-
PC. This remarkable result highlights how subtle changes to lipid structure can have striking
consequences for cellular robustness, and suggests a potential mechanism for osmoadaptation,
for which hopanoids have been shown to play a critical role in soil and plant associated bacteria
(15, 16). Indeed, multiple hopanoid-bearing bacteria have Δ11 as the monounsaturation site (17),

instead of ∆9 in eukaryotes. The hopanoid producing yeast, *Schizosaccharomyces japonicus*, also
possess a ∆12-desaturase (18, 19). Interestingly, in 2020, Chwastek et al. investigated the lipidome
of a hopanoid-bearing organism *Methylobacterium extorquens*, and found that the main
unsaturation position was ∆11 instead of ∆9, with an additional ∆5 unsaturation upregulated in cold-
adapted lipidome (11). Therefore, double bond position could represent a modifiable lipidomic
feature that cells can employ to homeostatically fine-tune the ordering effects of hopanoids.
164
For Chol-bearing organisms, double bond position does not have such a pronounced effect on lipid
ordering. This indifference might have alleviated evolutionary selection against ∆9 unsaturation in
early sterol-bearing organisms, providing more flexibility to produce lipids with double bond
positions optimized for orthogonal lipid-lipid or lipid-protein interactions. Our results, therefore,
suggest that a transition from hopanoid to sterol-containing lipidomes could have widened the
chemical landscape available for cells to explore for tuning membrane properties.

171
172   **Acknowledgments**
173
The authors thank Tomasz Czerniak, Lisa Junghans, Nataliya Safronova, Isaac Justice and Fabian
Rost for helpful discussions and feedback. This work was supported by the B CUBE of the
Technische Universität Dresden, a Maria Reich Fellowship (to H.N.A.N), a German Federal Ministry
of Education and Research (Bundesministerium für Bildung und Forschung - BMBF) grant (to
J.P.S., Project No. 03Z22EN12), and a Volkswagen Foundation "Life?" grant (to J.P.S., Project
No. 93090). E.L. and L.S. were supported by NIH award RO1GM120351. This work used Expanse2
at the San Diego Supercomputing Center through allocation BIO230093 from the Advanced
Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is
supported by National Science Foundation grants #2138259, #2138286, #2138307, #2137603, and
#2138296.
184

185   **Materials**
186   ∆6-, ∆9-, ∆11-PC and egg sphingomyelin were purchased from Avanti Polar Lipids. Chol and
187   palmitic acid were purchased from Sigma, and Dpop from Chiron. Stock concentrations of lipids
188   were measured by phosphate assay. Chol and Dpop were weighed out on a precision scale and
189   solubilized in a known volume of chloroform.
190

Methods
Monolayer. Chloroform solutions of pure lipids and mixtures were prepared at 0.2 mg/mL lipid
concentrations. Monolayers were prepared by injecting 15-30 µL of lipid solution onto an aqueous
subphase maintained at 20°C by a built-in temperature-controlled circulating water bath. The
subphase was comprised of 10 mM HEPES, 150 mM NaCl, pH 7. Isotherms were recorded using
a 70 cm^2 teflon Langmuir trough fitted with a motorized compression barrier equipped with pressure
sensor (Kibron DeltaPi).
The mean molecular area (MMAs) for each mixture were estimated from the averages of isotherms
from three monolayers that were prepared independently. Data were rounded down to the nearest
neighbor for condensation effect and free energy calculation. All isotherms were fitted to a
regression, and statistical significance was tested using manova with the 2 coefficients.
The condensation effect was calculated as follows:
203                                $$c = 100 - \frac{A_0}{X1A1 + X2A2} \ (\%)$$ (Equation 1)

Where c = % condensation, A_o = the MMA of the lipid mixture, X_1, X_2 = the mole fraction of lipid 1
and 2 in the mix, and A_1, A_2 = the MMAs of lipid 1 and 2 at surface pressures 30 mN/m. Error bars
were produced based on error propangation.

The ΔG was calculated by integrating the areas of lipid mixtures over pressures Π = 5, 10, 15, 20,
and 25 mN/m according to Grzybek et al (20). Error bars were produced based on error
propagation.
210
211 **Diplopterol model development.** A CHARMM compatible model for diplopterol (Dpop) was
212 developed using the automated atom typing and parameter assignment pipeline CGenFF.30
213 Charmm topology and parameter files are provided as Supplemental Material.
214
Simulation composition and construction. Simulation systems contained either DOPC or POPC
(unsaturation at either the Δ9 or Δ11 position) and one of either Dpop or Chol. All initial
configurations were built using the CHARMM-GUI webserver (21–23). Systems containing atypical
unsaturated chains (ie, Δ11) were generated by first building a binary mixture of the corresponding
Δ9 lipid (DOPC or POPC) with either Dpop or Chol, then "mutating" the unsaturated chain(s) to
move the double bond to the appropriate position, using a Charmm script provided by the Klauda
Lab.
All simulations contained approximately 550 lipids per leaflet and at least 50 TIP3P (24) water
molecules per lipid. All lipids were modeled with the CHARMM36 force-field (23), except Dpop
which was modeled using the CHARMM general force field, with atom types determined by the
paramchem server. (Gromacs topology file provided in the Supplemental Information.) Initial
dimensions in the membrane plane were about 17.5 nm x 17.5 nm, containing approximately
270,000 atoms.
Four different binary mixtures were simulated: Δ9-DOPC: Dpop, Δ11-DOPC: Dpop, Δ9-POPC:
Dpop, Δ11 POPC: Dpop. Each binary mixture was simulated at four different compositions: 95:5,
85:15, 70:30, 50:50. Each binary system was simulated for 500 nsec of production simulation as
described below. Four additional controls were simulated without any sterol or hopanoid, each for
50 nsec of production simulation as described below: Δ9-DOPC, Δ11-DOPC, Δ9-POPC, Δ11-
POPC.
234
Equilibration and production simulations. Each system was prepared individually for production
simulation through a series of 6 minimization and heating steps as provided by the CHARMM-GUI
equilibration protocol: (i) steepest descent to minimize the initial configuration; (ii) 125,000 steps of
leapfrog dynamics with a 1 fsec timestep and velocities reassigned every 500 steps; (iii) 125,000
steps of leapfrog dynamics with a 1 fsec timestep, pressure controlled by the Parinello-Rahman
barostat (25) and velocities reassigned every 500 steps, then a total of 750,000 steps of leapfrog
dynamics with a 2 fsec timestep and hydrogen positions constrained by LINCS (26), pressure
controlled by the Parinello-Rahman barostat (25), and velocities reassigned every 500 steps.
During equilibration, double bonds were restrained in the cis configuration to prevent isomerization;
these restraints are gradually reduced during the final three stages of the equilibration protocol.
Production simulations (NPT ensemble) were integrated with leapfrog using the Parinello-Rahman
(25) barostat to control pressure (time constant 5 psec; compressibility 4.5e−5 bar−1; coupled
anisotropically to allow independent fluctuation of the in-plane and normal directions) and
temperature controlled using Nose-Hoover38,39 (time constant 1 psec) at a temperature of 298K.
Hydrogens were constrained with LINCS (expansion order 4), a 2 fsec timestep was used, short
range electrostatics were computed directly within 1.2 nm, and long-range electrostatics were
computed every timestep using particle mesh Ewald (27, 28) with a grid spacing of 1 Å and cubic
interpolation. Long range dispersion was smoothly truncated over 10-12 nm using a force-switch
cutoff scheme. Simulations were performed with Gromacs 2020.4.
254
255 **Calculation of simulation observables.** The distribution of angles between either Dpop or Chol
256 and the membrane normal was computed, defining the orientation of both by a vector from atom
257 C24 to atom O3. The locations of methyl groups in Dpop or Chol along the direction normal to the
258 membrane were recorded and compiled into histograms with a bin size of 0.87 Å. Deuterium order
259 parameters were obtained from the simulations via

260
$$S_{CD} = \frac{1}{2}\langle 3cos^2(\phi) - 1\rangle \qquad \text{(Equation 2)}$$

where ɸ is the angle between the C-H bond vectors and the membrane normal at each position
along the hydrocarbon chain. The area per lipid was calculated using the Voronoi construction
implemented in MEMBPLUGIN (29). The location of each lipid is defined by the center of geometry
of the C2, C21, and C31 atoms and the location of the Chol/ Dpop was defined by the O3 atom,
and then a Voronoi construction is built around these points in the plane parallel to the membrane
surface.
267
Cell culture. *Mesoplasma florum* L1 strains were grown in a modified, lipid-free SP4 media with
components as follows (per 1L): Bacto Tryptone 10g, Bacto Peptone 5.3g, PPLO 3.5g, BSA 5.95g,
Yeastoleate 2g, D-Glucose 5g, sodium bicarbonate 3.15g, L-Glutamine 0.05g, Penicillin G-sodium
salt 0.645g, phenol red 11 mg, pH to 7.0. Lipid diet was added separately prior to passaging at
concentration: Dpop, Chol 5 mg/L, egg sphingomyelin 25 mg/L, palmitic acid 10 mg/L, Δ9- and Δ11-
PC 12.5 mg/L for the corresponding diets. Cells were grown in glass flasks and incubated at 30°C
with shaking at 60 rpm. Growth was recorded using phenol red media pH detection through
absorbance at 562nm using a 10mm cuvette (DeNovix DS-11 FX+). Growth rate was defined as
negative of the slope of the linearly fitted trendline in the indicative range of phenol red (OD_{562nm}
from 0.75 – 0.4).
278
Membrane incorporation. Cells were collected in early exponential stage and centrifuged (5000
rcf, 7 min, 30°C). Supernatant were discarded and cell pellet was washed with wash buffer (200
mM NaCl, 25 mM HEPES, 1% glucose, pH 7.0) and centrifuged (5000 rcf, 7 min, 30°C). The
collected pellet was then subjected to a Bligh Dyer extraction (30). Briefly, the pellet was
homogenized in a mixture of water:chloroform:methanol in 0.8:1:2 ratio and sonicated for 2
minutes. Subsequently, water and chloroform were added in 1:1 ratio. The mixture was centrifuged
at 2000 rcf for 30 seconds in a mini centrifuge to promote phase separation. The lower, organic
fraction containing lipids was collected and transferred to a fresh tube. The total lipid extract was
then deposited on a silica gel plate (Supelco) and placed in a glass chamber. Chromatography was
performed using chloroform as the running phase. After the run, the plate was dried and stained
using 8 % copper sulphate in 3% phosphoric acid solution and heated until visible bands were
observed. Images were captured using a GelDoc (Biozym Azure c600) and analysed using ImageJ.
291
Membrane osmotic shock. Cells were collected in early exponential stage and centrifuged (5000
rcf, 7 min, 30°C). The collected pellet was resuspended in a serial dilution of 0%, 20%, 40%, 60%,
80% and 100% of wash buffer (200 mM NaCl, 25 mM HEPES, 1% glucose, pH 7.0). The
suspension was stained with 10 uM propidium iodide and added to a 96-well plate. Fluoresence
emission was recorded using a Tecan Spark fluorescence reader, with excitation at 529 – 549 nm
and emission at 609 – 629 nm. The fraction of cell lysed was calculated by normalizing the signal
of each sample to the 0% and 100% wash buffer sample. Figure 5C represented the fraction of cell
lysed in 80% of wash buffer.
300
301

References

1. K. Yamaguchi, M. Kinoshita, Highly stable lipid membranes from archaebacterial extremophiles. *Prog. Polym. Sci.* **18**, 763–804 (1993).
2. O. G. Mouritsen, M. J. Zuckermann, What's so special about cholesterol? *Lipids* **39**, 1101–1113 (2004).
3. B. J. Belin, *et al.*, Hopanoid lipids: From membranes to plant-bacteria interactions. *Nature Reviews Microbiology* **16**, 304–315 (2018).
4. J. J. Brocks, *et al.*, Biomarker evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea. *Nature* **437**, 866–870 (2005).
5. S. Racolta, P. B. Juhl, D. Sirim, J. Pleiss, The triterpene cyclase protein family: A systematic analysis. *Proteins: Structure, Function, and Bioinformatics* **80**, 2009–2019 (2012).
6. J. P. Sáenz, *et al.*, Hopanoids as functional analogues of cholesterol in bacterial membranes. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 11971–11976 (2015).
7. J. P. Sáenz, E. Sezgin, P. Schwille, K. Simons, Functional convergence of hopanoids and sterols in membrane ordering. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 14236–14240 (2012).
8. H. Martinez-Seara, *et al.*, Interplay of unsaturated phospholipids and cholesterol in membranes: Effect of the double-bond position. *Biophysical Journal* **95**, 3295–3305 (2008).
9. V. Baby, *et al.*, Inferring the Minimal Genome of Mesoplasma florum by. *mSystems* **3**, 1–14 (2018).
10. J. H. Lorent, *et al.*, Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nat Chem Biol* **16**, 644–652 (2020).
11. G. Chwastek, *et al.*, Principles of Membrane Adaptation Revealed through Environmentally Induced Bacterial Lipidome Remodeling. *Cell Reports* **32** (2020).
12. J. A. Urbina, *et al.*, Molecular order and dynamics of phosphatidylcholine bilayer membranes in the presence of cholesterol, ergosterol and lanosterol: a comparative study using 2H-, 13C- and 31P-NMR spectroscopy. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1238**, 163–176 (1995).
13. A. M. Smondyrev, M. L. Berkowitz, Molecular Dynamics Simulation of the Structure of Dimyristoylphosphatidylcholine Bilayers with Cholesterol, Ergosterol, and Lanosterol. *Biophysical Journal* **80**, 1649–1658 (2001).
14. K. Sabatini, J.-P. Mattila, P. K. J. Kinnunen, Interfacial Behavior of Cholesterol, Ergosterol, and Lanosterol in Mixtures with DPPC and DMPC. *Biophysical Journal* **95**, 2340–2355 (2008).
15. E. M. Tookmanian, B. J. Belin, J. P. Sáenz, D. K. Newman, The role of hopanoids in fortifying rhizobia against a changing climate. *Environmental Microbiology* **23**, 2906–2918 (2021).
16. E. Tookmanian, *et al.*, Hopanoids Confer Robustness to Physicochemical Variability in the Niche of the Plant Symbiont Bradyrhizobium diazoefficiens. *Journal of Bacteriology* **204**, e00442-21 (2022).
17. A. J. Fulco, Fatty acid metabolism in bacteria. *Progress in Lipid Research* **22**, 133–160 (1983).
18. M. Makarova, *et al.*, Delineating the Rules for Structural Adaptation of Membrane-Associated Proteins to Evolutionary Changes in Membrane Lipidome. *Curr Biol* **30**, 367-380.e8 (2020).
19. J. Bouwknegt, *et al.*, A squalene–hopene cyclase in Schizosaccharomyces japonicus represents a eukaryotic adaptation to sterol-limited anaerobic environments. *Proceedings of the National Academy of Sciences* **118**, e2105225118 (2021).
20. M. Grzybek, J. Kubiak, A. \Lach, M. Przyby\lo, A. F. Sikorski, A raft-associated species of phosphatidylethanolamine interacts with cholesterol comparably to sphingomyelin. A Langmuir-Blodgett monolayer study. *PLoS ONE* **4** (2009).

21. S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: A web-based graphical user interface for
CHARMM. *Journal of Computational Chemistry* **29**, 1859–1865 (2008).
22. , CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and
CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field | Journal of
Chemical Theory and Computation (August 29, 2023).
23. E. L. Wu, *et al.*, CHARMM-GUI Membrane Builder toward realistic biological membrane
simulations. *Journal of Computational Chemistry* **35**, 1997–2004 (2014).
24. D. J. Price, C. L. Brooks, A modified TIP3P water potential for simulation with Ewald
summation. *J Chem Phys* **121**, 10096–10103 (2004).
25. M. Parrinello, A. Rahman, Polymorphic transitions in single crystals: A new molecular
dynamics method. *Journal of Applied Physics* **52**, 7182–7190 (1981).
26. , LINCS: A linear constraint solver for molecular simulations - Hess - 1997 - Journal of
Computational Chemistry - Wiley Online Library (August 29, 2023).
27. T. Darden, D. York, L. Pedersen, Particle mesh Ewald: An N·log(N) method for Ewald sums
in large systems. *The Journal of Chemical Physics* **98**, 10089–10092 (1993).
28. , A smooth particle mesh Ewald method | The Journal of Chemical Physics | AIP Publishing
(August 29, 2023).
29. R. Guixà-González, *et al.*, MEMBPLUGIN: studying membrane complexity in VMD.
Bioinformatics **30**, 1478–1480 (2014).
30. E. G. Bligh, W. J. Dyer, A RAPID METHOD OF TOTAL LIPID EXTRACTION AND
PURIFICATION.
375

Fig 1. (A) Chemical structure of di-unsaturated PC isomers, Chol and Dpop and (B) Isotherms of the corresponding lipids at 20 °C, *** (F=12.4, p < 0.0005).

Fig 2. PC isomers interact differently with Chol and Dpop. While Chol ordering of PC increases as double bond position is shifted further from the headgroup, Dpop only exhibits an ordering effect with Δ11-PC. (A) Isotherms of PC isomers mixed with Chol (2:1) at 20°C. ** ($F=7.75$, $p < 0.005$) manova (B) Isotherms of PC mixture with Dpop (2:1) at 20°C. ** ($F=8.18$, $p < 0.005$) manova (C) Condensing effect of Chol and Dpop on PC isomers calculated at 30 mN/m. Error bar represents standard deviation. n.s. ($p > 0.5$) * ($p < 0.05$) unpaired t-test (D) Energy of interaction (ΔG_{mix}) of lipid pairs during compression. Error bar represents standard deviation. n.s. ($p > 0.5$) * ($p < 0.05$) ** ($p < 0.005$) unpaired t-test.

Fig 3. Molecular dynamic simulations show Dpop orders 18:1 Δ11 more efficiently than 18:1 Δ9.
(A) Dpop ordering effect with saturated DPPC were confirmed by S_{CD}. (B) Dpop ordered saturated chains most efficiently, then 18:1 Δ11 and the least for 18:1 Δ9. (C) Dpop's ordering effect was similar regardless of sn chain position, but depended on double bond position, with Δ11 a greater ordering effect than Δ9.

Fig 4. Overlapping distribution of Dpop's methyl groups and PC's double bond correspond to a reduced ordering effect. (A) The double bond in Δ9-PC overlaps with Dpop M2, preventing efficient lipid packing. (B) No Dpop methyl group overlaps with the double bond of Δ11-PC.

Fig 5. Dpop and Δ11-PC enhance robustness of *Mesoplasma florum* to hypoosmotic shock compared with Dpop and Δ9-PC. (A) Chol and Dpop were incorporated into Mesoplasma membranes according to their respective diets (B) Growth rate of *M. florum* on different diets. ** (F=24.4, p < 0.005) * (F=10.6, p < 0.05) Analyses were performed using one-sided anova with Tukey post hoc test (C) Membrane robustness reflected by the fraction of cell lysed when subjected to hypoosmotic shock. n.s. (F=1.47, p<0.5) ** (F=16.6, p < 0.005) Analyses were performed using one-sided anova with Tukey post hoc test.