Emergence of *Prochlorococcus* in the Tonian oceans and the initiation of Neoproterozoic oxygenation

Hao Zhang[1,2^], Sishuo Wang[2^], Tianhua Liao[2], Sean A. Crowe[3], Haiwei Luo[2,4,5*]

[1]Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen 518000, China

[2]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[3]Departments of Microbiology and Immunology, and Earth, Ocean, and Atmospheric Sciences, University of British Columbia, Vancouver, British Columbia, Canada

[4]Earth and Environmental Sciences Programme, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[5]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[^] These authors contributed equally to this study

[*]**Corresponding author:** Haiwei Luo

Email: haiweiluo@cuhk.edu.hk

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences - Microbiology; Physical Sciences - Earth, Atmospheric, and Planetary Sciences

Key words: *Prochlorococcus*, Neoproterozoic oxygenation event, molecular dating, plastid, cyanobacteria

Abstract

Prochlorococcus are the smallest and most abundant photosynthetic organisms on Earth,

contributing up to 50% of the chlorophyll in the oligotrophic oceans. Despite being important

in regulating the carbon cycle in today's ocean, the ecological significance of

Prochlorococcus in Earth's history remains elusive. Our new robustly calibrated molecular

clock reveals that *Prochlorococcus* emerged in the deep photic zone of the Tonian oceans.

The classical antenna for light harvesting in Cyanobacteria, the phycobilisome, was replaced

in *Prochlorococcus* by the chlorophyll☐based light☐harvesting complex enabling more

efficient use of blue light that penetrated to deeper water. Importantly, *Prochlorococcus*

colonization of deep water enhanced access to phosphate, which was rich in upwelled

seawater, but likely scarce in the Tonian surface ocean, promoting expansion of

Prochlorococcus and associated photosynthetic oxygen production. Colonization of deeper

waters would also have improved access to ammonium, leading to the neutral loss of nitrate

utilization genes. Our research thus documents the conspicuous emergence of new

photosynthetic bacterial lineages in the run-up to the Neoproterozoic oxygenation event,

implying an additional layer of eco-evolutionary complexity during this pivotal interval in

Earth's history.

Significance (max 120 words allowed)

Despite being important in today's oceans, the ecological significance of *Prochlorococcus* in the past remains elusive. By 'borrowing' fossils from photosynthetic eukaryotes and leveraging the sequential Bayesian dating approach, we have developed a new pipeline for dating the evolution of *Prochlorococcus*, which systematically quantifies the uncertainties in using different fossil calibrations, molecular datasets, species tree topologies and molecular clock models. Our analyses pin-pointed the origin of *Prochlorococcus* to the phosphorous-limited Tonian oceans. The changes in the photosynthetic apparatus in early *Prochlorococcus* enabled them to colonize the phosphorus-rich deep photic niches, which would have led to increased primary productivity and oxygen release, likely contributing to the Neoproterozoic Oxygenation Event.

Introduction

Prochlorococcus host more than half of the chlorophyll biomass in oligotrophic oceans (Partensky and Garczarek, 2010). As key primary producers in the modern ocean, *Prochlorococcus* fix as much as four gigatons of carbon each year and are the foundation of the marine carbon cycle and food web (Flombaum et al., 2013). The success of *Prochlorococcus* in today's oceans has been attributed to multiple physiological features. Importantly, *Prochlorococcus* use divinyl chlorophyll (DVChl) for harnessing light energy (Ralf and Repeta, 1992). DVChl harvests blue light much more efficiently than the more common monovinyl chlorophyll used by other cyanobacterial lineages and it thus enables *Prochlorococcus* to thrive in the deepest layers of the euphotic zone, where blue light

dominates (Ito and Tanaka, 2011). *Prochlorococcus* is also the smallest photosynthetic group

on Earth (Partensky et al., 1999). As a result, their high surface-to-volume ratio provides

enhanced nutrient acquisition efficiency, which together with effective blue-light absorption

promotes photosynthesis by *Prochlorococcus* in oligotrophic tropical and subtropical oceans

(Partensky et al., 1999). In today's ocean, *Prochlorococcus* is also the dominant phototroph in

more nutrient-rich, oxygen-depleted anoxic marine zones (AMZ) in the eastern tropical North

and South Pacific Oceans (Goericke et al., 2000; Lavin et al., 2010). Since the AMZ lineages

represent the earliest-split branches of *Prochlorococcus*, it has been proposed that

Prochlorococcus emerged in low-oxygen environments and, by extension, contributed to

early ocean oxygenation (Ulloa et al., 2021).

Despite the ecological prominence of *Prochlorococcus* in both modern and likely the

past oceans, the divergence time of *Prochlorococcus* remains contentious. Different relaxed

clock analyses based on 16S and 23S rRNA genes yielded divergence times of the total

Prochlorococcus group at ~200 Mya (Boden et al., 2021), ~500 Mya (Sánchez-Baracaldo,

2015) and ~800 Mya (Sánchez-Baracaldo et al., 2014). The use of protein-coding genes gave

divergence times of the total *Prochlorococcus* group at ~400 Mya (Fournier et al., 2021),

~500 Mya (Sánchez-Baracaldo et al., 2017) and ~1,000 Mya (Schirrmeister et al., 2015). The

discrepancies in *Prochlorococcus* divergence times are likely caused by variable use of

fossils and other time calibrations and the use of alternative gene sets, clock models, and tree

topologies (Warnock et al., 2012; Duchêne et al., 2014; dos Reis et al., 2015) (see

Supplementary Discussion). Despite their importance, these factors were rarely tested

rigorously thus making it difficult to evaluate the accuracy of age estimates in previous

studies. Importantly, even if all these factors are well tested and controlled, the rarity of

cyanobacterial fossils and the absence of a maximum boundary of the available

cyanobacterial fossils poses an appreciable challenge in determining the antiquity of

Prochlorococcus (Zhang et al., 2021). Indeed, applying maximum age constraints at the

calibration nodes strongly impacts posterior age estimates of uncalibrated lineages (Hedges et

al., 2018; Morris et al., 2018; Wang and Luo, 2021). However, informative maximum age

constraints are typically missing from microbial fossils and can only be found in some animal

and plant fossils. Moreover, the recent availability of several genome sequences of

uncultivated basal *Prochlorococcus* lineages (Ulloa et al., 2021) requires re-estimation of the

antiquity of *Prochlorococcus*, which should also include an evaluation of the factors that

influence the accuracy of posterior ages.

Here, we leverage the abundant plant and algal fossils and the well-established plant

plastid endosymbiosis theory to develop a new pipeline that systematically tests the factors

that influence the accuracy of posterior ages and thus yields robust estimates of

Prochlorococcus antiquity. The plant plastid endosymbiosis theory states that the origin of all

plastids in photosynthetic eukaryotes, except for the photosynthetic amoeba *Paulinella*

(Marin et al., 2005), can be traced back to an ancient primary endosymbiosis involving a

eukaryote and a cyanobacterium (Gray, 1992; Bhattacharya and Medlin, 1995; Keeling, 2013;

Ponce-Toledo et al., 2017). Plant plastid endosymbiosis theory thus ties the evolutionary

histories of cyanobacteria to those of photosynthetic eukaryotes. Eukaryotic fossils are indeed

being increasingly used in dating the evolution of Cyanobacteria (Shih et al., 2016; Sánchez-

Baracaldo et al., 2017; Fournier et al., 2021). Our strategy builds on these by: 1) using more

fossil-based age constraints (including maximum age constraints) on eukaryotic lineages with

more complete taxonomic sampling including lineages that have undergone secondary

endosymbiosis; and 2) applying a sequential Bayesian dating approach to better constrain the

divergence time of eukaryotic lineages thereby propagating more accurate time information

to Cyanobacteria, including *Prochlorococcus*. These joint analyses led us to conclude that

Prochlorococcus arose in the Tonian ocean. Further, the use of an evolutionary genetic proxy

implies that *Prochlorococcus* was born through a founder effect, and this strengthens the idea

that *Prochlorococcus* emerged in the deep photic zone, a unique niche separated from upper

waters where its *Synechococcus* ancestors thrived.

Results

<u>*Prochlorococcus* originated in the Tonian ocean</u>

We implemented a sequential Bayesian dating method that takes advantage of the

abundant fossil records (Fig. 1a; Fig. S1a) available from photosynthetic eukaryotes to

calibrate the evolution of Cyanobacteria. The last common ancestor (LCA) of

Prochlorococcus (denoted as "Proch-AMZI/II/III-LCA") clearly emerged within the Tonian

period at 878 Mya (95% HPD: 987-767 Mya) (Fig. 1b). Meanwhile, the LCA of

Prochlorococcus clades HL, LL and AMZI/II (denoted as "Proch-AMZI/II-LCA") evolved at

787 Mya (95% HPD: 896-687 Mya), which coincided with the early stage of the

Neoproterozoic Oxygenation Event (NOE) (Fig. 1b). In this case, the branch leading to the

LCA of *Prochlorococcus* HL, LLI, and LLII/III clades (denoted as "SBE-LCA" to keep

consistency with (Zhang et al., 2021)) spanned the time that encompassed the duration of the

Neoproterozoic Snowball Earth events (645-635 Mya for Marinoan glaciation and 717-659

Mya for Sturtian glaciation), supporting the main conclusion of the previous study, which

used the traditional cyanobacterial fossil-based strategy (Zhang et al., 2021).

Note that the sequential Bayesian method used here is different from the commonly used

secondary dating method (using time estimates from previous studies as calibrations)

(Heckman et al., 2001; Aoki et al., 2013; Chriki-Adeeb and Chriki, 2016) for two reasons.

First, the time prior used in sequential dating analysis follows a specific probability

distribution, but secondary calibrations ignore the uncertainties on age estimates. Second, the

sequential dating analysis contains two steps each based on a distinct set of molecular data,

while secondary dating analysis largely relies on the same molecular dataset (gene alignments)

(dos Reis et al., 2018).

In our implementation, the posterior ages of eukaryotic lineages derived from the first-

step of the sequential analysis were used as the time priors to calibrate Cyanobacteria

evolution where few time constraints are available. Since genome-scale dating of eukaryotic

lineages is very slow with fully partitioned molecular data and is prone to have reduced

coverage probability (the probability that the 95% HPD interval of posterior ages contains the

true age) due to the large number of partitions (Angelis et al., 2018), we first clustered the

eukaryotic gene families into 1-, 3-, 5-, 7-, and 9-partitions based on their evolutionary rates

(see details in Supplementary Methods). In general, the posterior ages of eukaryotic lineages

were consistent, except for the case where a single partition and a single substitution model

was used, which did not fit the data well as indicated by its larger Akaike Information

Criterion (AIC) value (Fig. S1b). In all other cases, the crown group of Archaeplastida was

estimated to evolve at ~1.60 Gya, which is consistent with the estimates from other studies

(Parfrey et al., 2011; Betts et al., 2018; Wang and Luo, 2021) (Fig. S1b). By comparing the

distributions of the posterior ages of eukaryotic nodes in the first-step of sequential analysis

to the effective prior on the corresponding nodes in the second-step of analysis (see details in

Supplementary Methods), we found that the two distributions in dating analyses were nearly

identical when 5-, 7- and 9-partition data were used (Fig. S2), suggesting that the sequential

Bayesian approach works well for these analyses. We therefore adopted the evolutionary

timeline of eukaryotes based on the 5-partition data (lowest AIC value and high similarity

between posterior ages and effective prior) and applied the posterior ages of eukaryotic

lineages as time prior at the overlapping nodes in a larger phylogenomic tree that includes

both photosynthetic eukaryotes and cyanobacteria (Fig. S1c).

To validate the evolutionary timeline of *Prochlorococcus*, we performed a series of tests

by using different molecular data, clock models, fossil calibrations and species tree

topologies. The posterior ages estimated with all these alternative settings are largely

consistent with that estimated with the focal strategy (detailed above) and fully support the

origin of *Prochlorococcus* (Proch-AMZI/II/III-LCA) in the Tonian period, followed by the

emergence of Proch-AMZI/II-LCA in the early stages of the NOE (Fig. S3). Moreover, the

branch leading to the SBE-LCA encompasses the Snowball Earth events in all the dating

analyses (Fig. S4; see Supplementary Results). Our study also results in an estimate of

Cyanobacteria age. While this is not the focus here, the antiquity of Cyanobacteria has been

an important research topic and under heavy debate (Schirrmeister et al., 2015; Schirrmeister

et al., 2016; Fournier et al., 2021; Sánchez-Baracaldo et al., 2022). We therefore included a

discussion (see Supplementary Discussion).

Prochlorococcus genomic changes

We inferred the genes gained and lost upon the emergence of *Prochlorococcus* first

without using the SAGs to avoid the false prediction of genomic events due to the relatively

low completeness of the SAGs (e.g., 81.9% for AMZI-B-ETNP). Our reconstructions rely on

the gene tree-species tree reconciliation approach, in which a best-fit reconciliation tool was

selected through a comprehensive simulation-based benchmarking analysis [see Fig. S4 in

(Zhang et al., 2023)]. We then used the same reconstruction methods to validate these

evolutionary events when SAGs were included.

As the only phytoplankton group using divinyl chlorophyll (DVChl) for harvesting light

energy (Ralf and Repeta, 1992), *Prochlorococcus* lost the gene *bciB* that inhibits the

conversion of DVChl to MVChl (monovinyl chlorophyll) at Proch-AMZI/II/III-LCA, thereby

promoting the accumulation of DVChl in their membranes (Ito et al., 2008). Compared to the

MVChl used by *Synechococcus*, DVChl more efficiently absorbs blue light that penetrates to

deeper waters than other photosynthetically active radiation (Ito and Tanaka, 2011). Likewise,

the gain of the *PcCao* gene for the synthesis of chlorophyll b (Satoh and Tanaka, 2006) in

Proch-AMZI/II/III-LCA also enables *Prochlorococcus* ancestors to efficiently harvest blue

light at exceedingly low intensities characteristic of deep waters (Hess et al., 2001) (Fig. 1b).

These genomic changes would have enabled *Prochlorococcus* ancestors to explore the deep

photic zone, a niche where its *Synechococcus* ancestor would not thrive.

Unlike most cyanobacteria, which use the phycobilisome as the photosynthetic antenna,

the main light-harvesting antenna of *Prochlorococcus* is made up of prochlorophyte

chlorophyll-binding (Pcb) protein (Biller et al., 2015). By reconstructing the gene

evolutionary paths with *Prochlorococcus* SAGs included, we found that the phycobilisome

genes (*apcACDEF*, *cpcEFG*, and *cpeCES*) were present in both Proch-AMZI/II/III-LCA and

Proch-AMZI/II-LCA and lost at SBE-LCA. Using the same approach, we found that the

cholorophyll-binding protein (encoded by *pcbD*) was obtained at Proch-AMZI/II-LCA. The

replacement of the photosynthetic antenna thus did not co-occur with the emergence of

Prochlorococcus. We note that gene absence in SAGs could be due to the incomplete nature

of the SAG genomes, however, in this case, simultaneous absence of the *pcbD* gene in all the

five SAGs that are more than 80% complete seems unlikely.

Despite the absence of the nitrate utilization genes in most *Prochlorococcus* isolates,

some uncultivated *Prochlorococcus* lineages contain these genes (Martiny et al., 2009;

Berube et al., 2015; Berube et al., 2019). By including *Prochlorococcus* SAGs that contain

the assimilatory nitrate transporter gene (*nrtB*) and the assimilatory nitrate reductase gene

(*narB*) in our reconstructions, we inferred the presence of both genes in Proch-AMZI/II/III-

LCA and Proch-AMZI/II-LCA and the losses of these genes at SBE-LCA (Fig. 1b). In

addition to the nitrate utilization genes, we inferred the losses of molybdopterin biosynthesis

genes at SBE-LCA. Since these genes are known to co-locate with the nitrate reductase genes

in *Synechococcus* (Rubio et al., 1998; Palenik et al., 2003), they may function as the cofactor

of nitrate reductase in *Prochlorococcus*. Therefore, the loss of nitrate reductase in

Prochlorococcus likely rendered the molybdopterin dispensable and thus led to the losses of

molybdopterin biosynthesis genes (*moaABCDE*, *mobA* and *moeA*) at SBE-LCA.

Selection efficiency was reduced upon the emergence of *Prochlorococcus*

The method for inferring selection efficiency in deep time was developed (Zhang, 2000)

and recently improved (Luo et al., 2017), which involves nonsynonymous substitutions only

and compares the rate of nonsynonymous substitutions leading to replacements of

physicochemically dissimilar amino acids (i.e., radical changes; d_R) to the rate of

nonsynonymous substitutions leading to replacements of physicochemically similar amino

acids (i.e., conservative changes; d_C). Since radical changes are more likely to be deleterious

than conservative changes (E. Zuckerkandl, 1965; Dayhoff, 1972), an excess of the radical

changes in a deeply branching lineage compared to its sister lineage suggests random fixation

of deleterious mutations by genetic drift in the former. Using this method, we found a

significant increase of the d_R/d_C ratio across the genomic regions in the branch leading to

Proch-AMZI/II/III-LCA relative to the branch leading to the LCA of *Synechococcus* clade 5.1

(Fig. 2a), indicating that the emergence of *Prochlorococcus* was accompanied by a significant

reduction of the efficiency of purifying selection, allowing for an accelerated accumulation of

deleterious mutations through genetic drift.

Discussion

Geochemical context that supports the emergence of *Prochlorococcus* in the Tonian period

At the 95% HPD, *Prochlorococcus* emerged between 989 to 770 Mya, encompassing

almost the entire Tonian period. A Tonian, pre-Sturtian, age for the emergence of the

Prochlorococcus lineage would have been set against the backdrop of dynamic ocean

chemistry characterized by a variably oxygenated surface ocean that was co-populated by

diverse microbial eukaryotes, including phytoplankton and the earliest metazoans (Trower et

al., 2023) (Fig. 1c). Multi-proxy data, collected from geographically diverse sites, converge

on Tonian deep oceans that were pervasively anoxic across 60-80% of the ocean floor, or

more (Tahata et al., 2015; Lau et al., 2017). The deep anoxic oceans were predominantly

ferruginous in nature, albeit with evidence for transient, spatially restricted euxinia, and

possibly even brief (<0.5 Myr) episodes of pervasive deep ocean oxygenation (Stolper and

Keller, 2018; Tostevin and Mills, 2020). Shallow waters, by contrast, were modestly, though

perhaps increasingly, oxygenated across the Tonian with evidence for the episodic and

persistent intrusion of anoxic deep waters into the surface oceans (Zhang et al., 2022;

Clarkson et al., 2023), with corresponding implications for nutrient cycling and availability

and for marine life in the euphotic zone.

The strongest differentiators between the earliest *Prochlorococcus* (i.e., Proch-

AMZI/II/III-LCA) and its *Synechococcus* ancestors are adaptation to low-light and loss of

capacity for nitrate assimilation, which imply both a deeper photosynthetic niche and a likely

shift to use ammonium as the primary nitrogen source. A key driver of *Prochlorococcus*

emergence was thus likely a competitive advantage to dwelling in deeper waters. A well-

known advantage emerges under conditions of strong P-limitation, whereby deep-dwelling

low-light adapted phototrophs gain first access to upwelling phosphorus (Jones et al., 2015;

Ozaki et al., 2019). The low-light advantage conferred on the earliest *Prochlorococcus* cells

through the adoption of divinyl chlorophyll would have enhanced their access to P, relative to

contemporary algae and their *Synechococcus* ancestors. It would also have increased access

to NH_4^+ upwelling from deeper anoxic waters obviating the need for nitrate assimilation

(Michiels et al., 2017) (Fig. 1c).

In this way, the emergence of the *Prochlorococcus* would have restricted the flux of

nutrients to shallower waters and limited the contributions of higher-light adapted

phytoplankton like algae to primary production. This is supported through the biomarker

record, which indicates a dominance of cyanobacterial primary production until after the

Sturtian glaciation (Brocks, 2018). It is also supported by the Tonian N-isotope record, which

implies a limited contribution from nitrate assimilation to primary production at this time

(Kang et al., 2023). Importantly, the adaptation of the *Prochlorococcus* LCA to lower-light

would also allow it to better compete with low-light adapted anoxygenic phototrophs that

would likely have populated anoxic regions of the Tonian euphotic zone with strong potential

to cause ocean oxygenation (Johnston et al., 2009a; Jones et al., 2015; Ozaki et al., 2019).

Competition between oxygenic photosynthetic cyanobacteria and anoxygenic phototrophs

that oxidize ferrous iron (photoferrotophs), is known to limit photosynthetic oxygen

production (Jones et al., 2015; Ozaki et al., 2019). In this way, the capacity of the

Prochlorococcus LCA to grow in deeper waters could have enhanced its ability to access

upwelling phosphorus thereby enhancing the oxygenic fraction of total photosynthesis with

potential to shift oxygen production towards runaway oxygenation (Ozaki et al., 2019).

<u>Evolutionary mechanisms underlying the origin of *Prochlorococcus*</u>

We propose that the colonization of the deep photic zone by *Prochlorococcus* started by a

seed population with only a few individuals that obtained the ability to use divinyl

chlorophylls. Therefore, the reduced effective population size (N_e) upon the emergence of

Prochlorococcus was likely caused by a "founder effect", which suggests that a new

population was established by a few individuals from a larger ancestral population (Barton et

al., 1984). Note that this mechanism is different from a population bottleneck, which has been

used to explain the reduced N_e occurring on the branch leading to SBE-LCA (Zhang et al.,

2021). The latter conclusion is confirmed here with the inclusion of new SAG genomes, as

the d_R/d_C values in the branch leading to SBE-LCA were significantly elevated compared to

the those in the branches leading to *Prochlorococcus* LLIV and AMZ clades (Fig. 2b)

<u>Evolutionary drivers leading to the loss of phycobilisome and nitrate reductase in early</u>

<u>*Prochlorococcus*</u>

295   The phycobilisome constitutes as much as 50%-60% of the soluble proteins in

Synechococcus (Grossman et al., 1993). The loss of the phycobilisome was thought to reduce

nitrogen (N) investments by at least 40% in *Prochlorococcus* (Ting et al., 2002). Therefore,

the losses of phycobilisome genes was once considered to be favored by *Prochlorococcus*,

which conferred them advantages in inhabiting oligotrophic oceans (Ting et al., 2002).

However, since the colonization of *Prochlorococcus* at the deep photic zone of the Tonian

ocean conferred them advantages in acquiring the limiting nutrients like ammonium and

phosphate upwelled from the deep ocean, the loss of the phycobilisome in *Prochlorococcus*

was unlikely driven by selection for metabolic efficiency. Instead, since chlorophyll□based

light□harvesting complex gradually became the primary photosynthetic antenna in

Prochlorococcus, the phycobilisome genes were more likely subject to relaxed purifying

selection and thus neutral losses.

The absence of nitrate utilization genes (*narB* and *nrtB*) in cultured *Prochlorococcus*

strains has been attributed to biased taxon sampling, since the *Prochlorococcus* were

primarily cultured from ocean regions where P instead of N is the primarily limiting nutrient

(Berube et al., 2015). However, phylogenetic analysis that included both *Prochlorococcus*

and *Synechococcus* showed topological congruence of the *narB* gene tree with the species

tree, suggesting vertical inheritance of the gene (Berube et al., 2019). By including

Prochlorococcus SAGs in the present study, we confirmed the presence of the nitrate

utilization genes in early ancestral *Prochlorococcus* (Proch-AMZI/II/III-LCA). The

mechanism underlying the loss of nitrate utilization genes in *Prochlorococcus* was once

attributed to relaxed selection efficiency due to the low level of nitrate caused by the

intensive denitrification and anammox (anaerobic ammonium oxidation) activity (Canfield et

al., 2008; Johnston et al., 2009b). However, since *Prochlorococcus* originated in the deep

photic zone where ammonium would have been supplied through upwelling (Fig. 1c)

(Michiels et al., 2017), the loss of the nitrate utilization genes in *Prochlorococcus* is more

likely the result of superfluous nitrogen sources and a switch to uptake of ammonium instead

of nitrate, which was again a neutral process.

Implications for Neoproterozoic biogeochemical cycling

In accordance with the previous study (Ulloa et al., 2021), our analysis suggests that

Prochlorococcus diverged before transitioning from the use of the phycobilisome to the

chlorophyll□based light□harvesting complex. Therefore, while the divinyl pigment synthesis

gene was obtained at the earliest *Prochlorococcus* (i.e., Proch-AMZI/II/III-LCA) to enable

their exploration of the deep photic zone, our analyses imply that *Prochlorococcus* would not

have strongly influenced the Neoproterozoic Earth system until the emergence of Proch-

AMZI/II-LCA lineage during which the photosynthetic antenna was replaced. Our dating

analysis shows that the emergence of this lineage indeed coincided with the early stages of

Neoproterozoic Oxygenation Event (NOE).

The NOE is widely accepted as the second stage of Earth's protracted oxygenation,

during which atmospheric oxygen rose to near modern levels (Och and Shields-Zhou, 2012;

Lyons et al., 2014). This rise in atmospheric oxygen ultimately promoted the emergence of

large and complex animal life (Knoll and Nowak, 2017). The initiation of the NOE has been

previously attributed to the rise of eukaryotic algae (Lyons et al., 2014; Braakman et al., 2017;

Brocks et al., 2017). Our results, instead, suggest that early *Prochlorococcus* may have

played a role in enhancing oxygen production, with potential to initiate the NOE. This

hypothesis could be further tested through geochemical analyses that help refine estimates for

Tonian ocean oxygen contents and through biogeochemical models that can quantify the

potential effects of *Prochlorococcus* emergence on the global carbon and oxygen cycles.

Concluding remarks

We have integrated molecular clocks, evolutionary genetic analyses, and comparative

genomics along with knowledge of paleo-geochemistry to reconstruct the early evolution of

Prochlorococcus and infer their possible impact on the evolution of the Neoproterozoic Earth

system. The abundant fossil calibrations 'borrowed' from photosynthetic eukaryotes and the

new molecular dating pipeline provide robust dates for the origin of *Prochlorococcus* in the

Tonian oceans. Comparative genomics suggests that gains and losses of a few important

genes conferred the early ancestral *Prochlorococcus* with an unprecedented ability to absorb

the blue light that effectively penetrates water, empowering early *Prochlorococcus* to

colonize the deep photic zone, a niche distinctly different from the better illuminated surface

waters supporting other phytoplankton groups including early Eukaryotes and its

Synechococcus ancestor. The use of the evolutionary genetic proxy, d_R/d_C, implies that the

earliest *Prochlorococcus* had a highly reduced effective population size compared to its

Synechococcus ancestor, likely reflecting a founder effect and strengthening the idea that the

deep photic zone was an ecological niche uniquely accessible to the earliest *Prochlorococcus*.

This niche differentiation allowed the earliest *Prochlorococcus* to avoid fierce competition

with its *Synechococcus* ancestor and other phytoplankton groups and gave them easy access

to phosphate that otherwise strongly limited primary productivity in the Tonian oceans. Under

these favorable conditions, the early *Prochlorococcus* would have grown in population size,

promoting further evolution and the loss of the phycobilisome, which was replaced by a

chlorophyll-based light-harvesting system ultimately leading to enhanced oxygen production

and the possible initiation of the Neoproterozoic Oxygenation Event.

Materials and Methods

The nuclear-encoded and plastid-encoded protein sequences of eukaryotic representatives

were obtained from Dicots PLAZA (5.0) database and NCBI Refseq release of plastid

database and were used for the first-step and the second-step sequential Bayesian dating

analysis, respectively. The genomic sequences of Cyanobacteria (including *Prochlorococcus*

SAGs), Vampirovibrionia and Sericytochromatia were also used for the second-step

sequential dating analysis, which were obtained from GenBank and Integrated Microbial

Genomes (IMG) database (Supplementary Data 1).

For the first-step sequential dating analysis, the orthologous gene families shared by

eukaryotic species were identified by searching against a pre-compiled eukaryotic nuclear

gene dataset (Strassert et al., 2021) and were partitioned by using Gaussian Mixture Model

(GMM) clustering method based on their evolutionary rates. The posterior ages estimated

with these partitioned data were approximated to one of the skew-normal, skew-t and Gamma

distributions and then mapped back to the overlapping nodes in the species tree containing

both eukaryotes and Cyanobacteria as the time prior (Table S2).

For the second-step sequential dating analysis, the orthologous gene families were firstly

identified by searching against a pre-compiled plastid marker gene dataset (Ponce-Toledo et

al., 2017) and then sorted by ΔLL (a proxy for the extent to which a marker gene rejects

vertical evolution) and evolutionary rate difference (Supplementary Data 2) (Fig. S5). We

employed the program "mcmc3r" to select the gene family-specific clock model by following

published studies (McGowen et al., 2019; Wang and Luo, 2021). The calibrations and the

species tree topology used in the present study were also adapted from publications (see

details in Supplementary Methods). For those that remain disputed, we took all the

possibilities in our analysis for comparison (Fig. S3). We also tested the necessity for using

sequential Bayesian dating approach and hard bound calibration strategy in our analysis (Fig.

S6) and tested the convergence of the posterior age estimates (Fig. S7).

The gains and losses of *Prochlorococcus* orthologous gene families were inferred with a

gene tree-species tree reconciliation approach according to our recent studies (Zhang et al.,

2021; Zhang et al., 2023). The reduction of *Prochlorococcus* selection efficiency in ancient

time was inferred by comparing the genome-wide d_R/d_C values between the target lineage and

its sister lineage based on our recent study (Luo et al., 2017) (see details in Supplementary

Methods).

Code availability

The custom scripts used to analyze the data are available in the online GitHub repository

(https://github.com/luolab-cuhk/Prochl-NOE).

Acknowledgements

This work is supported by the Natural Science Foundation of China (42293294), the Hong

Kong Research Grants Council (RGC) General Research Fund (GRF) (14110820), the

Natural Science Foundation of Guangdong Province, China (2022A1515010844 to HZ), the

China Postdoctoral Science Foundation (2021M702296 to HZ), and the CUHK Direct Grant

(4053551).

References

Angelis, K., Álvarez-Carretero, S., Dos Reis, M., and Yang, Z. (2018) An Evaluation of Different Partitioning Strategies for Bayesian Estimation of Species Divergence Times. *Syst Biol* **67**: 61-77.

Aoki, S., Ito, M., Iwasaki, W.J.M.b., and evolution (2013) From β-to α-proteobacteria: the origin and evolution of rhizobial nodulation genes nodIJ. **30**: 2494-2508.

Barton, N.H., Charlesworth, B.J.A.r.o.e., and systematics (1984) Genetic revolutions, founder effects, and speciation. **15**: 133-164.

Berube, P.M., Rasmussen, A., Braakman, R., Stepanauskas, R., and Chisholm, S.W. (2019) Emergence of trait variability through the lens of nitrogen assimilation in Prochlorococcus. *eLife* **8**: e41043.

Berube, P.M., Biller, S.J., Kent, A.G., Berta-Thompson, J.W., Roggensack, S.E., Roache-Johnson, K.H. et al. (2015) Physiology and evolution of nitrate acquisition in Prochlorococcus. *The ISME journal* **9**: 1195.

Betts, H.C., Puttick, M.N., Clark, J.W., Williams, T.A., Donoghue, P.C., Pisani, D.J.N.e., and evolution (2018) Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. **2**: 1556-1562.

Bhattacharya, D., and Medlin, L.J.J.P. (1995) Phylogeny of plastids A review based on comparisons of small subunit ribosomal RNA coding regions. **31**: 487-496.

Biller, S.J., Berube, P.M., Lindell, D., and Chisholm, S.W. (2015) Prochlorococcus: the structure and function of collective diversity. *Nature Reviews Microbiology* **13**: 13.

Boden, J.S., Konhauser, K.O., Robbins, L.J., and Sánchez-Baracaldo, P. (2021) Timing the evolution of antioxidant enzymes in cyanobacteria. *Nature Communications* **12**: 1-12.

Braakman, R., Follows, M.J., and Chisholm, S.W. (2017) Metabolic evolution and the self-organization of ecosystems. *Proceedings of the National Academy of Sciences* **114**: E3091-E3100.

Brocks, J.J. (2018) The transition from a cyanobacterial to algal world and the emergence of animals. *Emerging Topics in Life Sciences* **2**: 181-190.

Brocks, J.J., Jarrett, A.J., Sirantoine, E., Hallmann, C., Hoshino, Y., and Liyanage, T. (2017) The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**: 578.

Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., and Strauss, H. (2008) Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. *Science* **321**: 949-952.

Chriki-Adeeb, R., and Chriki, A.J.E.B. (2016) Estimating divergence times and substitution rates in Rhizobia. **12**: EBO. S39070.

Clarkson, M.O., Sweere, T.C., Chiu, C.F., Hennekam, R., Bowyer, F., and Wood, R.A.J.E.-S.R. (2023) Environmental controls on very high δ238U values in reducing sediments: Implications for Neoproterozoic seawater records. 104306.

Dayhoff, M.O. (1972) A model of evolutionary change in proteins. *Atlas of protein sequence and structure* **5**: 89-99.

dos Reis, M., Gunnell, G.F., Barba-Montoya, J., Wilkins, A., Yang, Z., and Yoder, A.D. (2018) Using phylogenomic data to explore the effects of relaxed clocks and calibration strategies on divergence time estimation: primates as a test case. *Syst Biol* **67**: 594-615.

dos Reis, M., Thawornwattana, Y., Angelis, K., Telford, Maximilian J., Donoghue, Philip C.J., and Yang, Z. (2015) Uncertainty in the Timing of Origin of Animals and the Limits of Precision in Molecular Timescales. *Curr Biol* **25**: 2939-2950.

Duchêne, S., Lanfear, R., and Ho, S.Y.W. (2014) The impact of calibration and clock-model choice on molecular estimates of divergence times. *Mol Phylogen Evol* **78**: 277-289.

E. Zuckerkandl, L.P. (1965) Evolutionary divergence and convergence in proteins. In *Evolving genes and*
proteins: Elsevier, pp. 97-166.

Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincón, J., Zabala, L.L., Jiao, N. et al. (2013) Present and future
global distributions of the marine Cyanobacteria Prochlorococcus and Synechococcus. *Proceedings of the*
National Academy of Sciences **110**: 9824-9829.

Fournier, G., Moore, K., Rangel, L., Payette, J., Momper, L., and Bosak, T. (2021) The Archean origin of
oxygenic photosynthesis and extant cyanobacterial lineages. *Proceedings of the Royal Society B* **288**: 20210675.

Goericke, R., Olson, R., and Shalapyonok, A.J.D.S.R.P.I.O.R.P. (2000) A novel niche for Prochlorococcus sp. in
low-light suboxic environments in the Arabian Sea and the Eastern Tropical North Pacific. **47**: 1183-1205.

Gray, M.W.J.I.r.o.c. (1992) The endosymbiont hypothesis revisited. **141**: 233-357.

Grossman, A.R., Schaefer, M.R., Chiang, G.G., and Collier, J.L. (1993) The phycobilisome, a light-harvesting
complex responsive to environmental conditions. *Microbiol Mol Biol Rev* **57**: 725-749.

Heckman, D.S., Geiser, D.M., Eidell, B.R., Stauffer, R.L., Kardos, N.L., and Hedges, S.B.J.s. (2001) Molecular
evidence for the early colonization of land by fungi and plants. **293**: 1129-1133.

Hedges, S.B., Tao, Q., Walker, M., and Kumar, S.J.P.o.t.N.A.o.S. (2018) Accurate timetrees require accurate
calibrations. **115**: E9510-E9511.

Hess, W.R., Rocap, G., Ting, C.S., Larimer, F., Stilwagen, S., Lamerdin, J., and Chisholm, S.W. (2001) The
photosynthetic apparatus of Prochlorococcus: insights through comparative genomics. *Photosynthesis Res* **70**:
53-71.

Ito, H., and Tanaka, A. (2011) Evolution of a divinyl chlorophyll-based photosystem in Prochlorococcus.
Proceedings of the National Academy of Sciences **108**: 18014-18019.

Ito, H., Yokono, M., Tanaka, R., and Tanaka, A. (2008) Identification of a novel vinyl reductase gene essential
for the biosynthesis of monovinyl chlorophyll in Synechocystis sp. PCC6803. *J Biol Chem* **283**: 9002-9011.

Johnston, D.T., Wolfe-Simon, F., Pearson, A., and Knoll, A.H. (2009a) Anoxygenic photosynthesis modulated
Proterozoic oxygen and sustained Earth's middle age. *Proc Natl Acad Sci U S A* **106**: 16925-16929.

Johnston, D.T., Wolfe-Simon, F., Pearson, A., and Knoll, A.H. (2009b) Anoxygenic photosynthesis modulated
Proterozoic oxygen and sustained Earth's middle age. *Proceedings of the National Academy of Sciences* **106**:
16925-16929.

Jones, C., Nomosatryo, S., Crowe, S.A., Bjerrum, C.J., and Canfield, D.E.J.G. (2015) Iron oxides, divalent
cations, silica, and the early earth phosphorus crisis. **43**: 135-138.

Kang, J., Gill, B., Reid, R., Zhang, F., and Xiao, S. (2023) Nitrate limitation in early Neoproterozoic oceans
delayed the ecological rise of eukaryotes. *Science Advances* **9**: eade9647.

Keeling, P.J. (2013) The number, speed, and impact of plastid endosymbioses in eukaryotic evolution. *Annu Rev*
Plant Biol **64**: 583-607.

Knoll, A.H., and Nowak, M.A. (2017) The timetable of evolution. *Science Advances* **3**: e1603076.

Lau, K.V., Macdonald, F.A., Maher, K., and Payne, J.L. (2017) Uranium isotope evidence for temporary ocean
oxygenation in the aftermath of the Sturtian Snowball Earth. *Earth and Planetary Science Letters* **458**: 282-292.

Lavin, P., González, B., Santibáñez, J.F., Scanlan, D.J., and Ulloa, O.J.E.m.r. (2010) Novel lineages of
Prochlorococcus thrive within the oxygen minimum zone of the eastern tropical South Pacific. **2**: 728-738.

Luo, H., Huang, Y., Stepanauskas, R., and Tang, J. (2017) Excess of non-conservative amino acid changes in
marine bacterioplankton lineages with reduced genomes. *Nature microbiology* **2**: 17091.

Lyons, T.W., Reinhard, C.T., and Planavsky, N.J. (2014) The rise of oxygen in Earth's early ocean and
atmosphere. *Nature* **506**: 307.

Marin, B., M. Nowack, E.C., and Melkonian, M. (2005) A Plastid in the Making: Evidence for a Second
Primary Endosymbiosis. *Protist* **156**: 425-432.

Martiny, A.C., Kathuria, S., and Berube, P.M. (2009) Widespread metabolic potential for nitrite and nitrate
assimilation among Prochlorococcus ecotypes. *Proceedings of the National Academy of Sciences* **106**: 10787-
10792.

McGowen, M.R., Tsagkogeorga, G., Álvarez-Carretero, S., dos Reis, M., Struebig, M., Deaville, R. et al. (2019)
Phylogenomic Resolution of the Cetacean Tree of Life Using Target Sequence Capture. *Syst Biol* **69**: 479-501.

Michiels, C.C., Darchambeau, F., Roland, F.A., Morana, C., Llirós, M., García-Armisen, T. et al. (2017) Iron-
dependent nitrogen cycling in a ferruginous lake and the nutrient status of Proterozoic oceans. *Nature*
Geoscience **10**: 217-221.

Morris, J.L., Puttick, M.N., Clark, J.W., Edwards, D., Kenrick, P., Pressel, S. et al. (2018) The timescale of early
land plant evolution. **115**: E2274-E2283.

Och, L.M., and Shields-Zhou, G.A. (2012) The Neoproterozoic oxygenation event: Environmental perturbations
and biogeochemical cycling. *Earth-Sci Rev* **110**: 26-57.

Ozaki, K., Thompson, K.J., Simister, R.L., Crowe, S.A., and Reinhard, C.T.J.N.c. (2019) Anoxygenic
photosynthesis and the delayed oxygenation of Earth's atmosphere. **10**: 3026.

Palenik, B., Brahamsha, B., Larimer, F., Land, M., Hauser, L., Chain, P. et al. (2003) The genome of a motile
marine Synechococcus. **424**: 1037-1042.

Parfrey, L.W., Lahr, D.J., Knoll, A.H., and Katz, L.A.J.P.o.t.N.A.o.S. (2011) Estimating the timing of early
eukaryotic diversification with multigene molecular clocks. **108**: 13624-13629.

Partensky, F., and Garczarek, L. (2010) Prochlorococcus: advantages and limits of minimalism. *Annual Review*
of Marine Science **2**: 305-331.

Partensky, F., Hess, W.R., and Vaulot, D. (1999) Prochlorococcus, a marine photosynthetic prokaryote of global
significance. *Microbiol Mol Biol Rev* **63**: 106-127.

Ponce-Toledo, R.I., Deschamps, P., López-García, P., Zivanovic, Y., Benzerara, K., and Moreira, D. (2017) An
Early-Branching Freshwater Cyanobacterium at the Origin of Plastids. *Curr Biol* **27**: 386-391.

Ralf, G., and Repeta, D.J. (1992) The pigments of Prochlorococcus marinus: The presence of divinylchlorophyll
a and b in a marine procaryote. *Limnol Oceanogr* **37**: 425-433.

Rubio, L.M., Flores, E., and Herrero, A.J.J.o.b. (1998) The narA locus of Synechococcus sp. strain PCC 7942
consists of a cluster of molybdopterin biosynthesis genes. **180**: 1200-1206.

Sánchez-Baracaldo, P. (2015) Origin of marine planktonic cyanobacteria. *Scientific reports* **5**: 17418.

Sánchez-Baracaldo, P., Ridgwell, A., and Raven, J.A. (2014) A neoproterozoic transition in the marine nitrogen
cycle. *Curr Biol* **24**: 652-657.

Sánchez-Baracaldo, P., Raven, J.A., Pisani, D., and Knoll, A.H. (2017) Early photosynthetic eukaryotes
inhabited low-salinity habitats. *Proceedings of the National Academy of Sciences*: 201620089.

Sánchez-Baracaldo, P., Bianchini, G., Wilson, J.D., and Knoll, A.H. (2022) Cyanobacteria and biogeochemical
cycles through Earth history. *Trends Microbiol* **30**: 143-157.

Satoh, S., and Tanaka, A. (2006) Identification of chlorophyllide a oxygenase in the Prochlorococcus genome by
a comparative genomic approach. *Plant Cell Physiol* **47**: 1622-1629.

Schirrmeister, B.E., Gugger, M., and Donoghue, P.C. (2015) Cyanobacteria and the Great Oxidation Event:
evidence from genes and fossils. *Palaeontology* **58**: 769-785.

Schirrmeister, B.E., Sanchez-Baracaldo, P., and Wacey, D. (2016) Cyanobacterial evolution during the
Precambrian. *International Journal of Astrobiology* **15**: 187-204.

Shih, P.M., Hemp, J., Ward, L.M., Matzke, N.J., and Fischer, W.W. (2016) Crown group Oxyphotobacteria
postdate the rise of oxygen. *Geobiology* **15**: 19-29.
Stolper, D.A., and Keller, C.B.J.N. (2018) A record of deep-ocean dissolved O2 from the oxidation state of iron
in submarine basalts. **553**: 323-327.
Strassert, J.F.H., Irisarri, I., Williams, T.A., and Burki, F. (2021) A molecular timescale for eukaryote evolution
with implications for the origin of red algal-derived plastids. *Nat Commun* **12**: 1879.
Tahata, M., Sawaki, Y., Yoshiya, K., Nishizawa, M., Komiya, T., Hirata, T. et al. (2015) The marine
environments encompassing the Neoproterozoic glaciations: evidence from C, Sr and Fe isotope ratios in the
Hecla Hoek Supergroup in Svalbard. *Precambrian Research* **263**: 19-42.
Ting, C.S., Rocap, G., King, J., and Chisholm, S.W. (2002) Cyanobacterial photosynthesis in the oceans: the
origins and significance of divergent light-harvesting strategies. *Trends Microbiol* **10**: 134-142.
Tostevin, R., and Mills, B.J.J.I.F. (2020) Reconciling proxy records and models of Earth's oxygenation during
the Neoproterozoic and Palaeozoic. **10**: 20190137.

Trower, E.J., Gutoski, J.R., Wala, V.T., Mackey, T.J., and Simpson, C. (2023) Tonian Low-Latitude Marine

Ecosystems Were Cold Before Snowball Earth. *Geophysical Research Letters* **50**: e2022GL101903.
Ulloa, O., Henríquez-Castillo, C., Ramírez-Flandes, S., Plominsky, A.M., Murillo, A.A., Morgan-Lang, C. et al.
(2021) The cyanobacterium Prochlorococcus has divergent light-harvesting antennae and may have evolved in a
low-oxygen ocean. *Proceedings of the National Academy of Sciences* **118**.
Wang, S., and Luo, H. (2021) Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nature*
Communications **12**: 3324.
Warnock, R.C., Yang, Z., and Donoghue, P.C.J.B.l. (2012) Exploring uncertainty in the calibration of the
molecular clock. **8**: 156-159.
Zhang, F., Stockey, R.G., Xiao, S., Shen, S.-z., Dahl, T.W., Wei, G.-Y. et al. (2022) Uranium isotope evidence
for extensive shallow water anoxia in the early Tonian oceans. *Earth and Planetary Science Letters* **583**: 117437.
Zhang, H., Hellweger, F.L., and Luo, H.J.b. (2023) Genome reduction occurred in early Prochlorococcus with
an unusually low effective population size. 2023.2006. 2025.546417.
Zhang, H., Sun, Y., Zeng, Q., Crowe, S.A., and Luo, H. (2021) Snowball Earth, population bottleneck and
Prochlorococcus evolution. *Proceedings of the Royal Society B: Biological Sciences* **288**: 20211956.
Zhang, J. (2000) Rates of conservative and radical nonsynonymous nucleotide substitutions in mammalian
nuclear genes. *J Mol Evol* **50**: 56-68.
570

Figure Legends

Fig. 1 **The evolutionary timeline of *Prochlorococcus* estimated with plastid-based**

strategy. (a) The diagram shows the phylogenetic relations among Cyanobacteria and

eukaryotic species. The green solid line, black solid line and green dashed line in the

cladogram leading to the tip of oxygenic Cyanobacteria, anoxygenic Vampirovibrionia and

Sericytochromatia and eukaryotic species, respectively. The calibrated nodes in molecular

clock analysis are marked with orange circle (see calibration justifications in Supplementary

Methods). (b) (left) The *Prochlorococcus* evolutionary timeline estimated with the focal

molecular dating strategy using sequential Bayesian dating approach based on the 5-partition

eukaryotic genome-scale data (in the first step dating analysis) and fully partitioned 12-gene

dataset "T30" (in the second step dating analysis) under independent rate clock model. The

posterior age distribution is provided next to the ancestral node. The atmospheric oxygen

level at the geological time scale is adapted from Lyons et al., 2014, which is represented by

the percent of present atmospheric oxygen level (PAL). The vertical bars with green, orange

and blue colors represent the time of Tonian, the time of NOE, and the time of Sturtian (left)

and Marinoan (right) glaciation, respectively. (right) Phyletic pattern of key gene families.

Solid square, solid circle and open circle next to each analyzed genome represent multi-copy

gene family, single-copy gene family and absence of the gene family, respectively, in the

extant genomes. Gene families marked with asterisk (*) were consistently estimated to be

gained or lost by using both AnGST and GeneRax. (c) Diagram illustrating the

biogeochemical environments when *Prochlorococcus* arose from its *Synechococcus* ancestor.

Fig. 2　Diagram illustrating the schemes and the results of d_R/d_C comparison. The

genome-wide means of d_R/d_C values at the ancestral branches ('Target') leading to (a) Proch-

AMZI/II/III-LCA and (b) SBE-LCA compared with that at the sister lineages ('Control').

The d_R/d_C values were classified based on the physicochemical classification of the amino

acids by charge or by volume and polarity, and were either GC-corrected by codon frequency

(blue), GC-corrected by amino acid (AA) frequency (red) or uncorrected (gray). Error bars of

d_R/d_C represent the standard error of the mean.

Fig. 1

Fig. 2

(a)

- ■ GC-corrected (codon frequency)
- ■ GC-corrected (AA frequency)
- ■ Uncorrected

Proch-AMZI/II/III-LCA

HL II
HL I
LL I
LL II/III
LL IV
AMZ
Syne 5.1
Syne 5.2

Target
Control
Ref.

← SBE-LCA
← Proch-AMZI/II/III-LCA

Charge

Volume&Polarity

Target   Control

(b) SBE-LCA

HL II
HL I
LL I
LL II/III
LL IV
AMZ
Syne 5.1
Syne 5.2

Target
Control
Reference

Charge

Volume&Polarity

Target   Control