1

Emergence of *Prochlorococcus* in the Tonian oceans and the initiation of Neoproterozoic

oxygenation

Hao Zhang[1,2^], Sishuo Wang[2^], Tianhua Liao[2], Sean A. Crowe[3], Haiwei Luo[2,4,5*]

[1]Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen 518000, China

[2]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[3]Departments of Microbiology and Immunology, and Earth, Ocean, and Atmospheric Sciences, University of British Columbia, Vancouver, British Columbia, Canada

[4]Earth and Environmental Sciences Programme, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[5]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[^] These authors contributed equally to this study

[*]**Corresponding author:** Haiwei Luo

Email: haiweiluo@cuhk.edu.hk

Competing Interest Statement: The authors declare no competing interests.

Key words: *Prochlorococcus*, Neoproterozoic oxygenation event, molecular dating, plastid, cyanobacteria

Abstract

Prochlorococcus are the smallest and most abundant photosynthetic organisms on Earth,

contributing up to 50% of the chlorophyll in the oligotrophic oceans. Despite being important

in regulating the carbon cycle in today's ocean, the ecological significance of

Prochlorococcus in Earth's history remains elusive. Our new robustly calibrated molecular

clock reveals that *Prochlorococcus* emerged in the deep photic zone of the Tonian oceans.

The classical antenna for light harvesting in Cyanobacteria, the phycobilisome, was replaced

in *Prochlorococcus* by the chlorophyll□based light□harvesting complex enabling more

efficient use of blue light that penetrated to deeper water. Importantly, *Prochlorococcus*

colonization of deep water enhanced access to phosphate, which was rich in upwelled

seawater, but likely scarce in the Tonian surface ocean, promoting expansion of

Prochlorococcus, displacement of incumbent low-light adapted anoxygenic photoferrotrophs,

and associated increases in photosynthetic oxygen production. Colonization of deeper waters

would also have improved access to ammonium, leading to the neutral loss of nitrate

utilization genes. Our research thus documents the conspicuous emergence of new

photosynthetic bacterial lineages in the run-up to the Neoproterozoic oxygenation event,

implying an additional layer of eco-evolutionary complexity during this pivotal interval in

Earth's history.

Introduction

Prochlorococcus host more than half of the chlorophyll biomass in oligotrophic oceans (Partensky and Garczarek, 2010). As key primary producers in the modern ocean, Prochlorococcus fix as much as four gigatons of carbon each year and are the foundation of the marine carbon cycle and food web (Flombaum et al., 2013). The success of Prochlorococcus in today's oceans has been attributed to multiple physiological features. Importantly, Prochlorococcus use divinyl chlorophyll (DVChl) for harnessing light energy (Ralf and Repeta, 1992). DVChl harvests blue light much more efficiently than the more common monovinyl chlorophyll used by other cyanobacterial lineages and it thus enables Prochlorococcus to thrive in the deepest layers of the euphotic zone, where blue light dominates (Ito and Tanaka, 2011). Prochlorococcus is also the smallest photosynthetic group on Earth (Partensky et al., 1999). As a result, their high surface-to-volume ratio provides enhanced nutrient acquisition efficiency, which together with effective blue-light absorption promotes photosynthesis by Prochlorococcus in oligotrophic tropical and subtropical oceans (Partensky et al., 1999). In today's ocean, Prochlorococcus is also the dominant phototroph in more nutrient-rich, oxygen-depleted anoxic marine zones (AMZ) in the eastern tropical North and South Pacific Oceans (Goericke et al., 2000; Lavin et al., 2010). Since the AMZ lineages represent the earliest-split branches of Prochlorococcus, it has been proposed that Prochlorococcus emerged in low-oxygen environments and, by extension, contributed to early ocean oxygenation (Ulloa et al., 2021).

The emergence of an early branch of Prochlorococcus (named SBE-LCA) during the Cryogenian based on our recent study (Zhang et al. 2021) implies the origin of the total

Prochlorococcus group earlier in the Proterozoic Eon. Such an earlier origin then suggests

that *Prochlorococcus* might have contributed to the cyanobacterial dominated primary

production that supported marine biogeochemical cycles and underpinned dynamic ocean

chemistry in the run-up to the Sturtian Snowball Earth glaciation. This was an important

period in Earth's history that ultimately led to an Earth system succession and the rise of

eukaryotes with algae emerging as key primary producers before the end of the Cryogenian

(Brocks et al. 2017). Deciphering the potential role, if any, that *Prochlorococcus* might have

played depends critically on more precise age dates for the divergence of the total

Prochlorococcus group from its *Synechococcus* ancestors.

By far, however, the divergence time of *Prochlorococcus* remains contentious. Different

relaxed clock analyses based on 16S and 23S rRNA genes yielded divergence times of the

total *Prochlorococcus* group at ~200 Mya (Boden et al., 2021), ~500 Mya (Sánchez-

Baracaldo, 2015) and ~800 Mya (Sánchez-Baracaldo et al., 2014). The use of protein-coding

genes gave divergence times of the total *Prochlorococcus* group at ~400 Mya (Fournier et al.,

2021), ~500 Mya (Sánchez-Baracaldo et al., 2017) and ~1,000 Mya (Schirrmeister et al.,

2015). The discrepancies in *Prochlorococcus* divergence times are likely caused by variable

use of fossils and other time calibrations and the use of alternative gene sets, clock models,

and tree topologies (Warnock et al., 2012; Duchêne et al., 2014; dos Reis et al., 2015) (see

Supplementary Discussion). Despite their importance, these factors were rarely tested

rigorously thus making it difficult to evaluate the accuracy of age estimates in previous

studies. Importantly, even if all these factors are well tested and controlled, the rarity of

cyanobacterial fossils and the absence of a maximum boundary of the available

cyanobacterial fossils poses an appreciable challenge in determining the antiquity of

Prochlorococcus (Zhang et al., 2021). Indeed, applying maximum age constraints at the

calibration nodes strongly impacts posterior age estimates of uncalibrated lineages (Hedges et

al., 2018; Morris et al., 2018; Wang and Luo, 2021). However, informative maximum age

constraints are typically missing from microbial fossils and can only be found in some animal

and plant fossils. Moreover, the recent availability of several genome sequences of

uncultivated basal *Prochlorococcus* lineages (Ulloa et al., 2021) requires re-estimation of the

antiquity of *Prochlorococcus*, which should also include an evaluation of the factors that

influence the accuracy of posterior ages.

Here, we leverage the abundant plant and algal fossils and the well-established plant

plastid endosymbiosis theory to develop a new pipeline that systematically tests the factors

that influence the accuracy of posterior ages and thus yields robust estimates of

Prochlorococcus antiquity. The plant plastid endosymbiosis theory states that the origin of all

plastids in photosynthetic eukaryotes, except for the photosynthetic amoeba *Paulinella*

(Marin et al., 2005), can be traced back to an ancient primary endosymbiosis involving a

eukaryote and a cyanobacterium (Gray, 1992; Bhattacharya and Medlin, 1995; Keeling, 2013;

Ponce-Toledo et al., 2017). Plant plastid endosymbiosis theory thus ties the evolutionary

histories of cyanobacteria to those of photosynthetic eukaryotes. Eukaryotic fossils are indeed

being increasingly used in dating the evolution of Cyanobacteria (Shih et al., 2016; Sánchez-

Baracaldo et al., 2017; Fournier et al., 2021). Our strategy builds on these by: 1) using more

fossil-based age constraints (including maximum age constraints) on eukaryotic lineages with

more complete taxonomic sampling including lineages that have undergone secondary

endosymbiosis; and 2) applying a sequential Bayesian dating approach to better constrain the

divergence time of eukaryotic lineages thereby propagating more accurate time information

to Cyanobacteria, including *Prochlorococcus*. These joint analyses led us to conclude that

Prochlorococcus arose in the Tonian oceans. Further, the use of an evolutionary genetic proxy

implies that *Prochlorococcus* was born through a founder effect, and this strengthens the idea

that *Prochlorococcus* emerged in the deep photic zone, a unique niche separated from upper

waters where its *Synechococcus* ancestors thrived.

Results

<u>*Prochlorococcus* originated in the Tonian ocean</u>

We implemented a sequential Bayesian dating method that takes advantage of the

abundant fossil records (Fig. 1a; Fig. S1a) available from photosynthetic eukaryotes to

calibrate the evolution of Cyanobacteria. The last common ancestor (LCA) of

Prochlorococcus (denoted as "Proch-AMZI/II/III-LCA") clearly emerged within the Tonian

period at 878 Mya (95% HPD: 987-767 Mya) (Fig. 1b). Meanwhile, the LCA of

Prochlorococcus clades HL, LL and AMZI/II (denoted as "Proch-AMZI/II-LCA") evolved at

787 Mya (95% HPD: 896-687 Mya), which coincided with the early stage of the

Neoproterozoic Oxygenation Event (NOE) (Fig. 1b). In this case, the branch leading to the

LCA of *Prochlorococcus* HL, LLI, and LLII/III clades (denoted as "SBE-LCA" to keep

consistency with (Zhang et al., 2021)) spanned the time that encompassed the duration of the

Neoproterozoic Snowball Earth events (645-635 Mya for Marinoan glaciation and 717-659

Mya for Sturtian glaciation), supporting the main conclusion of the previous study, which

used the traditional cyanobacterial fossil-based strategy (Zhang et al., 2021).

Note that the sequential Bayesian method used here is different from the commonly used

secondary dating method (using time estimates from previous studies as calibrations)

(Heckman et al., 2001; Aoki et al., 2013; Chriki-Adeeb and Chriki, 2016) for two reasons.

First, the time prior used in sequential dating analysis follows a specific probability

distribution, but secondary calibrations ignore the uncertainties on age estimates. Second, the

sequential dating analysis contains two steps each based on a distinct set of molecular data,

while secondary dating analysis largely relies on the same molecular dataset (gene alignments)

(dos Reis et al., 2018).

In our implementation, the posterior ages of eukaryotic lineages derived from the first-

step of the sequential analysis were used as the time priors to calibrate Cyanobacteria

evolution where few time constraints are available. Since genome-scale dating of eukaryotic

lineages is very slow with fully partitioned molecular data and is prone to have reduced

coverage probability (the probability that the 95% HPD interval of posterior ages contains the

true age) due to the large number of partitions (Angelis et al., 2018), we first clustered the

eukaryotic gene families into 1-, 3-, 5-, 7-, and 9-partitions based on their evolutionary rates

(see details in Supplementary Methods). In general, the posterior ages of eukaryotic lineages

were consistent, except for the case where a single partition and a single substitution model

was used, which did not fit the data well as indicated by its larger Akaike Information

Criterion (AIC) value (Fig. S1b). In all other cases, the crown group of Archaeplastida was

estimated to evolve at ~1.60 Gya, which is consistent with the estimates from other studies

(Parfrey et al., 2011; Betts et al., 2018; Wang and Luo, 2021) (Fig. S1b). By comparing the

distributions of the posterior ages of eukaryotic nodes in the first-step of sequential analysis

to the effective prior on the corresponding nodes in the second-step of analysis (see details in

Supplementary Methods), we found that the two distributions in dating analyses were nearly

identical when 5-, 7- and 9-partition data were used (Fig. S2), suggesting that the sequential

Bayesian approach works well for these analyses. We therefore adopted the evolutionary

timeline of eukaryotes based on the 5-partition data (lowest AIC value and high similarity

between posterior ages and effective prior) and applied the posterior ages of eukaryotic

lineages as time prior at the overlapping nodes in a larger phylogenomic tree that includes

both photosynthetic eukaryotes and cyanobacteria (Fig. S1c).

To validate the evolutionary timeline of *Prochlorococcus*, we performed a series of tests

by using different molecular data, clock models, fossil calibrations and species tree

topologies. The posterior ages estimated with all these alternative settings are largely

consistent with that estimated with the focal strategy (detailed above) and fully support the

origin of *Prochlorococcus* (Proch-AMZI/II/III-LCA) in the Tonian period, followed by the

emergence of Proch-AMZI/II-LCA in the early stages of the NOE (Fig. S3). Moreover, the

branch leading to the SBE-LCA encompasses the Snowball Earth events in all the dating

analyses (Fig. S4; see Supplementary Results). Our study also results in an estimate of

Cyanobacteria age, which has been an important research topic and under heavy debate

(Schirrmeister et al., 2015; Schirrmeister et al., 2016; Fournier et al., 2021; Sánchez-

Baracaldo et al., 2022). In this case, we found that, while the posterior ages of the crown

Cyanobacteria group likely influenced by the root maximum age, the origin time of

Prochlorococcus (Proch-AMZI/II/III-LCA) was consistently estimated at ~900 Mya (Fig. S5;

see Supplementary Discussion).

Prochlorococcus genomic changes

We inferred the genes gained and lost upon the emergence of *Prochlorococcus* first

without using the SAGs to avoid the false prediction of genomic events due to the relatively

low completeness of the SAGs (e.g., 81.9% for AMZI-B-ETNP). Our reconstructions rely on

the gene tree-species tree reconciliation approach, in which a best-fit reconciliation tool was

selected through a comprehensive simulation-based benchmarking analysis [see Fig. S4 in

(Zhang et al., 2023)]. We then used the same reconstruction methods to validate these

evolutionary events when SAGs were included.

As the only phytoplankton group using divinyl chlorophyll (DVChl) for harvesting light

energy (Ralf and Repeta, 1992), *Prochlorococcus* lost the gene *bciB* that inhibits the

conversion of DVChl to MVChl (monovinyl chlorophyll) at Proch-AMZI/II/III-LCA, thereby

promoting the accumulation of DVChl in their membranes (Ito et al., 2008). Compared to the

MVChl used by *Synechococcus*, DVChl more efficiently absorbs blue light that penetrates to

deeper waters than other photosynthetically active radiation (Ito and Tanaka, 2011). Likewise,

the gain of the *PcCao* gene for the synthesis of chlorophyll b (Satoh and Tanaka, 2006) in

Proch-AMZI/II/III-LCA also enables *Prochlorococcus* ancestors to efficiently harvest blue

light at exceedingly low intensities characteristic of deep waters (Hess et al., 2001) (Fig. 1b).

These genomic changes would have enabled *Prochlorococcus* ancestors to explore the deep

photic zone, a niche where its *Synechococcus* ancestor would not thrive.

Unlike most cyanobacteria, which use the phycobilisome as the photosynthetic antenna,

the main light-harvesting antenna of *Prochlorococcus* is made up of prochlorophyte

chlorophyll-binding (Pcb) protein (Biller et al., 2015). By reconstructing the gene

evolutionary paths with *Prochlorococcus* SAGs included, we found that the phycobilisome

genes (*apcACDEF*, *cpcEFG*, and *cpeCES*) were present in both Proch-AMZI/II/III-LCA and

Proch-AMZI/II-LCA and lost at SBE-LCA. Using the same approach, we found that the

cholorophyll-binding protein (encoded by *pcbD*) was obtained at Proch-AMZI/II-LCA. The

replacement of the photosynthetic antenna thus did not co-occur with the emergence of

Prochlorococcus. We note that gene absence in SAGs could be due to the incomplete nature

of the SAG genomes, however, in this case, simultaneous absence of the *pcbD* gene in all the

five SAGs that are more than 80% complete seems unlikely.

Despite the absence of the nitrate utilization genes in most *Prochlorococcus* isolates,

some uncultivated *Prochlorococcus* lineages contain these genes (Martiny et al., 2009;

Berube et al., 2015; Berube et al., 2019). By including *Prochlorococcus* SAGs that contain

the assimilatory nitrate transporter gene (*nrtB*) and the assimilatory nitrate reductase gene

(*narB*) in our reconstructions, we inferred the presence of both genes in Proch-AMZI/II/III-

LCA and Proch-AMZI/II-LCA and the losses of these genes at SBE-LCA (Fig. 1b). In

addition to the nitrate utilization genes, we inferred the losses of molybdopterin biosynthesis

genes at SBE-LCA. Since these genes are known to co-locate with the nitrate reductase genes

in *Synechococcus* (Rubio et al., 1998; Palenik et al., 2003), they may function as the cofactor

of nitrate reductase in *Prochlorococcus*. Therefore, the loss of nitrate reductase in

Prochlorococcus likely rendered the molybdopterin dispensable and thus led to the losses of

molybdopterin biosynthesis genes (*moaABCDE*, *mobA* and *moeA*) at SBE-LCA.

<u>Selection efficiency was reduced upon the emergence of *Prochlorococcus*</u>

The method for inferring selection efficiency in deep time was developed (Zhang, 2000)

and recently improved (Luo et al., 2017), which involves nonsynonymous substitutions only

and compares the rate of nonsynonymous substitutions leading to replacements of

physicochemically dissimilar amino acids (i.e., radical changes; d_R) to the rate of

nonsynonymous substitutions leading to replacements of physicochemically similar amino

acids (i.e., conservative changes; d_C). Since radical changes are more likely to be deleterious

than conservative changes (E. Zuckerkandl, 1965; Dayhoff, 1972), an excess of the radical

changes in a deeply branching lineage compared to its sister lineage suggests random fixation

of deleterious mutations by genetic drift in the former. Using this method, we found a

significant increase of the d_R/d_C ratio across the genomic regions in the branch leading to

Proch-AMZI/II/III-LCA relative to the branch leading to the LCA of *Synechococcus* clade 5.1

(Fig. 2a), indicating that the emergence of *Prochlorococcus* was accompanied by a significant

reduction of the efficiency of purifying selection, allowing for an accelerated accumulation of

deleterious mutations through genetic drift.

Discussion

<u>Geochemical context that supports the emergence of *Prochlorococcus* in the Tonian period</u>

At the 95% HPD, *Prochlorococcus* emerged between 989 to 770 Mya, encompassing

almost the entire Tonian period. A Tonian, pre-Sturtian, age for the emergence of the

Prochlorococcus lineage would have been set against the backdrop of dynamic ocean

chemistry characterized by a variably oxygenated surface ocean that was co-populated by

diverse microbial eukaryotes, including phytoplankton and the earliest metazoans (Trower et

al., 2023) (Fig. 1c). Multi-proxy data, collected from geographically diverse sites, converge

on Tonian deep oceans that were pervasively anoxic across 60-80% of the ocean floor, or

more (Tahata et al., 2015; Lau et al., 2017). The deep anoxic oceans were predominantly

ferruginous in nature, albeit with evidence for transient, spatially restricted euxinia, and

possibly even brief (<0.5 Myr) episodes of pervasive deep ocean oxygenation (Stolper and

Keller, 2018; Tostevin and Mills, 2020). Shallow waters, by contrast, were modestly, though

perhaps increasingly, oxygenated across the Tonian with evidence for the episodic and

persistent intrusion of anoxic deep waters into the surface oceans (Zhang et al., 2022;

Clarkson et al., 2023), with corresponding implications for nutrient cycling and availability

and for marine life in the euphotic zone.

The strongest differentiators between the earliest *Prochlorococcus* (i.e., Proch-

AMZI/II/III-LCA) and its *Synechococcus* ancestors are adaptation to low-light and loss of

capacity for nitrate assimilation, which imply both a deeper photosynthetic niche and a likely

shift to use of ammonium as the primary nitrogen source. A key driver of *Prochlorococcus*

emergence was thus likely a competitive advantage to dwelling in deeper waters. A well-

known advantage emerges under conditions of strong P-limitation, whereby deep-dwelling

low-light adapted phototrophs gain first access to upwelling phosphorus (Jones et al., 2015;

Ozaki et al., 2019). The low-light advantage conferred on the earliest *Prochlorococcus* cells

through the adoption of divinyl chlorophyll would have enhanced their access to P, relative to

contemporary algae and their *Synechococcus* ancestors. It would also have increased access

to NH_4^+ upwelling from deeper anoxic waters obviating the need for nitrate assimilation

(Michiels et al., 2017) (Fig. 1c).

In this way, the emergence of the *Prochlorococcus* would have restricted the flux of

nutrients to shallower waters and limited the contributions of higher-light adapted

phytoplankton like algae to primary production. This is supported through the biomarker

record, which indicates a dominance of cyanobacterial primary production until after the

Sturtian glaciation (Brocks, 2018). It is also supported by the Tonian N-isotope record, which

implies a limited contribution from nitrate assimilation to primary production at this time

(Kang et al., 2023). Importantly, the adaptation of the *Prochlorococcus* LCA to lower-light

would also allow it to better compete with low-light adapted anoxygenic phototrophs that

would likely have populated anoxic regions of the Tonian euphotic zone with strong potential

to cause ocean oxygenation (Johnston et al., 2009; Jones et al., 2015; Ozaki et al., 2019).

Competition between oxygenic photosynthetic cyanobacteria and anoxygenic phototrophs

that oxidize ferrous iron (photoferrotophs), is known to limit photosynthetic oxygen

production (Jones et al., 2015; Ozaki et al., 2019). Whereas the accumulation of O_2 produced

through photosynthesis ultimately depends on organic matter burial (Berner, 1991), oxygen

production fluxes through organic matter burial scale with the fraction of total photosynthetic

production that is oxygenic rather than anoxygenic (Johnston et al., 2009; Ozaki et al., 2019).

In this way, the capacity of the *Prochlorococcus* LCA to grow in deeper waters would have

enhanced its ability to access upwelling phosphorus thereby increasing the oxygenic fraction

of total photosynthesis with potential to initiate a positive feedback on oxygenation (Ozaki et

al., 2019).

<u>Evolutionary mechanisms underlying the origin of *Prochlorococcus*</u>

We propose that the colonization of the deep photic zone by *Prochlorococcus* started by a

seed population that obtained the ability to use divinyl chlorophylls. Therefore, the reduced

effective population size (N_e) upon the emergence of *Prochlorococcus* was likely caused by a

"founder effect", which suggests that a new population was established by a few individuals

from a larger ancestral population (Barton et al., 1984). Note that this mechanism is different

from a population bottleneck, which has been used to explain the reduced N_e occurring on the

branch leading to SBE-LCA (Zhang et al., 2021). The latter conclusion is confirmed here

with the inclusion of new SAG genomes, as the d_R/d_C values in the branch leading to SBE-

LCA were significantly elevated compared to the those in the branches leading to

Prochlorococcus LLIV and AMZ clades (Fig. 2b)

<u>Evolutionary drivers leading to the loss of phycobilisome and nitrate reductase in early</u>

<u>*Prochlorococcus*</u>

The phycobilisome constitutes as much as 50%-60% of the soluble proteins in

Synechococcus (Grossman et al., 1993). The loss of the phycobilisome was thought to reduce

nitrogen (N) investments by at least 40% in *Prochlorococcus* (Ting et al., 2002). Therefore,

the losses of phycobilisome genes was once considered to be favored by *Prochlorococcus*

conferred advantages to inhabiting oligotrophic oceans (Ting et al., 2002). However, since the

colonization of *Prochlorococcus* in the deep photic zone of the Tonian ocean conferred them

advantages to acquiring the limiting nutrients like ammonium and phosphate upwelled from

the deep ocean, the loss of the phycobilisome in *Prochlorococcus* was unlikely driven by

selection for metabolic efficiency. Instead, as the chlorophyll□based light□harvesting

complex gradually became the primary photosynthetic antenna in *Prochlorococcus*, the

phycobilisome genes were more likely subject to relaxed purifying selection and thus neutral

losses.

The absence of nitrate utilization genes (*narB* and *nrtB*) in cultured *Prochlorococcus*

strains has been attributed to biased taxon sampling, since the *Prochlorococcus* were

primarily cultured from ocean regions where P instead of N is the primarily limiting nutrient

(Berube et al., 2015). However, phylogenetic analysis that included both *Prochlorococcus*

and *Synechococcus* showed topological congruence of the *narB* gene tree with the species

tree, suggesting vertical *narB* gene inheritance (Berube et al., 2019). By including

Prochlorococcus SAGs in our study, we confirmed the presence of the nitrate utilization

genes in early ancestral *Prochlorococcus* (Proch-AMZI/II/III-LCA). The mechanism

underlying the loss of nitrate utilization genes in *Prochlorococcus* was once attributed to

relaxed selection efficiency due to the low level of nitrate caused by intensive denitrification

and anammox (anaerobic ammonium oxidation) activity (Canfield et al., 2008; Johnston et al.,

2009). However, since *Prochlorococcus* originated in the deep photic zone where ammonium

would likely have been supplied through upwelling (Fig. 1c) (Michiels et al., 2017), the loss

of the nitrate utilization genes in *Prochlorococcus* is more likely the result of superfluous

nitrogen sources and a switch to uptake of ammonium instead of nitrate, which was again a

neutral process.

<u>Implications for Neoproterozoic biogeochemical cycling</u>

In accordance with previous research (Ulloa et al., 2021), our analysis suggests that

Prochlorococcus diverged before transitioning from the use of the phycobilisome to the

chlorophyll☐based light☐harvesting complex. Therefore, while the divinyl pigment synthesis

gene was obtained at the earliest *Prochlorococcus* (i.e., Proch-AMZI/II/III-LCA) to enable

their exploration of the deep photic zone, our analyses imply that *Prochlorococcus* would not

have strongly influenced the Neoproterozoic Earth system until the emergence of Proch-

AMZI/II-LCA lineage during which the photosynthetic antenna was replaced. Our dating

analysis shows that the emergence of this lineage indeed broadly coincides with geochemical

signals for the early stages of Neoproterozoic Oxygenation Event (NOE).

The NOE is widely considered the second stage of Earth's protracted oxygenation history,

during which atmospheric and marine oxygen concentrations rose to levels exceeding those

that characterized the proceeding mid-Proterozoic (Och and Shields-Zhou, 2012; Lyons et al.,

2014). This rise in atmospheric oxygen ultimately promoted the emergence of large and

complex animal life (Knoll and Nowak, 2017). The initiation of the NOE has been previously

attributed to the rise of eukaryotic algae (Lyons et al., 2014; Braakman et al., 2017; Brocks et

al., 2017). Our results, instead, suggest that early *Prochlorococcus* may have played a role in

enhancing oxygen production, with potential to initiate the NOE, which was likely ultimately

accelerated through increasing efficiency in the biological carbon pump driven by a

progressively a larger role for eukaryotes. Such an elevated role for eukaryotes is clearly

marked in the geologic record through an increase in sterane-hopane ratios following the

Sturtian glaciation (Brocks et al., 2017). Pre-Sturtian, Tonian oxygenation, was thus likely

driven by bacterially dominated carbon cycling, the timing of which strongly coincides with

the emergence of low-light adapted *Prochlorococcus*, their ensuing colonization of the deep

photic zone, and capacity to displace incumbent anoxygenic photoferrotrophs thereby

increasing oxygen production. This hypothesis could be further tested through geochemical

analyses that help refine estimates for Tonian ocean oxygen contents and through

biogeochemical models that can quantify the potential effects of *Prochlorococcus* emergence

on the global carbon and oxygen cycles.

Concluding remarks

We have integrated molecular clocks, evolutionary genetic analyses, and comparative

genomics along with knowledge of paleo-geochemistry to reconstruct the early evolution of

Prochlorococcus and infer their possible impact on the evolution of the Neoproterozoic Earth

system. The abundant fossil calibrations 'borrowed' from photosynthetic eukaryotes and the

new and robust molecular dating pipeline pinpoint the origin of *Prochlorococcus* to the

Tonian oceans. Comparative genomics suggests that gains and losses of a few important

genes conferred the early ancestral *Prochlorococcus* with an unprecedented ability to absorb

the blue light that effectively penetrates water, empowering early *Prochlorococcus* to

colonize the deep photic zone, a niche distinctly different from the better illuminated surface

waters supporting other phytoplankton groups including its *Synechococcus* ancestor and early

Eukaryotes. The use of the evolutionary genetic proxy, d_R/d_C, implies that the earliest

Prochlorococcus had a highly reduced effective population size compared to its

Synechococcus ancestor, likely reflecting a founder effect and strengthening the idea that the

deep photic zone was an ecological niche uniquely accessible to the earliest *Prochlorococcus*.

This niche differentiation allowed the earliest *Prochlorococcus* to avoid fierce competition

with its *Synechococcus* ancestor and other phytoplankton groups and gave them easy access

to phosphate that otherwise strongly limited primary productivity in the Tonian oceans. Under

these favorable conditions, the early *Prochlorococcus* would have grown in population size,

promoting further evolution and the loss of the phycobilisome, which was replaced by a

chlorophyll-based light-harvesting system ultimately leading to enhanced oxygen production

and the possible initiation of the Neoproterozoic Oxygenation Event.

Materials and Methods

The nuclear-encoded and plastid-encoded protein sequences of eukaryotic representatives

were obtained from Dicots PLAZA (5.0) database and NCBI Refseq release of plastid

database and were used for the first-step and the second-step sequential Bayesian dating

analysis, respectively. The genomic sequences of Cyanobacteria (including *Prochlorococcus*

SAGs), Vampirovibrionia and Sericytochromatia were also used for the second-step

sequential dating analysis, which were obtained from GenBank and Integrated Microbial

Genomes (IMG) database (Supplementary Data 1).

For the first-step sequential dating analysis, the orthologous gene families shared by

eukaryotic species were identified by searching against a pre-compiled eukaryotic nuclear

gene dataset (Strassert et al., 2021) and were partitioned by using Gaussian Mixture Model

(GMM) clustering method based on their evolutionary rates. The posterior ages estimated

with these partitioned data were approximated to one of the skew-normal, skew-t and Gamma

distributions and then mapped back to the overlapping nodes in the species tree containing

both eukaryotes and Cyanobacteria as the time prior (Table S2).

For the second-step sequential dating analysis, the orthologous gene families were firstly

identified by searching against a pre-compiled plastid marker gene dataset (Ponce-Toledo et

al., 2017) and then sorted by ΔLL (a proxy for the extent to which a marker gene rejects

vertical evolution) and evolutionary rate difference (Supplementary Data 2) (Fig. S6). We

employed the program "mcmc3r" to select the gene family-specific clock model by following

published studies (McGowen et al., 2019; Wang and Luo, 2021). The calibrations and the

species tree topology used in the present study were also adapted from publications (see

details in Supplementary Methods). For those that remain disputed, we took all the

possibilities in our analysis for comparison (Fig. S3). We also tested the necessity for using

sequential Bayesian dating approach and hard bound calibration strategy in our analysis (Fig.

S7) and tested the convergence of the posterior age estimates (Fig. S8).

The gains and losses of *Prochlorococcus* orthologous gene families were inferred with a

gene tree-species tree reconciliation approach according to our recent studies (Zhang et al.,

2021; Zhang et al., 2023) (Supplementary Data 3). The reduction of *Prochlorococcus*

selection efficiency in ancient time was inferred by comparing the genome-wide d_R/d_C values

between the target lineage and its sister lineage based on our recent study (Luo et al., 2017)

(see details in Supplementary Methods).

Code availability

The custom scripts used to analyze the data are available in the online GitHub repository

(https://github.com/luolab-cuhk/Prochl-NOE).

Acknowledgements

This work is supported by the Natural Science Foundation of China (42293294), the Hong

Kong Research Grants Council (RGC) General Research Fund (GRF) (14110820), the

Natural Science Foundation of Guangdong Province, China (2022A1515010844 to HZ), the

China Postdoctoral Science Foundation (2021M702296 to HZ), and the CUHK Direct Grant

(4053551).

References

Angelis, K., Álvarez-Carretero, S., Dos Reis, M., and Yang, Z. (2018) An Evaluation of Different Partitioning
Strategies for Bayesian Estimation of Species Divergence Times. *Syst Biol* **67**: 61-77.

Aoki, S., Ito, M., Iwasaki, W.J.M.b., and evolution (2013) From β-to α-proteobacteria: the origin and evolution
of rhizobial nodulation genes nodIJ. **30**: 2494-2508.

Barton, N.H., Charlesworth, B.J.A.r.o.e., and systematics (1984) Genetic revolutions, founder effects, and
speciation. **15**: 133-164.

Berner, R.A. (1991) A model for atmospheric CO_2 over Phanerozoic time. *American Journal of Science* **291**:
339-376.

Berube, P.M., Rasmussen, A., Braakman, R., Stepanauskas, R., and Chisholm, S.W. (2019) Emergence of trait
variability through the lens of nitrogen assimilation in Prochlorococcus. *eLife* **8**: e41043.

Berube, P.M., Biller, S.J., Kent, A.G., Berta-Thompson, J.W., Roggensack, S.E., Roache-Johnson, K.H. et al.
(2015) Physiology and evolution of nitrate acquisition in Prochlorococcus. *The ISME journal* **9**: 1195.

Betts, H.C., Puttick, M.N., Clark, J.W., Williams, T.A., Donoghue, P.C., Pisani, D.J.N.e., and evolution (2018)
Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. **2**: 1556-1562.

Bhattacharya, D., and Medlin, L.J.J.P. (1995) Phylogeny of plastids A review based on comparisons of small
subunit ribosomal RNA coding regions. **31**: 487-496.

Biller, S.J., Berube, P.M., Lindell, D., and Chisholm, S.W. (2015) Prochlorococcus: the structure and function of
collective diversity. *Nature Reviews Microbiology* **13**: 13.

Boden, J.S., Konhauser, K.O., Robbins, L.J., and Sánchez-Baracaldo, P. (2021) Timing the evolution of
antioxidant enzymes in cyanobacteria. *Nature Communications* **12**: 1-12.

Braakman, R., Follows, M.J., and Chisholm, S.W. (2017) Metabolic evolution and the self-organization of
ecosystems. *Proceedings of the National Academy of Sciences* **114**: E3091-E3100.

Brocks, J.J. (2018) The transition from a cyanobacterial to algal world and the emergence of animals. *Emerging*
Topics in Life Sciences **2**: 181-190.

Brocks, J.J., Jarrett, A.J., Sirantoine, E., Hallmann, C., Hoshino, Y., and Liyanage, T. (2017) The rise of algae in

Cryogenian oceans and the emergence of animals. *Nature* **548**: 578.

Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., and Strauss, H. (2008) Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. *Science* **321**: 949-952.

Chriki-Adeeb, R., and Chriki, A.J.E.B. (2016) Estimating divergence times and substitution rates in Rhizobia. **12**: EBO. S39070.

Clarkson, M.O., Sweere, T.C., Chiu, C.F., Hennekam, R., Bowyer, F., and Wood, R.A.J.E.-S.R. (2023) Environmental controls on very high δ238U values in reducing sediments: Implications for Neoproterozoic seawater records. 104306.

Dayhoff, M.O. (1972) A model of evolutionary change in proteins. *Atlas of protein sequence and structure* **5**: 89-99.

dos Reis, M., Gunnell, G.F., Barba-Montoya, J., Wilkins, A., Yang, Z., and Yoder, A.D. (2018) Using phylogenomic data to explore the effects of relaxed clocks and calibration strategies on divergence time estimation: primates as a test case. *Systematic biology* **67**: 594-615.

dos Reis, M., Thawornwattana, Y., Angelis, K., Telford, Maximilian J., Donoghue, Philip C.J., and Yang, Z. (2015) Uncertainty in the Timing of Origin of Animals and the Limits of Precision in Molecular Timescales. *Current Biology* **25**: 2939-2950.

Duchêne, S., Lanfear, R., and Ho, S.Y.W. (2014) The impact of calibration and clock-model choice on molecular estimates of divergence times. *Molecular Phylogenetics and Evolution* **78**: 277-289.

E. Zuckerkandl, L.P. (1965) Evolutionary divergence and convergence in proteins. In *Evolving genes and proteins*: Elsevier, pp. 97-166.

Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincón, J., Zabala, L.L., Jiao, N. et al. (2013) Present and future global distributions of the marine Cyanobacteria Prochlorococcus and Synechococcus. *Proceedings of the National Academy of Sciences* **110**: 9824-9829.

Fournier, G., Moore, K., Rangel, L., Payette, J., Momper, L., and Bosak, T. (2021) The Archean origin of oxygenic photosynthesis and extant cyanobacterial lineages. *Proceedings of the Royal Society B* **288**: 20210675.

Goericke, R., Olson, R., and Shalapyonok, A.J.D.S.R.P.I.O.R.P. (2000) A novel niche for Prochlorococcus sp. in low-light suboxic environments in the Arabian Sea and the Eastern Tropical North Pacific. **47**: 1183-1205.

Gray, M.W.J.I.r.o.c. (1992) The endosymbiont hypothesis revisited. **141**: 233-357.

Grossman, A.R., Schaefer, M.R., Chiang, G.G., and Collier, J.L. (1993) The phycobilisome, a light-harvesting complex responsive to environmental conditions. *Microbiology and Molecular Biology Reviews* **57**: 725-749.

Heckman, D.S., Geiser, D.M., Eidell, B.R., Stauffer, R.L., Kardos, N.L., and Hedges, S.B.J.s. (2001) Molecular evidence for the early colonization of land by fungi and plants. **293**: 1129-1133.

Hedges, S.B., Tao, Q., Walker, M., and Kumar, S.J.P.o.t.N.A.o.S. (2018) Accurate timetrees require accurate calibrations. **115**: E9510-E9511.

Hess, W.R., Rocap, G., Ting, C.S., Larimer, F., Stilwagen, S., Lamerdin, J., and Chisholm, S.W. (2001) The photosynthetic apparatus of Prochlorococcus: insights through comparative genomics. *Photosynthesis Research* **70**: 53-71.

Ito, H., and Tanaka, A. (2011) Evolution of a divinyl chlorophyll-based photosystem in Prochlorococcus. *Proceedings of the National Academy of Sciences* **108**: 18014-18019.

Ito, H., Yokono, M., Tanaka, R., and Tanaka, A. (2008) Identification of a novel vinyl reductase gene essential for the biosynthesis of monovinyl chlorophyll in Synechocystis sp. PCC6803. *Journal of Biological Chemistry* **283**: 9002-9011.

Johnston, D.T., Wolfe-Simon, F., Pearson, A., and Knoll, A.H. (2009) Anoxygenic photosynthesis modulated

Proterozoic oxygen and sustained Earth's middle age. *Proceedings of the National Academy of Sciences* **106**:
16925-16929.

Jones, C., Nomosatryo, S., Crowe, S.A., Bjerrum, C.J., and Canfield, D.E.J.G. (2015) Iron oxides, divalent
cations, silica, and the early earth phosphorus crisis. **43**: 135-138.

Kang, J., Gill, B., Reid, R., Zhang, F., and Xiao, S. (2023) Nitrate limitation in early Neoproterozoic oceans
delayed the ecological rise of eukaryotes. *Science Advances* **9**: eade9647.

Keeling, P.J. (2013) The number, speed, and impact of plastid endosymbioses in eukaryotic evolution. *Annu Rev*
Plant Biol **64**: 583-607.

Knoll, A.H., and Nowak, M.A. (2017) The timetable of evolution. *Science Advances* **3**: e1603076.

Lau, K.V., Macdonald, F.A., Maher, K., and Payne, J.L. (2017) Uranium isotope evidence for temporary ocean
oxygenation in the aftermath of the Sturtian Snowball Earth. *Earth and Planetary Science Letters* **458**: 282-292.

Lavin, P., González, B., Santibáñez, J.F., Scanlan, D.J., and Ulloa, O.J.E.m.r. (2010) Novel lineages of
Prochlorococcus thrive within the oxygen minimum zone of the eastern tropical South Pacific. **2**: 728-738.

Luo, H., Huang, Y., Stepanauskas, R., and Tang, J. (2017) Excess of non-conservative amino acid changes in
marine bacterioplankton lineages with reduced genomes. *Nature microbiology* **2**: 17091.

Lyons, T.W., Reinhard, C.T., and Planavsky, N.J. (2014) The rise of oxygen in Earth's early ocean and
atmosphere. *Nature* **506**: 307.

Marin, B., M. Nowack, E.C., and Melkonian, M. (2005) A Plastid in the Making: Evidence for a Second
Primary Endosymbiosis. *Protist* **156**: 425-432.

Martiny, A.C., Kathuria, S., and Berube, P.M. (2009) Widespread metabolic potential for nitrite and nitrate
assimilation among Prochlorococcus ecotypes. *Proceedings of the National Academy of Sciences* **106**: 10787-
10792.

McGowen, M.R., Tsagkogeorga, G., Álvarez-Carretero, S., dos Reis, M., Struebig, M., Deaville, R. et al. (2019)
Phylogenomic Resolution of the Cetacean Tree of Life Using Target Sequence Capture. *Systematic Biology* **69**:
479-501.

Michiels, C.C., Darchambeau, F., Roland, F.A., Morana, C., Llirós, M., García-Armisen, T. et al. (2017) Iron-
dependent nitrogen cycling in a ferruginous lake and the nutrient status of Proterozoic oceans. *Nature*
Geoscience **10**: 217-221.

Morris, J.L., Puttick, M.N., Clark, J.W., Edwards, D., Kenrick, P., Pressel, S. et al. (2018) The timescale of early
land plant evolution. **115**: E2274-E2283.

Och, L.M., and Shields-Zhou, G.A. (2012) The Neoproterozoic oxygenation event: Environmental perturbations
and biogeochemical cycling. *Earth-Sci Rev* **110**: 26-57.

Ozaki, K., Thompson, K.J., Simister, R.L., Crowe, S.A., and Reinhard, C.T. (2019) Anoxygenic photosynthesis
and the delayed oxygenation of Earth's atmosphere. *Nature communications* **10**: 3026.

Palenik, B., Brahamsha, B., Larimer, F., Land, M., Hauser, L., Chain, P. et al. (2003) The genome of a motile
marine Synechococcus. **424**: 1037-1042.

Parfrey, L.W., Lahr, D.J., Knoll, A.H., and Katz, L.A.J.P.o.t.N.A.o.S. (2011) Estimating the timing of early
eukaryotic diversification with multigene molecular clocks. **108**: 13624-13629.

Partensky, F., and Garczarek, L. (2010) Prochlorococcus: advantages and limits of minimalism. *Annual Review*
of Marine Science **2**: 305-331.

Partensky, F., Hess, W.R., and Vaulot, D. (1999) Prochlorococcus, a marine photosynthetic prokaryote of global
significance. *Microbiol Mol Biol Rev* **63**: 106-127.

Ponce-Toledo, R.I., Deschamps, P., López-García, P., Zivanovic, Y., Benzerara, K., and Moreira, D. (2017) An

Early-Branching Freshwater Cyanobacterium at the Origin of Plastids. *Curr Biol* **27**: 386-391.

Ralf, G., and Repeta, D.J. (1992) The pigments of Prochlorococcus marinus: The presence of divinylchlorophyll

a and b in a marine procaryote. *Limnology and Oceanography* **37**: 425-433.

Rubio, L.M., Flores, E., and Herrero, A.J.J.o.b. (1998) The narA locus of Synechococcus sp. strain PCC 7942

consists of a cluster of molybdopterin biosynthesis genes. **180**: 1200-1206.

Sánchez-Baracaldo, P. (2015) Origin of marine planktonic cyanobacteria. *Scientific reports* **5**: 17418.

Sánchez-Baracaldo, P., Ridgwell, A., and Raven, J.A. (2014) A neoproterozoic transition in the marine nitrogen

cycle. *Current Biology* **24**: 652-657.

Sánchez-Baracaldo, P., Raven, J.A., Pisani, D., and Knoll, A.H. (2017) Early photosynthetic eukaryotes

inhabited low-salinity habitats. *Proceedings of the National Academy of Sciences*: 201620089.

Sánchez-Baracaldo, P., Bianchini, G., Wilson, J.D., and Knoll, A.H. (2022) Cyanobacteria and biogeochemical

cycles through Earth history. *Trends in Microbiology* **30**: 143-157.

Satoh, S., and Tanaka, A. (2006) Identification of chlorophyllide a oxygenase in the Prochlorococcus genome by

a comparative genomic approach. *Plant and cell physiology* **47**: 1622-1629.

Schirrmeister, B.E., Gugger, M., and Donoghue, P.C. (2015) Cyanobacteria and the Great Oxidation Event:

evidence from genes and fossils. *Palaeontology* **58**: 769-785.

Schirrmeister, B.E., Sanchez-Baracaldo, P., and Wacey, D. (2016) Cyanobacterial evolution during the

Precambrian. *International Journal of Astrobiology* **15**: 187-204.

Shih, P.M., Hemp, J., Ward, L.M., Matzke, N.J., and Fischer, W.W. (2016) Crown group Oxyphotobacteria

postdate the rise of oxygen. *Geobiology* **15**: 19-29.

Stolper, D.A., and Keller, C.B.J.N. (2018) A record of deep-ocean dissolved O2 from the oxidation state of iron

in submarine basalts. **553**: 323-327.

Strassert, J.F.H., Irisarri, I., Williams, T.A., and Burki, F. (2021) A molecular timescale for eukaryote evolution

with implications for the origin of red algal-derived plastids. *Nat Commun* **12**: 1879.

Tahata, M., Sawaki, Y., Yoshiya, K., Nishizawa, M., Komiya, T., Hirata, T. et al. (2015) The marine

environments encompassing the Neoproterozoic glaciations: evidence from C, Sr and Fe isotope ratios in the

Hecla Hoek Supergroup in Svalbard. *Precambrian Research* **263**: 19-42.

Ting, C.S., Rocap, G., King, J., and Chisholm, S.W. (2002) Cyanobacterial photosynthesis in the oceans: the

origins and significance of divergent light-harvesting strategies. *Trends Microbiol* **10**: 134-142.

Tostevin, R., and Mills, B.J.J.I.F. (2020) Reconciling proxy records and models of Earth's oxygenation during

the Neoproterozoic and Palaeozoic. **10**: 20190137.

Trower, E.J., Gutoski, J.R., Wala, V.T., Mackey, T.J., and Simpson, C. (2023) Tonian Low-Latitude Marine

Ecosystems Were Cold Before Snowball Earth. *Geophysical Research Letters* **50**: e2022GL101903.

Ulloa, O., Henríquez-Castillo, C., Ramírez-Flandes, S., Plominsky, A.M., Murillo, A.A., Morgan-Lang, C. et al.

(2021) The cyanobacterium Prochlorococcus has divergent light-harvesting antennae and may have evolved in a

low-oxygen ocean. *Proceedings of the National Academy of Sciences* **118**.

Wang, S., and Luo, H. (2021) Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nature*

Communications **12**: 3324.

Warnock, R.C., Yang, Z., and Donoghue, P.C.J.B.l. (2012) Exploring uncertainty in the calibration of the

molecular clock. **8**: 156-159.

Zhang, F., Stockey, R.G., Xiao, S., Shen, S.-z., Dahl, T.W., Wei, G.-Y. et al. (2022) Uranium isotope evidence

for extensive shallow water anoxia in the early Tonian oceans. *Earth and Planetary Science Letters* **583**: 117437.

Zhang, H., Hellweger, F.L., and Luo, H.J.b. (2023) Genome reduction occurred in early Prochlorococcus with

an unusually low effective population size. 2023.2006. 2025.546417.

Zhang, H., Sun, Y., Zeng, Q., Crowe, S.A., and Luo, H. (2021) Snowball Earth, population bottleneck and

Prochlorococcus evolution. *Proceedings of the Royal Society B: Biological Sciences* **288**: 20211956.

Zhang, J. (2000) Rates of conservative and radical nonsynonymous nucleotide substitutions in mammalian

nuclear genes. *J Mol Evol* **50**: 56-68.

Figure Legends

Fig. 1   The evolutionary timeline of *Prochlorococcus* estimated with plastid-based

strategy. (a) The diagram shows the phylogenetic relations among Cyanobacteria and

eukaryotic species. The green solid line, black solid line and green dashed line in the

cladogram leading to the tip of oxygenic Cyanobacteria, anoxygenic Vampirovibrionia and

Sericytochromatia and eukaryotic species, respectively. The calibrated nodes in molecular

clock analysis are marked with orange circle (see calibration justifications in Supplementary

Methods). (b) (left) The *Prochlorococcus* evolutionary timeline estimated with the focal

molecular dating strategy using sequential Bayesian dating approach based on the 5-partition

eukaryotic genome-scale data (in the first step dating analysis) and fully partitioned 12-gene

dataset "T30" (in the second step dating analysis) under independent rate clock model. The

posterior age distribution is provided next to the ancestral node. The atmospheric oxygen

level at the geological time scale is adapted from Lyons et al., 2014, which is represented by

the percent of present atmospheric oxygen level (PAL). The vertical bars with green, orange

and blue colors represent the time of Tonian, the time of NOE, and the time of Sturtian (left)

and Marinoan (right) glaciation, respectively. (right) Phyletic pattern of key gene families.

Solid square, solid circle and open circle next to each analyzed genome represent multi-copy

gene family, single-copy gene family and absence of the gene family, respectively, in the

extant genomes. Gene families marked with asterisk (*) were consistently estimated to be

gained or lost by using both AnGST and GeneRax. (c) Diagram illustrating the

biogeochemical environments when *Prochlorococcus* arose from its *Synechococcus* ancestor.

Fig. 2   **Diagram illustrating the schemes and the results of d_R/d_C comparison.** The

genome-wide means of d_R/d_C values at the ancestral branches ('Target') leading to (a) Proch-

AMZI/II/III-LCA and (b) SBE-LCA compared with that at the sister lineages ('Control').

The d_R/d_C values were classified based on the physicochemical classification of the amino

acids by charge or by volume and polarity, and were either GC-corrected by codon frequency

(blue), GC-corrected by amino acid (AA) frequency (red) or uncorrected (gray). Error bars of

d_R/d_C represent the standard error of the mean.

Fig. 1

Fig. 2