Myeloid-mesenchymal crosstalk drives Arg1-dependent profibrotic metabolism via ornithine in lung fibrosis

Authors

Preeti Yadav[1,2], Javier Gómez Ortega[1,2], Prerna Dabral[3,4], Whitney Tamaki[5,6], Charles Chien[7,8], Kai-chun Chang[9,10], Nivedita Biswas[1,2], Sixuan Pan[10], Julia Nilsson[4], Xiaoyang Yin[1,2,11], Aritra Bhattacharyya[1,2,12], Kaveh Boostanpour[1,2], Tanay Jujaray[1,2], Jasper Wang[1,2], Tatsuya Tsukui[1,13], Chris Molina[1,13], Vincent Auyeung[1,13], Dean Sheppard[1,13], Baosheng Li[11], Mazharul Maishan[1], Hiroki Taenaka[1], Michael A. Matthay[1], Rieko Muramatsu[14], Lenka Maliskova[15], Arnab Ghosh[15], Walter L. Eckalbar[15], Ari B. Molofsky[4], Stanley J. Tamaki[15], Trever Bivona[3,4,16], Adam R. Abate[10], Allon Wagner[7,8,17], Satish K. Pillai[3,4], Paul J. Wolters[1], Kevin M. Tharp[18], Mallar Bhattacharya[1,2§]

Affiliations

[1]Division of Pulmonary, Critical Care, Allergy, and Sleep, Department of Medicine, University of California, San Francisco, San Francisco, CA, USA

[2]Sandler Asthma Basic Research Center, University of California, San Francisco, San Francisco, CA, USA

[3]Vitalant Research Institute, San Francisco, California.

[4]Department of Laboratory Medicine, University of California, San Francisco, San Francisco, California, USA.

[5]Department of Medicine, University of California, San Francisco, San Francisco, CA 94158, USA.

[6]Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, San Francisco, CA 94158, USA.

[7]Department of Electrical Engineering and Computer Sciences, University of California, Berkeley, CA 94720, USA

[8]Center for Computational Biology, University of California, Berkeley, CA 94720, USA

[9]Twist Bioscience, 681 Gateway Blvd, South San Francisco, CA, USA

[10]Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, San Francisco, CA, USA

[11]Department of Radiation Oncology, Shandong Cancer Hospital and Institute, Shandong First Medical University and Shandong Academy of Medical Sciences, Jinan, China

[12]Biological Science Division, Indian Statistical Institute, Kolkata, India.

[13]Cardiovascular Research Institute; University of California, San Francisco, San Francisco, CA, USA

[14]Department of Molecular Pharmacology, National Institute of Neuroscience, National Center of Neurology and Psychiatry, Kodaira, Japan

[15]UCSF CoLabs, University of California, San Francisco, San Francisco, CA 94143, USA

[16]Chan-Zuckerberg Biohub, San Francisco, CA 94158, USA.

[17]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA

[18]Cancer Metabolism and Microenvironment Program, NCI-Designated Cancer Center, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA, USA

§ Correspondence: Mallar Bhattacharya
         Email: mallar.bhattacharya@ucsf.edu
         Phone: 415-514-1018
         Address: 513 Parnassus Avenue, HSE-201P, San Francisco, CA 94143, USA

Declaration of Interests

The authors have declared that no conflict of interest exists.

<u>Abstract</u>

Idiopathic pulmonary fibrosis (IPF) is a disease of progressive lung remodeling and collagen deposition

that leads to respiratory failure. Myeloid cells are abundant in IPF lung and in murine lung fibrosis, but

their functional effects are incompletely understood. Using mouse and human lung models, we show

that ornithine produced by myeloid cells expressing Arg1 serves as a substrate for proline and collagen

synthesis by lung fibroblasts. The predominant Arg1-expressing myeloid cells in mouse lung were

macrophages, but in IPF lung, high-dimensional imaging revealed ARG1 to be expressed mainly in

neutrophils. Arg1 inhibition suppressed both ornithine levels and collagen expression in cultured,

precision-cut IPF lung slices and in murine lung fibrosis. These results were confirmed in macrophage-

specific Arg1 KO mice. Furthermore, we find that this pathway is regulated by cell-to-cell crosstalk,

starting with purinergic signaling: Fibroblast eATP receptor P2rx4 was necessary for fibroblast IL-6

expression, which in turn was necessary for Arg1 expression by myeloid cells. Taken together, our

findings define an immune-mesenchymal circuit that governs profibrotic metabolism in lung fibrosis.

<u>Keywords</u>

Myeloid Arginase 1; Fibroblasts; Fibrosis; IPF; Ornithine

Introduction

Pathologic fibrosis is a destructive response to tissue injury that results from the deposition of excess fibrillar collagen by activated fibroblasts. In idiopathic pulmonary fibrosis (IPF), it is believed that dysfunction of the epithelium leads to the recruitment and activation of multiple cell types, including most importantly fibroblasts and myeloid cells (1). These cells have been found to be in close proximity in the fibrotic niche (2, 3), but it remains unclear how they interact to induce the fibrotic phenotype. One approach is to define and test the functional role of diffusible factors that have been detected in fibrosis and that might subserve crosstalk between the various cell types leading to collagen production by fibroblasts.

We recently considered the role of extracellular ATP (eATP), a damage-associated molecular pattern (DAMP) molecule that is extensively present in the IPF lung, as revealed in the bronchoalveolar lavage from patients for example (4). eATP's role in fibrosis remains incompletely understood. Although eATP activates receptor-mediated transcriptional responses by ligating multiple purinergic receptors, in lung fibroblasts we reported P2rx4 as the critical purinergic receptor: Fibroblast-specific deletion of the eATP receptor P2rx4 decreased lung fibrosis in mice (5). However, the paracrine signaling effects of this pathway are not known.

Here we explore the effect of P2rx4 signaling on cell-to-cell crosstalk in fibrosis. Recent studies have revealed a heterogeneity of fibroblast states following injury, and inflammatory fibroblasts have been detected as a distinct subset of mesenchymal cells that express secreted inflammatory factors such as chemokines and cytokines (6-8). How these signals from inflammatory fibroblasts promote fibrosis is unknown. We hypothesized that eATP-P2rx4 signaling may stimulate expression of these diffusible signals from fibroblasts after injury, initiating profibrotic crosstalk with other cells in the fibrotic niche.

Studies with murine macrophage-fibroblast cocultures revealed that macrophage Arg1 expression was dependent on P2rx4 signaling in fibroblasts. Arg1+ macrophages were increased after bleomycin-induced lung fibrosis in mice; in IPF lung, ARG1+ cells were also increased, in the neutrophil compartment. In both murine and human lung systems, Arg1 inhibition decreased lung fibrosis.

94   Furthermore, Arg1 was induced by fibroblast expression of the cytokine IL-6 in a P2rx4-dependent

95   manner. Mechanistically, Arg1 drove fibroblast collagen synthesis via the production of the noncoding

96   amino acid ornithine, which served as a precursor for the collagen building block proline. These findings

97   indicate a critical profibrotic role for a metabolic pathway arising from cellular crosstalk in the

98   development of lung fibrosis.

Results

Arg1+ cells localize to the fibrotic niche. We recently reported that mice with deletion of the eATP

receptor P2rx4 in fibroblasts were protected from bleomycin injury-induced lung fibrosis (5). To screen

for macrophage genes regulated by fibroblast eATP-P2rx4 signaling, we used an *in vitro* system: Either

WT or P2rx4 KO fibroblasts were cocultured with WT lung macrophages (**Supplemental Figure 1A**),

followed by single cell RNAseq. Both cell types were isolated from murine lungs 7 days after injury to

approximate cellular states during the early fibrotic period, defined by increased fibroblast expression of

Col1a1 and Col3a1 (**Supplemental Figure 1B**). Fibroblasts were used directly, whereas macrophages

were treated for 48 hours in culture with Csf1 to maintain macrophage identity prior to coculture. After 5

days of coculture, cells were submitted for scRNAseq.

The most downregulated macrophage gene when fibroblast P2rx4 was deleted was Arg1, a

marker of alternatively activated macrophages (**Figure 1A**). Analysis of lung macrophages in a

published scRNAseq time course following bleomycin lung injury (9) showed maximal *Arg1* in the "C2",

transitional macrophage compartment that we previously demonstrated was localized to the fibrotic

niche after injury(2) (**Figure 1B**). C2 cells expressed higher levels of Arg1 and also the "Fab5" marker

genes that were recently found to be associated with a core pro-fibrotic macrophage program (10)

(**Supplemental Figure 1C-E**), and they were transcriptomically similar to recruited and interstitial

macrophages (11) (**Supplemental Figure 1F**). Interestingly, the macrophages from our cocultures also

expressed higher levels of interstitial macrophage than alveolar macrophage genes (**Supplemental

Figure 1G**). To determine whether Arg1+ cells localized to areas of fibrosis, we injured mice expressing

reporter alleles for Arg1 and for the fibroblast marker Col1a1. Confocal imaging revealed that Arg1+

cells were in proximity to clusters of activated fibroblasts in the fibrotic niche (**Figure 1C**).

To further profile Arg1 expression in myeloid cell subsets in mice, we performed flow cytometry

with Arg1 reporter mice. This analysis confirmed that Arg1 was expressed 10 days after bleomycin

injury in macrophages expressing CD11b and CD64, markers consistent with monocyte-derived

macrophages (**Supplemental Figure 2A-C**). Canonical neutrophils expressing Ly6G did not express

Arg1, which we confirmed by immunofluorescence for the neutrophil marker S100A8 (**Supplemental

Figure 2D). However, we did note emergence of a small myeloid population of Arg1+ cells expressing

Ly6G that also expressed CD64 and CD11c, though this population was about tenfold smaller in

number than moMacs (**Supplemental Figure 2E**). Taken together, these data confirm expression of

Arg1 in myeloid cells in the mouse lung fibrotic niche.

To determine the relevance of ARG1 to IPF, we first interrogated a published dataset of BAL

cell gene expression by microarray (12). This analysis confirmed higher ARG1 expression in IPF

compared to healthy controls (**Figure 1D**). We then found increased ARG1+ cells by

immunofluorescence of IPF lung explants acquired at the time of clinical transplantation compared with

deceased donor lungs not known to have lung disease (**Figure 1E; Supplemental Table 1**). In human

lung, ARG1 has been found to be expressed in neutrophils, based on data from lung cancer specimens

(13-15). However, deep phenotyping in IPF lung tissue of ARG1-expressing cell types has not been

undertaken. Therefore, to comprehensively profile the cell types expressing ARG1 in IPF, we

performed multiplexed ion beam imaging (MIBI) (16-19), an antibody-based imaging technique in which

isotopically pure elemental metal reporters are conjugated to antibodies and detected spatially in

combination with mass spectrometry. We applied MIBI to sections from 5 IPF explanted lungs using

anti-ARG1 antibody and 33 other antibodies, selecting markers primarily for the purpose of defining

major immune cell types. These data showed that ARG1+ cells predominantly expressed the

neutrophil-specific markers CD66B and MPO but not the macrophage markers CD68, CD206, and

CD163 (**Figure 2A-C; Supplemental Table 2**). Importantly, neutrophils have been associated with

increased mortality in IPF (20-23). However, their function in fibrogenesis has not been well studied.

Arg1 regulates availability of ornithine, a pro-fibrotic substrate. To understand Arg1's function, we

considered first that ornithine generated from arginine by Arg1 can serve as a substrate for the

synthesis of proline (24), a major constituent of collagen. Remarkably, we found that ornithine was

markedly increased both in mouse lung 14 days after bleomycin injury compared to steady state and in

IPF lung lysates compared to healthy controls (**Figure 3A**). To test the effects of Arg1 and ornithine in

fibrosis, we first used the small-molecule Arg1 inhibitor CB-1158 (25). We confirmed the on-target effect

of CB-1158 by measuring lung ornithine after treatment: In both bleomycin-injured mice and precision-

cut lung slices (PCLS) from explanted IPF lungs, the inhibitor decreased ornithine levels (**Figure 3B**).

Next, we treated bleomycin-injured mice with CB-1158 during the fibrotic period. Importantly, we found

that lung fibrosis was significantly decreased and body weight recovery improved with Arg1 inhibition

(**Figure 3C; Supplemental Figure 2F**). We then prepared PCLS from explanted IPF lungs and treated

them with CB-1158 for 24 hours, followed by immunoblot for COL1A1 in the RIPA buffer-soluble

fraction representing the more soluble, newly synthesized collagen (26). COL1A1 expression measured

by immunoblot of lung slice lysates was decreased by CB-1158 treatment compared to untreated

controls, indicating suppression of collagen synthesis with ARG1 inhibition (**Figure 3D**). Retention of

cells expressing ARG1 and neutrophil markers MPO and CD66B was confirmed in PCLS by

immunofluorescence (**Supplemental Figure 3A**). Finally, taking a genetic approach with macrophage-

specific Arg1 KO mice in which Cre-mediated deletion efficiency was approximately 55%

(**Supplementary Figure 3B**), we found that lung fibrosis was significantly decreased after injury in KO

mice compared to controls (**Figure 3E**). Taken together, these data indicate that Arg1 determines the

pathologic accumulation of collagen in lung fibrosis.

We then tested the effect of ornithine on collagen expression. First, we found that exogenous

ornithine increased fibroblast collagen expression by immunofluorescence in monocultured fibroblasts

from both mouse and human lung (**Figure 3F**). To test the profibrotic potential of ornithine *in vivo*, we

treated mice with ornithine by oral gavage, either at steady state or during the fibrotic phase after

bleomycin injury. Ornithine treatment increased lung collagen production in the setting of injury but not

at steady state, consistent with increased cellular demand for collagen synthetic precursors including

proline during fibrogenesis (**Figure 3G**). We then found that macrophage-fibroblast coculture increased

collagen expression compared to fibroblast monoculture, an effect that could be blocked with CB-1158

or fibroblast P2rx4 KO and enhanced with ornithine (**Figure 3H-I**). Furthermore, ornithine could rescue

the CB-1158-dependent decreased collagen expression in cocultures (**Supplemental Figure 3C**).

Finally, fibroblast coculture with Arg1-specific KO macrophages was associated with decreased

collagen expression compared with WT macrophages, an effect that was not significantly augmented

with CB-1158 (**Figure 3J**). Taken together, these results indicate a direct profibrotic effect of ornithine

in vitro and *in vivo*.

Ornithine is converted by ornithine aminotransferase (OAT) to glutamate-5-semialdehyde, which

spontaneously cyclizes to pyrroline-5-carboxylate and is then reduced by P5C reductases to proline.

Notably, OAT inhibition decreased coculture-dependent augmentation of collagen expression

(**Supplemental Figure 3C**). Interestingly, coculture also increased α-SMA expression, but this effect

was not augmented by ornithine or suppressed by OAT inhibition (**Supplemental Figure 3D)**. These

data indicate a dependence of collagen synthesis on Arg1 function and ornithine metabolism. Finally,

we directly tested the hypothesis that Arg1-mediated ornithine production was necessary for increasing

fibroblast proline content, using the approach of macrophage-fibroblast coculture followed by isolation

of fibroblasts for LC-MS to detect proline. Consistent with our hypothesis, Arg1 inhibition with CB-1158

significantly decreased fibroblast proline content in cocultures (**Figure 3K**). Taken together, these

results suggest that macrophage Arg1 drives fibrosis by producing ornithine, which serves as a

substrate for fibroblast proline synthesis, augmenting collagen expression.

Arg1 is regulated by IL-6. To determine how macrophage Arg1 expression is regulated by signaling

within the fibrotic niche, we considered that our previously described coculture data (**Figure 1A**)

indicated that P2rx4 signaling in fibroblasts was necessary for Arg1 expression in cocultured

macrophages. Therefore, we performed analysis of ligand-receptor interactions via CellChat (27) and

Ingenuity Pathways Analysis. This analysis revealed suppression of IL-6 signaling in the P2rx4 KO

condition (**Figure 4A**). We also found that P2rx4 KO fibroblasts expressed less IL-6 mRNA compared

to WT (**Figure 4B**). IL-6 is an important profibrotic factor and is a therapeutic target in clinical lung

fibrosis (28-31). We first confirmed an increase of Arg1 in cultured mouse lung macrophages treated

with IL-6 (**Figure 4C**). *In vivo*, we noted that IL-6 KO mice had decreased macrophage numbers in the

lung, consistent with a chemotactic effect of IL-6 (**Supplemental Figure 3E**). Nonetheless, even

accounting for this lower cell count, Arg1 lavage levels were disproportionately decreased in IL-6 KO

mice, as indicated by bronchoalveolar lavage Arg1 concentrations normalized to CD11b+CD64+

macrophage count and by qPCR measurement of *Arg1* relative to *Gapdh* in flow-sorted CD11b+CD64+

macrophages (**Figure 4D**). Of note, IL-6 KO mice had decreased lung fibrosis after bleomycin lung

injury (**Supplemental Figure 3F**). We then analyzed scRNAseq time course data for multiple lung cell

types and found that IL-6 was indeed highly expressed in fibroblasts relative to other cell types (**Figure**

4E). Furthermore, IL-6 was notably expressed in the emergent, disease-associated inflammatory and

fibrotic fibroblasts observed in recent time course data (8) (**Figure 4F**).

To test whether IL-6 could induce ARG1 in human lung, we first tested PCLS from explanted

IPF lungs. Treatment with the IL-6R blocking antibody tocilizumab for 24 hours decreased ARG1

expression in IPF PCLS lysates (**Figure 5A**). To confirm that paracrine interaction between ARG1+ and

IL-6+ cells was spatially plausible, we analyzed spatial transcriptomic data of IPF samples. For this

purpose, we defined a proximity statistic that allowed us to quantitatively compare the probabilities of

detecting an IL6+ fibroblast versus an IL6- fibroblast, within a given distance from an ARG1+ cell

(**Methods**). Interestingly, at a distance from ARG1+ cells consistent with potential paracrine signaling

(25 microns), fibroblasts were significantly more likely to express IL-6 than not to express IL-6 (**Figure**

5B). These data are consistent with paracrine IL-6 signaling from fibroblasts to macrophages resulting

in ARG1 induction in IPF.

We then interrogated published datasets for neutrophils from lung tissue to test whether ARG1+

neutrophils were more likely to express IL6R and therefore to support IL6 signaling. While genome-

wide transcriptomic profiling of neutrophils from human fibrotic lungs has not yet been reported in the

literature, as an alternative, we analyzed aggregate data from available non-small cell lung cancer

(NSCLC) tumor microenvironment studies from 309 patients across 19 datasets (32). This analysis

revealed a significant correlation between ARG1 and IL6R expression in neutrophils, consistent with

the hypothesis that ARG1 expression in neutrophils is driven by IL-6 receptor ligation and signaling

(**Figure 5C).**

Finally, we cocultured human peripheral blood-derived CD15+CD16+CD66B+CD14- neutrophils

(**Supplemental Figure 3G**) with human lung fibroblasts. After 24 hours of coculture, we removed the

neutrophils, which are nonadherent cells, and measured collagen expression in the fibroblasts by

immunofluorescence. Notably, neutrophil coculture increased fibroblast collagen expression compared

to fibroblast monoculture, and this increase could be blocked by IL6-R blockade with tocilizumab

(**Figure 5D**). Notably, ARG1 and ornithine levels in the conditioned media were suppressed by

tocilizumab (**Figure 5E-F**). Taken together, these analyses support the relevance of fibroblast IL-6-

dependent ARG1 expression in the human lung fibrotic niche.

eATP signaling induces fibroblast IL-6. To directly test whether the eATP receptor P2rx4 regulates

fibroblast IL-6 expression, we considered that myeloid cells including both macrophages and

neutrophils can be a source of eATP (33). Thus, we performed scRNAseq of fibroblasts cocultured with

macrophages or cultured alone. Interestingly, coculture increased fibroblast expression of IL-6 (**Figure**

6A), and GO analysis confirmed that the GO pathway "Cellular Response to ATP" was enriched in

fibroblasts in coculture compared to monoculture (**Figure 6B**). Importantly, IL-6 was higher in the

conditioned media of WT macrophages cocultured with WT versus P2rx4 knockout (KO) fibroblasts

(**Figure 6C**). Furthermore, direct treatment of cultured murine lung fibroblasts with ATPγS, a

nonhydrolyzable form of ATP, increased IL-6 in the conditioned media of WT but not P2rx4 KO lung

fibroblasts, and this effect was blocked by inhibition of p38 Map Kinase, which has been shown to

mediate signaling downstream of P2rx4 (34, 35) (**Figure 6D**). To evaluate these results *in vivo*, we

tested mice with fibroblast-specific P2rx4 knockout (KO) after bleomycin injury. We found reduced IL-6

levels in the bronchoalveolar lavage fluid of KO compared to WT mice (**Figure 6E**). Furthermore,

siRNA KD of P2RX4 in human lung fibroblasts decreased IL-6 expression in response to ATPγS

(**Figure 6F**). We also found that P2rx4 expression increased post-bleomycin injury (**Supplemental**

Figure 3H), consistent with increased functional effects of this pathway during fibrogenesis. Finally,

coculture of human peripheral blood-derived neutrophils with human lung fibroblasts increased IL-6 in

the conditioned media compared to fibroblasts or neutrophils alone. This effect could be inhibited by

treatment with a small molecule inhibitor of P2rx4 (BAY-1797(36); **Figure 6G**). Taken together, our

findings indicate that, in both murine and human systems, eATP-P2rx4 signaling regulates fibroblast IL-

6 expression, which in turn induces myeloid Arg1 resulting in ornithine loading of fibroblasts for proline

synthesis and pathologic collagen expression.

<u>Discussion</u>

For the first time to our knowledge, we demonstrate that Arg1 and its enzymatic product ornithine have

a profibrotic effect by driving fibroblast proline synthesis, a key substrate for collagen expression.

Imaging studies indicated that ARG1 is expressed more highly in IPF than in healthy human lung

explants. Studies using both genetic deletion and chemical inhibition demonstrated the profibrotic effect

of Arg1 in murine lung injury and in IPF precision-cut lung slices. Remarkably, we found that the

reaction product of Arg1, ornithine, was increased in fibrotic lung tissue from mice and IPF patients.

These data build on the recent findings that ornithine was the only amino acid to be elevated in plasma

of 100 IPF patients compared to 300 healthy controls (37) and that the expression of ornithine

aminotransferase, which converts ornithine to the proline precursor P5C, was correlated with areas of

fibrosis in IPF lung (38). Furthermore, we found that ornithine had a profibrotic effect: Ornithine directly

increased collagen expression *in vivo* and in cultured murine and human lung fibroblasts. Metabolic

labeling studies revealed that ornithine was a substrate for proline synthesis.

Our findings also address how this profibrotic amino acid metabolism was initiated via paracrine

crosstalk. In both murine and human fibrotic lung and coculture systems, myeloid Arg1 expression was

dependent on IL-6 expression in fibroblasts, which in turn was regulated by eATP-P2rx4 signaling. The

recent literature has revealed a subset of fibroblasts expressing inflammatory genes in fibrosis,

including cytokines (7, 8). A pathologic function of fibroblast-derived cytokine expression is elucidated

by our discovery that fibroblast IL-6 is necessary for Arg1 expression in myeloid cells, with the

profibrotic consequence of ornithine loading of fibroblasts leading to increased collagen production.

The role of type 2 inflammation in fibrosis deserves mention with respect to Arg1. In

schistosomiasis-induced fibrosis models, where Arg1 deletion led to an increase in type 2 inflammation

and consequently increased fibrosis because of an accumulation of arginine, which supports CD4 T cell

proliferation (39). However, in IPF and in murine bleomycin-induced pulmonary fibrosis, type 2

inflammation does not appear to be an important driver of fibrosis: Inhibition of type 2 inflammation with

dual IL-4 and IL-13 blockade failed to show efficacy in a clinical trial of IPF (40). Furthermore, in the

bleomycin-induced sterile injury model that we have used, the onset of type 2 inflammation is later than

the period when significant collagen deposition has already occurred and is therefore less relevant to

fibrogenesis (41). Therefore, we conclude that type 2 inflammation may be less important in the model.

The myeloid cell type expressing Arg1 varied by species. In mice, we found that Arg1 was

predominantly expressed in CD11b+CD64+ macrophages and not in canonical Ly6G+ neutrophils.

However, there was Arg1 expression in a relatively smaller population of Ly6G+CD64+CD11c+ cells,

which may be an intermediate inflammatory myeloid lineage similar to a profile that others have

recently described in the setting of murine influenza infection (42). Future studies should more deeply

profile these latter cells and ARG1+ cells from IPF, which expressed neutrophil markers, to determine

whether they are phenotypically similar. Nonetheless, in both species we confirmed a P2RX4 – IL-6 –

ARG1, myeloid-mesenchymal circuit that is functionally important in lung fibrosis. Notably, blood and

bronchoalveolar lavage neutrophilia has been associated with fibrosis progression and worse prognosis

in IPF (20-23). Our work highlights ARG1 and ornithine metabolism as a neutrophil-dependent pathway

that should be developed as a potential therapeutic target for IPF.

There are limitations that should be considered when interpreting our results. First, our

metabolomic studies indicate conversion of ornithine to proline, a key substrate for collagen synthesis;

however, we note that ornithine conversion to polyamines with profibrotic potential remains a further

possibility not yet addressed by our results. Second, Arg2 is another arginase that could also contribute

to ornithine loading of fibroblasts, and future studies should focus on how both arginases are regulated

across the temporal phases of fibrosis. Third, our findings are agnostic with respect to the site of

ornithine generation; we detected Arg1 both in both lysates and conditioned media of IL-6-treated

macrophages, consistent with the possibility of both intracellular ornithine production and export as well

as extracellular production, as per published reports (43, 44). Finally, regarding sources of eATP, we

concede that it is likely that multiple sources of eATP exist in the fibrotic niche, such as dying or

dysfunctional cells, in addition to myeloid cells themselves. Our results nonetheless reveal that the

DAMP signal eATP triggers expression of fibroblast IL-6, which induces profibrotic Arg1 in neighboring

myeloid cells.

In conclusion, we show the dependence of lung fibrosis on ornithine deriving from myeloid Arg1

in mouse models and in functional studies of IPF lung. The importance of understanding the

contribution of amino acid metabolism to fibrosis has been recognized in recent years (45), and our

findings highlight the role of its immune regulation in the synthesis of pathologic collagen. These

studies increase enthusiasm for targeting ARG1 as a therapeutic approach in IPF.

Methods

Sex as a biological variable. For preclinical in vivo studies, we used both female and male mice, and experimental groups were balanced with respect to sex. Human lung samples were acquired from both males and females.

Human lung tissues. IPF lung samples were obtained as explants at the time of lung transplantation. Deceased donor-control lungs not known to have lung disease were made available by Donor Network West. Demographic data with respect to ethnicity and race (**Supplemental Table 1**) were derived from the electronic medical record and classified as per NIH notice NOT-OD-15-089.

Mice. Pdgfrb-Cre(46), Lysm-Cre (47), P2rx4 floxed (48), Arg1 floxed (49), Col1a1-EGFP (50), Arg1-RFP-CreERT2 (51), and Arg1-YFP (52) mice were available to the investigative team, and IL-6 KO (53), WT, and R26-LSLS-TdTomato mice were obtained from JAX. Col1a1-GFP mice were obtained from David Brenner (50). All mice were maintained on a C57BL/6 background and were maintained in specific pathogen-free animal barrier facility at the University of California, San Francisco. All experiments were performed on 6-8 weeks old, sex-matched mice.

Mouse lung injury. For lung injury, mice anesthetized with isoflurane were instilled intratracheally with bleomycin (Fresenius, 3U/kg). In the case of Arg1 inhibition, bleomycin-injured mice were treated daily from day 9 to day 15 post-injury with 100 mg/kg CB-1158 (Numidargistat dihydrochloride, HY-101979A, MedChemExpress) dissolved in water, by oral gavage. In the case of ornithine treatment, mice were treated twice daily by gavage with ornithine 2 g/kg dissolved in 100 mL of water.

Murine macrophage-fibroblast coculture. We prepared cocultures of macrophages and fibroblasts that were isolated from the lungs of mice 7 days after bleomycin lung injury. To make single cell suspensions, lungs were harvested from euthanized mice and minced with scissors, and the tissue was resuspended in RPMI with 0.2% Collagenase (10103586001, Roche), 2000 U/ml DNase I (4716728001, Roche), and 0.1 mg/ml Dispase II (4942078001, Sigma). The suspension was incubated for 1 hour at 37°C and then passed through a 70 μm filter (130-110-916, MACS SmartStrainers) to obtain single cells. The cells were washed twice with 1X PBS pH 7.4 (10010023, Gibco). For macrophage isolation, a total of 10^7 cells were resuspended in 80μl PBS buffer supplemented with

0.5% BSA (0332, VWR Life Science) and 2mM EDTA (E57040, Research Products International). 20μl

CD11b microbeads were used per 10^7 cells. Macrophages were obtained by positive selection by

binding to CD11b microbeads and passing through MACS columns (130-049-601, Miltenyi Biotech).

Isolated CD11b$^+$ macrophages were cultured in DMEM supplemented with 10% FBS, 1% penicillin-

streptomycin (GIBCO), and 20 ng/ml M-CSF (315-02, Peprotech) for 2 days. Primary mouse lung

fibroblasts were freshly isolated after magnetic bead-based negative selection of epithelial cells

(118204, Biotin anti-mouse CD326 Epcam, Clone: G8.8, BioLegend), endothelial cells (102404, Biotin

anti-mouse CD31, clone: 390), immune cells (103104, Biotin anti-mouse CD45, clone: 30-F11,

BioLegend), pericytes and smooth muscle cells (134716, Biotin anti-mouse CD146, clone: ME-9F1,

BioLegend), and red blood cells (116204, Biotin anti-mouse Ter119, clone: TER-119, BioLegend) with

biotinylated antibodies and Dynabeads (MyOne Streptavidin T1, 65601, Thermo Fisher Scientific) as

previously described (7). Fibroblasts were cultured for 5 days prior to trypsinization and addition to

macrophages for coculture at a 1:1 ratio in DMEM complete media for 5 further days.

Human lung fibroblast isolation. Human lung tissue from deceased donors not used for transplant

was resected and minced in Hanks' Balanced Salt Solution (HBSS) buffer supplemented with 0.2%

Collagenase (10103586001, Roche), 2000 U/ml DNase I (4716728001, Roche), 0.1 mg/ml Dispase II

(4942078001, Sigma), and 1% Penicillin-streptomycin for 1.5 hours at 37°C and 5% CO2. 1X

Amphotericin B (15290026, Gibco) was added to the dissociation solution for the last 30 min. Digested

lung tissue was lysed further with a MACS tissue dissociator (gentleMACS Dissociator, Miltenyi Biotec)

using gentleMACS C tubes (130-093-237, Miltenyi Biotec) at mLUNG-01 setting. The suspension was

then passed through a 70 μm filter to obtain single cells. Cells were resuspended in PBS with 0.5%

BSA and 2mM EDTA. For the negative selection of human fibroblasts, the following antibodies were

used at a 1:200 ratio: epithelial cells (324216, Biotin anti-human Epcam, clone:9C4, BioLegend),

endothelial cells (13-0319-82, Biotin anti-human CD31, clone: WM-59, Invitrogen), immune cells

(368534, Biotin anti-human CD45, clone: 2D1, BioLegend), and pericytes and smooth muscle cells

(361036, Biotin anti-human CD146, clone: P1H12, BioLegend). Fibroblasts were cultured in DMEM

supplemented with 10% serum, 1% penicillin-streptomycin, and 1% amphotericin B for a total of 5 days.

Human neutrophil isolation and coculture with fibroblasts. Whole blood (10ml) from heathy donors

was collected in BD Vacutainer® k-EDTA Tubes (Vitalant, San Francisco, CA) and used for primary

neutrophil isolation as previously described (54). Briefly, 7ml blood was layered on top of 7ml of

PolymorphPrep (Cosmo Bio Usa Inc, AXS-1114683) and centrifuged at 500 x *g* for 35 min at room

temperature without braking. The peripheral blood mononuclear cells and plasma layers were aspirated

and the neutrophil layer was collected. The cells were washed with PBS and centrifuged at 400g for 5

minutes. The cell pellet was resuspended in 3ml ACK lysis buffer (Thermo Fisher, A1049201) and

centrifuged at 400g for 5 minutes. Neutrophil purity was 95%, confirmed by flow cytometry

(**Supplemental Figure 3F;** Antibodies: CD15 clone W6D3, Biolegend cat# 323039; CD66B clone

G10F5, Biolegend cat# B221034; CD16 clone 3G8, BD Biosciences cat# 560474; CD14 Clone 63D3,

Biolegend cat# 367118). The cells were finally resuspended in RPMI supplemented with 10% FBS and

counted. Neutrophils were then added to human lung fibroblasts for 24 hours of coculture in RPMI with

10% FBS, with or without P2RX4 inhibitor BAY-1797 (1 μm, 7573, Tocris) or IL6-R blocking antibody

tocilizumab (100 ng/mL, HY-P9917, MedChemExpress). The conditioned media were retained for

analysis after separation of the cellular fraction by centrifugation, and fibroblasts were fixed and

permeabilized for immunofluorescence analysis.

Flow cytometry and sorting. *Cocultured cells:* Cocultures were trypsinized with 0.25% trypsin and

resuspended in 1X PBS with 0.5% BSA and 2mM EDTA. Single-cell suspensions were pre-stained with

Fc blocker for 10 min in ice followed by staining at 1:300 with anti-CD64 (139304, PE anti-mouse

CD64, clone:X54-5/7.1, BioLegend) and anti-Pdgfra (25-1401-82, PE-Cy7, anti-mouse CD140a, clone:

APA5, Invitrogen) antibodies for 40 min. DAPI was used to distinguish dead cells. *Mouse lung cell*

suspensions: Whole lung single cell suspensions were prepared by harvesting lung lobes into 5 ml

HBSS with 40µl Liberase Tm (0.1 U/ml, 5401127001, Roche) and 20µl DNAse 1 (10mg/ml, Roche,

Cat# 10104159001), followed by automated tissue dissociation (GentleMacs; Miltenyi Biotec) and

tissue digestion for 30 min at 37°C on a shaker. Digested samples were processed on the GentleMacs

using the "lung2" program, passed through 70µm filters, and washed, followed by red blood cell lysis

and final suspension in FACS buffer. Cells were counted using a NucleoCounter (Chemometic). All

samples were stained in 96-well V-bottom plates. Single cell samples were first incubated with

antibodies to surface antigens for 30 min at 4°C in 50µl staining volume. Flow cytometry was performed

on BD LSRFortessa X-20. Fluorochrome compensation was performed with single-stained UltraComp

eBeads (Invitrogen, Cat# 01-2222-42). Samples were FSC-A/SSC-A gated to exclude debris, followed

by FSC-H/FSC-A gating to select single cells and Draq7 viability dye (Biolegend) to exclude dead cells.

Monocyte-derived macrophages (moMacs) were identified as $CD19^-$, $Ly6G^-$, $NK1.1^-$, $Siglec-F^-$, $CD11b^+$,

and $CD64^+$. Arginase1-positive cells were identified by the presence of eYFP. Data were analyzed

using FlowJo software (TreeStar, USA) and compiled using Prism (GraphPad Software). Monoclonal

antibodies used were: anti-CD45 (30-F11, Biolegend), anti-CD11b (M1/70, Biolegend), anti-CD11c

(N418, Biolegend), anti-NK1.1 (PK136, Biolegend), anti-CD19 (6D5, Biolegend), anti-CD64 (X54-5/7,

eBiosciences), anti-Ly6G (1A8, Biolegend), anti-SiglecF (E50-2440), anti-I-A/I-E (MHCII) (M5/114.15.2,

Biolegend), and anti-CCR2 (475301, BD BioSciences). In the case of quantifying and sorting

CD45+CD11b+CD64+ monocyte-derived macrophages in bleomycin-injured WT and IL-6 KO mouse

lung cell suspensions, 50,000 microbeads (Invitrogen) were added to each sample for quantification of

absolute live CD45+CD11b+CD64+ cells. Lavage cell pellets were added to the suspensions prior to

sorting. Cells were sorted on a FACS Aria2.

Quantitative Real-Time PCR analysis. Human lung fibroblasts were lysed in 300µl TRIzol reagent

(10296010, Ambion life technologies) to obtained RNA. 600 ng of RNA per sample was used to prepare

cDNA using iScript reverse transcriptase supermix (Bio-Rad, 1708841). qPCR was performed for the

target genes using SYBR Green super mix (Applied Biosystems, 4309155). Gene expression was

normalized to 18S rRNA. The following primers were used:

P2RX4 Forward: GAGATTCCAGATGCGACCACT

P2RX4 Reverse: ACCCGTTGAAAGCTACGCAC

18S rRNA Forward: GTAACCCGTTGAACCCCATT

18S rRNA Reverse: CCATCCAATCGGTAGTAGCG

In the case of sorted CD45+CD11b+CD64+ murine lung monocyte-derived macrophages, mRNA was

isolated form pelleted cells as above, and qPCR was performed. Gene expression was normalized to

Gapdh. The following primers were used:

Arg1 Forward: CTCCAAGCCAAAGTCCTTAGAG

Arg1 Reverse : AGGAGCTGTCATTAGGGACATC

Gapdh Forward: AGTATGACTCCACTCACGGCAA

Gapdh Reverse: TCTCGCTCCTGGAAGATGGT

siRNA Knockdown. Healthy human fibroblasts were treated with 5μM siRNA (Dharmacon, D-001810-

01-05 ON-TARGET plus No-targeting siRNA) control or (Dharmacon, L-006285-00-0010 ON-TARGET

plus Human P2RX4 SMARTpool) using DharmaFECT in serum-free media for 12 hours, followed by

supplementing with DMEM with 10% serum. Cells were incubated for an additional 48 hours. Treated

cells were harvested for either RNA isolated or stimulated with 100μM ATPγS for 12 hours and culture

supernatant was collected for Human IL6 ELISA assay.

IL-6 ELISA. Bronchoalveolar lavage fluid from mice or conditioned media from cultured murine cells

were collected for measurement of IL-6 with a Duoset ELISA kit (R&D Systems, DY406-05).

Conditioned media from cultured human cells were collected and IL-6 was measured using Duoset

ELISA kit (R&D Systems, DT206-05). In both cases, the capture antibody was coated overnight in a 96-

well plate at room temperature. Wells were aspirated and washed in ELISA wash buffer and blocked by

adding reagent diluent consisting of 1% BSA in PBS. After 1 hour of room temperature incubation, wells

were again washed with wash buffer and samples or standards were added and incubated for 2 hours

at room temperature. Detection antibody was used along with streptavidin-HRP and substrate solution.

The reaction was stopped with stop solution, and optical density was measured using a microplate

reader at 450 nm.

Arginase 1 ELISA. Mouse: Arginase 1 levels were measured from conditioned media or from

bronchoalveolar lavage, using Mouse Arginase 1 ELISA Kit (ab269541, Abcam) based on the

manufacturer's protocol. Briefly, CD11b+ mouse lung cells were seeded in 12-well dishes for 2 days in

DMEM supplemented with M-Csf (315-02, Peprotech). Cells were exposed to murine IL-6 for 72 hours,

and culture supernatant and cell lysate were then collected. For *in vivo* experiments, bleomycin-injured

mice received bronchoalveolar lavage with 1 mL of PBS on day 14 after injury, and the noncellular

fraction was retained for analysis. For the ELISA, subsequently, 50 µl of undiluted samples were added

to each well, along with a 50 µl antibody cocktail comprising capture antibody and detection antibody

dissolved in antibody diluent. Plates were incubated for 1 hour at room temperature. Wells were

washed thrice with wash buffer and incubated with 100 µl Tetramethylbenzidine (TMB) substrate for 15

min. The reaction was stopped using 100µl stop solution, and OD was measured at 450nm. *For*

Human: Arginase 1 levels were measured from IPF precision-cut lung slice (PCLS) using the Human

Arginase 1 ELISA Kit (BMS2216, ThermoFisher), following the manufacturer's instructions. Briefly,

following 24 hours of culture with or without tocilizumab (100 ng/mL, HY-P9917, MedChemExpress),

PCLS were lysed in RIPA buffer with protease inhibitor, and 100 µl of standards or undiluted sample

was added to each well of a pre-coated 96-well plate and incubated for 2 hours at room temperature.

Wells were then washed four times with provided wash buffer before the addition of 100 µl of biotin-

conjugated anti-arginase 1 antibody to each well, followed by a 1-hour incubation. After another wash

step, 100 µl of streptavidin-horseradish peroxidase (HRP) was added and incubated for 30 minutes.

Wells were washed again and incubated with 100 µl of TMB substrate solution for 15 minutes. The

enzymatic reaction was stopped by adding 100 µl of stop solution, and optical density (OD) was

measured at 450 nm using a microplate reader.

Histology and Immunofluorescence imaging. *Mouse lung:* Arg1-RFP-creERT2 /Rosa26-LSL-

TdTomato/Col1a1-EGFP mice were injured for 14 days. Lungs from uninjured or injured mice were

inflated with a solution consisting of 30% sucrose solution mixed 1:1 with OCT (4585, OCT) and fixed in

4% formaldehyde at 4°C for 4 hours. Lungs were then washed and submerged in 30% sucrose solution

overnight at 4°C. Next, the tissue was incubated in the 1:1 solution of 30% sucrose and OCT overnight

followed by changing to OCT for 2 hours, and then a tissue block was frozen after embedding in OCT.

5 µm sections were cut from OCT-embedded tissue. For immunofluorescence imaging, frozen sections

were incubated with Mertk antibody (AF591, R&D Systems) or S100A8 antibody (AF3059, R&D

Systems) in staining buffer (1% BSA, 0.5% Triton x-100 in PBS), washed, and then incubated with

anti-rabbit secondary antibody (Invitrogen). Slides were washed with PBS and mounted on antifade

DAPI mounting media. *Human lung:* 5 µm thick sections from patient FFPE blocks were collected for

IHC staining. Slides were stained using Opal Manual IHC kit (PerkinElmer). After deparaffinization,

antigen retrieval was performed in AR buffer for 45 sec at 100% power followed by 15 min at 20%

power. After blocking slides were incubated with primary antibodies ARG1 (Cat# 93668, Cell Signaling

Technology) overnight at 4C. Polymer HRP was introduced to slides for 10 min, followed by signal

amplification using Opal570 (NEL810001KT,Akoya Biosciences) for 10 min at room temperature. The

slides were then counterstained with DAPI mounting media and scanned using with a 10x and 40x

objective of Leica inverted widefield microscope. *Cultured Cells:* Macrophages and fibroblasts or

fibroblasts alone, isolated from mouse lung as detailed above, were cultured on glass coverslips and in

some cases treated with CB-1158 (1 µM), ornithine (1 mM), or 1 µM OAT inhibitor 5-FMOrn

dihydrochloride (HY154021A, MedChemExpress). Cells were fixed with 4% PFA, permeabilized with

1% BSA, 0.5 % triton x-100 in PBS, and incubated with primary antibodies (Col1a1, #720265, Cell

Signaling Technology; α-SMA, A5228, Sigma) followed wash, secondary antibodies, further wash, and

mounting. Imaging was performed on a Leica SP8 laser scanning confocal microscope. Fluorescence

signal was quantified with Imaris software for cellular areas.

Western blot. Precision cut lung slices were obtained from human IPF tissue sections. Briefly, IPF

tissue was injected with 2% low melting agarose (50111, Lonza) and then submerged in ice-cold PBS

to allow solidification of agarose. 400 µm lung slices was generated using a Compresstome device (VF-

310-OZ, Precisionary Instruments). Slices were kept in complete DMEM media for 2 hours to allow the

release of agarose followed by changing to fresh complete DMEM with 10% FBS and 1% penicillin-

streptomycin. Lung slices were treated with 50 µM CB-1158 for 24 hours. After incubation slices were

minced using a tissue homogenizer. Cells were lysed in Pierce RIPA buffer (Thermo Fisher Scientific,

89901) with protease inhibitor cocktail (Thermo Fisher Scientific, 1861278). 10 µg of protein was run on

10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (Bio-rad, 4561034) and transferred to a

PVDF membrane (Thermo Fisher Scientific, 88520). The membrane was incubated with 1:1000 Col1a1

antibody (#720265, Cell Signaling Technology) overnight at 4°C. Blots were washed in 1X TBST and

incubated with peroxidase-conjugated goat anti-rabbit (1:20,000, Anaspec, AS28177) for 4 hr at 4°C.

These blots were developed using SuperSignal West Pico Chemiluminescent substrate (Thermo Fisher

Scientific, 34080) in ChemiDoc XRS+ gel imaging system (Bio-rad). Quantification of bands was done

using ImageJ.

Hydroxyproline assay. Mice were euthanized and lungs were excised and snap-frozen. Isolated lung

samples were homogenized and incubated with 50% trichloroacetic acid (T6399, Sigma) on ice for 20

min. Samples were then incubated in 12N HCL (A144, Fisher) overnight at 110°C. Dried pellets were

reconstituted in distilled water with constant shaking for 2 hours at room temperature. Samples were

then mixed with 1.4% Chloramine T (Sigma, 85739) and 0.5 M sodium acetate (Sigma, 241245) in 10%

2-propanol (Fisher, A416) and incubated with Ehrlich's solution (Sigma, 03891) for 15 min at 65°C.

Absorbance was quantified at 550 nm, and concentration was calculated using a standard curve of

commercial hydroxyproline (Sigma, H5534).

scRNAseq library preparation and sequencing. PIPseq T2 3' Single Cell RNA Kit (v3.0 – cultured

cells, v4.0 – directly sequenced cells) was used for pre-templated instant partitioning (PIP) to capture

single-cell mRNA transcripts with PIP beads according to manufacturer's protocol (Fluent Biosciences,

FB0001026). After 5 days of coculture, single cell suspension of cells was obtained by trypsinization

followed by washing in 1X PBS. Cells were washed in ice-cold PIPseq Cell Suspension Buffer (Fluent

Biosciences, FB0002440). Cells were counted and stained with Trypan blue to confirm >90% viability.

Single-cell library preparation was performed using the manufacturer-recommended default protocol

and settings. The sequencing libraries were submitted to UCSF Center for Advanced Technology

(Novaseq X) or Novogene (Novaseq 6000) for sequencing. The demultiplexed FASTQ files were

aligned to mouse genome (mm10) using PIPseeker 1.0.0 (Fluent Biosciences). After sequence

alignment, we observed around 50% of input cells being called when PIPseeker sensitivity level was

set to 3, which was near the inflection point of the barcode rank plot. The original FASTQ files, the

quality reports, and the expression matrix outputs of PIPseeker have been deposited in Gene

Expression Omnibus (GEO).

scRNAseq Data Analysis. The Seurat – Guided Clustering Tutorial (March 27, 2023) was followed to

convert our expression matrixes into Seurat objects (Seurat version 4.3.0) (55). For quality control, we

removed the cells with less than 200 genes or more than 10,000 genes and larger than 5%

mitochondrial content. Seurat objects were integrated following Seurat's Introduction to scRNAseq

Integration (March 27, 2023), selecting the top 2,000 variable features as integration anchors. Cell

doublets were removed with the package DoubletFinder (2.0.3) (56). Following Seurat – Guided

Clustering Tutorial (March 27, 2023), we selected the top 2000 highly variable genes to obtain the cell

UMAP coordinates and group the cells into clusters with a sensitivity of 0.5(55). Cell types were

annotated using the SingleR package 1.10.04(2) using the ImmGen database (57) as reference. The

Gene Set Enrichment Analysis (GSEA) was performed using the package enrichR 3.1(58) (FDR<0.05),

with gene ontology data taken from the database "GO_Biological_Process_2021"(59). Cell

communication pathways analysis was performed using the package CellChat (1.6.1) (27). Upstream

regulator prediction was done using the Ingenuity Pathway Analysis software (Qiagen) (60).

Differentially expressed genes for macrophages with $p < 0.05$ and average log2 fold change > 0.75

were used for analysis. Analysis results with $p < 0.05$ were considered significant.

Multiplexed Ion Beam Imaging and Analysis (MIBI). *Slide Preparation:* Serial 5 um FFPE sections

were cut onto one glass and one gold slide. Both slides were baked at 70* Celsius overnight and

deparaffinized in three washes of fresh Xylene and rehydrated in EtOH (2X 100% EtOH, 2X 90% EtOH,

1X 80% EtOH, 1X 70% EtOH) and distilled water (2X) washes. Washes were performed in (include

wash machine specs). An antigen retrieval slide chamber was prepared by diluting 10X Tris with EDTA

antigen retrieval buffer 1:10 in diH2O. The prepared slide chamber was added to a PT Module (add

specs) filled with PBS and preheated to 75ºC. The rehydrated slides were run in the preheated PT

module at 97ºC for 40 minutes, then cooled to 65ºC in the PT module. The prepared slide chamber was

then removed from the PT module and cooled at RT for 30 minutes. Slides were washed 2X in 1X TBS-

Tween. *Glass Slide IHC:* Tissues were encircled by PAP pen boarders and blocked with 5% donkey

serum diluted in TBS IHC wash buffer for 1 hour at RT in a moisture chamber. Wash buffer was

aspirated from the slide, and Arginase-1 primary antibody (Ionpath Cat#: 715001) was stained

overnight at 4ºC overnight in the moisture chamber. The following day, primary antibody was aspirated,

and slides were washed twice with 1X TBS-T for 5 minutes, blocked with 3% peroxide buffer for 15

minutes at RT, and washed again twice with 1X TBS-T for 5 minutes each wash before incubation with

anti-rabbit secondary antibody for 1 hour at RT. Slides were washed twice with 1X TBS-T for 5 minutes

before visualization with 100uL DAB for 5 minutes at RT. DAB reaction was stopped by tapping waste

into contained waste bin, and then washing the slide into a slide chamber filled with diH2O 3X for 30

seconds each wash. The slide was then stained with Hematoxylin for 1 minute at RT and washed with

tap water 2X for 30 seconds. The slide was then dehydrated by washing in EtOH (1X 70% EtOH, 1X

80% EtOH, 2X 95% EtOH, 2X 100% EtOH), and Xylene (2X) before cover slipping. *Gold Slide Staining:*

Gold slides were transferred the Sequenza Immunostaining Center Staining System (Electron

Microscopy Sciences, Hatfield, PA). Endogenous biotin-binding proteins with blocked with

(Avidin/biotin-blocking reagents) for 30 minutes at room temperature. 5% donkey serum was added to

the top of the chamber to wash out the avidin blocking reagents and block additional non-specific

antibody binding sites for 1 hour at RT. Primary and Secondary antibodies panels were assembled with

appropriate volumes of each titrated antibody and a final concentration of 0.05M EDTA. The complete

cocktails were filtered through a pre-wet 0.1 um Ultragree MC Spin Filter, and then the Primary

Antibody cocktail (**Supplemental Table 2**) was added to the Sequenza top chamber and incubated

overnight at 4ºC. The Secondary antibody cocktail was stored at 4ºC. The following day, the slides

were washed by adding 1X TBS-T to the Sequenza chamber 2X before adding the Secondary Antibody

cocktail to the Sequenza chamber for 1 hour at RT. Gold slides were removed from the Sequenza

chamber and washed 3X with 1X TBS-T for 5 minutes each wash, 1X with filtered 2% Glutaraldehyde

for 5 minutes, 3X with filtered 1X Tris pH 8.5, 2X with filtered diH2O, 1X with 70% EtOH, 1X with 80%

EtOH, 2X with 90% EtOH, and 2X with 100% EtOH. Slides were allowed to dry at RT for 10 minutes

before being stored in a vacuum chamber (Cat#) before MIBIscope analysis. *Image Acquisition and*

Processing: Gold slides were loaded into the MIBIscope (Ionpath, Menlo Park, CA) and FOVs were

selected by matching tissue topography to ROIs with Arginase-1+ Staining from the serial IHC glass

slide. FOVs were acquired at Fine resolution, with a dwell time of 1 second at a resolution of 0.39 um

per pixel. Image QC was performed by following the Angelo Lab toffy pipeline

(https://github.com/angelolab/toffy). Analysis was performed by following the Angelo Lab ark pipeline

(https://github.com/angelolab/ark-analysis).

Spatial Transcriptomics (10x Xenium). FFPE preserved sections of lung tissue were prepared for

spatial transcriptomics imaging on the Xenium platform by following 10x Genomics protocols

CG000582 Rev E and CG000584 Rev A. Briefly, protocol CG000582 prepares the slides for the

imaging run by first hybridizing the probe panel of choice. Here, the standard human lung panel

available from 10x Genomics (Cat #1000601) was supplemented with a custom panel specific for lung

disease states, including pulmonary fibrosis (**Supplemental Table 3**). Following probe hybridization,

annealed probes are ligated together to create circular fragments, then circularized probes are

amplified with a rolling circle PCR. After rolling circle amplification, slides are DAPI stained for nuclei

then placed in the Xenium Analyzer with run reagents following CG000584. The gene panel is

uploaded to the Xenium Analyzer and a primary image is take of the slides. All areas with visible tissue

are selected for imaging and the imaging run is started. The Xenium processes imaging data during the

run, identified cell boundaries and outputting image files along with cell by transcript by location

matrices for further downstream analysis by Seurat 10x Xenium protocol. *Proximity Analysis of ARG1+*

cells and IL6+ fibroblasts: The output files of Xenium Analyzer were converted to an AnnData object

that contained the cell centroid coordinates and raw transcript counts. A filter of 10 counts per cell and

5 cells per gene was applied to filter out low-quality cells and sparsely detected genes. Counts were

normalized to 1e4 and log-transformed. Cells were then grouped into one of the following three

categories: (1) ARG1+; (2) IL6+ARG1- and either CTHRC1 or COL1A1; and (3) IL6+ARG1- and

CTHRC1 or COL1A1 , with positive expression of a marker defined as having non-zero counts. *Co-*

occurrence probability ratio: The 'Analyze Xenium data' tutorial in Squidpy (version 1.5.0) (61) was

followed to compute the co-occurrence probability ratio of ARG1+ and fibroblasts as defined above. We

performed 50% random subsampling of the data 5 times to compute the statistic and generate

confidence intervals bounded by the minimum and maximum probability ratio generated by the

subsamples. The co-occurrence probability ratio was computed at 25 µm between cells. Using the

619 implementation in the 1.5.0 release of Squidpy, for each radial distance, the ratio of cells belonging to

620 category exp (exp being any of the three categories of cell defined above) out of all cells within the

621 given distance of an ARG1+ cell was averaged across all ARG1+ cells to compute the conditional

622 probability in the numerator P(exp | ARG1+), while the denominator P(exp) was computed similarly but

623 averaging probabilities that were computed by centering around every cell in the sample. Previous

624 releases of Squidpy implemented a version of this test in the function gr.co_occurrence that used

625 discrete interval bins (only including cells within two consecutive choices of radial distances). We chose

626 to use inclusive intervals (including all cells within a given radial distance) for a more robust calculation

627 of the co-occurrence probability ratio. We implemented our method into the codebase of Squidpy, with

628 the help of Squidpy's authors, and it is now the default implementation of the function

629 gr.co_occurrence, starting in release 1.5.0.

630 ***Analysis of published datasets.*** *Bleomycin time course:* Time series of single-cell data after

631 bleomycin lung injury were obtained from Tsukui et al (8) and Strunz et al (9). These samples were

632 processed and merged with Seurat (4.3.0) using the function SCTransform to correct for batch effects

633 and annotated with SingleR 1.10.044 using the ImmGen database as reference to identify cell types.

634 *Macrophage Annotation in vivo:* We annotated macrophages from Strunz et al. (9) according to the

635 macrophage subtypes (C1, C2, C3) defined by Aran et al (2). and by Li et al (11) using SingleR(2). *BAL*

636 *Microarray:* Microarray RNA data from bronchoalveolar cells of healthy individuals and IPF patients was

637 extracted from the GPL14550 dataset within the GSE70867(12) repository. The differentially expressed

638 genes between IPF and healthy samples were obtained by following the R workflow provided by the

639 NCBI GEO2R platform using GEOquery (2.66.0) and limma (3.54.2) packages. *Gene expression of*

640 *cultured mouse lung-derived macrophages:* The top markers of macrophage subtypes as defined by Li

641 et al. (11) and Aran et al.(2) were quantified in our CD11b+, CSF1-treated macrophages (from the WT-

642 WT coculture condition). *Analysis of lung tumor neutrophils:* A high-resolution single-cell atlas of tumor-

643 associated neutrophils in non-small cell lung cancer was obtained from (32). The scANVI algorithm-

644 based integrated NSCLC transcriptome atlas provided more than 1.2M cells from 19 studies and 309

645 patients. The metadata embedded with Leiden-clustering and cell-type annotations were utilized to

identify and subset neutrophils. The subset data was log transformed and scaled, followed by filtering

for cells expressing both ARG1 and IL6R. Pearson correlation coefficient, p-value and R-squared

values were calculated for the correlation between ARG1 and IL6R.

Ornithine Measurement. Cultured PCLS from IPF explanted lungs lysed in water, IPF lung samples

lysed in RIPA buffer, or mouse lung homogenized and lysed in RIPA buffer were used to measure

ornithine following manufacturer's protocol (IS I-1000R, Immusmol). Briefly, each sample was

precipitated with 50 µL precipitating reagents (provided by manufacturer) followed by extraction using

D-reagent (a derivatizing agent provided by manufacturer) for 2h at room temperature. 10 µL of the

prepared standards, controls and samples were incubated with 90 µL of antiserum overnight at 4 C.

Subsequently, plates were washed and briefly incubated with enzyme conjugate for 30min followed by

addition of substate. Absorbance was measured using microplate reader at 450nm.

LC-MS metabolomics. Primary murine lung macrophages and fibroblasts were cocultured with or

without 1 µM CB-1158. Fibroblasts were isolated by antibody-coated bead-based negative selection of

macrophages. Fibroblast extracts were used to calculate protein equivalents by resuspension in 0.2 M

NaOH, heated to 95 °C for 25 min, and determined via BCA (Pierce, 23225). Dried metabolites were

resuspended in 50% ACN:water and 1/10th of the volume was loaded onto a Luna 3 um NH2 100A

(150 × 2.0 mm) column (Phenomenex). The chromatographic separation was performed on a Vanquish

Flex (Thermo Scientific) with mobile phases A (5 mM NH_4AcO pH 9.9) and B (ACN) and a flow rate of

200 µL/min. A linear gradient from 15% A to 95% A over 18 min was followed by 9 min isocratic flow at

95% A and re-equilibration to 15% A. Metabolites were detected with a Thermo Scientific Q Exactive

mass spectrometer run with polarity switching (+3.5 kV / -3.5 kV) in full scan mode with an m/z range of

65-975. TraceFinder 4.1 (Thermo Scientific) was used to quantify the targeted metabolites by area

under the curve using expected retention time and accurate mass measurements (< 5 ppm). Values

were normalized to cell number and sample protein concentration. Relative amounts of metabolites

were calculated by summing up the values for all isotopologues of a given metabolite.

Study Approval. Experiments in mice were performed in accordance with approved protocols by the

University of California, San Francisco Institutional Animal Care and Use Committee. The studies with

human tissues described in this paper were conducted according to the principles of the Declaration of

Helsinki. Written, informed consent was obtained from all subjects, and the study was approved by the

University of San Francisco, California and Vitalant institutional review board. With respect to deceased

donor control explanted lungs, because samples were acquired from deceased individuals the study is

not considered human subjects research as per UCSF and NIH policy.

Statistics. One-way ANOVA followed by post hoc Sidak's multiple comparisons tests was used for

comparisons among more than two groups, and Student's t-test was used for comparison between two

groups. A p value less than 0.05 was considered significant. Analysis appropriately corrects for multiple

comparisons and repeated measures.

Data availability. Newly generated single-cell sequencing data will be made public on acceptance for

publication at GSE242510.

References

<u>References</u>

1. Wolters PJ, Blackwell TS, Eickelberg O, Loyd JE, Kaminski N, Jenkins G, et al. Time for a

change: is idiopathic pulmonary fibrosis still idiopathic and only fibrotic? *Lancet Respir Med.*

2018;6(2):154-60.

2. Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, et al. Reference-based analysis of lung single-

cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol.* 2019;20(2):163-72.

3. Joshi N, Watanabe S, Verma R, Jablonski RP, Chen CI, Cheresh P, et al. A spatially restricted

fibrotic niche in pulmonary fibrosis is sustained by M-CSF/M-CSFR signalling in monocyte-

derived alveolar macrophages. *Eur Respir J.* 2020;55(1).

4. Riteau N, Gasse P, Fauconnier L, Gombault A, Couegnat M, Fick L, et al. Extracellular ATP is a

danger signal activating P2X7 receptor in lung inflammation and fibrosis. *Am J Respir Crit Care*

Med. 2010;182(6):774-83.

5. Bhattacharyya A, Torre P, Yadav P, Boostanpour K, Chen TY, Tsukui T, et al. Macrophage

Cx43 Is Necessary for Fibroblast Cytosolic Calcium and Lung Fibrosis After Injury. *Front*

Immunol. 2022;13:880887.

6. Buechler MB, Pradhan RN, Krishnamurty AT, Cox C, Calviello AK, Wang AW, et al. Cross-

tissue organization of the fibroblast lineage. *Nature.* 2021;593(7860):575-9.

7. Tsukui T, Sun KH, Wetter JB, Wilson-Kanamori JR, Hazelwood LA, Henderson NC, et al.

Collagen-producing lung cell atlas identifies multiple subsets with distinct localization and

relevance to fibrosis. *Nat Commun.* 2020;11(1):1920.

8. Tsukui T, Wolters PJ, and Sheppard D. Alveolar fibroblast lineage orchestrates lung

inflammation and fibrosis. *Nature.* 2024;631(8021):627-34.

9. Strunz M, Simon LM, Ansari M, Kathiriya JJ, Angelidis I, Mayr CH, et al. Alveolar regeneration

through a Krt8+ transitional stem cell state that persists in human lung fibrosis. *Nat Commun.*

2020;11(1):3559.

10.   Fabre T, Barron AMS, Christensen SM, Asano S, Bound K, Lech MP, et al. Identification of a

broadly fibrogenic macrophage subset induced by type 3 inflammation. *Sci Immunol.*

2023;8(82):eadd8945.

11.   Li X, Mara AB, Musial SC, Kolling FW, Gibbings SL, Gerebtsov N, et al. Coordinated chemokine

expression defines macrophage subsets across tissues. *Nat Immunol.* 2024;25(6):1110-22.

12.   Prasse A, Binder H, Schupp JC, Kayser G, Bargagli E, Jaeger B, et al. BAL Cell Gene

Expression Is Indicative of Outcome and Airway Basal Cell Involvement in Idiopathic Pulmonary

Fibrosis. *Am J Respir Crit Care Med.* 2019;199(5):622-30.

13.   Zhang H, Zhu X, Friesen TJ, Kwak JW, Pisarenko T, Mekvanich S, et al. Annexin

A2/TLR2/MYD88 pathway induces arginase 1 expression in tumor-associated neutrophils. *J*

Clin Invest. 2022;132(22).

14.   Rotondo R, Barisione G, Mastracci L, Grossi F, Orengo AM, Costa R, et al. IL-8 induces

exocytosis of arginase 1 by neutrophil polymorphonuclears in nonsmall cell lung cancer. *Int J*

Cancer. 2009;125(4):887-93.

15.   Cane S, Barouni RM, Fabbi M, Cuozzo J, Fracasso G, Adamo A, et al. Neutralization of NET-

associated human ARG1 enhances cancer immunotherapy. *Sci Transl Med.*

2023;15(687):eabq6221.

16.   Angelo M, Bendall SC, Finck R, Hale MB, Hitzman C, Borowsky AD, et al. Multiplexed ion beam

imaging of human breast tumors. *Nat Med.* 2014;20(4):436-42.

17.   Keren L, Bosse M, Marquez D, Angoshtari R, Jain S, Varma S, et al. A Structured Tumor-

Immune Microenvironment in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam

Imaging. *Cell.* 2018;174(6):1373-87 e19.

18.   Keren L, Bosse M, Thompson S, Risom T, Vijayaragavan K, McCaffrey E, et al. MIBI-TOF: A

multiplexed imaging platform relates cellular phenotypes and tissue structure. *Sci Adv.*

2019;5(10):eaax5851.

19.   Rahim MK, Okholm TLH, Jones KB, McCarthy EE, Liu CC, Yee JL, et al. Dynamic CD8(+) T cell

responses to cancer immunotherapy in human regional lymph nodes are disrupted in metastatic

lymph nodes. *Cell.* 2023;186(6):1127-43 e18.

20.   Achaiah A, Fraser E, Saunders P, Hoyles RK, Benamore R, and Ho LP. Neutrophil levels

correlate with quantitative extent and progression of fibrosis in IPF: results of a single-centre

cohort study. *BMJ Open Respir Res.* 2023;10(1).

21.   Pitre T, Lupas D, Ebeido I, Colak A, Modi M, Kachkovski GV, et al. Prognostic factors

associated with mortality in acute exacerbations of idiopathic pulmonary fibrosis: A systematic

review and meta-analysis. *Respir Med.* 2024;222:107515.

22.   Kono M, Miyashita K, Hirama R, Oshima Y, Takeda K, Mochizuka Y, et al. Prognostic

significance of bronchoalveolar lavage cellular analysis in patients with acute exacerbation of

interstitial lung disease. *Respir Med.* 2021;186:106534.

23.   Kinder BW, Brown KK, Schwarz MI, Ix JH, Kervitsky A, and King TE, Jr. Baseline BAL

neutrophilia predicts early mortality in idiopathic pulmonary fibrosis. *Chest.* 2008;133(1):226-32.

24.   Tran DH, Kesavan R, Rion H, Soflaee MH, Solmonson A, Bezwada D, et al. Mitochondrial

NADP(+) is essential for proline biosynthesis during cell growth. *Nat Metab.* 2021;3(4):571-85.

25.   Steggerda SM, Bennett MK, Chen J, Emberley E, Huang T, Janes JR, et al. Inhibition of

arginase by CB-1158 blocks myeloid cell-mediated immune suppression in the tumor

microenvironment. *J Immunother Cancer.* 2017;5(1):101.

26.   Wei Y, Dong W, Jackson J, Ho TC, Le Saux CJ, Brumwell A, et al. Blocking LOXL2 and

TGFbeta1 signalling induces collagen I turnover in precision-cut lung slices derived from

patients with idiopathic pulmonary fibrosis. *Thorax.* 2021;76(7):729-32.

27.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, et al. Inference and analysis

of cell-cell communication using CellChat. *Nat Commun.* 2021;12(1):1088.

28.   Roofeh D, Lin CJF, Goldin J, Kim GH, Furst DE, Denton CP, et al. Tocilizumab Prevents

Progression of Early Systemic Sclerosis-Associated Interstitial Lung Disease. *Arthritis

Rheumatol.* 2021;73(7):1301-10.

29.   Group RC. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a

randomised, controlled, open-label, platform trial. *Lancet.* 2021;397(10285):1637-45.

30.   Khanna D, Lin CJF, Furst DE, Goldin J, Kim G, Kuwana M, et al. Tocilizumab in systemic

sclerosis: a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Respir Med.*

2020;8(10):963-74.

31.   Narvaez J, J LL, Alegre Sancho JJ, Molina-Molina M, Nolla JM, and Castellvi I. Effectiveness

and safety of tocilizumab for the treatment of refractory systemic sclerosis associated interstitial

lung disease: a case series. *Ann Rheum Dis.* 2019;78(11):e123.

32.   Salcher S, Sturm G, Horvath L, Untergasser G, Kuempers C, Fotakis G, et al. High-resolution

single-cell atlas reveals diversity and plasticity of tissue-resident neutrophils in non-small cell

lung cancer. *Cancer Cell.* 2022;40(12):1503-20 e8.

33.   Idzko M, Ferrari D, and Eltzschig HK. Nucleotide signalling during inflammation. *Nature.*

2014;509(7500):310-7.

34.   Trang T, Beggs S, Wan X, and Salter MW. P2X4-receptor-mediated synthesis and release of

brain-derived neurotrophic factor in microglia is dependent on calcium and p38-mitogen-

activated protein kinase activation. *J Neurosci.* 2009;29(11):3518-28.

35.   Le Guilcher C, Garcin I, Dellis O, Cauchois F, Tebbi A, Doignon I, et al. The P2X4 purinergic

receptor regulates hepatic myofibroblast activation during liver fibrogenesis. *J Hepatol.*

2018;69(3):644-53.

36.   Werner S, Mesch S, Hillig RC, Ter Laak A, Klint J, Neagoe I, et al. Discovery and

Characterization of the Potent and Selective P2X4 Inhibitor N-[4-(3-Chlorophenoxy)-3-

sulfamoylphenyl]-2-phenylacetamide (BAY-1797) and Structure-Guided Amelioration of Its

CYP3A4 Induction Profile. *J Med Chem.* 2019;62(24):11194-217.

37.   Summer R, Todd JL, Neely ML, Lobo LJ, Namen A, Newby LK, et al. Circulating metabolic

profile in idiopathic pulmonary fibrosis: data from the IPF-PRO Registry. *Respir Res.*

2024;25(1):58.

38.   Lee JU, Song KS, Hong J, Shin H, Park E, Baek J, et al. Role of lung ornithine

aminotransferase in idiopathic pulmonary fibrosis: regulation of mitochondrial ROS generation

and TGF-beta1 activity. *Exp Mol Med.* 2024;56(2):478-90.

39.   Pesce JT, Ramalingam TR, Mentink-Kane MM, Wilson MS, El Kasmi KC, Smith AM, et al.

Arginase-1-expressing macrophages suppress Th2 cytokine-driven inflammation and fibrosis.

PLoS Pathog. 2009;5(4):e1000371.

40.   Raghu G, Richeldi L, Crestani B, Wung P, Bejuit R, Esperet C, et al. SAR156597 in idiopathic

pulmonary fibrosis: a phase 2 placebo-controlled study (DRI11772). *Eur Respir J.* 2018;52(6).

41.   Wilson MS, Madala SK, Ramalingam TR, Gochuico BR, Rosas IO, Cheever AW, et al.

Bleomycin and IL-1beta-mediated pulmonary fibrosis is IL-17A dependent. *J Exp Med.*

2010;207(3):535-52.

42.   Ruscitti C, Abinet J, Marechal P, Meunier M, de Meeus C, Vanneste D, et al. Recruited atypical

Ly6G(+) macrophages license alveolar regeneration after lung injury. *Sci Immunol.*

2024;9(98):eado1227.

43.   Czystowska-Kuzmicz M, Sosnowska A, Nowis D, Ramji K, Szajnik M, Chlebowska-Tuz J, et al.

Small extracellular vesicles containing arginase-1 suppress T-cell responses and promote tumor

growth in ovarian carcinoma. *Nat Commun.* 2019;10(1):3000.

44.   Zhang H, Liu J, Qu D, Wang L, Wong CM, Lau CW, et al. Serum exosomes mediate delivery of

arginase 1 as a novel mechanism for endothelial dysfunction in diabetes. *Proc Natl Acad Sci U*

S A. 2018;115(29):E6927-E36.

45.   Roque W, and Romero F. Cellular metabolomics of pulmonary fibrosis, from amino acids to

lipids. *Am J Physiol Cell Physiol.* 2021;320(5):C689-C95.

46.   Foo SS, Turner CJ, Adams S, Compagni A, Aubyn D, Kogata N, et al. Ephrin-B2 controls cell

motility and adhesion during blood-vessel-wall assembly. *Cell.* 2006;124(1):161-73.

47.   Clausen BE, Burkhardt C, Reith W, Renkawitz R, and Forster I. Conditional gene targeting in

macrophages and granulocytes using LysMcre mice. *Transgenic Res.* 1999;8(4):265-77.

48.   Ozaki T, Muramatsu R, Sasai M, Yamamoto M, Kubota Y, Fujinaka T, et al. The P2X4 receptor

is required for neuroprotection via ischemic preconditioning. *Sci Rep.* 2016;6:25893.

49.   El Kasmi KC, Qualls JE, Pesce JT, Smith AM, Thompson RW, Henao-Tamayo M, et al. Toll-like

receptor-induced arginase 1 in macrophages thwarts effective immunity against intracellular

pathogens. *Nat Immunol.* 2008;9(12):1399-406.

50.   Yata Y, Scanga A, Gillan A, Yang L, Reif S, Breindl M, et al. DNase I-hypersensitive sites

enhance alpha1(I) collagen gene expression in hepatic stellate cells. *Hepatology.*

2003;37(2):267-76.

51.   Schneider C, Lee J, Koga S, Ricardo-Gonzalez RR, Nussbaum JC, Smith LK, et al. Tissue-

Resident Group 2 Innate Lymphoid Cells Differentiate by Layered Ontogeny and In Situ

Perinatal Priming. *Immunity.* 2019;50(6):1425-38 e5.

52.   Reese TA, Liang HE, Tager AM, Luster AD, Van Rooijen N, Voehringer D, et al. Chitin induces

accumulation in tissue of innate immune cells associated with allergy. *Nature.*

2007;447(7140):92-6.

53.   Kopf M, Baumann H, Freer G, Freudenberg M, Lamers M, Kishimoto T, et al. Impaired immune

and acute-phase responses in interleukin-6-deficient mice. *Nature.* 1994;368(6469):339-42.

54.   Caudrillier A, Kessenbrock K, Gilliss BM, Nguyen JX, Marques MB, Monestier M, et al. Platelets

induce neutrophil extracellular traps in transfusion-related acute lung injury. *J Clin Invest.*

2012;122(7):2661-71.

55.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated

analysis of multimodal single-cell data. *Cell.* 2021;184(13):3573-87 e29.

56.   McGinnis CS, Murrow LM, and Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell

RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* 2019;8(4):329-37 e4.

57.   Heng TS, Painter MW, and Immunological Genome Project C. The Immunological Genome

Project: networks of gene expression in immune cells. *Nat Immunol.* 2008;9(10):1091-4.

58.   Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: interactive and

collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics.* 2013;14:128.

59. Gene Ontology C, Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, et al. The Gene

Ontology knowledgebase in 2023. *Genetics.* 2023;224(1).

60. Kramer A, Green J, Pollard J, Jr., and Tugendreich S. Causal analysis approaches in Ingenuity

Pathway Analysis. *Bioinformatics.* 2014;30(4):523-30.

61. Palla G, Spitzer H, Klein M, Fischer D, Schaar AC, Kuemmerle LB, et al. Squidpy: a scalable

framework for spatial omics analysis. *Nat Methods.* 2022;19(2):171-8.

62. Satija R, Farrell JA, Gennert D, Schier AF, and Regev A. Spatial reconstruction of single-cell

gene expression data. *Nat Biotechnol.* 2015;33(5):495-502.

Acknowledgements

This work was supported by funding from the US Department of Defense (Grant W81XWH2110417 to

MB), the UCSF Bakar Aging Research Institute (Investigator Grant to MB), Longevity Impetus Grants

from Norn Group (MB and JGO), a UCSF Bakar Aging Research Institute investigator award (MB), the

Nina Ireland Program for Lung Health Innovative Grant Program (PY), and NIH (1R01HL142701-01 to

ABM and R01AI172754 SKP). The authors would like to acknowledge the staff within the Biological

Imaging Development CoLab (BIDC) at UCSF Parnassus Heights, particularly Kyle Marchuk and Austin

Edwards, for their support in microscopy experiments. Sequencing performed at the UCSF CAT was

supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01. We are grateful to Dr. Clifford

Lowell for his valuable comments and suggestions on the manuscript. We thank Giovanni Palla and

Nathan Levy for help in extending the Squidpy package.

Author contributions

PY performed or assisted in all the experiments and in figure preparation. JGO performed single cell

sequencing experiments and analysis and figure preparation. KC and SP performed single cell library

preparation and sequencing, supervised by ARA. NB and AB assisted in functional experiments with

lung tissues and cells. PD assisted in human neutrophil isolation and coculture experiments under the

supervision of SKP. XY assisted PY in coculture experiments and microscopy under the supervision of

BL. JN performed flow cytometry of mouse lung cells supervised by ABM. KB, TJ, and JW performed

mouse breeding and development of crosses for experiments. TT and DS assisted in analysis of single

cell sequencing data from mouse lung fibroblasts. MM and HT provided deceased donor control lung

tissues supervised by MAM. LM, AG, CM, and VA performed Xenium spatial transcriptomic analysis

supervised by WE. CC and AW designed and implemented spatial proximity analysis of Xenium data.

RM generated and provided P2x4 floxed mice. WT and ST performed MIBI analysis under the guidance

of TB. PJW provided lung explants from IPF patients and helped to interpret associated imaging data.

KMT designed and performed metabolomic assays, assisted by PY. MB conceived of the work,

supervised experimental planning and execution, and wrote the manuscript with input from PJW and

KMT.

Figure 1: Arginase 1-expressing cells localize to the fibrotic niche in murine and human lung.

A) Volcano plot for macrophages from scRNAseq of macrophage-fibroblast cocultures for WT macrophages and fibroblasts versus WT macrophages cocultured with P2rx4 KO (*Pdgfrb-Cre: P2rx4 f/f*) fibroblasts. Significance was determined by Wilcoxon Rank Sum test corrected for multiple comparisons by Bonferroni method. Color-labeled genes had p<0.05 and absolute value of log2 fold change >0.75. Data represent n=2 separate cocultures.

B) Analysis of macrophages from bleomycin lung injury (data reanalyzed from[7]). Top left: Annotation of macrophages from multiple time points, according to C1, C2, or C3 macrophage annotation (as described in (7): C1=alveolar macrophages; C2=transitional monocyte-derived macrophages; C3=monocyte derived macrophages). Top right: Proportions of C1, C2, and C3 across time. Bottom left: Feature plot of Arg1 expression. Bottom right: Violin plot of Arg1 expression according to cluster (C1, C2, and C3). ***p<0.001 by Student's t-test.

C) Immunofluorescence imaging of tamoxifen-induced Arg1-RFP-CreERT2: LSL-tdTomato-
Col1a1-GFP mice at 14 days after bleomycin injury. Quantification shows total Arg1+ cells
inside and outside contiguous Col1a1-GFP+ areas. *p<0.05 by Student's t-test. +/- SEM.
D) *ARG1* expression by microarray analysis of gene expression of BAL cells from healthy (20
patients) and IPF (112 patients) from GSE70867(12). ****p<0.0001 by Mann-Whitney test. +/-
SEM.
E) Representative immunofluorescence of human healthy control and IPF lung sections (n=4 and
n=6, respectively) for ARG1. The quantification shows the proportion of ARG1+ cells (i.e.,
ARG1+ / total DAPI count) for each sample. *p<0.05 by Student's t-test. +/- SEM.
902
903

Figure 2: Arginase 1 is expressed predominantly by neutrophils in IPF.

A) Schematic of multiplexed ion beam imaging (MIBI) and heatmap of scaled marker expression from MIBI of sections from n=5 IPF explanted lungs.

B) MIBI UMAPs by cluster and individual markers.

910   C) MIBI images for individual markers within a representative field of view of IPF lung.
911

Figure 3: Arginase 1 regulates lung collagen via ornithine production.

A) Ornithine concentration measured in lysates of mouse lung (N=5 uninjured and N=6 injured mice), healthy human donor lung (N=9) and patient IPF lung explants (N=10). ****p<0.0001, ***p<0.001 by Student's t-test. +/- SEM.

B) Ornithine concentration with and without CB-1158 treatment of lungs of bleomycin-injured mice (N=8 in each condition) and cultured IPF PCLS (N=16 slices from a total of 3 patients). ****p<0.0001 by Student's t-test. +/- SEM.

C) Hydroxyproline assay for collagen content of lungs from injured and uninjured WT mice treated with or without Arg1 inhibitor CB-1158. Mice were dosed with 100 mg/kg of CB-1158 twice daily during days 9-15 post bleomycin. N=6, 7, 8 mice, left to right. ****$p<0.0001$ by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. Median, upper, and lower quartiles indicated by dashed lines.

D) Representative immunoblot for COL1A1 from lysates of precision-cut IPF lung slices cultured for 24 hours with or without Arg1 inhibitor CB-1158. Quantitation is for slices from a total of n=3 patients. ***$p<0.001$ by Student's t-test. +/- SEM.

E) Hydroxyproline assay for collagen content of lungs from mice 21 days after bleomycin injury. N=10 and 9 mice, left to right. *$p<0.05$ by Mann-Whitney test. Median, upper, and lower quartiles indicated by dashed lines.

F) *Left:* Col1a1 immunofluorescence of monocultured mouse lung fibroblasts with or without ornithine treatment. Quantification is for n=3 separate cultures each. *$p<0.05$ by Student's t-test. *Right:* COL1A1 immunofluorescence of monocultured human lung fibroblasts with or without ornithine treatment. Quantification is for n=3 separate cultures each. *$p<0.05$ by Student's t-test. +/- SEM.

G) Hydroxyproline assay for collagen content of lungs from injured WT mice treated with or without ornithine (2 g/kg) twice daily by ornithine gavage during days 7 through 20. N=8, 7 mice, left to right. ***$p<0.001$, ****$p<0.0001$ by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. Median, upper, and lower quartiles indicated by dashed lines.

H) Representative samples of Col1a1 immunofluorescence of mouse lung macrophage-fibroblast cocultures from WT or fibroblast-specific P2rx4 KO (Pdgfrb-Cre: P2rx4 f/f) mice treated with or without CB-1158 or ornithine.

I) Quantitation of Mean Fluorescence Intensity (MFI) from (H). N=3 biological replicates per condition. ****$p<0.0001$ by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. +/- SEM.

J) Representative samples of Col1a1 immunofluorescence of mouse lung macrophage-fibroblast cocultures treated with or without CB-1158. "Mac$_{\Delta Arg1}$" indicates macrophages isolated from Lysm-Cre: Arg1 f/f mice, whereas "Mac" indicates Arg1 f/f controls. **$p<0.01$, ***$p<0.0001$ by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. +/- SEM.

K) Relative quantities of proline and hydroxy-proline in lysates of murine primary lung fibroblasts isolated after coculture with macrophages, with or without CB-1158 inhibitor treatment, quantified by liquid chromatography-mass spectrometry. N=3 biological replicates for macrophages and fibroblasts, respectively. ****$p<0.0001$ by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. +/- SEM.

960
Figure 4: IL-6 is necessary for Arginase 1 expression after bleomycin injury in mice.
A) Left: Ingenuity Pathways Analysis of predicted upstream regulators for macrophages cultured
with WT relative to P2rx4 KO fibroblasts. Right: CellChat plot comparing WT and P2rx4
conditions. Significance was determined by Wilcoxon test with $p<0.05$. Data represent n=2
separate cocultures.
B) Violin plot for IL-6 expression in WT relative to P2rx4 KO fibroblasts. Line shows median values.
Data represent n=2 separate cocultures.
C) Arg1 ELISA of cell lysates and conditioned media collected from mouse lung macrophage
monoculture 72 hours after IL-6 treatment. N=3 biological replicates per condition. ***$p<0.001$,
****$p<0.0001$ by Student's t-test. +/- SEM.
D) Left: Arg1 ELISA of bronchoalveolar lavage fluid normalized to lung CD11b+CD64+
macrophage count taken at day 14 from bleomycin-injured WT or IL6 KO mice. Right: Arg1
qPCR of CD11b+CD64+ macrophages. N=5 mice per condition. **$p<0.01$ by Student's t test. +/-
SEM.
E) Quantification of *Il6* expression by cell type in cells sequenced by scRNAseq directly after
isolation from the lung at steady state and multiple time points after injury (reanalysis of merged
data(8, 9) normalized by Sctransform(62)).
F) UMAP of fibroblast subtypes (left) and IL-6 feature plots at steady state (center) and 7 days
after bleomycin injury (right), from Tsukui et al.(8)

980
Figure 5: IL-6 is necessary for Arginase 1 expression in IPF lung and in cocultures of human
blood-derived neutrophils and lung fibroblasts.
A) ARG1 ELISA of lysates of precision-cut IPF lung slices cultured for 24 hours with or without IL-
6R blocking antibody tocilizumab. Data represent N=3 patients, with 3 slices per condition per
patient assayed. ****p<0.0001 by Student's t-test. +/- SEM.
B) 10x Xenium spatial analysis: Left: Representative field of view showing cells expressing ARG1,
EPCAM, and IL6+ fibroblasts (defined by co-expression of either COL1A1 or CTHRC1). Right:
Proximity analysis of Xenium data: Data shown are for the probability ratio, P (exp |
Arg1+)/P(exp), the probability of encountering an IL6+ fibroblast, normalized by fibroblast

frequency, at the radial distance of 25 μm from Arg1+ cells. *p<0.05 by paired Student's t-test.
Data are for N=9 separate patients. +/- SEM.

C) Correlation scatter plot of tissue-resident neutrophils expressing IL-6R and ARG1. Represents
integrated scRNAseq data of non-small cell lung cancer (NSCLC) from 19 datasets (32).

D) COL1A1 immunofluorescence of human lung fibroblasts that were either monocultured, or
cocultured with neutrophils with or without IL-6R blocking antibody (tocilizumab) treatment.
Quantification is for n=3 separate cultures each. Donor 1 and Donor 2 represent separate
donors for both neutrophils and fibroblasts.**p<0.01 by 2-way ANOVA followed by Sidak's
multiple comparisons test. +/- SEM.

E) ARG1 ELISA of conditioned media from cocultures of human blood-derived neutrophils and
fibroblasts from healthy human explanted lung. ***p<0.001, ****p<0.0001 by 2-way ANOVA
followed by Sidak's multiple comparisons test. +/- SEM.

F) Ornithine concentration of conditioned media from the same cocultures as in (E). ****p<0.0001
by 2-way ANOVA followed by Sidak's multiple comparisons test. +/- SEM.

1004

Figure 6: P2rx4 is necessary for IL-6 expression in mouse and human lung fibroblasts.

A) Top Left: UMAP plot of scRNAseq for macrophage-fibroblast cocultures with SingleR-based cell type annotation(2) shown. Data represent n=2 separate cultures for each condition. Top Right: UMAP plot for data showing sample of origin ("Fib" = fibroblast monoculture, "Fib + Mac" = fibroblast coculture with macrophages). Below Left: Feature plot showing *Il6* expression. Below Right: Violin plot of *Il6 expression*. ***p<0.001 by Wilcoxon Rank Sum test corrected for multiple comparisons by Bonferroni method.

B) Gene set enrichment analysis (GSEA) of fibroblast single cell transcriptomes in coculture with macrophages compared to fibroblast monoculture using GO "Response to stimulus" pathways.

C) IL-6 ELISA of conditioned media from macrophage-fibroblast cocultures with or without fibroblast-specific P2rx4 deletion. N=5 five biological replicates per condition. ****p<0.0001 by 1-way ANOVA with post hoc Sidak's multiple comparisons tests. +/- SEM.

D) IL-6 ELISA of conditioned media from monocultured mouse lung fibroblasts from WT mice, with and without ATPγS and SB203580 (p38 Map Kinase inhibitor) treatment or from fibroblast-specific P2rx4 KO (Pdgfrb-Cre: P2rx4 f/f) mice, with ATPγS treatment. N=8, 8, 5, 3 biological

replicates per condition. *p<0.05, ***p<0.001, ****p<0.0001 by 1-way ANOVA with post hoc
Sidak's multiple comparisons tests. +/- SEM.
E) IL-6 ELISA from bronchoalveolar lavage from mice with or without fibroblast-specific P2rx4
deletion. N=7, 7, 6 biological replicates, left to right. **p<0.01, *p<0.05 by 1-way ANOVA with
post hoc Sidak's multiple comparisons tests. +/- SEM.
F) IL-6 ELISA for conditioned media from human donor lung fibroblast monocultures with P2RX4
siRNA KD or scrRNA, +/- ATPγS treatment (*left*, n=3 per condition. ***p<0.001, ****p<0.0001 by
1-way ANOVA with post hoc Sidak's multiple comparisons tests; quantification of KD by qPCR
on *right*, n=3 per condition, ***p<0.001 by Student's t-test). +/- SEM.
G) IL-6 ELISA for conditioned media from human donor lung fibroblast monocultures, human blood
derived neutrophil monocultures, or cocultures with or without P2RX4 inhibition with BAY-1797.
Conditioned media were collected after 24 hours of culture. Each point re[presents a separate
technical replicate. Neutrophils were derived from 2 separate blood donors, and fibroblasts were
derived from 2 separate healthy lung donors. ***p<0.001, ****p<0.0001 by 2-way ANOVA
followed by Sidak's multiple comparisons test. Donor 1 and Donor 2 represent separate donors
for both neutrophils and fibroblasts. +/- SEM.