***Mycobacterium tuberculosis* requires conditionally essential metabolic pathways**

for infection

Alisha M. Block[1], Parker C. Wiegert[1], Sarah B. Namugenyi[1,#a], and Anna D. Tischler[1*]

[1] Department of Microbiology and Immunology, University of Minnesota, Twin Cities

Campus, Minneapolis, MN, USA

[#a] Current Address: Infectious Disease Laboratory, Public Health Laboratory, Minnesota

Department of Health, St. Paul, Minnesota, USA

* Corresponding author

E-mail: tischler@umn.edu (ADT)

Short title: *M. tuberculosis* metabolic pathways essential for infection

ABSTRACT

15
New drugs are needed to shorten and simplify treatment of tuberculosis caused

by *Mycobacterium tuberculosis*. Metabolic pathways that *M. tuberculosis* requires for

growth or survival during infection represent potential targets for anti-tubercular drug

development. Genes and metabolic pathways essential for *M. tuberculosis* growth in

standard laboratory culture conditions have been defined by genome-wide genetic

screens. However, whether *M. tuberculosis* requires these essential genes during

infection has not been comprehensively explored because mutant strains cannot be

generated using standard methods. Here we show that *M. tuberculosis* requires

functional phenylalanine (Phe) and *de novo* purine and thiamine biosynthetic pathways

for mammalian infection. We used a defined collection of *M. tuberculosis* transposon

(Tn) mutants in essential genes, which we generated using a custom nutrient-rich

medium, and transposon sequencing (Tn-seq) to identify multiple central metabolic

pathways required for fitness in a mouse infection model. We confirmed by individual

retesting and complementation that mutations in *pheA* (Phe biosynthesis) or *purF*

(purine and thiamine biosynthesis) cause death of *M. tuberculosis* in the absence of

nutrient supplementation *in vitro* and strong attenuation in infected mice. Our findings

show that Tn-seq with defined Tn mutant pools can be used to identify *M. tuberculosis*

genes required during mouse lung infection. Our results also demonstrate that *M.*

tuberculosis requires Phe and purine/thiamine biosynthesis for survival in the host,

implicating these metabolic pathways as prime targets for the development of new

antibiotics to combat tuberculosis.

AUTHOR SUMMARY

38
Mycobacterium tuberculosis causes more than 10 million new cases of active

tuberculosis (TB) disease and 1.6 million deaths worldwide each year. Individuals with

active TB must take a combination of four antibiotics for a minimum of 6-9 months to

cure the infection. New anti-tubercular drugs are needed to simplify TB treatment and

combat drug resistance. Here, we describe a novel collection of *M. tuberculosis* mutants

lacking metabolic pathways essential for growth in standard laboratory conditions. Using

these mutants, a mouse infection model, and deep sequencing we identified those

metabolic pathways that *M. tuberculosis* also requires during infection. We find that *M.*

tuberculosis mutants that cannot synthesize purine nucleotides, riboflavin, or certain

amino acids are unable to grow in mice. We also find that mutant strains which cannot

synthesize purine nucleotides or the amino acid phenylalanine die rapidly in laboratory

cultures without nutrient supplementation, suggesting that new drugs targeting these

pathways would kill *M. tuberculosis*. Overall, our work reveals multiple metabolic

pathways that *M. tuberculosis* requires during infection, which could be pursued as new

targets for development of anti-tubercular drugs.

INTRODUCTION

Mycobacterium tuberculosis caused more deaths worldwide in 2021 than any

other bacterial infectious agent, in part due to the complexity of tuberculosis (TB)

treatment, which requires 6-9 months of therapy with multiple antibiotics [1, 2].

Multidrug-resistant *M. tuberculosis* (MDR-TB) accounts for ~4.2% of TB infections and

is more challenging to treat, requiring use of less effective second-line agents [1, 3].

Development of new anti-tubercular drugs will be critical to reduce the length of TB

treatment and to combat antibiotic resistance. New drug regimens hold some promise

for reducing TB treatment duration and for treating MDR-TB [4-6]. However, defining

diverse new targets for TB drug development will be necessary to counteract antibiotic

resistance, which has already been documented for all existing drugs and all new or

repurposed TB drugs in the clinical pipeline [7]. Metabolic pathways that *M. tuberculosis*

requires for *in vitro* growth and that are highly vulnerable to inhibition have been

proposed as novel TB drug targets [8-11]. However, these metabolic pathways may not

be required for *M. tuberculosis* growth during infection due to differences in nutrient

availability in host tissues as compared to the *in vitro* growth medium. Defining the

metabolic pathways that *M. tuberculosis* requires during mammalian infection will be

essential to prioritize targets for TB drug development.

M. tuberculosis genes required during mammalian infection have been defined

by genome-wide transposon (Tn) mutant screens in the mouse infection model [12-17].

However, these studies excluded analysis of genes that are essential for *M. tuberculosis*

growth in standard culture conditions *in vitro* [11], as it is not possible to generate

insertional mutations in these genes. In a few cases, auxotrophic *M. tuberculosis* mutant

strains generated on medium supplemented with exogenous nutrients were used to

demonstrate that specific central metabolic pathways are essential for growth in mice.

These include pathways for synthesis of the amino acids Met, Arg, Pro and Trp [13, 18-

21]. Other metabolic pathways such as coenzyme A, biotin, and trehalose synthesis

were shown to be essential during mouse infection using *M. tuberculosis* conditional

gene expression strains [22-24]. Each of these pathways are being pursued as potential

targets for TB drug development.

However, not all metabolic pathways essential for *M. tuberculosis* growth *in vitro*

are also required during mammalian infection. For example, the *nadABC* genes, which

encode enzymes for *de novo* NAD+ synthesis, are essential *in vitro* but dispensable in

the host [25, 26]. *M. tuberculosis* can scavenge the nicotinamide precursor from host

tissue to synthesize NAD+ using an alternative salvage pathway [26]. This example

highlights the importance of determining whether genes that *M. tuberculosis* requires for

in vitro growth are also necessary during infection. Of the 625 genes annotated as

essential for optimal growth of *M. tuberculosis* on standard Middlebrook medium [11],

only a few have been directly tested to determine whether they are also required during

infection because deletion mutant strains lacking these genes cannot be generated

under standard culture conditions.

We previously developed an arrayed Tn mutant library using MtbYM rich

medium, a custom medium that contains many additional carbon and nitrogen sources,

amino acids, nucleotide precursors, co-factors and vitamins as compared to standard

Middlebrook medium [27, 28]. Based on genome-wide transposon-sequencing (Tn-seq)

analysis, 118 genes annotated as essential for optimal growth on Middlebrook 7H10 are

non-essential on MtbYM rich [27]. These include genes required for synthesis of amino

acids (Arg, Trp, Ile, Val, Leu, Glu, Pro, Ser, Tyr, Gln, Phe, Met, Cys), vitamins and co-

factors (riboflavin, folates, pantothenate, NAD), and purine nucleotides, as well as

certain catabolic pathways, including glycolysis [27]. We reasoned that our unique

collection of Tn mutant strains selected on MtbYM rich medium could be exploited to

rapidly assess whether *M. tuberculosis* requires these conditionally essential metabolic

pathways for growth and survival during infection.

Here, we show that *M. tuberculosis* requires functional phenylalanine (Phe)
biosynthesis and *de novo* purine nucleotide and thiamine biosynthesis pathways for
growth in the mammalian host. We screened our collection of Tn mutants in
conditionally essential genes in a mouse infection model using transposon sequencing
(Tn-seq) and identified multiple central metabolic pathways that *M. tuberculosis* requires
for fitness in host lung and spleen tissues. We confirm that mutation of *pheA* (Phe
synthesis) or *purF* (purine/thiamine synthesis) causes death of *M. tuberculosis* in the
absence of exogenous nutrient supplementation *in vitro* and in the lungs of aerosol-
infected mice. Our results demonstrate roles for multiple metabolic pathways in *M.*
tuberculosis fitness during infection and implicate the Phe and purine/thiamine
biosynthetic pathways as novel targets for development of anti-tubercular drugs.
119
RESULTS
M. tuberculosis genes that are essential for growth in standard Middlebrook
medium have been defined by saturating Tn mutagenesis screens [11]. We previously
constructed an arrayed Tn mutant library in the *M. tuberculosis* Erdman strain using
MtbYM rich medium, which contains many additional carbon sources, amino acids,
metabolic intermediates, vitamins and co-factors that are not present in Middlebrook
medium [27, 28]. Our library contains 48 Tn mutants in genes annotated as "essential"
in Middlebrook medium that are non-essential in MtbYM rich medium [11, 27, 28]. We
refer to these genes as Middlebrook-Essential (M-ES) genes. To determine whether *M.*
tuberculosis requires these M-ES genes for survival in the host, we isolated Tn mutants
in M-ES genes from our arrayed library and screened them for fitness defects in mice
using transposon sequencing (Tn-seq).
Tn insertions in genes annotated as essential for growth on Middlebrook medium
can be isolated on MtbYM rich medium.
Our arrayed *M. tuberculosis* Erdman Tn library made on MtbYM rich medium
contains a total of 105 unique Tn insertions in 48 M-ES genes (**S1 Table**). Of these, 20
mutants in 17 unique M-ES genes harbored the Tn insertion near the 5' or 3' end of the
gene (within the first 5% or last 95% of the coding sequence). As these Tn insertions
were unlikely to disrupt gene function, these mutants were excluded from this study

(**S1C Table**, Tn mutants excluded). We recovered 45 of the Tn mutants in M-ES genes

from their mapped locations in our arrayed library on MtbYM rich agar and confirmed

the Tn insertion site by PCR (**S1A and S1B Table**).

We previously determined that some mutants in our MtbYM rich Tn library have

defects in production of the outer membrane lipid phthiocerol dimycocerosate (PDIM)

due to secondary mutations, unlinked to the Tn [28]. The M-ES Tn mutants might be

PDIM deficient due to similar secondary mutations. As *M. tuberculosis* requires PDIM

for virulence [17], we confirmed that the M-ES Tn mutants were PDIM-proficient prior to

screening for fitness defects in mice. We used PCR and Sanger sequencing to test the

M-ES Tn mutants for point mutations in PDIM biosynthesis genes that we identified in

other Tn mutants from our library. This analysis identified 6 M-ES Tn mutants with

mutations in genes required for PDIM production (**S1B Table**, unusable Tn mutants).

We conducted whole genome sequencing on the remaining M-ES Tn mutants, which

identified additional mutations predicted to prevent PDIM production in 17 Tn mutants

and one mutant with multiple Tn insertions (**S1B Table**, unusable Tn mutants). These

mutants were excluded from further analysis. We recovered Tn mutants in 19 M-ES

genes which contained a single Tn insertion and no mutations in genes required for

PDIM production, based on whole genome sequencing (**S1A Table**, Tn mutants in M-

ES library). These M-ES Tn mutants, representing various metabolic pathways and

essential processes (**Table 1**), were selected for analysis of growth *in vitro* and fitness

defects in mice.

Table 1. Tn mutants in M-ES genes in the Tn library screened in mice.

Gene	Annotation	Putative Function
purF	amidophosphoriboysltransferase	*de novo* purine & thiamine biosynthesis
purM	phosphoribosyl-aminoimidazole synthase	*de novo* purine & thiamine synthesis
purQ	phosphoribosyl-formylglycinamidine synthetase I	*de novo* purine & thiamine synthesis
pheA	prephenate dehydratase	Phe biosynthesis
trpG	anthranilate synthase component II	Trp biosynthesis
trpB	tryptophan synthase beta chain	Trp biosynthesis
proC	pyrroline-5-carboxylate reductase	Pro biosynthesis
gltB	ferredoxin-dependent glutamate synthetase	Glu biosynthesis

ilvA	threonine dehydratase	Ile biosynthesis
panC	pantoate-beta-alanine ligase	pantothenate biosynthesis
ribA2	GTP cyclohydrolase II	riboflavin biosynthesis
ribG	bifunctional riboflavin synthesis protein	riboflavin biosynthesis
pabB	*para*-aminobenzoic acid (PABA) synthase component I	PABA biosynthesis
fhaA	conserved protein with forkhead-associated domain	regulation of cell wall synthesis
pstP	serine-threonine phosphatase	regulation of cell wall synthesis
ftsQ	cell division protein	cell division
ERDMAN_3321	possible acyl transferase	unknown function
ERDMAN_2739	Obg GTPase	ribosome assembly
ERDMAN_4254	LigA DNA ligase	DNA damage repair

Tn mutants in central metabolic pathways have growth defects in Middlebrook 7H9 medium.

To determine whether the M-ES genes disrupted by Tn insertion are required for *M. tuberculosis* growth in standard culture conditions, we conducted growth curves in Middlebrook 7H9 medium. Tn mutants in genes encoding various central metabolic pathway enzymes exhibited severe growth defects. These included Tn mutants in genes required for purine and thiamine biosynthesis (*purF*, *purM*, *purQ*; **Fig 1A**), biosynthesis of the amino acids Phe, Trp, Pro, Glu and Ile (*pheA*, *trpB*, *trpG*, *proC*, *gltB*, *ilvA*; **Figs 1B and 1C**), and riboflavin or *para*-amino benozoic acid (PABA) biosynthesis (*ribA2*, *ribG*, *pabB*; **Fig 1D**). The *panC*::Tn mutant was included in our study as a positive control, and as expected it also failed to grow in Middlebrook 7H9 (**Fig 1D**). *M. tuberculosis* requires PanC for synthesis of pantothenate (vitamin B5), an essential precursor in coenzyme A biosynthesis, in the absence of exogenous pantothenate [29]. These data demonstrate that each of these M-ES genes is essential for growth of *M. tuberculosis* Erdman in standard Middlebook medium, consistent with their annotation as essential for growth on Middlebrook 7H10 based on saturating Tn mutagenesis [11].

Tn mutants in several other putative M-ES genes had no growth defects in 7H9 (**Figs 1E and 1F**). Four Tn mutants harbored insertions within non-essential domains of M-ES genes. PstP and FhaA regulate *M. tuberculosis* cell wall synthesis [30, 31] and have domains predicted to be non-essential by saturating Tn mutagenesis [11]. The *pstP*::Tn and *fhaA*::Tn mutants both harbor the Tn insertion in a non-essential region of

the gene and grew normally (**Fig 1E**). *M. tuberculosis* requires the membrane protein

FtsQ to regulate cell division [32]. Our *ftsQ*::Tn mutant harbors the Tn at 185 bp from

the 5' end, in a region predicted to be non-essential [11]. The *ftsQ*::Tn mutant exhibited

no growth defect (**Fig 1E**), suggesting that truncated FtsQ lacking 62 amino acids of the

N-terminal cytosolic domain can sustain normal cell division *in vitro.* Our data are

consistent with the observation that the FtsQ periplasmic domain complements the cell

division defects of an FtsQ-depleted strain [32]. *ERDMAN_3321* (*rv3034*), which

encodes a putative acetyl transferase, is annotated as essential, but Tn insertions near

the 5' end of the gene confer a growth advantage [11]. The *ERDMAN_3321*::Tn mutant

harbors the Tn insertion 151 bp from the 5' end of the gene, within this non-essential

region (**Fig 1F**).

We also isolated two mutants with Tn insertions in *M. tuberculosis* Erdman genes

with annotations suggesting essentiality. *ERDMAN_2739* is annotated as *obg* [33]; in *M.*

tuberculosis H37Rv *obgE* (r*v2440c*) encodes an essential GTPase that controls

ribosome assembly [11, 34, 35]. *ERDMAN_4254* is annotated as a DNA ligase [33], with

possible similarity to the essential DNA ligase *ligA* (*rv3014*) [11]. Whole genome

sequencing revealed that the *ERDMAN_2739*::Tn insertion is within *pe_pgrs43* and the

ERDMAN_4254::Tn insertion is within *espK*, both of which are non-essential genes [11].

Thus, it is not surprising that these mutants did not exhibit growth defects (**Fig 1F**).

These data indicate that a subset of the Tn mutants we isolated do not disrupt functions

essential for *M. tuberculosis* growth in standard Middlebrook medium.

***M. tuberculosis* requires many central metabolic pathways for fitness in mice.**

To identify those M-ES Tn mutants with fitness defects in mice, we created and

screened a library of Tn mutants in M-ES genes (**S1A Table**). We included in the M-ES

Tn library five positive control Tn mutants in genes known to be required for *M.*

tuberculosis fitness in mice: *panC*::Tn [29] and four independent Tn insertions in *phoP*,

which encodes a response regulator essential for virulence [36, 37]. We also included

five negative control Tn mutants in genes known to be dispensable for *M. tuberculosis*

growth and survival in mice: *rpfA* and *rpfE* that encode redundant resuscitation

promoting factors [38], *pstB* and *pstC1* that encode subunits of an alternate Pst

phosphate transporter [39], and *tadA* that encodes a Flp pilin assembly ATPase [40].

Each of these negative control Tn mutants were predicted to be PDIM-proficient based

on whole genome sequencing (**S1A Table**). The Tn mutants were grown individually in

MtbYM rich liquid medium and equivalent amounts of each mutant were combined to

generate the M-ES Tn library (**Fig 2A**). Bacteria in the M-ES Tn library were collected in

triplicate for genomic DNA extraction as a control to confirm equal representation of

each Tn mutant and aliquots of the library were frozen for use in experiments (**Fig 2A**).

To screen the M-ES Tn library for fitness defects in mice, the library was grown in

MtbYM rich liquid, pelleted in triplicate for genomic DNA extraction as an input control

(Liquid control), plated on MtbYM rich agar as a control for recovery on plates (Plate

control), and used to infect C57BL/6J mice by intravenous injection (**Fig 2A**). The

intravenous route was used to ensure adequate representation of all Tn mutants in both

lung and spleen tissues. At days 1, 7, 21 and 42 post-infection, groups of mice (*n*=3)

were euthanized and surviving *M. tuberculosis* CFU were recovered from lung and

spleen homogenates by plating on MtbYM rich agar (**Fig 2A**). We noted high variability

in *M. tuberculosis* CFU recovered from the lungs at days 7 and 42 (**Fig 2B**). For

analysis of Tn mutant fitness by Tn-seq, we selected plates containing at least 10^4 *M.*

tuberculosis CFU for genomic DNA extraction. Since only one mouse each at the day 7

and day 42 time points had sufficient CFU in the lungs for Tn-seq analysis (**Fig 2B**), the

lung samples were analyzed only at days 1 and 21. Sufficient bacteria were recovered

from the spleens at all time points for Tn-seq analysis (**Fig 2B**).

To determine Tn mutant fitness, the triplicate genomic DNA samples from the

library input, liquid input, plate input, and tissue homogenates plated on MtbYM rich

agar (**Fig 2A**) were subjected to Tn-seq analysis and read counts for each Tn insertion

site were tabulated (**S2 Table**). To determine relative Tn mutant fitness, we used

TnseqDiff, which identifies conditionally essential genes between conditions based on

the relative frequencies of Tn-seq sequencing reads at each Tn insertion site [41]. We

previously used this approach to identify Tn mutants with altered fitness in low-

complexity *M. tuberculosis* Tn mutant libraries treated with antibiotics *in vitro* [28].

Complete TnseqDiff analyses are provided in **S3 Table**.

No statistically significant fitness defects were observed in comparisons between

the original M-ES Tn library and the MtbYM rich liquid culture that was used to infect the

mice (**Fig 2C**, **S1A Fig**), suggesting that all Tn mutants grow equally well in MtbYM rich

medium. In TnseqDiff comparisons between the original M-ES Tn library and the input

plated on MtbYM rich agar, several M-ES Tn mutants showed slight, but statistically

significant fitness defects (*purF*::Tn, *purM*::Tn, *purQ*::Tn, *gltB*::Tn, *panC*::Tn; **Fig 2C**,

S1B Fig). These Tn mutants may have growth defects on MtbYM rich agar compared to

other mutants in the M-ES Tn library, which causes their under-representation after

recovery on plates. These data suggest that most M-ES Tn mutants are efficiently

recovered on MtbYM rich agar and that any fitness defects observed following mouse

infection are due to inability to survive within the mouse rather than a competitive

disadvantage on MtbYM rich agar.

To determine relative Tn mutant fitness at day 1 post-infection, we compared Tn

mutant abundance in plated mouse tissues to the input plated on MtbYM agar (D1

lungs/plate or D1 spleen/plate) using TnseqDiff. Three Tn mutants (*gltB*::Tn, *pheA*::Tn,

panC::Tn) were significantly under-represented in the lungs at day 1 post-infection (**Figs**

2C and 2D). Attenuation of *panC*::Tn was expected, as *panC* was essential for *M.*

tuberculosis growth in BALB/c mice [29]. We observed similarly reduced fitness of the

gltB::Tn and *pheA*::Tn mutants in the spleen at day 1 post-infection (**Fig 2C**, **S1C Fig**).

These data suggest that Phe biosynthesis (PheA) and Glu biosynthesis (GltB) are

essential for *M. tuberculosis* survival within host tissues at the earliest stage of infection.

To determine Tn mutant fitness in the lungs and spleens at later times post-

infection, the day 7, day 21 or day 42 outputs were compared to the day 1 input from

the corresponding tissue using TnseqDiff. We identified 14 Tn mutants that exhibited

significantly reduced fitness in the lungs at day 21 post-infection (**Figs 2C and 2E**).

These included Tn mutants in genes encoding enzymes required for purine and

thiamine biosynthesis (*purF*::Tn, *purM*::Tn, *purQ*::Tn), amino acid biosynthesis

(*pheA*::Tn, *trpB*::Tn, *trpG*::Tn, *proC*::Tn), riboflavin synthesis (*ribA2*::Tn, *ribG*::Tn) and

PABA synthesis (*pabB*::Tn), as well as the *panC*::Tn and *phoP*::Tn positive controls.

Attenuation of the *trpB*::Tn and *trpG*::Tn mutants was expected as another gene in the

Trp biosynthesis pathway (*trpE*) was previously implicated in *M. tuberculosis* growth in

C57BL/6 mice [13].

We observed a similar pattern of attenuation in the spleens at day 21. Most Tn

mutants that were attenuated in lungs also exhibited significant fitness defects in the

spleen (**Figs 2C and 2F**). One exception is *pabB*::Tn, which was significantly attenuated

in lungs but not spleens (**Figs 2C, 2E and 2F**). The Tn mutants that exhibited significant

fitness defects in the spleen at day 21 were also significantly attenuated in the spleen at

days 7 and 42 post-infection (**Fig 2C; S1D and S1E Figs**). The *pheA*::Tn mutant was

significantly attenuated in lungs and spleens at all time points, despite significantly

reduced fitness at day 1 (**Fig 2C**). In contrast, the *gltB*::Tn mutant was attenuated in

lungs and spleens, but this did not achieve statistical significance, possibly because the

gltB::Tn mutant was at much lower abundance in the day 1 normalization controls (**Fig**

2C, S1C Fig). Overall, these data suggest that *M. tuberculosis* requires multiple central

metabolic pathways, particularly purine nucleotide and thiamine biosynthesis, riboflavin

biosynthesis, and biosynthesis of specific amino acids (Phe, Trp, Pro, Glu) to replicate

within host tissues.

While most M-ES Tn mutants with *in vitro* growth defects were attenuated in

mice, the *ilvA*::Tn mutant was a notable exception. The *ilvA*::Tn mutant did not replicate

in Middlebrook 7H9 *in vitro* (**Fig 1C**), but exhibited no statistically significant fitness

defects in mouse lungs or spleens at any time point (**Fig 2C**). These data indicate that

M. tuberculosis does not require the *de novo* Ile biosynthesis during mouse infection

and suggest that it can acquire sufficient Ile from the host.

Purine and thiamine auxotrophy is cidal for *M. tuberculosis in vitro* and in mice.

To confirm the results of our Tn-seq screen, we selected two Tn mutants for

individual retesting. We selected *purF*::Tn as a representative mutant with defects in

biosynthesis of purine and thiamine. PurF catalyzes the first committed step of *de novo*

purine nucleotide biosynthesis, which also produces a metabolic precursor required for

the *de novo* synthesis of thiamine (vitamin B1) (**Fig 3A**) [42]. In *M. tuberculosis*, purine

nucleotides can be produced either *de novo* or via a salvage pathway from

hypoxanthine, which is converted to the purine nucleotide precursor inosine

monophosphate (IMP) by hypoxanthine-guanine phosphoribosyltransferase (Hpt) [43].

We confirmed that the *purF*::Tn mutant fails to grow in standard Middlebrook 7H9

medium and that the growth defect can be alleviated with exogenous hypoxanthine and

thiamine (**Fig 3B**), the two components of MtbYM rich predicted to bypass *de novo*

purine biosynthesis [27]. Exogenous hypoxanthine alone partially rescued growth of the

purF::Tn mutant, but full growth restoration only occurred with both nutrients added (**S2**

Fig). The growth defect of the *purF*::Tn mutant in 7H9 was also fully complemented by

purF expressed from a plasmid (pMV-*purF*, **Fig 3B**). The *purF*::Tn mutant lost viability in

unsupplemented 7H9, with viable CFU below the limit of detection (1 CFU/ml) after 21

days of incubation (**Fig 3C**). Viability of the *purF*::Tn mutant was restored by either

exogenous hypoxanthine and thiamine or by complementation with pMV-*purF* (**Fig 3C**).

These data demonstrate that loss of PurF function causes death of *M. tuberculosis* in

the absence of nutrient supplementation.

To determine if *M. tuberculosis* requires PurF during infection, we infected

C57BL/6J mice by aerosol with WT Erdman, *purF*::Tn or the *purF*::Tn pMV-*purF*

complemented strain. All strains were confirmed to produce the PDIM lipid that is

essential for virulence using thin layer chromatography of [14]C-propionate labelled lipid

extracts (**S3 Fig**). The *purF*::Tn mutant established infection in the lungs at day 1 post-

infection, but it failed to replicate, was cleared from the lungs by day 42 post-infection,

and failed to disseminate to the spleen (**Figs 3D and 3E**). Attenuation of the *purF*::Tn

mutant was partially complemented by pMV-*purF*. The *purF*::Tn pMV-*purF* strain

replicated in the lungs and disseminated to spleen, but significantly fewer CFU were

recovered compared to the WT control (**Figs 3D and 3E**). Expression of *purF* from the

pMV-*purF* plasmid may be insufficient to support *M. tuberculosis* growth in host tissues.

Alternatively, the *purF*::Tn insertion may be polar on the 3' *purM* gene, which is also a

M-ES gene [11]. Reduced expression of *purF* or *purM* could prevent normal *M.*

tuberculosis replication in mice, in which demand for purine nucleotides and/or thiamine

may be higher as compared to *in vitro* culture. Overall, our data demonstrate that the Tn

insertion in *purF*, not a secondary mutation, causes attenuation in mice. These data

indicate that *M. tuberculosis* cannot obtain sufficient purine nucleotide precursors and/or

thiamine to support growth in host tissues and implicate the purine and thiamine

biosynthetic pathways as TB drug targets.

***M. tuberculosis* requires Phe biosynthesis for survival *in vitro* and in mice.**

We selected the *pheA*::Tn mutant for retesting as the role of Phe biosynthesis in

M. tuberculosis fitness either *in vitro* or in the host has not previously been tested and

the Phe biosynthesis pathway is absent in humans, making it an attractive drug target

[44]. PheA is a prephenate dehydratase required for both pathways of Phe synthesis

(**Fig 4A**). PheA catalyzes the first step in Phe synthesis from prephenate [45], which is

produced from chorismate by chorismate mutase [44]. Two *M. tuberculosis* proteins

have chorismate mutase activity (Rv1885c and Rv0948c), but Rv1885c is secreted

while Rv0948c is cytosolic [46-48]. Only *rv0948c* is a M-ES gene [11], suggesting that it

functions as the chorismate mutase in Phe synthesis. In plants and some bacteria, Phe

synthesis can also proceed via an L-arogenate intermediate [44]. L-arogenate has been

detected in *M. tuberculosis* [49], and the aminotransferase Rv1178c may synthesize L-

arogenate from chorismate, as this enzyme activity has been demonstrated *in vitro* [50].

PheA is predicted to act as the dehydratase to convert L-arogenate to Phe.

We confirmed that the *pheA*::Tn mutant does not grow in 7H9 medium unless the

medium is supplemented with casamino acids, which contains Phe (**Fig 4B**). The

growth defect of the *pheA*::Tn mutant in 7H9 was also fully complemented with plasmid-

encoded *pheA* (pMV-*pheA*, **Fig 4B**). The *pheA*::Tn mutant also lost viability in 7H9 in

the absence of casamino acid supplementation, indicating that an inability to acquire or

produce Phe is cidal to *M. tuberculosis* (**Fig 4C**). This survival defect was also fully

complemented by pMV-*pheA* (**Fig 4C**). These data demonstrate that blocking Phe

synthesis causes death of *M. tuberculosis* in the absence of exogenous nutrient

supplementation.

To confirm attenuation of the *pheA*::Tn mutant, C57BL/6J mice were infected by

aerosol with WT Erdman, *pheA*::Tn, or the *pheA*::Tn pMV-*pheA* complemented strain.

The *pheA*::Tn and *pheA*::Tn pMV-*pheA* strains both produce the PDIM lipid (**Fig S3**).

Despite using a higher inoculating dose, we observed significantly fewer *pheA*::Tn

bacteria in the lungs at 24 hr post-infection compared to either the WT or complemented

controls (**Fig 4D**). This is consistent with the significant reduction in fitness we observed

for the *pheA*::Tn mutant at day 1 post-infection compared to the plated input control in

our Tn-seq screen (**Figs 2C & 2D**). These data suggest that PheA is required for *M.*

tuberculosis to establish infection in the lungs. The *pheA*::Tn mutant was also incapable

of replicating in mouse lungs, was cleared to below the limit of detection from lung

tissues by 42 days post-infection, and failed to disseminate to the spleen (**Figs 4D &**

4E). The *pheA*::Tn pMV-*pheA* complemented strain replicated in the lungs (**Fig 4D**) and

disseminated to spleen (**Fig 4E**) similarly to the WT control. These data show that

attenuation of the *pheA*::Tn mutant is due to the Tn insertion in *pheA* and not a

secondary mutation. These data also demonstrate that *M. tuberculosis* requires Phe

biosynthesis to survive within the host, suggest that Phe is not available in sufficient

quantities in host tissues to support *M. tuberculosis* growth, and indicate that PheA is a

potential target for development of new TB drugs.

DISCUSSION

Most metabolic pathways that are essential for *M. tuberculosis* growth in

standard *in vitro* culture conditions have not previously been evaluated to determine

their importance during infection because mutant strains lacking these metabolic

functions cannot be generated using standard methods. Here we use a Tn mutant

collection generated using a nutrient-enriched growth medium to demonstrate that

several metabolic pathways essential for growth in standard *in vitro* culture conditions

are also essential in the host using a mouse infection model. Our data show that *M.*

tuberculosis requires *de novo* purine and thiamine biosynthesis as well as Phe

biosynthesis for replication in the lungs and dissemination to the spleen in aerosol-

infected mice. In addition, we show that an inability to synthesize Phe or purine and

thiamine causes death of *M. tuberculosis* in the absence of exogenous nutrients,

suggesting that inhibitors of these biosynthetic pathways would be bacteriocidal.

Therefore, our data implicate the Phe and purine/thiamine biosynthesis pathways as

promising targets for development of novel anti-tubercular drugs.

Phe biosynthesis has been explored as a target for anti-tubercular drug

development because mammals lack *de novo* Phe synthesis, which is expected to limit

the toxicity of compounds targeting this pathway [44]. Our data support PheA as a

potential drug target since our *pheA*::Tn mutant exhibited strong attenuation in the

mouse model, suggesting that *M. tuberculosis* cannot acquire sufficient Phe from the

host. Phe biosynthesis also requires conversion of chorismate to prephenate by

chorismate mutase [44]. Inhibitors of *M. tuberculosis* chorismate mutase have been

identified, but most of these compounds had high minimal inhibitory concentrations

(MICs), suggesting that they have low whole cell permeability [51-53]. Our data suggest

that *M. tuberculosis* can import Phe, since growth of the *pheA*::Tn mutant *in vitro* was

restored by amino acid supplementation. Therefore, it may be possible to design PheA

or chorismate mutase inhibitors with improved permeability properties for use as anti-

tubercular agents.

Purine nucleotide biosynthesis is critical for the virulence of many pathogens and

has also been explored as a drug development target [54]. Although mammals also

synthesize purine nucleotides *de novo*, the enzymes are structurally distinct, suggesting

that specific inhibitors of bacterial enzymes can be designed [42]. Our data demonstrate

that *M. tuberculosis* requires PurF, which catalyzes the first committed step of *de novo*

purine nucleotide and thiamine co-factor biosynthesis, for growth in the host. Our data

suggest that both purine and thiamine biosynthesis are critical functions of PurF, as *in*

vitro growth was fully restored only by adding exogenous thiamine and hypoxanthine.

Attenuation of the *purF*::Tn mutant in mice was only partially reversed by

complementation, despite full restoration of *in vitro* growth in the absence of nutrient

supplementation. This suggests that the demand for purine and/or thiamine

biosynthesis is greater within host tissues and that even partial loss of PurF function can

limit *M. tuberculosis* growth. Collectively, our data suggest that *M. tuberculosis* cannot

acquire sufficient purine nucleotides and/or thiamine from host tissues to support its

growth.

Our data also indicate that *M. tuberculosis* requires other enzymes in the purine

and thiamine biosynthetic pathway for growth in the host, as *purM*::Tn and *purQ*::Tn

mutants were highly attenuated in our Tn-seq screen. Our data are consistent with a

prior report that a *M. tuberculosis purC* mutant was cleared from the tissues of

intravenously-infected mice [55]. Our data are also consistent with the observation that

depletion of GuaB2, which is required for *de novo* synthesis of guanine nucleotides,

prevents *M. tuberculosis* replication in aerosol-infected mice [56]. Loss of either PurC or

GuaB2 function is predicted to prevent production of guanine nucleotides, but not

thiamine. Whether an inability to produce thiamine contributes to the attenuation of the

purF::Tn mutant remains to be determined. In addition, it is unclear whether *M.*

tuberculosis requires *de novo* purine and/or thiamine synthesis only for replication

during acute infection or also for persistence in host tissues during chronic infection.

Our future studies will explore these questions using *M. tuberculosis* strains that

conditionally express enzymes specific to the thiamine or purine biosynthetic pathways

or that are required for both pathways, like PurF. Overall, our data are consistent with

prior studies indicating that *M. tuberculosis* requires *de novo* purine biosynthesis for

growth in the host and support further development of inhibitors targeting this pathway.

Nucleotide synthesis inhibitors, which include approved cancer

chemotherapeutics, have shown efficacy against other bacterial pathogens, but can be

toxic to mammalian cells [54]. Structures of *M. tuberculosis* PurN, PurH, and PurF,

which are involved in *de novo* purine synthesis, revealed structural distinctions from the

equivalent human enzymes, suggesting that specific inhibitors could be designed

against these proteins [57-59]. *Mycobacterium abscessus* PurC is also structurally

distinct from the human ortholog, and PurC inhibitors that prevent *M. tuberculosis*

growth *in vitro* were recently described [60]. An indazole sulfonamide compound that

inhibits GuaB2 was cidal against replicating *M. tuberculosis in vitro* but lacked activity in

infected mice due to either low drug concentration in lung lesions, slower growth of *M.*

tuberculosis in chronically infected mice, or high levels of guanine in host tissues that

antagonize the drug [61]. It may be possible to develop improved drugs targeting other

enzymes in the purine biosynthetic pathway that would disrupt both purine and thiamine

biosynthesis. As *de novo* purine biosynthesis includes 10 enzymatic steps, it represents

a pathway rich in targets for the development of novel anti-tubercular agents.

The *de novo* thiamine biosynthesis pathway is also a promising target for

development of novel antimicrobials given that it is absent in humans, who must obtain

thiamine in the diet [62]. Inhibitors of thiamine biosynthesis with *in vitro* efficacy have

been reported for *Salmonella* and *Pseudomonas* [63, 64]. Inhibitors of *M. tuberculosis*

ThiE, which catalyzes the final step in production of the active co-enzyme thiamine

pyrophosphate, have also been identified, one of which potently inhibited *in vitro* growth

[65]. However, this study did not test whether the compound specifically targeted ThiE

through either selection of resistant mutants or use of hyper-morphic strains with

reduced ThiE expression. Further studies will be needed to determine if *M. tuberculosis*

requires thiamine synthesis during infection and to identify the steps in the biosynthetic

pathway that are most vulnerable to inhibition.

Loss of either PheA or PurF function causes rapid death of *M. tuberculosis* in the

absence of exogenous nutrient supplementation, suggesting that inhibitors targeting

these enzymes would be bacteriocidal. The *pheA*::Tn mutant died *in vitro* at a rate

comparable to a methionine auxotroph (Δ*metA*) without amino acid supplementation

[18]. The *purF*::Tn mutant was sterilized to below the limit of detection after only 21 days

of culture without hypoxanthine and thiamine. This is a faster rate of death in the

absence of nutrient supplementation than any other auxotrophic *M. tuberculosis* strains

that have been characterized, including Arg biosynthesis mutants, which are sterilized

by Arg deprivation [18, 19]. However, the molecular mechanisms that cause death of

either the *pheA*::Tn or *purF*::Tn mutants remain to be determined.

For other auxotrophic *M. tuberculosis* mutants, loss of viability was correlated

with multiple effects on microbial physiology. For example, the Δ*metA* mutant cannot

produce either Met or *S*-adenosylmethionine (SAM), an essential co-factor in one-

carbon metabolism [18]. Similarly, death of Arg biosynthesis mutants deprived of Arg

was correlated with increased production of endogenous reactive oxygen species and

DNA damage, in addition to reduced intracellular Arg [19]. Reduced Phe production in

the *pheA*::Tn mutant may deplete chorismate because chorismate mutase is feedback

inhibited by Phe [66]. Chorismate is also required as a precursor for synthesis of Trp,

menaquinones and the mycobactin siderophore, so its depletion would be expected to

affect multiple processes [47]. Loss of PurF function may be cidal in the absence of

nutrient supplementation due to reduced production of both purine nucleotides, which

are needed for DNA synthesis, RNA synthesis and energy storage [54], and thiamine,

which is needed for activity of multiple enzymes involved in central carbon metabolism

and branched chain amino acid biosynthesis [62]. Alternatively, the *purF*::Tn mutant

may be hypersusceptible to exogenous stress, including oxygen limitation, as previously

reported for a *Mycobacterium smegmatis purF* mutant [67]. The mechanisms by which

these mutations cause death of *M. tuberculosis* will be explored in our future studies.

In addition to the *purF*::Tn and *pheA*::Tn mutants that we selected for follow-up

studies, we identified several other Tn mutants that exhibited strong fitness defects in

mice. These included Tn insertions in genes required for riboflavin synthesis (*ribA2*,

ribG), PABA synthesis (*pabB*), and synthesis of the amino acids Pro and Glu (*proC,*

gltB). Our data suggest that *M. tuberculosis* cannot acquire these nutrients in sufficient

quantities from host tissues and must synthesize them *de novo*. To prioritize these

conditionally essential metabolic pathways as targets for drug development, we will

consider whether the pathway is present in mammalian cells, whether loss of the

pathway is cidal *in vitro* in the absence of nutrient supplementation, and whether the

pathway is vulnerable to inhibition.

Riboflavin, Glu and PABA synthesis pathways are absent from mammalian cells,

so compounds targeting these pathways would be expected to have low toxicity. We

can use our Tn mutants to assess whether starvation for these nutrients is cidal for *M.*

tuberculosis in vitro, as we have done for the *pheA*::Tn and *purF*::Tn mutants. PABA is a

metabolic precursor in folate biosynthesis. A *M. tuberculosis* PabB inhibitor has been

reported to overcome resistance to anti-folate drugs [68], indicating that compounds

targeting PabB can be developed. Vulnerability of *M. tuberculosis* to loss of gene

function *in vitro* has been estimated based upon reduced fitness with transcriptional

inhibition using CRISPR interference (CRISPRi) [9]. A genome-wide CRISPRi screen

revealed that *ribA2* and *gltB* were among the genes most vulnerable to inhibition, with

vulnerability indices in the top 4% of all of *M. tuberculosis* genes [9]. Many other M-ES

genes that we analyzed including *pheA*, *purF* and other genes required for purine

biosynthesis were also highly vulnerable, with vulnerability indices similar to the targets

of existing anti-TB drugs [9]. Our data indicate that riboflavin, Glu and PABA synthesis

are all required for acute *M. tuberculosis* infection and should be considered as potential

targets for development of new anti-tubercular agents. Future studies will address

whether these genes are also essential during chronic *M. tuberculosis* infection using

conditional expression strains.

While our data indicate that *M. tuberculosis* requires most of the M-ES genes that

we analyzed for survival in mice, *ilvA* was an exception. IlvA catalyzes the first step in

de novo synthesis of isoleucine (Ile), a branched chain amino acid (BCAA) [69]. BCAA

biosynthesis has been actively investigated as a target for antimicrobial development

because these pathways are absent in mammals [70]. In addition, *M. tuberculosis ilvA* is

highly vulnerable to inhibition *in vitro* [9]. However, our data suggest that IlvA would be a

poor drug target because the *ilvA*::Tn mutant was not attenuated in mice, despite strong

in vitro growth inhibition. Our data suggest that *M. tuberculosis* can obtain sufficient Ile

from host tissues to support its growth. Our data are consistent with the observation that

a *M. tuberculosis* mutant lacking *ilvB1*, which is required for synthesis of all BCAAs (Ile,

Leu, Val), was not cleared from lung or spleen tissues in intravenously-infected mice,

despite loss of viability *in vitro* without exogenous BCAA [71]. Indeed *M. tuberculosis*

can acquire all BCAAs directly from infected macrophages [72]. Our data highlight the

importance of directly testing whether the biosynthetic pathways essential for *in vitro*

growth are also necessary during infection prior to pursuing these pathways as drug

development targets.

Finally, our work highlights a new strategy for identifying *M. tuberculosis* mutants

that are attenuated during mouse infection using defined pools of Tn mutants and

quantification of Tn mutant fitness with deep sequencing. Most genome-wide screens of

M. tuberculosis Tn mutants in the mouse model used complex Tn mutant pools, which

enabled analysis of Tn mutant fitness only in mouse spleens due to a strict colonization

bottleneck in the lungs [12-14, 16]. Initial Tn mutant screens conducted with signature-

tagged mutagenesis technology used low-complexity Tn mutant pools to identify *M.*

tuberculosis Tn mutants specifically attenuated in the lungs [17, 73, 74]. A few recent

studies have examined fitness of deletion mutants with sequence barcodes and

quantification of mutant fitness in the lungs by sequencing [75, 76]. However, to our

knowledge this is the first study to use low-complexity Tn mutant pools and Tn-seq to

identify *M. tuberculosis* mutants attenuated in mouse lungs. A similar strategy combined

with high-dose aerosol infection could be used to identify *M. tuberculosis* genes that are

required for aerosol transmission and for fitness within lung tissues.

Our work is also the first to determine role of multiple essential central metabolic

pathways in *M. tuberculosis* pathogenesis. Our results directly demonstrate that *M.*

tuberculosis requires both *de novo* Phe synthesis and *de novo* purine/thiamine

synthesis for replication and survival in infected mice, implicating these pathways as

prime targets for development of new TB drugs. Our Tn-seq screen identified IlvA as a

conditionally essential gene that is not required for *M. tuberculosis* survival in the host,

suggesting that BCAA biosynthesis pathways should be de-prioritized for drug

development. Our screen also revealed several other metabolic pathways, including

synthesis of other amino acids (Glu, Pro), PABA and riboflavin that *M. tuberculosis*

requires for growth in host tissues. Overall, our results widen the scope of potential

targets for TB drug development.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *M. tuberculosis* Erdman wild-type and

derivative strains were grown aerobically at 37°C in Middlebrook 7H9 (Difco) liquid

medium supplemented with 10% albumin-dextrose-saline (ADS), 0.5% glycerol, and

0.1% Tween-80 or on Middlebrook 7H10 (Difco) agar supplemented with 10% oleic

acid-albumin-dextrose-catalase (OADC; BD Biosciences) and 0.5% glycerol unless

otherwise noted. Frozen stocks of *M. tuberculosis* strains were made by growing

cultures to late-exponential phase, adding glycerol to 15% final concentration, and

storing at -80°C. Liquid cultures of M-ES Tn mutants or the M-ES Tn library were grown

in MtbYM rich medium (MtbYM) pH 6.6 [27] supplemented with 10% OADC and 0.05%

tyloxapol or on MtbYM agar plates [27], unless otherwise noted. Antimicrobials were

used at the following concentrations: kanamycin (Kan) 25 µg/ml for agar or 15 µg/ml for

liquid, hygromycin (Hyg) 50 µg/ml, cycloheximide 100 µg/ml.

Recovery of Tn mutants from the arrayed Tn mutant library. Tn mutants were

recovered from the mapped location in our arrayed Tn mutant library [28] by streaking

on MtbYM agar containing Kan and incubating at 37°C for at least three weeks. Up to

four individual colonies were picked and grown in 10 ml of MtbYM broth containing Kan

at 37°C with aeration. The Tn insertion site was confirmed by PCR using a gene-specific

primer 5' or 3' of the TA site and a primer specific to the Tn (**Table S4**) followed by

Sanger sequencing of the PCR product. M-ES Tn mutants were screened for secondary

point mutations that disrupt PDIM production known to be present in our Tn library by

PCR and Sanger sequencing. We screened for mutations in *ppsD* and *ppsE* previously

identified in other Tn mutants [28] and for additional common mutations in *ppsB, ppsC*

and *ppsE* identified by whole-genome sequencing of M-ES Tn mutants (**Table S4**).

Whole genome sequencing. The whole genome of each M-ES Tn mutant included in

the M-ES library was sequenced by short-read Illumina sequencing to identify mutations

in the phthiocerol dimycocerosate (PDIM) locus that could result in PDIM deficiency. Tn

mutants were grown to an OD_{600} of 1.0 in MtbYM broth; genomic DNA (gDNA) was

extracted by the CTAB-lysozyme method [77] and cleaned with the Genomic DNA

concentrator and cleanup kit 25 (Zymo). gDNA was submitted to SeqCenter (formerly

Microbial Genome Sequencing Center, MiGS) (Pittsburgh, PA) for library preparation

and Illumina sequencing (151 bp paired-end output, 400 Mbp, 2.67 M reads per

sample). To generate a consensus sequence for each strain, paired reads were mapped

to the *M. tuberculosis* Erdman reference genome (NC_020559.1) using the "map to

reference" function in Geneious 2021 software (Biomatters, Ltd.) as previously

described [28]. To identify single nucleotide polymorphisms (SNPs) in Tn mutant

genomes, the WT Erdman and Tn mutant consensus sequences were aligned in

Geneious using the "Align Whole Genomes" function with the default Mauve Genome

parameters, as described [28]. Tn mutant genomes were only examined for SNPs in the

PDIM locus (*tesA – lppX*). SNPs identified in genes required for PDIM synthesis were

confirmed by PCR amplification and Sanger sequencing (**Table S4**). Tn mutants with

confirmed SNPs in the PDIM locus were excluded from the study. To confirm Tn

insertion sites in the Tn mutants, reads were mapped to the *Himar1* Tn sequence in

Geneious as described above and sequences adjacent to the Tn were compared to the

M. tuberculosis Erdman reference genome. Mutants with more than one Tn insertion

were excluded from the study.

Creation and screening of the M-ES Tn library by Tn-seq. We created our M-ES Tn

mutant library using Tn mutants in genes predicted to be conditionally essential in

Middlebrook 7H9 medium but not MtbYM rich medium (M-ES genes) [11, 27], with a

single Tn insertion and no mutations in the PDIM locus based on whole-genome

sequencing (**Table S1**). For positive controls, we isolated Tn mutants in genes known to

be required for *M. tuberculosis* virulence (*panC*::Tn and *phoP*::Tn) [29, 36, 37]. For

negative controls, we isolated Tn mutants in genes known to be dispensable during

mouse infection (*rpfA*::Tn, *rpfE*::Tn, *tadA*::Tn, *pstB*::Tn, *pstC1*::Tn) [38-40]. Each Tn

mutant was grown individually in MtbYM broth until late-logarithmic phase (OD_{600} ~1.0)

and mixed in equal abundance to make the M-ES Tn library. Frozen stocks of the M-ES

Tn library were made by adding glycerol to 15% final concentration and freezing 1 ml

aliquots. The remaining Tn library culture was aliquoted into three 10 ml cultures and

pelleted by centrifugation (4100 x*g*, 10 min) for gDNA extraction as a total library

(Library) control. For the Tn-seq screen, the Tn library was grown from a frozen stock to

mid-exponential phase (OD_{600} = 0.5) in MtbYM broth, washed once with PBS containing

0.05% Tween-80 (PBS-T), and resuspended at OD_{600} = 0.05 in PBS-T. Three 9 ml

samples of the diluted M-ES Tn library (~10^7 CFU) were pelleted by centrifugation (4100

x*g*, 10 min) for gDNA extraction as an input (Liquid) control. The diluted M-ES Tn library

was serially diluted and plated on MtbYM agar to enumerate total bacterial CFU and to

isolate a plated input (Plate) control.

C57BL/6J mice 7 weeks of age were purchased from Jackson Laboratory, USA.

Mice were injected with 200 μl of the diluted M-ES Tn library (~2 x10^5 CFU) via the

lateral tail vein to deliver ~10,000 CFU to the mouse lungs. Groups of mice (*n*=3) were

euthanized by CO_2 overdose at days 1, 7, 21, and 42 post-infection. Lungs and spleens

were collected, homogenized in PBS-T, serially diluted, and plated on MtbYM agar

containing Kan and cycloheximide to enumerate total bacterial CFU in the sample and

to recover at least 10^4 CFU per plate for Tn-seq analysis. Plates were incubated at 37˚C

with 5% CO_2 until the biomass on the agar was confluent, up to four weeks. Plates with

at least 10^4 CFU for the Plate controls and mouse tissue homogenates were flooded

with 2 ml of GTE buffer [77], and gently scraped with a plastic 10 μl loop to loosen the

biomass. Bacteria were collected by centrifugation (4100 x*g*, 10 min). Genomic DNA

was extracted from all input controls (Library, Liquid, Plate) and from plated mouse lung

and spleen homogenate samples by the CTAB-lysozyme method [77], cleaned with the

Genomic DNA clean and concentrator kit (Zymo), and submitted to the University of

Minnesota Genomics Center (UMGC) for Tn-seq library preparation and Illumina

sequencing.

Transposon sequencing (Tn-seq) and data analysis. Tn-seq was performed as

previously described [27]. *M. tuberculosis* gDNA was fragmented with a Covaris S220

ultrasonicator and a whole genome library was prepared using the TruSeq Nano library

preparation kit (Illumina). Library fragments containing Tn junctions were PCR-amplified

from the whole genome library using the Tn-specific primer Mariner_1R_TnSeq_noMm

and Illumina p7 primer (**Table S4**). The amplified products were uniquely indexed to

allow sample pooling and multiplexed sequencing. A total of 27 Mariner-enriched Tn-seq

libraries were created. All libraries were pooled and sequenced on a NextSeq 2K P1

2x150-bp run (Illumina) with 20% PhiX target spiked in to account for the low diversity of

Tn-seq libraries. Approximately 117 M pass filter reads were generated for the lane. All

expected barcodes were detected and reads were balanced (Mean reads ≈ 2.5 M) with

mean quality scores for all libraries ≥Q30.

Tn-seq analysis was done as previously described [28]. Sequencing reads were

filtered to remove reads without the Tn sequence "GGACTTATCAGCCAACCTGT". The

5' Illumina adaptor sequences were trimmed using BBDuk

(https://sourceforge.net/projects/bbmap/). Each trimmed read was cut to 30 bases and

sequences not starting with TA were removed. Remaining reads were mapped to the *M.*

tuberculosis Erdman genome (NC_020559.1) using HISAT2. Mapped reads were

counted at each TA insertion site to generate read count tables for TnseqDiff analysis.

TnseqDiff normalized the read counts using the default trimmed mean of M values

(TMM) normalization method [78, 79], and then determined conditional essentiality for

each TA insertion site between experimental conditions. TnseqDiff calculated the fold

change and corresponding two-sided *P*-value for each TA insertion site [41]. All *P*-values

were adjusted for multiple testing using the Benjamini-Hochberg procedure in TnseqDiff.

The cut-off values for statistical significance were set at a fold-change of > $\log_2$ ±2 and

an adjusted *P*-value < 0.025.

Tn mutant complementation. Complementation vectors pMV-*purF* and pMV-*pheA*

were in made in the integrating plasmid pMV306hyg [28] by Gibson assembly. Each

gene was PCR-amplified (Phusion; New England Biolabs) along with ~300 bases 5' of

the translation start site to include the native promoter using primers designed with 16-

20 bases complementary to the gene of interest and 18-22 bases that overlap the NcoI

site in pMV306hyg (**Table S4**). PCR products were purified (Qiagen PCR purification kit)

and assembled with pMV306hyg that was linearized by digestion with NcoI using HiFi

DNA Assembly Master Mix (New England Biolabs) according to the manufacturer's

instructions followed by Sanger sequencing of the cloned gene. The *purF*::Tn and

pheA::Tn mutants were electroporated with the corresponding complementation vector

as described [80]. Each Tn mutant was grown in 7H9 with the appropriate supplements

added (60 μM thiamine and 150 μM hypoxanthine for *purF*::Tn or 0.5% casamino acids

for *pheA*::Tn). Transformants were selected on Middlebrook 7H10 agar containing Kan

and Hyg without supplements. The complementing plasmid was confirmed by PCR

(**Table S4**).

Growth curves. Each M-ES Tn mutant in the M-ES Tn library and WT Erdman were

grown individually in MtbYM broth to mid-exponential phase (OD_{600} = 0.4-0.6), pelleted

by centrifugation (2850 x*g*, 10 min) and washed three times with PBS-T before diluting

to OD_{600} = 0.01 in 7H9. Optical density (OD_{600}) was measured every 2-3 days.

To confirm conditional essentiality of *purF*::Tn in 7H9, WT, *purF*::Tn and *purF*::Tn

pMV-*purF* were grown to mid-exponential phase (OD_{600} = 0.4-0.6) in 7H9 supplemented

with 60 μM thiamine and 150 μM hypoxanthine (7H9 +thiam +hypo), pelleted by

centrifugation (2850 x*g*, 10 min) and washed three times with PBS-T before diluting to

OD_{600} = 0.01 in 7H9 or 7H9 +thiam +hypo. The OD_{600} was measured every 2-3 days.

Surviving bacteria were enumerated by serially diluting in PBS-T and plating on

Middlebrook 7H10 agar with 10% OADC and 0.5% glycerol. 7H10 agar was additionally

supplemented with 60 μM thiamine and 150 μM hypoxanthine for recovery of the

purF::Tn mutant. At later time points, bacteria in 1 ml of the *purF*::Tn 7H9 culture were

collected by centrifugation (3000 x*g*, 10 min), resuspended in 100 μl of PBS-T and

plated. Plates were incubated for at least 4 weeks at 37˚C before enumerating CFU.

To confirm conditional essentiality of *pheA*::Tn in 7H9, WT, *pheA*::Tn and

pheA::Tn pMV-*pheA* were grown to mid-exponential phase (OD_{600} = 0.4-0.6) in 7H9

supplemented with 0.5% casamino acids (7H9 +cas), pelleted by centrifugation (2850

x*g*, 10 min) and washed three times with PBS-T before diluting to OD_{600} = 0.01 in 7H9 or

7H9 +cas. The OD_{600} was measured every 2-3 days. Surviving bacteria were

enumerated by serially diluting in PBS-T and plating on Middlebrook 7H10 agar with

10% OADC and 0.5% glycerol. 7H10 agar was additionally supplemented with 0.5%

casamino acids for recovery of the *pheA*::Tn mutant. Plates were incubated for at least

4 weeks at 37˚C before enumerating CFU.

Phthiocerol dimycocerosate (PDIM) labeling and detection. PDIM production was

measured by a radiolabeling and thin layer chromatography (TLC) method [81].

Cultures grown to mid-logarithmic phase in 10 ml of 7H9 broth with appropriate

supplements (7H9 +thiam +hypo for *purF*::Tn, 7H9 +cas for *pheA*::Tn) were labeled for

48 hr with 10 µCi of [1-^{14}C] propionic acid, sodium salt (American Radiolabeled

Chemicals, Inc; specific activity 50-60 mCi/mmol) prior to apolar lipid extraction and

detection of the PDIM lipid by TLC as described [28].

Mouse aerosol infection. For retesting individual Tn mutant and complemented

strains, male and female C57BL/6J mice 6-8 weeks of age were purchased from

Jackson Laboratory, USA, and infected via the aerosol route using an Inhalation

Exposure System (GlasCol) as described [82]. The nebulizer was loaded with a

bacterial suspension in PBS-T at OD_{600} = 0.007 for WT and complemented strains or

OD_{600} = 0.01 for the *purF*::Tn or *pheA*::Tn mutant strains to deliver ~100 CFU to the

lungs. Groups of mice (*n* = 6, 3 female and 3 male) were euthanized by CO_2 overdose

at 1, 7, 21, and 42 days post-infection. Bacterial CFU were enumerated by plating

serially diluted lung and spleen homogenates on 7H10 agar with cycloheximide. The

7H10 agar was supplemented with 60 µM thiamine and 150 µM hypoxanthine for

recovery of the *purF*::Tn mutant or with 0.5% casamino acids for recovery of the

pheA::Tn mutant. Colonies were counted after at least 4 weeks of incubation at 37˚C.

Animal ethics statement. All animal protocols used in this study were reviewed and

approved by the University of Minnesota Institutional Animal Care and Use Committee

(IACUC) under protocol numbers 1912-37660A and 2210-40465A. The University of

Minnesota's NIH Animal Welfare Assurance Number is D16-00288 (A3456-01),

expiration date 04/30/2024. All animal experiments were done in strict accordance with

the recommendations in the Guide for the Care and Use of Laboratory Animals of the

National Institutes of Health [83].

Statistical analysis. A student's unpaired t-test (two-tailed) was used for pairwise

comparisons between WT, mutant and complemented strains. *P* values were calculated

using GraphPad Prism 8.0 (GraphPad Software, Inc). P values < 0.05 were considered

statistically significant.

Custom scripts and code. All custom scripts and R code used for Illumina sequence

read processing are available at https://github.com/bloc0078/umn-tischler-tnseq.

Custom R code used for TnseqDiff analysis is provided in the supplementary material

S1 Files.

Data availability. All raw sequencing data from whole genome sequencing and Tn-seq

experiments underlying the results reported are publicly available in FASTA format at

the NCBI Sequence Read Archive (BioProject PRJNA1006392). All raw numerical and

image data underlying the figures are provided in the supplementary materials.

ACKNOWLEDGEMENTS

This work was completed in part with resources, including instrumentation and

staff, at the University of Minnesota Genomics Center (UMGC; RRIC SCR_012413) and

the University of Minnesota Imaging Centers (UIC: RRID SCR_020997). We thank Drs.

Yusuke Minato and Anthony Baughn for developing the custom MtbYM rich media and

for helpful discussions. We thank Dr. Lili Zhao for providing the updated R code for

TnseqDiff analysis.

REFERENCES

1.   WHO. Global tuberculosis report 2022. Geneva: World Health Organization;
2022.
2.   WHO. WHO consolidated guidlines on tuberculosis. Module 4: treatment - drug-
susceptible tuberculosis treatment. Geneva: World Health Organization; 2022.
3.   WHO. WHO consolidated guidelines on tuberculosis. Module 4: treatment - drug-
resistant tuberculosis treatment, 2022 update. Geneva: World Health Organization;
2022.
4.   Dorman SE, Nahid P, Kurbatova EV, Phillips PPJ, Bryant K, Dooley KE, et al.
Four-month rifapentine regimens with or without moxifloxacin for tuberculosis. N Engl J
Med. 2021;384:1705-18.
5.   Conradie F, Diacon AH, Ngubane N, Howell P, Everitt D, Crook AM, et al.
Treatment of highly drug-resistant pulmonary tuberculosis. N Engl J Med.
2020;382:893-902.
6.   Conradie F, Bagdasaryan TR, Borisov S, Howell P, Mikiashvili L, Ngubane N, et
al. Bedaquiline-Pretomanid-Linezolid regimens for drug-resistant tuberculosis. N Engl J
Med. 2022;387:810-23.

7.   Kadura S, King N, Nakhoul M, Zhu H, Theron G, Köser CU, et al. Systematic
review of mutations associated with resistance to the new and repurposed
Mycobacterium tuberculosis drugs bedaquiline, clofazimine, linezolid, delamanid and
pretomanid. J Antimicrob Chemother. 2020;75:2031-43.
8.   Johnson EO, LaVerriere E, Office E, Stanley M, Meyer E, Kawate T, et al. Large-
scale chemical-genetics yields new *M. tuberculosis* inhibitor classes. Nature.
2019;571:72-8.
9.   Bosch B, DeJesus MA, Poulton NC, Zhang W, Engelhart CA, Zaveri A, et al.
Genome-wide gene expression tuning reveals diverse vulnerabilities of *M. tuberculosis*.
Cell. 2021;184:4579-92.
10.   Tomasi FG, Rubin EJ. Failing upwards: Genetics-based strategies to improve
antibiotic discovery and efficacy in *Mycobacterium tuberculosis*. Front Cell Infect
Microbiol. 2022;12:932556.
11.   DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, et al.
Comprehensive essentiality analysis of the *Mycobacterium tuberculosis* genome via
saturating transposon mutagenesis. mBio. 2017;8:e02133-16.
12.   Sassetti CM, Rubin EJ. Genetic requirements for mycobacterial survival during
infection. Proc Natl Acad Sci USA. 2003;100:12989-94.
13.   Zhang YJ, Reddy MC, Ioerger TR, Rothchild AC, Dartois V, Schuster BM, et al.
Tryptophan biosynthesis protects mycobacteria from CD4 T-cell-mediated killing. Cell.
2013;155:1296-308.
14.   Nambi S, Long JE, Mishra BB, Baker R, Murphy KC, Olive AJ, et al. The
oxidative stress network of *Mycobacterium tuberculosis* reveals coordination between
radical detoxification systems. Cell Host Microbe. 2015;17:829-37.
15.   Payros D, Alonso H, Malaga W, Volle A, Mazeres S, Dejean S, et al. Rv0180c
contributes to *Mycobacterium tuberculosis* cell shape and to infectivity in mice and
macrophages. PLoS Pathog. 2021;17:e1010020.
16.   Smith CM, Baker RE, Proulx MK, Mishra BB, Long JE, Park SW, et al. Host-
pathogen genetic interactions underlie tuberculosis susceptiblity in genetically diverse
mice. eLife. 2022;11:e74419.
17.   Cox JS, Chen B, McNeil M, Jacobs WRJ. Complex lipid determines tissue-
specific replication of *Mycobacterium tuberculosis* in mice. Nature. 1999;402:79-83.
18.   Berney M, Berney-Meyer L, Wong KW, Chen B, Chen M, Kim J, et al. Essential
roles of methionine and S-adenosylmethionine in the autarkic lifestyle of *Mycobacterium
tuberculosis*. Proc Natl Acad Sci USA. 2015;112:10008-13.
19.   Tiwari S, van Tonder AJ, Vilcheze C, Mendes V, Thomas SE, Malek A, et al.
Arginine-deprivation-induced oxidative damage sterlizes *Mycobacterium tuberculosis*.
Proc Natl Acad Sci USA. 2018;115:9779-84. doi: doi: 10.1073/pnas.1808874115.
20.   Smith DA, Parish T, Stoker NG, Bancroft GJ. Characterization of auxotrophic
mutants of *Mycobacterium tuberculosis* and their potential as vaccine candidates. Infect
Immun. 2001;69:1142-50.
21.   Wellington S, Nag PP, Michalska K, Johnston SE, Jedrzejczak RP, Kaushik VK,
et al. A small-molecule allosteric inhibitor of *Mycobacterium tuberculosis* tryptophan
synthase. Nat Chem Biol. 2017;13:943-50.

22.   Evans JC, Trujillo C, Wang Z, Eoh H, Ehrt S, Schnappinger D, et al. Validation of
CoaBC as a bactericidal target in the coenzyme A pathway of *Mycobacterium*
tuberculosis. ACS Infect Dis. 2016;2:958-68.
23.   Park SW, Klotzsche M, Wilson DJ, Boshoff HI, Eoh H, Manjunatha U, et al.
Evaluating the sensitivity of *Mycobacterium tuberculosis* to biotin deprivation using
regulated gene expression. PLoS Pathog. 2011;7:e1002264.
24.   Korte J, Alber M, Trujillo CM, Syson K, Kollwer-Brandl H, Deenen R, et al.
Trehalose-6-phosphate-mediated toxicity determines essentiality of OtsB2 in
Mycobacterium tuberculosis in vitro and in mice. PLoS Pathog. 2016;12:e1006043.
25.   Boshoff HIM, Xu X, Tahlan K, Dowd CS, Pethe K, Camacho LR, et al.
Biosynthesis and recycling of nicotinamide cofactors in *Mycobacterium tuberculosis*: an
essential role for NAD in nonreplicating bacilli. J Biol Chem. 2008;283:19329-41.
26.   Vilcheze C, Weinrick B, Wong KW, Chen B, Jacobs WR, Jr. NAD+ auxotrophy is
bacteriocidal for the tubercle bacilli. Mol Microbiol. 2010;76:365-77.
27.   Minato Y, Gohl DM, Thiede JM, Chacón JM, Harcombe WR, Maruyama F, et al.
Genomewide assessment of *Mycobacterium tuberculosis* conditionally essential
metabolic pathways. mSystems. 2019;4:e00070-19.
28.   Block AM, Namugenyi SB, Palani NP, Brokaw AM, Zhang L, Beckman KB, et al.
Mycobacterium tuberculosis requires the outer membrane lipid phthiocerol
dimycocerosate for starvation-induced antibiotic tolerance. mSystems. 2023;8:e00699-
22.
29.   Sambandamurthy VK, Wang X, Chen B, Russell RG, Derrick S, Collins FM, et al.
A pantothenate auxotroph of *Mycobacterium tuberculosis* is highly attenuated and
protects mice against tuberculosis. Nat Med. 2002;8:1171-4.
30.   Gee CL, Papavinasasundaram KG, Blair SR, Baer CE, Falick AM, King DS, et al.
A phosphorylated pseudokinase complex controls cell wall synthesis in mycobacteria.
Sci Signal. 2012;5:ra7.
31.   Shamma F, Rego EH, Boutte CC. Mycobacterial serine/threonine phosphatase
PstP is phosphoregulated and localized to mediate control of cell wall metabolism. Mol
Microbiol. 2022;118:47-60.
32.   Jain P, Malakar B, Khan MZ, Lochab S, Singh A, Nandicoori VK. Delineating
FtsQ-mediated regulation of cell division in *Mycobacterium tuberculosis*. J Biol Chem.
2018;293:12331-49.
33.   Miyoshi-Akiyama T, Matsumura T, Iwai H, Funatogawa K, Kirikae T. Complete
annotated genome sequence of *Mycobacterium tuberculosis* Erdman. J Bacteriol.
2012;194:2770.
34.   Sasindran SJ, Saikolappan S, Scofield VL, Dhandayuthapani S. Biochemical and
physiological characterization of the GTP-binding protein Obg of *Mycobacterium*
tuberculosis. BMC Microbiol. 2011;11:43.
35.   Kint C, Verstraeten N, Hofkens J, Fauvert M, Michiels J. Bacterial Obg proteins:
GTPases at the nexus of protein and DNA synthesis. Crit Rev Microbiol. 2014;40:207-
24.
36.   Pérez E, Samper S, Bordas Y, Guilhot C, Gicquel B, Martín C. An essential role
for *phoP* in *Mycobacterium tuberculosis* virulence. Mol Microbiol. 2001;41:179-87.

37. Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffé M, Smith I. The
Mycobacterium tuberculosis PhoPR two-component system regulates genes essential
for virulence and complex lipid biosynthesis. Mol Microbiol. 2006;60:312-30.
38. Tufariello JM, Jacobs WR, Jr., Chan J. Individual *Mycobacterium tuberculosis*
resuscitation-promoting factor homologues are dispensable for growth in vitro and in
vivo. Infect Immun. 2004;72:515-26.
39. Tischler AD, Leistikow RL, Ramakrishnan P, Voskuil MI, McKinney JD.
Mycobacterium tuberculosis phosphate uptake system component PstA2 is not required
for gene regulation or virulence. PLoS One. 2016;11:e0161467.
40. Mann KM, Pride AC, Flentie K, Kimmey JM, Weiss LA, Stallings CL. Analysis of
the contribution of MTP and the predicted Flp pilus genes to *Mycobacterium*
tuberculosis pathogenesis. Microbiol. 2016;162:1784-96.
41. Zhao L, Anderson MT, Wu W, Mobley HL, Bachman MA. TnseqDiff: Identification
of conditionally essential genes in transposon sequencing studies. BMC Bioinformatics.
2017;18:326.
42. Zhang Y, Morar M, Ealick SE. Structural biology of the purine biosynthetic
pathway. Cell Mol Life Sci. 2008;65:3699-724.
43. Warner DF, Evans JC, Mizrahi V. Nucleotide metabolism and DNA replication.
Microbiol Spectr. 2014;2:MGM2-0001-2013.
44. Parthsarathy A, Cross PJ, Dobson RCJ, Adams LE, Savka MA, Hudson AO. A
three-ring circus: metabolism of the three proteogenic aromatic amino acids and their
role in the health of plants and animals. Front Mol Biosci. 2018;5:29.
45. Prakash P, Pathak N, Hasnain SE. *pheA* (Rv3838c) of *Mycobacterium*
tuberculosis encodes an allosterically regulated monofunctional prephanate
dehydratase that requires both catalytic and regulatory domains for optimum activity. J
Biol Chem. 2005;280:20666-71.
46. Prakash P, Aruna B, Sardesai AA, Hasnain SE. Purified recombinant hypothetical
protein coded by open reading frame Rv1885c of *Mycobacterium tuberculosis* exhibits a
monofunctional AroQ class of periplasmid chorismate mutase activity. J Biol Chem.
2005;280:19641-8.
47. Sasso S, Ramakrishnan C, Gamper M, Hilvert D, Kast P. Characterization of the
secreted chorismate mutase from the pathogen *Mycobacterium tuberculosis*. FEBS J.
2005;272:375-89.
48. Kim S-K, Reddy SK, Nelson BC, Vasquez GB, Davis AS, Howard AJ, et al.
Biochemical and structural characterization of the secreted chorismate mutase
(Rv1885c) from *Mycobacterium tuberculosis* H37Rv: an *AroQ enzyme not regulated by
the aromatic amino acids. J Bacteriol. 2006;188:8638-48.
49. Lim J, Lee JJ, Lee S-K, Kim SH, Eum S-Y, Eoh H. Phosphoenolpyruvate
depletion mediates both growth arrest and drug tolerance of *Mycobacterium*
tuberculosis in hypoxia. Proc Natl Acad Sci USA. 2021;118:e2105800118.
50. Graindorge M, Giustini C, Kraut A, Moyet L, Curien G, Matringe M. Three
different classes of aminotransferases evolved prephenate aminotransferase
functionality in arogenate-competent microorganisms. J Biol Chem. 2014;289:3198-208.
51. Shukla S, Rao RN, Bhuktar H, Edwin RE, Jamma T, Medishetti R, et al. Wang
resin catalysed sonochemical synthesis of pyrazolo [4,3-*d*] pyrimidinones and 2,3

dihydroquinazolin-4(1*H*)-ones: Identification of chorismate mutase inhibitors having effects on *Mycobacterium tuberculosis* cell viability. Bioorgan Chem. 2023;134:106542.

52.   Alokam R, Jeankumar VU, Sridevi JP, Matikonda SS, Peddi S, Alvala M, et al. Identification and structure-activity relationship study of carvacrol derivatives as *Mycobacterium tuberculosis* chorismate mutase inhibitors. J Enzyme Inhib Med Chem. 2014;29:547-54.

53.   Jeankumar VU, Alokam R, Sridevi JP, Suryadevara P, Matikonda SS, Peddi S, et al. Discovery and structure optimization of a series of isatin derivatives as *Mycobacterium tuberculosis* chorismate mutase inhibitors. Chem Biol Drug Dis. 2014;83:498-506.

54.   Goncheva MI, Chin D, Heinrichs DE. Nucleotide biosynthesis: the base of bacterial pathogenesis. Trends Microbiol. 2022;30:793-804.

55.   Jackson M, Phalen SW, Lagranderie M, Ensergueix D, Chavarot P, Marchal G, et al. Persistence and protective efficacy of a *Mycobacterium tuberculosis* auxotroph vaccine. Infect Immun. 1999;67:2867-73.

56.   Singh V, Donini S, Pacitto A, Sala C, Hartkoorn RC, Dhar N, et al. The inosine monophosphate dehydrogenase, GuaB2, is a vulnerable new bactericidal drug target for tuberculosis. ACS Infect Dis. 2017;3:5-17.

57.   Zhang Z, Caradoc-Davies TT, Dickson JM, Baker EN, Squire CJ. Structures of glycinamide ribonucleotide transformylase (PurN) from *Mycobacterium tuberculosis* reveal a novel dimer with relevance to drug discovery. J Mol Biol. 2009;389:722-33.

58.   Le Nours J, Bulloch EMM, Zhang Z, Greenwood DR, Middleditch MJ, Dickson JMJ, et al. Structural analyses of a purine biosynthetic enzyme from *Mycobacterium tuberculosis* reveal a novel bound nucleotide. J Biol Chem. 2011;286:40706-16.

59.   Chen C, Liu Z, Liu L, Wang J, Jin Q. Structural characterization of glycinamide-RNase-transformylase T from *Mycobacterium tuberculosis*. Emerg Microbes Infect. 2020;9:58-66.

60.   Charoensutthivarakul S, Thomas SE, Curran A, Brown KP, Belardinelli JM, Whitehouse AJ, et al. Development of inhibitors of SAICAR synthetase (PurC) from *Mycobacterium abscessus* using a fragment-based approach. ACS Infect Dis. 2022;8:296-309.

61.   Park Y, Pacitto A, Bayliss T, Cleghorn LAT, Wang Z, Hartman T, et al. Essential but not vulnerable: indazole sulfonamides targeting inosine monophosphate dehydrogenase as potential leads against *Mycobacterium tuberculosis*. ACS Infect Dis. 2017;3:18-33.

62.   Du Q, Wang H, Xie J. Thiamin (vitamin B1) biosynthesis and regulation: a rich source of anti-microbial drug targets? Int J Biol Sci,. 2011;7:41-52.

63.   Zilles JL, Croal LR, Downs DM. Action of the thiamine antagonist bacimethrin on thiamine biosynthesis. J Bacteriol. 2000;182:5606-10.

64.   Kim HJ, Lee H, Lee Y, Choi I, Ko Y, Lee S, et al. The ThiL enzyme is a valid antibacterial target essential for both thiamine biosynthesis and salvage pathways in *Pseudomonas aeruginosa*. J Biol Chem. 2020;295:10081-91.

65.   Khare G, Kar R, Tyagi AK. Identification of inhibitors against *Mycobacterium tuberculosis* thiamin phosphate synthase, an important target for the development of anti-TB drugs. PLoS One. 2011;67:e22441.

66.   Sasso S, Ökvist M, Roderer K, Gamper M, Codoni G, Krengel U, et al. Structure
and function of a complex between chorismate mutase and DAHP synthase: efficiecy
boost for the junior partner. EMBO J. 2009;28:2128-42.

67.   Keer J, Smeulders MJ, Williams HD. A *purF* mutant of *Mycobacterium smegmatis*
has impaired survival during oxygen-starved stationary phase. Microbiol. 2001;147:473-
81.

68.   Thiede JM, Kordus SL, Turman BJ, Buonomo JA, Aldrich CC, Minato Y, et al.
Targeting intracellular *p*-aminobenzoic acid production potentiates the anti-tubercular
action of antifolates. Sci Rep. 2016;6:38083.

69.   Sharma R, Keshari D, Singh KS, Singh SK. Biochemical and functional
characterization of MRA_1571 of *Mycobacterium tuberculosis* H37Ra and effect of its
down-regulation on survival in macrophages. Biochem Biophys Res Commun.
2017;487:892-7.

70.   Amorim Franco TM, Blanchard JS. Bacterial branched-chain amino acid
biosynthesis: structures, mechanisms, and drugability. Biochemistry. 2017;56:5849-65.

71.   Awasthy D, Gaonkar S, Shandil RK, Yadav R, Bharath S, Marcel N, et al.
Inactivtion of the *ilvB1* gene in *Mycobacterium tuberculosis* leads to branched-chain
amino acid auxotrophy and attenuation of virulence in mice. Microbiol. 2009;155:2978-
87.

72.   Borah K, Beyb M, Theorell A, Wu H, Basu P, Mendum TA, et al. Intracellular
Mycobacterium tuberculosis exploits multiple host nitrogen sources during growth in
human macrophages. Cell Rep. 2019;29:3580-91.

73.   Camacho LR, Ensergueix D, Perez E, Gicquel B, Guilhot C. Identification of a
virulence gene cluster of *Mycobacterium tuberculosis* by signature-tagged transposon
mutagenesis. Mol Microbiol. 1999;34:257-67.

74.   Hisert KB, Kirksey MA, Gomez JE, Sousa AO, Cox JS, Jacobs WRJ, et al.
Identification of *Mycobacterium tuberculosis* counterimmune (*cim*) mutants in
immunodeficient mice by differential screening. Infect Immun. 2004;72:5315-21.

75.   Bellerose MM, Proulx MK, Smith CM, Baker RE, Ioerger TR, Sassetti CM.
Distinct bacterial pathways influence the efficacy of antibiotics against *Mycobacterium*
tuberculosis. mSystems. 2020;5:e00396-20.

76.   Koh E-I, Oluoch PO, Ruecker N, Proulx MK, Soni V, Murphy KC, et al. Chemical-
genetic interaction mapping links carbon metabolism and cell wall structure to
tuberculosis drug efficacy. Proc Natl Acad Sci USA. 2022;119:e2201632119.

77.   Larsen MH, Biermann K, Tandberg S, Hsu T, Jacobs WR, Jr. Genetic
manipulation of *Mycobacterium tuberculosis*. Curr Protocols Microbiol. 2007:10A.2.1-
A.2.21.

78.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26:139-40.

79.   Robinson MD, Oshlack A. A scaling normalization method for differential
expression analysis of RNA-seq data. Genome Biol. 2010;11:R25.

80.   Tischler AD, Leistikow RL, Kirksey MA, Voskuil MI, McKinney JD. *Mycobacterium*
tuberculosis requires phosphate-responsive gene regulation to resist host immunity.
Infect Immun. 2013;81:317-28.

81.   Kirksey MA, Tischler AD, Siméone R, Hisert KB, Uplekar S, Guilhot C, et al. Spontaneous phthiocerol dimycocerosate-deficient variants of *Mycobacterium tuberculosis* are susceptible to gamma interferon-mediated immunity. Infect Immun. 2011;79:2829-38.

82.   Ramakrishnan P, Aagesen AM, McKinney JD, Tischler AD. *Mycobacterium tuberculosis* resists stress by regulating PE19 expression. Infect Immun. 2016;84:735-46.

83.   National Research Council. Guide for the care and use of laboratory animals, 8th ed. Washington, DC: National Academies Press; 2011.

FIGURE LEGENDS

Fig 1. *M. tuberculosis* Tn mutants in genes encoding central metabolic enzymes exhibit growth defects in Middlebrook 7H9 medium. Wild-type *M. tuberculosis* Erdman and M-ES Tn mutant strains were grown in MtbYM rich medium, washed twice in PBS-T, and diluted to OD_{600} = 0.01 in Middlebrook 7H9 medium. Growth was monitored by measuring the OD_{600}. Tn mutants are grouped according to predicted function: **(A)** purine and thiamine metabolism (*purF*, *purM*, *purQ*), **(B-C)** amino acid biosynthesis (*pheA*, *trpB*, *trpG*, *proC*, *gltB*, *ilvA*), **(D)** pantothenate, riboflavin, and *p*-amino benzoic acid (PABA) biosynthesis (*panC*, *ribA2*, *ribG*, *pabB*), **(E-F)** Tn insertions within non-essential gene regions (*pstP, fhaA, ftsQ, ERDMAN_3321*) or misannotated genes (*ERDMAN_2739, ERDMAN_4254*). Data represent the mean ± standard error of three biological replicates.

Fig 2. Tn-seq screen of the *M. tuberculosis* M-ES Tn library identifies multiple central metabolic pathways required for fitness in mice. (A) M-ES Tn-seq screen methods. Individual Tn mutants grown in MtbYM rich medium were mixed in equal abundance to create the M-ES Tn library. Triplicate M-ES Tn library samples were collected for Tn-seq (Library control) and the remainder was aliquoted and frozen for experiments. The M-ES Tn library was grown in MtbYM rich broth, then washed and diluted in PBS-T to OD_{600} = 0.05. Triplicate samples of the diluted M-ES Tn library were collected (Liquid control) and the diluted M-ES Tn library was plated on MtbYM rich agar in triplicate (Plate control) for Tn-seq. Mice were injected via the lateral tail vein with ~10^5 CFU of the M-ES Tn library to seed at least 10^4 CFU in the lungs and spleen. Mice (*n*=3) were euthanized at days 1, 7, 21, and 42. Lung and spleen homogenates were plated on MtbYM rich agar to recover surviving Tn mutants for Tn-seq. **(B)** *M.*

tuberculosis CFU recovered from lungs (blue) and spleens (green). **(C)** Heat map of $\log_2$ fold change in Tn mutant abundance between the indicated conditions determined by TnseqDiff analysis of Tn-seq data. Positive values (red) indicate a relative fitness advantage; negative values (blue) indicate a relative fitness defect. **(D-F)** Volcano plots of TnseqDiff analysis of Tn-seq data for the M-ES Tn mutant library for **(D)** lungs at day 1 compared to the Plate control; **(E)** lungs at day 21 compared to lungs at day 1; or **(F)** spleens at day 21 compared to spleens at day 1. Dashed lines indicate cutoffs for statistical significance of ± 2 $\log_2$ fold change and adjusted *P* value <0.025. Tn mutants meeting these significance cutoffs are colored and labeled.

Fig 3. Loss of *purF* causes death of *M. tuberculosis* in the absence of nutrient supplementation *in vitro* and in mice. (A) Pathway for *de novo* synthesis of purine nucleotides and thiamine in *M. tuberculosis*. Solid lines indicate a single step; dashed lines indicate multiple steps. Proteins that catalyze the first five steps are indicated. Intermediate abbreviations: PPP, pentose phosphate pathway; PRPP, 5-phosphoribosyl pyrophosphate; PRA, 5-phospho-D-ribosylamine; GAR, 5'-phosphoribosylglycinamide; FGAR, 5'-phosphoribosyl-N-formylglycinamide; FGAM, 5'-phosphoribosyl N-formylglycinamidine; AIR, 5-aminoimidazole ribotide. **(B-C)** Strains grown in 7H9 supplemented with 60 µM thiamine and 150 µM hypoxanthine were washed in PBS-T and diluted to OD_{600} = 0.01 in 7H9 or 7H9 supplemented with 60 µM thiamine and 150 µM hypoxanthine. Bacterial growth and survival were measured by **(B)** optical density at 600 nm or **(C)** serial dilutions and plating to recover viable CFU. **(D-E)** C57BL/6J mice were infected by the aerosol with the indicated strains. Mice (*n*=6) were euthanized at days 1, 7, 21, and 42 post-infection. Lung **(D)** and spleen **(E)** homogenates were serially diluted and plated to recover viable CFU. In **(C-E)**, WT and *purF*::Tn pMV-*purF* were plated on 7H10 agar; *purF*::Tn was plated on 7H10 agar supplemented with 60 µM thiamine and 150 µM hypoxanthine. Data represent the mean ± standard error of three biological replicates **(B-C)** or six animals **(D-E)**. Asterisks indicate *P*-value < 0.05; n.d. indicates not detected (detection limit = 1 CFU).

Fig 4. *M. tuberculosis* requires Phe synthesis for survival *in vitro* in the absence of amino acid supplementation and in mice. (A) Phenylalanine biosynthesis in *M. tuberculosis*. Single arrows represent one-way reactions. Double arrows represent

reversible reactions. Proteins predicted to catalyze each step are indicated. **(B-C)**

Strains grown in 7H9 supplemented with 0.5% casamino acids were washed in PBS-T

and diluted to OD_{600} = 0.01 in 7H9 or 7H9 supplemented with 0.5% casamino acids.

Bacterial growth and survival were measured by **(B)** optical density at 600 nm and **(C)**

serial dilutions and plating to recover viable CFU. **(D-E)** C57BL/6J mice were infected

by aerosol with the indicated strains. Mice (*n*=6) were euthanized at days 1, 7, 21, and

42 post-infection. Lung **(D)** and spleen **(E)** homogenates were serially diluted and plated

to recover viable CFU. In **(C-E)**, WT and *pheA*::Tn pMV-*pheA* were plated on 7H10

agar; *pheA*::Tn was plated on 7H10 agar with 0.5% casamino acids. Data represent the

mean ± standard error of three biological replicates **(B-C)** or six animals **(D-E)**. Asterisks

indicate *P*-value < 0.05; n.d. indicates not detected (detection limit = 1 CFU).

1070

SUPPORTING INFORMATION

S1 Table. *M. tuberculosis* Tn mutants in M-ES genes recovered from an arrayed

Tn mutant library.

S2 Table. Tn-seq read counts at each Tn insertion site for all experimental

conditions.

S3 Table. Complete TnseqDiff analysis to identify Tn mutants with differential

fitness across experimental conditions. Comparisons between conditions are

indicated in each tab. Bolded text indicates Tn mutants that met statistical significance

cutoffs (log_2 fold change > ± 2, adjusted *P* value <0.025.

S4 Table. Oligonucleotide primers.

S1 Fig. TnseqDiff analyses to identify Tn mutants with fitness defects in *in vitro*

input controls and mouse spleens. Volcano plots of TnseqDiff statistical analyses of

Tn-seq data to determine relative Tn mutant fitness in (**A**) infection input liquid culture

(Liquid) vs. M-ES Tn library control (Library), **(B)** plate recovery of infection culture

(Plate) vs. M-ES Tn library control (Library), **(C)** spleens at day 1 post-infection (D1

Spleen) vs. plated input control (Plate)**, (D)** spleens at day 7 vs. day 1 post-infection,

and **(E)** spleens at day 42 vs. day 1 post-infection. Dashed lines indicate cutoffs for

statistical significance of ± 2 log_2 fold change and adjusted P value of <0.025. Tn

mutants meeting these significance cutoffs are colored and labeled.

S2 Fig. The *M. tuberculosis purF*::Tn mutant requires exogenous hypoxanthine and thiamine for *in vitro* growth. Strains were grown in 7H9 supplemented with 60 μM thiamine and 150 μM hypoxanthine, washed with PBS-T, and diluted to OD_{600} = 0.05 in 7H9 broth (wild-type and *purF*::Tn), 7H9 with 150 μM hypoxanthine (*purF*::Tn +hypoxanthine), or 7H9 with 60 μM thiamine and 150 μM hypoxanthine (*purF*::Tn +hypoxanthine +thiamine). Growth was measured by optical density at 600 nm. Data represent means ± standard errors of biological triplicate cultures.

S3 Fig. *M. tuberculosis* strains tested individually in mice are PDIM-proficient. Thin-layer chromatographic analysis of apolar lipids extracted from cultures of WT Erdman (1) *purF*::Tn (2), *pheA*::Tn (3), *purF*::Tn pMV-*purF* (4), and *pheA*::Tn pMV-*pheA* (5) labelled with [^{14}C]-propionate, which is preferentially incorporated into PDIM.

S1 Data. Supporting data files for all graphical figures. This supporting data file includes the individual numerical values used to generate the graphs in Fig 1, Fig 2B, Fig 2C, Fig 3, Fig 4, and S2 Fig and the results of statistical analyses on these data. Data for each figure are provided in separate tabs. Raw data used for generating the volcano plots in Figs 2D-F and S1 Fig are provided in S3 Table.

S2 Data. Raw images. The raw images used to generate S3 Fig are provided.

S1 Files. **Updated R code for TnseqDiff.**