Title:

Association of secondary metabolite gene clusters with host-specific lineages of the cereal blast fungus *Pyricularia oryzae*

Khyati Mehta[1,$], Jorge C. Navarro-Muñoz[2,#], Sayali Bakore[3], Jérôme Collemare[2,*] and Rajesh Patkar[1,3,*]

[1] Bharat Chattoo Genome Research Centre, Department of Microbiology and Biotechnology Centre, The Maharaja Sayajirao University of Baroda, Vadodara, India

[2] Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands

[3] Department of Biosciences & Bioengineering, Indian Institute of Technology Bombay, Mumbai, India

[$] Current address: Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands

[#] Current address: Wageningen University and Research, Wageningen, the Netherlands

[*] Author for correspondence – Rajesh Patkar - rajeshpatkar@iitb.ac.in and Jérôme Collemare - j.collemare@wi.knaw.nl

Abstract

Fungal plant pathogens constantly evolve and deploy novel peptide and metabolite effectors

to break down plant resistance and adapt to new host plants. The blast fungal pathogen

Pyricularia oryzae is a single species subdivided into multiple host-specific lineages that have

evolved through gain and/or loss of virulence and/or effector related genes through

chromosomal rearrangement. Here, we mined 68 genomes of *P. oryzae*, belonging to six host-

specific lineages, to identify secondary metabolite (SM) biosynthetic gene clusters (BGCs)

likely associated with potential metabolite effectors involved in host specialization. A

similarity network analysis grouped a total of 4501 BGCs into 283 gene cluster families

(GCFs), based on the content and architecture of the BGCs. While most of the GCFs were

present in all the *P. oryzae* lineages, two (BGC-O1 and BGC-O2) were found specifically in

the *Oryza* lineage and one (BGC-TLE) was found in the lineage specific to *Triticum, Lolium*

and *Eleusine* hosts. Further analysis of the phylogenetic relationships between core

biosynthetic genes confirmed that BGC-O1, which comprises a reducing polyketide synthase

gene (MGG_08236) and four putative tailoring genes, was present only in the *Oryza* lineage.

Importantly, most genes, including MGG_08236, from the BGC-O1 were expressed

specifically during pathogenesis. We propose that the *Oryza* lineage-specific BGC-O1

produces a metabolite effector likely involved in specialization of *P. oryzae* to the rice host.

In addition, we identified five SM genes under positive or balancing selection only in the

Oryza lineage, suggesting a role in the interaction with rice specifically. Our findings

highlight the importance of further mining novel metabolite effectors in specialization and

virulence of the blast fungus to different cereal hosts.

Keywords

Biosynthetic diversity, population genomics, *Magnaporthe oryzae,* gene cluster family,

polyketide synthase, effector, polymorphism, divergence

Background

Many plant pathogenic fungi are adapted to infect specific host plants. The emergence of new pathogenic strains, as a result of host range expansion or specialization to a new host, can cause severe disease outbreaks (Giraud et al., 2010). It is crucial to understand the molecular basis of such host specialization to increase the durability of crop resistance to diseases (Giraud et al., 2010; Gladieux et al., 2014). The filamentous fungus *Pyricularia oryzae* (synonym *Magnaporthe oryzae*) is the causative agent of the economically important cereal blast disease worldwide, and is a model organism to study host-pathogen interactions (Ebbole, 2007; Talbot, 2003). Although capable of infecting several grasses, *P. oryzae* is classified as a single species, subdivided into several host-specialized lineages with limited primary host range (Gladieux et al., 2018).

Host selection pressure could be one of the factors that shape the virulence of the population of a pathogen, whereby specific virulence-related genes that encode secreted proteins and/or enzymes for secondary metabolite production - collectively known as effectors – are either gained or lost (Hartmann et al., 2018; Zhong et al., 2016). Effectors manipulate the host physiology or immunity to the pathogen's advantage, and thus are major determinants of specialization on a host (Sánchez-Vallet et al., 2018). Indeed, gain- or loss-of-function mutation in avirulence (AVR) genes of *P. oryzae* has resulted in overcoming host resistance or non-host resistance in case of a new host. For example, a widespread cultivation of wheat cultivar IAC-5 carrying the *RWT3* resistance gene led to a loss of function of the corresponding *PWT3* AVR gene in *P. oryzae* and subsequent emergence of wheat blast (Inoue et al., 2017). Similarly, although gain of the AVR gene *PWL1* led to a loss of pathogenesis on

weeping lovegrass in the *Eleusine*-lineage of *P. oryzae*, it conferred virulence on the new

host, finger millet (Asuke et al., 2020). Further, *AVR-Pita*, flanked by retrotransposons, has

been found to be translocated to multiple genomic locations, including supernumerary

chromosomes, as a result of dynamic adaptation towards its corresponding resistance gene

Pita in certain rice varieties (Chuma et al., 2011).

Fungal secondary metabolites (SMs) have also been shown to exhibit important effector

functions during plant-fungal interactions (Collemare et al., 2019). For example, ACE1, a

secondary metabolite (SM) produced by *P. oryzae*, confers avirulence towards the rice

cultivars carrying the corresponding resistance gene *Pi33* (Böhnert et al., 2004). Various

fungal pathogens, including *P. oryzae*, are also known to produce analogues of plant

hormones such as jasmonic acid, gibberellin and cytokinins, to manipulate plant growth or

subvert plant hormone-based defense signalling pathways during invasion (Patkar & Naqvi,

2017; Shen et al., 2018). A derivative of jasmonic acid (12-hydroxyjasmonic acid) secreted by

P. oryzae suppresses the host innate immunity specifically in rice but not in barley or wheat

(Patkar et al., 2015). Many other fungal SMs such as host-selective toxins (HST) also act as

virulence factors – for example, HC-toxin in *Cochliobous carbonum*, T-toxin in *C.*

heterostrophus, AM-toxin and ACR-toxin in *Alternaria alternata* (Izumi et al., 2012; Lim &

Hooker, 1971; Tsuge et al., 2013; Walton, 2006). These HSTs are essential for pathogenesis

on specific hosts carrying the corresponding susceptible gene. The blast fungus produces SMs

such as tenuazonic acid and pyriculol during pathogenesis (Chen & Qiang, 2017; Jacob et al.,

2017).

The availability of an increasing number of fungal genomes and advances in computational

biology facilitates the exploration of molecular determinants of host specialization or host

range expansion. For instance, population genomics of the wheat fungal pathogen

Zymoseptoria tritici has revealed a possible role of SMs in pathogen lineage diversification

and adaptation to wheat host (Hartmann et al., 2018). In *P. oryzae* isolates, specialization to

rice host is likely associated with a small number of lineage-specific gene families, which

could include genes involved in biosynthesis of SMs and secreted proteins of unknown

functions (Chiapello et al., 2015). While most studies have focused on the effector function of

certain secreted proteins in the blast fungus, the link between diverse SM biosynthetic gene

clusters (BGCs) and *P. oryzae* host-specific lineages has not been explored.

Here, we embarked on the exploration of genomes from the different lineages of the blast

fungus, in search for SM BGCs associated with host specialization. Using similarity network,

comparative genomics and phylogenetic analyses, we identified a candidate BGC that is

likely involved in pathogenesis and specialization to rice and weeping lovegrass host plants.

Results

Genome sequencing of host-specific strains of *P. oryzae*

In this study, we used 68 *P. oryzae* genomes isolated from 6 different host plants, rice (*Oryza*

sativa, n = 26), finger millet *(Eleusine sp.*, n = 14), foxtail millet *(Setaria sp.*, n = 7), wheat

(*Triticum aestivum*, n = 12), perennial ryegrass *(Lolium sp.*, n = 8) and weeping lovegrass

(*Eragrostis curvula*, n = 1) (Fig. 1A, Supplementary Table S1). These genomes included 15

field isolates that were collected from infected rice, finger millet and foxtail millet plants

grown in different parts of India (Fig. 1A, Supplementary Table S1) and sequenced using

Illumina technology (Supplementary Table S2). The remaining 53 genomes were obtained

from publicly available genome sequences (Supplementary Table S1, Fig. 1A). Additionally,

sequence data from three *P. grisea* strains previously isolated from crabgrass (*Digitaria sp.)*

was used as an outgroup (Supplementary Table S1). Sixteen publicly available assemblies

were obtained from long-read sequencing technology and were highly-contiguous. We

performed new gene prediction for all 71 assemblies and analysis of BUSCO genes indicated

high-quality gene prediction and completeness of the assemblies (with minimum 91.5%,

maximum 98.6% and average 96.6% BUSCO-score; Supplementary Table S1).

We used 2655 BUSCO genes present in the 68 *P. oryzae* genomes to build a phylogenomic

tree (Supplementary dataset S4). The topology of the obtained species tree confirms that *P.*

oryzae consists of multiple genetic lineages specialized on different hosts, including *Oryza,*

Setaria, Eleusine, Eragrostis, Triticum and *Lolium* (Fig. 1B). This tree is concordant with the

previously reported population structure (Gladieux et al., 2018). A similar phylogenomic

analysis, using additional three genomes from *P. grisea* (*Digitaria* isolates), showed that *P.*

grisea is a separate species from *P. oryzae*, which is consistent with a previously reported

study (Supplementary Fig. S1; Supplementary dataset S7; Gladieux et al., 2018).

We performed infection assays to check the pathogenesis of 16 Indian isolates of *P. oryzae* on

three hosts – rice, finger millet and foxtail millet. We found that all isolates but one (*P. oryzae*

MOS1 strain) induced clear blast lesions on their host-of-origin, while they triggered no or

few symptoms on non-hosts (Supplementary Fig. S2). In several cases, small dark necrotic

spots were observed, indicative of a hypersensitive response caused by recognition of the

strain (Supplementary Fig. S2). Intriguingly, MOS1 and MOS4, two strains isolated from

infected rice leaf tissue, were found to be genetically similar to that of the *Eleusine* lineage

and not related to any lineage, respectively (Fig. 1B). The whole-plant infection assay using

the MOS1 strain showed differential virulence pattern, where inoculated finger millet and rice

plants developed highly-susceptible and moderately-resistant lesions, respectively, similar to

other isolates from finger millet (Supplementary Fig. S2). This result indicates that MOS1 is

indeed an *Eleusine* lineage strain which was isolated from infected rice tissue. It is possible

that the strain MOS1 could be evolving to adapt to rice host, or it randomly landed on the

collected rice leaf and benefited from a co-infection by an *Oryza* lineage strain. Further

studies are required to understand whether it was associated with any co-infection of the

concerned rice plant tissue with two genetically distinct *P. oryzae* strains. Further, although

genetically not similar to any specific lineage, the MOS4 strain showed similar virulence

pattern – a few susceptible lesions and small necrotic spots - as that by the *Oryza* lineage

strains MOS3 and MOS5 (Supplementary Fig. S2). These results suggest that MOS4 strain

defines a separate lineage that can infect rice, yet it may have another primary host that

remains to be determined.

Genome mining identified lineage-specific biosynthetic gene clusters in *P. oryzae*

The identification of genomic regions that contain SM BGCs was performed in all 71

genomes using antiSMASH (Blin et al., 2021). This analysis resulted in predicting a total of

4224 BGCs, with an average number of 59 BGCs per strain (Supplementary Table S3). The

predicted BGCs were classified based on the core biosynthetic genes, such as genes encoding

polyketide synthases (PKSs), non-ribosomal peptide synthetases (NRPSs) or terpene cyclases

(TCs) (Supplementary Fig. S3, Supplementary Table S3). All the host-specific lineages

contained similar number of BGCs from all different classes of BGCs, with type I PKSs being

the most abundant (Supplementary Fig. S3). Overall, these analyses highlight the great

potential of *P. oryzae* in producing a diverse set of SMs, some of which could be involved in

virulence and/or host specialization.

In order to understand the role of SM genes during divergence of the *Oryza* lineages, we

assessed natural selection by determining the polymorphism within populations and

divergence between the *Oryza*, *Triticum* and *Eleusine* populations in SM core biosynthetic

genes shared by at least ten isolates from each host-specific lineage. We found that most SM

core biosynthetic genes are either under strong or relaxed purifying selection in all three

lineages (Supplementary Table S4). For instance, core genes in the melanin and ACE1

biosynthetic pathways are under strong purifying selection, which is consistent with their key

roles in the interaction of the fungus on any host species (Chumley & Valent, 1990; Fudal et

al., 2007; Vy et al., 2024). Only four SM core enzyme genes (MGG_00806, MGG_01951,

MGG_03401 and MGG_04775) were found to be under balancing selection in the *Oryza*

lineage, and two of these genes were found to be upregulated in appressoria during host

penetration (Oh et al., 2008; unpublished data). Interestingly, one gene MGG_09239 is under

positive selection in the *Oryza* lineage only (Supplementary Table S4). These observations

suggest that only a handful of SM genes likely have a role during interaction with rice

specifically. Especially, genes under balancing selection could be involved in the arms race

with the rice host and represent interesting targets for future functional studies (Brunner &

McDonald, 2018; Oliva et al., 2015; Van Der Hoorn, 2002).

To determine if any putative BGC is associated with the ability to infect specific host plants,

we performed a network similarity analysis using BiG-SCAPE (Navarro-Muñoz et al., 2019).

This analysis included the 4224 predicted BGCs and 277 characterized BGCs from the

MIBiG database (Kautsar et al., 2019) used as a reference. The 4501 BGCs were clustered

into 283 gene cluster families (GCFs)/subnetworks, of which 180 are singletons, including

160 from the reference characterized BGCs (Fig. 2). Twelve out of 91 GCFs present in

Pyricularia strains comprised a characterized BGC, including largely conserved ones for

DHN melanin, epipyriculol, alternapyrone, squalestatin and cytochalasins in *P. oryzae*, and

betaneone in *P. grisea* (Fig. 2; Supplementary Fig. S4-S8). Forty-four GCFs are found in

several *P. oryzae* lineages as well as in *P. grisea*, most of which remain uncharacterized.

Interestingly, 13 and 14 out of 91 GCFs appeared to be specific to *P. grisea* and *P. oryzae*,

respectively (Fig. 2). Of the *P. oryzae*-specific GCFs, three were potentially associated with

host specific lineages. While BGC-O1 and BGC-O2 were found mostly specific to the *Oryza*

lineage, BGC-TLE was found specific to *Tritici*, *Lolium, Eleusine* and *Eragrostis* lineages

(Fig. 2), which all share a common origin and diverged from the *Oryza* and *Setaria* lineages

(Fig. 1). BGC-O1 consists of genes encoding a reducing PKS and tailoring enzymes (see

below), whereas BGC-O2 is restricted to a truncated PKS gene alone that is likely not

functional (Supplementary Fig. S9). In the case of BGC-TLE, although the standalone core

biosynthetic NRPS gene therein is present in all the strains, the gene structure appeared

different (Supplementary Fig. S10). Manual curation of the NRPS gene showed the deletion

of 557 bp in *Triticum* and *Lolium* lineages corresponding to the second exon in the *Eleusine*

lineage (CD156 strain). Further, the presence of a stop codon within the adenylation domain

of the NRPS gene in the *Triticum* and *Lolium* lineages suggests pseudogenization. BGC-TLE

is thus likely not functional in those strains (Supplementary Fig. S11). Yet, it could be either

important for virulence on *Eleusine* host plants only, or its product might have an avirulence

effector-like role in *Triticum* and *Lolium* hosts. Thus, out of three candidate BGCs, we further

investigated BGC-O1 as potentially involved in specialization on rice and weeping lovegrass

hosts.

A novel polyketide biosynthetic gene cluster specifically found in rice/Eragrostis-

infecting lineage of *P. oryzae*

BGC-O1 was found in 23 out of the 24 *Oryza* lineage strains used in the study, as well as in

one *Eragrostis* lineage strain (Fig. 3). This GCF appeared to contain two different networks,

and comparison of the loci indeed clearly differentiated two putative BGCs - one conserved in

all lineages, and BGC-O1 restricted to the *Oryza* and *Eragrostis* lineages (Fig. 3). BGC-O1

contains a novel reducing PKS gene (MGG_08236) as the core biosynthetic gene along with

flanking tailoring genes, including one methyl transferase (MGG_15107), two cytochrome

P450 monooxygenases (MGG_12496 and MGG_12497) and one Co-A transferase

(MGG_15108).

To assess if only the BGC-O1 cluster region was specific to the *Oryza* lineage, we checked

whether flanking genomic regions of BGC-O1 were conserved or variable in different

lineages. We mapped representative high-quality assemblies from each lineage with the

reference genome assembly of the 70-15 strain from *Oryza* lineage to assess synteny at global

level. We found that BGC-O1 was located in the sub-telomeric region of chromosome 2

(NC_017850.1), and approximately 528 kb downstream of the *ACE1* gene cluster in the

reference strain 70-15. The macrosynteny analysis between 70-15 and GUY11, both strains

from *Oryza* lineage, showed conservation of ~134 kb upstream and ~117 kb downstream

flanking regions, along with the BGC-O1 cluster (Fig. 4A). The ~10 kb-long immediate

flanking regions were found to be swapped between the GUY11 and 70-15 genomes (Fig.

4A). Similarly, comparison between 70-15 and FR13 strains revealed a partial conservation

(~14 kb) of BGC-O1, with an ~9.5 kb region from BGC-O1 in 70-15 strain translocated

further downstream on the same contig in the FR13 strain (Fig. 4A). Further, while ~102 kb

upstream flanking region was syntenic in 70-15 and FR13, the downstream flanking region

showed a few structural rearrangements (Fig. 4A). In contrast, comparison with non-*Oryza*

lineages, like *Eleusine* and *Setaria*, showed a significant loss of synteny in the sub-telomeric

region of chromosome 2. While the upstream flanking region (~97 kb upstream of BGC-O1 in

70-15) was found to be syntenic to a corresponding region in MZ5-1-6, CD156 and US71, the

BGC-O1 cluster and downstream sequences were absent in the genomes of these lineages

(Fig. 4B).

Importantly, BGC-O1 did not show any similarity with any of the reference BGCs in MIBiG,

suggesting it to be a BGC, likely involved in production of a novel metabolite.

BGC-O1 likely originates from a duplication event in the common ancestor of *P. oryzae*

and *Colletotrichum eremochloae*

To understand the evolutionary history of BGC-O1, we searched for homologues of the

MGG_08236 PKS protein within *Pezizomycotina* from the MycoCosm repository (Grigoriev

et al., 2014). Only two orthologues could be retrieved from the fungus *Colletotrichum*

eremochloae. A phylogenetic analysis showed that the PKS 670826 (scaffold_178) from *C.*

eremochloae is an orthologue of MGG_08236 PKS, forming an outgroup to the *P. oryzae*

lineage (Fig. 5A). The other paralogue in *C. eremochloae*, 679399 (scaffold_5), belongs to a

clade that is restricted to the *Colletotrichum* genus and forms a sister clade to the

MGG_08236 PKS clade. Comparison of the genomic loci of both homologues in *C.*

eremochloae and BGC-O1 in *P. oryzae* showed that these two clusters shared the PKS and

two cytochrome P450 genes with BGC-O1 (Fig. 5B), while the orthologues of

methyltransferase and Co-A transferase genes were present elsewhere as single copies –

671186 and 576843 – on scaffold_185 and scaffold_10, respectively, in the *C. eremochloae*

genome. The presence of the two similar BGCs in *C. eremochloae* suggests that BGC-O1

likely originated from an ancestral duplication event. While *Pyricularia* and *Colletotrichum*

diverged ~250 MYA (calculated using TimeTree; Kumar et al., 2022), BGC-O1 is retained

only in *P. oryzae* and *C. eremochloae*, suggesting a role of respective host selection pressure

in its retention.

BGC-O1 genes are expressed specifically during host invasion

The presence of the PKS MGG_08236 gene was checked by PCR using genomic DNA of *P.*

oryzae strains from rice and finger millet host plants (Fig. 6A). The MGG_08236 gene was

present in all the *Oryza* strains studied, except for MOS3, which was identified as the only

Oryza lineage strain lacking the BGC-O1 based on the *in-silico* analysis. The MGG_08236

ORF was absent in all the *Eleusine* strains used as well as in the MOS1 and MOS4 strains,

which are placed outside of *Oryza* lineage in the phylogenetic analysis (Fig. 1B and 6A).

We then studied the expression of the PKS (MGG_08236) and three predicted tailoring genes

– methyl transferase (MGG_15107), CoA-transferase (MGG_15108) and cytochrome P450

(MGG_12496) of the BGC-O1 locus during different stages of infection, using RT-PCR.

Total RNA was isolated from fungal vegetative mycelia grown in complete medium, rice

leaves inoculated with *P. oryzae* and incubated for different time-points, and uninoculated

rice leaves, to assess the levels of transcripts of the aforementioned genes relative to those of

β-Tubulin as an internal control. While expression of the core PKS gene was undetected in

mycelium and at 12 hpi (pre-invasive stage), its transcripts started accumulating during

pathogenic development, with significantly elevated expression from 24 hpi till 96 hpi (Fig.

6B). The expression pattern of MGG_15107 and MGG_15108 tailoring genes partly

correlated with that of the PKS, (Fig. 6B). In contrast, MGG_12496 showed a different

expression pattern, where the transcript levels were similar during both vegetative growth and

host invasion (Fig. 6B), suggesting it is not co-regulated with other genes in the predicted

BGC-O1. This result shows that the MGG_08236 PKS gene is specifically expressed during

pathogenesis on rice and potentially has a key role to play during host colonization.

Altogether, our *in-silico* analyses identified a novel PKS gene cluster in the *Oryza*- and

Eragrostis-specific lineages, which likely played a key role in shaping specialization of the

blast fungus to rice and weeping lovegrass host plants.

Discussion

Population genomics identifies BGCs with potential roles in virulence and host

specialization in *P. oryzae*

We identified three lineage-specific BGCs in the blast fungal pathogen *P. oryzae* using a

population genomics approach. While two of them appeared to encode non-functional

enzymes, the *Oryza* and *Eragrostis*-specific BGC, named BGC-O1, appears functional and is

present in almost all the genomes from *Oryza* (23 out of 24) and *Eragrostis* (1 out of 1) lineages studied here. Our population-genomics-based finding of the *Oryza* lineage-specific BGC (BGC-O1) is supported by gene expression analyses. Indeed, we found that the *Oryza*-lineage-specific PKS (MGG_08236) transcript accumulated specifically during penetration and colonization of the rice host. Recent transcriptomics-based studies have shown that the expression of the PKS encoded in BGC-O1 (MGG_08236) and neighboring tailoring genes (methyl transferase, MGG_15107 and Co-A transferase, MGG_15108) was induced during the initial biotrophic colonization stage of blast disease (Jeon et al., 2020; Yan et al., 2023). Importantly, the expression of MGG_08236, along with the other pathogenesis-related genes such as the SM gene cluster *ACE1* and protein effectors SLP2, BAS2, BAS3 and AVR-Pi9, was recently found to be under the control of the bZIP transcription factor Bip1 (Lambou et al., 2024). Bip1 is found to be regulating expression of a specific set of genes involved in early biotrophic stage of blast infection (Lambou et al., 2024). These findings strongly suggest that the SM produced by BGC-O1 could exhibit an effector-like function. Many plant fungal pathogens such as *Colletotrichum higginsianum, Fusarium graminearum, Zymoseptoria tritici*, and *P. oryzae* are known to produce a number of SMs during pathogenesis, especially during host penetration and colonization (Collemare, Billard, et al., 2008; Dallery et al., 2017; Harris et al., 2016; Jeon et al., 2020; Palma-Guerrero et al., 2017; Patkar et al., 2015; Rudd et al., 2015; Yan et al., 2023). To study if the BGC-O1 has any role in virulence in a host-specific manner, deletion of the *PKS* gene (MGG_08236) was attempted in an *Oryza*-specific isolate. Despite using different approaches and a number of attempts, generation of a deletion mutant has been unsuccessful so far, possibly due to its presence in the sub-telomeric region of chromosome 2 of the reference strain 70-15. A similar difficulty was faced in generating deletion mutant at the *ACE1* locus (Collemare, Pianfetti, et al., 2008).

Nonetheless, the MOS3 strain from *Oryza* lineage, which lacks the BGC-O1, failed to infect

rice host as effectively as B157, MOS2 or MOS6 (Fig. 6A; Supplementary Fig. S2A). This

could possibly explain the key role of BGC-O1 in specialization to rice host.

While only three BGCs were present in host-specific lineages, most of the other BGCs were

found in all lineages. However, such apparently conserved BGCs can show variation in their

tailoring gene content. Such differences could result in the production of distinct SMs with

potential effector function in a host specific manner. Alternatively, conservation of BGCs in

different lineages could indicate a key role in virulence on cereals. Thus, analyses

investigating variations among conserved SM-producing BGCs across different *P. oryzae*

lineages could be useful in identifying the likely diversity in the SMs produced and their

association with specific hosts.

Conservation of BGC-O1 in an unstable genomic context

We found that the *Oryza*-specific cluster BGC-O1 is located in the sub-telomeric region on

the chromosome 2 (NC_017850.1) of the reference strain 70-15. While the genomic regions

covering the BGC-O1 and its flanking sequences were highly syntenic in strains from the

Oryza lineage, only the upstream flanking region of this locus was found conserved in all the

other lineages. The chromosome ends tend to have substantially higher levels of

polymorphism because they frequently undergo rearrangements due to the presence of AT-

rich repeats and transposable elements (Farman & Kim, 2005; Rahnama et al., 2020; Starnes

et al., 2012). In the absence of sexual reproduction, asexually propagating pathogens likely

utilize chromosomal rearrangements as one of the mechanisms of host-specialization as

shown in *Verticillium dahliae* (Jonge et al., 2013). Similarly, the regions adjacent to a

telomere in *P. oryzae* are reported to be highly polymorphic and enriched with genes involved

in interaction with its host plants (Rahnama et al., 2021). Gain and/or loss of effector genes, lineage-specific gene families and chromosomal rearrangements are likely the major evolutionary mechanisms involved in such host specificity and adaptation of *P. oryzae* (Chiapello et al., 2015; Chuma et al., 2011; Gómez Luciano et al., 2019; Jonge et al., 2013; Yoshida et al., 2016). Given that BGC-O1 is also flanked by transposons in most strains of the *Oryza* lineage of *P. oryzae* (Fig. 3), we hypothesize that genomic alterations, most likely through deletions, may have resulted in the loss of the BGC in other host-specialized lineages. We further found that BGC-O1 is located on a mini-chromosome (UEMA03000009.1) in the FR13 isolate, which also carries the *ACE1* BGC. Large-scale genomic rearrangements around the loci of several virulence-related genes (*ACE1* and *AVR-Pik*) likely led to the emergence of supernumerary mini-chromosome in the rice isolate FR-13, facilitating the adaptive evolution of the blast fungus (Langner et al., 2021). Thus, retention of BGC-O1 in *Oryza* and *Eragrostis* lineages, despite its localization in unstable sub-telomeric regions, suggests a significant role of host selection pressure in its conservation therein. The most likely selection pressure to retain BGC-O1 relates to the ability to successfully infect a host, which is also indicated by the specific expression profile during infection. The presence of homologous BGCs in *C. eremochloae*, the causal agent of anthracnose disease on centipede-grass turf in southern United States (Crouch & Tomaso-Peterson, 2012), suggests a particular function for BGC-O1 in infecting certain cereals only (*Oryza*, *Eragrostis*, *Eremochloa*). Further investigations are needed to determine if the *C. eremochloae* BGC is expressed during infection and whether it produces the same metabolite as BGC-O1 in *P. oryzae*.

Conclusions

Our findings highlight the importance of population genomics-based studies in identifying

secondary metabolite BGCs and the corresponding genomic rearrangements, likely driven by

the host selection pressure, which could lead to a host-specialized lineage of the blast fungal

pathogen.

Methods

Fungal strains and culture conditions

We isolated 15 field strains of *P. oryzae* from different geographic locations in India and from

three different host plants – rice (*Oryza sativa*), finger millet *(Eleusine coracana)* and foxtail

millet (*Setaria italica*). Infected tissues with blast lesions were collected and used for single

spore isolation. Leaf tissues with blast disease lesions were surface sterilized using 70%

ethanol and 1% sodium hypochlorite, followed by three successive washes with sterile

distilled water. Surface-sterilized leaf tissues were incubated under micro-humid conditions

maintained in a Petri dish for 24-48 h to induce sporulation. Spores were suspended in sterile

water and spread evenly on 2% water agar plate. Single germinating spores were picked up

under a microscope, and inoculated on Prune Agar (PA: 1 g/L yeast extract, 2.5 g/L lactose,

2.5 g/L sucrose, 0.04% prune juice, pH 6.5; Soundararajan et al., 2004) plates and incubated

at 28 °C. Vegetative growth of the fungus on PA plates was continued for 10 days at 28 °C,

with initial 3-day incubation under dark conditions, followed by 7-day incubation under

constant illumination for conidiation. Genomic DNA was extracted by grinding in liquid

nitrogen the fungal biomass obtained from vegetative culture grown in a complete medium

(CM: 6 g/L yeast extract, 6 g/L casein enzyme hydrolysate, 10 g/L sucrose) for ~3 days at 28

°C, followed by the standard protocols as described earlier (Dellaporta et al., 1983).

393

Genome sequencing and assemblies

Newly isolated strains were sequenced for whole genome using a paired-end sequencing

approach at >50X depth on Illumina HiSeq2500 platform at AgriGenome Labs Pvt. Ltd.,

India. Short raw reads were processed using Trimmomatic v0.38 at a threshold for the

minimum read length of 80 bp for paired-end reads (parameters PE, Leading:10, Trailing:10,

Slidingwindow:4:20, Minlen:80; Bolger et al., 2014). *De novo* sequence assemblies were

constructed using CLC Genomics Workbench v11.0 with default parameters (Word size:23,

Bubble Size:50, Minimum contig length:500, Mismatch cost:2, Insertion cost:3, Deletion

cost:3, Length fraction:0.5, Similarity fraction:0.8). Additionally, we used publicly available

genome sequences of *P. oryzae* strains from NCBI (Supplementary table S1). The overall

quality of each genome assembly was evaluated using BUSCO v5.2.2 (Simão et al., 2015)

with the Sordariomycetes dataset. We carried out gene predictions on all the assemblies using

Augustus v3.3.3 (Stanke et al., 2006) with Magnaporthe_grisea as a species model.

Construction of species tree

2655 BUSCO proteins conserved in 68 *P. oryzae* strains were retrieved from the BUSCO

analysis and aligned using Mafft v7.475 (parameters -reorder; Katoh et al., 2002). TrimAL

v1.4.1 (parameters -automated1; Capella-Gutiérrez et al., 2009) was used to remove poorly

aligned regions (Supplementary dataset S1, S2). A maximum likelihood tree was generated

using IQ-TREE v2.1.2 (Minh et al., 2020) with partition models (Chernomor et al., 2016;

Lanfear et al., 2014), model finder (Kalyaanamoorthy et al., 2017) and ultrafast bootstrap

(Hoang et al., 2018; parameters -m MF -p partition.nex -bb 1000) (Supplementary dataset S3,

S4). Similarly, a species tree including 3 *Pyricularia grisea* and 68 *P. oryzae* strains was

generated using a total of 2557 BUSCO proteins (Supplementary dataset S5-S7). Both trees

were visualized and annotated using iTOL (Letunic & Bork, 2021).

Prediction of BGCs and similarity network analysis

BGCs were predicted using antiSMASH v6.0.1 (Blin et al., 2021; parameters -taxon fungi -

minimal -cluster_hmmer -pfam2go). All the predicted BGCs and reference fungal BGCs from

the MIBiG v2.1 (Kautsar et al., 2019) repository were included to perform a similarity

network analysis using BiG-SCAPE v1.1.0 (Navarro-Muñoz et al., 2019) with glocal mode

and with different cut-off values (parameters –include_singletons –mix –no_classify –clans-

off –cutoffs 0.3 0.4 0.5 0.7). A cut-off value of 0.5 was found to be appropriate, as the

relevant BGCs showed similarity with the reference BGCs from MIBiG known for

metabolites such as DHN (melanin), epipyriculol, alternapyrone, squalestatin S1 and

cytochalasans. The network files were visualized with Cytoscape v3.8.2 (Shannon et al.,

2003). Visualization of BGC conservation was done with Clinker (Gilchrist & Chooi, 2021).

Gene models belonging to BGC-O1 family were manually curated (Supplementary dataset

S8).

Statistics for natural selection

To test for natural selection, 40 SM core biosynthetic genes shared by at least ten isolates

from each host-specific lineage (*Oryza*, *Eleusine* and *Triticum*) were used to calculate

polymorphism and divergence (Supplementary Table S4). For each gene, the nucleotide

coding region in the different strains were aligned using Mafft v7.475 (parameters -reorder;

Katoh et al., 2002). DnaSP v6 (Rozas et al., 2017) was used to calculate the levels of

polymorphism within population (Pi(a)/Pi(s)) and divergence between the populations

(Ka/Ks).

Synteny analysis

To identify the syntenic genomic regions between any genome assembly and the reference

70-15 strain assembly, global alignments were generated using nucmer (parameters –c 100 --

maxmatch) program from MUMMER v3.23 package (Kurtz et al., 2004). Alignments were

further filtered using delta-filter utility with length >10 kb and percent identity >80%

(parameters -l 10000 -i 80) to retrieve continuous syntenic blocks. The show-coords utility

was used to extract alignment coordinates in tabular format and the plots were generated in R

using KaryoploteR package (Gel & Serra, 2017).

Core gene phylogeny

Homologs of MGG_08236 were identified in *Pezizomycotina* from the Joint Genome Institute

MycoCosm repository (Grigoriev et al., 2014) using BLASTp. The resulting Blast hits were

filtered for top 3 hits of each species. Amino acid sequences were aligned using MAFFT

v7.475 (Katoh et al., 2002) and poorly aligned regions were removed with TrimAl v1.4.1

(Capella-Gutiérrez et al., 2009). A phylogenetic tree was constructed using Fasttree v2.1.11

(parameters -lg; Price et al., 2010) to identify orthologues and closest paralogues. The

identified orthologous protein sequences were used to build a phylogenetic tree using IQ-

TREE v2.1.2 (Minh et al., 2020) with LG substitution model and ultrafast bootstrap (Hoang et

al., 2018), and the SH-like approximate likelihood-ratio test (Guindon et al., 2010) for branch

support (parameters -mset LG -bb 1000 -alrt 1000) (Supplementary dataset S9-S11). The tree

was visualized and annotated using iTOL (Letunic & Bork, 2021).

Whole-plant infection assay

Conidia were harvested as per the protocol described earlier (Patkar et al., 2012) from

different strains collected in India. Whole-plant infection assay was carried out by spraying 10

mL spore suspension ($\sim 10^5$ conidia/mL) containing 0.05% w/v gelatin onto ~ 4-week-old

host plants, rice cv. CO-39 and finger millet cv. GN-4, and ~3-week-old foxtail millet cv.

SIA-3088 plants, followed by incubation at 28–30 °C under humid conditions, initially for 24

h in the dark and then for 4–6 days under 14 h light and 10 h dark cycles. Development of

disease symptoms was monitored regularly during the entire incubation period and recorded

around 5-8 days post inoculation depending on the appearance of lesions on the leaves. The

infection assays were carried out with at least three biological replicates.

Gene expression analysis by semi-quantitative RT-PCR

The expression profiles of four genes (MGG_08236, MGG_15107, MGG_15108 and

MGG_12496) were studied during different stages of infection on ~ 4-week-old rice plants

(blast susceptible variety PB-1509). Briefly, 20 μL conidial suspension ($\sim 10^5$ conidia/mL

with 0.01% Tween-20) of B157 strain from *Oryza* lineage was drop-inoculated on to the

surface-sterilized detached leaf blades placed on water agar plates containing 2 μg/mL

Kinetin, followed by incubation under dark (10 h) and light (14 h) cycles at 25 °C. Samples

were collected by excising inoculated portions on the leaf blades, at 12-, 24-, 48-, 72- and 96-

hours post inoculation (hpi). Three-day-old vegetative mycelia, grown in liquid CM, was used

as a control condition. A sample from mock-inoculation (leaf blades inoculated with only

0.01% Tween-20) was used as another experimental control to check any non-specific

amplification during RT-PCR. Total RNA was extracted from all samples using TRIzol®

reagent (Invitrogen, USA) as per the manufacturer's instructions. Two μg of total RNA each

was used for cDNA synthesis using iScript™ cDNA Synthesis Kit (Bio-Rad, USA).

Oligonucleotide primers used for RT-PCR for each gene are listed in Supplementary Table

S5. Optimized thermal cycling conditions for RT-PCR were as follows: initial denaturation

step at 95 °C for 5 min, followed by 30 cycling reactions each at 95 °C (denaturation step) for

30 s, 60 °C (annealing step) for 30 s and 72 °C (extension step); and the final extension step at

72 °C for 5 min. The *MGG_08236* ORF was amplified from the genomic DNA of *P. oryzae*

strains from different lineages using the same thermal cycling conditions as above, except for

the extension step, which was set at 1 min in each cycle.

Abbreviations

ACE1	Avirulene Conferring Enzyme1
AVR	Avirulence
BGC	Biosynthetic Gene Cluster
BGC-O1	Biosynthetic Gene Cluster specific to *Oryza* lineage - 1
BGC-O2	Biosynthetic Gene Cluster specific to *Oryza* lineage - 2
BGC-TLE	Biosynthetic Gene Cluster specific to *Triticum*, *Lolium* and *Eleusine* lineages
BiG-SCAPE	Biosynthetic Gene Similarity Clustering and Prospecting Engine
BUSCO	Benchmarking Universal Single-Copy Orthologs
CM	Complete Media
GCF	Gene Cluster Family
hpi	hours post inoculation
HST	Host Selective Toxin
MIBiG	Minimum Information about a Biosynthetic Gene cluster
NCBI	National Centre for Biotechnology Information
NRPS	NonRibosomal Peptide Synthetase
ORF	Open Reading Frame
PA	Prune Agar
PCR	Polymerase Chain Reaction
PKS	PolyKetide Synthase
RT-PCR	Reverse Transcriptase Polymerase Chain Reaction
SM	Secondary Metabolites
TC	Terpene Cyclases

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Raw sequence data for the newly sequenced Indian strains under this study have been made

accessible at SRA database under the BioProject ID PRJNA973860. All datasets generated

for this study are included in the article/Supplementary Material. All the Supplementary

Datasets can be accessed through a link:

https://drive.google.com/drive/folders/18Uw6663e87ZyXRDEO-

2aJpcoNuB7padZ?usp=share_link

Competing interests

The authors declare that they have no competing interests.

Funding

Prof. Bharat Chattoo acquired the grant (BT/PR6949/AGIII/103/862/2012) from DBT, GoI.

KM was supported by the DBT (GoI)-funded grant (BT/PR6949/AGIII/103/862/2012) and

partly by the intramural University Research Scholarship from The M. S. University of

Baroda. KM received EMBO - Scientific Exchange Grant (SEG no. 8866) to conduct part of

this work at the Westerdijk Institute, The Netherlands. RP was awarded by SERB, GoI

(EMR/2017/005303) and the Seed Grant, IIT Bombay.

Authors' contributions

KM collected samples and contributed to the isolation of the Indian strains of blast fungus and

phenotyping experiments. KM conducted formal analyses on diversity in BGCs, JC and JCN-

M helped with the investigation of the analyses. KM, RP and JC conducted image analyses

and statistical analyses. JCN-M provided important in-house scripts for similarity network

analysis. SB helped KM with the gene expression analysis. KM wrote the original draft of the

manuscript. All authors contributed to the review and editing of the manuscript. KM, RP and

JC designed the study. RP provided the resources for the study. RP and JC supervised the

study.

Acknowledgements

We acknowledge late Prof. Bharat B. Chattoo for his guidance and providing sophisticated

laboratory facility at the Bharat Chattoo Genome Research Centre (BCGRC). We also fondly

remember late Dr. Johannes Manjrekar for the useful scientific discussions during this work.

We thank Dr. Malali Gowda and Dr. T. R. Sharma for providing blast-infected tissues for

some of the isolates used in this study. We thank the BCGRC group at MSU for useful

discussions. We thank Dr. Noppol Kobmoo (Biotec) for his useful insights on natural

selection and population genetics analyses.

Some sequence data were produced by the US Department of Energy (DOE) Joint Genome

Institute (https://www.jgi.doe.gov/) in collaboration with the user community. We especially

thank Dr. Rytas Vilgalys and Dr. Francis Martin for using the *Diaporthaceae* sp. PMI_573

genome generated within the framework of the 1000 Fungal Genome projects (CSP 1974).

We also thank Dr. Riccardo Baroncelli for letting us use the genomes of *Colletotrichum*

melonis CBS 134730 (Colme1) and *Colletotrichum tamarilloi* CBS 129955 (Colta1).

Benefit-sharing section statement

A research collaboration established with scientists from Westerdijk Fungal Biodiversity

Institute, The Netherlands, was helpful in *in-silico* analyses of the secondary metabolism

related gene clusters in the blast fungal pathogen. All collaborators are included as co-authors.

The results of research, addressing a priority concern, will be shared with the broader

scientific community.

References

Asuke, S., Tanaka, M., Hyon, G.-S., Inoue, Y., Vy, T. T. P., Niwamoto, D., Nakayashiki, H., & Tosa, Y. (2020). Evolution of an *Eleusine* -Specific Subgroup of *Pyricularia oryzae* Through a Gain of an Avirulence Gene. *Molecular Plant-Microbe Interactions®*, *33*(2), 153–165. https://doi.org/10.1094/MPMI-03-19-0083-R

Blin, K., Shaw, S., Kloosterman, A. M., Charlop-Powers, Z., Van Wezel, G. P., Medema, M. H., & Weber, T. (2021). antiSMASH 6.0: Improving cluster detection and comparison capabilities. *Nucleic Acids Research*, *49*(W1), W29–W35. https://doi.org/10.1093/nar/gkab335

Böhnert, H. U., Fudal, I., Dioh, W., Tharreau, D., Notteghem, J. L., & Lebrun, M. H. (2004). A putative polyketide synthase/peptide synthetase from Magnaporthe grisea signals pathogen attack to resistant rice. *Plant Cell*, *16*(9), 2499–2513. https://doi.org/10.1105/tpc.104.022715

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Brunner, P. C., & McDonald, B. A. (2018). Evolutionary analyses of the avirulence effector AvrStb6 in global populations of *Zymoseptoria tritici* identify candidate amino acids involved in recognition. *Molecular Plant Pathology*, *19*(8), 1836–1846. https://doi.org/10.1111/mpp.12662

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, *25*(15), 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Chen, S., & Qiang, S. (2017). Recent advances in tenuazonic acid as a potential herbicide. *Pesticide Biochemistry and Physiology*, *143*, 252–257. https://doi.org/10.1016/j.pestbp.2017.01.003

Chernomor, O., Von Haeseler, A., & Minh, B. Q. (2016). Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Systematic Biology*, *65*(6), 997–1008. https://doi.org/10.1093/SYSBIO/SYW037

Chiapello, H., Mallet, L., Guérin, C., Aguileta, G., Amselem, J., Kroj, T., Ortega-Abboud, E., Lebrun, M.-H., Henrissat, B., Gendrault, A., Rodolphe, F., Tharreau, D., & Fournier, E. (2015). Deciphering Genome Content and Evolutionary Relationships of Isolates from the Fungus Magnaporthe oryzae Attacking Different Host Plants. *Genome Biology and Evolution*, *7*(10), 2896–2912. https://doi.org/10.1093/gbe/evv187

Chuma, I., Isobe, C., Hotta, Y., Ibaragi, K., Futamata, N., Kusaba, M., Yoshida, K., Terauchi, R., Fujita, Y., Nakayashiki, H., Valent, B., & Tosa, Y. (2011). Multiple translocation of the AVR-Pita effector gene among chromosomes of the rice blast fungus Magnaporthe oryzae and related species. *PLoS Pathogens*, *7*(7), e1002147. https://doi.org/10.1371/journal.ppat.1002147

Chumley, F. G., & Valent, B. (1990). Genetic Analysis of Melanin-Deficient, Nonpathogenic Mutants of Magnaporthe grisea. *Molecular Plant-Microbe Interactions*, *3*(3), 135. https://doi.org/10.1094/MPMI-3-135

Collemare, J., Billard, A., Böhnert, H. U., & Lebrun, M.-H. (2008). Biosynthesis of secondary metabolites in the rice blast fungus Magnaporthe grisea: The role of hybrid PKS-NRPS in pathogenicity. *Mycological Research*, *112*(2), 207–215. https://doi.org/10.1016/j.mycres.2007.08.003

Collemare, J., O'Connell, R., & Lebrun, M. H. (2019). Nonproteinaceous effectors: The terra incognita of plant–fungal interactions. *New Phytologist*, *223*(2), 590–596. https://doi.org/10.1111/nph.15785

Collemare, J., Pianfetti, M., Houlle, A.-E., Morin, D., Camborde, L., Gagey, M.-J., Barbisan, C., Fudal, I., Lebrun, M.-H., & Böhnert, H. U. (2008). *Magnaporthe grisea* avirulence gene *ACE1* belongs to an infection-specific gene cluster involved in secondary metabolism. *New Phytologist*, *179*(1), 196–208. https://doi.org/10.1111/j.1469-8137.2008.02459.x

Crouch, J. A., & Tomaso-Peterson, M. (2012). Anthracnose disease of centipedegrass turf caused by Colletotrichum eremochloae, a new fungal species closely related to Colletotrichum sublineola. *Mycologia*, *104*(5), 1085–1096. https://doi.org/10.3852/11-317

Dallery, J.-F., Lapalu, N., Zampounis, A., Pigné, S., Luyten, I., Amselem, J., Wittenberg, A. H. J., Zhou, S., de Queiroz, M. V., Robin, G. P., Auger, A., Hainaut, M., Henrissat, B., Kim, K.-T., Lee, Y.-H., Lespinet, O., Schwartz, D. C., Thon, M. R., & O'Connell, R. J. (2017). Gapless genome assembly of Colletotrichum higginsianum reveals chromosome structure and association of transposable elements with secondary metabolite gene clusters. *BMC Genomics*, *18*(1), 667. https://doi.org/10.1186/s12864-017-4083-x

Dellaporta, S. L., Wood, J., & Hicks, J. B. (1983). A plant DNA minipreparation: Version II. *Plant Molecular Biology Reporter*, *1*(4), 19–21. https://doi.org/10.1007/BF02712670

Ebbole, D. J. (2007). Magnaporthe as a model for understanding host-pathogen interactions. *Annual Review of Phytopathology*, *45*, 437–456. https://doi.org/10.1146/annurev.phyto.45.062806.094346

Farman, M. L., & Kim, Y. S. (2005). Telomere hypervariability in Magnaporthe oryzae. *Molecular Plant Pathology*, *6*(3), 287–298. https://doi.org/10.1111/J.1364-3703.2005.00285.X

Fudal, I., Collemare, J., Böhnert, H. U., Melayah, D., & Lebrun, M.-H. (2007). Expression of Magnaporthe grisea Avirulence Gene ACE1 Is Connected to the Initiation of Appressorium-Mediated Penetration. *Eukaryotic Cell*, *6*(3), 546–554. https://doi.org/10.1128/EC.00330-05

Gel, B., & Serra, E. (2017). karyoploteR: an R/Bioconductor package to plot customizable genomes displaying arbitrary data. *Bioinformatics*, *33*(19), 3088–3090. https://doi.org/10.1093/BIOINFORMATICS/BTX346

Gilchrist, C. L. M., & Chooi, Y. H. (2021). clinker & clustermap.js: Automatic generation of gene cluster comparison figures. *Bioinformatics*, *37*(16), 2473–2475. https://doi.org/10.1093/BIOINFORMATICS/BTAB007

Giraud, T., Gladieux, P., & Gavrilets, S. (2010). Linking the emergence of fungal plant diseases with ecological speciation. *Trends in Ecology & Evolution*, *25*(7), 387–395. https://doi.org/10.1016/J.TREE.2010.03.006

Gladieux, P., Condon, B., Ravel, S., Soanes, D., Maciel, J. L. N., Nhani, A., Chen, L., Terauchi, R., Lebrun, M., Tharreau, D., Mitchell, T., Pedley, K. F., Valent, B., Talbot, N. J., Farman, M., & Fournier, E. (2018). Gene Flow between Divergent Cereal- and Grass-Specific Lineages of the Rice Blast Fungus Magnaporthe oryzae. *mBio*, *9*(1), 1–19. https://doi.org/10.1128/mBio.01219-17

Gladieux, P., Ropars, J., Badouin, H., Branca, A., Aguileta, G., De Vienne, D. M., Rodríguez De La Vega, R. C., Branco, S., & Giraud, T. (2014). Fungal evolutionary genomics

provides insight into the mechanisms of adaptive divergence in eukaryotes. *Molecular Ecology*, *23*(4), 753–773. https://doi.org/10.1111/mec.12631

Gómez Luciano, L. B., Tsai, I. J., Chuma, I., Tosa, Y., Chen, Y. H., Li, J. Y., Li, M. Y., Lu, M. Y. J., Nakayashiki, H., & Li, W. H. (2019). Blast Fungal Genomes Show Frequent Chromosomal Changes, Gene Gains and Losses, and Effector Gene Turnover. *Molecular Biology and Evolution*, *36*(6), 1148–1161. https://doi.org/10.1093/molbev/msz045

Grigoriev, I. V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otillar, R., Riley, R., Salamov, A., Zhao, X., Korzeniewski, F., Smirnova, T., Nordberg, H., Dubchak, I., & Shabalov, I. (2014). MycoCosm portal: Gearing up for 1000 fungal genomes. *Nucleic Acids Research*, *42*(D1), D699–D704. https://doi.org/10.1093/NAR/GKT1183

Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., & Gascuel, O. (2010). New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Systematic Biology*, *59*(3), 307–321. https://doi.org/10.1093/SYSBIO/SYQ010

Harris, L. J., Balcerzak, M., Johnston, A., Schneiderman, D., & Ouellet, T. (2016). Host-preferential Fusarium graminearum gene expression during infection of wheat, barley, and maize. *Fungal Biology*, *120*(1), 111–123. https://doi.org/10.1016/j.funbio.2015.10.010

Hartmann, F. E., McDonald, B. A., & Croll, D. (2018). Genome-wide evidence for divergent selection between populations of a major agricultural pathogen. *Molecular Ecology*, *27*(12), 2725–2741. https://doi.org/10.1111/mec.14711

Hoang, D. T., Chernomor, O., Von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, *35*(2), 518–522. https://doi.org/10.1093/MOLBEV/MSX281

Inoue, Y., Vy, T. T. P., Yoshida, K., Asano, H., Mitsuoka, C., Asuke, S., Anh, V. L., Cumagun, C. J. R., Chuma, I., Terauchi, R., Kato, K., Mitchell, T., Valent, B., Farman, M., & Tosa, Y. (2017). Evolution of the wheat blast fungus through functional losses in a host specificity determinant. *Science (New York, N.Y.)*, *357*(6346), 80–83. https://doi.org/10.1126/science.aam9654

Izumi, Y., Ohtani, K., Miyamoto, Y., Masunaka, A., Fukumoto, T., Gomi, K., Tada, Y., Ichimura, K., Peever, T. L., & Akimitsu, K. (2012). A Polyketide Synthase Gene, *ACRTS2* , Is Responsible for Biosynthesis of Host-Selective ACR-Toxin in the Rough Lemon Pathotype of *Alternaria alternata*. *Molecular Plant-Microbe Interactions®*, *25*(11), 1419–1429. https://doi.org/10.1094/MPMI-06-12-0155-R

Jacob, S., Grötsch, T., Foster, A. J., Schüffler, A., Rieger, P. H., Sandjo, L. P., Liermann, J. C., Opatz, T., & Thines, E. (2017). Unravelling the biosynthesis of pyriculol in the rice blast fungus magnaporthe oryzae. *Microbiology (United Kingdom)*, *163*(4), 541–553. https://doi.org/10.1099/mic.0.000396

Jeon, J., Lee, G. W., Kim, K. T., Park, S. Y., Kim, S., Kwon, S., Huh, A., Chung, H., Lee, D. Y., Kim, C. Y., & Lee, Y. H. (2020). Transcriptome profiling of the rice blast fungus magnaporthe oryzae and its host oryza sativa during infection. *Molecular Plant-Microbe Interactions*, *33*(2), 141–144. https://doi.org/10.1094/MPMI-07-19-0207-A

Jonge, R. de, Bolton, M., Kombrink, A., Berg, G. V. D. van den, Yadeta, K. A., & Thomma, B. (2013). Extensive chromosomal reshuffling drives evolution of virulence in an asexual pathogen. *Genome Research*. https://doi.org/10.1101/GR.152660.112

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A., & Jermiin, L. S. (2017). ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods 2017 14:6*, *14*(6), 587–589. https://doi.org/10.1038/nmeth.4285

Katoh, K., Misawa, K., Kuma, K. I., & Miyata, T. (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066. https://doi.org/10.1093/NAR/GKF436

Kautsar, S. A., Blin, K., Shaw, S., Navarro-Muñoz, J. C., Terlouw, B. R., van der Hooft, J. J. J., van Santen, J. A., Tracanna, V., Suarez Duran, H. G., Pascal Andreu, V., Selem-Mojica, N., Alanjary, M., Robinson, S. L., Lund, G., Epstein, S. C., Sisto, A. C., Charkoudian, L. K., Collemare, J., Linington, R. G., … Medema, M. H. (2019). MIBiG 2.0: A repository for biosynthetic gene clusters of known function. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkz882

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher, G., & Hedges, S. B. (2022). TimeTree 5: An Expanded Resource for Species Divergence Times. *Molecular Biology and Evolution*, *39*(8), msac174. https://doi.org/10.1093/molbev/msac174

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., & Salzberg, S. L. (2004). Versatile and open software for comparing large genomes. *Genome Biology*, *5*(2), R12. https://doi.org/10.1186/gb-2004-5-2-r12

Lambou, K., Tag, A., Lassagne, A., Collemare, J., Clergeot, P.-H., Barbisan, C., Perret, P., Tharreau, D., Millazo, J., Chartier, E., Vries, R. P. D., Hirsch, J., Morel, J.-B., Beffa, R., Kroj, T., Thomas, T., & Lebrun, M.-H. (2024). The bZIP transcription factor BIP1 of the rice blast fungus is essential for infection and regulates a specific set of appressorium genes. *PLOS Pathogens*, *20*(1), e1011945. https://doi.org/10.1371/journal.ppat.1011945

Lanfear, R., Calcott, B., Kainer, D., Mayer, C., & Stamatakis, A. (2014). Selecting optimal partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology*, *14*(1), 1–14. https://doi.org/10.1186/1471-2148-14-82

Langner, T., Harant, A., Gomez-Luciano, L. B., Shrestha, R. K., Malmgren, A., Latorre, S. M., Burbano, H. A., Win, J., & Kamoun, S. (2021). Genomic rearrangements generate hypervariable mini-chromosomes in host-specific isolates of the blast fungus. *PLOS Genetics*, *17*(2), e1009386. https://doi.org/10.1371/journal.pgen.1009386

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*, *49*(W1), W293–W296. https://doi.org/10.1093/NAR/GKAB301

Lim, S. M., & Hooker, A. L. (1971). Southern Corn Leaf Blight: Genetic Control of Pathogenicity and Toxin Production in Race T and Race O of *Cochliobolus heterostrophus*. *Genetics*, *69*(1), 115–117. https://doi.org/10.1093/genetics/69.1.115

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A., Lanfear, R., & Teeling, E. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*, *37*(5), 1530–1534. https://doi.org/10.1093/molbev/msaa015

Navarro-Muñoz, J. C., Selem-Mojica, N., Mullowney, M. W., Kautsar, S. A., Tryon, J. H., Parkinson, E. I., De Los Santos, E. L. C., Yeong, M., Cruz-Morales, P., Abubucker, S., Roeters, A., Lokhorst, W., Fernandez-Guerra, A., Cappelini, L. T. D., Goering, A. W., Thomson, R. J., Metcalf, W. W., Kelleher, N. L., Barona-Gomez, F., & Medema, M. H. (2019). A computational framework to explore large-scale biosynthetic diversity. *Nature Chemical Biology*, 1–9. https://doi.org/10.1038/s41589-019-0400-9

Oh, Y., Donofrio, N., Pan, H., Coughlan, S., Brown, D. E., Meng, S., Mitchell, T., & Dean, R.
A. (2008). Transcriptome analysis reveals new insight into appressorium formation
and function in the rice blast fungus Magnaporthe oryzae. *Genome Biology*, *9*(5), R85.
https://doi.org/10.1186/gb-2008-9-5-r85
Oliva, R. F., Cano, L. M., Raffaele, S., Win, J., Bozkurt, T. O., Belhaj, K., Oh, S.-K., Thines,
M., & Kamoun, S. (2015). A Recent Expansion of the RXLR Effector Gene *Avrblb2*
Is Maintained in Global Populations of *Phytophthora infestans* Indicating Different
Contributions to Virulence. *Molecular Plant-Microbe Interactions®*, *28*(8), 901–912.
https://doi.org/10.1094/MPMI-12-14-0393-R
Palma-Guerrero, J., Ma, X., Torriani, S. F. F., Zala, M., Francisco, C. S., Hartmann, F. E.,
Croll, D., & McDonald, B. A. (2017). Comparative Transcriptome Analyses in
Zymoseptoria tritici Reveal Significant Differences in Gene Expression Among
Strains During Plant Infection. *Molecular Plant-Microbe Interactions: MPMI*, *30*(3),
231–244. https://doi.org/10.1094/MPMI-07-16-0146-R
Patkar, R. N., Benke, P. I., Qu, Z., Constance Chen, Y. Y., Yang, F., Swarup, S., & Naqvi, N.
I. (2015). A fungal monooxygenase-derived jasmonate attenuates host innate
immunity. *Nature Chemical Biology*, *11*(9), 733–740.
https://doi.org/10.1038/nchembio.1885
Patkar, R. N., & Naqvi, N. I. (2017). Fungal manipulation of hormone-regulated plant
defense. *PLoS Pathogens*, *13*(6), 10–14. https://doi.org/10.1371/journal.ppat.1006334
Patkar, R. N., Ramos-Pamplona, M., Gupta, A. P., Fan, Y., & Naqvi, N. I. (2012).
Mitochondrial β-oxidation regulates organellar integrity and is necessary for conidial
germination and invasive growth in *Magnaporthe oryzae*: Mitochondrial β-oxidation
in *Magnaporthe*. *Molecular Microbiology*, *86*(6), 1345–1363.
https://doi.org/10.1111/mmi.12060
Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – Approximately Maximum-
Likelihood Trees for Large Alignments. *PLoS ONE*, *5*(3), e9490.
https://doi.org/10.1371/journal.pone.0009490
Rahnama, M., Novikova, O., Starnes, J. H., Zhang, S., Chen, L., & Farman, M. L. (2020).
Transposon-mediated telomere destabilization: A driver of genome evolution in the
blast fungus. *Nucleic Acids Research*, gkaa287. https://doi.org/10.1093/nar/gkaa287
Rahnama, M., Wang, B., Dostart, J., Novikova, O., Yackzan, D., Yackzan, A., Bruss, H.,
Baker, M., Jacob, H., Zhang, X., Lamb, A., Stewart, A., Heist, M., Hoover, J., Calie,
P., Chen, L., Liu, J., & Farman, M. L. (2021). Telomere Roles in Fungal Genome
Evolution and Adaptation. *Frontiers in Genetics*, *12*, 676751.
https://doi.org/10.3389/fgene.2021.676751
Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J. C., Guirao-Rico, S., Librado, P., Ramos-
Onsins, S. E., & Sánchez-Gracia, A. (2017). DnaSP 6: DNA Sequence Polymorphism
Analysis of Large Data Sets. *Molecular Biology and Evolution*, *34*(12), 3299–3302.
https://doi.org/10.1093/molbev/msx248
Rudd, J. J., Kanyuka, K., Hassani-Pak, K., Derbyshire, M., Andongabo, A., Devonshire, J.,
Lysenko, A., Saqi, M., Desai, N. M., Powers, S. J., Hooper, J., Ambroso, L., Bharti,
A., Farmer, A., Hammond-Kosack, K. E., Dietrich, R. A., & Courbot, M. (2015).
Transcriptome and metabolite profiling of the infection cycle of Zymoseptoria tritici
on wheat reveals a biphasic interaction with plant immunity involving differential
pathogen chromosomal contributions and a variation on the hemibiotrophic lifestyle
definition. *Plant Physiology*, *167*(3), 1158–1185.
https://doi.org/10.1104/pp.114.255927

Sánchez-Vallet, A., Fouché, S., Fudal, I., Hartmann, F. E., Soyer, J. L., Tellier, A., & Croll, D. (2018). The Genome Biology of Effector Gene Evolution in Filamentous Plant Pathogens. *Annual Review of Phytopathology*, *56*(1), 21–40. https://doi.org/10.1146/annurev-phyto-080516-035303

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., & Ideker, T. (2003). Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Research*, *13*(11), 2498–2504. https://doi.org/10.1101/GR.1239303

Shen, Q., Liu, Y., & Naqvi, N. I. (2018). Fungal effectors at the crossroads of phytohormone signaling. *Current Opinion in Microbiology*, *46*, 1–6. https://doi.org/10.1016/J.MIB.2018.01.006

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(19), 3210–3212. https://doi.org/10.1093/BIOINFORMATICS/BTV351

Soundararajan, S., Jedd, G., Li, X., Ramos-pamplon, M., Chua, N. H., & Naqvi, N. I. (2004). Woronin body function in Magnaporthe grisea is essential for efficient pathogenesis and for survival during nitrogen starvation stress. *The Plant Cell*, *16*(June), 1564–1574. https://doi.org/10.1105/tpc.020677.2001

Stanke, M., Schöffmann, O., Morgenstern, B., & Waack, S. (2006). Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics*, *7*(1), 1–11. https://doi.org/10.1186/1471-2105-7-62/TABLES/2

Starnes, J. H., Thornbury, D. W., Novikova, O. S., Rehmeyer, C. J., & Farman, M. L. (2012). Telomere-Targeted Retrotransposons in the Rice Blast Fungus *Magnaporthe oryzae*: Agents of Telomere Instability. *Genetics*, *191*(2), 389–406. https://doi.org/10.1534/genetics.111.137950

Talbot, N. J. (2003). On the Trail of a Cereal Killer: Exploring the Biology of Magnaporthe grisea. *Annual Review of Microbiology*, *57*(1), 177–202. https://doi.org/10.1146/annurev.micro.57.030502.090957

Tsuge, T., Harimoto, Y., Akimitsu, K., Ohtani, K., Kodama, M., Akagi, Y., Egusa, M., Yamamoto, M., & Otani, H. (2013). Host-selective toxins produced by the plant pathogenic fungus *Alternaria alternata*. *FEMS Microbiology Reviews*, *37*(1), 44–66. https://doi.org/10.1111/j.1574-6976.2012.00350.x

Van Der Hoorn, R. (2002). Balancing selection favors guarding resistance proteins. *Trends in Plant Science*, *7*(2), 67–71. https://doi.org/10.1016/S1360-1385(01)02188-4

Vy, T. T. P., Inoue, Y., Asuke, S., Chuma, I., Nakayashiki, H., & Tosa, Y. (2024). The ACE1 secondary metabolite gene cluster is a pathogenicity factor of wheat blast fungus. *Communications Biology*, *7*(1), 812. https://doi.org/10.1038/s42003-024-06517-7

Walton, J. D. (2006). HC-toxin. *Phytochemistry*, *67*(14), 1406–1413. https://doi.org/10.1016/j.phytochem.2006.05.033

Yan, X., Tang, B., Ryder, L. S., MacLean, D., Were, V. M., Eseola, A. B., Cruz-Mireles, N., Ma, W., Foster, A. J., Osés-Ruiz, M., & Talbot, N. J. (2023). The transcriptional landscape of plant infection by the rice blast fungus *Magnaporthe oryzae* reveals distinct families of temporally co-regulated and structurally conserved effectors. *The Plant Cell*, koad036. https://doi.org/10.1093/plcell/koad036

Yoshida, K., Saunders, D. G. O., Mitsuoka, C., Natsume, S., Kosugi, S., Saitoh, H., Inoue, Y., Chuma, I., Tosa, Y., Cano, L. M., Kamoun, S., & Terauchi, R. (2016). Host

specialization of the blast fungus Magnaporthe oryzae is associated with dynamic gain
and loss of genes linked to transposable elements. *BMC Genomics*, *17*(1), 370.
https://doi.org/10.1186/s12864-016-2690-6
Zhong, Z., Norvienyeku, J., Chen, M., Bao, J., Lin, L., Chen, L., Lin, Y., Wu, X., Cai, Z.,
Zhang, Q., Lin, X., Hong, Y., Huang, J., Xu, L., Zhang, H., Chen, L., Tang, W.,
Zheng, H., Chen, X., … Nielsen, R. (2016). Directional Selection from Host Plants Is
a Major Force Driving Host Specificity in Magnaporthe Species. *Scientific Reports*, *6*,
25591. https://doi.org/10.1038/srep25591
852

 Figure captions

Figure 1. *Pyricularia oryzae* **worldwide populations are organized in host-specific**
lineages. A) Geographic distribution of sequenced *P. oryzae* strains. The color-coded dots
denote the host plant of origin of the strains. B) Phylogenetic tree constructed based on
concatenation of 2655 BUSCO genes present in all 68 *P. oryzae* genomes used in this study.
Colored shapes in the background depict different host-specific genetic lineages.

Figure 2. Similarity network analysis of biosynthetic gene clusters (BGCs) from
Pyricularia oryzae **and** *Pyricularia grisea.* A BiG-SCAPE analysis with a cutoff c0.5 depicts
similarity of 4224 BGCs from *P. oryzae* or *P. grisea* with 277 reference BGCs from MIBiG
database. Each dot represents a BGC and is color-coded according to the lineage. Gene cluster
families (GCF; subnetworks) marked with green boxes share significant homology with
reference MIBiG BGCs (grey-colored circle). GCFs marked with red boxes are found to be
unique to host-specific lineages. The length of the gray lines is proportional to the genetic
distance between BGCs. Singletons are shown as individual dots at the bottom.

Figure 3. BGC-O1 is predominantly present in *Oryza* **lineage of** *Pyricularia oryzae*. BiG-
SCAPE network showing BGC-O1 present in 23 genomes from *Oryza* and one genome from
Eragrostis lineages. Conservation of the BGC in selected strains is depicted using Clinker.
Vertical bars besides the name of strain are colored according to the lineages. Red and gray
dashed boxes separate BGC-O1 and BGC-O1-like clusters, respectively.

Figure 4. Genomic localization of BGC-O1 and synteny between the *Oryza* **and non-**
Oryza **lineages.** A) Synteny analysis using pairwise comparison within *Oryza* lineage.
Genomes of GUY11 and FR13 aligned individually with reference genome 70-15. The
syntenic BGC-O1 locus (orange) and the flanking regions (blue) on chromosome 2 of 70-15
are depicted. The differential lengths of BGC-O1 in different isolates are marked with a bar
and corresponding length in kilobases. The black arrowhead depicts genomic rearrangement
(swapping) in the flanking ~10 Kb region. The red arrowhead marks the rearrangement of ~9
Kb region of the BGC-O1 in FR13. B) Synteny analyses of representative genome assemblies
belonging to *Setaria* (US71), *Eleusine* (MZ5-1-6 and CD156) and *Oryza* (70-15) lineages.
The BGC-O1 locus (orange) and flanking region (blue) sequences from US71, MZ5-1-6 and
CD156 aligned with that of 70-15 are depicted.

Figure 5. Evolutionary origin of the reducing polyketide synthase (PKS) MGG_08236
gene. A) Maximum-likelihood phylogenetic tree using protein sequences of reducing PKS
genes from BGC-O1 and related gene cluster families, as well as homologues retrieved from
MycoCosm repository. Tip labels depicting sequences from *Oryza* and *Eragrostis*-specific
clade are marked with blue and pink background, respectively. The *Colletotrichum*
eremocloae ortholog (jgi.p_Coler1_670826), closer to MGG_08236, is denoted with pink
label; whereas the more distant paralogue (jgi.p_Coler1_679399) is shown in orange. Gray
shaded triangles denote collapsed clades with distant sequences. Branches were supported by
> 95% Bootstrap values indicated with gray circles at the nodes. The tree is rooted at

midpoint. B) Comparative analysis, using Clinker tool, of the BGC-O1 or BGC-O1-like loci
from three *Oryza*-specific and *C. eremochloae* genomes.

Figure 6. BGC-O1 genes are specifically expressed during pathogenic development. A)
Gel electrophoresis of PCR products displaying presence or absence of ORF region of rPKS
MGG_08236 from genomic DNA of the indicated *P. oryzae* strains. B157, MOS2, MOS3 and
MOS5 belong to the *Oryza* lineage (labelled in blue); whereas, the MOS1, MEC1, MEC4,
MEC5, MEC6 and MEA1 belong to the *Eleusine* lineage (labelled in red). * - the only *Oryza*
strain that lacked the BGC-O1 in *in-silico* analysis. ** - the *Oryza* strains placed outside
Oryza lineage. Arrowheads corresponds to the 1 Kb size of band from Marker. B) RT-PCR
gel depicting expression of genes MGG_08236 (Polyketide synthase), MGG_15107 (Methyl
transferase), MGG_15108 (Co-A transferase) and MGG_12496 (Cytochrome P450) in *Oryza*-
specific strain B157 at different stages of rice infection (12 to 96 hours post inoculation) and
during vegetative growth (mycelium) in complete medium.

Supplementary Information

Supplementary tables:

Supplementary Table S1: *Pyricularia oryzae* and *Pyricularia grisea* genomes used in this study.
Supplementary Table S2: *De novo* assembly statistics and gene prediction content of *Pyricularia oryzae* strains sequenced in this study.
Supplementary Table S3: Number of BGCs predicted using fungiSMASH in this study.
Supplementary Table S4: Summary of population genetic variation in 40 core biosynthetic genes in *Oryza*, *Eleusine* and *Triticum* lineages of *P. oryzae*.
Supplementary Table S5: Oligonucleotide primers used to study the relative expression of BGC-O1 genes.

Supplementary figures:

Supplementary Figure S1: *Pyricularia oryzae* and *Pyricularia grisea* are evolutionarily distinct species adapted to different hosts.
Supplementary Figure S2: Pathogenicity tests of *P. oryzae* strains isolated from different hosts and geographic locations in India.
Supplementary Figure S3: Occurrence of BGCs associated with specific classes of SM in individual genomes of *P. oryzae* and *P. grisea*.
Supplementary Figure S4: Homology of melanin-associated *P. oryzae* and *P. grisea* gene cluster family with reference BGCs in MIBiG database.
Supplementary Figure S5: Analysis of homology between *ACE1* gene cluster family in *P. oryzae* and *P. grisea* and reference BGCs in MIBiG database.
Supplementary Figure S6: Analysis of homology between putative epipyriculol-associated gene cluster family in *P. oryzae* and *P. grisea* and reference BGCs in MIBiG database.
Supplementary Figure S7: Homology of putative Squalestatin S1-associated *P. oryzae* and *P. grisea* gene cluster family with reference BGCs in MIBiG database.
Supplementary Figure S8: Analysis of homology between Tenuazonic acid-associated gene cluster families in host-adapted *P. oryzae* and *P. grisea* isolates.
Supplementary Figure S9: BGC-O2 is predominantly present in *Oryza*-specific lineage of *P. oryzae*.
Supplementary Figure S10: BGC-TLE is predominantly present in *Triticum*, *Lolium* and *Eleusine*-specific lineages of *P. oryzae*.
Supplementary Figure S11: Comparison of core biosynthetic gene region of BGC-TLE shows the pseudogenization in NRPS gene belonging to *Triticum* lineage.

Supplementary datasets:

Supplementary dataset S1: Amino acid sequences of 2655 BUSCOs present in 68 strains of *P. oryzae*.
Supplementary dataset S2: Amino acid sequence alignments of 2655 BUSCOs present in 68 strains of *P. oryzae*.

Supplementary dataset S3: Best partitioning scheme used to construct Maximum-likelihood
tree for *P. oryzae* strains.
Supplementary dataset S4: Maximum-likelihood tree based on 2655 BUSCOs present in 68
strains of *P. oryzae*.
Supplementary dataset S5: Amino acid sequences of 2557 BUSCOs present in 71 strains of
P. oryzae and *P. grisea*.
Supplementary dataset S6: Amino acid sequence alignments of 2557 BUSCOs present in 71
strains of *P. oryzae* and *P. grisea*.
Supplementary dataset S7: Maximum-likelihood tree based on 2557 BUSCOs present in 71
strains of *P. oryzae* and *P. grisea*.
Supplementary dataset S8: Manually curated gene models from *Oryza*-lineage-specific
BGC-O1.
Supplementary dataset S9: Amino acid sequences of the subset of rPKS (MGG_08236)
homologs in *Pezizomycotina* group.
Supplementary dataset S10: Amino acid sequence alignments of the subset of rPKS
(MGG_08236) homologs in *Pezizomycotina* group.
Supplementary dataset S11: Maximum-likelihood tree based on the subset of rPKS
(MGG_08236) homologs in *Pezizomycotina* group.

A

Host of origin: ● *Oryza* ● *Eleusine* ● *Setaria* ● *Triticum* ● *Lolium* ● *Eragrostis* ● *Digitaria*

B

Tree scale: 0.001

BGC-O1
Oryza specific

Genes

- Polyketide synthase | MGG_08236
- beta-xylosidase | MGG_12489
- Universal minicircle sequence binding protein | MGG_08239
- MGG_12491
- MGG_12492
- Methyl transferase | MGG_15107
- CoA-transferase | MGG_15108
- Cytochrome P450 | MGG_12496
- Cytochrome P450 | MGG_12497
- Glycosyl transferase | MGG_08232
- RNAse HI domain-containing protein | GUY11_02554, GUY11_02548
- transposase | GUY11_02553
- Retrotransposons-Gypsy family | Sar2_05502,Sar2_05503
- Zinc finger domain-containing protein | Sar2_05501
- BLM10_mid and DUF3437 domain-containing protein | MGG_04342
- Transferase | MGG_04340
- MGG_04339
- Potassium uptake protein | MGG_04338
- Catalase/peroxidase | MGG_04337
- DUF3445 domain-containing protein | MGG_04336, MGG_04335
- FAD binding domain-containing protein | MGG_04334

Lineages

● *Oryza* ● *Eleusine* ● *Setaria* ● *Triticum* ● *Lolium* ● *Eragrostis* ● *Digitaria* ● Known fungal BGCs

A

GUY11
MQOP01000012.1

BGC-O1
45 kb

70-15
NC_017850.1

48 kb

FR13
UEMA03000009.1

42 kb

FR13
UEMA03000012.1

B

US71
UCNY03000003.1

70-15
NC_017850.1

BGC-O1
48 kb

MZ5-1-6
CP034205.1

CD156
UELZ03000008.1

Lineages
● *Oryza* ● *Eleusine* ● *Setaria*

A

Tree scale: 0.1 ⊢——⊣

jgi|Colgr1|2442|GLRG 11878T0
jgi|Coler1|679399|MIX14804 34 33
jgi|Colso1|621565|MIX19503 23 27
jgi|Colsu1|687791|estExt Genemark1.C 10058
jgi|Colzo1|733567|estExt Genemark1.C 3150005
jgi|Gloci1|1841336|e gw1.15.201.1
jgi|Color1|8761|CORC01 00678T0
jgi|Colph1|455038|fgenesh1 pg.33 28
jgi|Colgo1|716101|estExt fgenesh1 pg.C 230218
jgi|Gloac1|1398789|estExt Genewise1Plus.C 520057
jgi|Colfi1|281529|mRNA6349
jgi|Colac2|797178|estExt Genemark1.C 1740014
jgi|Colsi1|979924|mrna 8044
jgi|Colny1|1022492|mrna 14212
jgi|Colme1|1833|mrna 1833
jgi|Colpa1|2449|mrna 12266
jgi|Colab1|4797|mrna 4797
jgi|Colco1|15579|mrna 8055
jgi|Collup1|10609|mrna 10609
jgi|Collu1|410842|e gw1.30.48.1
jgi|Colcu1|7861|mrna 7861
jgi|Colsp1|6070|mrna 3976
jgi|Colta1|1723|mrna 14004
jgi|Coler1|670826|MIX6231 187 8
G17.region45 L0 CDS8
IC17.region53 L0 CDS2
IE1K.region54 L0 CDS1
IB33.region53 L0 CDS2
RMgDl.region02 L0 CDS3
GUY11.region37 L0 CDS6
70-15.region24 L0 CDS6 | MGG_08236
P131.region20 L0 CDS1
MOS2.region14 L0 CDS2
Y34.region17 L0 CDS1
MG01.region08 L0 CDS1
jgi|MagorFR13 1|2134543
B157.region05 L0 CDS1
IA1.region54 L0 CDS1
FR13.region46 L0 CDS5
jgi|Magor1|5062
jgi|MagorTH3 1|1461275
FJ81278.region55 L0 CDS10
jgi|MagorGY11 1|1536307
IB49.region51 L0 CDS3
MOS5.region03 L0 CDS2
FJ72ZC7.region16 L0 CDS7
FJ98099.region18 L0 CDS16
PH14.region61 L0 CDS5
FJ98099.region18 L0 CDS9
ML33.region50 L0 CDS2
Sar2.region53 L0 CDS7
FJ98099.region17 L0 CDS7
AV1.region06 L0 CDS5

B