Identification of amnion progenitor-like cells at the amnion-epiblast boundary in the primate peri-gastrula

AUTHORS: Nikola Sekulovski[1], Lauren N. Juga[1], Chari L. Cortez[2], Michael Czerwinski[2,3], Allison E. Whorton[1], Jason R. Spence[2,3], Jenna K. Schmidt[4], Thaddeus G. Golos[4,5,6], Deborah L. Gumucio[2,3], Chien-Wei Lin[7] and Kenichiro Taniguchi[1,8,†]

AFFILIATIONS:
[1]Department of Cell Biology, Neurobiology and Anatomy, Medical College of Wisconsin, Milwaukee, WI 53226, USA
[2]Department of Cell and Developmental Biology, University of Michigan Medical School, Ann Arbor, MI 48109, USA
[3]Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI 48109, USA
[4]Wisconsin National Primate Research Center (WNPRC), Madison, WI, USA
[5]Department of Obstetrics and Gynecology, University of Wisconsin - Madison School of Medicine, Madison, WI USA
[6]Department of Comparative Biosciences, University of Wisconsin - Madison School of Veterinary Medicine, Madison, WI, USA
[7]Division of Biostatistics, Medical College of Wisconsin, Milwaukee, WI 53226, USA
[8]Department of Pediatrics, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[†]Corresponding author

CONTACTS:
Kenichiro Taniguchi
8701 Watertown Plank Road
B4245 BSB CB062
Milwaukee, WI 53226
ktaniguchi@mcw.edu

ABSTRACT

Amniogenesis is triggered in a collection of pluripotent epiblast cells as the human embryo implants. To gain insights into the critical but poorly understood transcriptional machinery governing amnion fate determination, we examined the evolving transcriptome of a developing human pluripotent stem cell-derived amnion model at the single cell level. This analysis revealed a previously unrecognized CLDN10$^+$ amnion progenitor cell type and enabled the identification of several markers specific to the different stages of lineage progression. Strikingly, we found that expression of CLDN10 is restricted to the amnion-epiblast boundary region in peri-gastrula cynomolgus macaque embryos, bolstering the notion that, at this stage, the amnion-epiblast boundary is a site of active amniogenesis. Bioinformatic analysis of published primate peri-gastrula single cell sequencing data further confirmed that CLDN10 is expressed in cells progressing to amnion. Overall, this study identifies a previously unrecognized CLDN10$^+$ amnion progenitor-like cell population and validates the presence of these fate transitioning cells at the boundary of the developing amnion.

INTRODUCTION

Amniogenesis is a critical event that is initiated during implantation in humans, eventually leading to the formation of an amniotic sac structure that surrounds and protects the developing embryo (Enders et al., 1983; Enders et al., 1986; Miki et al., 2005; Sasaki et al., 2016; Shahbazi and Zernicka-Goetz, 2018; Taniguchi et al., 2019). At implantation, the embryo (referred to at this time as a blastocyst) contains three morphologically and molecularly distinct cell types: a collection of unpolarized pluripotent epiblast cells (precursors to both the embryo proper and the amniotic ectoderm), a surrounding layer of polarized trophectoderm (a placental tissue precursor) and an underlying extraembryonic primitive endoderm (a yolk sac precursor). Upon implantation, the pluripotent epiblast cells undergo apico-basal polarization to form a cyst with a central lumen, the future amniotic cavity (Carleton et al., 2022). This event is followed by the fate transition of pluripotent epiblast cells at the uterine-proximal pole of the cyst to squamous amniotic ectoderm, forming a sharp boundary between amnion and pluripotent epiblast portions of the cyst (Shahbazi and Zernicka-Goetz, 2018; Shao and Fu, 2022; Taniguchi et al., 2019). This structure, known as the amniotic sac, represents the substrate for the next essential steps of embryonic development (e.g., primitive streak formation, neural specification).

Recent studies using early human and monkey embryos have provided a basic understanding of the transcriptomic characteristics of early primate amniogenesis (Bergmann et al., 2022; Nakamura et al., 2016; Nakamura et al., 2017; Sasaki et al., 2016; Tyser et al., 2021; Yang et al., 2021). Yet, experimental dissection of the molecular mechanisms involved in this process is difficult in these *in vivo* models. To enable molecular and cellular investigations, we and others developed human pluripotent stem cell-derived amnion models and showed that BMP signaling plays a crucial role in initiating amniogenesis (Chen et al., 2021; Overeem et al., 2023; Shao et al., 2017a; Shao et al., 2017b; Zheng et al., 2019). Recently, we further explored the BMP-dependent amniogenic transcriptional cascade, identifying several distinct transcriptional stages, as well as a requirement for TFAP2A-dependent transcription in regulating amnion fate progression (Sekulovski et al., 2023). However, changes in cellular differentiation states during amnion lineage progression remain largely unexplored.

In this study, we used single cell RNA sequencing (scRNA-seq) analysis to examine the dynamics in gene expression that accompany amnion differentiation of hPSC (herein referred to as the hPSC-amnion) cultured in a soft gel environment, called Gel-3D (Shao et al., 2017a). In this Gel-3D amnion model, BMP signaling, activated by a mechanosensitive cue provided by the soft substrate, is a critical initiator of human amniogenesis (Shao and Fu, 2022; Shao et al., 2017a; Taniguchi et al., 2019). Interestingly, the data reveal two likely contiguous amniogenic cell states: an amnion progenitor (AmP) state as well as an advanced amnion (advAm) state, each of which shows transcriptional similarities to distinct cell types in a Carnegie stage 7 human embryo (Tyser et al., 2021). Importantly, we identify a cohort of markers specific to each of these two amnion states, and validate selected markers for their expression in the amnion from cynomolgus macaque (*Macaca fascicularis*) embryos. Strikingly, we show that CLDN10, a marker of the AmP population, exhibits a restricted amniotic expression pattern at the boundary between the amnion and the epiblast of the cynomolgus

macaque peri-gastrula. This further supports our findings from a separate study
(Sekulovski et al., 2023), which show that the amnion-epiblast boundary is a site of
active amniogenesis in the macaque peri-gastrula. Together, this study provides
important single cell-level insight into human amnion fate progression and presents
additional evidence for the presence of amnion progenitor-like cells in primate embryos
undergoing gastrulation.
98
RESULTS AND DISCUSSION
Previously, we described culture conditions (Gel-3D) in which plated hPSC form
polarized cysts initially composed of columnar pluripotent cells, which then
spontaneously undergo squamous morphogenesis and begin to express amnion
markers (Shao et al., 2017a). Specifically, singly dissociated hPSC are plated densely
on a soft gel substrate to form aggregates and are allowed to grow for 24 hours (hrs) in
the presence of a ROCK inhibitor (Y-27632). After the first 24 hrs (day 1, d1), the ROCK
inhibitor is removed (a trigger to initiate lumenal cyst formation (Hamed et al., 2023;
Shao et al., 2017a; Taniguchi et al., 2015)), and a diluted extracellular matrix gel overlay
(2% v/v, e.g., Geltrex) is added to provide an additional 3D cue. Immunofluorescent (IF)
staining at this time shows cuboidal cells surrounding a central lumen; these cells
express NANOG, but not TFAP2A (**Fig. 1A**).
Interestingly, 24 hrs after adding the gel overlay (d2), a subset of cells within the
cysts display reduced expression of NANOG and SOX2, while TFAP2A expression
becomes weakly activated (**Fig. 1A**, arrowhead in d2). By d3, NANOG is no longer
detectable. Moreover, while many cells exhibit prominent TFAP2A expression, some
cells are only weakly positive for TFAP2A (**Fig. 1A**, arrowheads). This pattern of
TFAP2A expression is consistent with the previous observation that amniogenesis
initiates focally on the pluripotent cysts and then spreads laterally to form fully
squamous cysts (**Fig. 1A**, (Shao et al., 2017a; Shao et al., 2017b)). These molecular
changes are accompanied by a decrease in nuclear aspect ratio (a measure of
epithelial cell morphology, (Shao et al., 2017a; Townshend et al., 2020)) over time,
revealing more flattened (squamous) shapes by d3 and d4 (**Fig. 1B**). These results
provide further molecular and structural evidence that the transition from pluripotent to
amnion cell types occurs progressively over the cyst, starting from focal initiation sites
(Shao et al., 2017a; Shao et al., 2017b). As expected for such a spreading fate
conversion, d2 and d3 cysts contain cells representing several differentiation states.
To explore transcriptional programs of the fate transitioning cells during early
stages of human amnion development in this model, we performed scRNA-seq analysis
of d3 hPSC-amnion cysts, in which molecularly distinct amniotic cell types co-exist
within same cyst (**Fig. 1A**, d3). d3 hPSC-amnion cysts were dissociated into single
cells, followed by single cell isolation and labeling using the Fluidigm C3 system, cDNA
amplification and sequencing followed by demultiplexing (see Material and Methods for
additional details). This scRNA-seq dataset was examined using Seurat (Butler et al.,
2018) for data filtering, regression for genes associated with cell cycle progression,
normalization, variable gene selection and subsequent unsupervised clustering of cells
(see Materials and Methods). The data yield two distinct populations among 379 cells,
which, after marker and lineage analyses described below, are labeled AmP (amnion
progenitor, sage color) and advAm (advanced amnion, salmon color). Transcriptomic

features of these cells are visualized in Uniform Manifold Approximation and Projection
(UMAP) plots (**Fig. 1C**). Analysis of cell cycle markers reveals that cells of the AmP
population are predominantly in S phase, while in advAm cells, G1 phase markers
predominate (**Fig. S1**).
To broadly characterize the transcriptional state of each population, we examined
the expression of known amnion markers (Roost et al., 2015; Shao et al., 2017a; Yang
et al., 2021; Zhao et al., 2021; Zheng et al., 2022). Consistent with the amniotic lineage,
pan-amnion markers (*ISL1*, *TFAP2A* and *GATA3*, **Fig. 1D**, expression superimposed on
the UMAP plot in **Fig. 1C**) are expressed in cells of both populations. In contrast,
GABRP, *VTCN*1 and *IGFBP3* – differentiated amnion markers expressed in the first-
trimester human amnion (Roost et al., 2015) and detected in other human amnion
models (Shao et al., 2017a; Yang et al., 2021; Zhao et al., 2021; Zheng et al., 2022) –
are enriched primarily in salmon-colored cells (**Fig. 1E**), indicating that this population
contains cells characteristic of more advanced amnion states (advAm). Indeed,
pseudotime analysis (Cao et al., 2019) identifies distinct changes in gene expression
dynamics within the cell population, with cells on the left side of the plot showing
characteristics of less mature cells and those on the right side exhibiting more
differentiated profiles (**Fig. 1F**). In agreement with this idea, when RNA-velocity analysis
(velocyto, (La Manno et al., 2018)) was applied to examine lineage relationships among
the two populations based on the relative abundance of unspliced and spliced mRNA in
each cell, the majority of the RNA vector trajectories were directed toward advAm cells
on the right side of the plot (**Fig. 1G**, RNA velocity superimposed on the UMAP plot).
These results suggest that the cells marked in sage color in **Fig. 1C**, which do not
express later amnion markers, may be previously unrecognized precursor cell types
(AmP) that give rise to more mature amniotic cells.
In a recent study, we performed a detailed meta-analysis of the Tyser *et al.*
scRNA-seq dataset from a human embryo staged at Carnegie stage 7 (CS7) (Tyser et
al., 2021) and showed that the rod-shaped cluster of cells annotated as "Ectoderm" is
comprised of lineage progressing, lineage committed, and fully differentiated amnion
cell types (Sekulovski et al., 2023). The Tyser *et al*. UMAP plot is reproduced in **Fig. 1H**
with the original coordinates and annotations. Here, we show that *pan*-amnion markers
such as *GATA3, TFAP2A and ISL1* are broadly expressed in the rod-shaped cell
population marked "Ectoderm" (reproduced in **Fig. 1I**, see **Fig. S2** for uncropped
expression plots). Cells expressing the advAm markers, *GABRP*, *VTCN*1 and *IGFBP3*,
are found at the distal tip of this rod-shaped Ectoderm population (**Fig. 1J**). To examine
which cells in the CS7 human embryo share transcriptomic similarities to AmP or
advAm cells, the scRNA-seq dataset from the Tyser *et al.* CS7 human embryo was
combined with our d3 hPSC-amnion model and normalized using an integration feature
based on canonical correlation analysis in Seurat (Hao et al., 2021) (**Fig. 1K-M**).
Though the combined UMAP plot of this integrated dataset is different in shape from the
original Tyser *et al.* UMAP, it is clear that the cells from our d3 hPSC-amnion dataset
map to the regions marked by Tyser *et al*. as Epiblast, Primitive Streak and Ectoderm.
Moreover, while AmP cells are populated across the Tyser Epiblast and Primitive Streak
clusters, advAm cells overlap substantially with Tyser Ectoderm cells (**Fig. 1L,M**),
further validating the human amnion-like transcriptional characteristics of Gel-3D
amnion models (expression of selected markers shown in **Fig. S3**). Together, these

results strongly support our notion that AmP markers characterize a progenitor-like cell
population with transcriptional features that are more similar to pluripotent and primitive
streak cell types, and that further differentiated amnion cells can be identified using
advAm markers.
To identify genes that are unique to each cluster, unsupervised differential gene
enrichment analysis was performed (**Table 1**, $p < 0.05$, selected advAm cluster markers
shown in **Fig. 2A**). As expected, the advAm cluster shows enriched expression of
additional known differentiated amnion markers (e.g., *HIF-2A/EPAS1, IGFBP3, GABRP*)
as well as previously uncharacterized genes (**Table 1**, e.g., *ADAMTS18* and *SESN3*).
To confirm these findings in an *in vivo* system, we used RNA *in situ* hybridization to
examine the amniotic expression of selected differentiated markers (*GABRP*,
ADAMTS18 and *SESN3*, co-stained for *ISL1*) in a cynomolgus macaque (*Macaca*
fascicularis) embryo staged between Carnegie stages (CS) 12 and 13, an
organogenesis stage (**Fig. 2B,C**). Expression of all three markers is observed in the
amnion of this embryo, supporting the conclusion that the markers found in advAm cells
are valid markers of differentiated amnion. Interestingly, while *GABRP* and *SESN3*
show expression throughout the amnion, *ADAMTS18* expression is restricted to a
fraction of cells within the amnion (**Fig. 2C**), perhaps labeling a previously unrecognized
amnion sub-population.
The differential gene enrichment analysis also revealed that the AmP cells
(marked in sage in **Fig. 1C**) exhibit high expression of genes including *FABP5, IFITM1*
and *CLDN10* (**Fig. 3A** (dotplot), **3B** (UMAP)). Among these markers, only *CLDN10* (a
gene encoding Claudin-10, a component of tight junction) shows an enriched
expression in the "Primitive Streak" cells directly adjacent to the "Ectoderm" cells in the
Tyser *et al.* UMAP (**Fig. 3B**, uncropped image in **Fig. S4A**); we previously showed that
this subpopulation of the Primitive Streak cells displays weak *TBXT* expression, and
appears to include intermediate cells transitioning to amnion (Sekulovski et al., 2023), a
notion supported by a recent species comparison study by others (Ton et al., 2023). A
similar *TBXT* expression pattern is seen in a scRNA-seq dataset from d14 cynomolgus
macaque peri-gastrula (**Fig. S4B,C**, (Yang et al., 2021)). Indeed, CLDN10 enriched
cells are also seen within the $TBXT^{low}$ cell population in the d14 cynomolgus macaque
embryos (**Fig. S4C**), suggesting that CLDN10$^+$ progenitor cells might play a role in
amniogenesis at earlier timepoints.
Using IF staining in our hPSC-amnion model, we validated a dynamic expression
pattern of CLDN10 over time (**Fig. 3C**). In d2 hPSC-amnion cysts, CLDN10 expression
is restricted to the small fraction of cells that are weakly positive for TFAP2A (**Fig. 3C**,
arrowhead). At d3, CLDN10 is co-expressed with TFAP2A in several, but not all cells. In
contrast, most cells at d4 retain abundant TFAP2A expression but do not express
CLDN10 (**Fig. 3C**). Importantly, our analysis for nuclear aspect ratio (NAR) shows that,
at d2, CLDN10$^+$ cells (NAR = 0.54 ± 0.17 STDEV, n = 39) are significantly more
squamous compared to CLDN10$^-$ cells (NAR = 0.97 ± 0.31 STDEV, n = 87, $p<0.001$).
While previous studies implicated *CLDN10* as an amnion marker (Zheng et al., 2019;
Zheng et al., 2022), a detailed analysis was not performed. Thus, these data present
evidence suggesting that CLDN10 marks an early amnion state, but its expression is
extinguished in more differentiated cells. Consistent with this result, CLDN10 displays a

similar transient expression pattern in a newly established Glass-3D^{+BMP} hPSC-amnion
model (Sekulovski et al., 2023).
Interestingly, in PASE (post-implantation amniotic sac embryoids), an *in vitro*
model of human amniotic sac formation (Shao et al., 2017b), CLDN10 is localized
specifically to cells at the boundary between squamous amnion and columnar
pluripotent cells (**Fig. 3D**, arrowheads), an intriguing finding which suggests that cells
with a unique progenitor-like state might be present in the developing primate amniotic
sac *in vivo*. To test this, we analyzed CLDN10 expression in two cynomolgus macaque
embryos displaying the amniotic sac (**Fig. 3E**, staged between CS6 and CS7; **Fig. 3F**,
staged at CS7). Strikingly, both show restricted expression of CLDN10 at the boundary
separating amnion and epiblast tissues. These results are consistent with our recent
findings indicating that this boundary is likely a site of active amniogenesis in the
primate peri-gastrula (Sekulovski et al., 2023). Importantly, cynomolgus macaque
embryos from later stages (**Fig. 4A,B**, CS8; **Fig. 4C,D**, CS9) also show CLDN10
positive cells at the amnion-embryo boundary. In the CS8 embryo, CLDN10$^+$ cells are
seen at the tip of the head fold, a region spanning the anterior most neuroectoderm and
the nascent mediolateral surface ectoderm (**Fig. 4A,B** – inset (ii)) as well as the
posterior region of the embryo-amnion boundary (**Fig. 4A,B** – inset (i)). Moreover, in the
CS9 embryo, while a collection of cells expressing weak CLDN10 is seen posteriorly
(**Fig. 4C,D** – inset (iii)), CLDN10 expression is highly prominent anteriorly in the cells at
the boundary of the amnion and the developing mediolateral placode (**Fig. 4C,D** – inset
(i)). Together, these results establish the presence of previously unrecognized CLDN10$^+$
amnion-embryo boundary population during gastrulation as well as during early
organogenesis in the cynomolgus macaque.
In summary, we have profiled the transcriptomic signatures of lineage
progressing cells in an *in vitro* model of hPSC-amnion development at the single cell
level. Key results were validated *in vivo*, using early monkey embryo sections and two
published single cell RNA sequencing datasets from early human and cynomolgus
macaque embryos (Tyser et al., 2021; Yang et al., 2021). These analyses reveal a
previously unrecognized CLDN10$^+$ amnion progenitor-like cell population that is
localized at the boundary between the amnion and the epiblast of the amniotic sac in
cynomolgus macaque peri-gastrula. Strikingly, our recent study (Sekulovski et al., 2023)
provided evidence for an active amniogenic transcriptional cascade at the
epiblast/amnion boundary in primate peri-gastrula embryos; this present work reveals
CLDN10 as a key marker for this boundary cell population. Given the role of Claudin
proteins in controlling tight junction properties and epithelial characteristics (Angelow et
al., 2008), CLDN10 might be critical for organizing the boundary structure containing
shape/fate-transitioning cells flanked by squamous amnion or columnar/pseudostratified
pluripotent cells at each end. Molecularly, we speculate that the polarized distribution of
CLDN10 expression, which varies across embryo stages, might be linked to active BMP
signaling in each vicinity. We have previously shown that PASE are posteriorlized and
display high BMP signaling output (Shao et al., 2017b). At CS7, anterior BMP signaling
is likely inhibited by secreted BMP antagonists (e.g., Chordin, Noggin) derived from the
axial mesendoderm, while the posterior boundary is in close proximity to the primitive
streak with active BMP signaling (Anderson et al., 2002; Arnold and Robertson, 2009;
Shawlot et al., 1999; Yang et al., 2010). At CS8, CLDN10 appears to increase

anteriorly, perhaps due to increased BMP signaling output as the forebrain tissue
expands anteriorly away from the midline tissue (e.g., prechordal plate) that secretes
BMP antagonists (Anderson et al., 2002; Dobreva et al., 2012; Geng and Oliver, 2009;
Muenke and Beachy, 2000; Nishioka et al., 2005). By CS9, BMP signaling output may
weaken in the extending primitive streak, while strengthening anteriorly as BMP levels
in this region increase (Dobreva et al., 2012). Further analyses of dynamic BMP
signaling gradients will be required to test these hypotheses. Together, this study
provides an important single cell transcriptomic resource for future investigation of the
amniogenic transcriptional cascade in developing primate embryos.
284
MATERIALS AND METHODS
hESC lines
Human embryonic stem cell line H9 was used in this study (WA09, P30, P48, WiCell;
NIH registration number: 0062). All protocols for the use of hPSC lines were approved
by the Human Stem Cell Research Oversight Committee at the Medical College of
Wisconsin and the Human Pluripotent Stem Cell Research Oversight Committee at the
University of Michigan. All hPSC lines were maintained in a feeder-free system for at
least 20 passages and authenticated as karyotypically normal at the indicated passage
number. Karyotype analysis was performed at Cell Line Genetics. All hPSC lines tested
negative for mycoplasma contamination (LookOut Mycoplasma PCR Detection Kit,
Sigma-Aldrich). In summary, hESC were maintained in a feeder-free culture system with
mTeSR1 medium, or with 50%/50% mix of mTeSR1 and mTeSR plus (STEMCELL
Technologies). hESC were cultured on 1% (v/v) Geltrex (Thermo Fisher Scientific), or
with Cultrex SCQ (Bio-techne) coated 6 well plates (Nunc). Cells were passaged as
small clumps every 4 to 5 days with Dispase (Gibco). All cells were cultured at 37°C
with 5% CO_2. Media was changed every day. hESC were visually checked every day to
ensure the absence of spontaneously differentiated, mesenchyme-like cells in the
culture. Minor differentiated cells were scratched off the plate under a dissecting scope
once identified. The quality of all hESC lines was periodically examined by
immunostaining for pluripotency markers and successful differentiation to three germ
layer cells.
306
Gel-3D hPSC-amnion formation assays
Methods for these assays have been previously described (Shao et al., 2017a; Shao et
al., 2017b) with exceptions that 60-70uL of undiluted ECM gel solution is hand-streaked
on ice cold 22mmx22mm square coverslips, and that, in addition to Geltrex (Life
Technologies), Cultrex Ultimatrix (Bio-Techne) was also used. Singly dissociated cells
were prepared using Accutase (Sigma-Aldrich) and were plated on ECM gel-coated
coverslips at 25,000 cells/cm^2 in mTeSR1 or in 50%/50% mTeSR1/mTeSR plus mix
medium in the presence of 10µM Y-27632 (STEMCELL Technologies). After 24 hr, cells
were then incubated in media containing 2% Geltrex/Ultimatrix overlay without Y-27632
with daily media changes (note that 1% gel overlay was used between d2 to d4).
317
Cynomolgus macaque
Animals

The female and male cynomolgus macaques were housed and cared for at the Wisconsin
National Primate Research Center (WNPRC). All procedures were performed in
accordance with the NIH Guide for the Care and Use of Laboratory Animals and under
approval of the University of Wisconsin College of Letters and Sciences and Vice
Chancellor Office for Research and Graduate Education Institutional Animal Care and
Use Committee (protocol g005061).
326
Animal breeding and pregnancy detection
Beginning on day 8 post-onset of menses, the female was housed with a compatible
male and monitored for breeding. Animals were pair-housed until day 16-20 post-onset
of menses. A 2-3 mL blood draw was performed daily from day 8 post-onset of menses
until day 16 to assess the timing of ovulation based on the estradiol peak and rise in
progesterone in serum. Serum samples were analyzed for estradiol (25 μL) and
progesterone (20 µL) using a cobas e411 analyzer equipped with
ElectroChemiLuminescence technology (Roche, Basal, Switzerland) according to
manufacturer instructions. Results were determined via a calibration curve which was
instrument-generated by 2-point calibration using traceable standards and a master
curve provided via the reagent barcode. Inter-assay coefficient of variation (CV) was
determined by assaying aliquots of a pool of rhesus plasma. For estradiol, the limit of
quantitation (LOQ) was 25 pg/mL, the intra-assay CV was 2.02%, and the inter-assay
CV was 5.05%. For progesterone, the LOQ was 0.2 ng/mL, the intra-assay CV was
1.37%, and the inter-assay CV was 4.63%. A transabdominal ultrasound was performed
to detect pregnancy as early as 14 days post-ovulation. The ultrasound measurements
in combination with the timing of ovulation were used to estimate the day of conception
and gestational age of the pregnancy.
345
Terminal perfusion uterine collection, paraffin embedding, sectioning and staining
The pregnant females were sedated with intramuscular ketamine (>15 mg/kg) followed
by IV sodium pentobarbital (>35 mg/kg) and then perfused with 4% paraformaldehyde
(PFA) via the left ventricle. The entire uterus and cervix were removed. The serosa and
superficial myometrium were scored for better fixative penetration and to denote dorsal
and ventral orientation. Tissues were fixed in 4% PFA with constant stirring and solution
changes were performed every 24 hrs for a total of 72 hrs. The uterus was serially
sectioned from the fundus to the internal cervical os into 4 mm slices using a dissection
box. Cassettes with tissue sections were transferred into 70% ethanol, routinely
processed in an automated tissue processor and embedded in paraffin for histological
analyses (5 µm sections). Fixed uterine tissue were cut in 5 µm thick cross-sections,
mounted on slides, deparaffinized in xylene and rehydrated in an ethanol series. Antigen
retrieval was performed by boiling in citrate buffer. Sections were blocked 4% goat serum
in PBS at RT for at least 3-hr. Subsequent immunolocalization was performed using
commercially available primary antibodies, incubated overnight at 4 °C in 4% serum.
Immunofluorescent detection was performed using secondary antibodies tagged with a
fluorescent dye (fluorophores excitation = 488, 555, and 647nm), and counterstained with
DAPI. Negative controls were performed in which the primary antibody was substituted
with the same concentration of normal IgG of the appropriate isotype. Images were
obtained with a Zeiss LSM980 microscope.

Confocal microscopy of fixed samples
Confocal Images of fixed samples were acquired using a Nikon-A1 (Nikon) and a Zeiss LS980 laser scanning confocal microscopes. Non-3D images were generated using Zen, FIJI (NIH) and Photoshop (Adobe).

Immunostaining
hPSC-amnion cysts grown on the coverslip were rinsed with PBS (Gibco) twice, fixed with 4% paraformaldehyde (Sigma) for 60 min, then rinsed with PBS three times, and permeabilized with 0.1% SDS (Sigma, in 1x PBS) solution for 60 min. The samples were blocked in 4% heat-inactivated goat serum (Gibco) or 4% normal donkey serum (Gibco) in PBS overnight at 4°C. The samples were incubated with primary antibody solution prepared in blocking solution at 4°C for 48 hr, washed three times with PBS (30 min each), and incubated in blocking solution with goat or donkey raised Alexa Fluor-conjugated secondary antibodies (Thermo Fisher), at room temperature for 24 hours. Counter staining was performed using Hoechst 33342 (nucleus, Thermo Fisher Scientific), Alexa-Fluor-conjugated wheat germ agglutinin (membrane, Thermo Fisher Scientific) and Phalloidin (F-ACTIN, Thermo Fisher). All samples were mounted on slides using 90% glycerol (in 1x PBS). When mounting hPSC-cysts samples, modeling clay was used as a spacer between coverslip and slide to preserve lumenal cyst morphology. Antibodies for IF staining are found in **Table S2**.

RNA isolation and labeling
Single cell RNA sequencing
hPSC-amnion samples were rinsed twice with DMEM/F12, were briefly treated with 0.05% Trypsin (Life Technologies) for 1.5-min to initiate ECM gel digestion and were rinsed twice with DMEM/F12. Subsequently, the samples were incubated with Accutase (Sigma) for 10-min and were dissociated into single cells by gentle agitation. The scRNA-seq samples were prepared in the University of Michigan Rogel Cancer Center Single Cell Analysis Core. Cells were spun down for 5-min at 300xg, and the cell pellet was resuspended for Fluidigm C1 analysis (single cell preparation, single cell isolation and barcoding performed according to the manufacture's instructions). Libraries were sequenced (Illumina HiSeq 4000 H4K paired-end 150 cycle) followed by quality control (adaptor trimming, de-duplication), alignment and demultiplexing based on manufactured indices.

Bioinformatics – single cell RNA sequencing dataset analysis
Analysis of the d3 amnion scRNA-seq dataset was performed using the Seurat R package (v.4.2.1, (Hao et al., 2021)). Cells were filtered out if the total number of detected genes was less than 2,600 or if the expression percentage from mitochondrial genes was greater than 15%. Expression values were then log-normalized with scaling factor 10,000 for cell-level normalization, and further centered and scaled across genes. Principal component analysis (PCA) with top 30 PCs was performed prior to embedding 379 cells into two dimensional space with UMAP. Cell clusters were identified using FindClusters function based on a shared nearest neighbor modularity optimization

411   clustering algorithm (resolution set as 1). The cluster marker genes were identified
412   using FindAllMarkers function (Wilcoxon Rank Sum test).
413
Mapping and gene expression analyses of Tyser et al. Carnegie Stage 7 human embryo
single cell RNA sequencing dataset
The original UMAP coordinates and annotations for all 1,195 cells were used in **Fig. 1H**
as well as in the accompanying expression plots. The processed data were downloaded
from http://www.human-gastrula.net, which was used to generate a UMAP plot (using
DimPlot function in R package Seurat), as well as to perform expression analyses
(FeaturePlot function in Seurat).
421
422   *Integration and gene expression analyses of d3 amnion and Tyser et al. Carnegie Stage*
423   *7 human embryo single cell RNA sequencing datasets*
424   The two single cell datasets were integrated using a canonical correlation analysis
425   based approach implemented in the IntegrateData function with 4,000 anchor features
426   and 20 dimensions for anchor weighting in Seurat R package (v4.2.1).
427
Integration and gene expression analyses of Yang et al. GD14 cynomolgus macaque
embryo single cell RNA sequencing dataset
The processed data were downloaded from GSE148683, and the ensemble genome
build Macaca_fascicularis_5.0 release 96 was used to identify human orthogonal gene
symbols. Datasets from two distinct GD14 embryos (Yang et al., 2021) were integrated
using 5,000 anchor features and 10 dimensions in the IntegrateData function in Seurat
R package (v.4.2.1) package. Six general cell populations (epiblast, transition,
mesoderm, amnion, endoderm and extraembryonic mesenchyme) were identified using
FindClusters function (resolution as 0.4).
437
Pseudotime
Pseudotime analysis was performed using R package Monocle 3 (Cao et al., 2019).
UMAP was used to learn the principal graph using reverse graph embedding algorithm.
To decide the start of the trajectory, given that we defined that AmP is the initial lineage
in this dataset, we selected the node with the lowest value of the first UMAP component
as the root node. The estimated pseudotime was overlayed onto the original UMAP
coordinates.
445
RNA velocity (velocyto) analysis
After alignment to human genome hg19, the BAM files were used to generate a loom
file using the function run_smartseq2 in Python package velocyto (v0.17) with default
parameters to create spliced and unspliced gene count matrices. R package velocyto.R
(Python package v.0.6 (Bergen et al., 2020) was used to estimate inferred RNA
velocities of single cells and we visualized the velocities on the original UMAP
embedding.
453
FIGURES AND FIGURE LEGENDS
Figure 1. Single-cell transcriptomic signatures of d3 human amnion model

(A) Confocal images of d1, d2, d3 and d4 hPSC-amnion cysts stained for TFAP2A, an
amnion marker, and NANOG, a pluripotency marker. White arrowheads indicate cells
that display weak TFAP2A expression. Scale bars = 20μm.
(B) Quantitation for nuclear aspect ratio, a measure of epithelial cell morphology, of
developing hPSC-amnion at indicated timepoints, revealing a gradual squamous
morphogenesis. * indicates p < 0.05.
(C) A UMAP plot displaying the single cell transcriptomes of 379 d3 hPSC-amnion cells
with two identified clusters (AmP – sage color; advAm – salmon color).
(D,E) Expression of *pan*-amnion (*ISL1*, *TFAP2A* and *GATA3*) as well as late/advanced
amnion (*GABRP*, *VTCN1*, *IGFBP3*) markers superimposed onto the UMAP plot.
(F) Pseudotime progression (Monocle 3, indicated by the gradient scale) is
superimposed onto the UMAP plot.
(G) RNA velocity analysis (velocyto) of the d3 hPSC-amnion dataset, showing predicted
lineage progression trajectories from AmP (left) to advAm (right) populations.
(H) A UMAP plot displaying the original single cell transcriptome coordinates of the CS7
human embryo described in Tyser *et al.*, shown with the original annotations. The inset
(**i**) is used for expression analyses in (I) and (J).
(I,J) Expression of *pan*-amnion (*ISL1*, *TFAP2A* and *GATA3*) as well as late/advanced
amnion (*GABRP*, *VTCN1*, *IGFBP3*) markers, superimposed onto the Tyser *et al.* UMAP
plot.
(K-M) A UMAP plot showing the integrated single cell transcriptomes of d3 hPSC-
amnion and Tyser *et al.* CS7 embryo datasets, revealing close overlap of hPSC-amnion
cells with Tyser Epiblast, Primitive Streak and Ectoderm/Amnion cells (shown in (L) with
annotations in Tyser *et al.*, magnified in (M); AmP and advAm coloring is identical to
(C)). Note that a small number of hPSC-amnion cells show an overlap with Advanced
Mesoderm cells, possibly due to cardiac differentiation triggered by high BMP input.
482
Figure 2. Validation of selected advAm markers in a cynomolgus macaque
embryo staged between CS12 and CS13
(A) A dotplot of selected known and previously unrecognized late/advanced amnion
markers.
(B,C) CS12/13 cynomolgus macaque embryo sections were stained for indicated
markers (B: *GABRP* – green, *SESN3* – red, *ISL1* – magenta; C: *ADAMTS18* – green,
SESN3 – red, *ISL1* – magenta) using RNA *in situ* hybridization. (i) – (vii) indicate insets:
individual channels are shown in gray scale to aid visualization (from left to right:
GABRP, SESN3, ISL1 and merge in (B), *ADAMTS18, SESN3, ISL1* in (C)), and dotted
lines indicate a layer of amniotic epithelium and mesenchyme. Most amnion cells are
positive for *GABRP*, *SESN3* and *ISL1*. *ADAMTS18* expression is restricted to a fraction
of cells. In (iv), there are four *ADAMTS18*high cells, while *ADAMTS18* is largely
undetected in the amnion in (v). In (vi) and (vii), most cells are *ADAMTS18*high. Fig. S3E
provides a bird's-eye view of the CS12/13 embryo sample in (B) in which uterus and
placenta are readily seen.
FB; forebrain, HB, hindbrain; FG, foregut; C; cardiac tissue; NT, neural tube; S, somite.
* indicates an *ISL1*+ tissue that is likely of genital tubercle lineage. Scale bars = 500μm.
500

Figure 3. CLDN10 is an AmP marker expressed in the amnion-epiblast boundary cells of the PASE model as well as cynomolgus macaque peri-gastrula
(A) A dotplot of selected AmP markers.
(B) Expression of *CLDN10* superimposed onto the d3 hPSC-amnion as well as in Tyser *et al.* CS7 human embryo UMAP plots.
(C) Confocal micrographs of developing hPSC-amnion harvested at indicated timepoints, stained with indicated markers. A white arrowhead indicates a CLDN10/TFAP2A double-positive cell. CLDN10 expression is extinguished by d4. Membrane was stained using wheat germ agglutinin.
(D) Optical sections of PASE stained with indicated markers. White arrowheads indicate CLDN10 staining at the amnion-epiblast boundary.
(E,F) Confocal images of cynomolgus macaque embryos staged at CS6/7 (E) and CS7 (F) stained using indicated markers. CLDN10 is expressed at the transitioning boundary cells between the amnion and the epiblast. Insets indicate magnified regions. White arrowheads in (F – ii) indicate selected boundary cells with membrane CLDN10 staining. Clear membrane CLDN10 staining is not detected on the anterior side of the boundary cells (F – iii). Fig. S4H provides a bird's-eye view of the CS7 embryo in (F) with uterus and placental tissues. Scale bars = 20μm (C,D), 200μm (E), 500μm (F).

Figure 4. CLDN10 labels the amnion-epiblast boundary cells of cynomolgus macaque embryos undergoing the initial stages of organogenesis
(A-D) Confocal images of cynomolgus macaque embryos staged at CS8 (A,B) and CS9 (C,D) stained using indicated markers. Insets indicate magnified regions. CLDN10 is expressed at the anterior and posterior amnion-embryo boundaries in the CS8 embryo displaying the head-fold formation (A,B). The anterior head-fold region is magnified in (ii in (A) and (B)). White arrowheads indicate a squamous cell with a clear membrane CLDN10 staining (A - i). Brackets in (A) and (B) indicate regions with membrane CLDN10 staining. In the CS9 embryo (early organogenesis stage in (C) and (D)), while some posterior boundary CLDN10 staining is seen (seen in (iii)), the anterior CLDN10 staining is highly prominent at the amnion-anterior surface ectoderm border (in (i)). CLDN10 is also enriched in the dorsal foregut labeled by ISL1/2 and SOX2 but not SOX17 (indicated by * in (ii)). Fig. S4I and S4J provide bird's-eye views of the CS8 ((A) and (B)) and CS9 ((C) and (D)) embryos. HT, heat tube; FB, forebrain; YS, yolk sac. White arrowheads indicate the foregut pocket, while white arrows indicate the hindgut pocket. Bracket indicates somites in (C) and (D). Scale bars = 200μm (A,B), 500μm (C,D).

Figure S1.
(A) A UMAP plot displaying the single cell transcriptomes of 379 d3 hPSC-amnion cells with two identified clusters (AmP – sage color; advAm – salmon color).
(B) Predicted cell cycle (default parameter in Seurat) superimposed onto the d3 amnion UMAP plot.
(C) Venn diagrams showing a fraction of G1, G2/M or S cells in AmP (left) and advAm (right).

Figure S2.

547 (A) A UMAP plot displaying the original single cell transcriptome coordinates of the CS7
548 human embryo described in Tyser *et al.*, shown with the original annotations.
549 (B,C) Expression of indicated *pan*-amnion (B) and late/advanced amnion (C) markers
550 superimposed onto the UMAP plot.
551
Figure S3.
(A,B) UMAP plots displaying the Tyser *et al.* CS7 human embryo and d3 hPSC-amnion
single cell RNA sequencing integrated dataset (cell annotations shown in A, and sample
type annotations shown in B)
(C,D) Expression of indicated *pan*-amnion (C) and late/advanced amnion (D) markers
superimposed onto the UMAP plot.
(E) An overview image of the CS12/13 embryo implantation site shown in Fig. 2B. Only
nuclear staining is shown in grayscale. Scale bar = 2mm.
560
Figure S4.
(A) Expression of CLDN10 superimposed onto the uncropped Tyser *et al.* CS7 human
embryo UMAP plot.
(B) A UMAP plot displaying the GD14 cynomolgus macaque single cell transcriptome
with six identified general clusters (epiblast light blue; transition – purple; amnion – gold;
mesoderm – green; endoderm – teal; extraembryonic mesoderm – salmon). Arrow
indicates the likely trajectory of amnion differentiation based on the data in Yang *et al.*
as well as the expression analysis in (C). Note that, although not identical to the
published UMAP plot in Yang *et al.*, general characteristics are well recapitulated.
Trophectoderm cells have been omitted as performed in Yang *et al.*.
(C) Expression of indicated markers superimposed onto the Yang *et al.* UMAP plot.
Advanced amnion markers (EPAS1 and GABRP) are expressed at the tip of the
TFAP2A and ISL1 double-positive amnion cluster. CLDN10 positive cells show a weak
TBXT expression level.
(D,E) Individual channels of the whole embryo images in Fig. 4A and 4B, respectively.
(F,G) Individual channels of the whole embryo images in Fig. 4C and 4D, respectively.
(H-J) Overview images of the CS7 (H, from Fig. 3F), CS8 (I, Fig. 4A) and CS9 (J, Fig.
4C) embryo implantation sites (DNA signal shown in gray scale, Scale bar = 2mm).
579
580 **Table 1. Lists of differentially expressed genes in AmP (493) and advAm (563)**
581 **clusters**
582
583 **Table 2. List of antibodies and RNAscope probes used in this study**
584

ACKNOWLEDGEMENTS
We thank the Wisconsin National Primate Research Center (WNPRC) Veterinary,
Scientific Protocol Implementation, Pathology and Animal Services staff for providing
animal care, and assisting in procedures including breeding, pregnancy monitoring, and
sample collection. The contents of this manuscript are solely the responsibility of the
authors and do not represent the official views of the NIH.
Funding: This work was supported by NIH grants R01-HD098231, R01-DK089933,
P51 OD011106 (to the WNPRC) as well as by MCW CBNA Start-up funds. Specifically,

R01-DK089933 and MCW CBNA Start-up funds were used to perform experiments
using PASE. **Competing interests:** The authors declare no competing interest. **Data**
and materials availability: All data needed to evaluate the conclusions in the paper are
present in the paper and the Supplementary Materials. The raw data, unfiltered count
matrix and processed count matrix will be available to the database of Genotypes and
Phenotypes (dbGaP) upon publication. Further information and requests for reagents
may be directed to Kenichiro Taniguchi.
600

REFERENCES

Anderson, R.M., Lawrence, A.R., Stottmann, R.W., Bachiller, D., and Klingensmith, J.
(2002). Chordin and noggin promote organizing centers of forebrain development in the
mouse. Development *129*, 4975-4987.
Angelow, S., Ahlstrom, R., and Yu, A.S. (2008). Biology of claudins. Am J Physiol Renal
Physiol *295*, F867-876.
Arnold, S.J., and Robertson, E.J. (2009). Making a commitment: cell lineage allocation
and axis patterning in the early mouse embryo. Nature reviews Molecular cell biology
10, 91-103.
Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA
velocity to transient cell states through dynamical modeling. Nature biotechnology *38*,
1408-1414.
Bergmann, S., Penfold, C.A., Slatery, E., Siriwardena, D., Drummer, C., Clark, S.,
Strawbridge, S.E., Kishimoto, K., Vickers, A., Tewary, M*., et al.* (2022). Spatial profiling
of early primate gastrulation in utero. Nature *609*, 136-143.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species.
Nature biotechnology *36*, 411-420.
Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos,
S., Christiansen, L., Steemers, F.J*., et al.* (2019). The single-cell transcriptional
landscape of mammalian organogenesis. Nature *566*, 496-502.
Carleton, A.E., Duncan, M.C., and Taniguchi, K. (2022). Human epiblast
lumenogenesis: From a cell aggregate to a lumenal cyst. Seminars in cell &
developmental biology *131*, 117-123.
Chen, K., Zheng, Y., Xue, X., Liu, Y., Resto Irizarry, A.M., Tang, H., and Fu, J. (2021).
Branching development of early post-implantation human embryonic-like tissues in 3D
stem cell culture. Biomaterials *275*, 120898.
Dobreva, M.P., Lhoest, L., Pereira, P.N., Umans, L., Camus, A., Chuva de Sousa
Lopes, S.M., and Zwijsen, A. (2012). Periostin as a biomarker of the amniotic
membrane. Stem Cells Int *2012*, 987185.
Enders, A.C., Hendrickx, A.G., and Schlafke, S. (1983). Implantation in the rhesus
monkey: initial penetration of endometrium. The American journal of anatomy *167*, 275-
298.
Enders, A.C., Schlafke, S., and Hendrickx, A.G. (1986). Differentiation of the embryonic
disc, amnion, and yolk sac in the rhesus monkey. The American journal of anatomy
177, 161-185.
Geng, X., and Oliver, G. (2009). Pathogenesis of holoprosencephaly. J Clin Invest *119*,
1403-1413.

Hamed, M.M., Taniguchi, K., and Duncan, M.C. (2023). Monitoring Effects of Membrane
Traffic Via Changes in Cell Polarity and Morphogenesis in Three-Dimensional Human
Pluripotent Stem Cell Cysts. Methods in molecular biology *2557*, 83-98.
Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee,
M.J., Wilk, A.J., Darby, C., Zager, M*., et al.* (2021). Integrated analysis of multimodal
single-cell data. Cell *184*, 3573-3587 e3529.
La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V.,
Lidschreiber, K., Kastriti, M.E., Lonnerberg, P., Furlan, A*., et al.* (2018). RNA velocity of
single cells. Nature *560*, 494-498.
Miki, T., Lehmann, T., Cai, H., Stolz, D.B., and Strom, S.C. (2005). Stem cell
characteristics of amniotic epithelial cells. Stem cells *23*, 1549-1559.
Muenke, M., and Beachy, P.A. (2000). Genetics of ventral forebrain development and
holoprosencephaly. Current opinion in genetics & development *10*, 262-269.
Nakamura, T., Okamoto, I., Sasaki, K., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y.,
Nakamura, S., Yamamoto, T., and Saitou, M. (2016). A developmental coordinate of
pluripotency among mice, monkeys and humans. Nature *537*, 57-62.
Nakamura, T., Yabuta, Y., Okamoto, I., Sasaki, K., Iwatani, C., Tsuchiya, H., and
Saitou, M. (2017). Single-cell transcriptome of early embryos and cultured embryonic
stem cells of cynomolgus monkeys. Scientific data *4*, 170067.
Nishioka, N., Nagano, S., Nakayama, R., Kiyonari, H., Ijiri, T., Taniguchi, K., Shawlot,
W., Hayashizaki, Y., Westphal, H., Behringer, R.R*., et al.* (2005). Ssdp1 regulates head
morphogenesis of mouse embryos by activating the Lim1-Ldb1 complex. Development
132, 2535-2546.
Overeem, A.W., Chang, Y.W., Moustakas, I., Roelse, C.M., Hillenius, S., Helm, T.V.,
Schrier, V.F.V., Goncalves, M., Mei, H., Freund, C*., et al.* (2023). Efficient and scalable
generation of primordial germ cells in 2D culture using basement membrane extract
overlay. Cell Rep Methods *3*, 100488.
Roost, M.S., van Iperen, L., Ariyurek, Y., Buermans, H.P., Arindrarto, W., Devalla, H.D.,
Passier, R., Mummery, C.L., Carlotti, F., de Koning, E.J*., et al.* (2015). KeyGenes, a
Tool to Probe Tissue Differentiation Using a Human Fetal Transcriptional Atlas. Stem
cell reports *4*, 1112-1124.
Sasaki, K., Nakamura, T., Okamoto, I., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y.,
Nakamura, S., Shiraki, N., Takakuwa, T*., et al.* (2016). The Germ Cell Fate of
Cynomolgus Monkeys Is Specified in the Nascent Amnion. Developmental cell *39*, 169-
185.
Sekulovski, N., Wettstein, J.C., Carleton, A.E., Taniguchi, L.E., Ma, X., Rao, S.,
Schmidt, J.K., Golos, T.G., Lin, C.-W., and Taniguchi, K. (2023). Temporally resolved
early BMP-driven transcriptional cascade during human amnion specification. eLife *12:*
RP89367.
Shahbazi, M.N., and Zernicka-Goetz, M. (2018). Deconstructing and reconstructing the
mouse and human early embryo. Nature cell biology.
Shao, Y., and Fu, J. (2022). Engineering multiscale structural orders for high-fidelity
embryoids and organoids. Cell stem cell *29*, 722-743.
Shao, Y., Taniguchi, K., Gurdziel, K., Townshend, R.F., Xue, X., Yong, K.M.A., Sang, J.,
Spence, J.R., Gumucio, D.L., and Fu, J. (2017a). Self-organized amniogenesis by

human pluripotent stem cells in a biomimetic implantation-like niche. Nature materials
16, 419-425.
Shao, Y., Taniguchi, K., Townshend, R.F., Miki, T., Gumucio, D.L., and Fu, J. (2017b).
A pluripotent stem cell-based model for post-implantation human amniotic sac
development. Nature communications *8*, 208.
Shawlot, W., Wakamiya, M., Kwan, K.M., Kania, A., Jessell, T.M., and Behringer, R.R.
(1999). Lim1 is required in both primitive streak-derived tissues and visceral endoderm
for head formation in the mouse. Development *126*, 4925-4932.
Taniguchi, K., Heemskerk, I., and Gumucio, D.L. (2019). Opening the black box: Stem
cell-based modeling of human post-implantation development. The Journal of cell
biology *218*, 410-421.
Taniguchi, K., Shao, Y., Townshend, R.F., Tsai, Y.H., DeLong, C.J., Lopez, S.A.,
Gayen, S., Freddo, A.M., Chue, D.J., Thomas, D.J.*, et al.* (2015). Lumen Formation Is
an Intrinsic Property of Isolated Human Pluripotent Stem Cells. Stem cell reports *5*, 954-
962.
Ton, M.N., Keitley, D., Theeuwes, B., Guibentif, C., Ahnfelt-Ronne, J., Andreassen,
T.K., Calero-Nieto, F.J., Imaz-Rosshandler, I., Pijuan-Sala, B., Nichols, J.*, et al.* (2023).
An atlas of rabbit development as a model for single-cell comparative genomics. Nature
cell biology *25*, 1061-1072.
Townshend, R.F., Shao, Y., Wang, S., Cortez, C.L., Esfahani, S.N., Spence, J.R.,
O'Shea, K.S., Fu, J., Gumucio, D.L., and Taniguchi, K. (2020). Effect of Cell Spreading
on Rosette Formation by Human Pluripotent Stem Cell-Derived Neural Progenitor Cells.
Front Cell Dev Biol *8*, 588941.
Tyser, R.C.V., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and Srinivas,
S. (2021). Single-cell transcriptomic characterization of a gastrulating human embryo.
Nature *600*, 285-289.
Yang, R., Goedel, A., Kang, Y., Si, C., Chu, C., Zheng, Y., Chen, Z., Gruber, P.J., Xiao,
Y., Zhou, C.*, et al.* (2021). Amnion signals are essential for mesoderm formation in
primates. Nature communications *12*, 5126.
Yang, Y.P., Anderson, R.M., and Klingensmith, J. (2010). BMP antagonism protects
Nodal signaling in the gastrula to promote the tissue interactions underlying mammalian
forebrain and craniofacial patterning. Hum Mol Genet *19*, 3030-3042.
Zhao, C., Reyes, A.P., Schell, J.P., Weltner, J., Ortega, N.M., Zheng, Y., Björklund,
Å.K., Rossant, J., Fu, J., Petropoulos, S.*, et al.* (2021). Reprogrammed blastoids
contain amnion-like cells but not trophectoderm. bioRxiv, 2021.2005.2007.442980.
Zheng, Y., Xue, X., Shao, Y., Wang, S., Esfahani, S.N., Li, Z., Muncie, J.M., Lakins,
J.N., Weaver, V.M., Gumucio, D.L.*, et al.* (2019). Controlled modelling of human
epiblast and amnion development using stem cells. Nature *573*, 421-425.
Zheng, Y., Yan, R.Z., Sun, S., Kobayashi, M., Xiang, L., Yang, R., Goedel, A., Kang, Y.,
Xue, X., Esfahani, S.N.*, et al.* (2022). Single-cell analysis of embryoids reveals lineage
diversification roadmaps of early human development. Cell stem cell *29*, 1402-1419
e1408.
726

Figure 1

Figure 2

Figure 3

Figure 4