Temporally resolved single cell transcriptomics in a human model of

amniogenesis

AUTHORS: Nikola Sekulovski[1,¶], Amber E. Carleton[1,¶], Anusha A. Rengarajan[1,¶],

Chien-Wei Lin[2, ¶], Lauren N. Juga[1], Allison E. Whorton[1], Jenna K. Schmidt[3], Thaddeus

G. Golos[3,4,5], and Kenichiro Taniguchi[1,6,7,†]

AFFILIATIONS:

[1]Department of Cell Biology, Neurobiology and Anatomy, Medical College of Wisconsin,

Milwaukee, WI 53226, USA

[2]Division of Biostatistics, Institute for Health and Equity, Medical College of Wisconsin,

Milwaukee, WI 53226, USA

[3]Wisconsin National Primate Research Center (WNPRC), Madison, WI, USA

[4]Department of Obstetrics and Gynecology, University of Wisconsin - Madison School of

Medicine, Madison, WI USA

[5]Department of Comparative Biosciences, University of Wisconsin - Madison School of

Veterinary Medicine, Madison, WI, USA

[6]Department of Pediatrics, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[7]Cancer Center, Medical College of Wisconsin, Milwaukee, WI 53226, USA

[¶]These authors contributed equally to this work

[†]Corresponding author

CONTACT:

Kenichiro Taniguchi

8701 Watertown Plank Road

B4245 BSB CB062

Milwaukee, WI 53226

ktaniguchi@mcw.edu

ABSTRACT

Amniogenesis is triggered in a collection of pluripotent epiblast cells as the human

embryo implants. To gain insights into the critical but poorly understood transcriptional

machinery governing amnion fate determination, we examined the evolving

transcriptome of a developing human pluripotent stem cell-derived amnion model at the

single cell level. This analysis revealed several continuous amniotic fate progressing

states with state-specific markers, which include a previously unrecognized CLDN10[+]

amnion progenitor state. Strikingly, we found that expression of CLDN10 is restricted to

the amnion-epiblast boundary region in the human post-implantation amniotic sac

model as well as in a peri-gastrula cynomolgus macaque embryo, bolstering the

growing notion that, at this stage, the amnion-epiblast boundary is a site of active

amniogenesis. Bioinformatic analysis of published primate peri-gastrula single cell

sequencing data further confirmed that CLDN10 is expressed in cells progressing to

amnion. Additionally, our loss of function analysis shows that CLDN10 promotes

amniotic but suppresses primordial germ cell-like fate. Overall, this study presents a

comprehensive amniogenic single cell transcriptomic resource and identifies a

previously unrecognized CLDN10[+] amnion progenitor population at the amnion-epiblast

boundary of the primate peri-gastrula.

INTRODUCTION

Amniogenesis is initiated during implantation in humans, and leads to the formation of an amniotic sac structure that surrounds and protects the developing embryo (Enders et al., 1983; Enders et al., 1986; Miki et al., 2005; Sasaki et al., 2016; Shahbazi and Zernicka-Goetz, 2018; Taniguchi et al., 2019). At implantation, the embryo (referred to at this time as a blastocyst) contains three morphologically and molecularly distinct cell types: 1) a collection of unpolarized pluripotent epiblast cells, precursors to both the embryo proper and the amniotic ectoderm, 2) a surrounding layer of polarized trophectoderm, a placental tissue precursor, and 3) an underlying extraembryonic primitive endoderm, a yolk sac precursor. Upon implantation, the pluripotent epiblast cells initiate epithelial polarization to form a cyst with a central lumen, the future amniotic cavity (Carleton et al., 2022; Shao and Fu, 2022). This event is followed by the fate transition of pluripotent epiblast cells that are in close proximity to the uterus to squamous amniotic ectoderm, forming a clear boundary between amnion and pluripotent epiblast territories of the cyst (Shahbazi and Zernicka-Goetz, 2018; Shao and Fu, 2022; Taniguchi et al., 2019). This asymmetric amnion-epiblast structure, known as the amniotic sac, provides the foundation for the next essential developmental steps (e.g., primitive streak formation, neural specification).

Recent studies using early human and monkey embryos have provided a basic understanding of the transcriptomic characteristics of early primate amniogenesis (Bergmann et al., 2022; Nakamura et al., 2016; Nakamura et al., 2017; Sasaki et al., 2016; Tyser et al., 2021; Yang et al., 2021). Yet, experimental dissection of the molecular mechanisms involved in this process is difficult in these *in vivo* models. To

enable molecular and cellular investigations, we and others developed human

pluripotent stem cell (hPSC)-derived amnion models and showed that BMP signaling

plays a crucial role in initiating amniogenesis (Chen et al., 2021; Overeem et al., 2023;

Shao et al., 2017a; Shao et al., 2017b; Zheng et al., 2019). Recently, we further

explored the BMP-dependent amniogenic transcriptional cascade, identifying several

distinct transcriptional stages, as well as a requirement for TFAP2A-dependent

transcription in regulating amnion fate progression (Sekulovski et al., 2024). However,

changes in cellular differentiation states during amnion lineage progression remain

largely unexplored.

In this study, we used single cell RNA sequencing (scRNA-seq) analysis to

examine the dynamics in gene expression that accompany amnion differentiation of

hPSC (herein referred to as the hPSC-amnion) cultured in a soft gel environment, called

Gel-3D (Shao et al., 2017a). While no exogenous BMP is added in this Gel-3D amnion

model, BMP signaling is activated in the cells by a mechanosensitive cue provided by

the soft substrate, thereby initiating human amniogenesis (Shao and Fu, 2022; Shao et

al., 2017a; Taniguchi et al., 2019). Interestingly, our data reveal contiguous amniogenic

cell states: pluripotency-exiting, early progenitor, late progenitor, specified and maturing,

each of which shows transcriptional similarities to distinct cell types in a Carnegie stage

7 human embryo (Tyser et al., 2021). Our data also reveal the presence of two non-

amniotic cell types in the Gel-3D system: primordial germ cell-like and advanced

mesoderm-like cells. Importantly, we identify a cohort of markers specific to each of

these amnion lineage progressing states, and validate selected markers for their

expression in the amnion from cynomolgus macaque (*Macaca fascicularis*) embryos.

Strikingly, we show that CLDN10, a marker of the progenitor population, exhibits a

restricted amniotic expression pattern at the boundary between the amnion and the

epiblast of the cynomolgus macaque peri-gastrula. This further supports our findings

from a separate study (Sekulovski et al., 2024), which show that the amnion-epiblast

boundary is a site of active amniogenesis in the macaque peri-gastrula. Furthermore,

loss of CLDN10 results in the formation of primordial germ cell-like cells at the expense

of amnion cells. Together, this study provides important single cell-level insight into

human amnion fate progression and presents additional evidence for the presence of

amnion progenitor cells in primate embryos undergoing gastrulation.

RESULTS

Previously, we described culture conditions (Gel-3D) in which plated hPSC form

polarized cysts initially composed of columnar pluripotent cells, which then

spontaneously undergo squamous morphogenesis and begin to express amnion

markers (Shao et al., 2017a). Specifically, singly dissociated hPSC are plated densely

on a soft gel substrate to form aggregates and are allowed to grow for 24 hours (hrs) in

the presence of a ROCK inhibitor (Y-27632). After the first 24 hrs (day 1, d1), the ROCK

inhibitor is removed (a trigger to initiate lumenal cyst formation (Hamed et al., 2023;

Shao et al., 2017a; Taniguchi et al., 2015)), and a diluted extracellular matrix gel overlay

(2% v/v, e.g., Geltrex) is added to provide an additional 3D cue. Immunofluorescent (IF)

staining at this time shows cuboidal cells surrounding a central lumen; these cells

express NANOG, but not TFAP2A (**Fig. 1A**).

Interestingly, 24 hrs after adding the gel overlay (d2), a subset of cells within the

cysts displays reduced expression of NANOG, while TFAP2A expression becomes

weakly activated (**Fig. 1A**, arrowhead in d2). By d3, NANOG is no longer detectable.

Moreover, while many cells exhibit prominent TFAP2A expression, some cells are only

weakly positive for TFAP2A (**Fig. 1A**, arrowheads). This pattern of TFAP2A expression

is consistent with the previous observation that amniogenesis initiates focally on the

pluripotent cysts and then spreads laterally to form fully squamous cysts (**Fig. 1A**,

(Sekulovski et al., 2024; Shao et al., 2017a; Shao et al., 2017b)). These molecular

changes are accompanied by a decrease in nuclear aspect ratio (a measure of

epithelial cell morphology) over time, revealing more flattened (squamous) shapes by

d3 and d4 (**Fig. 1B**, quantitation strategy shown in **Fig. S1A**, and previously established

in Shao et al., 2017a; Townshend et al., 2020)). These results provide further molecular

and structural evidence that the transition from pluripotent to amnion cell types occurs

progressively over the cyst, starting from focal initiation sites.

To explore transcriptional programs of the fate transitioning cells during early

stages of human amnion development in this model, we performed time-course scRNA-

seq analysis of developing hPSC-amnion cysts, harvested at d1, d2, d3 and d4. hPSC-

amnion cysts were dissociated into single cells, followed by single cell isolation and

labeling using the 10x Genomics Chromium system, cDNA amplification and

sequencing followed by demultiplexing (see Material and Methods for additional details).

This scRNA-seq dataset was examined using Seurat (Butler et al., 2018) for data

filtering, regression for genes associated with cell cycle progression, normalization,

variable gene selection and subsequent unsupervised clustering of cells (see Materials

and Methods). The data yield seven distinct populations among 8,765 cells (d1 = 1,359

(salmon), d2 = 1,546 (sage), d3 = 2,352 (light blue), d4 = 3,508 cells (purple), **Fig. 1C**),

which, after marker and lineage analyses described below, are labeled pluripotency-

exiting (salmon color, 1,363 cells), early progenitor (brown, 479 cells), late progenitor

(sage, 1,045 cells), specified (green, 1,366 cells) and maturing (purple, 2,795 cells),

PGC-LC (primordial germ cell-like cells, blue, 1,055 cells) and advM-LC (advanced

mesoderm-like cells, magenta, 662, **Fig. 1D**). Transcriptomic features of these cells are

visualized in Uniform Manifold Approximation and Projection (UMAP) plots (**Fig. 1C,D**).

To broadly characterize the transcriptional state of each population, we examined

the expression of known amnion markers (Roost et al., 2015; Shao et al., 2017a; Yang

et al., 2021; Zhao et al., 2024; Zheng et al., 2022). Consistent with the amniotic lineage,

early amnion markers (*ID2*, *GATA3* and *TFAP2A*, **Fig. 1E**, expression superimposed on

the UMAP plot in **Fig. 1C**) are widely expressed starting at d1 in pluripotency-exiting

cells that also show broad expression of pluripotency markers (**Fig. S1B**), while

intermediate markers that label specified amnion (*ISL1*, *HAND1*, *DLX5*, (Sekulovski et

al., 2024; Yang et al., 2021; Zhao et al., 2024)) show abundant expression by d3 (**Fig.**

1F). Late amnion genes such as *GABRP*, *VTCN*1 and *IGFBP3* (expressed in the first-

trimester human amnion (Roost et al., 2015) and also in other human amnion models

(Sekulovski et al., 2024; Shao et al., 2017a; Yang et al., 2021; Zhao et al., 2024; Zheng

et al., 2022)) are primarily enriched in the d3 and d4 populations (**Fig. 1G**). Interestingly,

most of the d2 cells show the expression of early, but not specified or pluripotency

(SOX2, NANOG), genes, suggesting that the d2 cell population may contain

transitioning cell types that give rise to specified amnion (brown and sage cells in **Fig.**

1D). Next, RNA-velocity analysis (Bergen et al., 2020) was applied to examine lineage

relationships based on the relative abundance of unspliced and spliced mRNA in each

cell. Indeed, the majority of the RNA-vector trajectories are directed from the cells in the

early progenitor (brown), late progenitor (sage), to specified (green) states (**Fig. 1H**).

In a recent study, we performed a detailed meta-analysis of the Tyser *et al.*

scRNA-seq dataset from a human embryo staged at Carnegie stage 7 (CS7) (Tyser et

al., 2021) and showed that the rod-shaped cluster of cells annotated as "Ectoderm" is

comprised of lineage progressing, lineage committed, and fully differentiated amnion

cell types (Sekulovski et al., 2024). The Tyser *et al*. UMAP plot is reproduced in **Fig.**

S1C with the original coordinates and annotations (expression of late markers, *GABRP*,

*VTCN*1 and *IGFBP3*, found at the distal tip of the rod-shaped "Ectoderm" population,

Fig. S1C-E). To examine which cells in the CS7 human embryo share transcriptomic

similarities to amnion progressing cells in Gel-3D, the scRNA-seq datasets from the

Tyser *et al.* CS7 human embryo and our d1-d4 Gel-3D time-course samples were

combined and normalized using an integration feature based on canonical correlation

analysis in Seurat (Hao et al., 2021) (**Fig. 1I-K**). Although the combined UMAP plot of

this integrated dataset shows some changes in shape from our original UMAP, it is clear

that "Epiblast", "Primitive Streak" and "Ectoderm" populations in the Tyser *et al.* dataset

closely map to the amnion progressing populations in our time-course Gel-3D dataset

(**Fig. 1I**, see **Fig. S1F** for mapping of each population). Indeed, the amniotic Tyser

"Ectoderm" cells primarily overlap with Gel-3D "specified" and "maturing" cell

populations, consistent with their amniotic fate; most Tyser "Epiblast" cells are seen in

the "pluripotency-exiting" population (**Fig. 1J,K**, **Fig. S1F**). Interestingly, Tyser "Primitive

Streak" cells are populated across several Gel-3D amniogenic cell states (**Fig. 1J,K**,

Fig. S1F), overlapping with the early (brown) and late (sage) progenitor populations,

suggesting that some Tyser "Primitive Streak" cells may be actively transitioning, and

that our early and late progenitor cells may display transcriptomic characteristics of fate

transitioning cells in human peri-gastrula.

Additionally, we identify two non-amniotic cell types that show transcriptional

characteristics of PGC- (blue) and advM- (magenta) like cells (**Fig. 1D**). The Tyser-Gel-

3D integration analysis shows that the blue cells overlap with the Tyser "Primitive

Streak" cells that abundantly express PGC markers (e.g., *PRDM1*, *SOX17*, *NANOS3*,

PRDM14, *NANOG*, *XACT*, **Fig. 1L**). Several Tyser "Advanced Mesoderm" cells are

seen in the magenta cells, and our marker analysis shows that developed mesoderm

makers (*HAND2*, *GATA4*, *PITX2*, *ISL1*, *HAND1*) are enriched (**Fig. 1M**, summary of

additional markers found in **Fig. 1N**). Strikingly, the RNA velocity trajectories are

directed from the interface of the early/late amnion progenitor populations to the PGC-

LC (**Fig. 1H**), suggesting their close transcriptomic characteristics. These findings

support a growing notion that amnion and PGC progressing cells initially share a

common intermediate lineage (Castillo-Venzor et al., 2023; Chen et al., 2019; Xiao et

al., 2024; Zheng et al., 2022). The advM-LC population may have formed due to high

BMP signaling (Schultheiss et al., 1997; Tsaytler et al., 2023; van Wijk et al., 2007).

To identify genes that are unique to each cluster, unsupervised differential gene

enrichment analysis was performed (**Table 1**, adjusted p-value < 0.05). As expected,

pluripotency markers are seen in the pluripotency-exiting cluster, and the specified and

maturing clusters show enriched expression of known late amnion markers as well as

previously uncharacterized genes (*ADAMTS18*, *SESN3* (**Fig. 1N**) – RNA *in situ*

hybridization in a cynomolgus macaque (*Macaca fascicularis*) embryo staged between

CS12 and CS13, an organogenesis stage, shown in **Fig. S2A-C**).

The most differentially expressed genes in early (brown) and late (sage)

progenitor populations are *TBXT* (T-box transcription factor T) and *CLDN10* (a gene

encoding Claudin-10, a component of the tight junction), respectively (**Fig. 2A,B**, top

three genes plotted in **Fig. 2C**). Recently, we showed that amnion lineage progression

in Glass-3D^{+BMP}, another model of human amniogenesis, traverses an intermediate

transcriptional phase weakly expressing *TBXT* (*TBXT*^{low}) before specified markers (e.g.,

ISL1, *HAND1*, *DLX5*) are expressed, revealing lineage progressing characteristics of

the cells in the *TBXT*^{low} transcriptional phase (Sekulovski et al., 2024). The same study

also showed that cells displaying transcriptomic characteristics consistent with fate

progression from *TBXT*^{low} to *ISL1*$^+$/*HAND1*$^+$ specified state are also seen in the Tyser *et*

al. CS7 human embryo as well in the Yang *et al.* d14 cynomolgus macaque embryo

datasets (Sekulovski et al., 2024). In Gel-3D, while most cells in the early progenitor

population express some level of *TBXT*, only a few cells express abundant *TBXT* (**Fig.**

2A). Indeed, our IF time-course analysis shows that cells with weak TBXT expression

are seen at d2 in several cells that are not yet expressing ISL1 (**Fig. 2D**, arrowheads);

TBXT expression is largely missing in d3 and d4 cysts (**Fig. 2D**), confirming the

presence of an early and transient *TBXT*^{low} state in developing Gel-3D hPSC-amnion

cysts.

Prominent *CLDN10* expression is seen throughout the late progenitor population

(*CLDN10*^{high}), but is rapidly diminished in the specified cells (**Fig. 2B**). Although at lower

levels, some *CLDN10* expression is observed in the pluripotency-exiting and early

progenitor states (*CLDN10*^{low}, **Fig. 2B,C**), consistent with our previous bulk-RNA

sequencing analysis showing that low *CLDN10* expression is seen in pluripotent cells

(Shao et al., 2017a). Using IF staining in Gel-3D hPSC-amnion model, we validated a

dynamic expression pattern of CLDN10 over time (**Fig. 2D,E**). Despite broadly

expressed at transcript levels in hPSC-amnion cysts at d2 (**Fig. 2B**), CLDN10

expression is seen in the small fraction of cells that are weakly positive for TBXT and

TFAP2A (**Fig. 2D,E**, arrowheads, quantitation in **Fig. 2F**). At d3, CLDN10 is co-

expressed with TFAP2A in several, but not all cells (**Fig. 2E**). Further, ISL1^{high} cells do

not show clear CLDN10 membrane staining (**Fig. 2D**). In contrast, most cells at d4

retain abundant TFAP2A and ISL1 expression but do not express CLDN10 (**Fig. 2D,E**).

Importantly, our analysis for nuclear aspect ratio (NAR) shows that, at d2, CLDN10^{+}

cells (mean NAR = 0.54 ± 0.17 STDEV, n = 39) are significantly more squamous

compared to CLDN10^{-} cells (NAR = 0.97 ± 0.31 STDEV, n = 87, p<0.001). While

previous studies implicated *CLDN10* as an amnion marker (Zheng et al., 2019; Zheng et

al., 2022), a detailed analysis was not performed. Thus, these data present evidence

suggesting that CLDN10^{high} marks a later transient progenitor state that follows an

earlier *TBXT*^{low}/*CLDN10*^{low} progenitor state, but CLDN10 expression is extinguished in

more differentiated cells. Single cell transcriptomic characteristics consistent with this

lineage progression from the *TBXT*^{low}/*CLDN10*^{low} state to the *CLDN10*^{high} state is also

seen in the Tyser *et al.* CS7 human embryo as well in the Yang *et al.* d14 cynomolgus

macaque embryo datasets (**Fig. S3A-C**).

Interestingly, in PASE (post-implantation amniotic sac embryoids), an *in vitro*

model of human amniotic sac formation (Shao et al., 2017b), CLDN10 is localized

specifically to cells at the boundary between squamous amnion and columnar

pluripotent cells (**Fig. 3A** , arrowheads). To test whether CLDN10$^+$ progenitor-like

boundary cells are present in the developing primate amniotic sac *in vivo*, we analyzed

CLDN10 expression in a cynomolgus macaque embryo displaying the amniotic sac

(**Fig. 3B**, staged between CS6 and CS7). Strikingly, CLDN10 expression is exclusively

seen at the boundary separating amnion and epiblast tissues. These results are

consistent with our recent findings indicating that this boundary is likely a site of active

amniogenesis (i.e., epiblast cells at the boundary actively undergo amnion specification)

in the primate peri-gastrula (Sekulovski et al., 2024). Importantly, a cynomolgus

macaque embryo from a later stage (**Fig. 3C,D**, CS10) also shows CLDN10 positive

cells at the amnion-embryo boundary. In the CS10 embryo, while a collection of cells

expressing weak CLDN10 is seen posteriorly (**Fig. 3C,D** – inset (iii)), CLDN10

expression is highly prominent anteriorly in the cells at the boundary of the amnion and

the developing mediolateral placode (**Fig. 3C,D** – inset (i), additional images in **Fig.**

S3D,E). Together, these results establish the presence of previously unrecognized

CLDN10$^+$ amnion-embryo boundary population that primarily gives rise to amnion

during gastrulation as well as during early organogenesis in the cynomolgus macaque.

To investigate the role of CLDN10, H9 hESC lacking CLDN10 were cultured in

the Gel-3D condition (**Fig. 4A-D**, see **Fig. S4** for generation of *CLDN10*-KO lines).

Morphologically, while *CLDN10*-KO cyst formation is largely similar to controls by d3, by

d4, squamous morphogenesis as well as overall cyst formation are disrupted.

Interestingly, our IF analysis shows that, in the KO background, SOX2 expression

gradually reduces overtime similar to controls, but cells expressing ISL1 are reduced by

d4 (**Fig. 4A,B**). Given the presence of PGC-LC in Gel-3D (**Fig. 1D,L**), we next

examined the expression of PGC markers. Strikingly, while very few cells are

NANOG$^+$/SOX17$^+$ PGC-LC in controls at d4, NANOG/SOX17 double positive cells are

seen prematurely at d2 as well as at d3 (**Fig. 4C,D**). These results suggest that

CLDN10 functions to promote amnion fate progression, and prevent emergence of the

PGC-like lineage.

DISCUSSION

Together, this study presents a temporally resolved single cell transcriptomic resource

for future investigations of the amniogenic transcriptional cascade in developing primate

embryos. In summary, we have profiled the transcriptomic signatures of lineage

progressing cells in an *in vitro* model of hPSC-amnion development, and identified 1)

five contiguous amnion progressing states (outlined in **Fig. 4E**) including two previously

unrecognized amnion progenitor populations, 2) an amnion progenitor population

labeled by CLDN10 at the boundary between the amnion and the epiblast of the primate

amniotic sac (model schematic in **Fig. 4F**), and 3) the role of CLDN10 in maintaining

amnion fate progression.

Our time-course scRNA-seq analysis showed that PGC-LC are present in our

Gel-3D model, and that some of the amnion progenitor population may contribute to

PGC-LC, supporting a growing notion that amnion and PGC progressing cells initially

share a common lineage (Castillo-Venzor et al., 2023; Chen et al., 2019; Xiao et al.,

2024; Zheng et al., 2022), and providing detailed transcriptomic insights into this lineage

diversification step that occurs in the Gel-3D system (**Fig. 4E**). Moreover, the loss of

CLDN10, a marker of the late amnion progenitor and a tight junction component, leads

to increased PGC-LC formation, while amnion formation is reduced. Interestingly, a

recent study by Vasic *et al.* showed that reduced level of TJP1 (also known as ZO-1) in

unconfined hPSC monolayer colonies treated with BMP4 (a system that undergoes

gastrulation-like patterning (Joy et al., 2021) similar to the micropatterned gastruloid

model, (Heemskerk et al., 2017; Warmflash et al., 2014) leads to increased emergence

of a germ cell lineage (Vasic et al., 2023). Therefore, our results present additional

evidence for the role of tight junction formation in suppressing PGC lineages, and

demonstrate a key role of CLDN10 in maintaining amnion fate progression in progenitor

cells. Given that Claudin proteins control tight junction properties and epithelial

characteristics (Angelow et al., 2008), CLDN10 might also be critical for organizing a

boundary structure containing shape/fate-transitioning cells flanked by squamous

amnion or columnar/pseudostratified pluripotent cells on each side.

Previous studies established that BMP is a major trigger of amniogenesis. What

other potential mechanisms could help to initially induce and maintain the half amnion-

half epiblast structure? Trophectoderm may play some roles in triggering amnion

specification given that these cells directly overlay the nascent amnion during

implantation. Indeed, molecular characterization of early cynomolgus macaque embryos

by Sasaki *et al.* (Sasaki et al., 2016) have suggested that the trophectoderm may

provide some cue (e.g., secreted ligands) to the underlying epiblast cells. Also, a recent

study by Pedroza *et al.* showed that co-culturing hPSC on a monolayer of human

trophoblast stem cells helps initiate amniogenesis (Pedroza et al., 2023).

Furthermore, visceral endoderm might play an important role in defining the zone

in which amniogenesis can occur, helping to shape the amnion-epiblast boundary

territory. Studies have established that visceral endoderm plays a major role in embryo

patterning by expressing secreted BMP antagonists (Chambers et al., 2009; Perea-

Gomez et al., 2002; Shawlot et al., 1999), even at very early stages of implantation

(Bergmann et al., 2022; Sasaki et al., 2016). Therefore, it is possible that, during

implantation, epiblast cells that are in close proximity to visceral endoderm contribute to

embryo proper because the presence of secreted BMP antagonists prevents them from

undergoing amniogenesis. However, more distal epiblast cells at the uterine-proximal

pole of the epiblast cyst, which are farther away from the BMP antagonist source, can

respond to BMP signaling and form amnion. The amnion-epiblast boundary territory

might be the most distal region at which amniogenesis can take place, and the balance

between the rate of epiblast proliferation and the rate of amniogenesis at the boundary

likely contributes to maintaining the ratio between amnion and epiblast cells in

implanting embryos, as well as in peri-gastrula.

Recently, single cell transcriptomics analysis has emerged as a valuable tool to

ground findings from *in vitro* models with natural as well as cultured primate embryo

systems (e.g., (Yang et al., 2021; Zhao et al., 2024; Zheng et al., 2022)). To aid in

additional investigations, several culture systems have been developed to generate

amnion, and single cell transcriptomic analyses have been performed in several of the

systems (e.g., (Chen et al., 2019; Minn et al., 2020; Overeem et al., 2023; Rostovskaya

et al., 2022)). Future transcriptomic comparisons of the amnion in each of these models

will enable us to gain additional insights into amniogenic mechanisms.

MATERIALS AND METHODS

hESC lines

Human embryonic stem cell line H9 was used in this study (WA09, P30, P48, WiCell;

NIH registration number: 0062). All protocols for the use of hPSC lines were approved

by the Human Stem Cell Research Oversight Committee at the Medical College of

Wisconsin and the Human Pluripotent Stem Cell Research Oversight Committee at the

University of Michigan. All hPSC lines were maintained in a feeder-free system for at

least 20 passages and authenticated as karyotypically normal at the indicated passage

number. Karyotype analysis was performed at Cell Line Genetics. All hPSC lines tested

negative for mycoplasma contamination (LookOut Mycoplasma PCR Detection Kit,

Sigma-Aldrich). In summary, hESC were maintained in a feeder-free culture system with

mTeSR1 medium, or with 50%/50% mix of mTeSR1 and mTeSR plus (STEMCELL

Technologies). hESC were cultured on 1% (v/v) Geltrex (Thermo Fisher Scientific), or

with Cultrex SCQ (Bio-techne) coated 6 well plates (Nunc). Cells were passaged as

small clumps every 4 to 5 days with Dispase (Gibco). All cells were cultured at 37°C

with 5% CO_2. Media was changed every day. hESC were visually checked every day to

ensure the absence of spontaneously differentiated, mesenchyme-like cells in the

culture. Minor differentiated cells were scratched off the plate under a dissecting scope

once identified. The quality of all hESC lines was periodically examined by

immunostaining for pluripotency markers and successful differentiation to three germ

layer cells. Similar methods were previously used in (Sekulovski et al., 2024; Wang et

al., 2021).

Gel-3D hPSC-amnion formation assays

Methods for these assays have been previously described (Shao et al., 2017a; Shao et

al., 2017b) with exceptions that 60-70uL of undiluted ECM gel solution is hand-streaked

on ice cold 22mmx22mm square coverslips, and that, in addition to Geltrex (Life

Technologies), Cultrex Ultimatrix (Bio-Techne) was also used. Singly dissociated cells

were prepared using Accutase (Sigma-Aldrich) and were plated on ECM gel-coated

coverslips at 25,000 cells/cm^2 in mTeSR1 or in 50%/50% mTeSR1/mTeSR plus mix

medium in the presence of 10µM Y-27632 (STEMCELL Technologies). After 24 hr, cells

were then incubated in media containing 2% Geltrex/Ultimatrix overlay without Y-27632

with daily media changes (note that 1% gel overlay was used between d2 to d4).

DNA constructs

piggyBac-CRISPR/Cas9 (pBACON) constructs

A piggyBac-CRISPR/Cas9 (pBACON) vector that contains SpCas9-T2A-puro and hU6-

gRNA expression cassettes flanked by piggyBac transposon terminal repeat elements

(pBACON-puro), which allows subcloning of annealed oligos containing gRNA

sequence at *BbsI* site, has been previously described (Shao et al., 2017b; Townshend

et al., 2020; Wang et al., 2021). gRNA targeting genomic sites and oligo sequences to

generate pBACON-puro-h*CLDN10*-ICL1 (primers: CRISPR_hCLDN10ICL#1_s and

CRISPR_hCLDN10ICL#1_as) are found in **Fig. S4** and **Table 2**; these were designed

using a publicly available tool

(https://www.idtdna.com/site/order/designtool/index/CRISPR_CUSTOM). Similar

methods were previously used in several publications (e.g., (Sekulovski et al., 2024;

Wang et al., 2021)).

piggyBac-based transgenic and genome edited hESC lines

PB constructs (3μg) and pCAG-ePBase (Lacoste et al., 2009) (1μg) were co-transfected

into H9 hESC (70,000 cells cm^{-2}) using GeneJammer transfection reagent (Agilent

Technologies). To enrich for successfully transfected cells, drug selection (puromycin,

2μg/mL for 4 days) was performed 48- to 72-hrs after transfection. hESC stably

expressing each construct maintained the expression of pluripotency markers.

During pBACON-based genome editing, puro-selected cells were cultured at low

density (300 cells cm^{-2}) for clonal selection. Established colonies were manually picked

and expanded for screening indel mutations using PCR amplification of a region

spanning the targeted gRNA region (genomic DNA isolated using DirectPCR Lysis

Reagent (Tail) (VIAGEN), primers: hCLDN10, Geno_hCLDN10CDS2_RI_fw and

Geno_hCLDN10CDS2_NI_rv), which were subcloned into pPBCAG-GFP (Chen and

LoTurco, 2012) at EcoRI and NotI sites, and sequenced (Seq-3'TR-pPB-Fw). At least

12 to 15 bacterial colonies were sequenced (Sanger sequencing) to confirm genotypic

clonality. Control cells are wild-type, unedited H9 hESC in all loss-of-function

experiments.

Cynomolgus macaque

Animals

The female and male cynomolgus macaques were housed and cared for at the Wisconsin

National Primate Research Center (WNPRC). All procedures were performed in

accordance with the NIH Guide for the Care and Use of Laboratory Animals and under

approval of the University of Wisconsin College of Letters and Sciences and Vice

Chancellor Office for Research and Graduate Education Institutional Animal Care and

Use Committee (protocol g005061).

Animal breeding and pregnancy detection

Beginning on day 8 post-onset of menses, the female was housed with a compatible

male and monitored for breeding. Animals were pair-housed until day 16-20 post-onset

of menses. A 2-3 mL blood draw was performed daily from day 8 post-onset of menses

until day 16 to assess the timing of ovulation based on the estradiol peak and rise in

progesterone in serum. Serum samples were analyzed for estradiol (25 μL) and

progesterone (20 µL) using a cobas e411 analyzer equipped with

ElectroChemiLuminescence technology (Roche, Basal, Switzerland) according to

manufacturer instructions. Results were determined via a calibration curve which was

instrument-generated by 2-point calibration using traceable standards and a master

curve provided via the reagent barcode. Inter-assay coefficient of variation (CV) was

determined by assaying aliquots of a pool of rhesus plasma. For estradiol, the limit of

quantitation (LOQ) was 25 pg/mL, the intra-assay CV was 2.02%, and the inter-assay

CV was 5.05%. For progesterone, the LOQ was 0.2 ng/mL, the intra-assay CV was

1.37%, and the inter-assay CV was 4.63%. A transabdominal ultrasound was performed

to detect pregnancy as early as 14 days post-ovulation. The ultrasound measurements

in combination with the timing of ovulation were used to estimate the day of conception

and gestational age of the pregnancy.

Terminal perfusion uterine collection, paraffin embedding, sectioning and staining

The pregnant females were sedated with intramuscular ketamine (>15 mg/kg) followed

by IV sodium pentobarbital (>35 mg/kg) and then perfused with 4% paraformaldehyde

(PFA) via the left ventricle. The entire uterus and cervix were removed. The serosa and

superficial myometrium were scored for better fixative penetration and to denote dorsal

and ventral orientation. Tissues were fixed in 4% PFA with constant stirring and solution

changes were performed every 24 hrs for a total of 72 hrs. The uterus was serially

sectioned from the fundus to the internal cervical os into 4 mm slices using a dissection

box. Cassettes with tissue sections were transferred into 70% ethanol, routinely

processed in an automated tissue processor and embedded in paraffin for histological

analyses (5 μm sections). Fixed uterine tissue were cut in 5 μm thick cross-sections,

mounted on slides, deparaffinized in xylene and rehydrated in an ethanol series. Antigen

retrieval was performed by boiling in citrate buffer. Sections were blocked 4% goat serum

in PBS at RT for at least 3-hr. Subsequent immunolocalization was performed using

commercially available primary antibodies, incubated overnight at 4 °C in 4% serum.

Immunofluorescent detection was performed using secondary antibodies tagged with a

fluorescent dye (fluorophores excitation = 488, 555, and 647nm), and counterstained with

DAPI. Negative controls were performed in which the primary antibody was substituted

with the same concentration of normal IgG of the appropriate isotype. Images were

obtained with a Zeiss LSM980 microscope.

Note that similar methods were used in (Sekulovski et al., 2024).

Confocal microscopy of fixed samples

Confocal Images of fixed samples were acquired using a Nikon-A1 (Nikon) and a Zeiss

LS980 laser scanning confocal microscopes. Non-3D images were generated using

Zen, FIJI (NIH) and Photoshop (Adobe).

Immunostaining

hPSC-amnion cysts grown on the coverslip were rinsed with PBS (Gibco) twice, fixed

with 4% paraformaldehyde (Sigma) for 60 min, then rinsed with PBS three times, and

permeabilized with 0.1% SDS (Sigma, in 1x PBS) solution for 60 min. The samples

were blocked in 4% heat-inactivated goat serum (Gibco) or 4% normal donkey serum

(Gibco) in PBS overnight at 4°C. The samples were incubated with primary antibody

solution prepared in blocking solution at 4°C for 48 hr, washed three times with PBS (30

min each), and incubated in blocking solution with goat or donkey raised Alexa Fluor-

conjugated secondary antibodies (Thermo Fisher), at room temperature for 24 hours.

Counter staining was performed using Hoechst 33342 (nucleus, Thermo Fisher

Scientific), Alexa-Fluor-conjugated wheat germ agglutinin (membrane, Thermo Fisher

Scientific) and Phalloidin (F-ACTIN, Thermo Fisher). All samples were mounted on

slides using 90% glycerol (in 1x PBS). When mounting hPSC-cysts samples, modeling

clay was used as a spacer between coverslip and slide to preserve lumenal cyst

morphology. Antibodies for IF staining are found in **Table 2**. Similar staining method

was previously used in (Shao et al., 2017a; Shao et al., 2017b).

RNA isolation and Single cell RNA sequencing

Gel-3D hPSC-amnion samples were rinsed once with cold 1x PBS prior to treating with

Cultrex Organoid Harvesting solution (R&D Systems) to non-enzymatically

depolymerize Ultimatrix (gel bed, R&D systems), which were then dissociated into

single cell suspensions using the Neural Tissue Dissociation Kit (P) (Miltenyi Biotech,

130-092-628) following the manufacturer's protocol with the exception that gentle

agitations were performed manually with a P1000 pipette and all incubations were

performed at 10°C. Cells were filtered through 70μm cell strainers (SP Bel-Art,

136800070), rinsed with ice cold 1x PBS and counted before proceeding with the 10x

Chromium Next Single Cell 3' v3.1 platform (single cell preparation, single cell isolation

and barcoding performed according to the manufacturer's instructions). Libraries were

sequenced (Illumina NovaSeq 6000 S4 flowcell, the DNA services laboratory of the Roy

J. Carver Biotechnology Center at the University of Illinois at Urbana-Champaign)

followed by quality control (adaptor trimming, de-duplication), alignment and

demultiplexing based on manufactured indices using Partek Flow software (Partek Inc.

St. Louis, MO, USA, CellRanger).

Bioinformatics – single cell RNA sequencing dataset analysis

Analysis of the d1-4 time-course amnion scRNA-seq dataset was performed using the

Seurat R package (v.4.2.1, (Hao et al., 2021)). For quality control, cells were filtered out

if the total number of detected genes was less than 1,500, if the expression percentage

from mitochondrial genes was less than 3% or greater than 15%, or if the total number

of transcripts was greater than 30,000. Expression values were then log-normalized

with scaling factor 10,000 for cell-level normalization, and further centered and scaled

across genes. Principal component analysis (PCA) was performed prior to embedding

8,765 cells into two-dimensional space with UMAP using top 15 PCs. Cell clusters were

identified using FindNeighbors and FindClusters functions based on a shared nearest

neighbor modularity optimization clustering algorithm (resolution set as 0.2). In addition,

the "early progenitor" cluster was identified within the original "progenitor" cluster

(containing both early and later progenitors) by a gating strategy that sets a cut-off

based on the normalized expression levels of TBXT (more than 0.4) and CLDN10 (less

than 3.8). Nearby TBXT$^-$ cells within the TBXT$^+$ domain were also included in the "early

progenitor" cluster using CellSelector function because those cells are likely due to

dropouts. The cluster marker genes were identified using FindAllMarkers function

(Wilcoxon Rank Sum test) with adjusted p values less than 0.05.

Mapping and gene expression analyses of Tyser et al. Carnegie Stage 7 human embryo

single cell RNA sequencing dataset

The original UMAP coordinates and annotations for all 1,195 cells were used in **Fig.**

S1C as well as in the accompanying expression plots. The processed data were

downloaded from http://www.human-gastrula.net, which was used to generate a UMAP

plot (using DimPlot function in R package Seurat), as well as to perform expression

analyses (FeaturePlot function in Seurat).

Integration and gene expression analyses of the d1-4 amnion and Tyser et al. Carnegie

Stage 7 human embryo single cell RNA sequencing datasets

The two single cell datasets were integrated using a canonical correlation analysis

based approach implemented in the IntegrateData function with 4,000 anchor features

and 50 dimensions for anchor weighting in Seurat R package (v4.2.1).

Integration and gene expression analyses of Yang et al. GD14 cynomolgus macaque

embryo single cell RNA sequencing dataset

The processed data were downloaded from GSE148683, and the ensemble genome

build Macaca_fascicularis_5.0 release 96 was used to identify human orthogonal gene

symbols. Datasets from two distinct GD14 embryos (Yang et al., 2021) were integrated

using 5,000 anchor features and 10 dimensions in the IntegrateData function in Seurat

R package (v.4.2.1) package: trophoblast cells were removed from the dataset prior to

integration. Six general cell populations (epiblast, transition, mesoderm, amnion,

endoderm and extraembryonic mesenchyme) were identified using FindClusters

function (resolution as 0.4).

RNA velocity (scvelo) analysis

From the aligned BAM files, a loom file was generated for each d1-d4 dataset

respectively using the function run10x mode in software velocyto (v0.17) with default

parameters to create count matrices made of spliced and unspliced read counts (with

Human genome annotation hg19). scVelo (Python package v.0.2.5 (Bergen et al., 2020)

was then used to analyze the loom files and examine RNA velocity. Inferred RNA

velocity was overlaid onto the UMAP embedding (created by Seurat pipeline as

described above) and visualized by stream plot (with default parameters except

smoothness was set to 1).

FIGURES AND FIGURE LEGENDS

Figure 1. Single-cell transcriptomic signatures of developing Gel-3D human

amnion model

(A) Confocal images of d1, d2, d3 and d4 hPSC-amnion cysts stained for TFAP2A, an

amnion marker, NANOG, a pluripotency marker, as well as for DNA (blue) and

membrane (magenta, using wheat germ agglutinin). White arrowheads indicate cells

that display weak TFAP2A expression. Scale bars = 20μm.

(B) Quantitation for nuclear aspect ratio, a measure of epithelial cell morphology, of

developing hPSC-amnion at indicated timepoints, revealing a gradual squamous

morphogenesis (quantitation method in **Fig. S1A**). * indicates p < 0.05.

(C,D) A UMAP plot displaying the single cell transcriptomes of 1,359 d1 (salmon), 1,546

d2 (sage), 2,352 d3 (light blue) and 3,508 d4 Gel-3D cells (C) with seven identified cell

populations (D, pluripotency-exiting (salmon, 1,363 cells), early progenitor (brown, 479

cells), late progenitor (sage, 1,045 cells), specified (green, 1,366 cells) and maturing

(purple, 2,795 cells), PGC-LC (primordial germ cell-like cells, blue, 1,055 cells) and

advM-LC (advanced mesoderm-like cells, magenta, 662)).

(E-G) Expression of known early (E, *ID2*, *GATA3*, *TFAP2A*), specified (F, *ISL1*, *HAND1*,

DLX5) and late (G, *GABRP*, *VTCN1*, *IGFBP3*) amnion markers superimposed onto the

UMAP plot.

(H) RNA velocity analysis of the time-course Gel-3D dataset, showing predicted lineage

progression trajectories.

(I-K) A UMAP plot showing the integrated single cell transcriptomes of d1-d4 Gel-3D

and Tyser *et al.* CS7 embryo datasets, revealing close overlap of Gel-3D amnion

progressing cells with Tyser Epiblast, Primitive Streak and Ectoderm/Amnion cells

(shown in (I) with original sample, (J) with Gel-3D, and (K) with Tyser *et al.* annotations

(individual Tyser annotations plotted in **Fig. S1F**)).

(L,M) Expression of primordial germ cell (L, *PRDM1*, *SOX17*, *NANOS3*) and advanced

mesoderm (M, *HAND2*, *GATA4*, *PITX2*) markers.

(N) Summary of marker expression.

Figure 2. Transient expression of TBXT and CLDN10 marks amnion progenitor

populations

(A,B) Expression of *TBXT* (A) and *CLDN10* (B) superimposed onto the time-course Gel-

3D UMAP plot.

(C) Expression summary of top three most differentially expressed genes in early

progenitor and late progenitor populations.

(D,E) Confocal micrographs of developing hPSC-amnion harvested at indicated

timepoints, stained with indicated markers. (D) Weak TBXT expression is present at d2

(white arrowheads), but is diminished by d3. (E) A white arrowhead indicates a

CLDN10/TFAP2A double-positive cell. CLDN10 expression is extinguished by d4.

Membrane was stained using wheat germ agglutinin. Scale bars = 20μm

(F) Quantitation of CLDN10 expressing cells per cyst in Gel-3D overtime (n = 25 cysts

per timepoint).

Figure 3. CLDN10 is expressed in amnion progenitor cells at the amnion-epiblast

boundary of the PASE model as well as cynomolgus macaque peri-gastrula

(A) Optical sections of PASE stained with indicated markers. White arrowheads indicate

CLDN10 staining at the amnion-epiblast boundary. Weak CLDN10 signal in the

pluripotent territory is likely due to weak CLDN10 expression in the pluripotent cells;

brighter signal in the pluripotent territory is likely caused by the boundary cell

extensions.

(B) Confocal images of a cynomolgus macaque embryo staged at CS6/7 stained using

indicated markers. CLDN10 is expressed at the transitioning boundary cells between

the amnion and the epiblast. Insets indicate magnified regions.

(C,D) Confocal images of cynomolgus macaque embryos staged at CS10 stained using

indicated markers. Insets indicate magnified regions. In the CS10 embryo (early

organogenesis stage), while some posterior boundary CLDN10 staining is seen (seen in

(iii)), the anterior CLDN10 staining is highly prominent at the amnion-anterior surface

ectoderm border (in (i)). CLDN10 is also enriched in the dorsal foregut labeled by ISL1/2

and SOX2 but not SOX17 (indicated by * in (ii)). **Fig. S3F** provides a bird's-eye view of

the CS10 embryo. HT, heat tube; FB, forebrain; YS, yolk sac. White arrowheads

indicate the foregut pocket, while white arrows indicate the hindgut pocket.

Scale bars = 20μm (A), 200μm (B), 500μm (C,D).

Figure 4. CLDN10 is critical for maintaining amnion fate progression in the

progenitor population.

(A-D) Confocal images of control (A,C) and *CLDN10*-KO (B,D) hPSC grown in Gel-3D,

harvested at indicated timepoints and stained with indicated markers. CLDN10

expression is not seen in the KO (B). In the absence of *CLDN10*, PGC-LC formation is

seen early in d2 cysts, and cyst organization is disrupted by d4. Scale bars = 50μm.

(E) Flow chart outlining five continuous amniogenic lineage progressing states with

representative markers.

(F) A schematic representation of primate peri-gastrula near the primitive streak

(generated based on images in **Fig. 3B**, purple cells indicate primitive streak-derived

disseminating cells). Transcriptional signatures consistent with amniotic lineage

progression (indicated by dotted arrow) are observed at the amnion-epiblast boundary:

progressing from SOX2[+] epiblast cells (blue), CLDN10[+] transitioning boundary cells

(green), and, then, to ISL1[+] lineage specified amnion (orange).

Figure S1. Related to Figure 1

(A) Confocal images of hPSC-amnion stained with indicated markers, outlining nuclear

aspect ratio quantitation.

(B) Expression of pluripotency markers, SOX2, NANOG and POU5F1, superimposed

onto the Gel-3D UMAP plot.

(C) A UMAP plot displaying the original single cell transcriptome coordinates of the CS7

human embryo described in Tyser *et al.*, shown with the original annotations.

(D,E) Expression of indicated broad (D) and late (E) amnion markers superimposed

onto the Tyser UMAP plot.

(F) UMAP plots showing the integrated single cell transcriptomes of d1-d4 Gel-3D and

Tyser *et al.* CS7 embryo datasets with individual Tyser annotations.

Figure S2. Validation of selected specified and maturing markers in a cynomolgus

macaque embryo staged between CS12 and CS13

(A,B) CS12/13 cynomolgus macaque embryo sections were stained for indicated

markers (A: *GABRP* – green, *SESN3* – red, *ISL1* – magenta; B: *ADAMTS18* – green,

SESN3 – red, *ISL1* – magenta) using RNA *in situ* hybridization. (i) – (vii) indicate insets:

individual channels are shown in gray scale to aid visualization (from left to right:

GABRP, *SESN3*, *ISL1* and merge in (A), *ADAMTS18*, *SESN3*, *ISL1* in (B)), and dotted

lines indicate a layer of amniotic epithelium and mesenchyme. Most amnion cells are

positive for *GABRP*, *SESN3* and *ISL1*. *ADAMTS18* expression is restricted to a fraction

of cells. In (iv), there are four *ADAMTS18*^{high} cells, while *ADAMTS18* is largely

undetected in the amnion in (v). In (vi) and (vii), most cells are *ADAMTS18*^{high}.

FB; forebrain, HB, hindbrain; FG, foregut; C; cardiac tissue; NT, neural tube; S, somite.

* indicates an *ISL1*$^{+}$ tissue that is likely of genital tubercle lineage. Scale bars = 500μm.

(C) An overview image of the CS12/13 embryo implantation site shown in (A). Only

nuclear staining is shown in grayscale. Scale bar = 2mm.

Figure S3. Additional expression analyses

(A) Expression of *TBXT* and *CLDN10* superimposed onto the uncropped Tyser *et al.*

CS7 human embryo UMAP plot.

(B) A UMAP plot displaying the GD14 cynomolgus macaque single cell transcriptome

with six identified general clusters (epiblast – light blue; transition – purple; amnion –

gold; mesoderm – green; endoderm – teal; extraembryonic mesoderm – salmon). Arrow

indicates the likely trajectory of amnion differentiation based on the data in Yang *et al.*

as well as the expression analysis in (C). Note that, although not identical to the

published UMAP plot in Yang *et al.*, general characteristics are well recapitulated.

Trophectoderm cells have been omitted as performed in Yang *et al.*.

(C) Expression of indicated markers superimposed onto the Yang *et al.* UMAP plot. Late

amnion markers (EPAS1 and GABRP) are expressed at the tip of the TFAP2A and ISL1

double-positive amnion cluster. CLDN10 positive cells show a weak TBXT expression

level.

(D,E) Individual channels of the whole embryo images in **Fig. 3C** and **3D**, respectively.

(F) An overview image of the CS10 (**Fig. 3C**) embryo implantation sites (DNA signal

shown in gray scale, Scale bar = 2mm).

Figure S4. Validation of two distinct *CLDN10*-KO hPSC lines

(A) Sequenced genotyping results of *CLDN10*-KO #1 and #2 lines. KO lines #1 and #2

have similar phenotype, both displaying an increased formation of PGC-LC and

defective cyst organization. The second coding sequence of *CLDN10* was targeted.

(B) Representative sequence traces for each mutations.

Table 1. Lists of differentially expressed genes in pluripotency-exiting (1506), early progenitor (692), late progenitor (697), specified (944), maturing (868), PGC-LC (1181) and advM-LC (746) clusters

Table 2. List of antibodies, RNAscope probes, primers and plasmids used in this study

ACKNOWLEDGEMENTS

We thank the Wisconsin National Primate Research Center (WNPRC) Veterinary, Scientific Protocol Implementation, Pathology and Animal Services staff for providing animal care, and assisting in procedures including breeding, pregnancy monitoring, and sample collection. A special thanks to Michele Schotzko, Sara Shaw and Drs. Heather Simmons and Puja Basu for their help in generating the macaque specimens. The contents of this manuscript are solely the responsibility of the authors and do not represent the official views of the NIH. We thank Dr. Deborah Gumucio for insightful comments to the manuscript, as well as the Roy J. Carver Biotechnology Center at the University of Illinois at Urbana-Champaign for sequence services.

Funding: This work was supported by NIH grants R01-HD098231 (K.T.), P51 OD011106 (to the WNPRC) as well as by MCW CBNA Start-up funds, Advancing a Healthier Wisconsin (AHW) Endowment (16003-5520766, N.S.) and the Lalor Foundation Postdoctoral Fellowship (N.S.). Specifically, MCW CBNA Start-up funds

were used to perform experiments using PASE. **Competing interests:** The authors

declare no competing interest. **Data and materials availability**: All data needed to

evaluate the conclusions in the paper are present in the paper and the Supplementary

Materials. The raw data, unfiltered count matrix and processed count matrix will be

available to the database of Genotypes and Phenotypes (dbGaP) upon publication.

Further information and requests for reagents may be directed to Kenichiro Taniguchi.

REFERENCES

Angelow, S., Ahlstrom, R., and Yu, A.S. (2008). Biology of claudins. Am J Physiol Renal
Physiol *295*, F867-876.
Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA
velocity to transient cell states through dynamical modeling. Nature biotechnology *38*,
1408-1414.
Bergmann, S., Penfold, C.A., Slatery, E., Siriwardena, D., Drummer, C., Clark, S.,
Strawbridge, S.E., Kishimoto, K., Vickers, A., Tewary, M*., et al.* (2022). Spatial profiling
of early primate gastrulation in utero. Nature *609*, 136-143.
Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating
single-cell transcriptomic data across different conditions, technologies, and species.
Nature biotechnology *36*, 411-420.
Carleton, A.E., Duncan, M.C., and Taniguchi, K. (2022). Human epiblast
lumenogenesis: From a cell aggregate to a lumenal cyst. Seminars in cell &
developmental biology *131*, 117-123.
Castillo-Venzor, A., Penfold, C.A., Morgan, M.D., Tang, W.W., Kobayashi, T., Wong,
F.C., Bergmann, S., Slatery, E., Boroviak, T.E., Marioni, J.C*., et al.* (2023). Origin and
segregation of the human germline. Life Sci Alliance *6*.
Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and
Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by dual
inhibition of SMAD signaling. Nature biotechnology *27*, 275-280.
Chen, D., Sun, N., Hou, L., Kim, R., Faith, J., Aslanyan, M., Tao, Y., Zheng, Y., Fu, J.,
Liu, W*., et al.* (2019). Human Primordial Germ Cells Are Specified from Lineage-Primed
Progenitors. Cell reports *29*, 4568-4582 e4565.
Chen, F., and LoTurco, J. (2012). A method for stable transgenesis of radial glia lineage
in rat neocortex by piggyBac mediated transposition. Journal of neuroscience methods
207, 172-180.
Chen, K., Zheng, Y., Xue, X., Liu, Y., Resto Irizarry, A.M., Tang, H., and Fu, J. (2021).
Branching development of early post-implantation human embryonic-like tissues in 3D
stem cell culture. Biomaterials *275*, 120898.

Enders, A.C., Hendrickx, A.G., and Schlafke, S. (1983). Implantation in the rhesus monkey: initial penetration of endometrium. The American journal of anatomy *167*, 275-298.

Enders, A.C., Schlafke, S., and Hendrickx, A.G. (1986). Differentiation of the embryonic disc, amnion, and yolk sac in the rhesus monkey. The American journal of anatomy *177*, 161-185.

Hamed, M.M., Taniguchi, K., and Duncan, M.C. (2023). Monitoring Effects of Membrane Traffic Via Changes in Cell Polarity and Morphogenesis in Three-Dimensional Human Pluripotent Stem Cell Cysts. Methods in molecular biology *2557*, 83-98.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M.*, et al.* (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587 e3529.

Heemskerk, I., Burt, K., Miller, M., Chhabra, S., Guerra, M.C., and Warmflash, A. (2017). Morphogen dynamics control patterning in a stem cell model of human embryo. bioRxiv *http://dx.doi.org/10.1101/202366*.

Joy, D.A., Libby, A.R.G., and McDevitt, T.C. (2021). Deep neural net tracking of human pluripotent stem cells reveals intrinsic behaviors directing morphogenesis. Stem cell reports *16*, 1317-1330.

Lacoste, A., Berenshteyn, F., and Brivanlou, A.H. (2009). An efficient and reversible transposable system for gene delivery and lineage-specific differentiation in human embryonic stem cells. Cell stem cell *5*, 332-342.

Miki, T., Lehmann, T., Cai, H., Stolz, D.B., and Strom, S.C. (2005). Stem cell characteristics of amniotic epithelial cells. Stem cells *23*, 1549-1559.

Minn, K.T., Fu, Y.C., He, S., Dietmann, S., George, S.C., Anastasio, M.A., Morris, S.A., and Solnica-Krezel, L. (2020). High-resolution transcriptional and morphogenetic profiling of cells from micropatterned human ESC gastruloid cultures. eLife *9*.

Nakamura, T., Okamoto, I., Sasaki, K., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y., Nakamura, S., Yamamoto, T., and Saitou, M. (2016). A developmental coordinate of pluripotency among mice, monkeys and humans. Nature *537*, 57-62.

Nakamura, T., Yabuta, Y., Okamoto, I., Sasaki, K., Iwatani, C., Tsuchiya, H., and Saitou, M. (2017). Single-cell transcriptome of early embryos and cultured embryonic stem cells of cynomolgus monkeys. Scientific data *4*, 170067.

Overeem, A.W., Chang, Y.W., Moustakas, I., Roelse, C.M., Hillenius, S., Helm, T.V., Schrier, V.F.V., Goncalves, M., Mei, H., Freund, C.*, et al.* (2023). Efficient and scalable generation of primordial germ cells in 2D culture using basement membrane extract overlay. Cell Rep Methods *3*, 100488.

Pedroza, M., Gassaloglu, S.I., Dias, N., Zhong, L., Hou, T.J., Kretzmer, H., Smith, Z.D., and Sozen, B. (2023). Self-patterning of human stem cells into post-implantation lineages. Nature *622*, 574-583.

Perea-Gomez, A., Vella, F.D., Shawlot, W., Oulad-Abdelghani, M., Chazaud, C., Meno, C., Pfister, V., Chen, L., Robertson, E., Hamada, H.*, et al.* (2002). Nodal antagonists in the anterior visceral endoderm prevent the formation of multiple primitive streaks. Developmental cell *3*, 745-756.

Roost, M.S., van Iperen, L., Ariyurek, Y., Buermans, H.P., Arindrarto, W., Devalla, H.D., Passier, R., Mummery, C.L., Carlotti, F., de Koning, E.J.*, et al.* (2015). KeyGenes, a

Tool to Probe Tissue Differentiation Using a Human Fetal Transcriptional Atlas. Stem
cell reports *4*, 1112-1124.
Rostovskaya, M., Andrews, S., Reik, W., and Rugg-Gunn, P.J. (2022). Amniogenesis
occurs in two independent waves in primates. Cell stem cell *29*, 744-759 e746.
Sasaki, K., Nakamura, T., Okamoto, I., Yabuta, Y., Iwatani, C., Tsuchiya, H., Seita, Y.,
Nakamura, S., Shiraki, N., Takakuwa, T.*, et al.* (2016). The Germ Cell Fate of
Cynomolgus Monkeys Is Specified in the Nascent Amnion. Developmental cell *39*, 169-
185.
Schultheiss, T.M., Burch, J.B., and Lassar, A.B. (1997). A role for bone morphogenetic
proteins in the induction of cardiac myogenesis. Genes & development *11*, 451-462.
Sekulovski, N., Wettstein, J.C., Carleton, A.E., Juga, L.N., Taniguchi, L.E., Ma, X., Rao,
S., Schmidt, J.K., Golos, T.G., Lin, C.W.*, et al.* (2024). Temporally resolved early bone
morphogenetic protein-driven transcriptional cascade during human amnion
specification. eLife *12*.
Shahbazi, M.N., and Zernicka-Goetz, M. (2018). Deconstructing and reconstructing the
mouse and human early embryo. Nature cell biology.
Shao, Y., and Fu, J. (2022). Engineering multiscale structural orders for high-fidelity
embryoids and organoids. Cell stem cell *29*, 722-743.
Shao, Y., Taniguchi, K., Gurdziel, K., Townshend, R.F., Xue, X., Yong, K.M.A., Sang, J.,
Spence, J.R., Gumucio, D.L., and Fu, J. (2017a). Self-organized amniogenesis by
human pluripotent stem cells in a biomimetic implantation-like niche. Nature materials
16, 419-425.
Shao, Y., Taniguchi, K., Townshend, R.F., Miki, T., Gumucio, D.L., and Fu, J. (2017b).
A pluripotent stem cell-based model for post-implantation human amniotic sac
development. Nature communications *8*, 208.
Shawlot, W., Wakamiya, M., Kwan, K.M., Kania, A., Jessell, T.M., and Behringer, R.R.
(1999). Lim1 is required in both primitive streak-derived tissues and visceral endoderm
for head formation in the mouse. Development *126*, 4925-4932.
Taniguchi, K., Heemskerk, I., and Gumucio, D.L. (2019). Opening the black box: Stem
cell-based modeling of human post-implantation development. The Journal of cell
biology *218*, 410-421.
Taniguchi, K., Shao, Y., Townshend, R.F., Tsai, Y.H., DeLong, C.J., Lopez, S.A.,
Gayen, S., Freddo, A.M., Chue, D.J., Thomas, D.J.*, et al.* (2015). Lumen Formation Is
an Intrinsic Property of Isolated Human Pluripotent Stem Cells. Stem cell reports *5*, 954-
962.
Townshend, R.F., Shao, Y., Wang, S., Cortez, C.L., Esfahani, S.N., Spence, J.R.,
O'Shea, K.S., Fu, J., Gumucio, D.L., and Taniguchi, K. (2020). Effect of Cell Spreading
on Rosette Formation by Human Pluripotent Stem Cell-Derived Neural Progenitor Cells.
Front Cell Dev Biol *8*, 588941.
Tsaytler, P., Liu, J., Blaess, G., Schifferl, D., Veenvliet, J.V., Wittler, L., Timmermann,
B., Herrmann, B.G., and Koch, F. (2023). BMP4 triggers regulatory circuits specifying
the cardiac mesoderm lineage. Development *150*.
Tyser, R.C.V., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and Srinivas,
S. (2021). Single-cell transcriptomic characterization of a gastrulating human embryo.
Nature *600*, 285-289.

van Wijk, B., Moorman, A.F., and van den Hoff, M.J. (2007). Role of bone
morphogenetic proteins in cardiac differentiation. Cardiovasc Res *74*, 244-255.
Vasic, I., Libby, A.R.G., Maslan, A., Bulger, E.A., Zalazar, D., Krakora Compagno, M.Z.,
Streets, A., Tomoda, K., Yamanaka, S., and McDevitt, T.C. (2023). Loss of TJP1
disrupts gastrulation patterning and increases differentiation toward the germ cell
lineage in human pluripotent stem cells. Developmental cell *58*, 1477-1488 e1475.
Wang, S., Lin, C.W., Carleton, A.E., Cortez, C.L., Johnson, C., Taniguchi, L.E.,
Sekulovski, N., Townshend, R.F., Basrur, V., Nesvizhskii, A.I.*, et al.* (2021). Spatially
resolved cell polarity proteomics of a human epiblast model. Sci Adv *7*.
Warmflash, A., Sorre, B., Etoc, F., Siggia, E.D., and Brivanlou, A.H. (2014). A method to
recapitulate early embryonic spatial patterning in human embryonic stem cells. Nature
methods *11*, 847-854.
Xiao, Z., Cui, L., Yuan, Y., He, N., Xie, X., Lin, S., Yang, X., Zhang, X., Shi, P., Wei, Z.*,*
et al. (2024). 3D reconstruction of a gastrulating human embryo. Cell *187*, 2855-2874
e2819.
Yang, R., Goedel, A., Kang, Y., Si, C., Chu, C., Zheng, Y., Chen, Z., Gruber, P.J., Xiao,
Y., Zhou, C.*, et al.* (2021). Amnion signals are essential for mesoderm formation in
primates. Nature communications *12*, 5126.
Zhao, C., Plaza Reyes, A., Schell, J.P., Weltner, J., Ortega, N.M., Zheng, Y., Björklund,
Å.K., Baque-Vidal, L., Sokka, J., Torokovic, R.*, et al.* (2024). A comprehensive human
embryogenesis reference tool using single-cell RNA sequencing data. bioRxiv *doi:*
10.1101/2021.05.07.442980.
Zheng, Y., Xue, X., Shao, Y., Wang, S., Esfahani, S.N., Li, Z., Muncie, J.M., Lakins,
J.N., Weaver, V.M., Gumucio, D.L.*, et al.* (2019). Controlled modelling of human
epiblast and amnion development using stem cells. Nature *573*, 421-425.
Zheng, Y., Yan, R.Z., Sun, S., Kobayashi, M., Xiang, L., Yang, R., Goedel, A., Kang, Y.,
Xue, X., Esfahani, S.N.*, et al.* (2022). Single-cell analysis of embryoids reveals lineage
diversification roadmaps of early human development. Cell stem cell *29*, 1402-1419
e1408.
864

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1

Figure S2

A

GABRP *SESN3* *ISL1* MERGE

CS12/13 cynomolgus macaque

i

ii

iii

B

ADAMTS18 *SESN3* *ISL1* MERGE

CS12/13 cynomolgus macaque

iv

v

vi

vii

C

Figure S3

Figure S4

A   **Green: PAM**   **Red: Target sequence**   **--- : Deletion**

CLDN10-WT   1. ATGAAGTGTACCAAAGTCGGAGGCTCCGATAAAGCCAAAGCTAAA
                2. ATGAAGTGTACCAAAGTCGGAGGCTCCGATAAAGCCAAAGCTAAA

CLDN10-KO #1 1. ATGAAGTGTAC- - - - - - - - - - - - -TCCGATAAAGCCAAAGCTAAA (13bp deletion)
                2. ATGAAGTGTACCAAA- -CGGAGGCTCCGATAAAGCCAAAGCTAAA (2bp deletion)

CLDN10-KO #2 1. ATGAAGTGTACCAAA- TCGGAGGCTCCGATAAAGCCAAAGCTAAA (1bp deletion)
                2. ATGAAGTGTACCAAA- TCGGAGGCTCCGATAAAGCCAAAGCTAAA (1bp deletion)

B