Non-feature-specific elevated responses and feature-specific backward replay in human brain induced by visual sequence exposure

Tao He[1,2,3,4], Xizi Gong[3,4], Qian Wang[3,4], Xinyi Zhu[3,4], Yunzhe Liu[5], Fang Fang[3,4,6,7*]

[1] Center for the Cognitive Science of Language, Beijing Language and Culture University, Beijing 100083, People's Republic of China

[2] Key Laboratory of Language Cognitive Science (Ministry of Education), Beijing Language and Culture University, Beijing 100083, People's Republic of China

[3] School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing 100871, People's Republic of China

[4] IDG/McGovern Institute for Brain Research, Peking University, Beijing 100871, People's Republic of China

[5] Chinese Institute for Brain Research, Beijing 102206, People's Republic of China

[6] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, People's Republic of China

[7] Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing 100871, People's Republic of China

Corresponding author:

Fang Fang, Ph.D. Professor

School of Psychological and Cognitive Sciences

Peking University

Beijing, P.R. China

Tel & Fax: +86-10-62756932   Email: ffang@pku.edu.cn

Web: www.psy.pku.edu.cn/english/people/faculty/professor/fangfang/index.htm

ABSTRACT

The ability of cortical circuits to adapt in response to experience is a fundamental

property of the brain. After exposure to a moving dot sequence, flashing a dot as cue at

the starting point of the sequence can induce successive elevated responses even in

the absence of the sequence. This cue-triggered elevated responses have been

demonstrated to play a crucial role in predicting future events in dynamic environments.

However, temporal sequences we are exposed usually contain rich feature information.

It remains unknown whether the elevated responses are feature specific and, more

crucially, how the brain organizes this sequence information after exposure. To address

these questions, participants were exposed to a predefined sequence of four motion

directions for about 30 min and subsequently presented with the start or end motion

direction of the sequence as a cue. Surprisingly, we found that the cue-triggered

elevated responses were not specific to a particular motion direction. Interestingly, the

motion direction information was spontaneously reactivated and the motion sequence

was backward replayed in a time-compressed manner. These effects were marginally

significant even with brief exposure. Notably, no replay events were observed when the

second or third motion direction of the sequence served as a cue. Further analyses

revealed that activity in the medial temporal lobe (MTL) preceded the ripple power

increase in visual cortex at replay onset, implying a coordinated relationship between

the activities in the MTL and visual cortex. Together, we demonstrate that visual

sequence exposure could induce two-fold brain plasticity that may simultaneously serve

for different functional purposes. The non-feature-specific elevated responses may

facilitate general processing of upcoming stimuli, whereas the feature-specific backward

replay may underpin passive learning of visual sequence.

INTRODUCTION

The capacity of cortical circuits to undergo plasticity in response to experience is a fundamental feature of the brain (Buonomano and Merzenich, 1998; Costandi, 2016; Li, 2016). This plasticity can be induced not only by active, task-dependent training, such as visual perceptual learning (Watanabe and Sasaki, 2015; Chen et al., 2016; Lu and Dosher, 2022), but also by passive, repetitive exposure (Gutnisky et al., 2009; Sasaki et al., 2010; Ekman et al., 2017). For instance, after repeated exposure to a moving dot sequence, flashing a dot (i.e., cue) at the starting point of the sequence elicits successive elevated neural responses along the motion path in visual cortex, akin to those induced by the actual moving dot sequence in both humans (Ekman et al., 2017, 2023; Lu et al., 2020) and animals (Eagleman and Dragoi, 2012; Xu et al., 2012). This cue-triggered reactivation is thought to be driven by expectation and is prediction-related (Ekman et al., 2017, 2023) since it has been shown to facilitate the prediction of upcoming stimuli and affect the perception of sensory information (Gutnisky et al., 2009; Baker et al., 2014; Pojoga et al., 2020).

Rather than simple white dot sequences used in previous studies (Xu et al., 2012; Ekman et al., 2017, 2023; Lu et al., 2020), temporal sequences to which we are exposed in daily life usually contain rich feature information. However, it remains unknown whether the cue-triggered elevated responses are feature-specific. On the one hand, if they are not feature-specific, these elevated responses may simply reflect general cortical preparedness for any upcoming stimuli. On the other hand, if they are indeed specific to a particular feature in the sequence, the elevated responses could

only facilitate the processing of specific future events, according to the view that

expectation leads to sharper neural tuning and therefore facilitates the processing of

expected stimuli (Kok et al., 2012). For instance, expectation increased the prestimulus

baseline of sensory neurons tuned to the expected stimulus (Wyart et al., 2012; Kok et

al., 2014) and elicited preactivation of the stimulus template in both visual (Kok et al.,

2017) and auditory (Demarchi et al., 2019) cortices. It should be noted that all these

prediction-related feature-specific activities arose from static contexts. Whether

prediction-related responses in dynamic temporal contexts (e.g., visual sequence) are

feature-specific remains unclear.

After exposure to a feature-contained temporal sequence, another important

question is how the feature information in the sequence is encoded and organized in

neural activity following the cue. Previous studies have demonstrated that, after training

with a temporal sequence, memory consolidation (Gridchyn et al., 2020; Gillespie et al.,

2021) or learning process (Igata et al., 2021) has been linked to a neural pattern known

as replay. Replay refers to the sequential reactivation of neural activity patterns

associated with the trained sequence in both sleep (Lee and Wilson, 2002) and awake

(Foster and Wilson, 2006) states and is believed to be crucial for memory consolidation

and learning (Carr et al., 2011; Liu et al., 2021b, 2022). During replay, the neural

representation of the trained sequence was temporally compressed and displayed in

either forward or backward direction (Carr et al., 2011; Joo and Frank, 2018; Nour et al.,

2021; Liu et al., 2022; McFadyen et al., 2023). These replay events frequently coincide

with sharp-wave ripples (SWRs), which are high-frequency (150–220 Hz) oscillations

(O'Keefe and Nadel, 1978; Bush et al., 2022) detected in the local field potentials of the

hippocampus (Buzsáki, 1986, 2015). To date, replay was exclusively found when

participants performed a task with sequences, e.g., nonlocal reinforcement learning (Liu

et al., 2021b) and episodic memory retrieval (Wimmer et al., 2020). It remains unclear

whether simple visual exposure to temporal sequences could also trigger replay events

in humans.

In the current study, we used magnetoencephalography (MEG) to investigate

whether the cue-triggered elevated brain responses following visual sequence exposure

are feature-specific and, more importantly, to understand how the feature information in

the exposed sequence is encoded and organized in the brain after exposure. To

address these questions, participants were initially exposed to a predetermined motion

sequence. We subsequently decoded the motion direction information during the blank

period after presenting only the first or last motion direction in the sequence as a cue.

Surprisingly, we found that the cue-triggered elevated responses were not specific to a

particular motion direction. Interestingly, the motion direction information was

spontaneously reactivated during the blank period and the motion sequence was

backward replayed in a time-compressed manner. This backward replay was identified

even with brief exposure. However, neither forward nor backward replay was detected

when an intermediate motion direction in the sequence was presented as a cue. Lastly,

through MEG source reconstruction analysis, we found that medial temporal lobe (MTL)

activation preceded visual cortical activation, implying that the replay events observed in

visual cortex may be triggered by activities in the MTL.

RESULTS

The visual stimuli used in the study were four random-dot kinematograms

(RDKs). All dots in an RDK moved in a direction (i.e., 0°, 90°, 180°, or 270°) with 100%

coherence. In Experiment 1, we included three trial conditions: full sequence trial, start-

only trial, and end-only trial (**Figure 1A**). In a full sequence trial, participants were

exposed to a predefined sequence of the four RDKs in either clockwise (i.e., 0° → 90°

→ 180° → 270°) or counterclockwise in the center of the screen, with a 300 ms of inter-

stimulus interval (ISI) between every two RDKs. In a start- or end-only trial, the first or

last RDK of the sequence was presented at the beginning of the trial, followed by a 2.4

s of blank period. Participants were instructed to complete three successive phases:

functional localizer phase, exposure phase, and main phase (**Figure 1B**). The functional

localizer data were used to train models to decode each motion direction in the

sequence. During this phase, one of the four RDKs was randomly presented for 1 s in

each trial. During the exposure phase, the participants were exposed to only full

sequence trials for about 30 min. In the main phase, 50%, 25%, and 25% trials were full

sequence, start- and end-only trials, respectively. The full sequence trials served as

topping-up exposure to maintain the exposure effect.

Cue-triggered elevated responses are not feature-specific

We first measured the event-related field (ERF) activity evoked by RDKs in the

three trial conditions using all occipital gradiometer sensors (see Methods). Figure 1C

showed the evoked response by full sequence trials (left panel, cluster-based

permutation test with cluster forming threshold t > 3.1 and 5000 permutations).

Remarkably, start- and end-only trials elicited similar wave-like responses as the full

sequence trials, despite the absence of stimuli following the first RDK (**Figure 1C**,

middle and right panels, cluster-based permutation test with cluster forming threshold

t > 3.1 and 5000 permutations). To further quantify the ERF peak amplitudes in the three

conditions while mitigating baseline confounds, we calculated each peak amplitude by

using the 300 ms blank period just preceding the onset of the corresponding RDK as

baseline (see Methods). We found that the four successive RDKs in the full sequence

trials evoked four comparable peaks after stimulus onset (**Figure 1C**, inset figure in the

left panel; two-tailed t-test; all $ts_{(17)} > 6.8072$, all ps $< 10^{-5}$). Interestingly, in start- and

end-only trials, we could still observe significant peaks during the periods corresponding

to the intervals of the second and third RDKs in the full sequence trials (**Figure 1C**,

inset figures in the middle and right panels; start-only condition, two-tailed t-test; all

$ts_{(17)} > 4.5790$, all ps $< 10^{-3}$; end-only condition, two-tailed t-test; all $ts_{(17)} > 6.0140$, all ps

$< 10^{-4}$), although there was no significant peak during the period corresponding to the

interval of the last RDK (start-only condition, two-tailed t-test; $t_{(17)} = 1.3221$, $p = 0.2036$;

end-only condition, two-tailed t-test; $t_{(17)} = 0.2473$, $p = 0.8076$). These results

demonstrate cue-triggered elevated brain responses following visual exposure to the

RDK sequences, resembling previous studies using simple white dot sequences

(Ekman et al., 2017; Lu et al., 2020).

Figure 1. Stimuli, experimental procedure and evoked responses in Experiment 1

(A) Participants were presented with RDKs in three conditions. In the full sequence trial

condition, four successive RDKs were presented. In the start- or end-only condition,

only the first or last RDK in the sequence was presented at the beginning of the trial. **(B)**

During MEG scanning, the participants were presented with two functional localizer runs

(i.e., functional localizer phase) before providing any sequence information. Next, they

were exposed with four full sequence runs (i.e., exposure phase). Finally, in the main

phase, full sequence, start- and end-only trials were presented in a pseudo-randomized

order. **(C)** Evoked brain responses as a function of time relative to trial onset in the full

sequence, start- and end-only trial conditions, respectively. Bold black lines at the

bottom indicate temporal clusters in which they reached significance when compared to

the pre-stimulus baseline. Inset figures at the top-right corner show the peak amplitudes

during the four corresponding RDK intervals after baseline correction.

Given that we observed the elevated responses even in the absence of stimuli

following the cue, next we examined whether these responses were specific to a

particular feature (i.e., motion direction). Specifically, we asked if we could successfully

decode motion directions in start- and end-only trials, particularly during the blank

periods that corresponding to the intervals of the second and third RDKs in full

sequence trials. To this end, we applied a time-resolved decoding analysis. For each

participant, we trained a one-versus-rest Lasso logistic regression model using the

functional localizer data to classify the neural activity pattern elicited by each motion

direction in the main phase. MEG signals from 72 occipital sensors were selected as

features in the training model. To validate the reliability of our data, we first used a

leave-one-out cross-validation scheme on the localizer data to independently estimate

the decoding accuracy at each time point. At the group level, decoding accuracies

peaked at 411 ms, 464 ms, 444 ms, and 434 ms after stimulus onset for motion

directions 0° (55.37% ± 1.21), 90° (58.10% ± 1.14), 180° (54.07% ± 1.47), and 270°

(53.64% ± 1.34), respectively. There were no significant differences among the four

motion directions in terms of either latency ($F(3, 51) = 0.375$, $p = 0.7716$, $\eta_p^2 = 0.0159$)

or decoding accuracy ($F(3, 51) = 2.757$, $p = 0.0517$, $\eta_p^2 = 0.091$) at the peak time point.

Thereafter, the trained classifier was applied to the MEG signals in the three types of

trials in the main phase. Finally, decoding probability (Turner et al., 2023) for each

motion direction was calculated at each time point at the group level (**Figure 2**). Here,

we used decoding probability because it provides a time-resolved decoding preference

for each motion direction, rather than only a decoded label (e.g., 0°, 90°, 180°, or 270°).

In full sequence trials, the motion direction information could be successfully

decoded, as evidenced by the highest decoding probability during the interval

corresponding to the presentation of the respective RDK (**Figure 2A and 2B**, left

panels). In start- and end-only trials, we could also reliably decode the motion direction

of the first RDK (i.e., the cue) after stimulus onset (**Figure 2A and 2B**, middle and right

panels, only the decoding probability of the first RDK surpass the peak-level

significance threshold obtained from a nonparametric permutation test, FWE corrected

across time). Surprisingly, however, subsequent motion direction information was

absent at the group level during the post-cue blank period, where the cue-triggered

elevated responses were previously observed. Together, these results demonstrate that

the cue-triggered elevated response induced by visual sequence exposure was not

consistently specific to a particular feature across participants and trials.

Figure 2. Time-resolved decoding probability for each motion direction

(A) Visualization of the time-resolved decoding probability in the three trial conditions. Each row shows the decoding probabilities for the four motion directions at that time point and each column indicates one of the four motion directions. **(B)** The line plots of the time-resolved decoding probability for the three conditions. Each colored line shows the time course of the decoding probability for each motion direction. For the start- and end-only conditions, we calculated the permutation threshold estimated by randomly shuffling of the labels and re-decoding, only the decoding probability of the cue surpassed the threshold.

Time-compressed backward replay of exposed motion sequence

How is the motion direction information encoded and organized in the brain

during the post-cue blank period? Clearly, the motion direction representation is not

time-locked to the onset of the cue in either start- or end-only trials. However, it is

possible that individual motion directions are encoded in the MEG signals in a

spontaneous way but are sequentially organized (Liu et al., 2019).

To test this possibility, we trained four decoding models using the functional

localizer data to capture the neural features of the four motion directions. Here, we

employed four decoding models because our aim was to obtain four feature-specific

classifiers. Each classifier was designed to be sensitive to only one motion direction,

thereby being useful for quantifying the evidence of feature-specific sequence in

subsequent steps. The models used a one-versus-rest Lasso logistic regression

algorithm, with MEG signals from 72 occipital sensors as features. We then selected the

time point with the highest decoding accuracy, which was estimated during the

validation of the functional localizer data mentioned above, as the optimal time point for

each participant and motion direction, since these time points are believed to carry the

richest feature information (Mo et al., 2019). For subsequent analyses, we used

classifiers trained at their respective optimal time point. Finally, we obtained four

classifiers, each yielding significant decoding probabilities only when the stimulus

matched the motion direction that the classifier was trained to detect (**Figure 3A**). We

did not find any spatial correlation between any two trained classifiers (highest

correlation $r < 0.12$, Figure S1A). Since each participant and motion direction had an

optimal time point at different latencies, for visualization at the group level, we circularly

shifted the decoding probability over time and aligned their optimal time points to an

arbitrary time point (here, 200 ms after stimulus onset) for each participant and motion

direction, respectively (**Figure 3A**).

Having established the classifiers for each participant and motion direction, we

then applied the classifiers to the MEG signals during the post-cue blank period in start-

and end-only conditions to estimate the reactivation probability (i.e., decoding

probability) for each motion direction at each time point (Liu et al., 2019; Wimmer et al.,

2020; Nour et al., 2021). Example trials in the start- and end-only conditions are shown

in Figure S2A. The results revealed that motion direction reactivations occurred

sparsely during the post-cue blank period, rather than crowded within the intervals

corresponding to the RDK presentations in the full sequence condition, suggesting that

the motion direction information might be reactivated in a spontaneous way. Next, we

applied temporally delayed linear modeling (TDLM) to quantify the evidence whether

and how the spontaneous reactivations follow the order of the exposed sequence

(**Figure 3D**). This algorithm includes two-level regression analyses; a first-level

regression that quantifies the evidence for each pairwise state-to-state transition, and a

second-level regression that evaluates the extent to which the reactivation patterns

align with a particular order of interest. Finally, we defined "sequenceness" as a metric

of forward (i.e., $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$) or backward (i.e., $4 \rightarrow 3 \rightarrow 2 \rightarrow 1$) replay (Kurth-

Nelson et al., 2016; Liu et al., 2019, 2021a; Nour et al., 2021).

Figure 3. Illustration of replay analysis pipeline: Temporally delayed linear modeling (TDLM)

(A) The decoding probabilities were aligned at 200 ms post-stimulus onset according to their corresponding optimal time point per participant and motion direction. Each colored line indicates the decoding results of a classifier applied to a motion direction dataset. The dashed horizontal line indicates the permutation threshold estimated by random shuffling of the labels and re-decoding. **(B)** Regression models were trained for each participant and motion direction using MEG signals from the functional localizer data.

(C) Classifiers were next applied to MEG signals during the post-cue blank period to

derive a decoded [time x state] reactivation matrix. An example of backward sequential

reactivation of stimuli is shown on the left. **(D)** Using TDLM, we quantified the evidence

for sequential replay of motion directions during the post-cue blank period in start- and

end-only conditions. We initially performed a time-lagged regression to create a

regression coefficient matrix [states x states] for each time lag by regressing each

lagged predictor matrix X(Δt) onto the state reactivation matrix, Y (i.e., 1st level GLM

analysis). The degree to which the reactivations systematically follow a transition matrix

of interest was measured by "sequenceness" (i.e., 2nd level GLM analysis). We tested

the magnitude of this effect at each time lag independently for all transition lags up to

600 ms. The dashed line represents the corrected nonparametric statistical significance

threshold (see Methods for details). The green area indicates the lags when the

evidence of sequenceness in the backward direction exceeded the permutation

threshold.

As shown in **Figure 4A** and **4B**, in the start-only condition, we found evidence of

backward replay of the exposed motion sequence (i.e., the replay sequence was 270°

→ 180° → 90° → 0° when the exposed motion sequence was 0° → 90° → 180° →

270°) during the post-cue blank period, with a peak transition lag at 28–40 ms (maximal

effect at 32 ms-lag: $\beta = -0.0202 \pm 0.002$, $p < 1/24 \approx 0.042$, peak-level significance

threshold derived from a nonparametric permutation test, FWE corrected across lags,

Figure 4B). For visualization, an example of backward replay of the motion sequence

was illustrated in **Figure 4A**. This effect was observed in most participants (Figure S3A).

In the end-only condition, we also found evidence of backward replay during the post-

cue blank period, with a peak transition lag at 28–36 ms (maximal effect at 32 ms-lag: β

= −0.0145 ± 0.0016, $p < 1/24 \approx 0.042$, peak-level significance threshold obtained from a

nonparametric permutation test, FWE corrected across lags; **Figure 4E**). **Figure 4D**

shows an example of backward replay of the motion sequence in this condition, found in

most participants (Figure S3A). Note that the time lag in the horizontal axis in **Figure 4B**

and **4E** indicates the interval between the onsets of every two items in the replayed

motion sequence. We found that the replayed sequence was approximately 10 times

faster than the evoked activity sequence in the full sequence condition.

To further examine the period during which the backward replay occurred most

frequently, we divided the blank period (2.4 s) into 4 stages, each lasting 600 ms. In

start-only trials, we found that the backward replay predominantly appeared within the

third stage of the blank period (1.2 – 1.8 s after the onset of the blank period, Wilcoxon

signed-rank test, $p = 0.0108$), but there was no significant sequenceness in the other

three stages (first stage, $p = 0.7112$; second stage, $p = 0.5566$; fourth stage, $p = 0.6791$;

Figure 4C). In end-only trials, the backward replay was more likely to occur during the

second and third stages of the blank period, although the results approached, but did

not reach, significance (Wilcoxon signed-rank test, second stage, $p = 0.0936$; third

stage, $p = 0.0642$; **Figure 4F**), In contrast, the replay was not frequently observed

during the first and last stages (Wilcoxon signed-rank test, first stage, $p = 0.9479$; fourth

stage, $p = 0.4997$). Finally, in a control analysis, we conducted a comprehensive

examination of all 24 potential sequences. Only the backward replay was detected in

both start- and end-only conditions (Figure S4).

Figure 4. Backward replay in both start- and end-only conditions

(A and **D)** Examples of backward sequential reactivation in start- (A) and end- (D) only conditions from a representative participant. Each row represents the reactivation probabilities for the four motion directions at that time point and each column indicates one of the four motion directions. **(B and E)** Backward replay of the exposed motion sequence with peak transition lags at 28–40 ms in the start-only condition (B) and at 28 – 36 ms in the end-only condition (E). Horizontal dashed lines represent corrected significance levels from a nonparametric permutation test at the 2nd-level GLM analysis of TDLM. **(C and F)** Backward replay of the exposed motion sequence predominantly appeared at 1.2 – 1.8 s in the start-only condition (C) and 0.6 – 1.8 s in the end-only condition (F) after the onset of the blank period.

Backward replay is cue-dependent and depends on the amount of exposure

So far, we have demonstrated that the cue-triggered elevated responses induced

by the motion sequence exposure were not motion direction specific. However, the

motion information was spontaneously reactivated and the motion sequence was

backward replayed in a time-compressed manner. It remains unknown whether the

observed backward replay of the motion sequence was cue-dependent or not. In other

words, does the replay of the motion sequence occur irrespective of which item of the

sequence is presented as a cue? To address this question, we conducted Experiment 2,

mirroring the design of Experiment 1 but with a different cue. Instead of flashing the first

or last RDK, we presented the second or third RDK as a cue (2^{nd}-only or 3^{rd}-only

condition, see Figure S5A and Methods) to examine whether these two non-terminal

cues could induce replay during the post-cue blank period. We found no evidence of

either forward or backward replay in either condition (**Figure 5A**; maximal nonsignificant

effect at 32 ms-lag: 2^{nd}-only condition, $\beta = -0.0024 \pm 0.0014$; 3^{rd}-only condition, $\beta =$

-0.0018 ± 0.0015).

Figure 5. Backward replay is cue-dependent and depends on the amount of

exposure

(A) No evidence for replay was found in either the 2nd-only (left) or 3rd-only (right)

condition in Experiment 2. Horizontal dashed lines have the same meaning as those in

Figure 4B and 4E. **(B)** In Experiment 3, immediately after the functional localizer

phase, participants entered the main phase without the exposure phase. In the end-

only condition, backward replay was observed; however, in the start-only condition, no

such replay was observed.

Another interesting question is how changing the amount of exposure would

affect the replay events during the blank period. To explore this issue, we conducted

Experiment 3, also similar to Experiment 1 except for the removal of the exposure

phase. Accordingly, only full sequence trials (50% trials) in the main phase served for

exposure (see Figure S5B and Methods). In the start-only condition, we found a

numerical trend of backward replay at transition lags at 20–40 ms, although it did not

reach statistical significance (**Figure 5B**, left panel; maximum effect at 28 ms-lag: β =

-0.0139 ± 0.0021). In the end-only condition, the evidence for backward replay just

reached the significance level at transition lags at 20 – 40 ms (**Figure 5B**, right panel;

maximal effect at 24 ms-lag: $\beta = -0.0147 \pm 0.0019$, $p < 1/24 \approx 0.042$, using the peak-

level significance threshold from a nonparametric permutation test, FWE-corrected

across lags).

Power increase in replay-associated sharp-wave ripple frequencies

Previous studies on rodents and humans showed that replay events were

associated with increased high-frequency ripple power (Buzsáki, 2015; Liu et al., 2019).

To investigate whether such replay-associated power increase could be observed in our

study, we performed a time-frequency analysis using combined data from start- and

end-only conditions in Experiment 1. We first identified putative replay onsets during the

post-cue blank period in each trial, using probabilities decoded from MEG signals in the

occipital region. Replay onsets were determined by choosing time points with a high

(>95[th] percentile) probability for backward replay with a 32 ms-lag transition (maximal

replay effect at the group level for both start- and end-only conditions, Figure 4B and

4E; see Methods for details). Using the MEG signals recorded from whole brain

sensors, we found a transient ripple power increase at 120–180 Hz at the onset of

replay events, compared to the baseline period of 50–100 ms prior to replay onset

(**Figure 6A**; cluster-based permutation test with cluster forming threshold t > 3.1 and

5000 permutations).

Figure 6. Replay events align with ripple power, with source activation in the MTL

preceding activation in visual cortex

(A) Top: In Experiment 1, a time-frequency decomposition of sensor-level data revealing

a brief increase in high-frequency oscillatory power at replay onset. Bottom: a cluster-

based permutation test (cluster forming threshold, t > 3.1; number of permutations =

5000) could identify a significant cluster around 140 Hz. **(B)** Source localization of

ripple-band power 30 ms before replay onset showing significant activation in the MTL

(peak Montreal Neurological Institute [MNI] coordinates: X = 21, Y = 21, Z = 13.

Neurological orientation.) **(C)** Source localization of ripple-band power at replay onset

showing significant activation in visual cortex (peak MNI coordinates: X = 20, Y = 8, Z =

17). **(D)** The activation time course of the MTL at its peak MNI coordinate is shown in

red, whereas that of visual cortex at its peak MNI coordinate is displayed in green. The

MTL reached its peak activation before visual cortex.

Visual cortical activation lags MTL activation

Since replay events were obtained using decoding probabilities from occipital

sensors only, we next examined whether the observed replay onsets were related to

power increase within visual cortex. Using a linearly constrained minimum variance

(LCMV) beamforming algorithm (Van Veen et al., 1997), we first epoched the data using

replay onsets combined from both start- and end-only conditions and then beamformed

the broadband power into the source space (see Methods for details). We found that the

power increase at replay onset was localized to sources in bilateral visual cortex

(**Figure 6C**). Moreover, the right MTL, including the hippocampus and amygdala, was

associated with power increase prior to the onset of replay events (**Figure 6B**).

Specifically, activation (i.e., power increase) in the MTL occurred 30 ms earlier than that

in visual cortex ($p < 0.05$, corrected, whole brain nonparametric permutation test). For

display purposes, we extracted activations from the 10 most activated voxels within the

MTL and visual cortex, respectively, and plotted the time courses of their broadband

power. Peak activation in visual cortex at replay onset was preceded by peak activation

in the MTL (**Figure 6D**), implying an information flow from the MTL to visual cortex.

DISCUSSION

We found the co-existence of time-locked, non-feature-specific elevated

responses and non-time-locked, feature-specific backward replay after visual sequence

exposure in human brain. The feature-specific backward replay occurred in a time-

compressed manner and could be triggered only by the first or last stimulus of the

sequence. Interestingly, even brief exposure to the sequence could still induce a trend

for the backward replay. Finally, we observed that MTL activity preceded the ripple

power increase in visual cortex at replay onset.

Our study provides two new findings in the fields of vision and learning. First,

different from many previous studies showing that expectation-based responses in

static contexts are feature-specific (Wyart et al., 2012; Kok et al., 2014, 2017), we show

here that prediction-related elevated responses in the dynamic temporal context are not.

This non-feature-specific elevated responses potentially facilitate the general

processing of any upcoming stimuli, rather than stimuli with a specific feature. Second,

contrary to the prevailing notion that replay in learning and memory requires lengthy

training with a task, our study shows that even brief exposure to a visual sequence is

sufficient to induce replay. The replay we unveiled here may play a critical role in

memorizing the visual sequence.

Regarding the two-fold neural consequences following the visual sequence

exposure, an interesting question is whether the elevated responses and the backward

replay share the same neural origin, for instance, originating from the hippocampus or

other brain areas. For the elevated responses, the cue serves to provide temporal

information for upcoming stimuli but without detailed feature information, thereby

priming the cortices for activation and facilitating the general processing of future

events. Previous animal studies (Xu et al., 2012; Gavornik and Bear, 2014) showed that

this process can be implemented through a simple local synaptic mechanism in visual

cortex (Xu et al., 2012) without top-down guidance. Moreover, Ekman et al. (2023)

recently found no functional relationship between activities in V1 and the hippocampus

when exposing human participants with a white dot sequence, further suggesting that

the elevated responses may indeed originate in visual cortex.

Different from the elevated responses, the replay events are likely initiated by the

hippocampus. This speculation is supported by two key findings. First, the replay of the

motion direction information manifested in a backward direction and a time-compressed

manner. Second, the replay is observed regardless of whether the cue is the first or last

RDK in the sequence. Both of these two properties cannot be explained by a simple

local synaptic mechanism (Xu et al., 2012) or a pattern completion-like mechanism

(Hindy et al., 2016; Ekman et al., 2017; Kok and Turk-Browne, 2018) within visual cortex

alone. Instead, the replay entails reorganizing the motion direction sequence in the

brain. Therefore, we propose the involvement of active communication between the

hippocampus and visual cortex in the occurrence of replay events, consistent with the

view that the hippocampus encodes relationships among stimuli whereas visual cortex

primarily acts as a platform for the manifestation of replay events (Whittington et al.,

2020). Previous studies on memory consolidation considered the exchange of

information between these two regions critical for facilitating replay events through

hippocampal-neocortical circuits (Buzsáki, 1996; Ji and Wilson, 2007; Carr et al., 2011;

Ólafsdóttir et al., 2016; Buch et al., 2021). Functionally, replay events offer a mechanism

for transferring recent experience from the hippocampus to the cortex, enabling the

encoding of stimulus relationships in the cortex (Marr and Brindley, 1971; Alvarez and

Squire, 1994; Redish and Touretzky, 1998; Dimakopoulos et al., 2022).

A natural behavioral consequence of visual sequence exposure is visual

sequence learning (Baker et al., 2014; Finnie et al., 2021; Ekman et al., 2023). How is

the learning implemented through hippocampus-dependent replay in visual cortex?

Three key processes are considered here. First, the hippocampus is involved in

encoding relationships among stimuli (Staresina and Davachi, 2009; Turk-Browne et al.,

2009; Hsieh et al., 2014; Garvert et al., 2017); in fact, the ability to encode such

relationships drastically decreases when the hippocampus is damaged (Chun and

Phelps, 1999; Hannula et al., 2006; Konkel et al., 2008; Schapiro et al., 2014; Finnie et

al., 2021). Second, visual cortical areas act as a "cognitive blackboard" where task-

relevant features are highlighted through feedback connections (Roelfsema and de

Lange, 2016). Such a blackboard can be flexibly written or edited. Third, there are

strong bidirectional connections between the hippocampus and sensory cortices

(Eichenbaum et al., 2007; Henke, 2010). As proposed by the hippocampal-cortical

backward projection model (Rolls, 2000), sequential reactivations of feature information

initially generated in the hippocampus can be quickly and accurately sent back to the

sensory cortices, consistent with the findings of Ji and Wilson, (2007). A recent study

also provided direct evidence that visual sequence plasticity is impaired when the

hippocampus is damaged (Finnie et al., 2021), supporting the hypothesis of functional

feedback information flow.

Why does the replay manifest in a reverse order? To date, there is no consensus

on this issue. In rodents, a seminal study of replay during the awake state showed that

when animals stopped at the end of a rewarded pathway, place cells were reactivated in

the reverse order of the previously experienced direction (Foster and Wilson, 2006).

However, a later study revealed that awake replay could occur in either a forward or

backward direction relative to behavioral experience (Diba and Buzsáki, 2007).

Similarly, in humans, both directions have been observed in different nonspatial

cognitive tasks (Liu et al., 2019, 2021b; Wimmer et al., 2020). Forward replay may be

associated with planning (Ólafsdóttir et al., 2015; Gillespie et al., 2021), providing

information pertaining to the assessment of future pathways (Diba and Buzsáki, 2007).

Backward replay may be more related to experience, as often observed at the end of

runs when animals consume a reward (Foster and Wilson, 2006; Ambrose et al., 2016).

Nevertheless, the exact function of the replay direction remains mysterious, as both

forward and backward replays are modulated by task demands (Ólafsdóttir et al., 2017).

Thus, the underlying neural mechanisms of backward replay in visual cortex remain to

be investigated.

Finally, we also found that replay occurrence is modulated by the cue. This result

highlights the importance of the start and end points of the sequence in the replay. One

fascinating proposal is that the replay event is sensitive to sequence boundaries, as

indicated by the role of the start or end point of the sequence as salient boundaries that

anchor place cell firing (Rivard et al., 2004). Accordingly, previous studies have shown

that when rats are trained to run along a linear track starting at different points, place

fields tend to be anchored to either the start or end of the journey (Gothard et al., 1996;

Redish et al., 2000), suggesting that a boundary is essential to sequence integrity and

may play a pivotal role in triggering replay onset. An alternative explanation to these

findings posits that the onset of the second or third stimulus in the sequence reinstate

the neural representations of a partial visual sequence (Ekman et al., 2023). For

example, for a given sequence, A → B → C → D, flashing the third stimulus (i.e., C)

only trigger a backward reactivation of the sequence giving C → B → A during the blank

period.

Taken together, we found that simple visual sequence exposure could

concurrently induce two-fold brain plasticity, that is, non-feature-specific elevated

responses and feature-specific backward replay in the human visual cortex. We

speculate that the non-feature-specific elevated responses may enhance general

processing of upcoming visual stimuli, whereas the feature-specific backward replay

may subserve visual sequence learning and memory. These findings significantly

advance our understanding of the task-independence and the multifaceted nature of

brain plasticity in response to visual experience.

METHODS

Participants

A total of 59 healthy participants (29 females) were involved in the three

experiments (Experiment 1: n = 21, 11 females, 22.1 ± 2.61 years; Experiment 2: n =

18, 10 females, 23.56 ± 3.29 years; Experiment 3: n = 20, 10 females, 20.65 ± 2.62

years). In Experiment 1, data from three participants were excluded before analyses, as

two showed large head motion (>20 mm) and the behavioral performance of one was at

chance level. In Experiment 3, data from two participants were excluded before

analyses because of large head motion (>20 mm). All participants were recruited in

exchange for monetary compensation (100 RMB/h). They reported normal or corrected-

to-normal vision and had no history of neurological disorders. They were naive to the

purposes of the study. The experiments reported here were carried out in accordance

with the guidelines expressed in the Declaration of Helsinki. All participants provided

written informed consent in accordance with the procedures and protocols approved by

the Human Subject Review Committee of Peking University.

Task

Experiment 1

Visual stimuli were RDKs with 100% coherence. All dots in an RDK moved in the

same direction (luminance: 2.86 cd/ m^2; diameter: 0.1°; speed: 8°/s) and were

presented against a gray background (luminance: 16.7 cd/m^2). At any one moment, 400

dots were visible within a 9° circular aperture and moved in one of the four directions: 0°

(right), 90° (up), 180° (left), and 270° (down). Each participant completed three phases

successively in the MEG scanner, namely, the functional localizer phase, the exposure

phase, and the main phase (Figure 1C). In the functional localizer phase, each trial

started with the presentation of an RDK for 1 s followed by a 1–1.5 s intertrial interval

(ITI). Participants did not need to perform any task in this phase. The motion direction of

the RDK was randomly chosen from the four directions. Each participant performed two

functional localizer runs and, each run comprised 100 trials, resulting in a total of 50

trials per motion direction. The localizer data were used to train motion direction

classification models. This phase took approximately 10 min.

In the exposure phase, we showed participants the four RDKs in either clockwise

or counterclockwise order (e.g., 0° → 90° → 180° → 270°, Figure 1A and 1C); each was

displayed for 400 ms, followed by 300 ms of a blank screen with fixation only. Therefore,

the total sequence lasted for 2.8 s. The order was counterbalanced among participants,

but once decided, it was fixed for each participant. Participants were instructed to detect

an oddball RDK by pressing a button, i.e., in 20% trials, dots in one of the four RDKs

traveled at a faster speed (9°/s). Each participant completed four runs of 50 trials.

In the main phase, participants were presented with trials in three different

conditions: full sequence condition (50% trials), start-only condition (25% trials), and

end-only condition (25% trials). Trials in the full sequence condition were identical to

those in the exposure phase which served as "topping-up" exposure to maintain the

exposure effect, similar to "topping-up" adaptation in visual adaptation studies (Fang et

al., 2005). In the start- and end-only conditions; however, we only presented the first

RDK (start-only condition) or the last RDK (end-only condition) of the full sequence. For

a given run, the order of the three conditions was pseudorandomized with the restriction

that the start- and end-only trials were always preceded or followed by a full sequence

trial. Participants performed an identical oddball detection task as during the exposure

phase. The oddball occurred in 10% of full sequence trials. Finally, each participant

completed four runs of 48 trials, yielding a total of 96 full sequence trials, 48 start-only

trials and 48 end-only trials.

 Experiments 2 and 3

In Experiment 2, we only presented the second or the third RDK as a cue at the

start of the trial, referred to as the 2nd-only and 3rd-only conditions, respectively. The

procedure of Experiment 2 was similar to that of Experiment 1 except that the start- and

end-only conditions were replaced with the 2nd-only and 3rd-only conditions. Experiment

3 followed the same procedure as Experiment 1 except that the exposure phase was

removed.

QUANTIFICATION AND STATISTICAL ANALYSIS

MEG acquisition and preprocessing

Neuromagnetic signals were recorded continuously at 1000 Hz with a 306-

channel (204 planar gradiometers; 102 magnetometers), whole-head MEG system

(Elekta Neuromag TRIUX) in a magnetically shielded room. Before scanning, four-

headed position indicator coils attached to the scalp determined the head position

with respect to the sensor array. Coil location was digitized with respect to three

anatomical landmarks (nasion and preauricular points) with a 3D digitizer

(Polhemus Isotrak system). Participants sat upright inside the scanner, while the

stimuli were projected onto a screen suspended in front of them. Participants

responded using a MEG-compatible button box held in their right hand.

To reduce noise from external environment, the temporal extension of signal-

space separation (tSSS) method was applied at the preprocessing stage using the

Elekta Neuromag MaxFilter software (Taulu and Simola, 2006). MEG signals were

high-pass filtered at 0.5 Hz using a first-order IIR filter to remove slow-drifts. Data

were then downsampled from 1000 Hz to 250 Hz for sequenceness analysis or 500

Hz for time-frequency analysis. Excessively noisy segments and sensors were

automatically removed before independent component analysis (FastICA,

http://research.ics.aalto.fi/ica/fastica) and performed to remove artifacts including

cardiac signals, eye movements and blinks, and environmental noise. Artifact

components were removed by visual inspection of spatial topography, time course,

kurtosis of the time course, and frequency spectrum of all components. Only 72

sensors (including both gradiometers and magnetometers) covering the occipital

lobe, labeled as "Occipital" in the MEG data acquisition system were used for MEG

data analyses, except for source localization. The sensor selection was primarily

motivated by the main objective of the study, examining replay events in visual

cortex.

Event-related fields

To calculate the event-related fields (ERFs), MEG epochs around trial onset

were segmented for each trial and baseline-corrected using the mean activity in the

time window of [−0.3 s, 0] before trial onset. Only 48 planar gradiometers of

"Occipital" sensors were used to calculate the ERFs. We then averaged the planar-

combined ERF activity for each condition. To avoid any potential baseline influence

due to the short interval between every two RDKs, we further calculated the ERF

peak amplitude by using the mean activity in the time window of [−0.3 s, 0] before

the onset of the corresponding RDK as baseline. Subsequently, peak amplitudes

during the four corresponding RDK intervals were calculated as the difference

between the peak and its corresponding baseline.

Decoding analysis

Decoding analysis was performed to classify the neural activity patterns

elicited by the motion directions of the four RDKs in the main phase. A one-versus-

rest Lasso-regularized logistic regression model was trained using the MEG signals

from the 72 occipital sensors in the functional localizer phase. Specifically, we

trained a five-class classifier, including four classes from trials in which the four

RDKs were presented, and an additional class comprising an equivalent amount of

null data extracted from the 1–1.5 s ITI. Null data were included to reduce the

spatial correlation among the classes, thereby concurrently lowering the decoding

probabilities for all classes (Liu et al., 2021a; Nour et al., 2021). Class weights were

balanced by adjusting inversely proportional to class frequencies (i.e., trial

numbers) in the training procedure. Additionally, every 5 trials were averaged within

the same class to reduce noises. We then applied the trained classifier to the MEG

signals at each time point in the main phase. Finally, decoding probabilities were

averaged across each condition and motion direction to yield a time course of

decoding probability.

Optimal time point of motion direction representation

The optimal time point of motion direction representation was considered as the

time point with the highest decoding accuracy. To index the optimal time point of each

motion direction for each participant, we conducted time-resolved motion direction

decoding analysis on the functional localizer data using Lasso-regularized logistic

regression models. A leave-one-trial-out cross-validation procedure was used to train

the classifier to determine one of the four motion directions, yielding a decoding

accuracy at each time point for each participant. Finally, the time point with the highest

decoding accuracy was independently extracted for each participant and each motion

direction. These time points are referred to as optimal time points and were used for

sequenceness analysis.

Sequenceness measure

To identify sequenceness during the post-cue blank period in each trial, we

trained models to detect transient spontaneous neural reactivation of each motion

direction representation. Therefore, one-versus-rest Lasso-regularized logistic

regression models were trained separately for each motion direction using the functional

localizer data. MEG signals from the 72 occipital sensors obtained throughout all

scanning sessions were used to train the decoding models. For each motion direction,

we trained a binomial classifier, using positive instances from trials in which that feature

(i.e., one motion direction) was presented and negative instances from trials in which all

other features (i.e., the other three motion directions) were presented, together with an

equivalent amount of null data from the 1–1.5 s ISI. The sensor distributions of beta

estimates and the spatial correlation among the classifiers are shown in Figure S1.

The sequenceness analysis pipeline is illustrated in Figure 3. We first applied the

trained classifiers with the functional localizer data at the optimal time point to MEG

signals at each time point during the blank period, to generate a [time x state]

reactivation probability matrix for each trial. The TDLM framework was then used to

quantify evidence for sequential reactivations consistent with the inferred task transition

structure (Liu et al., 2021b, 2021a; Nour et al., 2021).

TDLM is a multiple linear regression approach to quantify the extent to which a

lagged reactivation time course of state i, $(X(\Delta t)_i, \Delta t$ indicates lag time) can predict the

reactivation time course of state j, (X_j). Two steps were included in this pipeline. First,

we performed multiple separate regressions using each state's reactivation time course

as a dependent variable ($j \in [1: 4]$) and the historical (i.e., time-lagged) reactivation time

courses of all states ($i \in [1: 4]$) as predictor variables.

$$X_j = \sum_{i=1}^{4} X(\Delta t)_i * \beta(\Delta t)_{i,j} + C \tag{1}$$

The regression coefficients from Equation 1 quantify the evidence for each

empirical state $\rightarrow$ state reactivation pattern at a given time lag, Δt. For example, $\beta(\Delta t)_{ij}$ is

the coefficient capturing the unique variance in X_j explained by $X(\Delta t)_i$. All such first-level

coefficients were placed in a lag-specific [4 x 4] empirical transition matrix *B*,

representing evidence for all possible state-to-state transitions at a specific lag.

In the second step, we quantified evidence supporting that the empirical (lag-

specific) transition matrix, *B*, can be predicted by the sequences of interest, i.e., the

exposed motion sequence. All transitions of interest were specified in a model transition

matrix (1 for transitions of interest and 0 otherwise), separately for forward (as in the

visual experience) (T_F), and backward (opposite to the visual experience) direction (T_B).

Then, the strength of all sequences of interest was measured by:

$$\boldsymbol{B} = \sum_{r=1}^{4} Z_r * T_r \qquad (2)$$

where $\boldsymbol{B}$ is the [state x state] empirical transition matrix, T_r is the [state x state] predictor

transition matrix (for regressor r), predicting the empirical transitions $\boldsymbol{B}$. We included four

predictor matrices: (1) T_F: transitions as in the motion sequence in the forward direction

(transitions corresponding to [A → B → C → D]), (2) T_B: transitions opposite to the

experience in the backward direction ([D → C → B → A], T_B is the transpose of T_F), (3)

T_{auto}: self-transitions to control for autocorrelation ([4 x 4] identity matrix), and (4) T_{const}:

a constant matrix to model away the average of all transitions, ensuring that any weight

on T_F and T_B was not due to general background neural dynamics.

Z_r is the scalar regression coefficient quantifying evidence for the hypothesized

sequences in the empirical transitions, i.e., sequence strength. Note, this estimate of

sequence strength is a relative quantity. An estimate of zero for state i to state j, for

example, does not mean lack of replay from i → j but no stronger replay of i → j than

that of other transitions. Repeating the regression in Equation 2 at each time lag ($\Delta t = 4$,

8, 12, ..., 600 ms) results in both forward and backward sequence strength as a function

of time lag. Shorter lags indicate greater time compression, corresponding to faster

speed.

In the current study, sequenceness was defined from the contrast between the

evidence for sequential replay of task structure in the forward ([A → B → C → D])

versus backward ([D → C → B → A]) direction (i.e., Z_1–Z_2); thus, removing between-

participant variance in the sequential replay per se (which may arise secondary to task

engagement and measurement sensitivity (Liu et al., 2021a; Nour et al., 2021)). Positive

sequenceness values indicate replay in a predominantly forward direction, whereas

negative sequenceness values indicate replay in a predominantly backward direction.

For statistical inference, we used nonparametric permutation tests involving all

possible permutations of the stimulus labels at the second-level regression, equivalent

to permuting the rows and columns together of the transition matrices used to calculate

sequenceness. For each permutation, we calculated the peak absolute mean sequence

strength over participants and across lags (controlling for multiple comparisons across

lags). Sequence strength in unpermuted data was considered significant if its absolute

magnitude was >95% of the within-permutation peak.

Identifying replay onsets and analyzing time-frequency

Given that the maximal evidence for replay transitions at the group level was a

32 ms-lag, we next identified replay onsets. Replay onsets were defined as the times

when a strong reactivation of one stimulus (e.g., A) was followed by a strong

reactivation of the next stimulus (e.g., B) in the sequence, with a 32 ms-lag (Liu et al.,

2019; Nour et al., 2021). Specifically, we shifted the reactivation matrix X up to the time

lag Δt for the first time with maximal evidence for replay (i.e., 32 ms), obtained X(Δt).

$$Proj = X(\Delta t) \qquad (3)$$

705   We then multiplied X by the transition matrix P; corresponding to the

706   unscrambled sequences: X x P. The transition matrix P defines the mapping between

707   the task state corresponding to column i in X, and column i in Orig. Specifically, column i

708   in Orig is the reactivation time course of the state preceding state i in T.

$$Orig = X * P \qquad (4)$$

709   Next, we multiplied Proj and Orig element-wise, summing over the columns of

710   the resulting matrix; therefore, creating a [time x 1] vector, R, in which each element

711   (i.e., row) indicates the strength of replay with a 32 ms-lag at a given moment in time.

$$R_t = \sum_{i=1}^{4} Orig_{ti} * Proj_{ti} \qquad (5)$$

Finally, we identified putative replay event onsets by threshold R at its 95^{th}

percentile preceded by a 100 ms pre-onset baseline exhibiting a low probability of

replay at each time point.

We then epoched MEG data in the blank period surrounding each onset and

computed a frequency decomposition (wavelet transformation) in the time window of

−100–150 ms using all sensors. We aimed to find an evidence for the power increase in

the high-frequency (120–160 Hz) region of interest at replay onset, compared with a

pre-onset baseline (−100–−50 ms from onset), as mentioned in our previous study (Liu

et al., 2019). Finally, we generated two separate [time x frequency] matrices (i.e., using

forward and backward task transition matrices separately) by averaging estimates over

sensors and events, capturing the typical spectrally-resolved power change at replay

onset.

MEG source reconstruction

We identified the neural sources associated with increased ripple power at

putative replay onsets. Forward models were generated based on a single shell using

the superposition of basic functions that approximately correspond to the plane

tangential to the MEG sensor array. Linearly constrained minimum variance

beamforming (Van Veen et al., 1997) was used to reconstruct the epoched MEG data to

a grid in MNI space (grid step, 5 mm). The sensor covariance matrix for beamforming

was estimated using broadband power data across all frequencies. The baseline activity

was the mean activity averaged over $-100 - -50$ ms relative to replay onset. All

nonartifactual replay epochs were baseline-corrected at source level. We obtained

whole-brain results for voxels predicted by participant-specific ripple power at replay

onset. Nonparametric permutation tests were performed on the volume of interest to

compute the multiple comparison P values of clusters >10 voxels (whole-brain

corrected, cluster-defining threshold; $t = 3.1$, $n = 5000$ permutations).

Data and Code availability

739   Data used to generate the findings of this study will be available upon request (subject

740   to participant consent) to the Lead Contact. TDLM MATLAB Code available at:

741   https://github.com/YunzheLiu/TDLM. Custom computer code will be made available

742   upon request to the Lead Contact.

AUTHOR CONTRIBUTIONS

Conceptualization, T.H., Y.L., and F.F.; Investigation, T.H., Q.W., X.G., and X.Z.; Writing

– Original Draft, T.H.; Writing – Review & Editing, T.H., Q.W., X.Z., Y.L., and F.F.

ACKNOWLEDGEMENTS

This study was supported by National Science and Technology Innovation 2030 Major

Program (2022ZD0204802), the National Natural Science Foundation of China

(31930053), Beijing Natural Science Foundation (5244044), and the Young Scientists

Fund of the Humanities and Social Science Foundation of Ministry of Education of

China (23YJCZH071).

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Alvarez P, Squire LR (1994) Memory consolidation and the medial temporal lobe: a
simple network model. Proc Natl Acad Sci U S A 91:7041–7045.

Ambrose RE, Pfeiffer BE, Foster DJ (2016) Reverse Replay of Hippocampal Place Cells
Is Uniquely Modulated by Changing Reward. Neuron 91:1124–1136.

Baker R, Dexter M, Hardwicke TE, Goldstone A, Kourtzi Z (2014) Learning to predict:
Exposure to temporal sequences facilitates prediction of future events. Vision
Res 99:124–133.

Buch ER, Claudino L, Quentin R, Bönstrup M, Cohen LG (2021) Consolidation of
human skill linked to waking hippocampo-neocortical replay. Cell Rep 35:109193.

Buonomano DV, Merzenich MM (1998) CORTICAL PLASTICITY: From Synapses to
Maps. Annual Review of Neuroscience 21:149–186.

Bush D, Ólafsdóttir HF, Barry C, Burgess N (2022) Ripple band phase precession of
place cell firing during replay. Current Biology 32:64-73.e5.

Buzsáki G (1986) Hippocampal sharp waves: Their origin and significance. Brain
Research 398:242–252.

Buzsáki G (1996) The hippocampo-neocortical dialogue. Cereb Cortex 6:81–92.

Buzsáki G (2015) Hippocampal sharp wave-ripple: A cognitive biomarker for episodic
memory and planning. Hippocampus 25:1073–1188.

Carr MF, Jadhav SP, Frank LM (2011) Hippocampal replay in the awake state: a
potential substrate for memory consolidation and retrieval. Nat Neurosci 14:147–
153.

Chen N, Cai P, Zhou T, Thompson B, Fang F (2016) Perceptual learning modifies the
functional specializations of visual cortical areas. Proceedings of the National
Academy of Sciences 113:5724–5729.

Chun MM, Phelps EA (1999) Memory deficits for implicit contextual information in
amnesic subjects with hippocampal damage. Nat Neurosci 2:844–847.

Costandi M (2016) Neuroplasticity. MIT Press.

Demarchi G, Sanchez G, Weisz N (2019) Automatic and feature-specific prediction-
related neural activity in the human auditory system. Nat Commun 10:3440.

Diba K, Buzsáki G (2007) Forward and reverse hippocampal place-cell sequences
during ripples. Nat Neurosci 10:1241–1242.

Dimakopoulos V, Mégevand P, Stieglitz LH, Imbach L, Sarnthein J (2022) Information
flows from hippocampus to auditory cortex during replay of verbal working
memory items Griffiths TD, Colgin LL, eds. eLife 11:e78677.

Eagleman SL, Dragoi V (2012) Image sequence reactivation in awake V4 networks.
PNAS 109:19450–19455.

Eichenbaum H, Yonelinas AR, Ranganath C (2007) The Medial Temporal Lobe and
Recognition Memory. Annu Rev Neurosci 30:123–152.

Ekman M, Kok P, Lange FP de (2017) Time-compressed preplay of anticipated events
in human primary visual cortex. Nat Commun 8:15276.

Ekman M, Kusch S, de Lange FP (2023) Successor-like representation guides the
prediction of future events in human visual cortex and hippocampus Barense M,
Baker CI, Barron H, eds. eLife 12:e78904.

Fang F, Murray SO, Kersten D, He S (2005) Orientation-Tuned fMRI Adaptation in
Human Visual Cortex. Journal of Neurophysiology 94:4188–4195.

Finnie PSB, Komorowski RW, Bear MF (2021) The spatiotemporal organization of
experience dictates hippocampal involvement in primary visual cortical plasticity.
Current Biology 31:3996-4008.e6.

Foster DJ, Wilson MA (2006) Reverse replay of behavioural sequences in hippocampal
place cells during the awake state. Nature 440:680.

Garvert MM, Dolan RJ, Behrens TE (2017) A map of abstract relational knowledge in
the human hippocampal–entorhinal cortex Davachi L, ed. eLife 6:e17086.

Gavornik JP, Bear MF (2014) Learned spatiotemporal sequence recognition and
prediction in primary visual cortex. Nat Neurosci 17:732–737.

Gillespie AK, Maya DAA, Denovellis EL, Liu DF, Kastner DB, Coulter ME, Roumis DK,
Eden UT, Frank LM (2021) Hippocampal replay reflects specific past experiences
rather than a plan for subsequent choice. Neuron 109:3149-3163.e6.

Gothard KM, Skaggs WE, Moore KM, McNaughton BL (1996) Binding of hippocampal
CA1 neural activity to multiple reference frames in a landmark-based navigation
task. J Neurosci 16:823–835.

Gridchyn I, Schoenenberger P, O'Neill J, Csicsvari J (2020) Assembly-Specific
Disruption of Hippocampal Replay Leads to Selective Memory Deficit. Neuron
106:291-300.e6.

Gutnisky DA, Hansen BJ, Iliescu BF, Dragoi V (2009) Attention Alters Visual Plasticity
during Exposure-Based Learning. Current Biology 19:555–560.

Hannula DE, Tranel D, Cohen NJ (2006) The Long and the Short of It: Relational
Memory Impairments in Amnesia, Even at Short Lags. J Neurosci 26:8352–8359.

Henke K (2010) A model for memory systems based on processing modes rather than
consciousness. Nat Rev Neurosci 11:523–532.

Hindy NC, Ng FY, Turk-Browne NB (2016) Linking pattern completion in the
hippocampus to predictive coding in visual cortex. Nat Neurosci 19:665–667.

Hsieh L-T, Gruber MJ, Jenkins LJ, Ranganath C (2014) Hippocampal Activity Patterns
Carry Information about Objects in Temporal Context. Neuron 81:1165–1178.

Igata H, Ikegaya Y, Sasaki T (2021) Prioritized experience replays on a hippocampal
predictive map for learning. Proceedings of the National Academy of Sciences
118:e2011266118.

Ji D, Wilson MA (2007) Coordinated memory replay in the visual cortex and
hippocampus during sleep. Nature Neuroscience 10:100–107.

Joo HR, Frank LM (2018) The hippocampal sharp wave–ripple in memory retrieval for
immediate use and consolidation. Nat Rev Neurosci 19:744–757.

Kok P, Failing M, de Lange F (2014) Prior Expectations Evoke Stimulus Templates in
the Primary Visual Cortex. Journal of cognitive neuroscience 26.

Kok P, Jehee JFM, de Lange FP (2012) Less Is More: Expectation Sharpens
Representations in the Primary Visual Cortex. Neuron 75:265–270.

Kok P, Mostert P, Lange FP de (2017) Prior expectations induce prestimulus sensory
templates. PNAS 114:10473–10478.

Kok P, Turk-Browne NB (2018) Associative Prediction of Visual Shape in the
Hippocampus. J Neurosci 38:6888–6899.

Konkel A, Warren D, Duff M, Tranel D, Cohen N (2008) Hippocampal amnesia impairs
all manner of relational memory. Frontiers in Human Neuroscience 2 Available at:
https://www.frontiersin.org/article/10.3389/neuro.09.015.2008 [Accessed May 26,
2022].

Kurth-Nelson Z, Economides M, Dolan RJ, Dayan P (2016) Fast Sequences of Non-
spatial State Representations in Humans. Neuron 91:194–204.

Lee AK, Wilson MA (2002) Memory of Sequential Experience in the Hippocampus
during Slow Wave Sleep. Neuron 36:1183–1194.

Li W (2016) Perceptual Learning: Use-Dependent Cortical Plasticity. Annual Review of
Vision Science 2:109–130.

Liu Y, Dolan RJ, Higgins C, Penagos H, Woolrich MW, Ólafsdóttir HF, Barry C, Kurth-
854   Nelson Z, Behrens TE (2021a) Temporally delayed linear modelling (TDLM)
855   measures replay in both animals and humans Colgin LL, Kemere C, van der
856   Meer M, eds. eLife 10:e66917.

857   Liu Y, Dolan RJ, Kurth-Nelson Z, Behrens TEJ (2019) Human Replay Spontaneously
858   Reorganizes Experience. Cell 178:640-652.e14.

859   Liu Y, Mattar MG, Behrens TEJ, Daw ND, Dolan RJ (2021b) Experience replay is
860   associated with efficient nonlocal learning. Science 372 Available at:
861   https://science.sciencemag.org/content/372/6544/eabf1357 [Accessed May 21,
862   2021].

863   Liu Y, Nour MM, Schuck NW, Behrens TEJ, Dolan RJ (2022) Decoding cognition from
864   spontaneous neural activity. Nat Rev Neurosci:1–11.

865   Lu J, Luo L, Wang Q, Fang F, Chen N (2020) Cue-triggered activity replay in human
866   early visual cortex. Sci China Life Sci Available at: /doi/10.1007/s11427-020-
867   1726-5 [Accessed June 29, 2020].

868   Lu Z-L, Dosher BA (2022) Current directions in visual perceptual learning. Nat Rev
869   Psychol 1:654–668.

870   Marr D, Brindley GS (1971) Simple memory: a theory for archicortex. Philosophical
871   Transactions of the Royal Society of London B, Biological Sciences 262:23–81.

872   McFadyen J, Liu Y, Dolan RJ (2023) Differential replay of reward and punishment paths
873   predicts approach and avoidance. Nat Neurosci 26:627–637.

874   Mo C, Lu J, Wu B, Jia J, Luo H, Fang F (2019) Competing rhythmic neural
875   representations of orientations during concurrent attention to multiple orientation
876   features. Nat Commun 10:1–10.

877   Nour MM, Liu Y, Arumuham A, Kurth-Nelson Z, Dolan RJ (2021) Impaired neural replay
878   of inferred relationships in schizophrenia. Cell 184:4315-4328.e17.

879   O'Keefe J, Nadel L (1978) The hippocampus as a cognitive map. Oxford : New York:
880   Clarendon Press ; Oxford University Press.

881   Ólafsdóttir HF, Barry C, Saleem AB, Hassabis D, Spiers HJ (2015) Hippocampal place
882   cells construct reward related sequences through unexplored space Eichenbaum
883   H, ed. eLife 4:e06063.

Ólafsdóttir HF, Carpenter F, Barry C (2016) Coordinated grid and place cell replay
during rest. Nat Neurosci 19:792–794.

Ólafsdóttir HF, Carpenter F, Barry C (2017) Task Demands Predict a Dynamic Switch in
the Content of Awake Hippocampal Replay. Neuron 96:925-935.e6.

Pojoga SA, Kharas N, Dragoi V (2020) Perceptually unidentifiable stimuli influence
cortical processing and behavioral performance. Nat Commun 11:6109.

Redish AD, Rosenzweig ES, Bohanick JD, McNaughton BL, Barnes CA (2000)
Dynamics of Hippocampal Ensemble Activity Realignment: Time versus Space. J
Neurosci 20:9298–9309.

Redish AD, Touretzky DS (1998) The role of the hippocampus in solving the Morris
water maze. Neural Comput 10:73–111.

Rivard B, Li Y, Lenck-Santini P-P, Poucet B, Muller RU (2004) Representation of objects
in space by two classes of hippocampal pyramidal cells. J Gen Physiol 124:9–25.

Roelfsema PR, de Lange FP (2016) Early Visual Cortex as a Multiscale Cognitive
Blackboard. Annual Review of Vision Science 2:131–151.

Rolls ET (2000) Hippocampo-cortical and cortico-cortical backprojections. Hippocampus
10:380–388.

Sasaki Y, Nanez JE, Watanabe T (2010) Advances in visual perceptual learning and
plasticity. Nat Rev Neurosci 11:53–60.

Schapiro AC, Gregory E, Landau B, McCloskey M, Turk-Browne NB (2014) The
necessity of the medial temporal lobe for statistical learning. J Cogn Neurosci
26:1736–1747.

Staresina BP, Davachi L (2009) Mind the Gap: Binding Experiences across Space and
Time in the Human Hippocampus. Neuron 63:267–276.

Taulu S, Simola J (2006) Spatiotemporal signal space separation method for rejecting
nearby interference in MEG measurements. Phys Med Biol 51:1759–1768.

Turk-Browne NB, Scholl BJ, Chun MM, Johnson MK (2009) Neural evidence of
statistical learning: efficient detection of visual regularities without awareness. J
Cogn Neurosci 21:1934–1945.

Turner W, Blom T, Hogendoorn H (2023) Visual Information Is Predictively Encoded in
914   Occipital Alpha/Low-Beta Oscillations. J Neurosci 43:5537–5545.

915   Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A (1997) Localization of brain
916   electrical activity via linearly constrained minimum variance spatial filtering. IEEE
917   Trans Biomed Eng 44:867–880.

918   Watanabe T, Sasaki Y (2015) Perceptual learning: Toward a comprehensive theory.
919   Annu Rev Psychol 66:197–221.

920   Whittington JCR, Muller TH, Mark S, Chen G, Barry C, Burgess N, Behrens TEJ (2020)
921   The Tolman-Eichenbaum Machine: Unifying Space and Relational Memory
922   through Generalization in the Hippocampal Formation. Cell 183:1249-1263.e23.

923   Wimmer GE, Liu Y, Vehar N, Behrens TEJ, Dolan RJ (2020) Episodic memory retrieval
924   success is associated with rapid replay of episode content. Nature Neuroscience
925   23:1025–1033.

926   Wyart V, Nobre AC, Summerfield C (2012) Dissociable prior influences of signal
927   probability and relevance on visual contrast sensitivity. PNAS 109:3593–3598.

928   Xu S, Jiang W, Poo M, Dan Y (2012) Activity recall in a visual cortical ensemble. Nature
929   Neuroscience 15:449–455.

SUPPLEMENTARY INFORMATION

Figure S1. Sensor maps and spatial correlation of trained Lasso logistic

regression models (related to Figure 3)

(A) Left: Sensor map for each state decoding model in Experiment 1; MEG signals from

72 occipital sensors were selected as features while training the classifier. Right:

Correlation matrix among classifiers. No spatial correlation was found among trained

classifiers (highest correlation; $r < 0.12$). **(B and C)** Same as (A), except the sensor

maps and correlation matrix correspond to Experiments 2 (Panel B, highest correlation;

$r < 0.1$) and 3 (Panel C, highest correlation; $r < 0.1$), respectively.

Figure S2. Decoded feature representations in start- and end-only conditions

(related to Figures 3)

Time series of reactivation probability output from the four regression models on an example trial. Time zero corresponds to trial onset. Both participants S08 and S12 were exposed with a fixed motion sequence of 0° → 90° → 180° → 270°.

Figure S3. Sequenceness distribution across participants (related to Figures 4 and 5)

(A) In Experiment 1, of the 18 participants, 14 showed backward replay at 32 ms-lag for

the start-only condition (left), and 13 showed backward replay at 32 ms-lag for the end-

only condition (right). Each dot represents an individual participant. Backward and

forward sequenceness is denoted by blue and yellow dots, respectively. The inset

histogram shows the distribution of deviations from the unity line. **(B)** Same as

described in (A), but for the results of Experiment 2 in the 2nd-only (left) and 3rd-only

(right) conditions, respectively. **(C)** Same as described in (A) but for the results of

Experiment 3 in the start- (left) and end-only (right) conditions, respectively.

Figure S4. Sequenceness for each of the 24 possible orders (related to Figure 4)

(A) In the start-only condition of Experiment 1, only the backward sequence [4 → 3 → 2

→ 1] with lags at 28–40 ms was significantly different (gray panels). The horizontal

dashed lines represent significance thresholds derived from state label permutation at

the 2nd-level GLM matrix of temporally delayed linear modeling. **(B)** Same as described

in (A) but for the end-only condition in Experiment 1.

Figure S5. Stimuli and experimental procedures in Experiments 2 and 3 (related to

Figure 5)

(A) We presented the second (2nd-only condition) or third RDK (3rd-only condition) as a

cue in Experiment 2. **(B)** The procedure of Experiment 3 was identical to that of

Experiment 1 except for removing the exposure phase.