Non-feature-specific elevated responses and feature-specific backward replay in human brain induced by visual sequence exposure

Tao He[1,2,3,4], Xizi Gong[3,4], Qian Wang[3,4], Xinyi Zhu[3,4], Yunzhe Liu[5], Fang Fang[3,4,6,7*]

[1] Center for the Cognitive Science of Language, Beijing Language and Culture University, Beijing 100083, People's Republic of China

[2] Key Laboratory of Language Cognitive Science (Ministry of Education), Beijing Language and Culture University, Beijing 100083, People's Republic of China

[3] School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing 100871, People's Republic of China

[4] IDG/McGovern Institute for Brain Research, Peking University, Beijing 100871, People's Republic of China

[5] Chinese Institute for Brain Research, Beijing 102206, People's Republic of China

[6] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, People's Republic of China

[7] Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing 100871, People's Republic of China

Corresponding author:

Fang Fang, Ph.D. Professor

School of Psychological and Cognitive Sciences

Peking University

Beijing, P.R. China

Tel & Fax: +86-10-62756932   Email: ffang@pku.edu.cn

Web: www.psy.pku.edu.cn/english/people/faculty/professor/fangfang/index.htm

ABSTRACT

The ability of cortical circuits to adapt in response to experience is a fundamental

property of the brain. After exposure to a moving dot sequence, flashing a dot as a cue

at the starting point of the sequence can elicit successive elevated responses even in

the absence of the sequence. These cue-triggered elevated responses have been

shown to play a crucial role in predicting future events in dynamic environments.

However, temporal sequences we are exposed typically contain rich feature information.

It remains unknown whether the elevated responses are feature specific and, more

crucially, how the brain organizes sequence information after exposure. To address

these questions, participants were exposed to a predefined sequence of four motion

directions for about 30 min, followed by the presentation of the start or end motion

direction of the sequence as a cue. Surprisingly, we found that cue-triggered elevated

responses were not specific to any motion direction. Interestingly, motion direction

information was spontaneously reactivated, and the motion sequence was backward

replayed in a time-compressed manner. These effects were observed even after brief

exposure. Notably, no replay events were observed when the second or third motion

direction of the sequence served as a cue. Further analyses revealed that activity in the

medial temporal lobe (MTL) preceded the ripple power increase in visual cortex at the

onset of replay, implying a coordinated relationship between the activities in the MTL

and visual cortex. Together, these findings demonstrate that visual sequence exposure

induces two-fold brain plasticity that may simultaneously serve for different functional

purposes.   The   non-feature-specific   elevated   responses   may   facilitate   general

processing of upcoming stimuli, whereas the feature-specific backward replay may

underpin passive learning of visual sequence.

INTRODUCTION

The capacity of cortical circuits to undergo plasticity in response to experience is a fundamental feature of the brain (Buonomano and Merzenich, 1998; Costandi, 2016; Li, 2016). This plasticity can be induced not only by active, task-dependent training, such as visual perceptual learning (Watanabe and Sasaki, 2015; Chen et al., 2016; Lu and Dosher, 2022), but also by passive, repetitive exposure (Gutnisky et al., 2009; Sasaki et al., 2010; Ekman et al., 2017). For instance, after repeated exposure to a moving dot sequence, flashing a dot (i.e., cue) at the starting point of the sequence elicits successive elevated neural responses in visual cortex, akin to those induced by the actual moving dot sequence in both humans (Ekman et al., 2017, 2023; Lu et al., 2020) and animals (Eagleman and Dragoi, 2012; Xu et al., 2012). This cue-triggered reactivation is thought to be driven by expectations and is prediction-related (Ekman et al., 2017, 2023), as it facilitates the prediction of upcoming stimuli and influences the perception of sensory information (Gutnisky et al., 2009; Baker et al., 2014; Pojoga et al., 2020).

Unlike the simple white dot sequences used in previous studies (Xu et al., 2012; Ekman et al., 2017, 2023; Lu et al., 2020), temporal sequences to which we are exposed in daily life usually contain rich feature information. However, it remains unknown whether cue-triggered elevated responses are feature-specific. On the one hand, if these responses are not feature-specific, they may reflect a general state of cortical readiness for any upcoming stimuli. On the other hand, if these responses are indeed specific to a particular feature in the sequence, they would selectively facilitate

the processing of specific future events. This aligns with the view that expectation

sharpens neural tuning and enhances the processing of expected stimuli (Kok et al.,

2012). For instance, expectation has been shown to elevate the prestimulus baseline

activity of sensory neurons tuned to expected stimuli (Wyart et al., 2012; Kok et al.,

2014) and to preactivate stimulus templates in both the visual (Kok et al., 2017) and

auditory (Demarchi et al., 2019) cortices. Notably, however, all these prediction-related,

feature-specific activities were observed in static contexts. Whether prediction-related

responses in dynamic temporal contexts (e.g., visual sequences) are feature-specific

remains unclear.

After exposure to a feature-contained temporal sequence, another important

question is how the feature information in the sequence is encoded and organized in

neural activity following the cue. Previous studies have demonstrated that memory

consolidation (Carr et al., 2011; Gridchyn et al., 2020; Gillespie et al., 2021) or learning

process (Igata et al., 2021; Liu et al., 2021b, 2022) following training with a temporal

sequence is associated with a neural phenomenon known as replay. Replay refers to

the sequential reactivation of neural activity patterns associated with the trained

sequence in both sleep (Lee and Wilson, 2002) and awake (Foster and Wilson, 2006)

states. During replay, the neural representation of the trained sequence is temporally

compressed and can occur in either a forward or backward direction (Carr et al., 2011;

Joo and Frank, 2018; Nour et al., 2021; Liu et al., 2022; McFadyen et al., 2023). These

replay events frequently coincide with sharp-wave ripples (SWRs), which are high-

frequency (150–220 Hz) oscillations (O'Keefe and Nadel, 1978; Bush et al., 2022)

detected in hippocampal local field potentials (Buzsáki, 1986, 2015). To date, replay has

been observed in tasks involving sequences, such as nonlocal reinforcement learning

(Liu et al., 2021b) and episodic memory retrieval (Wimmer et al., 2020). It remains

unclear whether simple visual exposure to temporal sequences could also trigger replay

events in humans.

In the current study, we used magnetoencephalography (MEG) to investigate

whether cue-triggered elevated brain responses following visual sequence exposure are

feature-specific and, more importantly, to determine how feature information in the

exposed sequence is encoded and organized in the brain after exposure. To address

these questions, participants were initially exposed to a predetermined motion

sequence. We then decoded motion direction information during the blank period

following the presentation of either the first or last motion direction in the sequence as a

cue. Surprisingly, we found that cue-triggered elevated responses were not specific to

any motion direction. Interestingly, motion direction information was spontaneously

reactivated during the blank period, and the motion sequence was backward replayed in

a time-compressed manner. This backward replay was identified even after brief

exposure. However, neither forward nor backward replay was detected when an

intermediate motion direction in the sequence was presented as a cue. Lastly, MEG

source reconstruction analysis revealed that medial temporal lobe (MTL) activation

preceded visual cortical activation, implying that the replay events observed in visual

cortex may be triggered by activities in the MTL.

RESULTS

116  The visual stimuli used in the study were four random-dot kinematograms

117 (RDKs). All dots in an RDK moved in a single direction (i.e., 0°, 90°, 180°, or 270°) with

118 100% coherence. In Experiment 1, we included three trial conditions: full sequence

119 trials, start-only trials, and end-only trials (**Figure 1A**). In a full sequence trial,

120 participants were exposed to a predefined sequence of the four RDKs presented either

121 clockwise (i.e., 0° → 90° → 180° → 270°) or counterclockwise at the center of the

122 screen, with an inter-stimulus interval (ISI) of 300 ms between every two RDKs. In a

123 start- or end-only trial, the first or last RDK of the sequence was presented at the

124 beginning of the trial, followed by a blank period of 2.4 s. Participants were instructed to

125 complete three successive phases: functional localizer phase, exposure phase, and

126 main phase (**Figure 1B**). The functional localizer data were used to train models to

127 decode each motion direction in the sequence. During this phase, one of the four RDKs

128 was randomly presented for 1 s in each trial. During the exposure phase, participants

129 were exposed only to full sequence trials for about 30 min. In the main phase, 50%,

130 25%, and 25% of all trials were full sequence, start- and end-only trials, respectively.

131 The full sequence trials served as a topping-up exposure to maintain the exposure

132 effect.

133 **Cue-triggered elevated responses are not feature-specific**

134  We first measured the event-related field (ERF) activity evoked by RDKs in the

135 three trial conditions using all occipital gradiometer sensors (see Methods). **Figure 1C**

136 shows the evoked responses for full sequence trials (left panel, cluster-based

137 permutation test with a cluster forming threshold of t > 3.1 and 5000 permutations).

Remarkably, start- and end-only trials elicited similar wave-like responses as the full

sequence trials, despite the absence of stimuli following the first RDK (**Figure 1C**,

middle and right panels, cluster-based permutation test with a cluster forming threshold

of t > 3.1 and 5000 permutations). To further quantify the ERF peak amplitudes in the

three conditions while mitigating baseline confounds, we calculated each peak

amplitude using the 300 ms blank period just preceding the onset of the corresponding

RDK as the baseline (see Methods). We found that the four successive RDKs in the full

sequence trials evoked four comparable peaks after stimulus onset (**Figure 1C**, inset

figure in the left panel; two-tailed t-test; all $ts_{(17)}$ > 6.8072, all ps < 10^{-5}). Interestingly, in

start- and end-only trials, significant peaks were still observed during the periods

corresponding to the intervals of the second and third RDKs in the full sequence trials

(**Figure 1C**, inset figures in the middle and right panels; start-only condition, two-tailed t-

test; all $ts_{(17)}$ > 4.5790, all ps < 10^{-3}; end-only condition, two-tailed t-test; all $ts_{(17)}$ >

6.0140, all ps < 10^{-4}). However, no significant peak was observed during the period

corresponding to the interval of the last RDK (start-only condition, two-tailed t-test; $t_{(17)}$ =

1.3221, p = 0.2036; end-only condition, two-tailed t-test; $t_{(17)}$ = 0.2473, p = 0.8076).

These results demonstrate cue-triggered elevated neural responses following visual

exposure to the RDK sequences, resembling previous studies using simple white dot

sequences (Ekman et al., 2017; Lu et al., 2020).

Figure 1. Stimuli, experimental procedure and evoked responses in Experiment 1

(A) Participants were presented with RDKs in three conditions. In the full sequence trial condition, four successive RDKs were presented. In the start- or end-only condition, only the first or last RDK in the sequence was presented at the beginning of the trial. **(B)** During MEG scanning, the participants were presented with two functional localizer runs (i.e., functional localizer phase) before providing any sequence information. Next, they were exposed with four full sequence runs (i.e., exposure phase). Finally, in the main

phase, full sequence, start- and end-only trials were presented in a pseudo-randomized

order. **(C)** Evoked neural responses as a function of time relative to trial onset in the full

sequence, start- and end-only trial conditions, respectively. Bold black lines at the

bottom indicate temporal clusters in which they reached significance when compared to

the pre-stimulus baseline. Inset figures at the top-right corner show the peak amplitudes

during the four corresponding RDK intervals after baseline correction.

Given that we observed elevated responses even in the absence of stimuli

following the cue, we next examined whether these responses were specific to a

particular feature (i.e., motion direction). Specifically, we asked whether motion

directions could be successfully decoded in start- and end-only trials, particularly during

the blank periods corresponding to the intervals of the second and third RDKs in full

sequence trials. To this end, we applied a time-resolved decoding analysis. For each

participant, we trained a one-versus-rest Lasso logistic regression model using the

functional localizer data to classify the neural activity pattern elicited by each motion

direction in the main phase. MEG signals from 72 occipital sensors were selected as

features for the training model.

To validate the reliability of our model, we first used a leave-one-out cross-

validation scheme on the localizer data to independently estimate decoding accuracy at

each time point. At the group level, decoding accuracies peaked at 411 ms, 464 ms, 444

ms, and 434 ms after stimulus onset for motion directions 0° (55.37% $\pm$ 1.21), 90°

(58.10% $\pm$ 1.14), 180° (54.07% $\pm$ 1.47), and 270° (53.64% $\pm$ 1.34), respectively. There

were no significant differences among the four motion directions in terms of either

latency ($F(3, 51) = 0.375$, $p = 0.7716$, $\eta_p^2 = 0.0159$) or decoding accuracy ($F(3, 51) =$

2.757, $p = 0.0517$, $\eta_p^2 = 0.091$) at the peak time point. Next, we trained a model using

the localizer data averaged between 100 ms and 500 ms after stimulus onset. The

trained model was then applied to the MEG signals recorded in the three trial conditions

in the main phase. Finally, we calculated the decoding probability (Liu et al., 2019; Nour

et al., 2021; Turner et al., 2023) for each motion direction at each time point at the group

level (**Figure 2**). Here, decoding probability for each motion direction reflects the

likelihood that the decoded stimulus had a specific motion direction at a given time point

(see Methods). Therefore, it provides a time-resolved decoding preference for each

motion direction, rather than only a single decoded label (e.g., 0°, 90°, 180°, or 270°).

In full sequence trials, the motion direction information could be successfully

decoded, as evidenced by the highest decoding probability during the interval

corresponding to the presentation of the respective RDK (**Figure 2A and 2B**, left

panels). In start- and end-only trials, we could also reliably decode the motion direction

of the first RDK (i.e., the cue) after stimulus onset (**Figure 2A and 2B**, middle and right

panels, only the decoding probability of the first RDK surpass the peak-level

significance threshold obtained from a nonparametric permutation test, FWE corrected

across time). Surprisingly, however, subsequent motion direction information was

absent at the group level during the post-cue blank period, where the cue-triggered

elevated responses were previously observed (i.e., 0.4 – 2.8 s after stimulus onset).

Together, these results demonstrate that the cue-triggered elevated response induced

by visual sequence exposure was not consistently specific to a particular feature across

participants and trials.

Figure 2. Time-resolved decoding probability for each motion direction

(A) Visualization of the time-resolved decoding probability in the three trial conditions. Each row shows the decoding probabilities for the four motion directions at that time point and each column indicates one of the four motion directions. **(B)** The line plots of the time-resolved decoding probability for the three trial conditions. Each colored line shows the time course of the decoding probability for each motion direction. For the start- and end-only conditions, we calculated the permutation threshold estimated by randomly shuffling of the labels and re-decoding, only the decoding probability of the cue surpassed the threshold.

Time-compressed backward replay of exposed motion sequence

How is the motion direction information encoded and organized in the brain

during the post-cue blank period? Clearly, the motion direction representation is not

time-locked to the onset of the cue in either start- or end-only trials. However, it is

possible that individual motion directions are encoded in the MEG signals in a

spontaneous way but are sequentially organized (Liu et al., 2019).

To test this hypothesis, we trained four decoding models using the localizer data

to capture the neural features of the four motion directions. We employed four decoding

models because our aim was to build feature-specific classifiers, each sensitive to only

one motion direction. These classifiers were designed to quantify the evidence of

feature-specific sequence in subsequent analyses. The models used a one-versus-rest

Lasso logistic regression algorithm, with MEG signals from 72 occipital sensors as

features (**Figure 3A**). For each participant and motion direction, we then selected the

time point with the highest decoding accuracy, which was estimated during the

validation of the functional localizer data, as the optimal time point. These optimal time

points were chosen because they are believed to carry the richest feature information

(Mo et al., 2019). For subsequent analyses, we obtained four classifiers that were

trained on the localizer data at their respective optimal time points. Each classifier

yielded significant decoding probability only when the stimulus matched the motion

direction it was trained to detect (**Figure 3B**). We did not find any spatial correlation

between any two trained classifiers (highest correlation $r < 0.12$, Figure S1A). Since the

optimal time point varied across participants and motion directions, we circularly shifted

the decoding probabilities over time and aligned them to a common time point

(arbitrarily set to 200 ms after stimulus onset) for visualization at the group level (**Figure**

3B).

Having established the classifiers for each participant and motion direction, we

then applied the classifiers to the MEG signals during the post-cue blank period in start-

and end-only conditions to estimate the reactivation probability (i.e., decoding

probability) for each motion direction at each time point (**Figure 3C;** Liu et al., 2019;

Wimmer et al., 2020; Nour et al., 2021). Example trials for the start- and end-only

conditions are shown in Figure S2. The results revealed that motion direction

reactivations occurred sparsely during the post-cue blank period, rather than crowded

within intervals corresponding to the RDK presentations in the full sequence condition,

suggesting that motion direction information might be reactivated in a spontaneous way.

Next, we applied temporally delayed linear modeling (TDLM) to quantify whether and

how these spontaneous reactivations followed the order of the exposed sequence

(**Figure 3D**; Kurth-Nelson et al., 2016; Liu et al., 2019, 2021a; Nour et al., 2021). This

algorithm includes two-level regression analyses. The first-level regression quantifies

the evidence for each pairwise transition (e.g., $0° \rightarrow 90°$), resulting in an empirical

transition matrix. The second-level regression evaluates the extent to which this

empirical transition matrix aligns with a specific sequence of interest. Finally, we defined

"sequenceness" as a metric of forward (i.e., $0° \rightarrow 90° \rightarrow 180° \rightarrow 270°$) or backward

(i.e., $270° \rightarrow 180° \rightarrow 90° \rightarrow 0°$) replay (see Methods).

Figure 3. Illustration of replay analysis pipeline: Temporally delayed linear modeling (TDLM)

(A) A Lasso logistic regression model was trained for each participant and motion direction using MEG signals from the functional localizer data. **(B)** The four feature-specific models that were trained using functional localizer data. The decoding probabilities were aligned at 200 ms post-stimulus onset according to their corresponding optimal time point per participant and motion direction. Each colored line indicates the decoding results of a model applied to a motion direction dataset. The

dashed horizontal line indicates the permutation threshold estimated by random

shuffling of the labels and re-decoding. **(C)** The four models were next applied to MEG

signals during the post-cue blank period to derive a decoded reactivation matrix [time x

motion direction]. An illustration of backward sequential reactivations of motion direction

is shown on the left. Reactivation probabilities correspond to the decoding probabilities

derived from the four models. **(D)** Using TDLM, we quantified the evidence for

sequential replay of the motion sequence during the post-cue blank period in start- and

end-only conditions. We first performed a time-lagged regression to generate a [4 x 4]

empirical regression coefficient matrix for each time lag by regressing each lagged

predictor matrix, $X(\Delta t)$, onto the original reactivation matrix, Y (i.e., 1^{st} level GLM

analysis). Next, we used a second-level GLM analysis to evaluate the extent to which

the empirical transition matrix follows a model transition matrix (e.g., forward or

backward transitions) (i.e., 2^{nd} level GLM analysis). Finally, we calculated the difference

between the second-level regression coefficients for forward and backward transitions,

referred to as "sequenceness". We tested the magnitude of this "sequenceness" at each

time lag independently for all transition lags up to 600 ms. The dashed line represents

the corrected nonparametric statistical significance threshold. The green area indicates

the lags when the evidence of "sequenceness" in the backward direction exceeded the

permutation threshold.

As shown in **Figure 4A** and **4B**, in the start-only condition, we found evidence of

backward replay of the exposed motion sequence (i.e., the replay sequence was 270°

→ 180° → 90° → 0° when the exposed motion sequence was 0° → 90° → 180° →

270°) during the post-cue blank period, with a peak transition lag at 28 – 40 ms

(maximal effect at 32 ms-lag: β = −0.0202 ± 0.002, p < 1/24 ≈ 0.042, peak-level

significance threshold derived from a nonparametric permutation test, FWE corrected

across lags, **Figure 4B**). For visualization, an example of backward replay of the motion

sequence is illustrated in **Figure 4A**. This effect was observed in most participants

(Figure S3A). In the end-only condition, we also found evidence of backward replay

during the post-cue blank period, with a peak transition lag at 28–36 ms (maximal effect

at 32 ms-lag: β = −0.0145 ± 0.0016, p < 1/24 ≈ 0.042, peak-level significance threshold

obtained from a nonparametric permutation test, FWE corrected across lags; **Figure**

4E). **Figure 4D** shows an example of backward replay of the motion sequence in this

condition, found in most participants (Figure S3A). Note that the time lag in the

horizontal axis in **Figure 4B** and **4E** indicates the interval between the onsets of every

two items in the replayed motion sequence. We found that the replayed sequence was

approximately 10 times faster than the evoked activity sequence in the full sequence

condition.

To further examine the period during which the backward replay occurred most

frequently, we divided the blank period (2.4 s) into 4 stages, each lasting 600 ms. In

start-only trials, we found that the backward replay predominantly appeared within the

third stage of the blank period (1.2 – 1.8 s after the onset of the blank period, Wilcoxon

signed-rank test, p = 0.0108), but there was no significant sequenceness in the other

three stages (first stage, p = 0.7112; second stage, p = 0.5566; fourth stage, p = 0.6791;

Figure 4C). In end-only trials, the backward replay was more likely to occur during the

second and third stages of the blank period, although the results approached, but did

not reach, significance (Wilcoxon signed-rank test, second stage, p = 0.0936; third

stage, p = 0.0642; **Figure 4F**), In contrast, the replay was not frequently observed

during the first and last stages (Wilcoxon signed-rank test, first stage, p = 0.9479; fourth

stage, p = 0.4997). Finally, in a control analysis, we conducted a comprehensive

examination of all 24 potential sequences. Only the backward replay was detected in

both start- and end-only conditions (Figure S4).

Figure 4. Backward replay in both start- and end-only conditions

(A and **D)** Examples of backward sequential reactivation in start- (A) and end- (D) only

conditions from a representative participant. Each row represents the reactivation

probabilities for the four motion directions at that time point and each column indicates

one of the four motion directions. **(B and E)** Backward replay of the exposed motion

sequence with peak transition lags at 28 – 40 ms in the start-only condition (B) and at

28 – 36 ms in the end-only condition (E). Horizontal dashed lines represent corrected

significance levels from a nonparametric permutation test at the 2^{nd}-level GLM analysis

of TDLM. The sequenceness on the y-aixs is a unitless measure. The lag on the x-axis

represents the time lag (Δt) between sequential reactivations, rather than absolute time.

The red shaded areas indicate the lags when the evidence of sequenceness exceeded

the permutation-based significance threshold. **(C and F)** Backward replay of the

exposed motion sequence predominantly appeared at 1.2 – 1.8 s in the start-only

condition (C) and 0.6 – 1.8 s in the end-only condition (F) after the onset of the blank

period.

Backward replay is cue-dependent and depends on the amount of exposure

So far, we have demonstrated that the cue-triggered elevated responses induced

by the motion sequence exposure were not motion direction specific. However, motion

information was spontaneously reactivated and the motion sequence was backward

replayed in a time-compressed manner. It remains unknown whether the observed

backward replay of the motion sequence was cue-dependent. In other words, does the

replay of the motion sequence occur irrespective of which item of the sequence is

presented as a cue? To address this question, we conducted Experiment 2, mirroring

the design of Experiment 1 but with a different cue. Instead of flashing the first or last

RDK, we presented the second or third RDK as a cue (2^{nd}-only or 3^{rd}-only condition, see

Figure S5A and Methods) to examine whether these two non-terminal cues could

induce replay during the post-cue blank period. We found no evidence of either forward

or backward replay in either condition (**Figure 5A**; maximal nonsignificant effect at 32

ms-lag: 2^{nd}-only condition, β = −0.0024 ± 0.0014; 3^{rd}-only condition, β = −0.0018 ±

0.0015).

Figure 5. Backward replay is cue-dependent and depends on the amount of

exposure

(A) No evidence for replay was found in either the 2^{nd}-only (left) or 3^{rd}-only (right)

condition in Experiment 2. Horizontal dashed lines have the same meaning as those in

Figure 4B and 4E. **(B)** In Experiment 3, immediately after the functional localizer

phase, participants entered the main phase without the exposure phase. In the end-

only condition, backward replay was observed; however, in the start-only condition, no

such replay was observed.

Another interesting question is how varying the amount of exposure would affect

replay events during the blank period. To explore this issue, we conducted Experiment

3, also similar to Experiment 1 except for the removal of the exposure phase.

Accordingly, only full sequence trials (50% trials) in the main phase served for exposure

(see Figure S5B and Methods). In the start-only condition, we found a numerical trend

of backward replay at transition lags at 20–40 ms, although it did not reach statistical

significance (**Figure 5B**, left panel; maximum effect at 28 ms-lag: β = −0.0139 ±

0.0021). In the end-only condition, the evidence for backward replay just reached the

significance level at transition lags at 20 – 40 ms (**Figure 5B**, right panel; maximal effect

at 24 ms-lag: β = −0.0147 ± 0.0019, p < 1/24 $\approx$ 0.042, using the peak-level significance

threshold from a nonparametric permutation test, FWE-corrected across lags).

Power increase in replay-associated sharp-wave ripple frequencies

Previous studies on rodents and humans showed that replay events are

associated with increased high-frequency ripple power (Buzsáki, 2015; Liu et al., 2019).

To investigate whether such replay-associated power increase could be observed in our

study, we performed a time-frequency analysis using combined data from the start- and

end-only conditions in Experiment 1. We first identified putative replay onsets during the

post-cue blank period in each trial. Replay onsets were determined by choosing time

points with a high (>95[th] percentile) probability for backward replay with a 32 ms-lag

transition (maximal replay effect at the group level for both start- and end-only

conditions, Figure 4B and 4E; see Methods for details). Using the MEG signals

recorded from whole brain sensors, we found a transient ripple power increase at 120–

180 Hz at the onset of replay events, compared to the baseline period of 50–100 ms

prior to replay onset (**Figure 6A**; cluster-based permutation test with cluster forming

threshold t > 3.1 and 5000 permutations).

Figure 6. Replay events align with ripple power, with source activation in the MTL

preceding activation in visual cortex

(A) Top: In Experiment 1, a time-frequency decomposition of sensor-level data revealing

a brief increase in high-frequency oscillatory power at replay onset. Bottom: a cluster-

based permutation test (cluster forming threshold, t > 3.1; number of permutations =

5000) could identify a significant cluster around 140 Hz. **(B)** Source localization of

ripple-band power 30 ms before replay onset showing significant activation in the MTL

(peak Montreal Neurological Institute [MNI] coordinates: X = 21, Y = 21, Z = 13.

Neurological orientation.) **(C)** Source localization of ripple-band power at replay onset

showing significant activation in visual cortex (peak MNI coordinates: X = 20, Y = 8, Z =

17). **(D)** The activation time course of the MTL at its peak MNI coordinate is shown in

red, whereas that of visual cortex at its peak MNI coordinate is displayed in green. The

MTL reached its peak activation before visual cortex.

Visual cortical activation lags MTL activation

Since replay events were obtained based on decoding probabilities of motion

directions from occipital sensors only, we next examined whether the observed replay

onsets were related to power increase within visual cortex. Using a linearly constrained

minimum variance (LCMV) beamforming algorithm (Van Veen et al., 1997), we first

epoched the data using replay onsets combined from both start- and end-only

conditions and then beamformed the broadband power into the source space (see

Methods). We found that the power increase at replay onset was localized to sources in

bilateral visual cortex (**Figure 6C**). Moreover, the right MTL, including the hippocampus

and amygdala, was associated with power increase prior to the onset of replay events

(**Figure 6B**). Specifically, activation (i.e., power increase) in the MTL occurred 30 ms

earlier than that in visual cortex ($p < 0.05$, corrected, whole brain nonparametric

permutation test). For display purposes, we extracted activations from the 10 most

activated voxels within the MTL and visual cortex, respectively, and plotted the time

courses of their broadband power. Peak activation in visual cortex at replay onset was

preceded by peak activation in the MTL (**Figure 6D**), implying an information flow from

the MTL to visual cortex.

DISCUSSION

We found the co-existence of time-locked, non-feature-specific elevated

responses and non-time-locked, feature-specific backward replay after visual sequence

exposure in human brain. The feature-specific backward replay occurred in a time-

compressed manner and could be triggered only by the first or last stimulus of the

sequence. Interestingly, even brief exposure to the sequence could still induce a trend

for the backward replay. Finally, we observed that MTL activity preceded the ripple

power increase in visual cortex at replay onset.

Our study provides two new findings in the fields of vision and learning. First,

different from many previous studies showing that expectation-based responses in

static contexts are feature-specific (Wyart et al., 2012; Kok et al., 2014, 2017), we show

here that prediction-related elevated responses in the dynamic temporal context are not.

This non-feature-specific elevated responses potentially facilitate the general

processing of any upcoming stimuli, rather than stimuli with a specific feature. Second,

contrary to the prevailing notion that replay in learning and memory requires lengthy

training with a task, our study shows that even brief exposure to a visual sequence is

sufficient to induce replay. The replay we unveiled here may play a critical role in

memorizing the visual sequence.

Regarding the two-fold neural consequences following the visual sequence

exposure, the non-feature-specific elevated responses could be induced by rhythmic

entrainment. Substantial evidence supports that rhythmic stimulation can entrain neural

oscillations, which in turn facilitates predictions about future inputs and enhances the

brain's readiness for incoming stimuli (Lakatos et al., 2008, 2013; Herrmann et al., 2016;

Barne et al., 2022). For instance, recent findings demonstrate that even task-irrelevant

information could be more effectively decoded when presented at moments close to a

highly probable target presentation (Auksztulewicz et al., 2019). In our study, the

rhythmic presentation of the motion sequence may have entrained oscillatory activity in

the brain, leading to periodic activation of sensory cortices. This rhythmic entrainment

likely serves as a possible mechanism supporting the general facilitation of neural

processing for any upcoming stimuli, independent of specific stimulus features.

In contrast, feature-specific replay may operate through a different mechanism

driven by intrinsic neural processes rather than direct external stimuli. Therefore, this

intrinsic activity is not time-locked to stimulus onset, but instead manifests in a

spontaneous way. Furthermore, since replay often occurs during offline periods (e.g.,

rest or blank period), it allows the neural system to reactivate and consolidate past

experiences without interference from ongoing external inputs. As a result, feature-

specific replay may facilitate the integration of fragmented experiences into coherent

representations, thereby subserving visual sequence learning and memory.

Another interesting question regarding the two-fold neural responses is whether

the elevated responses and the backward replay share the same neural origin, for

instance, originating from the hippocampus or other brain areas. For the elevated

responses, the cue serves to provide temporal information for upcoming stimuli but

without detailed feature information, thereby priming the cortices for activation and

facilitating the general processing of future events. Previous animal studies (Xu et al.,

2012; Gavornik and Bear, 2014) showed that this process can be implemented through

a simple local synaptic mechanism in visual cortex (Xu et al., 2012) without top-down

guidance. Moreover, Ekman et al. (2023) recently found no functional relationship

between activities in V1 and the hippocampus when exposing human participants with a

white dot sequence, further suggesting that the elevated responses may indeed

originate in visual cortex.

Different from the elevated responses, the replay events are likely initiated by the

hippocampus. This speculation is supported by two key findings. First, the replay of the

motion direction information manifested in a backward direction and a time-compressed

manner. Second, the replay is observed regardless of whether the cue is the first or last

RDK in the sequence. Both of these two properties cannot be explained by a simple

local synaptic mechanism (Xu et al., 2012) or a pattern completion-like mechanism

(Hindy et al., 2016; Ekman et al., 2017; Kok and Turk-Browne, 2018) within visual cortex

alone. Instead, the replay entails reorganizing the motion direction sequence in the

brain. Therefore, we propose the involvement of active communication between the

hippocampus and visual cortex in the occurrence of replay events, consistent with the

view that the hippocampus encodes relationships among stimuli whereas visual cortex

primarily acts as a platform for the manifestation of replay events (Whittington et al.,

2020). Previous studies on memory consolidation considered the exchange of

information between these two regions critical for facilitating replay events through

hippocampal-neocortical circuits (Buzsáki, 1996; Ji and Wilson, 2007; Carr et al., 2011;

Ólafsdóttir et al., 2016; Buch et al., 2021). Functionally, replay events offer a mechanism

for transferring recent experience from the hippocampus to the cortex, enabling the

encoding of stimulus relationships in the cortex (Marr and Brindley, 1971; Alvarez and

Squire, 1994; Redish and Touretzky, 1998; Dimakopoulos et al., 2022).

A natural behavioral consequence of visual sequence exposure is visual

sequence learning (Baker et al., 2014; Finnie et al., 2021; Ekman et al., 2023). How is

the learning implemented through hippocampus-dependent replay in visual cortex?

Three key processes are considered here. First, the hippocampus is involved in

encoding relationships among stimuli (Staresina and Davachi, 2009; Turk-Browne et al.,

2009; Hsieh et al., 2014; Garvert et al., 2017); in fact, the ability to encode such

relationships drastically decreases when the hippocampus is damaged (Chun and

Phelps, 1999; Hannula et al., 2006; Konkel et al., 2008; Schapiro et al., 2014; Finnie et

al., 2021). Second, visual cortical areas act as a "cognitive blackboard" where task-

relevant features are highlighted through feedback connections (Roelfsema and de

Lange, 2016). Such a blackboard can be flexibly written or edited. Third, there are

strong bidirectional connections between the hippocampus and sensory cortices

(Eichenbaum et al., 2007; Henke, 2010). As proposed by the hippocampal-cortical

backward projection model (Rolls, 2000), sequential reactivations of feature information

initially generated in the hippocampus can be quickly and accurately sent back to the

sensory cortices, consistent with the findings of Ji and Wilson, (2007). A recent study

also provided direct evidence that visual sequence plasticity is impaired when the

hippocampus is damaged (Finnie et al., 2021), supporting the hypothesis of functional

feedback information flow.

Why does the replay manifest in a reverse order? To date, there is no consensus

on this issue. In rodents, a seminal study of replay during the awake state showed that

when animals stopped at the end of a rewarded pathway, place cells were reactivated in

the reverse order of the previously experienced direction (Foster and Wilson, 2006).

However, a later study revealed that awake replay could occur in either a forward or

backward direction relative to behavioral experience (Diba and Buzsáki, 2007).

Similarly, in humans, both directions have been observed in different nonspatial

cognitive tasks (Liu et al., 2019, 2021b; Wimmer et al., 2020). Forward replay may be

associated with planning (Ólafsdóttir et al., 2015; Gillespie et al., 2021), providing

information pertaining to the assessment of future pathways (Diba and Buzsáki, 2007).

Backward replay may be more related to experience, as often observed at the end of

runs when animals consume a reward (Foster and Wilson, 2006; Ambrose et al., 2016).

Nevertheless, the exact function of the replay direction remains mysterious, as both

forward and backward replays are modulated by task demands (Ólafsdóttir et al., 2017).

Thus, the underlying neural mechanisms of backward replay in visual cortex remain to

be investigated.

Finally, we also found that replay occurrence is modulated by the cue. This result

highlights the importance of the start and end points of the sequence in the replay. One

fascinating proposal is that the replay event is sensitive to sequence boundaries, as

indicated by the role of the start or end point of the sequence as salient boundaries that

anchor place cell firing (Rivard et al., 2004). Accordingly, previous studies have shown

that when rats are trained to run along a linear track starting at different points, place

fields tend to be anchored to either the start or end of the journey (Gothard et al., 1996;

Redish et al., 2000), suggesting that a boundary is essential to sequence integrity and

may play a pivotal role in triggering replay onset. An alternative explanation to these

findings posits that the onset of the second or third stimulus in the sequence reinstate

the neural representations of a partial visual sequence (Ekman et al., 2023). For

example, for a given sequence, A → B → C → D, flashing the third stimulus (i.e., C)

only trigger a backward reactivation of the sequence giving C → B → A during the blank

period.

Taken together, we found that simple visual sequence exposure could

concurrently induce two-fold brain plasticity, that is, non-feature-specific elevated

responses and feature-specific backward replay in the human visual cortex. We

speculate that the non-feature-specific elevated responses may enhance general

processing of upcoming visual stimuli, whereas the feature-specific backward replay

may subserve visual sequence learning and memory. These findings significantly

advance our understanding of the task-independence and the multifaceted nature of

brain plasticity in response to visual experience.

METHODS

Participants

A total of 59 healthy participants (29 females) were involved in the three

experiments (Experiment 1: n = 21, 11 females, 22.1 ± 2.61 years; Experiment 2: n =

18, 10 females, 23.56 ± 3.29 years; Experiment 3: n = 20, 10 females, 20.65 ± 2.62

years). In Experiment 1, data from three participants were excluded before analyses, as

two showed large head motion (>20 mm) and the behavioral performance of one was at

chance level. In Experiment 3, data from two participants were excluded before

analyses because of large head motion (>20 mm). All participants were recruited in

exchange for monetary compensation (100 RMB/h). They reported normal or corrected-

to-normal vision and had no history of neurological disorders. They were naive to the

purposes of the study. The experiments reported here were carried out in accordance

with the guidelines expressed in the Declaration of Helsinki. All participants provided

written informed consent in accordance with the procedures and protocols approved by

the Human Subject Review Committee of Peking University.

Task

Experiment 1

Visual stimuli were RDKs with 100% coherence. All dots in an RDK moved in the

same direction (luminance: 2.86 cd/ m^2; diameter: 0.1°; speed: 8°/s) and were

presented against a gray background (luminance: 16.7 cd/m^2). At any one moment, 400

dots were visible within a 9° circular aperture and moved in one of the four directions: 0°

(right), 90° (up), 180° (left), and 270° (down). Each participant completed three phases

successively in the MEG scanner, namely, the functional localizer phase, the exposure

phase, and the main phase (Figure 1C). In the functional localizer phase, each trial

started with the presentation of an RDK for 1 s followed by a 1–1.5 s intertrial interval

(ITI). Participants did not need to perform any task in this phase. The motion direction of

the RDK was randomly chosen from the four directions. Each participant performed two

functional localizer runs and, each run comprised 100 trials, resulting in a total of 50

trials per motion direction. The localizer data were used to train motion direction

classifiers. This phase took approximately 10 min.

In the exposure phase, we showed participants the four RDKs in either clockwise

or counterclockwise order (e.g., 0° → 90° → 180° → 270°, Figure 1A and 1C); each was

displayed for 400 ms, followed by 300 ms of a blank screen with a fixation only.

Therefore, the full sequence lasted for 2.8 s. The order was counterbalanced among

participants, but once decided, it was fixed for each participant. Participants were

instructed to detect an oddball RDK by pressing a button, i.e., in 20% trials, dots in one

of the four RDKs traveled at a faster speed (9°/s). Each participant completed four runs

of 50 trials.

In the main phase, participants were presented with trials in three different

conditions: full sequence condition (50% trials), start-only condition (25% trials), and

end-only condition (25% trials). Trials in the full sequence condition were identical to

those in the exposure phase which served as "topping-up" exposure to maintain the

exposure effect, similar to "topping-up" adaptation in visual adaptation studies (Fang et

al., 2005). In the start- and end-only conditions; however, we only presented the first

RDK (start-only condition) or the last RDK (end-only condition) of the full sequence. For

a given run, the order of the three conditions was pseudorandomized with the restriction

that the start- and end-only trials were always preceded or followed by a full sequence

trial. Participants performed an identical oddball detection task as during the exposure

phase. The oddball occurred only in 10% of full sequence trials. Finally, each participant

completed four runs of 48 trials, yielding a total of 96 full sequence trials, 48 start-only

trials, and 48 end-only trials.

Experiments 2 and 3

In Experiment 2, we only presented the second or third RDK as a cue at the start

of the trial, referred to as the 2^{nd}-only and 3^{rd}-only conditions, respectively. The

procedure of Experiment 2 was similar to that of Experiment 1 except that the start- and

end-only conditions were replaced with the 2^{nd}-only and 3^{rd}-only conditions. Experiment

3 followed the same procedure as Experiment 1 except that the exposure phase was

removed.

QUANTIFICATION AND STATISTICAL ANALYSIS

MEG acquisition and preprocessing

Neuromagnetic signals were recorded continuously at 1000 Hz with a 306-

channel (204 planar gradiometers; 102 magnetometers), whole-head MEG system

(Elekta Neuromag TRIUX) in a magnetically shielded room. Before scanning, four-

headed position indicator coils attached to the scalp determined the head position

with respect to the sensor array. Coil location was digitized with respect to three

anatomical landmarks (nasion and preauricular points) with a 3D digitizer

(Polhemus Isotrak system). Participants sat upright inside the scanner, while the

stimuli were projected onto a screen suspended in front of them. Participants

responded using a MEG-compatible button box held in their right hand.

To reduce noise from external environment, the temporal extension of signal-

space separation (tSSS) method was applied at the preprocessing stage using the

Elekta Neuromag MaxFilter software (Taulu and Simola, 2006). MEG signals were

high-pass filtered at 0.5 Hz using a first-order IIR filter to remove slow-drifts. Data

were then downsampled from 1000 Hz to 250 Hz for sequenceness analysis or 500

Hz for time-frequency analysis. Excessively noisy segments and sensors were

automatically removed before independent component analysis (FastICA,

http://research.ics.aalto.fi/ica/fastica) and performed to remove artifacts including

cardiac signals, eye movements and blinks, and environmental noise. Artifact

components were removed by visual inspection of spatial topography, time course,

kurtosis of the time course, and frequency spectrum of all components. Only 72

sensors (including both gradiometers and magnetometers) covering the occipital

lobe, labeled as "Occipital" in the MEG data acquisition system were used for MEG

data analyses, except for source localization. The sensor selection was primarily

motivated by the main objective of the study, examining replay events in visual

cortex.

Event-related fields

To calculate the event-related fields (ERFs), MEG epochs were segmented

around trial onset for each trial and baseline-corrected using the mean activity in

the time window of [−0.3 s, 0] before trial onset. Only 48 planar gradiometers of

"Occipital" sensors were used to calculate the ERFs. The planar-combined ERF

activity was then averaged for each condition. To minimize potential baseline

confounds caused by the short interval between every two RDKs, we further

calculated the ERF peak amplitude using the mean activity in the time window of

[−0.3 s, 0] before the onset of the corresponding RDK as the baseline.

Subsequently, peak amplitude during each of the four RDK intervals was calculated

as the difference between the peak and its corresponding baseline.

Multivariate pattern analyses

Multivariate pattern analyses (MVPA) were performed to classify the neural

activity patterns elicited by the motion directions of the four RDKs in the main

phase. A one-versus-rest Lasso-regularized logistic regression model was trained

using the MEG signals from the 72 occipital sensors in the functional localizer

phase. Specifically, we trained a five-class classifier, including four classes from

trials in which the four RDKs were presented, and an additional class comprising an

equivalent amount of null data extracted from the 1–1.5 s ITI. Null data were

included to reduce the spatial correlation among the classes, thereby concurrently

lowering the decoding probabilities for all classes (Liu et al., 2021a; Nour et al.,

2021). Class weights were balanced by adjusting inversely proportional to class

frequencies (i.e., trial numbers) in the training procedure. To reduce noise, MEG

signals were averaged across every five trials within the same class before

decoding.

The trained classifier was then applied to MEG signals at each time point in

the main phase, and decoding probabilities were computed as the outputs of the

classifier. Different from the conventional decoding accuracy, where the classifier

predicts a single label at each time point, decoding probabilities provide a likelihood

estimate for each class. In our study, the classifier outputs a 5-column matrix for all

trials, where each row represents a single trial and each column represents the

probability for one class (i.e., one of the four RDKs and blank ITI). The probabilities

in each row sum to 1, reflecting the relative likelihoods across all classes for that

trial. The highest probability determines the decoded label when computing

decoding accuracy. Finally, averaging these probabilities across trials yields five

values that indicate the overall likelihood of the predicted stimulus belonging to a

given class.

Optimal time point of motion direction representation

The optimal time point of motion direction representation was considered as the

time point with the highest decoding accuracy in the functional localizer data. To index

the optimal time point of each motion direction for each participant, we conducted a

time-resolved motion direction decoding analysis on the functional localizer data using

Lasso-regularized logistic regression models. A leave-one-trial-out cross-validation

procedure was used to train the classifier to determine one of the four motion directions,

yielding a decoding accuracy at each time point for each participant. Finally, the time

point with the highest decoding accuracy was independently extracted for each

participant and each motion direction. These time points are referred to as optimal time

points and were used for training feature-specific decoding models.

Sequenceness measure

To identify sequenceness during the post-cue blank period in each trial, we

trained models to detect transient spontaneous neural reactivation of each motion

direction representation. Therefore, a one-versus-rest Lasso-regularized logistic

regression model was trained separately for each participant and motion direction using

the functional localizer data. MEG signals from the 72 occipital sensors obtained

throughout all localizer scanning sessions were used to train the decoding models. As

our aim was to quantify the evidence of feature-specific sequence, for each motion

direction, we trained a binomial classifier, using positive instances from trials in which

that feature (e.g., 0°) was presented and negative instances from trials in which all other

features (e.g., 90°, 180°, and 270°) were presented, together with an equivalent amount

of null data from the 1–1.5 s ITI. The sensor distributions of beta estimates and the

spatial correlation among the classifiers are shown in Figure S1.

The analysis pipeline of sequenceness is illustrated in **Figure 3**. We first applied

the trained models to MEG signals at each time point during the blank period, to

generate a [time x motion direction] reactivation probability (i.e., decoding probability)

matrix for each trial (Figure S2). The TDLM framework was then used to quantify

evidence for sequential reactivations consistent with the exposed motion sequence (Liu

et al., 2021b, 2021a; Nour et al., 2021).

TDLM is a multiple linear regression approach to quantify the extent to which a

lagged reactivation time course of one motion direction i, (denoted as $X(\Delta t)_i$, where Δt

indicates lag time) can predict the reactivation time course of another motion direction j,

(denoted as X_j). Two steps were included in this pipeline. First, we performed multiple

separate regressions using the reactivation time course of each motion direction j

(where j ∈ [1: 4]) as the dependent variable. The predictors were the time-lagged

reactivation time courses of all four motion directions i (where i ∈ [1: 4]). The regression

model can be expressed as:

$$X_j = \sum_{i=1}^{4} X(\Delta t)_i * \beta(\Delta t)_{i,j} + C \tag{1}$$

The predictor $X(\Delta t)_i$ was a Δt-lagged copy of the reactivation time course of X_i.

The regression coefficients $\beta(\Delta t)_{i,j}$ quantified the strength of the empirical reactivation

pattern from motion direction i to motion direction j at a given time lag, Δt. For example,

if X_j represents the reactivation time course of 0° during the blank period in the main

phase, and X_i represents the reactivation time course of 90° during the same period,

then $\beta(\Delta t)_{i,j}$ is the coefficient that captures the unique variance in the reactivation time

course of 0° explained by the Δt-lagged reactivation time course of 90° (Figure 3D, 1st

level GLM analysis). Finally, all such first-level coefficients were placed in a lag-specific

[4 x 4] empirical transition matrix B, representing evidence for all possible transitions at

a specific lag.

In the second step, we quantified the extent to which this empirical transition

matrix B could be predicted by a model transition matrix reflecting the sequence of

interest, e.g., the exposed motion sequence (Figure 3D, 2nd level GLM analysis; 1 for

transitions of interest and 0 otherwise). We separately modeled forward transitions (T_F),

which followed the order of the motion sequence, and backward transitions (T_B), which

followed the reverse order. The strength of these sequences was measured by:

$$B = \sum_{r=1}^{4} Z_r * T_r \tag{2}$$

where B is the [4 x 4] empirical (lag-specific) transition matrix obtained from the

data, T_r is a [4 x 4] model transition matrix (for regressor r), and Z_r is the scalar

regression coefficient quantifying the extent to which the model transition matrix T_r

predict the empirical transitions, B. We included four model transition matrices as

regressors: (1) T_F: transitions as in the motion sequence in the forward direction

(transitions corresponding to [0° → 90° → 180° → 270°]), (2) T_B: transitions opposite to

the motion sequence in the backward direction ([270° → 180° → 90° → 0°], T_B is the

transpose of T_F), (3) T_{auto}: self-transitions to control for autocorrelation ([4 x 4] identity

matrix), and (4) T_{const}: a constant matrix to model away the average of all transitions,

ensuring that any weight on T_F and T_B was not due to general background neural

dynamics.

Notably, the estimate of sequence strength, Z_r, is a relative measure. For

instance, a Z_r value of zero for the transition from 0° to 90° does not indicate an

absence of replay for that transition; rather, it reflects that the strength of replay of 0° →

90° is not stronger than that of other transitions. Repeating the regression in Equation 2

at each time lag (Δt = 4, 8, 12, ..., 600 ms) results in both forward (i.e., Z_1) and

backward (i.e., Z_2) sequence strength as a function of time lag. Shorter lags indicate

greater time compression, corresponding to faster speeds.

In the current study, sequenceness was defined as the contrast between the

evidence for replay of the motion sequence in the forward direction ([0° → 90° → 180°

→ 270°]) versus the backward direction ([270° → 180° → 90° → 0°]). Specifically,

sequenceness was calculated as the difference between the regression coefficients for

forward and backward transitions (i.e., $Z_1 - Z_2$). This contrast control for between-

participant variance in the sequential replay per se, which may arise from factors such

as task engagement or measurement sensitivity (Liu et al., 2021a; Nour et al., 2021). As

sequenceness is derived from regression coefficients, it is inherently a unitless

measure. Positive sequenceness values indicate replay in a predominantly forward

direction, whereas negative sequenceness values indicate replay in a predominantly

backward direction.

For statistical inference, we used nonparametric permutation tests involving all

possible permutations of the stimulus labels at the second-level regression, equivalent

to permuting the rows and columns together of the transition matrices used to calculate

sequenceness. For each permutation, we calculated the peak absolute mean sequence

strength over participants and across lags (controlling for multiple comparisons across

lags). Sequence strength in unpermuted data was considered significant if its absolute

magnitude was >95% of the within-permutation peak.

Identifying replay onsets and analyzing time-frequency

Having identified that the replay transition at the group level was a 32 ms-lag, we

next identified replay onsets. Replay onset was defined as the time point when a strong

reactivation of one motion direction (e.g., 0°) was followed by a strong reactivation of

the next motion direction (e.g., 90°) in the sequence, with a 32 ms-lag (Liu et al., 2019;

Nour et al., 2021). We first generated a matrix Orig as

$$Orig = X * T \qquad (3)$$

where X is the [time x motion direction] reactivation matrix, and T is the backward

transition matrix. The transition matrix T defines the mapping between the motion

direction corresponding to column i in X, and column i in Orig. For example, column 1 in

X is the reactivation time course of the motion direction 0°, while column 1 in Orig is the

reactivation time course of the motion direction 90°. We then shifted each column of X

by Δt = 32 ms, to generate another matrix Proj,

$$Proj = X(\Delta t) \qquad (4)$$

769   where row i in Proj corresponds to row i + 32 ms in X. Next, we multiplied Proj

770   and Orig elementwise, summing over the columns of the resulting matrix; therefore,

771   creating a [time x 1] vector, R, in which each element (i.e., row) indicates the strength of

772   replay with a 32 ms-lag at a given time.

$$R_t = \sum_{i=1}^{4} Orig_{ti} * Proj_{ti} \qquad (5)$$

Finally, we identified putative replay event onsets by thresholding R at its 95th

percentile preceded by a 100 ms pre-onset baseline exhibiting a low probability of

replay at each time point.

We then epoched MEG data in the blank period surrounding each onset and

computed a frequency decomposition (wavelet transformation) in the time window of

−100 – 150 ms using all sensors. We aimed to find evidence for the power increase in

the high-frequency (120 – 160 Hz) region of interest at replay onset, compared with a

pre-onset baseline (−100 – −50 ms from onset), similar to a previous study (Liu et al.,

2019). Finally, we generated two separate [time x frequency] matrices (i.e., using

forward and backward transition matrices separately) by averaging estimates over

sensors and events, capturing the typical spectrum-resolved power change at replay

onset.

MEG source reconstruction

We identified the neural sources associated with increased ripple power at

putative replay onsets. Forward models were generated based on a single shell using

the superposition of basic functions that approximately correspond to the plane

tangential to the MEG sensor array. Linearly constrained minimum variance

beamforming (Van Veen et al., 1997) was used to reconstruct the epoched MEG data to

a grid in MNI space (grid step, 5 mm). The sensor covariance matrix for beamforming

was estimated using broadband power data across all frequencies. The baseline activity

was the mean activity averaged over −100–−50 ms relative to replay onset. All

nonartifactual replay epochs were baseline-corrected at source level. We obtained

whole-brain results for voxels predicted by participant-specific ripple power at replay

onset. Nonparametric permutation tests were performed on the volume of interest to

compute the multiple comparison p values of clusters >10 voxels (whole-brain

corrected, cluster-defining threshold; t = 3.1, n = 5000 permutations).

Data and Code availability

800 Data used to generate the findings of this study will be available upon request (subject

801 to participant consent) to the Lead Contact. TDLM MATLAB Code available at:

802 https://github.com/YunzheLiu/TDLM. Custom computer code will be made available

803 upon request to the Lead Contact.

AUTHOR CONTRIBUTIONS

805 Conceptualization, T.H., Y.L., and F.F.; Investigation, T.H., Q.W., X.G., and X.Z.; Writing

806 – Original Draft, T.H.; Writing – Review & Editing, T.H., Q.W., X.Z., Y.L., and F.F.

ACKNOWLEDGEMENTS

808 This study was supported by National Science and Technology Innovation 2030 Major

809 Program (2022ZD0204802), the National Natural Science Foundation of China

810 (T2421004, 31930053, 32400874), Beijing Natural Science Foundation (5244044), and

811 the Young Scientists Fund of the Humanities and Social Science Foundation of Ministry

812 of Education of China (23YJCZH071).

DECLARATION OF INTERESTS

814 The authors declare no competing interests.

REFERENCES

Alvarez P, Squire LR (1994) Memory consolidation and the medial temporal lobe: a simple network model. Proc Natl Acad Sci U S A 91:7041–7045.

Ambrose RE, Pfeiffer BE, Foster DJ (2016) Reverse Replay of Hippocampal Place Cells Is Uniquely Modulated by Changing Reward. Neuron 91:1124–1136.

Auksztulewicz R, Myers NE, Schnupp JW, Nobre AC (2019) Rhythmic Temporal Expectation Boosts Neural Activity by Increasing Neural Gain. J Neurosci 39:9806–9817.

Baker R, Dexter M, Hardwicke TE, Goldstone A, Kourtzi Z (2014) Learning to predict: Exposure to temporal sequences facilitates prediction of future events. Vision Res 99:124–133.

Barne LC, Cravo AM, de Lange FP, Spaak E (2022) Temporal prediction elicits rhythmic preactivation of relevant sensory cortices. European Journal of Neuroscience 55:3324–3339.

Buch ER, Claudino L, Quentin R, Bönstrup M, Cohen LG (2021) Consolidation of human skill linked to waking hippocampo-neocortical replay. Cell Rep 35:109193.

Buonomano DV, Merzenich MM (1998) CORTICAL PLASTICITY: From Synapses to Maps. Annual Review of Neuroscience 21:149–186.

Bush D, Ólafsdóttir HF, Barry C, Burgess N (2022) Ripple band phase precession of place cell firing during replay. Current Biology 32:64-73.e5.

Buzsáki G (1986) Hippocampal sharp waves: Their origin and significance. Brain Research 398:242–252.

Buzsáki G (1996) The hippocampo-neocortical dialogue. Cereb Cortex 6:81–92.

Buzsáki G (2015) Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. Hippocampus 25:1073–1188.

Carr MF, Jadhav SP, Frank LM (2011) Hippocampal replay in the awake state: a potential substrate for memory consolidation and retrieval. Nat Neurosci 14:147–153.

Chen N, Cai P, Zhou T, Thompson B, Fang F (2016) Perceptual learning modifies the
functional specializations of visual cortical areas. Proceedings of the National
Academy of Sciences 113:5724–5729.

Chun MM, Phelps EA (1999) Memory deficits for implicit contextual information in
amnesic subjects with hippocampal damage. Nat Neurosci 2:844–847.

Costandi M (2016) Neuroplasticity. MIT Press.

Demarchi G, Sanchez G, Weisz N (2019) Automatic and feature-specific prediction-
related neural activity in the human auditory system. Nat Commun 10:3440.

Diba K, Buzsáki G (2007) Forward and reverse hippocampal place-cell sequences
during ripples. Nat Neurosci 10:1241–1242.

Dimakopoulos V, Mégevand P, Stieglitz LH, Imbach L, Sarnthein J (2022) Information
flows from hippocampus to auditory cortex during replay of verbal working
memory items Griffiths TD, Colgin LL, eds. eLife 11:e78677.

Eagleman SL, Dragoi V (2012) Image sequence reactivation in awake V4 networks.
PNAS 109:19450–19455.

Eichenbaum H, Yonelinas AR, Ranganath C (2007) The Medial Temporal Lobe and
Recognition Memory. Annu Rev Neurosci 30:123–152.

Ekman M, Kok P, Lange FP de (2017) Time-compressed preplay of anticipated events
in human primary visual cortex. Nat Commun 8:15276.

Ekman M, Kusch S, de Lange FP (2023) Successor-like representation guides the
prediction of future events in human visual cortex and hippocampus Barense M,
Baker CI, Barron H, eds. eLife 12:e78904.

Fang F, Murray SO, Kersten D, He S (2005) Orientation-Tuned fMRI Adaptation in
Human Visual Cortex. Journal of Neurophysiology 94:4188–4195.

Finnie PSB, Komorowski RW, Bear MF (2021) The spatiotemporal organization of
experience dictates hippocampal involvement in primary visual cortical plasticity.
Current Biology 31:3996-4008.e6.

Foster DJ, Wilson MA (2006) Reverse replay of behavioural sequences in hippocampal
place cells during the awake state. Nature 440:680.

Garvert MM, Dolan RJ, Behrens TE (2017) A map of abstract relational knowledge in
the human hippocampal–entorhinal cortex Davachi L, ed. eLife 6:e17086.

Gavornik JP, Bear MF (2014) Learned spatiotemporal sequence recognition and
prediction in primary visual cortex. Nat Neurosci 17:732–737.

Gillespie AK, Maya DAA, Denovellis EL, Liu DF, Kastner DB, Coulter ME, Roumis DK,
Eden UT, Frank LM (2021) Hippocampal replay reflects specific past experiences
rather than a plan for subsequent choice. Neuron 109:3149-3163.e6.

Gothard KM, Skaggs WE, Moore KM, McNaughton BL (1996) Binding of hippocampal
CA1 neural activity to multiple reference frames in a landmark-based navigation
task. J Neurosci 16:823–835.

Gridchyn I, Schoenenberger P, O'Neill J, Csicsvari J (2020) Assembly-Specific
Disruption of Hippocampal Replay Leads to Selective Memory Deficit. Neuron
106:291-300.e6.

Gutnisky DA, Hansen BJ, Iliescu BF, Dragoi V (2009) Attention Alters Visual Plasticity
during Exposure-Based Learning. Current Biology 19:555–560.

Hannula DE, Tranel D, Cohen NJ (2006) The Long and the Short of It: Relational
Memory Impairments in Amnesia, Even at Short Lags. J Neurosci 26:8352–8359.

Henke K (2010) A model for memory systems based on processing modes rather than
consciousness. Nat Rev Neurosci 11:523–532.

Herrmann B, Henry MJ, Haegens S, Obleser J (2016) Temporal expectations and
neural amplitude fluctuations in auditory cortex interactively influence perception.
NeuroImage 124:487–497.

Hindy NC, Ng FY, Turk-Browne NB (2016) Linking pattern completion in the
hippocampus to predictive coding in visual cortex. Nat Neurosci 19:665–667.

Hsieh L-T, Gruber MJ, Jenkins LJ, Ranganath C (2014) Hippocampal Activity Patterns
Carry Information about Objects in Temporal Context. Neuron 81:1165–1178.

Igata H, Ikegaya Y, Sasaki T (2021) Prioritized experience replays on a hippocampal
predictive map for learning. Proceedings of the National Academy of Sciences
118:e2011266118.

Ji D, Wilson MA (2007) Coordinated memory replay in the visual cortex and
hippocampus during sleep. Nature Neuroscience 10:100–107.

Joo HR, Frank LM (2018) The hippocampal sharp wave–ripple in memory retrieval for
immediate use and consolidation. Nat Rev Neurosci 19:744–757.

Kok P, Failing M, de Lange F (2014) Prior Expectations Evoke Stimulus Templates in
the Primary Visual Cortex. Journal of cognitive neuroscience 26.

Kok P, Jehee JFM, de Lange FP (2012) Less Is More: Expectation Sharpens
Representations in the Primary Visual Cortex. Neuron 75:265–270.

Kok P, Mostert P, Lange FP de (2017) Prior expectations induce prestimulus sensory
templates. PNAS 114:10473–10478.

Kok P, Turk-Browne NB (2018) Associative Prediction of Visual Shape in the
Hippocampus. J Neurosci 38:6888–6899.

Konkel A, Warren D, Duff M, Tranel D, Cohen N (2008) Hippocampal amnesia impairs
all manner of relational memory. Frontiers in Human Neuroscience 2 Available at:
https://www.frontiersin.org/article/10.3389/neuro.09.015.2008 [Accessed May 26,
2022].

Kurth-Nelson Z, Economides M, Dolan RJ, Dayan P (2016) Fast Sequences of Non-
spatial State Representations in Humans. Neuron 91:194–204.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE (2008) Entrainment of
Neuronal Oscillations as a Mechanism of Attentional Selection. Science 320:110–
113.

Lakatos P, Musacchia G, O'Connel MN, Falchier AY, Javitt DC, Schroeder CE (2013)
The Spectrotemporal Filter Mechanism of Auditory Selective Attention. Neuron
77:750–761.

Lee AK, Wilson MA (2002) Memory of Sequential Experience in the Hippocampus
during Slow Wave Sleep. Neuron 36:1183–1194.

Li W (2016) Perceptual Learning: Use-Dependent Cortical Plasticity. Annual Review of
Vision Science 2:109–130.

Liu Y, Dolan RJ, Higgins C, Penagos H, Woolrich MW, Ólafsdóttir HF, Barry C, Kurth-
Nelson Z, Behrens TE (2021a) Temporally delayed linear modelling (TDLM)
measures replay in both animals and humans Colgin LL, Kemere C, van der
Meer M, eds. eLife 10:e66917.

Liu Y, Dolan RJ, Kurth-Nelson Z, Behrens TEJ (2019) Human Replay Spontaneously
Reorganizes Experience. Cell 178:640-652.e14.

Liu Y, Mattar MG, Behrens TEJ, Daw ND, Dolan RJ (2021b) Experience replay is
associated with efficient nonlocal learning. Science 372 Available at:
https://science.sciencemag.org/content/372/6544/eabf1357 [Accessed May 21,
2021].

Liu Y, Nour MM, Schuck NW, Behrens TEJ, Dolan RJ (2022) Decoding cognition from
spontaneous neural activity. Nat Rev Neurosci:1–11.

Lu J, Luo L, Wang Q, Fang F, Chen N (2020) Cue-triggered activity replay in human
early visual cortex. Sci China Life Sci Available at: /doi/10.1007/s11427-020-
1726-5 [Accessed June 29, 2020].

Lu Z-L, Dosher BA (2022) Current directions in visual perceptual learning. Nat Rev
Psychol 1:654–668.

Marr D, Brindley GS (1971) Simple memory: a theory for archicortex. Philosophical
Transactions of the Royal Society of London B, Biological Sciences 262:23–81.

McFadyen J, Liu Y, Dolan RJ (2023) Differential replay of reward and punishment paths
predicts approach and avoidance. Nat Neurosci 26:627–637.

Mo C, Lu J, Wu B, Jia J, Luo H, Fang F (2019) Competing rhythmic neural
representations of orientations during concurrent attention to multiple orientation
features. Nat Commun 10:1–10.

Nour MM, Liu Y, Arumuham A, Kurth-Nelson Z, Dolan RJ (2021) Impaired neural replay
of inferred relationships in schizophrenia. Cell 184:4315-4328.e17.

O'Keefe J, Nadel L (1978) The hippocampus as a cognitive map. Oxford : New York:
Clarendon Press ; Oxford University Press.

Ólafsdóttir HF, Barry C, Saleem AB, Hassabis D, Spiers HJ (2015) Hippocampal place
cells construct reward related sequences through unexplored space Eichenbaum
H, ed. eLife 4:e06063.

Ólafsdóttir HF, Carpenter F, Barry C (2016) Coordinated grid and place cell replay
during rest. Nat Neurosci 19:792–794.

Ólafsdóttir HF, Carpenter F, Barry C (2017) Task Demands Predict a Dynamic Switch in
the Content of Awake Hippocampal Replay. Neuron 96:925-935.e6.

Pojoga SA, Kharas N, Dragoi V (2020) Perceptually unidentifiable stimuli influence
cortical processing and behavioral performance. Nat Commun 11:6109.

Redish AD, Rosenzweig ES, Bohanick JD, McNaughton BL, Barnes CA (2000)
Dynamics of Hippocampal Ensemble Activity Realignment: Time versus Space. J
Neurosci 20:9298–9309.

Redish AD, Touretzky DS (1998) The role of the hippocampus in solving the Morris
water maze. Neural Comput 10:73–111.

Rivard B, Li Y, Lenck-Santini P-P, Poucet B, Muller RU (2004) Representation of objects
in space by two classes of hippocampal pyramidal cells. J Gen Physiol 124:9–25.

Roelfsema PR, de Lange FP (2016) Early Visual Cortex as a Multiscale Cognitive
Blackboard. Annual Review of Vision Science 2:131–151.

Rolls ET (2000) Hippocampo-cortical and cortico-cortical backprojections. Hippocampus
10:380–388.

Sasaki Y, Nanez JE, Watanabe T (2010) Advances in visual perceptual learning and
plasticity. Nat Rev Neurosci 11:53–60.

Schapiro AC, Gregory E, Landau B, McCloskey M, Turk-Browne NB (2014) The
necessity of the medial temporal lobe for statistical learning. J Cogn Neurosci
26:1736–1747.

Staresina BP, Davachi L (2009) Mind the Gap: Binding Experiences across Space and
Time in the Human Hippocampus. Neuron 63:267–276.

Taulu S, Simola J (2006) Spatiotemporal signal space separation method for rejecting
nearby interference in MEG measurements. Phys Med Biol 51:1759–1768.

Turk-Browne NB, Scholl BJ, Chun MM, Johnson MK (2009) Neural evidence of
statistical learning: efficient detection of visual regularities without awareness. J
Cogn Neurosci 21:1934–1945.

Turner W, Blom T, Hogendoorn H (2023) Visual Information Is Predictively Encoded in
Occipital Alpha/Low-Beta Oscillations. J Neurosci 43:5537–5545.

Van Veen BD, van Drongelen W, Yuchtman M, Suzuki A (1997) Localization of brain
electrical activity via linearly constrained minimum variance spatial filtering. IEEE
Trans Biomed Eng 44:867–880.

Watanabe T, Sasaki Y (2015) Perceptual learning: Toward a comprehensive theory.
Annu Rev Psychol 66:197–221.

Whittington JCR, Muller TH, Mark S, Chen G, Barry C, Burgess N, Behrens TEJ (2020)
The Tolman-Eichenbaum Machine: Unifying Space and Relational Memory
through Generalization in the Hippocampal Formation. Cell 183:1249-1263.e23.

Wimmer GE, Liu Y, Vehar N, Behrens TEJ, Dolan RJ (2020) Episodic memory retrieval
success is associated with rapid replay of episode content. Nature Neuroscience
23:1025–1033.

Wyart V, Nobre AC, Summerfield C (2012) Dissociable prior influences of signal
probability and relevance on visual contrast sensitivity. PNAS 109:3593–3598.

Xu S, Jiang W, Poo M, Dan Y (2012) Activity recall in a visual cortical ensemble. Nature
Neuroscience 15:449–455.

1006

 SUPPLEMENTARY INFORMATION

Figure S1. Sensor maps and spatial correlation of trained Lasso logistic regression models (related to Figure 3)

(A) Left: Sensor map for each state decoding model in Experiment 1; MEG signals from

72 occipital sensors were selected as features while training the classifier. Right:

Correlation matrix among classifiers. No spatial correlation was found among trained

classifiers (highest correlation; $r < 0.12$). **(B and C)** Same as (A), except the sensor

maps and correlation matrix correspond to Experiments 2 (Panel B, highest correlation;

$r < 0.1$) and 3 (Panel C, highest correlation; $r < 0.1$), respectively.

1017

Figure S2. Decoded feature representations in start- and end-only conditions (related to Figures 3)

Time series of reactivation probability output from the four regression models on an example trial. Time zero corresponds to trial onset. Both participants S08 and S12 were exposed with a fixed motion sequence of $0° \rightarrow 90° \rightarrow 180° \rightarrow 270°$.

1025

1026 **Figure S3. Sequenceness distribution across participants (related to Figures 4**

1027 **and 5)**

(A) In Experiment 1, of the 18 participants, 14 showed backward replay at 32 ms-lag for the start-only condition (left), and 13 showed backward replay at 32 ms-lag for the end-only condition (right). Each dot represents an individual participant. Backward and forward sequenceness is denoted by blue and yellow dots, respectively. The inset histogram shows the distribution of deviations from the unity line. **(B)** Same as described in (A), but for the results of Experiment 2 in the 2nd-only (left) and 3rd-only (right) conditions, respectively. **(C)** Same as described in (A) but for the results of Experiment 3 in the start- (left) and end-only (right) conditions, respectively.

1036

Figure S4. Sequenceness for each of the 24 possible orders (related to Figure 4)

(A) In the start-only condition of Experiment 1, only the backward sequence [4 → 3 → 2 → 1] with lags at 28–40 ms was significantly different (gray panels). The horizontal

1040   dashed lines represent significance thresholds derived from state label permutation at

1041   the 2^{nd}-level GLM matrix of temporally delayed linear modeling. **(B)** Same as described

1042   in (A) but for the end-only condition in Experiment 1.

1043

Figure S5. Stimuli and experimental procedures in Experiments 2 and 3 (related to

Figure 5)

(A) We presented the second (2nd-only condition) or third RDK (3rd-only condition) as a

cue in Experiment 2. **(B)** The procedure of Experiment 3 was identical to that of

Experiment 1 except for removing the exposure phase.