The role of the stress axis in mediating behavioural flexibility in a
social cichlid, *Neolamprologus pulcher*

Stefan Fischer[1,2], Zala Ferlinc[2], Katharina Hirschenhauser[3], Barbara Taborsky[4], Leonida Fusani[1,2] &
Sabine Tebbich[2]

[1]Konrad Lorenz Institute for Ethology, Department of Interdisciplinary Life Sciences, University of
Veterinary Medicine Vienna, Savoyenstrasse 1, 1160 Vienna, Austria

[2]Department of Behavioral and Cognitive Biology, University of Vienna, Djerassiplatz 1, 1030 Vienna,
Austria

[3]University College for Education of Upper Austria (PH OÖ), Kaplanhofstraße 40, 4020 Linz, Austria

[4]Division of Behavioural Ecology, University of Bern, Wohlenstrasse 50a, CH-3032 Hinterkappelen,
Switzerland

Correspondence and requests for materials should be addressed to S.F.
(email:Stefan.fischer@vetmeduni.ac.at)

ORCIDs:

0000-0001-8811-7518 (Stefan Fischer)

0000-0001-8710-9022 (Katharina Hirschenhauser)

0000-0003-1690-8155 (Barbara Taborsky)

0000-0001-8900-796X (Leonida Fusani)

0000-0003-3971-2139 (Sabine Tebbich)

Abstract

Behavioural flexibility plays a major role in the way animals cope with novel situations, and

physiological stress responses are adaptive and highly efficient mechanisms to cope with

unpredictable events. Previous studies investigating the role of stress responses in mediating

behavioural flexibility were mostly done in laboratory rodents using stressors and cognitive

challenges unrelated to the ecology of the species. To better understand how stress mediates

behavioural flexibility in a natural context, direct manipulation of the stress response and cognitive

tests in ecologically relevant contexts are needed. To this aim, we pharmacologically blocked

glucocorticoid receptors (GR) in adult cichlid fish using a minimally invasive application of a GR

antagonist. GRs blockade prevents the termination of the stress response, which we predicted to

impair behavioural flexibility. After the application of the GR antagonist, we repeatedly exposed fish

to a predator and tested their behavioural flexibility using a detour task, i.e. fish had to find a new,

longer route to the shelter when the shortest route was blocked. While the latencies to find the

shelter were not different between treatments, GR blocked fish showed more failed attempts during

the detour tasks than control fish. Furthermore, reduced performance was accompanied by an

increase of fear related behaviours. This suggests that blocking GRs plays an important role in

determining perceived fear and resulted in an impaired behavioural flexibility. We confirm a causal

role of terminating the stress response in mediating behavioural flexibility with potential

consequences for an effective and adaptive coping with changing environments.

Introduction

Stress is a ubiquitous feature of life and all organisms have evolved internal mechanisms to cope

with external stressors (Taborsky et al., 2021). Stressors can be abiotic or biotic, ranging from

extreme temperatures to predator attacks or social stressors such as aggressive conflicts. An

adequate response to a stressor, the so-called stress response, has major fitness consequences

because it is of utmost importance for the survival of the individual (Lemonnier et al., 2022; McEwen

and Wingfield, 2003; Romero et al., 2009; Sopinka et al., 2015).

When a stressor is experienced the adaptive stress response in vertebrates is generally characterised

by a physiological increase of circulating glucocorticoids (GCs) (Sapolsky et al., 2000). Cortisol is the

major GC and a main actor of the stress response in fish and most mammals (Sopinka et al., 2015).

Cortisol can cross the blood-brain barrier where it binds to high affinity mineralcorticoid receptors

and lower affinity glucocorticoid receptors (GRs) that are expressed across the brain (Joels et al.,

2008; Roozendaal et al., 2009). Cortisol acts on GRs to induce a negative feedback that reduces the

further release of cortisol, which slowly returns to baseline levels (Joels et al., 2008). This swift

increase and slow decrease of cortisol is an adaptive stress response leading to beneficial changes in

physiology, behaviour and cognition to successfully cope with the stressor. On the contrary, if

cortisol concentrations do not return to baseline levels after the stressor ends, prolonged elevation

of cortisol has adverse effects on health and behaviour particularly on cognitive abilities (Sandi and

Pinelo-Nava, 2007; Schwabe et al., 2012).

Behavioural flexibility is an important mechanism to cope with environmental change and requires

different cognitive abilities including learning and the seeking of alternative solutions (Tebbich et al.,

2010). GR activity has been shown to play an important role in mediating cognition and particularly

in regulating learning performances (for reviews see: de Kloet et al., 2005; Roozendaal et al., 2009;

Sandi and Pinelo-Nava, 2007). For example, laboratory rats receiving an application of a specific GR

antagonist (mifepristone) performed worse in a spatial learning task than rats receiving a control

manipulation (Oitzl and Dekloet, 1992). Conversely, an early-life treatment of the same GR

antagonist in a social cichlid (*Neolamprologus pulcher*), marginally enhanced learning but had no

effects on reversal learning, a commonly used proxy for behavioural flexibility (Reyes-Contreras and

Taborsky, 2022). Thus, more research is needed to better understand the causal role of GRs in

mediating cognition and in particular behavioural flexibility. Furthermore, it is not known whether a

short-term application of a GR antagonist immediately before a cognitive challenge is sufficient to

induce changes in behavioural flexibility.

Previous studies that tried to link mediators of the stress axis to behavioural flexibility in adult

individuals manipulated mostly type or duration of stressor (e.g. Bryce and Howland, 2015; Butts et

al., 2013; Calandreau et al., 2011; George et al., 2015; Graybeal et al., 2014; Graybeal et al., 2011;

Thai et al., 2013). However, we do not know whether the responsiveness of the stress axis to the

same stressor impairs or facilitates behavioural flexibility. Furthermore, most studies have been

conducted on laboratory rodents using stressors and cognitive tasks with limited ecological

relevance. Such studies are limited in their ability to reveal how adaptive stress responses mediate

behavioural flexibility in a changing environment. A more appropriate approach to study whether

adaptive stress responses determine behavioural flexibility is to (i) pharmacologically block GR

activity to verify the causal role of the stress response in mediating behavioural flexibility and (ii) use

ecologically relevant stressors and cognitive challenges.

In this work, we administered, minimally invasive, a GR antagonist (mifepristone) to adult *N. pulcher*

and tested their behavioural flexibility in a spatial task after being repeatedly exposed to a predator.

In this spatial task, individuals had to navigate through a maze to hide in a shelter after encountering

a predator. After displaying flight responses using the shortest route to the shelter, we tested each

individual`s ability to find a new route to the shelter when the shortest route was blocked. Our study

species *N. pulcher* is a highly social cichlid and in nature individuals live in large groups consisting of

dominant and subordinate individuals that are structured in a size-based, linear hierarchy (Wong and

Balshine, 2011). Individuals that live in more complex social environments often have enhanced

cognitive skills, including an enhanced behavioural flexibility (e.g. Amici et al., 2008; Ashton et al.,

2018; Bond et al., 2007; Johnson-Ulrich and Holekamp, 2020).

Based on the role that GRs play in the negative feedback mechanism to terminate stress responses

we formulated two predictions: (1) GR antagonist-treated fish show impaired behavioural flexibility,

particularly after repeated exposures to a predator. (2) Effects that are causally mediated by GRs are

specific to behavioural flexibility and the ability to cope with stress and do not affect other

behaviours such as the motivation of the fish to solve the task.

Methods

Subjects and housing

The experiment was conducted at the Konrad Lorenz Institute of Ethology, University of Veterinary

Medicine, Vienna, Austria, under the licence (ETK-101/06/2019). All experimental tanks (45L, 30 x 50

x 30 cm) were equipped with a 2 cm layer of sand, a biological filter, a heater and one-half clay

flowerpot that served as a shelter. The water temperature was kept at 27 ± 1°C with a light-dark

regime of 13:11 h and a dimmed-light phase of 10 min. During the experiment all fish were fed

commercial flake food 6 days a week.

Our study species was the cooperatively breeding cichlid *Neolamprologus pulcher* because stress

responses are well researched (Antunes et al., 2021a; Mileva et al., 2009) in this species, and it is a

model to study the evolution of vertebrate sociality highlighting their social and non-social

behavioural flexibility (Taborsky, 2016; Wong and Balshine, 2011). These fish are endemic to Lake

Tanganyika and groups always consist of a dominant breeding pair and related or unrelated helpers

(Balshine et al., 2001; Heg et al., 2005; Taborsky, 1984, 1985). Helpers assist the breeding pair in

raising the offspring by cleaning and fanning the eggs, cleaning the breeding shelters, and defending

the common territory (Grantner and Taborsky, 1998; Taborsky, 1984, 1985). Larger territories are divided into sub-territories where each group member occupies their own shelter. As an ecologically relevant stress stimulus, we used the piscivorous cichlid *Lepidiolamprologus elongatus* which occurs in sympatry with *N. pulcher* (Ochi and Yanagisawa, 1998). *L. elongatus* preys on all sizes of *N. pulcher*, except eggs, and is one of its most important predators (Groenewoud et al., 2016). Previous research indicates that *N. pulcher* can innately recognise this predator based on visual or olfactory cues (Fischer et al., 2014; Zöttl et al., 2013). In nature, groups of *N. pulcher* face a high predation pressure and individuals either attack or hide when predators intrude their territories, particularly when faced with a large predator such as *L. elongatus*. Sexually mature *N. pulcher* feed on zooplankton in the water column that requires them to stay far away from their shelters (Taborsky, 2016). Thus, flexibility in the way how individuals return to their shelter in case of danger is of utmost importance for their survival.

Our test fish were laboratory-reared first- and second-generation offspring of wild-caught fish. We randomly selected 35 male and 35 female adult fish (N=70, mean standard length [SL; from the tip of the snout to the end of the caudal peduncle]: 7.2 cm, [range: 5.6-8.2 cm]) from the laboratory stock, where they had been kept before in tanks separated by sex. The experiment was conducted in seven blocks over seven weeks with 10 test fish per block. Test fish were permanently marked using subcutaneous injections of Visible Implant Elastomer tags (VIE; Northwest 191 Marine Technology), at least three days before each block started, except for one block where test fish had been marked one day before the experiment started. To avoid any confounding effects, we excluded all test fish from this block from the behavioural analysis because some individuals had not fully recovered from the marking procedures which reduced the final sample size to N=60.

We also randomly chose adult *L. elongatus* fish (N = 10, mean SL: 12.68 cm [range: 10.3-14.7 cm]) from the laboratory stock population, where they had been housed in pairs or small groups. To avoid exposing predators to repeated stress from catching and transporting them to the presentation

tanks (see below), which could influence their behaviours, we used five individuals for the first three

blocks and another five individuals for the last four blocks. During the experiment, each predator

was kept in a separate 45L tank. These tanks were set-up close to the test fish but with a visual and

olfactory separation. Body sizes of predators were obtained after the last block of the experiment

was completed. All fish were kept under the same housing conditions and food regimes before and

during the experiment.

Experimental set-up

During the trials, we visually separated the experimental tanks using opaque dividers. The mazes

used to carry out the detour trials remained in the experimental tanks throughout all the

experimental phases (see below). The tanks were separated from the observer and the rest of the

room using a dark curtain to minimize any disturbances from the surrounding area. All trials were

recorded from above and from the front using two web cameras (Creative Live Cam Sync HD

webcam) placed inside the curtain and connected to a computer outside of the curtain. Prior to each

trial, a randomly chosen predator was transported to a small presentation tank (25 x 30 cm) filled

with water, that was placed on a small movable rack. The front of the presentation tank – the part

facing the experimental tanks – was covered with an opaque divider which could be moved with a

string by the observer from outside the curtain. The divider was lifted before each trial for 60s to

visually expose the test fish to the predator.

For all the trials, we used a transparent, U-shaped maze that prevented fish from using the maze as a

shelter. The walls of the maze were covered with a mesh-like patterning that was fine enough to

clearly see though. Halved clay flowerpots were used as shelters. The shelter could be reached using

two routes: either via the direct route where fish only entered the arm of the maze that contains the

shelter ('shelter arm', see Figure 1), or via a detour route where fish entered first the arm furthest

away from the shelter ('non-shelter arm', see Figure 1)and had to cross to the other side in the back

of the maze to enter the shelter arm. The middle arm of the maze was a dead end (Figure 1). The

area in front of the maze (30 x 10 cm) that faced towards the predator was used as a starting area

that could be separated from the rest of the tank with a removable opaque divider. Again, using a

string the observer could remotely move the divider with minimal disturbance to the fish from

outside the curtain. Biological filters and heaters were removed during the trials to ensure fish could

only seek shelter inside the flowerpot.

Experimental phases

Each block consisted of five distinctive experimental phases: a habituation phase (day 1, day 2), a

pre-treatment direct phase (day 3), a treatment immersion phase (day 4, day 5), a post-treatment

direct phase (day 6), and a detour phase (day 6).

Habituation. After fish were measured (SL) to the closest mm, each fish was individually transported

into one of the experimental tanks. Each tank included a maze, and a shelter was placed in the

shelter arm. Fish were allowed to habituate to the new environment for 48h. During the habituation

phase we removed the opaque dividers between the experimental tanks to allow fish to visually

interact with their neighbours.

Pre-treatment direct. After the habituation phase (day 3, 10a.m.), we tested whether the fish used

the shortest route to the shelter after being exposed to a brief predator presentation. First, the

observer gently guided the fish to the starting area using a small handheld net. Second, the observer

lowered the divider to separate the start area from the rest of the tank. Third, the observer added

an olfactory predator cue (15 ml of water from a predator tank) to the starting area and lifted the

divider inside the predator tank to visually expose the fish to the predator for 60s. Fourth, directly

afterwards, the observer lifted the divider separating the start area from the rest of the tank, and

the fish could enter the maze and seek shelter. In this phase, the entrance of the shelter faced

towards the starting area. Fish successfully completed this phase if they used the direct route to

reach the shelter within 60s without any detours. Fish that did not use the direct route (N=8) were

excluded from the experiment.

Immersion treatment. The subjects that successfully completed the previous phase (N=52),

underwent a 48h period of either a mifepristone (N=26, females=14) or a control immersion

treatment (N=26, females=13) starting on the next day (day 4, 8a.m.). Half of the females and half of

the males were randomly assigned to one of the two treatments, in a way that ensured that the

observer was blind to the treatments of the fish within each block. For this a second person (SF)

applied the treatments to the experimental tanks without the observer (ZF) present. The treatments

were applied in the form of water baths following Nyman et al. (2018). In short, we dissolved

mifepristone (RU 484) in dimethylsulfoxide at 50 mg ml-1, then serially diluted it in 0.1 M acetic acid

(1:10), phosphate-buffered saline (1:100), and finally, diluted it in distilled water for an immersion

concentration of 400 ng l^{-1}. Controls were prepared with diluents without mifepristone. Both

mifepristone and control solutions were prepared and stored in a -20C freezer one week prior to the

experiment. The fish were immersed in 2L of water inside 3L glass containers that were place inside

the experimental tank.

Post-treatment direct. After the immersion treatments (day 6, 9a.m.) we repeated the previous

direct phase to test whether (i) the fish memorised the shortest route to the shelter, and (ii) whether

the immersion treatments influenced the motivation or memory of fish. Seven out of 52 fish did not

take the direct route in this phase. The immersion treatments did not influence whether fish took

the short or the longer route (control: 4/26; mifepristone: 3/26). This time all fish proceeded to the

detour phase because they already used the direct route in the pre-treatment direct phase. This

ensured that (i) all fish that proceeded to the detour phase had used at least once the direct route

and (ii) the immersion treatments had no effect on the motivation of the fish to seek shelter.

Detour. For the detour trials (day 6, 10:30a.m.) we blocked the direct route to the shelter arm (Fig.

1) with a rectangular barrier (5 x 25 cm), made from the same material as the maze. The shelter

entrance was turned to face away from the starting area. This ensured that fish approached the

shelter from the same side irrespective of whether they entered the shelter arm via the direct or via

the longer route (Figure 1). We used the same procedures as described for the pre-treatment direct

phase. Each fish experienced 4 detour trials with a 2h recovery period between each trial.

Video analysis

Video recordings were analysed using BORIS (Friard and Gamba, 2016). One trial (block 3, detour 1)

was excluded from the analysis due to problems with the video quality. For each trial the following

information was recorded: (i) the type of the test (post-treatment direct/detour 1/detour 2/detour

3/detour 4), (ii) the latency to cross the shelter arm line, (iii) the latency to enter the shelter, (iv)

whether the fish entered the shelter or not and (v) the number of failed attempts. In the post-

treatment direct trials, we additionally recorded whether the fish used the shortest route to the

shelter or not. A failed attempt in the post-treatment direct trials was defined as either crossing the

middle arm line or the non-shelter arm line. A failed attempt in the detour trials was defined as

crossing the middle arm line, attempting, and trying to cross the blocked shelter arm line, or crossing

the non-shelter arm line but then returning back to the starting area without entering the shelter

arm. A fish was classified as having crossed into a new arm when both pectoral fins crossed the

virtual line separating the two arms. We started measuring the latencies immediately after the

predator exposure – i.e. immediately after the divider in the presentation tank was closed and the

divider in the experimental tank was lifted. If fish did not cross any arm line or entered the shelter

before the end of the experiment, we assigned maximum latencies (600s) in the respective trial.

Statistical analysis

Data were analysed using R 4.1.1. (R Core Team, 2021) with the packages "lme4" (Bates et al., 2015)

and "afex" (Singmann et al., 2021). We analysed treatment effects using Linear Models (LMs),

Generalized Linear Models (GLMs), Linear Mixed Models (LMMs) and Generalized Linear Mixed

Models (GLMMs). Residuals and Q/Q-plots of all LMs and LMMs were visually inspected, and the

distributions of residuals were compared to a normal distribution using Kolmogorov-Smirnov and

Shapiro tests. If residuals were not normally distributed, a log transformation was applied, and

residuals again checked. All GLMs and GLMMs were checked for over-dispersion, but none required

a correction. To obtain p-values, we used either the mixed() function in the package 'afex' with a

Satterthwaite's approximation for degrees of freedom, or the drop1() function to perform either a

type II anova (for LM and LMM) or a χ^2-test (for GLM and GLMM). Interactions, and control variables

(sex and body size of test fish) were stepwise removed if non-significant (Engqvist, 2005) and

reported p-values refer to the final model without non-significant interactions, and control variables.

An overview of all full models can be found in Supplementary Table S1.

Except for the analyses of the probability to enter the shelter with increasing numbers of predator

exposures (see below) we always included 'Treatment' (GR antagonist or control treatment), the

body size and sex of test fish in the models. When analysing the detour trials, we additionally

included test fish identity as a random effect to control for multiple observations on the same

individual and the number of the detour trial as a fixed effect. If the interaction 'Nr. of detour trial x

Treatment' was significant we compared the marginal means of both treatments within each detour

trial separately (four comparisons) with a Tukey correction for multiple testing. To analyse the

probability to enter the shelter with increasing numbers of predator presentations we used all trials

that were done after the fish received their respective treatments (post-treatment direct, detour 1,

detour 2, detour 3, detour 4). Fish identity was included as a random effect and the number of daily

predator presentations as a continuous covariate. Initially we constructed two models, one that

included a linear interaction between the number of daily predator presentations and treatment and

one that included a non-linear, polynomial relationship with a degree of 2. We compared both

models using a χ^2- goodness of fit test. To foster comparability, we kept the structure of the models

similar and did not include the body size or sex of fish in this analysis. To further investigate the

significant non-linear interaction, we constructed two additional models separately for GR blocked

and control fish. Both models included the identity of the fish as a random factor and the number of

daily predator presentations as a linear predictor or as a non-linear polynomial predictor with a

degree of 2.

Results

Does the application of a GR antagonist influence memory or motivation to seek shelter in the post-

treatment learning trials?

As expected, we found no behavioural differences between the GR blocked and control fish in the

post-treatment direct trials. There were no significant differences between the two immersion

treatments in the latency to cross the shelter line ($F_{1,50}=0.08$, $p=0.78$; see factor: 'Treatment' in Table

1a), the latency to enter the shelter ($F_{1,50}=0.1$, $p=0.77$; see factor 'Treatment' in Table 1b) or the

probability to show a failed attempt ($\chi^2=0.12$, $p=0.73$, N=52; see factor 'Treatment' in Table 2).

Does the application of a GR antagonist influence behavioural flexibility in the detour trials during

repeated exposures to a stressor?

GR blocked fish showed less behavioural flexibility during the detour trials than control fish. That is,

the application of a GR antagonist influenced the number of errors that fish made over time with

repeated exposures to predators ($\chi^2=23.96$, $p<0.01$, N=52; see interaction 'Nr. of detour trial x

Treatment' in Table 3, Figure 2). Post-hoc analysis revealed that GR blocked fish made significantly

more errors in the third detour trial compared to control fish (see Table 4 for pairwise comparisons).

In contrast, GR blocked fish did not show a longer latency to enter the shelter arm ($F_{1,50.11}=0.0$,

p=0.97, see factor 'Treatment' in Table 5a) or a longer latency to enter the shelter ($F_{1,48.98}=0.42$,

p=0.52, see factor 'Treatment' in Table 5b) than the control fish. Overall, all fish reduced the latency

to enter the shelter arm with increasing predator exposures ($F_{1,152.25}=3.78$, p=0.01; see factor 'Nr. of

detour trial' in Table 5a), whereas this was not the case for the latency to enter the shelter

($F_{1,152.18}=0.6$, p=0.61; see factor 'Nr. of detour trial' in Table 5b). Irrespective of the treatments, we

found a positive relationship between body size and the latency to enter the shelter ($F_{1,48.9}=5.1$,

p=0.03; see factor 'Body size' in Table 5b) indicating that larger fish took longer to enter the shelter.

Does the application of a GR antagonist determine how fish behaviourally cope with repeated

exposures to a stressor?

When analysing the probability of fish to enter the shelter with increasing numbers of daily predator

exposures, we found that a model containing a non-linear interaction between the number of daily

predator exposures and treatment explained the data significantly better than a linear interaction

term ($\chi2=15.37$, p<0.01). GR blocked fish showed a linear reduction in the probability to enter the

shelter with increasing numbers of daily predator exposures (z=-2.17, p=0.03, N=62; see factor 'Nr.

of predator exposure' in Table 6a). In contrast, control fish showed a non-linear, U-shaped

relationship between the probability to enter the shelter and the number of daily predator

exposures [z=3.26, p<0.01, N=26; see factor 'poly (Nr. of predator exposure, 2)' in Table 6b]. Here,

control fish entered the shelter more frequently during the initial and the last predator exposures

than during any of the intermediate predator exposures (Figure 3).

Discussion

In the present experiment we applied, minimally invasive, a GR antagonist or a control immersion to

groups of fish and subsequently tested their behavioural flexibility after repeated exposures to an

ecologically relevant stressor. Fish treated with a GR antagonist (1) made more errors and (2) had a

higher probability to enter the shelter during the intermediate compared to the initial and last

predator exposures. Furthermore, the application of a GR antagonist did not influence the individual

motivation to participate in the task or their spatial memory.

Our results show that GR play an important role for behavioural flexibility in a spatial task when

repeatedly exposed to a mild naturalistic stressor. Previous studies using laboratory rodents have

shown that GR play a causal role in determining memory formation (Oitzl and Dekloet, 1992;

Roozendaal and McGaugh, 1997), fear conditioning (Donley et al., 2005), and reversal learning

(Bryce and Howland, 2015). Taken together with the current study, these results indicate that GR

activity affects specific cognitive abilities across a diverse set of cognitive tasks. In contrast to

previous studies, we used an ecologically relevant task where fish had to find a new route to avoid a

predator. This revealed that the stress response and particularly the activity of GRs play an

important role in determining the way individuals cope with environmental change.

A slower retention of previously formed associations during an increasingly stressful situation might

be the underlying mechanism determining the observed behavioural differences between GR

blocked and control fish. Although GRs are distributed across the entire brain, GR activity in the

basolateral amygdala (BLA) and the ventral hippocampus (VH) have been specifically linked to the

retention of fear memory (Donley et al., 2005). Similarly, we found that GR blocked fish showed a

higher probability to enter the shelter during the intermediate predator presentations compared to

control fish. This might be an indicator of a higher perceived fear in GR blocked fish. We assume that

the higher number of fear related behaviours was associated with increased concentrations of

circulating GCs, in our case cortisol because previous studies showed that GCs are essential for fear

related behaviours (for review see Schulkin et al., 2005). Cortisol might have accumulated more in

GR blocked fish due to their reduced ability to terminate the stress response before encountering

the next stressful predator exposure. Interestingly, this increase in fear related behaviours and thus

cortisol was accompanied by a reduction in behavioural flexibility as shown by a higher number of

errors of GR blocked fish while searching for the new, longer route to the shelter.

Generally, the effects of GC concentrations on cognition are diverse and context dependent (for

reviews see de Kloet et al., 2005; Roozendaal et al., 2009; Sandi and Pinelo-Nava, 2007). In *N.*

pulcher experimentally elevating GCs during early life reduced social competence and behavioural

flexibility in a reversal learning task (Reyes-Contreras et al., 2019; Reyes-Contreras and Taborsky,

2022). In contrast, a study using the same GR antagonist in rainbow trout found that GR blocked fish

had significantly reduced GC concentrations after experiencing a single stressor compared to control

fish (Alderman et al., 2012). The current study employed adult fish and repeated predator exposures

and we cannot conclude whether GC concentrations have increased or decreased after repeated

exposures to a predator. Further investigations are needed to disentangle the complex interplay

between GCs, GRs and cognition particularly when repeatedly experiencing a stressor during

development or later in life.

Behavioural flexibility generally refers to adaptive changes in behaviour, in response to

environmental changes (Brown and Tait, 2014). The flexible adjustment of behaviour is thought to

involve several cognitively demanding components such as learning and memory (Tebbich et al.,

2010), and flexibility has also been used as evidence of complex cognition (Mikhalevich et al., 2017).

On an evolutionary perspective, complex cognition covaries with the increased demands of more

complex or unpredictable social ('social brain hypothesis' Shultz and Dunbar, 2007) or ecological

environments ('cognitive buffer hypothesis' Sol, 2009). For example, between species comparisons

of varying sociality, as well as within species comparisons of individuals that experience different

social environments showed that the demands of a more complex social life are important

evolutionary drivers of behavioural flexibility (Ashton et al., 2018; Bond et al., 2007). Similarly to the

social environment, a more complex ecological environment, for example exposure to predation

cues has been linked to a stronger learnt antipredator response and longer memory retention

(Ferrari, 2014; Ferrari et al., 2005; Mitchell et al., 2016).

In *N. pulcher*, social and ecological experiences during the first two month of life are of particular

relevance for the development of social and anti-predator behaviours, as well as for non-social

behavioural flexibility (Bannier et al., 2017; Fischer et al., 2017). These organisational effects are

most likely mediated by a life-long reprogramming of the physiological stress axis (Antunes et al.,

2021a; Antunes et al., 2021b; Nyman et al., 2018) although an early-life manipulation of the stress

axis using the same GR antagonist did not determine behavioural flexibility when adult individuals

were tested in a reversal learning paradigm (Reyes-Contreras and Taborsky, 2022). In contrast, our

study shows an activational effect on behavioural flexibility mediated by differences in the

immediate stress response to a predator. Future studies focusing on both the organizational and the

activational effects of the stress response in determining cognition and in particular behavioural

flexibility are needed to better understand the way how animals respond to naturalistic

environmental challenges.

In conclusion, our study provides novel insights into the way how animals adapt to environmental

change. In this study we used an ecologically relevant stressor and cognitive task to highlight the role

of GRs in mediating behavioural flexibility. The results of the current study may stimulate further

research to better understand how stress responses shape cognitive abilities and their associated

behaviours in a natural context.

Data availability

Data for this study are available at Figshare public repository:

https://doi.org/10.6084/m9.figshare.24083952

References

Alderman, S.L., McGuire, A., Bernier, N.J., Vijayan, M.M., 2012. Central and peripheral glucocorticoid
396   receptors are involved cortisol response to an acute stressor in rainbow trout. Gen. Comp.
397   Endocrinol. 176, 79-85. https://doi.org/10.1016/j.ygcen.2011.12.031

399   Amici, F., Aureli, F., Call, J., 2008. Fission-fusion dynamics, behavioral flexibility, and inhibitory
400   control in primates. Curr. Biol. 18, 1415-1419. https://doi.org/10.1016/j.cub.2008.08.020

402   Antunes, D.F., Reyes-Contreras, M., Glauser, G., Taborsky, B., 2021a. Early social experience has life-
403   long effects on baseline but not stress-induced cortisol levels in a cooperatively breeding
404   fish. Horm. Behav. 128, 104910. https://doi.org/10.1016/j.yhbeh.2020.104910

406   Antunes, D.F., Teles, M.C., Zuelling, M., Friesen, C.N., Oliveira, R.F., Aubin-Horth, N., Taborsky, B.,
407   2021b. Early social deprivation shapes neuronal programming of the social decision-making
408   network in a cooperatively breeding fish. Mol. Ecol. 30, 4132.
409   https://doi.org/10.1111/mec.16019

411   Ashton, B.J., Ridley, A.R., Edwards, E.K., Thornton, A., 2018. Cognitive performance is linked to group
412   size and affects fitness in Australian magpies. Nature 554, 364–367.
413   https://doi.org/10.1038/nature25503

415   Balshine, S., Leach, B., Neat, F., Reid, H., Taborsky, M., Werner, N., 2001. Correlates of group size in a
416   cooperatively breeding cichlid fish (*Neolamprologus pulcher*). Behav. Ecol. Sociobiol. 50, 134-
417   140. https://doi.org/10.1007/s002650100343

419   Bannier, F., Tebbich, S., Taborsky, B., 2017. Early experience affects learning performance and
420   neophobia in a cooperatively breeding cichlid. Ethology 123, 712-723.
421   https://doi.org/https://doi.org/10.1111/eth.12646

423   Bates, D., Maechler, M., Bolker, B.M., Walker, S., 2015. Fitting linear mixed-effects models using
424   {lme4}. J. Stat. Softw. 67, 1-48. https://doi.org/ https://doi.org/10.18637/jss.v067.i01

426   Bond, A.B., Kamil, A.C., Balda, R.P., 2007. Serial reversal learning and the evolution of behavioral
427   flexibility in three species of North American corvids (Gymnorhinus cyanocephalus,
428   Nucifraga columbiana, Aphelocoma californica). J. Comp. Psycho. 121, 372-379.
429   https://doi.org/10.1037/0735-7036.121.4.372

431   Brown, V.J., Tait, D.S., 2014. Behavioral Flexibility: Attentional Shifting, Rule Switching, and Response
432   Reversal, in: Stolerman, I., Price, L. (Eds.), Encyclopedia of Psychopharmacology. Springer,
433   Berlin, Heidelberg.

435   Bryce, C.A., Howland, J.G., 2015. Stress facilitates late reversal learning using a touchscreen-based
436   visual discrimination procedure in male Long Evans rats. Behav. Brain Res. 278, 21-28.
437   https://doi.org/10.1016/j.bbr.2014.09.027

439   Butts, K.A., Floresco, S.B., Phillips, A.G., 2013. Acute stress impairs set-shifting but not reversal
440   learning. Behav. Brain Res. 252, 222-229. https://doi.org/10.1016/j.bbr.2013.06.007

442   Calandreau, L., Bertin, A., Boissy, A., Arnould, C., Constantin, P., Desmedt, A., Guemene, D., Nowak,
443   R., Leterrier, C., 2011. Effect of one week of stress on emotional reactivity and learning and

444   memory performances in Japanese quail. Behav. Brain Res. 217, 104-110.
445   https://doi.org/10.1016/j.bbr.2010.10.004
446

447 de Kloet, E.R., Joels, M., Holsboer, F., 2005. Stress and the brain: from adaptation to disease. Nat.
448   Rev. Neurosci. 6, 463-475. https://doi.org/10.1038/nrn1683
449

450 Donley, M.P., Schulkin, J., Rosen, J.B., 2005. Glucocorticoid receptor antagonism in the basolateral
451   amygdala and ventral hippocampus interferes with long-term memory of contextual fear.
452   Behav. Brain Res. 164, 197-205. https://doi.org/10.1016/j.bbr.2005.06.020
453

454 Engqvist, L., 2005. The mistreatment of covariate interaction terms in linear model analyses of
455   behavioural and evolutionary ecology studies. Anim. Behav. 70, 967-971.
456   https://doi.org/10.1016/j.anbehav.2005.01.016
457

458 Ferrari, M.C.O., 2014. Short-term environmental variation in predation risk leads to differential
459   performance in predation-related cognitive function. Anim. Behav. 95, 9-14.
460   https://doi.org/10.1016/j.anbehav.2014.06.001
461

462 Ferrari, M.C.O., Trowell, J.J., Brown, G.E., Chivers, D.P., 2005. The role of learning in the
463   development of threat-sensitive predator avoidance by fathead minnows. Anim. Behav. 70,
464   777-784. https://doi.org/10.1016/j.anbehav.2005.01.009
465

466 Fischer, S., Bohn, L., Oberhummer, E., Nyman, C., Taborsky, B., 2017. Divergence of developmental
467   trajectories is triggered interactively by early social and ecological experience in a
468   cooperative breeder. Proc. Natl. Acad. Sci. U.S.A. 114, E9300-E9307.
469   https://doi.org/10.1073/pnas.1705934114
470

471 Fischer, S., Taborsky, B., Burlaud, R., Fernandez, A.A., Hess, S., Oberhummer, E., Frommen, J.G.,
472   2014. Animated images as a tool to study visual communication: a case study in a
473   cooperatively breeding cichlid. Behaviour 151, 1921-1942.
474   https://doi.org/10.1163/1568539x-00003223
475

476 Friard, O., Gamba, M., 2016. BORIS: a free, versatile open-source event-logging software for
477   video/audio coding and live observations. Methods Ecol. Evol. 7, 1325-1330.
478   https://doi.org/10.1111/2041-210x.12584
479

480 George, S.A., Rodriguez-Santiago, M., Riley, J., Abelson, J.L., Floresco, S.B., Liberzon, I., 2015.
481   Alterations in cognitive flexibility in a rat model of post-traumatic stress disorder. Behav.
482   Brain Res. 286, 256-264. https://doi.org/10.1016/j.bbr.2015.02.051
483

484 Grantner, A., Taborsky, M., 1998. The metabolic rates associated with resting, and with the
485   performance of agonistic, submissive and digging behaviours in the cichlid fish
486   *Neolamprologus pulcher* (Pisces: Cichlidae). J. Comp. Physiol. 168, 427-433.
487   https://doi.org/10.1007/s003600050162
488

489 Graybeal, C., Bachu, M., Mozhui, K., Saksida, L.M., Bussey, T.J., Sagalyn, E., Williams, R.W., Holmes,
490   A., 2014. Strains and stressors: an analysis of touchscreen learning in genetically diverse
491   mouse strains. Plos ONE 9, e87745. https://doi.org/10.1371/journal.pone.0087745
492

493   Graybeal, C., Feyder, M., Schulman, E., Saksida, L.M., Bussey, T.J., Brigman, J.L., Holmes, A., 2011.
494   Paradoxical reversal learning enhancement by stress or prefrontal cortical damage: rescue
495   with BDNF. Nat. Neurosci. 14, 1507-1509. https://doi.org/10.1038/nn.2954
496
497   Groenewoud, F., Frommen, J.G., Josi, D., Tanaka, H., Jungwirth, A., Taborsky, M., 2016. Predation risk
498   drives social complexity in cooperative breeders. Proc. Natl. Acad. Sci. U.S.A., 4104–4109.
499   https://doi.org/10.1073/pnas.1524178113
500
501   Heg, D., Brouwer, L., Bachar, Z., Taborsky, M., 2005. Large group size yields group stability in the
502   cooperatively breeding cichlid *Neolamprologus pulcher*. Behaviour 142, 1615-1641.
503   https://doi.org/10.1163/156853905774831891
504
505   Joels, M., Karst, H., DeRijk, R., de Kloet, E.R., 2008. The coming out of the brain mineralocorticoid
506   receptor. Trends Neurosci. 31, 1-7. https://doi.org/10.1016/j.tins.2007.10.005
507
508   Johnson-Ulrich, L., Holekamp, K.E., 2020. Group size and social rank predict inhibitory control in
509   spotted hyaenas. Anim. Behav. 160, 157-168.
510   https://doi.org/10.1016/j.anbehav.2019.11.020
511
512   Lemonnier, C., Bize, P., Boonstra, R., Dobson, F.S., Criscuolo, F., Viblanc, V.A., 2022. Effects of the
513   social environment on vertebrate fitness and health in nature: Moving beyond the stress
514   axis. Horm. Behav. 145, 105232. https://doi.org/10.1016/j.yhbeh.2022.105232
515
516   McEwen, B.S., Wingfield, J.C., 2003. The concept of allostasis in biology and biomedicine. Horm.
517   Behav. 43, 2-15. https://doi.org/10.1016/s0018-506x(02)00024-7
518
519   Mikhalevich, I., Powell, R., Logan, C., 2017. Is behavioural flexibility evidence of cognitive
520   complexity? How evolution can inform comparative cognition. Interface Focus 7, 20160121.
521   https://doi.org/10.1098/rsfs.2016.0121
522
523   Mileva, V.R., Fitzpatrick, J.L., Marsh-Rollo, S., Gilmour, K.M., Wood, C.M., Balshine, S., 2009. The
524   stress response of the highly social African cichlid *Neolamprologus pulcher*. Physiol.
525   Biochem. Zool. 82, 720-729. https://doi.org/10.1086/605937
526
527   Mitchell, M.D., Chivers, D.P., Brown, G.E., Ferrari, M.C.O., 2016. Living on the edge: how does
528   environmental risk affect the behavioural and cognitive ecology of prey? Anim. Behav. 115,
529   185-192. https://doi.org/10.1016/j.anbehav.2016.03.018
530
531   Nyman, C., Fischer, S., Aubin-Horth, N., Taborsky, B., 2018. Evolutionary conserved neural signature
532   of early life stress affects animal social competence. Proc. R. Soc. B-Biol. Sci. 285, 9.
533   https://doi.org/10.1098/rspb.2017.2344
534
535   Ochi, H., Yanagisawa, Y., 1998. Commensalism between cichlid fishes through differential tolerance
536   of guarding parents toward intruders. J. Fish Biol. 52, 985-996.
537   https://doi.org/10.1111/j.1095-8649.1998.tb00598.x
538
539   Oitzl, M.S., Dekloet, E.R., 1992. Selective Corticosteroid antagonists modulate specific aspects of
540   spatial orientation learning. Behav. Neurosci. 106, 62-71. https://doi.org/10.1037/0735-
541   7044.106.1.62
542
R Core Team, 2021. R: a language and environment for statistical computing.

Reyes-Contreras, M., Glauser, G., Rennison, D.J., Taborsky, B., 2019. Early-life manipulation of
cortisol and its receptor alters stress axis programming and social competence. Philos. Trans.
R. Soc. B-Biol. Sci. 374. https://doi.org/10.1098/rstb.2018.0119

Reyes-Contreras, M., Taborsky, B., 2022. Stress axis programming generates long-term effects on
cognitive abilities in a cooperative breeder. Proc. R. Soc. B-Biol. Sci. 289, 20220117.
https://doi.org/10.1098/rspb.2022.0117

Romero, L.M., Dickens, M.J., Cyr, N.E., 2009. The reactive scope model - a new model integrating
homeostasis, allostasis, and stress. Horm. Behav. 55, 375-389.
https://doi.org/10.1016/j.yhbeh.2008.12.009

Roozendaal, B., McEwen, B.S., Chattarji, S., 2009. Stress, memory and the amygdala. Nat. Rev.
Neurosci. 10, 423-433. https://doi.org/10.1038/nrn2651

Roozendaal, B., McGaugh, J.L., 1997. Glucocorticoid receptor agonist and antagonist administration
into the basolateral but not central amygdala modulates memory storage. Neurobiol. Learn.
Mem. 67, 176-179. https://doi.org/10.1006/nlme.1996.3765

Sandi, C., Pinelo-Nava, M.T., 2007. Stress and memory: behavioral effects and neurobiological
mechanisms. Neural Plasticity 2007, 2007: 78970. https://doi.org/10.1155/2007/78970

Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions.
Endocr. Rev. 21, 55-89. https://doi.org/10.1210/er.21.1.55

Schulkin, J., Morgan, M.A., Rosen, J.B., 2005. A neuroendocrine mechanism for sustaining fear.
Trends Neurosci. 28, 629-635. https://doi.org/10.1016/j.tins.2005.09.009

Schwabe, L., Joels, M., Roozendaal, B., Wolf, O.T., Oitzl, M.S., 2012. Stress effects on memory: an
update and integration. Neurosci. Biobehav. Rev. 36, 1740-1749.
https://doi.org/10.1016/j.neubiorev.2011.07.002

Shultz, S., Dunbar, R.I.M., 2007. The evolution of the social brain: anthropoid primates contrast with
other vertebrates. Proc. R. Soc. B-Biol. Sci. 274, 2429-2436.
https://doi.org/10.1098/rspb.2007.0693

Singmann, H., Bolker, B., Westfall, J., Aust, F., Ben-Shachar, M.S., 2021. afex: analysis of factorial
experiments, R package version 1.0-1 ed.

Sol, D., 2009. Revisiting the cognitive buffer hypothesis for the evolution of large brains. Biol. Lett. 5,
130-133. https://doi.org/10.1098/rsbl.2008.0621

Sopinka, N.M., Patterson, L.D., Redfern, J.C., Pleizier, N.K., Belanger, C.B., Midwood, J.D., Crossin,
G.T., Cooke, S.J., 2015. Manipulating glucocorticoids in wild animals: basic and applied
perspectives. Conserv. Physiol. 3, cov031. https://doi.org/10.1093/conphys/cov031

Taborsky, B., English, S., Fawcett, T.W., Kuijper, B., Leimar, O., McNamara, J.M., Ruuskanen, S., Sandi,
C., 2021. Towards an Evolutionary Theory of Stress Responses. Trends Ecol. Evol. 36, 39-48.
https://doi.org/10.1016/j.tree.2020.09.003

Taborsky, M., 1984. Broodcare helpers in the cichlid fish *Lamprologus brichardi*: their costs and
benefits. Anim. Behav. 32, 1236-1252. https://doi.org/10.1016/s0003-3472(84)80241-9
598

599 Taborsky, M., 1985. Breeder-helper conflict in a cichlid fish with broodcare helpers: an experimental
600   analysis. Behaviour 95, 45-75. https://doi.org/10.1163/156853985x00046
601

602 Taborsky, M., 2016. Cichlid fishes: A model for the integrative study of social behavior, in: Koenig,
603   W., Dickinson, J. (Eds.), Cooperative Breeding in Vertebrates: Studies of Ecology, Evolution,
604   and Behavior. Cambridge University Press, Cambridge.
605

606 Tebbich, S., Sterelny, K., Teschke, I., 2010. The tale of the finch: adaptive radiation and behavioural
607   flexibility. Philos. Trans. R. Soc. B-Biol. Sci. 365, 1099-1109.
608   https://doi.org/10.1098/rstb.2009.0291
609

610 Thai, C.A., Zhang, Y., Howland, J.G., 2013. Effects of acute restraint stress on set-shifting and reversal
611   learning in male rats. Cogn. Affect. Behav. Neurosci. 13, 164-173.
612   https://doi.org/10.3758/s13415-012-0124-8
613

614 Wong, M., Balshine, S., 2011. The evolution of cooperative breeding in the African cichlid fish,
615   *Neolamprologus pulcher*. Biol. Rev. 86, 511-530.
616   https://doi.org/https://doi.org/10.1111/j.1469-185X.2010.00158.x
617

618 Zöttl, M., Frommen, J.G., Taborsky, M., 2013. Group size adjustment to ecological demand in a
619   cooperative breeder. Proc. R. Soc. B-Biol. Sci. 280, 20122772.
620   https://doi.org/10.1098/rspb.2012.2772
621

Factors	Estimate ± SE	Num. D.F.	Den. D.F.	F-value	*P*-value
(a) Latency to enter the shelter arm					
Intercept	2.92 ± 0.31	-	-	-	-
Treatment	0.12 ± 0.44	1	50	0.08	0.78
(b) Latency to enter the shelter					
Intercept	3.29 ± 0.3	-	-	-	-
Treatment	0.12 ± 0.42	1	50	0.1	0.77

Table 1: Latencies recorded during the post-treatment direct trials. (a) The latency to enter the
shelter arm, and (b) the latency to enter the shelter were not significantly different in glucocorticoid
receptor antagonist treated or control treated fish. The dependent variables were log transformed
to obtain normally distributed residuals. Estimates are shown on a log scale and as differences to the
reference level 'control' for factor 'Treatment'. Body size and sex of test fish (N=52) did not predict
any of the latencies and were subsequently removed from the models. F-values and p-values refer to
comparisons of models with or without the factor of interest.

Factors	Estimate ± SE	X^2-value	P-value
Intercept	1.2 ± 0.47	-	-
Treatment	0.23 ± 0.68	0.12	0.73

Table 2: Probability of a failed attempt during the post-treatment direct trials. The probability of a failed attempt was not significantly different in glucocorticoid receptor antagonist treated or control treated fish. Estimates are shown on a logit scale and as differences to the reference level 'control' for factor 'Treatment'. Body size and sex of test fish (N=52) did not predict the probability of a failed attempt and were subsequently removed from the model. X^2 and p-values refer to comparisons of models with or without the factor of interest.

Factors	Estimate ± SE	X^2-value	*P*-value
Intercept	1.44 ± 0.18	-	-
Treatment	-0.26 ± 0.26	0.65	0.42
Nr. of detour trial	-*	288.56	**<0.01**
Nr. of detour trial x Treatment	-*	23.96	**<0.01**

*Estimates are shown in Table 4

Table 3: Number of failed attempts during the four detour trials. Fish treated with a glucocorticoid
receptor (GR) antagonist showed a significantly different learning curve during the detour trials
compared to control treated fish. (see significant interaction 'Nr. of detour trials x Treatment'). Post-
hoc tests revealed that GR blocked fish showed significantly more errors during detour trial 3 than
control fish (see Table 4 for the pairwise comparisons). Estimates are shown on a log scale and as
differences to the reference level 'control' for factor 'Treatment'. Estimates for the factor 'Nr. of
detour trials' and for the interaction 'Nr. of detour trials x Treatment' are presented as post-hoc
comparisons in Table 4. Body size and sex of test fish did not predict the number of a failed attempts
and were subsequently removed from the model. X^2 and p-values refer to comparisons of models
with or without the factor of interest. N=52 (207 observations in total). P-values < 0.05 are
highlighted in bold.

Contrast	Estimate ± SE	z-value	*P*-value
Detour 1			
Control vs Mifepristone	0.26 ± 0.26	1	0.31
Detour 2			
Control vs. Mifepristone	-0.62 ± 0.37	-1.7	0.09
Detour 3			
Control vs. Mifepristone	-1.15 ± 0.42	-2.72	**0.01**
Detour 4			
Control vs. Mifepristone	0.64 ± 0.41	1.57	0.12

Table 4: Posthoc comparisons with a Tukey correction of the significant interaction term in Table 3.
Fish treated with a glucocorticoid receptor antagonist made significantly more errors during detour
trail 3 leading to a significantly different error pattern during the detour trials (see Figure 2).
Estimates are shown on a log scale, the direction of comparison within a contrast is left to right and
the estimate value refers to the treatment at the right. N=52 (207 observations in total). P-values <
0.05 are highlighted in bold.

Factors	Estimate ± SE	Num. D.F.	Den. D.F.	F-value	*P*-value
(a) Latency to cross the shelter arm					
Intercept	4.92 ± 0.19	-	-	-	-
Treatment	0.01 ± 0.23	1	50.11	0.0	0.97
Nr. of detour trial	-	3	152.25	3.78	**0.01**
Detour 2	-0.49 ± 0.16	-	-	-	-
Detour 3	-0.39 ± 0.16	-	-	-	-
Detour 4	-0.36 ± 0.16	-	-	-	-
(b) Latency to enter the shelter					
Intercept	1.71 ± 1.6	-	-	-	-
Treatment	-0.14 ± 0.22	1	48.98	0.42	0.52
Body size	0.5 ± 0.22	3	48.9	5.1	**0.03**
Nr. of detour trial	-		152.18	0.6	0.61
Detour 2	-0.2 ± 0.2	-	-	-	-
Detour 3	-0.19 ± 0.17	-	-	-	-
Detour 4	-0.18 ± 0.17	-	-	-	-

Table 5: Latencies recorded during the detour trials. (a) The latency to enter the shelter arm, and (b)
the latency to enter the shelter was not significantly different in glucocorticoid receptor antagonist
treated or control treated fish. The dependent variables were log transformed to obtain normally
distributed residuals. Estimates are shown on a log scale and as differences to the reference levels
'control' for factor 'Treatment' and 'Detour 1' for factor 'Nr. of detour trial'. Sex of test fish did not
predict the latencies and was removed from the models, whereas body size was only dropped in (a).
F-values and p-values refer to comparisons of models with or without the factor of interest. N=52
(207 observation in total). P-values < 0.05 are highlighted in bold

Factors	Estimate ± SE	z-value	*P*-value
(a) GR blocked fish			
Intercept	2.07 ± 0.47	-	-
Nr. of predator exposure	-7.69 ± 3.54	-2.17	**0.03**
poly(Nr. of predator exposure, 2)	3.42 ± 3.18	1.08	0.28
(b) Control fish			
Intercept	1.8 ± 0.51		
Nr. of predator exposure	-4.23 ± 3.5	-1.21	0.23
poly(Nr. of predator exposure, 2)	11.87 ± 3.64	3.26	**<0.01**

Table 6: The probability to enter the shelter with increasing numbers of daily predator
presentations. Fish treated with a glucocorticoid receptor (GR) antagonist showed a significantly
different pattern over the course of the experiment than control treated test fish. Separate models
were performed only after confirming that a non-linear interaction term between the number of
daily predator presentations and treatment explained the data better than a linear relationship (see
Results section). (a) GR blocked fish show a linear decrease in the probability to enter the shelter
with increasing daily predator presentations, whereas (b) control fish show a significant U-shape
relationship (see also Figure 3). Estimates are shown on a logit scale. The factor 'Nr. of predator
exposure' was included as a continues variable either as a linear predictor or as a polynomial
predictor (poly) of degree 2. N=26 (a: 129 observations; b: 130 observations in total).

Figure legends

Figure 1: Experimental set-up consisting of one predator tank and a 45L experimental tank. The

predator tank contained a removable opaque divider that could be remotely lifted during the

predator presentations. The experimental tank was divided into five areas that are indicated with

letters (a)-(e): (a) The start area which could be separated by a removable opaque divider from the

rest of the experimental tank. (b) The non-shelter arm which served as the correct entrance during

the detour trials. (c) The middle arm leading to a dead end. (d) The shelter arm that always

contained a half clay flowerpot with the opening either facing towards the start area during the

learning trials or with the opening facing away from the start area during the detour trials (as

depicted here). (e) The back side arm which test fish had to use to enter the shelter arm in the

detour trials. The U-shaped maze consisted of transparent Perspex material. A mesh was glued on

top to make the maze clearly recognizable. Dashed lines indicate virtual lines that were used to

record latencies and failed attempts from the video recordings (see Methods section for more

details). The shortest route towards the shelter was either directly accessible (direct trials) or

blocked (detour trials) with a barrier made from the same material as the maze.

Figure 2: The number of failed attempts in the detour trials. Glucocorticoid receptor (GR) antagonist

treated fish showed a different pattern of failed attempts during the detour trials than control

treated fish. GR blocked fish made more errors particularly during detour trial 3. Individual

datapoints (smaller dots in the background) and predicted values (larger dots in the front) are shown

with 95% confidence intervals for control (red dots) and GR blocked (blue dots) fish.

Figure 3: Probability to enter the shelter with increasing numbers of daily predator presentations for

glucocorticoid receptor (GR) antagonist treated (red dots and line) and control treated (blue dots

and line) fish. GR blocked fish showed a constant decrease in the probability to enter the shelter

with successive predator presentations whereas control fish showed a nonlinear U-shape

relationship. Dots represent the raw data and lines the model prediction with standard errors

depicted as shaded areas.

740   Figure 1

745   Figure 2

751   Figure 3