Human Skeletal Muscle Fiber Heterogeneity Beyond Myosin Heavy Chains

Roger Moreno-Justicia[1]*, Thibaux Van der Stede[2,3]*, Ben Stocks[1]*, Jenni Laitila[4], Robert A Seaborne[4,5], Alexia Van de Loock[2], Eline Lievens[2], Diana Samodova[1], Leyre Marín-Arraiza[6], Oksana Dmytriyeva[1], Kim Van Vossel[2], Nurten Yigit[7], Jasper Anckaert[7], Anneleen Weyns[2], Ruud Van Thienen[2], Ronni E Sahl[4], Edmar Zanoteli[8], Michael W Lawlor[9,10], Michael Wierer[11], Pieter Mestdagh[7], Jo Vandesompele[7], Julien Ochala[4], Morten Hostrup[3], Wim Derave[2#], & Atul S Deshmukh[1#]

*Joint first-authors

[#]Co-corresponding authors (Atul S Deshmukh: atul.deshmukh@sund.ku.dk; Wim Derave: wim.derave@ugent.be)

[1]Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

[2]Department of Movement and Sports Sciences, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

[3]The August Krogh Section for Human Physiology, Department of Nutrition, Exercise and Sports, University of Copenhagen, Copenhagen, Denmark

[4]Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

[5]Centre for Human and Applied Physiological Sciences, King's College London, London, United Kingdom (present address).

[6]Novo Nordisk Foundation Center for Protein Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

[7]Department of Biomolecular Medicine, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium

[8]Department of Neurology, Faculdade de Medicina (FMUSP), Universidade de São Paulo, São Paulo, Brazil

[9]Department of Pathology, Medical College of Wisconsin, Milwaukee, WI, USA

[10]Diverge Translational Science Laboratory, Milwaukee, WI, USA

[11]Proteomics Research Infrastructure, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

Graphical Abstract

Abstract

Skeletal muscle is an inherently heterogenous tissue comprised primarily of myofibers, which

are historically classified into three distinct fiber types in humans: one "slow" (type 1) and two

"fast" (type 2A and type 2X), delineated by the expression of myosin heavy chain isoforms

(MYHs)[1]. However, whether discrete fiber types exist or whether fiber heterogeneity reflects

a continuum remains unclear. Furthermore, whether MYHs are the main classifiers of skeletal

muscle fibers has not been examined in an unbiased manner. Through the development and

application of novel transcriptomic and proteomic workflows, applied to 1050 and 1038 single

muscle fibers from human *vastus lateralis*, respectively, we show that MYHs are not the

principal drivers of skeletal muscle fiber heterogeneity. Instead, ribosomal heterogeneity and

cell junction proteins drive the majority of variance between skeletal muscle fibers in a

continual fashion, independent of slow/fast fiber type. Furthermore, whilst slow and fast fiber

clusters can be identified, described by their contractile and metabolic profiles, our data

challenge the concept that type 2X are phenotypically distinct to other fast fibers at an omics

level. Moreover, MYH-based classifications do not adequately describe the phenotype of

skeletal muscle fibers in one of the most common genetic muscle diseases, nemaline

myopathy, with fibers shifting towards a non-oxidative phenotype independently of MYH-

based fiber type. Instead, we characterize novel transcriptomic and proteomic features of

slow and fast skeletal muscle fibers, including identifying several muscle fiber type-specific

polypeptides, termed microproteins, encoded by transcripts annotated as non-coding RNA.

Overall, our data challenges the currently accepted model of multiple distinct fiber types

defined by the expression of specific MYHs, and proposes new avenues for skeletal muscle

research.

Main

Cellular heterogeneity is an inherent feature of all biological systems, allowing for cellular specialization to meet the diverse demands imposed upon tissues and cells[2]. The classic view of skeletal muscle fiber heterogeneity is that motoneurons define the typology of the fibers within a motor unit, with fiber types being defined by myosin heavy chain isoform (MYH) characteristics[1]. This was first based on its ATPase pH lability[3,4] and later by its molecular MYH expression[5]. Nonetheless, debate continues as to whether discrete fiber types exist (i.e. type 1, type 2a, and type 2x in humans) or whether fiber heterogeneity is better described as a continuum[1]. Furthermore, the current view of MYHs as primary classifiers of muscle fibers might be limited and heavily biased by early studies in rodents, displaying a different MYH expression profile and thus fiber type range than humans[1]. This debate has been further fueled by other limitations such as the identification and subsequent acceptance of the existence of "hybrid" fibers that express varying proportions of multiple MYHs simultaneously.

An unbiased exploration of skeletal muscle fiber diversity with powerful omics tools is thus heavily warranted, but challenging, in part owing to the multinucleated nature of skeletal muscle fibers. Nonetheless, both transcriptomics[6,7] and proteomics[8] technologies have recently experienced a sensitivity revolution due to various technological advances, making profiling of skeletal muscle at a single fiber resolution now possible. As a result, notable progress has already been made in describing the single muscle fiber diversity and responses to atrophic stimuli and aging[9-16]. Importantly, these technological advances lend themselves to being advantageous in the clinical setting, helping to describe in greater detail and precision the dysregulation associated with disease. For example, the underlying pathophysiology of nemaline myopathy, one of the most prevalent genetic muscle disorder (MIM 605355 and MIM 161800), is complex and convoluted[17,18], thus a deeper characterization of skeletal muscle fiber dysregulation may spur significant developments in our understanding of the disease.

Here, our further methodological development of transcriptome and proteome profiling of single skeletal muscle fibers manually isolated from human biopsy specimens, and their application to over a thousand fibers each, enabled us to investigate the true cellular heterogeneity of human skeletal muscle fibers. In doing so, we demonstrate the power of

muscle fiber phenotyping at the transcriptomic and proteomic level, identifying that

ribosomal heterogeneity is one of the primary sources of variation among muscle fibers.

Furthermore, with our proteomics workflow we characterize the clinical implications of

nemaline myopathies within single skeletal muscle fibers, revealing a coordinated shift

towards non-oxidative fibers independently of MYH-based fiber type.

Results

Development of high-sensitivity and high-throughput single muscle fiber transcriptomic and

proteomic pipelines

To investigate the heterogeneity of human skeletal muscle fibers, we developed two novel

workflows to enable transcriptome and proteome profiling of single skeletal muscle fibers (**Fig**

1A and 1 S1A). Several methodological steps were developed and optimized, from sample

storage and preservation of RNA and protein integrity to optimizing the throughput of each

method. This was achieved for transcriptome analysis by inserting sample-specific molecular

barcodes during the initial reverse transcription step, enabling pooling of 96 fibers for further

efficient downstream processing. Rich transcriptome data was further obtained by deeper

sequencing (± 1M reads per fiber) compared to conventional single cell methods[19]. For

proteomics, the combination of a short chromatographic gradient (21 minutes) and timsTOF

DIA-PASEF-based mass spectrometry was used to optimize proteome depth, whilst

maintaining a high-throughput[20,21]. To investigate skeletal muscle fiber heterogeneity in the

healthy state, the transcriptome was determined for 1050 individual fibers from 14 healthy

adult donors, whilst the proteome was determined for 1038 fibers from 5 healthy adult donors

(**Table S1)**. These datasets will be referred to as the 1000 fiber transcriptome and proteome

datasets, respectively, throughout this manuscript. Our approach detected a total of 27237

transcripts and 2983 proteins in the 1000 fiber transcriptomics and proteomics studies (**Fig**

1A, Table S2-3). After filtering for >1000 detected genes and for 50% valid values within each

fiber in both transcriptomics and proteomics datasets, respectively, downstream

bioinformatic analyses were performed on 925 and 974 fibers in the transcriptome and

proteome, respectively. On average 4257 ± 1557 genes and 2015 ± 234 proteins (mean ± SD)

were detected per fiber after filtering, with limited inter-individual variation (**Fig 1 S1B-C,**

Table S4-5). The intra-individual variation within a participant was more substantial however,

most likely due to differences in RNA/protein yield among fibers of different length and cross-

sectional area. For the majority of proteins (>2000), the coefficient of variation was below 20%

(**Fig 1 S1D**). Both methodologies captured a wide dynamic range of transcripts and proteins,

with features known to be important for muscle contraction being highly expressed (e.g.

ACTA1, MYH2, MYH7, TNNT1, TNNT3) (**Fig 1 S1E-F**). A large part of the detected features was

shared between the transcriptome and proteome datasets (**Fig 1 S1G**), alongside a reasonable

correlation (r = 0.52) in average UMI counts/LFQ intensities for these features (**Fig 1 S1H**).

Type 2X is not a distinct fiber type

We initially set out to define the MYH-based fiber type of each fiber using an optimized

methodology, leveraging the high sensitivity and dynamic range of MYH expression in the

omics datasets. Previous studies have used arbitrary cut-offs to assign a fiber as pure type 1,

type 2A, type 2X, or hybrid, based on a fixed percentage of expression for the different

MYHs[9,12,22]. We employed a different approach, in which we ranked the expression of each

fiber by the MYHs used for fiber typing: MYH7, MYH2 and MYH1, corresponding to type 1,

type 2A and type 2X fibers, respectively. We then mathematically calculated the bottom knee

for each of the resulting curves and used it as a threshold to assign a fiber as being positive

(above threshold) or negative (below threshold) for each MYH (**Fig 1B-D**). These data show

that MYH7 (**Fig 1B**) and MYH2 (**Fig 1C**) have a more pronounced on/off expression profile on

the RNA level, compared to the protein level. Indeed, at the protein level, very few fibers did

not express MYH7 and no fibers had 100% MYH2 expression. Next, we used the determined

expression thresholds to assign MYH-based fiber types for all fibers in each dataset. For

example, a MYH7$^+$/MYH2$^-$/MYH1$^-$ was assigned as type 1, and a MYH7$^-$/MYH2$^+$/MYH1$^+$ was

assigned as a hybrid type 2A/2X fiber (see **Table S6** for full description). When combining all

fibers, a very similar MYH-based fiber type distribution was observed at the RNA (**Fig 1E**) and

protein (**Fig 1F**) levels, with an expected inter-individual variation in relative MYH-based fiber

type composition (**Fig 1 S2A**). Most fibers were considered pure type 1 (34-35%) or type 2A

(36-38%), although a substantial number of hybrid 2A/2X fibers (16-19%) were also detected.

A striking discrepancy was that pure type 2X fibers could only be detected at the RNA but not

at the protein level, indicating that fast MYH expression is, at least in part, post-transcriptionally regulated.

Figure 1. Transcriptomic- and proteomic-based clustering reveals only two distinct fiber types. (A) Transcriptomic and proteomic workflow. **(B-D)** Dynamic range curves for MYH7, MYH2, and MYH1, with calculated assignment thresholds. **(E-F)** Distribution of fibers by expression of MYHs in transcriptomics and proteomics datasets. **(G-H)** Uniform manifold approximation and projection (UMAP) plots for transcriptomics and proteomics, colored by MYH-based fiber type. **(I-J)** Feature plots displaying the expression of MYH7, MYH2, and MYH1 in transcriptomics and proteomics.

We validated the MYH-based fiber typing from the proteomics data against an antibody-based

dot blot technique, in which both methodologies were in 100% agreement on the

identification of both pure type 1 and type 2A fibers (**Fig 1 S2B**). However, the increased

sensitivity of the proteomics-based approach was superior in identifying hybrid fibers and in

quantifying the proportion of each MYH within each fiber. These data demonstrate the

efficacy of using an unbiased high-sensitivity omics-based approach for the characterization

of skeletal muscle fiber types.

We then utilized the full depth of information that transcriptomics and proteomics provide to

classify fibers in an unbiased manner based on their whole transcriptome or proteome. Using

uniform manifold approximation and projection (UMAP) for dimension reduction of 6

principal components (**Fig 1 S3 A-B**), we were able to visualize the variation in fibers in the

transcriptome (**Fig 1G**) and proteome (**Fig 1H**). Interestingly, fibers did not cluster by

participant in either the transcriptomics or proteomics datasets (**Fig 1 S3C-D**), nor by test day

(**Fig 1 S3E**), indicating that intra-individual variance in skeletal muscle fibers outweighs inter-

individual variance. Two distinct clusters were apparent in the UMAP plots, which represented

"fast" and "slow" fibers (**Fig 1G-H**). MYH7$^+$ (slow) fibers clustered to the positive side of

UMAP1, and MYH2$^+$ and MYH1$^+$ (fast) fibers clustered to the negative side of UMAP1 (**Fig 1I-**

J). No distinction between the various fast MYH-based fiber types (i.e. type 2A, type 2X, or

hybrid 2A/2X) could be identified, however, suggesting that when the whole transcriptome or

proteome is taken into account, the expression of MYH1 is not discriminative. Given that we

detected no pure 2X fibers at the protein level (**Fig 1F**), these data question the validity of

MYH1 as a marker for fiber typing. Furthermore, whilst type 2X fibers can be identified at the

transcriptional level based solely on the quantification of MYHs, these fibers cannot be

discriminated from other fast fiber types when the whole transcriptome is considered.

Considerable skeletal muscle fiber heterogeneity beyond myosin heavy chain isoforms

As an initial exploration of fiber heterogeneity beyond MYHs, we assessed four established

slow fiber type-specific proteins: TPM3, TNNT1, MYL3, and ATP2A2[1]. In both transcriptomics

(**Fig 2 S1A**) and proteomics (**Fig 2 S1B**) approaches, the slow isoforms exhibited a high,

although not perfect, Pearson correlation coefficient with MYH7. Approximately 25 and 33%

of the fibers in the transcriptomics (**Fig 2 S1C**) and proteomics (**Fig 2 S1D**) approaches,

respectively, were not classified as pure slow fibers by all gene/protein isoforms. Thus, fiber

typing based on multiple gene/protein isoforms introduces additional complexity, even with

well-known proteins that are assumed to be fiber type-specific. This suggests that fiber typing

based on isoforms of a single family of genes/proteins is likely inadequate to capture the true

heterogeneity of skeletal muscle fibers.

To further investigate the omics-wide phenotypical variability between human skeletal muscle

fibers, we applied unbiased dimensionality reduction by principal component analysis (PCA)

to our data (**Fig 2A**). Similarly to the UMAP plot, neither participant nor test day influenced

the clustering of fibers at the PCA level (**Fig 2 S2A-C**). MYH-based fiber type was explained by

PC2 in both datasets, displaying clusters for slow type 1 fibers and a second cluster containing

the fast type 2A, type 2X and hybrid type 2A/2X fibers (**Fig 2A**). These two clusters were

bridged in both datasets by a small number of hybrid type 1/2A fibers. As expected, over-

representation analysis of the top PC drivers confirmed that PC2 is driven by contractile and

metabolic features (**Fig 2B & 2 S2D-E, Table S7-9**). In general, the MYH-based fiber types

adequately explain the continual variance along PC2, except for the so-called type 2X fibers,

which were spread across the entirety of the transcriptomic fast cluster.

Unexpectedly, MYH-based fiber type was only the second most important source of variability

(PC2), indicating that other biological factors (PC1) independent of MYH-based fiber type,

have an even greater role in regulating skeletal muscle fiber heterogeneity. Over-

representation analysis of the top drivers in PC1 indicated that variance within PC1 was

determined primarily by cell-cell adhesion and ribosomal content in the transcriptome, and

costamere and ribosomal proteins in the proteome (**Fig 2B & 2 S2D-E, Table S9**). In skeletal

muscle, the costamere connects the Z-disk to the sarcolemma and participates in force

transmission and signaling[23]. Annotating PCA plots with cell-cell adhesion (transcriptome, **Fig**

2C) and costamere (proteome, **Fig 2D**) features showed a strong shift to the left side of PC1,

suggesting an enrichment for these features in some fibers.

Figure 2. Skeletal muscle fiber heterogeneity beyond MYHs. (A) Principal component analysis (PCA) plots of transcriptomics and proteomics datasets colored by MYH-based fiber type. **(B)** Enrichment analysis of transcript and protein drivers of PC2 and PC1. **(C-D)** PCA plot colored by cell-cell adhesion gene ontology (GO) term in the transcriptome, and by costamere GO term in the proteome. Arrows represent transcript and protein drivers and their direction. **(E-F)** Uniform manifold approximation and projection (UMAP) feature plots of clinically relevant features displaying a slow/fast fiber type-independent gradient of expression. **(G-H)** Correlation between drivers of PC2 and PC1 between the transcriptome and proteome.

Closer inspection of fiber clustering at the UMAP level showed a MYH-based fiber type-independent gradient of expression for most features, rather than distinct subclusters of muscle fibers. This continuum holds true for several genes related to pathological conditions (**Fig 2E**), such as *CHCHD10* (neuromuscular disorders)[24], *SLIT3* (muscle loss)[25] and *CTDNEP1* (muscle disease)[26]. This same continuum was observed in the proteome, with proteins related to neurological diseases (UGDH)[27], insulin signaling (PHIP)[28] and transcription (HIST1H2AB) (**Fig 2F**). These data collectively show that slow/fast fiber type independent heterogeneity across fibers is of a continual nature, and no distinct subtypes of human skeletal muscle fibers beyond a slow-fast classification should be considered.

Fiber heterogeneity is post-transcriptionally regulated by ribosomal heterogeneity

Interestingly, drivers of PC2 showed a good correlation (r = 0.663) between the transcriptome and proteome (**Fig 2G**), indicating that the slow and fast fiber types, particularly the contractile and the metabolic profile of skeletal muscle fibers, are transcriptionally regulated. However, there was no correlation between the drivers of PC1 in the transcriptome and the proteome (r = -0.027) (**Fig 2H**). This suggests that slow/fast fiber type-independent variance is for a large part post-transcriptionally regulated. Since PC1 variance was largely explained by ribosomal gene ontology terms and given that ribosomes play a profound and specialized role in the cell by actively participating and influencing protein translation[29], we next set out to explore this unexpected ribosomal heterogeneity.

We first colored the proteomics PCA plot based on the relative abundance of the proteins within the "cytosolic ribosome" GOCC term (**Fig 3A**). Although the term was enriched on the positive side of PC1, and a slight gradient could be observed accordingly, ribosomal proteins were driving separation in both directions of PC1 (**Fig 3A**). Amongst the ribosomal proteins enriched in the negative direction of PC1 were RPL18, RPS18 and RPS13 (**Fig 3B**), whilst RPL31, RPL35 and RPL38 (**Fig 3C**) were major drivers in the positive direction of PC1. Interestingly, RPL38 and RPS13 are highly expressed in skeletal muscle when compared to other tissues (**Fig 3 S1A**). These distinct ribosomal signatures across PC1 could not be observed in the transcriptome (**Fig 3 S1B**), indicating that this phenomenon is post-transcriptionally regulated.

Figure 3. Ribosomal heterogeneity drives fiber type-independent heterogeneity. (A) Principal component analysis (PCA) plot colored by cytosolic ribosome gene ontology (GO) term in the proteome. Arrows represent protein drivers and their direction. **(B-C)** PCA feature plots of RPS13 and RPL38. **(D)** Unsupervised hierarchical clustering analysis of proteins constituting the cytosolic ribosome. **(E)** Structural model of the 80S ribosome (PDB: 4V6X) highlighting ribosomal proteins displaying variable abundance across skeletal muscle fibers. **(F)** Ribosomal proteins with variable stoichiometry located close to the mRNA exit tunnel.

The concept of ribosomal heterogeneity and specialization was previously introduced, where the presence of distinct subpopulations of ribosomes (ribosomal heterogeneity) can directly influence protein translation in different tissues[30] and cells[31] by selectively translating specific mRNA transcript pools[32] (ribosomal specialization). To identify sub-sets of ribosomal proteins that are co-expressed within skeletal muscle fibers, we performed an unsupervised hierarchical clustering analysis of ribosomal proteins within the proteome (**Fig 3D, Table S10**). As expected, ribosomal proteins did not cluster by MYH-based fiber type. However, we identified three distinct clusters of ribosomal proteins; the first of which (ribosomal_cluster_1)

was coregulated alongside RPL38, therefore elevated in fibers in the positive direction of PC1.

The second cluster (ribosomal_cluster_2) was coregulated alongside RPS13 and was elevated

in fibers in the negative direction of PC1. A third cluster (ribosomal_cluster_3) displayed no

coordinated differential expression within skeletal muscle fibers and could therefore be

considered "core" ribosomal proteins within skeletal muscle. Both ribosomal_cluster_1 and

ribosomal_cluster_2 contain ribosomal proteins previously demonstrated to regulate

selective translation (e.g. RPL10A, RPL38, RPS19 and RPS25) and functionally influence

development (e.g. RPL10A, RPL38)[32-36].

To visualize the position of the ribosomal proteins that display heterogeneity within the

ribosome, we utilized a structural model of the human 80S ribosome (Protein Data Bank:

4V6X)[37] (**Fig 3E**). Upon highlighting the ribosomal proteins that belong to the different

ribosomal clusters, they were not located in close proximity, showing that our approach did

not enrich for a particular area/section of the ribosome. Yet interestingly, ribosomal_cluster_2

contained a lower proportion of large ribosomal subunit proteins than ribosomal_cluster_1

and ribosomal_cluster_3 (**Fig 3 S1C**). We observed that the majority of the proteins that

display variable stoichiometry within skeletal muscle fibers are located on the surface of the

ribosome (**Fig 3E**), which is consistent with an ability to interact with internal ribosome entry

site (IRES) elements within distinct mRNA populations to coordinate selective translation[38,39].

Furthermore, numerous proteins that display variable stoichiometry within skeletal muscle

fibers are located close to functional regions, such as the mRNA exit tunnel (**Fig 3F**), which can

selectively regulate translation elongation and arrest of specific nascent peptides[40]. Overall,

our data identifies that skeletal muscle ribosomal protein stoichiometry displays

heterogeneity, driving variance between skeletal muscle fibers.

Slow and fast fiber signatures and their transcriptional regulators

We next set out to identify novel features of fast and slow skeletal muscle fibers and how

these are transcriptionally regulated. Comparing the fast and slow clusters defined in the

UMAPs of both datasets (**Fig 1G-H & inlays Fig 4A-B**), transcriptome and proteome analysis

yielded 1366 and 804 differentially abundant features, respectively (**Fig 4A-B, Table S11-14**).

Expected differences in sarcomeric (e.g. tropomyosin and troponin), excitation-contraction

coupling (SERCA isoforms) and energy metabolism-related (e.g. ALDOA and CKB) features were observed. In addition, transcripts and proteins regulating protein ubiquitination displayed differences between fast and slow fibers (e.g. *USP54, SH3RF2,* USP28 and USP48) (**Fig 4A-B**). Furthermore, the microprotein-encoding gene *RP11-451G4.2*, which has previously been shown to be differentially expressed between lamb muscle fiber types[41], was significantly up-regulated in slow skeletal muscle fibers (**Fig 4A**). Also at the single fiber level, clear differences could be observed for known features such as the metabolism-related isoforms of lactate dehydrogenase (LDHA and LDHB, **Fig 4C** and **Fig 4 S1A**)[42,43], as well as previously unknown fiber type-specific features (e.g. *IRX3*, *USP54*, USP28, and DPYSL3) (**Fig 4C**). There was a reasonable overlap of differentially expressed features (**Fig 4 S1B**) and correlation of fold changes between the transcriptomic and proteomic datasets, primarily driven by the large expression differences in sarcomeric features (**Fig 4 S1C**). Notably, some features (e.g. USP28, USP48, GOLGA4, AKAP13) showed strong post-transcriptional regulation with slow/fast fiber type-specific expression profiles only at the proteome level (**Fig 4 S1B**).

We next performed over-representation analysis on the differentially abundant genes and proteins (**Fig 4D, Table S15**). Enriched pathways of features that were differential in both datasets showed expected differences such as fatty acid beta-oxidation and ketone metabolic process (slow fibers), myofilament/muscle contraction (fast and slow fibers, respectively) and carbohydrate catabolic process (fast fibers). Serine/Threonine protein phosphatase activity was also enriched in fast fibers, driven by features such as phosphatase regulatory and catalytic subunits (PPP3CB, PPP1R3D, and PPP1R3A), known to regulate glycogen metabolism[44] (**Fig 4 S1D-E**). Other enriched pathways in fast fibers were Processing (P-) bodies (YTHDF3, TRIM21, LSM2) in the proteome (**Fig 4 S1F**), possibly related to post-transcriptional regulation[45], and transcription factor activity (*SREBPF1*, *RXRG*, *RORA*) in the transcriptome (**Fig 4 S1G**). Slow fibers showed enrichment for oxidoreductase activity (*BDH1*, *DCXR*, *TXN2*) (**Fig 4 S1H**), amide binding (*CPTP*, *PFDN2*, *CRYAB*) (**Fig 4 S1I**), extracellular matrix (*CTSD*, *ADAMTSL4*, *LAMC1*) (**Fig 4 S1J**) and receptor-ligand activity (*FNDC5*, *SPX*, *NENF*) (**Fig 4 S1K**).

Figure 4. Fast and slow skeletal muscle fiber signatures. (A-B) Volcano plots comparing the slow and fast clusters identified from the uniform manifold approximation and projection (UMAP) plots in Fig 1G-H. Colored dots represent significantly different transcripts or proteins at an FDR < 0.05, darker dots represent significantly different transcripts or proteins with a fold change > 2. **(C)** Feature plots of selected differentially expressed genes or proteins between slow and fast fibers. **(D)** Enrichment analysis of significantly different transcript and proteins. Overlap – enriched in both datasets, Transcriptome – enriched only in transcriptome, Proteome – enriched only in proteome. **(E)** Transcription factor motif enrichment analysis from the transcriptome using the TRANSFAC database. **(F)** Significantly regulated non-coding RNA transcripts. **(G)** Representative RNAscope images displaying

the slow/fast fiber-type specificity of LINC01405 and RP11-255P5.3, respectively. Scaling bar = 50 µm. **(H)** De novo microprotein identification workflow. **(I)** The microprotein LINC01405_ORF408:17441:17358 is specific to slow skeletal muscle fibers.

To get more insights into the transcriptional regulation of slow/fast fiber type signatures, we performed transcription factor binding motif enrichment using the TRANSFAC databases on the differentially expressed genes in the transcriptome dataset (**Table S16**). Many transcription factor binding motifs were significantly enriched between fast and slow fibers (**Fig 4E**). This included known transcription factors such as MEF2[46], enriched in the fast fibers, but also multiple novel transcription factors potentially involved in the fiber type specific gene program. TIEG1, known to regulate mitochondrial content[47], was the most enriched transcription factor within slow fibers and as such should be considered as an important regulator of skeletal muscle metabolism. In addition, PATZ, BTEB3 and multiple motifs from the SP family of transcription factors were enriched in slow muscle fibers. In fast skeletal muscle fibers, TFAP2A, ZXDL, FPM315, AP-2 and ETF are all novel transcription factors putatively regulating skeletal muscle fiber type.

Alongside known protein-coding genes, there exists a multitude of non-coding RNA biotypes, potentially involved in the regulation of human development and disease[48,49]. Several non-coding RNAs displayed fiber type specificity in the transcriptomics dataset (**Fig 4F & Table S17**), including *LINC01405*, which is very specific to slow fibers and is reported to be downregulated in muscle of mitochondrial myopathy patients[50]. Conversely, *RP11-255P5.3,* corresponding to the *lnc-ERCC5-5* gene (https://lncipedia.org/db/transcript/lnc-ERCC5-5:2)[51] displayed fast fiber type-specificity. Both *LINC01405* (https://tinyurl.com/x5k9wj3h) and *RP11-255P5.3* (https://tinyurl.com/29jmzder) display specificity to skeletal muscle (**Fig 4 S2A-B**) and have very few or no known contractile genes within a 1Mb genomic neighborhood, suggesting a specialized role in fiber type regulation rather than a regulatory role for a neighboring contractile gene. The slow/fast fiber type-specific specific expression profiles of *LINC01405* and *RP11-255P5.3* were independently validated using RNAscope (**Fig 4G**).

Recently, it has become apparent that numerous assumed non-coding transcripts code for transcribed microproteins, some of which regulate muscle function[52,53]. To identify novel

microproteins with potential fiber-type specificity, we searched our 1000 fiber proteomic dataset using a custom FASTA file containing sequences from the detected non-coding transcripts (n = 305) from the 1000 fiber transcriptome dataset (**Fig 4H**). This resulted in the identification of 197 novel microproteins arising from 22 distinct transcripts (**Table S18**, of which 71 microproteins display differential regulation between slow and fast skeletal muscle fibers (**Fig 4 S2C and Table S19**). Three microprotein products were identified for *LINC01405*, of which one displays a slow-fiber type specificity similar to its transcript (**Fig 4I and 4 S2D**). Thus, we identify *LINC01405* as a microprotein-encoding gene, which displays specificity for slow skeletal muscle fibers.

Nemaline myopathies induce a shift towards faster, less oxidative skeletal muscle fibers

Having developed a comprehensive workflow to characterize the proteome of single muscle fibers at scale and discovered novel regulators of fiber heterogeneity in the healthy state, we applied this pipeline to understand how nemaline myopathy impacts skeletal muscle fiber heterogeneity. Nemaline myopathy is a genetic muscular disorder that causes muscle weakness, resulting in a range of complications for the affected children including respiratory difficulties, scoliosis, and physical immobility[17,18]. Typically, in nemaline myopathy, pathogenic variants in genes such as actin alpha 1 (*ACTA1*) drive the fiber type composition towards slow fiber predominance, although this effect is heterogenous. The only clear exception is the troponin T1 (*TNNT1*) nemaline myopathy, in which a fast fiber predominance is seen. Thus, a deeper characterization of the heterogeneity behind the skeletal muscle fiber dysregulation observed in nemaline myopathies may help untangle the complex relationship between these diseases and muscle fiber types.

Muscle fibers isolated from patients with *ACTA1*- and *TNNT1*-mutation derived nemaline myopathies display substantial myofiber atrophy or hypotrophy, compared to healthy controls (n = 3 per group) (**Fig 5A, Table S20**), which provides considerable technical challenge arising from the limited material for proteomic analysis. Nonetheless, we were able to detect 2485 proteins from 272 skeletal muscle fibers. After filtering for a minimum of 1000 quantified proteins per fiber, downstream bioinformatic analyses were performed on 250 fibers. On average of 1573 ± 359 proteins were quantified per fiber after filtering (**Fig 5 S1A, Table S21-**

22). Importantly, only a modest reduction in proteome depth was apparent in samples from patients with nemaline myopathy, despite the markedly reduced fiber size. A wide dynamic range was apparent in the proteome, whilst the shared proteins in the control participants correlated well with those in the previous analysis of the 1000 fiber proteome study (**Fig 5 S1B-C**).

As nemaline myopathies influence the MYH-based fiber type proportions within skeletal muscle[17,18], we first investigated the MYH-based fiber type of our nemaline myopathy patients and controls. Fiber type was determined using the unbiased approach described previously for the 1000 fiber studies (**Fig 5 S1D-E**) and once again pure type 2X fibers could not be identified (**Fig 5B**). We observed a heterogeneous effect of nemaline myopathies on fiber type two patients with *ACTA1* mutations displaying an increased proportion of type 1 fibers, whilst two patients with *TNNT1*-nemaline myopathy displayed a reduced proportion of type 1 fibers (**Fig 5B**). Indeed, MYH2 and the fast troponin isoforms were downregulated in *ACTA1*-nemaline myopathy (TNNC2, TNNI2, and TNNT3), whilst MYH7 was downregulated in *TNNT1*-neamline myopathy (**Fig 5 S2A**). This is in line with previous reports of heterogeneous fiber type switching in nemaline myopathies[17,18]. We validated these findings using immunohistochemistry, finding type 1 fiber predominance in *ACTA1*-nemaline myopathy patients, while the opposite was observed in *TNNT1-nemaline myopathy* patients (**Fig 5A**).

At the whole proteome level, skeletal muscle fibers from *ACTA1*- and *TNNT1*-nemaline myopathy patients clustered away from the majority of control fibers, with TNNT1- nemaline myopathy fibers tending to be the most severely affected (**Fig 5C**). This was particularly apparent when we produced a PCA plot of pseudo-bulked fibers for each patient, with *TNNT1*-nemaline myopathy patients 2 & 3 lying furthest from the control samples (**Fig 5 S2B, Table S23**). To further understand how fibers from myopathy patients compare to the healthy condition, we capitalized on the depth of information provided by our analysis in 1000 fiber proteome study from healthy adult participants. We projected the fibers from our myopathy data set (both *ACTA1*- and *TNNT1*-nemaline myopathy patients and controls) onto the PCA determined from the 1000 fiber proteome study (**Fig 5D**). Control fibers displayed a similar distribution of MYH-based fiber type along PC2 as the 1000 fiber proteome study. However, the majority of fibers from nemaline myopathy patients shifted downwards along PC2,

445 overlapping with the healthy fast fibers, irrespective of their own MYH-based fiber type. Thus,

446 despite evidence for a fiber-type shift towards type 1 fibers in *ACTA1*-nemaline myopathy

447 patients when quantified using MYH-based approaches, both *ACTA1*- and *TNNT1*-nemaline

448 myopathies shift the skeletal muscle fiber proteome to resemble fast fibers.

 Figure 5. Nemaline myopathies induce a shift towards faster, less oxidative skeletal muscle fibers. (A)
Microscopy images displaying fiber atrophy or hypotrophy and divergent MYH-based fiber type predominance
in *ACTA1-* and *TNNT1*-nemaline myopathies (NM). **(B)** MYH-based fiber type proportions by participant. Scaling
bar = 100 μm. **(C)** Principal component analysis (PCA) plot of skeletal muscle fibers from nemaline myopathy
patients and controls. **(D)** Skeletal muscle fibers from nemaline myopathy patients and controls projected onto
PCA plot determined from the 1000 fibers analysed in Fig 2. **(E-G)** Volcano plots comparing the *ACTA1-* and
TNNT1-nemaline myopathies with control participants and between *ACTA1-* and *TNNT1*-nemaline myopathies.
Colored circles represent significantly different proteins at $\pi < 0.05$, darker dots represent significantly different
proteins at FDR < 0.05. **(H)** Enrichment analysis of significantly different proteins in the whole proteome and
within type 1 and type 2A fibers. **(I-J)** Principal component analysis (PCA) plot colored by extracellular matrix and
mitochondria gene ontology (GO) terms.

We next directly compared each patient group with the samples from healthy controls,
resulting in 256 and 552 differentially expressed proteins in *ACTA1-* and *TNNT1*-nemaline
myopathies, respectively (**Fig 5E-F & Fig 5 SC, Table S24**). Gene set enrichment analysis
identified the coordinated reduction of mitochondrial proteins (**Fig 5H & I, Table S25**).
Surprisingly, this was completely independent of MYH-based fiber type (**Fig 5H & Fig 5 S2D-G,**
Table S26), despite the divergent fiber type predominance between *ACTA1-* and *TNNT1-*
nemaline myopathies. Conversely, ribosomal proteins were largely unaffected by nemaline
myopathies, although RPS17 was downregulated in *ACTA1* nemaline myopathy (**Fig 5E**).

Enrichment analysis also identified the up-regulation of immune system process in both
ACTA1- and *TNNT1*-nemaline myopathies, as well as cell adhesion in *TNNT1*-nemaline
myopathy (**Fig 5H**). The enrichment of these extracellular terms, was reflected by extracellular
matrix proteins driving the PCA in the negative direction in both PC1 and PC2, i.e. towards the
most affected fibers (**Fig 5I**). Both patient groups overexpressed extracellular proteins involved
in immune response and the sarcolemma repair machinery, such as annexins (ANXA1, ANXA2,
ANXA5)[54,55] and their interactor S100A11[56] (**Fig 5 S2J-K**). This process has been previously
reported to be up-regulated in muscle dystrophies[57], yet to our knowledge it has not been
associated with nemaline myopathies before. A normal function of this process is required for
both sarcolemma membrane repair upon injury and for the fusion of new myoblasts to muscle
fibers[55,58]. Therefore, an up-regulation of this process in both patient groups suggests a
reparative response to damage caused by myofibril instability.

The effects of each nemaline myopathy were well correlated (r = 0.736) and displayed reasonable overlap (**Fig 5 S2A & B**), indicating that *ACTA1*- and *TNNT1*-nemaline myopathies induce similar effects on the proteome. Nonetheless, a number of proteins displayed regulation in only *ACTA1*- or *TNNT1*-nemaline myopathies (**Fig 5 S2A & S2C**). MFAP4, a pro-fibrotic protein, was one of the most up-regulated proteins in *TNNT1*-nemaline myopathy whilst unchanged in *ACTA1*-nemaline myopathy. SKIC8, a component of the PAF1C complex, which regulates the transcription of *HOX* genes[59], was down-regulated in *TNNT1*-nemaline myopathy but was unaffected in *ACTA1*-nemaline myopathy (**Fig 5 S2A**). Direct comparisons between *ACTA1*- and *TNNT1*-nemaline myopathies identify a greater effect of *TNNT1*-nemaline myopathies on the reduction of mitochondrial proteins and the increase of immune system proteins (**Fig 5G-H & Fig 5 S2C & S2H-I**). These data are consistent with the greater degree of atrophy/hypotrophy apparent in *TNNT1*- compared to *ACTA1*-nemaline myopathies (**Fig 5A**), indicating that *TNNT1*-nemaline myopathy is the more severe form of the disease.

These data demonstrate that single muscle fiber proteomics can illuminate clinical biology that is unapparent using targeted approaches such as immunoblotting. Furthermore, these data highlight the limitations of solely relying upon MYH-based fiber typing to describe phenotypic adaptations. Indeed, despite divergent MYH-based fiber type switching in actin and troponin nemaline myopathies, both nemaline myopathies uncouple the MYH-based fiber type from skeletal muscle fiber metabolism, shifting towards a faster, less oxidative muscle proteome.

Discussion

Cellular heterogeneity is important to enable tissues to meet a wide range of demands. In skeletal muscle, this has classically been described by fiber types characterized by varying degrees of force production and fatigability. It is apparent, however, that this explains just a fraction of the variation in skeletal muscle fibers, with far greater variability, complexity and nuance residing in skeletal muscle fibers than previously believed. With the advent and development in technologies, these elements that regulate skeletal muscle fibers are now able to be elucidated. Indeed, our data herein, questions the existence of the classical type 2X fiber type as a distinct sub-classification, whilst also identifying determinants of fiber type-

independent heterogeneity, which is predominantly regulated post-transcriptionally via ribosomal heterogeneity. By applying our proteomics workflow to samples from nemaline myopathy patients, we further evidence that MYH-based fiber typing does not capture the true heterogeneity of skeletal muscle, particularly when the system is perturbed. Indeed, nemaline myopathies induce a shift towards faster less oxidative fibers regardless of their MYH-based fiber type.

Skeletal muscle fibers have been classified since the 19[th] century[1,60]. However, recent omics-based analyses have enabled us to start understanding expression profiles and responses to various stimuli specific to distinct MYH-based fiber types[9-14,22,43,61]. We leveraged complimentary transcriptomics and proteomics workflows, and integrated the results, to explore the transcriptional and post-transcriptional regulation of human skeletal muscle fiber heterogeneity. This pipeline resulted in the observation that the existence of pure type 2X fibers should be questioned, especially at the protein level. Indeed, clustering-based analyses could only confidently identify two distinct clusters, which represented slow and fast skeletal muscle fibers regardless of their exact MYH composition. These findings indicate that we should revert to the view of muscle fiber classification from the early twentieth century and classify human skeletal muscle fibers not into three distinct classifications based on MYHs, but instead into just two clusters based on their metabolic and contractile properties[60].

Better still, we should consider muscle fiber heterogeneity in multiple dimensions. Over and above the variance within the contractile and metabolic signatures of skeletal muscle, we found that fibers could also be separated by features related to the cell junction and translation machinery. Indeed, we identified ribosomal heterogeneity across skeletal muscle fibers that drives a slow/fast-fiber type independent heterogeneity. Numerous ribosomal proteins displaying variable abundance across fibers regulate selective translation (e.g. RPL10A, RPL38, RPS19 and RPS25)[32-36]. RPL10A is required for efficient nuclear export of the nascent pre-60s complex and therefore ribosomal biogenesis[62]. RPL10A also directly binds internal ribosome entry site (IRES) elements to selectively regulate translation[32,63], whilst depletion of RPL10A in murine embryos is associated with increased translation of mRNAs associated with striated muscle contraction[34]. In RPL38 mutant embryos the translation of *HOX* transcripts is impaired[33], which are involved in cell development, tissue-specific

patterning and skeletal muscle plasticity[33,64]. Functionally, RPL10A and RPL38 mutant mice

exhibit impaired embryonic development[33,34], demonstrating the physiological relevance of

ribosomal heterogeneity.

The underlying reason for such vast slow/fast-fiber type independent heterogeneity amongst

fibers is not immediately obvious, though this could allude to a specialized spatial organization

within a muscle fascicle to enable an optimal response to specific forces and loads[65], or

specialized cellular or organ communication with other cell types within the muscle

microenvironment[66-68]. Indeed, ribosomal heterogeneity (via paralog substitution of RPL3 and

RPL3L) has been implicated in skeletal muscle hypertrophy[69]. Multi-omics and spatial

applications in concert with functional characteristics of single muscle fibers will further

advance our understanding of muscle biology on a multi-omics level[70]. Indeed, as we and

others continue to optimize single-cell and single muscle fiber omics-analyses towards ultra-

low sample input (as demonstrated here with the analysis of fibers from nemaline myopathy

patients), the possibility of combining functional and (multi-)omics approaches in a single

muscle fiber is tantalizingly close.

In analyzing the proteome of single muscle fibers from patients with nemaline myopathies,

we also demonstrate the utility, power, and applicability of single muscle fiber proteomics in

uncovering clinical pathophysiology in skeletal muscle. In addition to observing expected,

albeit divergent, differences in fiber-type proportions in *ACTA1*- and *TNNT1*-nemaline

myopathies compared to healthy controls[17], we also identified oxidative and extracellular

remodeling, which is uncoupled from this MYH-based fiber type switch. Fibrosis has

previously been reported for *TNNT1* nemaline myopathy[17]. However, our analysis builds upon

this to also identify the up-regulation of stress-related secreted proteins within the

extracellular space, such as annexins, involved in the sarcolemma repair machinery[54-56] in

fibers from both *ACTA1*- and *TNNT1*-nemaline myopathy patients. Overall, the up-regulation

of annexins in muscle fibers from nemaline myopathy patients may represent a cellular

response to rescue severely atrophying fibers.

Collectively, our data identifies and explains transcriptional and post-transcriptional drivers of

heterogeneity within skeletal muscle. In particular, we provide new knowledge of ribosomal

heterogeneity in skeletal muscle and how this drives cellular heterogeneity, as well providing

data to question long-standing dogmas within skeletal muscle physiology related to the

classical definitions of MYH-based fiber-types. In doing so we hope to reignite the debate and

ultimately redefine our understanding of skeletal muscle fiber classifications and

heterogeneity.

Methods

1000 fiber transcriptomics

Participant information. Fourteen participants (12 males / 2 females) volunteered to take part

in this study, which was approved by the Ethical Committee of Ghent University Hospital (BC-

10237), in agreement with the 2013 Declaration of Helsinki and registered on ClinicalTrials.gov

(NCT05131555). General characteristics of the participants can be found in **Table S1**. After oral

and written informed consent, participants were medically screened before final inclusion.

Participants were young (22-42 years old), healthy (no diseases and non-smoking) and

moderately physically active. Maximal oxygen uptake was determined as marker of physical

fitness during a graded incremental cycling test, as previously described[71].

Muscle biopsy collection. Muscle biopsies were collected in the rested and fasted state, on

three different days, separated by 14 days. As these samples were collected as part of a larger

study, on each of these days participants ingested a placebo (lactose), an H1-receptor

antagonist (540 mg fexofenadine) or an H2-receptor antagonist (40 mg famotidine) 40

minutes before muscle biopsy collection. We have previously shown that these histamine

receptor antagonists do not affect the resting skeletal muscle state[71], and no clustering was

apparent based on condition in our quality control plots (**Fig 1 S3E & Fig 2 S2B**). Dietary intake

was standardized 48 hours before each experimental day (41.4 kcal/kg body weight, 5.1 g/kg

body weight carbohydrates, 1.4 g/kg body weight protein and 1.6 g/kg body weight fat per

day), followed by a standardized breakfast on the morning of the experimental day (1.5 g/kg

bodyweight carbohydrates). Muscle biopsies of the *m. vastus lateralis* were then collected

after local anaesthesia (0.5 mL of 1% Xylocaïne without epinephrine) using the percutaneous

Bergström technique with suction[72]. The muscle samples were immediately submerged in

RNA*later* and stored at 4 °C until manual fiber dissection (max. 3 days).

Single fiber isolation. Freshly excised muscle fiber bundles were transferred to fresh RNA*later* in a petri dish. Individual muscle fibers were then manually dissected using a stereomicroscope and fine forceps. Twenty-five fibers were dissected per biopsy, with special care to select fibers from different sections of the biopsy. After dissection, each fiber was carefully submerged in 3 μL of lysis buffer (SingleShot Cell Lysis kit, Bio-rad), containing proteinase K and DNase enzymes to remove unwanted proteins and DNA. Next, cell lysis and protein/DNA removal was initiated by short vortexing, spinning liquid down in a microcentrifuge and incubation at room temperature (10 min). Lysates were then incubated for 5 min at 37 °C and 5 min at 75 °C in a thermocycler (T100, Bio-Rad), immediately followed by storage at -80 °C until further processing.

Sequencing library preparation. Illumina-compatible libraries from polyadenylated RNA were prepared from 2 μL of the muscle fiber lysates using the QuantSeq-Pool 3' mRNA-Seq library prep kit (Lexogen). Detailed methodology can be found in the manufacturer guidelines. The process was initiated by reverse transcription for first strand cDNA synthesis, during which Unique Molecular Identifiers (UMIs) and sample-specific i1 barcodes were introduced enabling sample pooling and reducing technical variability in the downstream process. Next, cDNA from 96 fibers was pooled and purified using magnetic beads, followed by RNA removal and second strand synthesis by random priming. Libraries were purified using magnetic beads, followed by addition of pool-specific i5/i7 indices and PCR amplification. A last purification step was performed, finalizing the Illumina-compatible libraries. A high sensitivity small DNA Fragment Analysis kit (Agilent Technologies, DNF-477-0500) was used to assess the quality of each library pool.

Illumina sequencing. The individual pools were further equimolarly (2 nM) pooled, based on Qubit-quantified concentrations. The final pool was subsequently sequenced with a NovaSeq S2 kit (1 x 100 nucleotides) with a loading of 2 nM (4% PhiX) in standard mode on a NovaSeq 6000 instrument.

Primary data processing. Our pipeline was based on the QuantSeq Pool data analysis pipeline from Lexogen (https://github.com/Lexogen-Tools/quantseqpool_analysis). First, the data was demultiplexed based on the i7/i5 indices with bcl2fastq2 (v2.20.0). The next demultiplexing

step was performed via idemux (v0.1.6) according to the i1 sample-specific barcodes in read

2, followed by extraction of UMI sequences with umi_tools (v1.0.1). Trimming of the reads

was then performed in multiple rounds with cutadapt (v3.4), with removal of too short reads

(length < 20) or reads consisting entirely of adapter sequences. Reads were then aligned to

the human genome with STAR (v2.6.0c), followed by BAM file indexing with SAMtools (v1.11).

Duplicates of reads were removed with umi_tools (v1.0.1). Finally, counting of the alignments

was performed with featureCounts from Subread (v2.0.3). At several intermediate steps

during the pipeline, quality control with performed with FastQC (v0.11.9).

Initial Seurat processing. All further bioinformatics processing and visualization was

performed in R (v4.2.3), primarily with the Seurat v4 workflow[73]. The individual UMI counts

and metadata matrices were thus transformed into a Seurat object. Genes with expression in

less than 30% of all fibers were removed. Low-quality samples were then removed based on

a minimum threshold of 1000 UMI counts and 1000 detected genes. This resulted in a total of

925 fibers that passed all quality control filtering steps. Normalization of UMI counts was

performed using the SCTransform v2 Seurat method[74], including all 7418 detected features

and regressing out participant variation. All relevant metadata can be found in **Table S27**.

Proteomics

Sample collection - 1000 fiber proteome

Participant information. Stored biobank muscle specimens were used for the purpose of the

present study (Clinicaltrials.gov identifier: NCT04048993). The specimens were collected from

five active and healthy male volunteers (aged 21-35 years) who gave their written and oral

informed consent with approval from the Science Ethics Committee of the Capital Region in

Denmark (H-1-2012-090) and complied with the guidelines of the 2013 Declaration of

Helsinki. General characteristics of the participants can be found in **Table S1.** Participants were

young, healthy (no diseases and non-smoking) and moderately physically active.

Muscle biopsy collection. Participants arrived in the morning after an overnight fast and

rested in the supine position for 1 hour. Then, local anesthesia (2-3 mL Xylocaine 2%; lidocaine

without epinephrine, AstraZeneca, Denmark) was applied under the skin above the fascia at

the belly of the *m. vastus lateralis* muscle. A muscle biopsy was sampled through a small 3-4

mm incision using a Bergström needle with suction. The muscle biopsy specimen was snap-

frozen in liquid nitrogen and stored at −80°C until analysis.

Single muscle fiber isolation. Muscle fibers were isolated from freeze-dried specimens as

previously described[22]. In brief, muscle biopsies were freeze-dried for 48 hours. Subsequently,

fibers were isolated in a humidity- and temperature-controlled room (humidity of 25%) using

fine forceps under a stereomicroscope. Approximately 200 single muscle fibers were isolated

for each biopsy, resulting in a total of 1038. To ensure the fibers settled at the bottom of the

tube, each fiber-containing tube underwent centrifugation at 20,000 g using a small

centrifuge. Next, fibers were resuspended in 15 µL of lysis buffer (1% sodium dodecyl sulfate

(SDC), 40 mM chloroacetamide (CAA), 10 mM dithiothreitol (DTT) in 50 mM Tris pH 8.5).

Sample collection – myopathy

Participant information. Six patients with severe nemaline myopathy were selected from our

nemaline myopathy study cohort. Three patients had pathogenic variants in *ACTA1*,

representing the conventional severe form, and three patients had pathogenic variants in

TNNT1, resulting in a rare, progressive form of nemaline myopathy. Three healthy individuals

with no history of neuromuscular disease were used as controls (**Table S20**).

Muscle biopsy collection. Healthy control participant biopsies (n = 3) were used from an

original study[75], and were therefore collected, snap frozen in liquid nitrogen and stored in -

80℃ as per that originally described. For the present study, a fragment of this stored biopsy

was dissected under sterile, frozen conditions before being prepared for single myofiber

isolation (detailed below). Acquisition of biopsies of healthy control patients was approved by

local ethics committee (Copenhagen and Frederiksberg) in Denmark (hs:h-15002266). Those

of myopathy patients were consented, stored, and used in accordance with the Human Tissue

Act under local ethical approval in United Kingdom (REC 13/NE/0373). All procedures were

carried out in accordance to the Declaration of Helsinki.

Single fiber isolation. Dissected fragments of muscle biopsy were placed in ice-cold, 22 micron

filtered relaxing solution (4 mM Mg-ATP, 1 mM free Mg $^{2+}$, $10^{-6.00}$ mM free Ca^{2+}, 20 mM

imidazole, 7 mM EGTA, 14.5 mM creatine phosphate, KCl to an iconic strength of 180 mM and

pH to 7.0)[76] for ~3 minutes before being immersed in fresh relaxing solution on a sterile petri

dish and mounted on ice under a dissection microscope for single fiber isolation. Fibers were

cleaved from the tissue/biopsy ensuring a variety of sample/biopsy locations were used and

that only single fibers were selected. Following isolation, fibers were manually moved to a

sterile 96-well plate containing 15 µL of lysis buffer (identical to that detailed above) where

the tweezers containing the fiber were submerged, spun and agitated in lysis buffer to ensure

the fibers dissociated from the tweezers. To ensure fibers were settled to the bottom of the

well, the 96-well plate was subjected to gentle vortexing and centrifugation (1,000 g).

Proteomics analysis

Sample preparation. Samples from both proteomics studies followed the same sample

preparation workflow. In order to extract the proteins, samples were boiled at 95°C in a

thermomixer with gentle shaking (800 rpm) and sonicated in a bioruptor instrument with 30

seconds on/off cycles for 15 minutes. A small 5 µL fraction of lysate from each sample was

saved for antibody-based fiber typing of the 1000 fiber samples. Next, samples were

processed following a modified version of the in-solution digestion sample preparation

protocol. In brief, total volume was adjusted to 50 µl by addition of digestion buffer, containing

50 mM Tris PH 8.5 buffer, an enzyme to protein ratio of 1:500 LysC (Wako) and a 1:100 enzyme

to protein ratio of trypsin (Promega). SMF lysates were digested overnight in a thermomixer

set to 37° C and 800 rpm. The next day, protein digestion was quenched by addition of 50 µl

of 2% trifluoroacetic acid (TFA) in isopropanol. Peptides were desalted using in-house

prepared single-use reverse-phase StageTips containing styrenedivinylbenzene reverse-phase

sulfonate (SDB-RPS) disks. Then, desalted peptides were loaded in Evotips (Evosep) following

manufacturer instructions prior LC-MS/MS analysis.

Proteomics library preparation. Fibers from the five healthy control individuals participating

in the 1000 fiber study were carefully dissected and combined in order to create a pooled

fiber lysate. Then, 200 µg of protein corresponding to each participant-specific lysate were

pooled together into one final protein lysate that was processed following the same sample

preparation workflow just described. 20 µg of desalted peptides were fractionated using High

pH Reverse-Phase Chromatography (HpH-RP). Fractionation was carried out on a Kinetex 2.6

µm EVO C18 100 Å, 150 x 0.3 mm column manufactured by Phenomenex and using an EASY-

nLC 1200 System (Thermo) operating at 1.5 µL/min. Separation was accomplished using a 62

min step-gradient starting from 3% to 60% solvent B (which consisted of 10 mM TEAB in 80%

acetonitrile) and solvent A (containing 10 mM TEAB in water). The total run time was 98 min,

which included wash and column equilibration. Throughout the fractionation, peptides were

eluted and collected every 60 s, obtaining 96 single fractions without concatenation. Finally.

200 ng of HpH-RP fractionated peptides were loaded, concentrated and desalted on Evotips

(Evosep) following the instructions provided by the manufacturer.

Liquid chromatography tandem mass spectrometry. Proteomics measurements were

performed using LC-MS instrumentation consisting of an Evosep One HPLC system (Evosep)[20]

coupled via electrospray ionization to a timsTOF SCP mass spectrometer (Bruker). Peptides

were separated utilizing 8 cm, 150 µM ID columns packed with C18 beads (1.5 µm) (Evosep).

Chromatographic separation was achieved by the '60 samples per day' method, followed by

electrospray ionization through a CaptiveSpray ion source and a 10 µm emitter into the MS

instrument. Single muscle fiber peptides were measured in DIA-PASEF mode following a

previously described method[21], while library fractions were measured using DDA-PASEF. In

brief, the DDA-PASEF scan range encompassed 100 - 1700 m/z for both MS and MS/MS, and

TIMS mobility range was set to $0.6 - 1.6$ (V cm^{-2}). Both TIMS ramp and accumulation times

were configured to 100 ms, and 10 PASEF ramps were recorded for a total cycle time of 1.17

s. The MS/MS target intensity and intensity threshold were defined as 20.000 and 1.000,

respectively. An exclusion list of 0.4 min for precursors within 0.015 m/z and 0.015 V cm^{-2}

width was also activated. For DIA-PASEF the scan range was established at 400-1000 (m/z),

the TIMS mobility range to 0.64-1.37 (V cm^{-2}), and ramp and accumulation times were both

set to 100 ms. A short-gradient method was used, which included 8 DIA-PASEF scans with

three 25 Da windows per ramp, resulting in an estimated cycle time of 0.95 sec.

MS data processing. Library files were processed using the MS-Fragger functionality within

Fragpipe v19.0 under the SpecLib workflow with default settings, including a minimum

peptide length of seven amino acids and maximum of two missed cleavages allowed[77]. Spectra

were searched against a human reviewed FASTA from Uniprot (March 2022, 20410 entries)

and the output library contained a total of 5350 protein groups and 84383 precursors. Sample

raw MS files were analyzed using DIA-NN version 1.8[78] in a library-based manner against the

MS library just described. Protein groups quantification was based on proteotypic peptides,

neural network was set to double-pass mode, quantification strategy was set to "Robust LC

(high accuracy)" and the match between runs options was enabled, the rest of parameters

remained as default, which included precursor FDR set to 1% and peptide length of 7-30 amino

acids.

Data processing. Further data analysis was performed under the R environment (R version

4.22). Both a metadata data frame containing sample and participant information and the

"PG_matrix.tsv" file from DIA-NN's output were loaded in RStudio. The 1000 fiber data frame

was filtered to remove samples with less than 50% valid protein intensity values, resulting in

a total of 974 fibers. Next, rows were filtered to remove proteins with less than 30% valid

values across samples, resulting in a total of 1685 proteins. Regarding the myopathy dataset,

after filtering samples for 50% valid values, the number of samples was 250. We included in

the analysis proteins that were quantified in 70% of the samples in at least one condition

(conditions: control, actin myopathy and troponin myopathy), resulting in a total of 1545

proteins. Both data frames were then log2 transformed and normalized using the

normalizeBetweenArrays() function from the limma package, with the method argument set

to quantile[79]. Then, batch correction through the *ComBat()* function from the sva package[80]

was applied to minimize the effect of the three technical batches originated during mass

spectrometry measurement. Finally, missing values were replaced by random numbers from

a Gaussian distribution with the default settings of the *tImpute()* function from the PhosR

package[81]. All relevant metadata for the 1000 fiber proteome and nemaline myopathy

datasets can be found in **Tables S28 and S29, respectively**.

Bioinformatics analysis

Transcriptome and proteome dynamic range. The expression/intensity for each gene/protein

was calculated relative to the total counts/intensity for each fiber. This value was then

averaged across fibers in each dataset and log10-transformed. The overlap of detected

features between both datasets were analyzed using the *VennDiagram* package (v 1.7.3).

Coefficient of variation – proteomics. For each of the 96 well plates used during the MS

measurement of the 1000 fiber study, one technical control sample was included in A1

position to monitor total ion current intensity and quality control of the runs (a total of eleven

technical controls). The coefficient of variation between proteins was calculated by dividing

the standard deviation by the mean of the LFQ intensities from each protein across technical

replicates and then multiplied by one hundred.

Correlation analyses. Mean log2 transformed transcript counts, protein intensities, and/or

fold change values across fibers were calculated, filtered for shared proteins/genes and

Pearson correlation was calculated.

Omics-based fiber typing. Normalized counts and raw LFQ intensities were retrieved from

well-described contractile proteins that have a slow (MYH7, TNNT1, TPM3, ATP2A2 and MYL3)

and fast (MYH2, MYH1, TNNT3, TPM1, ATP2A1 and MYL1) isoform[1]. For each isoform

combination, the relative expression of each was then calculated and samples were ordered

from high to low. The mathematical bottom knee for each curve was then determined using

the *barcodeRanks()* function in the DropletUtils package (v 1.18.1). This threshold was used

to assign fiber types as pure (type 1, type 2A or type 2X) or hybrid (hybrid 1/2A, hybrid 2A/2X

or hybrid 1/2X) (**Table S6**). For the features with only two isoforms, fibers were assigned as

'slow', 'fast' or 'hybrid'. To determine the overlap of the contractile features assigning a fiber

as being slow, upset plots were generated using the *upset()* function of the ComplexUpset

package (v 1.3.3), and then simplified to bar plots.

Principal Component Analysis (PCA). PCA was performed using the *RunPCA()* function of the

Seurat package. Scree plots were based on the *fviz_eig()* function of the factoextra package (v

1.0.7), which worked after PCA with *prcomp()*.

Seurat clustering. Uniform Manifold Approximation and Projection (UMAP) clustering was

performed based on the K-nearest neighbor graph with the first 6 dimensions as input for

both the transcriptome and proteome datasets (**Fig 1 S3D-E**). Feature plots were generated

using the *FeaturePlot()* function. UMAP plots were colored based on different criteria (MYH-

based fiber types, participant, test day) stored in the metadata.

Enrichment analysis. Genes and protein sets were processed to obtain lists of features that were differentially expressed or, in the case of the top PCA drivers, in the top 5% drivers for the positive and negative direction of the first and second principal component. Over-representation analysis was then performed on these features with the *enrichGO()* and *simplify()* function of the clusterProfiler package (v 4.6.2)[82] using all gene ontology terms. Obtained lists of significant terms were manually curated to extract interesting and relevant terms.

Hierarchical clustering of ribosomal proteins. Raw proteomics data was log2 transformed and filtered to contain proteins enlisted in the 'cytosolic ribosome' GO term, followed by Z-scoring prior heatmap visualization using the *pheatmap()* function from the Pheatmap package (v 1.0.12). The number of clusters was determined by visual inspection and assigning a value of 3 to the *cuttree*() function.

Differential expression analysis. To avoid artificially inflated p-values, which would arise from regarding every fiber as an independent replicate, we employed a pseudobulk differential expression analysis approach. We mathematically downsampled the total data points to one value per MYH-based fiber type per participant by aggregating (transcriptomics) or taking the median value (proteomics). Transcriptomics data were further processed using the DESeq2 pipeline (v 1.38.3)[83] with a '~ participant + fiber type' statistical model. 1000 fiber proteomics data was processed using the limma (v 3.54.2) workflow, fitting the data to a linear model defined as: '~ 0 + fiber type + participant', whereas the myopathy dataset was fitted to '~ 0 + condition' for the comparisons between conditions and '~ 0 + fiber type and condition' for the comparisons including fiber type. Fitted models were then subjected to gene ranking using an empirical Bayes method using *eBayes*() prior extracting the results through *topTable*(), with p-value adjustment set to Benjamini-Hochberg, both functions from the limma package. Threshold for significantly different genes/proteins was defined as adjusted p-values smaller than 0.05. For the nemaline myopathy dataset, the Xiao significance score was applied, which combines expression fold change and statistical significance[84]. Proteins with a Xiao score under 0.05 were regarded as differentially expressed between conditions.

Transcription factor motif enrichment. This analysis was performed using the gProfiler

webtool (https://biit.cs.ut.ee/gprofiler/gost), with the list of differentially expressed genes in

each slow/fast fiber type as input, ordered based on adjusted p-values (transcriptomics data).

Data was sourced from the TRANSFAC database.

Non-coding RNA. For the transcriptomics data, the biotype of each gene was determined

using the 'GENEBIOTYPE' column using the AnnotationDbi package (v 1.60.2) with the

EnsDb.Hsapiens.v86 database. Genomic location interrogation was performed using the UCSC

Human Genome Browser (https://genome.ucsc.edu). Tissue-specific gene expression of

interesting long non-coding RNAs was explored using the GTEx Portal database.

Microprotein identification.

Construction of putative lncRNA-encoded protein database

RNA sequences of the non-coding transcripts were extracted using the *getSequence() function*

from the biomaRt package (v 2.56.1)*, with 'transcript_exon_intron' as the *seqType.* Both

intergenic (lincRNA) and antisense long non-coding RNA (lncRNA) transcripts were utilized for

database construction. A six-frame-translation was used to translate the corresponding RNA

sequences into the proteins, as well as ORFfinder NCBI functionality

(https://www.ncbi.nlm.nih.gov/orffinder/) was used to extract open reading frames (ORFs)

from the transcripts. Minimal ORF length was set to 75 nucleotides, genetic code was set to

"Standard" and "Any sense codon" was used, as a start codon, to extract maximum number

of open reading frames. The obtained protein fasta file contained multiple entries for each

gene name, ensured by various combinations of: i) transcript identifiers, ii) ORF identifiers and

iii) start:stop codons.

Identification of lncRNA-encoded proteins

DIA raw MS data were analysed with Spectronaut v18 using an in-house generated sample-

specific fasta, comprised of the reviewed human proteome (proteome ID: UP000005640, 20

426 proteins, downloaded Sep 2023) and lncRNA-encoded protein sequences (125 223

proteins), in directDIA mode. The default settings were used unless otherwise noted. Data

filtering was set to   "Qvalue".   False discovery rate (FDR)   was set to 1%   at

peptide precursor level and 1% at protein level. Top3 peptide precursors were used for protein

quantification. The downstream data analysis was performed using in-house developed R

script.

PCA projection. The 1000 fiber and myopathy data sets were initially filtered to remove non-

overlapping proteins. Then, they were combined and normalized using the

normalizeBetweenArrays() function from the limma package. The normalization method used

was "quantile" to ensure that both data sets had the same distributions and were comparable.

Subsequently, the merged data set was divided back into the two separate data sets, namely

the 1000 fiber data set and the myopathy data set. For the 1000 fiber data set, PCA was

calculated using the *prcomp()* function. Moving on, the myopathy data set was multiplied by

the PC loadings obtained from the 1000 fiber dataset to generate its PCA projection. Finally,

the PCA projections from the myopathy samples were plotted on the top of the 1000 fiber

PCA visualization.

Muscle-specific ribosomal gene signature

The skeletal muscle-specific ribosomal gene signature, consisting of log2 fold change values

comparing the mRNA expression of ribosomal subunits in skeletal muscle compared to 52

other human tissues, was downloaded from Panda et al[85]. Log2 fold change values were

ranked to identify ribosomal proteins with the highest overexpression in human skeletal

muscle.

Structural Analysis of Ribosomal Proteins

The human 80S ribosome structural model (Protein Data Bank: 4V6X) was downloaded from

the Protein Data Bank website (RCSB PDB). Visualization and editing of the ribosomal

structure, and preparation of figures and movies were performed in UCSF ChimeraX[86].

Antibody-based fiber typing

Dot blot was conducted following a previously described protocol with a few modifications[22].

Initially, two identical PVDF membranes were activated using 96% ethanol and washed with

transfer buffer. Subsequently, the membranes were placed on wet filter paper with transfer

buffer until they dried. Next, 1 μL of fiber lysate was spotted at the same position on both

membranes, and the membranes were allowed to dry. Reactivation of the membranes was carried out using 96% ethanol, followed by gentle washing with TBST. The membranes were then blocked in TBST containing 5% skim milk for 15 minutes. After blocking, the membranes were washed three times with TBST and incubated with the primary antibody solution of either anti-MYH7 (A4.840) or anti-MYH2 (A4.74), both from Developmental Studies Hybridoma Bank (DSHB) at a dilution of 1:200 in TBST containing 1% skim milk for one hour. Subsequently, the membranes were gently washed three times with TBST and incubated with the secondary antibody (anti-mouse) at a dilution of 1:20,000 in TBST containing 1% skim milk for two hours. Finally, the membranes were washed three times for five minutes each with TBST and visualized using Immobilon Forte (Milipore) in a ChemiDoc XRS+ (Bio-Rad) imaging system.

RNAscope.

8μm sections from fixed-frozen human muscle biopsies were used for RNAscope labelling and subsequent immunohistochemistry (IHC). For detection of RP11-255P5.3 and LINC01405 commercially available RNAscope Multiplex Fluorescent Assay V2 (Advanced Cell Diagnostics) and probes against Lnc-ERCC5-5-C1 (# 1276851-C1), and LINC01405-C2 (# 549201-C2) (Advanced Cell Diagnostics), were used according to the manufacturer's protocols. To control tissue quality Positive 3-plex probe (# 320881) and negative 3-plex probe (# 320871) were used. To visualize different subtypes of muscle fibers, sections after RNAscope were blocked with 5% donkey serum and incubated with antibodies against MYH2 (A4.74-s (1:5); DSHB) and MYH7 (A4.840 (1:5); BSHD) overnight. After washing sections were incubated with Alexa Flour 488-conjugated Donkey Anti-Mouse IgG, Fcγ Subclass 1 and DyLightTM405-conjugated Donkey Anti-Mouse IgM secondary antibodies respectively and mounted with ProLong™ Diamond Antifade Mountant (Invitrogen). Slides were imaged using Zeizz Axio Observer microscope equipped with Axiocam 702 camera.

Immunostaining for muscle sections

Immunolabelling was performed on 10 μm cryosections, fixed in 4% PFA (10 min), permeabilized in 0.1% Triton X-100 (20 min) and blocked in 10% Normal Goat Serum (50062Z, Life Technologies) with 0.1% BSA (1 h). Sections were incubated o/n (4°C) with primary antibodies against MYH7 (mouse monoclonal A4.951, Santa Cruz, sc-53090, diluted 1:25) or

MYH2 (mouse monoclonal SC71, DSHB, 1:25), each combined with primary antibody against TNNT1 (rabbit polyclonal HPA058448, Sigma, diluted 1:500) in 5% goat serum with 0.1% BSA and 0.1% Triton X-100. Alexa Fluor Goat anti-Mouse 647 (A21237) was used as the secondary antibody for the MYHs and Alexa Fluor Donkey anti-Rabbit 488 (A11034) for the tropomyosin (Life Technologies, 1:500 each in 10% Normal Goat Serum). Fluorescent images were obtained with a 10x objective on a Zeiss Axio Observer 3 fluorescence microscope with a Colibri 5 led detector, combined with Zeiss Axiocam 705 mono camera, using Zen software (Zeiss). For visualization purposes, a selection of fibres were mounted on copper grids glued on a microscopy slide, and imaged under a stereomicroscope.

Acknowledgments

Mass spectrometry analyses were performed by the Proteomics Research Infrastructure (PRI) at the University of Copenhagen (UCPH), supported by the Novo Nordisk Foundation (NNF) (grant agreement number NNF19SA0059305). This work was generously funded by grants from the Novo Nordisk Foundation (NNF-0070539; to JO), the Foundation Building Strength For Nemaline Myopathy (to JO) and the Research Foundation Flanders (G080321N, 11B4220N and 1268023N; to WD, TVdS and EL, respectively). Several plots (Fig 1A, Fig 1S1A, Fig 5S2H) were made with BioRender.com. The authors would like to thank Leonidas Lundell & Lars Roed Ingerslev for their input on data analysis and visualization.

Author contributions

Conceptualization: RMJ, TVdS, BS, EL, PM, JV, JO, MH, WD, ASD

Methodology: RMJ, TVdS, BS, JL, RAS, DS, OD, AVdL, EL, LMA, KVV, MW, PM, JV, MH, WD, ASD

Software: RMJ, TVdS, BS, DS, LMA, JA

Validation: RMJ, TVdS, BS, JL, PM, JV, WD, ASD

Formal Analysis: RMJ, TVdS, BS, JL, DS, LMA

Investigation: RMJ, TVdS, BS, JL, RAS, DS, AVdL, EL, LMA, KVV, NY, AW, RVT, RES, EZ, MWL

Resources: RES, EZ, MWL, MW, PM, JV, JO, MH, WD, ASD

Data curation: RMJ, TVdS, BS, JL, DS, LMA, JA

Writing – Original Draft: RMJ, TVdS, BS, WD, ASD

Writing – Review and Editing: RMJ, TVdS, BS, JL, RAS, DS, OD, AVdL, EL, LMA, KVV, NY, JA, AW,

RVT, RES, EZ, MWL, MW, PM, JV, JO, MH, WD, ASD

Visualization: RMJ, TVdS, BS, JL, LMA, OD

Supervision: BS, PM, JV, JO, MH, WD, ASD

Project Administration: RMJ, TVdS, BS, WD, ASD

Funding Acquisition: TVdS, EL, PM, JV, JO, MH, WD, ASD

Supplemental Figures

Figure 1 S1. **Quality control of novel transcriptomic and proteomic workflows.** (**A**) Detailed schematic of

developed workflows. (**B-C**) Number of quantified genes/proteins by participant and sample in 1000 fiber

transcriptomics and proteomics. (**D**) Coefficient of variation by protein in 1000 fiber proteomics. (**E-F**) Dynamic

range of genes in 1000 fiber transcriptomics and of proteins in 1000 fiber proteomics. (**G**) Overlapping features

within 1000 fiber transcriptomics and proteomics. (**H**) Correlation between features within 1000 fiber

transcriptomics and proteomics.

1005

1006

1007 **Figure 1 S2. Higher sensitivity with proteomics compared to standard dot blotting.** (**A**) MYH-based fiber type

1008 proportions by participant. (**B**) Comparison between dot blot- and proteomics-based fiber typing.

1009

Figure 1 S3. Clustering of transcriptomic and proteomic data. (**A-B**) Scree plots of transcriptomics and proteomics. Colored box indicates the number of principal components used for clustering. (**C-D**) 1000 fiber transcriptomics and proteomics UMAP colored by participant. (**E**) 1000 fiber transcriptomics UMAP colored by trial day.

Figure 2 S1. Heterogeneity of fibers with well-known fiber type specific features. (**A-B**) Fiber type curves of established slow fiber type genes/proteins in 1000 fiber transcriptomics and proteomics. (**C-D**) Proportion of fibers determined to be slow via different numerical combinations of genes/proteins in 1000 fiber transcriptomics and proteomics.

Figure 2 S2. Additional data of Principal Component Analysis (PCA). (A-B) 1000 fiber transcriptomics PCA plot colored by participant or trial day. (**C**) 1000 fiber proteomics PCA plot colored by participant. (**D-E**) Top gene and protein drivers of PC1 and PC2.

1021   **Figure 3 S1. Ribosomal heterogeneity of skeletal muscle fibers.** (**A**) mRNA expression of ribosomal proteins in

1022   skeletal muscle compared to the whole body. Data derived from Panda et al[67]. (**B**) PCA plot colored by cytosolic

1023   ribosome gene ontology (GO) term in the transcriptome. Arrows represent protein drivers and their direction.

1024   (**C**) Proportions of ribosomal proteins in each cluster by small (40S) and large (60S) subunits.

Figure 4 S1. Gene and protein signature of fast and slow fibers. (**A**) Feature plots of LDH protein isoforms between slow and fast fibers. (**B**) Overlap of significantly different genes/proteins between slow and fast skeletal muscle fibers in 1000 fiber transcriptomics and proteomics. (**C**) Correlation between the log$_2$ fold change difference between slow and fast skeletal muscle fibers within 1000 fiber transcriptomics and proteomics. (**D-K**) Expression profiles of top drivers of selected enrichment terms within slow or fast fibers.

Figure 4 S2. Specific expression of non-coding RNA in skeletal muscle. (A-B) Expression profile of *LINC01405* and *RP11-255P5.3* in different tissues. (C) Volcano plots comparing microprotein abundance between the slow and fast clusters identified from the uniform manifold approximation and projection (UMAP) plots in Fig 1G-H. Colored dots represent significantly different microproteins at an FDR < 0.05. (D) Slow/fast fiber-type comparison of LINC01405_ORF52 14515:14294 and LINC01405_ORF503 18862:18770 microproteins.

1035   **Figure 5 S1. Quality control and fiber typing of nemaline myopathy dataset.** (**A**) Number of quantified proteins

1036   by participant and sample in nemaline myopathy proteomics. (**B**) Dynamic range of proteins in nemaline

1037   myopathy proteomics. (**C**) Correlation between features within 1000 fiber proteomics and healthy controls of

1038   nemaline myopathy proteomics. (**D**) Dynamic range curves for MYH7, MYH2, and MYH1, with calculated

1039   assignment thresholds. (**E**) Distribution of fibers by expression of MYHs in nemaline myopathy proteomics.

Figure 5 S2. Fiber type-independent remodelling in nemaline myopathy. (**A**) Correlation between the log$_2$ fold change difference between *ACTA1*- and *TNNT1*-nemaline myopathies with control participants. (**B**) PCA of nemaline myopathy patients (pseudobulked protein profile form single fibers). (**C**) Upset plot of significantly regulated proteins *ACTA1*- and *TNNT1*-nemaline myopathies. (**D-I**) Volcano plots comparing the *ACTA1*- and *TNNT1*-nemaline myopathies with control participants by MYH-based fiber type. (**J**) Abundance of annexin proteins in *ACTA1*- and *TNNT1*-nemaline myopathy. (**K**) Schematic of changes within the sarcolemma repair machinery in nemaline myopathy.

Supplementary Tables

Tables can be accessed here: https://drive.google.com/drive/folders/15j7U-Yj3_2y9bICxouWAhF4yUgXVHBHL

Table S1. Characteristics of participants of the 1000 fiber transcriptomics and proteomics datasets.

Table S2. Raw UMI counts matrix of the 1000 fiber transcriptomics dataset.

Table S3. Raw LFQ intensity matrix of the 1000 fiber proteomics dataset.

Table S4. Quantified transcripts (raw UMI counts) within the 1000 fiber transcriptomics dataset after filtering.

Table S5. Quantified proteins ($\log_2$ LFQ intensity) within the 1000 fiber proteomics dataset after filtering and imputation.

Table S6. Classification system for fiber typing based on myosin heavy chain isoform expression.

Table S7. PC1 and PC2 drivers within the 1000 fiber transcriptomics dataset.

Table S8. PC1 and PC2 drivers within the 1000 fiber proteomics dataset.

Table S9. Over-representation analysis of the top drivers (top 5%) in PC1 and PC2 within the 1000 fiber transcriptomics and proteomics datasets.

Table S10. Protein abundance of proteins within the GO term "Cytosolic Ribosome" in the 1000 fiber proteomics dataset.

Table S11. Pseudobulked transcript expression within slow and fast skeletal muscle fibers in the 1000 fiber transcriptomics dataset.

Table S12. Pseudobulked protein abundance within slow and fast skeletal muscle fibers in the 1000 fiber proteomics dataset.

Table S13. Differential expression results between slow and fast skeletal muscle fibers in the 1000 fiber transcriptomics dataset.

Table S14. Differential abundance results between slow and fast skeletal muscle fibers in the 1000 fiber proteomics dataset.

Table S15. Over-representation analysis of the differentially regulated transcripts and proteins between slow and fast skeletal muscle fibers in the 1000 fiber transcriptomics and proteomics datasets.

1091   **Table S16**. Transcription factor binding motif enrichment using the TRANSFAC databases on
1092   the differentially expressed transcripts between slow and fast fibers in the 1000 fiber
1093   transcriptomics dataset.
1094

1095   **Table S17**. Differential expression results between slow and fast skeletal muscle fibers in
1096   non-coding transcripts in the 1000 fiber transcriptomics dataset.
1097

1098   **Table S18**. Raw LFQ intensity matrix of the microprotein datset.
1099

1100   **Table S19**. Differential expression results between slow and fast skeletal muscle fibers in the
1101   microprotein dataset.
1102

1103   **Table S20**. Characteristics of patients within the nemaline myopathy proteomics dataset.
1104

1105   **Table S21**. Detected proteins within the nemaline myopathy proteomics dataset.
1106

1107   **Table S22**. Quantified proteins within the nemaline myopathy proteomics dataset after
1108   filtering and imputation.
1109

1110   **Table S23**. Pseudobulked protein abundance within the nemaline myopathy proteomics
1111   dataset.
1112

1113   **Table S24**. Differential protein abundance within *ACTA1*- and *TNNT1*-nemaline myopathies.
1114

1115   **Table S25**. Gene set enrichment analysis of the differential regulation within *ACTA1*- and
1116   *TNNT1*-nemaline myopathies.
1117

1118   **Table S26**. Differential protein abundance within *ACTA1*- and *TNNT1*-nemaline myopathies
1119   by MYH-based fiber type.
1120

1121   **Table S27**. Metadata for the 1000 fiber transcriptomics dataset.
1122

1123   **Table S28**. Metadata for the 1000 fiber proteomics dataset.
1124

1125   **Table S29**. Metadata for the nemaline myopathy proteomics dataset.

References

Schiaffino, S. & Reggiani, C. Fiber types in mammalian skeletal muscles. *Physiol Rev* **91**, 1447-1531 (2011).

Altschuler, S. J. & Wu, L. F. Cellular heterogeneity: do differences make a difference? *Cell* **141**, 559-563 (2010).

Engel, W. K. The essentiality of histo- and cytochemical studies of skeletal muscle in the investigation of neuromuscular disease. 1962. *Neurology* **12**, 778-784 (1962).

Brooke, M. H. & Kaiser, K. K. Three "myosin adenosine triphosphatase" systems: the nature of their pH lability and sulfhydryl dependence. *J Histochem Cytochem* **18**, 670-672 (1970).

Biral, D., Betto, R., Danieli-Betto, D. & Salviati, G. Myosin heavy chain composition of single fibres from normal human muscle. *Biochem J* **250**, 307-308 (1988).

Tang, F. *et al.* mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods* **6**, 377-382 (2009).

Islam, S. *et al.* Characterization of the single-cell transcriptional landscape by highly multiplex RNA-seq. *Genome Res* **21**, 1160-1167 (2011).

Brunner, A.-D. *et al.* Ultra-high sensitivity mass spectrometry quantifies single-cell proteome changes upon perturbation. *Mol Syst Biol* **18**, e10798 (2022).

Murgia, M. *et al.* Single muscle fiber proteomics reveals fiber-type-specific features of human muscle aging. *Cell Rep* **19**, 2396-2409 (2017).

Murgia, M. *et al.* Signatures of muscle disuse in spaceflight and bed rest revealed by single muscle fiber proteomics. *PNAS Nexus* **1**, pgac086 (2022).

Murgia, M. *et al.* Single muscle fiber proteomics reveals unexpected mitochondrial specialization. *EMBO rep* **16**, 387-395 (2015).

Murgia, M. *et al.* Protein profile of fiber types in human skeletal muscle: a single-fiber proteomics study. *Skeletal Muscle* **11**, 24 (2021).

Lang, F. *et al.* Single muscle fiber proteomics reveals distinct protein changes in slow and fast fibers during muscle atrophy. *J Proteome Res* **17**, 3333-3347 (2018).

Momenzadeh, A. *et al.* A complete workflow for high throughput human single skeletal muscle fiber proteomics. *J Am Soc Mass Spectrom* (2023).

Chemello, F. *et al.* Degenerative and regenerative pathways underlying Duchenne muscular dystrophy revealed by single-nucleus RNA sequencing. *Proc Natl Acad Sci USA* **117**, 29691-29701 (2020).

Lin, H. *et al.* Decoding the transcriptome of denervated muscle at single-nucleus resolution. *J Cachexia Sarcopenia Muscle* **13**, 2102-2117 (2022).

Sewry, C. A., Laitila, J. M. & Wallgren-Pettersson, C. Nemaline myopathies: a current view. *J Muscle Res Cell Motil* **40**, 111-126 (2019).

Laitila, J. & Wallgren-Pettersson, C. Recent advances in nemaline myopathy. *Neuromuscul Disord* **31**, 955-967 (2021).

Zhang, M. J., Ntranos, V. & Tse, D. Determining sequencing depth in a single-cell RNA-seq experiment. *Nat Commun* **11**, 774 (2020).

Bache, N. *et al.* A novel LC system embeds analytes in pre-formed gradients for rapid, ultra-robust proteomics. *Mol Cell Proteomics* **17**, 2284-2296 (2018).

Meier, F. *et al.* diaPASEF: parallel accumulation–serial fragmentation combined with data-independent acquisition. *Nat Methods* **17**, 1229-1236 (2020).

22   Deshmukh, A. S. *et al.* Deep muscle-proteomic analysis of freeze-dried human muscle
biopsies reveals fiber type-specific adaptations to exercise training. *Nat Commun* **12**,
304 (2021).

23   Pardo, J. V., Siliciano, J. D. & Craig, S. W. A vinculin-containing cortical lattice in
skeletal muscle: transverse lattice elements ("costameres") mark sites of attachment
between myofibrils and sarcolemma. *Proc Natl Acad Sci USA* **80**, 1008-1012 (1983).

24   Xiao, Y. *et al.* Loss of mitochondrial protein CHCHD10 in skeletal muscle causes
neuromuscular junction impairment. *Hum Mol Genet* **29**, 1784-1796 (2020).

25   Cho, H. J. *et al.* SLIT3 promotes myogenic differentiation as a novel therapeutic factor
against muscle loss. *J Cachexia Sarcopenia Muscle* **12**, 1724-1740 (2021).

26   Han, S. *et al.* Nuclear envelope phosphatase 1-regulatory subunit 1 (formerly
TMEM188) is the metazoan Spo7p ortholog and functions in the lipin activation
pathway. *J Biol Chem* **287**, 3123-3137 (2012).

27   Hengel, H. *et al.* Loss-of-function mutations in UDP-Glucose 6-Dehydrogenase cause
recessive developmental epileptic encephalopathy. *Nat Commun* **11**, 595 (2020).

28   Farhang-Fallah, J. *et al.* The pleckstrin homology (PH) domain-interacting protein
couples the insulin receptor substrate 1 PH domain to insulin signaling pathways
leading to mitogenesis and GLUT4 translocation. *Mol Cell Biol* **22**, 7325-7336 (2002).

29   Genuth, N. R. & Barna, M. The discovery of ribosome heterogeneity and its
implications for gene regulation and organismal life. *Mol Cell* **71**, 364-374 (2018).

30   Parks, M. M. *et al.* Variant ribosomal RNA alleles are conserved and exhibit tissue-
specific expression. *Sci Adv* **4**, eaao0665 (2018).

31   Slavov, N., Semrau, S., Airoldi, E., Budnik, B. & van Oudenaarden, A. Differential
stoichiometry among core ribosomal proteins. *Cell Rep* **13**, 865-873 (2015).

32   Shi, Z. *et al.* Heterogeneous ribosomes preferentially translate distinct subpools of
mRNAs genome-wide. *Mol Cell* **67**, 71-83.e77 (2017).

33   Kondrashov, N. *et al.* Ribosome-mediated specificity in Hox mRNA translation and
vertebrate tissue patterning. *Cell* **145**, 383-397 (2011).

34   Genuth, N. R. *et al.* A stem cell roadmap of ribosome heterogeneity reveals a
function for RPL10A in mesoderm production. *Nat Commun* **13**, 5491 (2022).

35   Landry, D. M., Hertz, M. I. & Thompson, S. R. RPS25 is essential for translation
initiation by the Dicistroviridae and hepatitis C viral IRESs. *Genes Dev* **23**, 2753-2764
(2009).

36   Horos, R. *et al.* Ribosomal deficiencies in Diamond-Blackfan anemia impair
translation of transcripts essential for differentiation of murine and human
erythroblasts. *Blood* **119**, 262-272 (2012).

37   Anger, A. M. *et al.* Structures of the human and Drosophila 80S ribosome. *Nature*
497, 80-85 (2013).

38   Spahn, C. M. *et al.* Cryo-EM visualization of a viral internal ribosome entry site bound
to human ribosomes: the IRES functions as an RNA-based translation factor. *Cell* **118**,
465-475 (2004).

39   Boehringer, D., Thermann, R., Ostareck-Lederer, A., Lewis, J. D. & Stark, H. Structure
of the hepatitis C virus IRES bound to the human 80S ribosome: Remodeling of the
HCV IRES. *Structure* **13**, 1695-1706 (2005).

40   Kannan, K., Vázquez-Laslop, N. & Mankin, A. S. Selective protein synthesis by
ribosomes with a drug-obstructed exit tunnel. *Cell* **151**, 508-520 (2012).

41   Armstrong, E. *et al.* Comparison of transcriptomic landscapes of different lamb
muscles using RNA-Seq. *PLoS One* **13**, e0200732 (2018).
42   Leberer, E. & Pette, D. Lactate dehydrogenase isozymes in type I, IIA and IIB fibres of
rabbit skeletal muscles. *Histochemistry* **80**, 295-298 (1984).
43   Rubenstein, A. B. *et al.* Single-cell transcriptional profiles in human skeletal muscle.
Sci Rep **10**, 229 (2020).
44   Lerín, C. *et al.* Regulation and function of the muscle glycogen-targeting subunit of
protein phosphatase 1 (GM) in human muscle cells depends on the COOH-terminal
region and glycogen content. *Diabetes* **52**, 2221-2226 (2003).
45   Luo, Y., Na, Z. & Slavoff, S. A. P-bodies: Composition, properties, and functions.
Biochemistry **57**, 2424-2431 (2018).
46   Chin, E. R. *et al.* A calcineurin-dependent transcriptional pathway controls skeletal
muscle fiber type. *Genes Dev* **12**, 2499-2509 (1998).
47   Kammoun, M. *et al.* Novel role of Tieg1 in muscle metabolism and mitochondrial
oxidative capacities. *Acta Physiologica* **228**, e13394 (2020).
48   Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding
RNAs and its biological functions. *Nat Rev Mol Cell Biol* **22**, 96-118 (2021).
49   Lorenzi, L. *et al.* The RNA Atlas expands the catalog of human non-coding RNAs. *Nat*
Biotech **39**, 1453-1465 (2021).
50   Wang, W. *et al.* Identification of miRNA, lncRNA and mRNA-associated ceRNA
networks and potential biomarker for MELAS with mitochondrial DNA A3243G
mutation. *Sci Rep* **7**, 41639 (2017).
51   Volders, P.-J. *et al.* LNCipedia 5: towards a reference set of human long non-coding
RNAs. *Nucleic Acids Res* **47**, D135-D139 (2019).
52   Matsumoto, A. *et al.* mTORC1 and muscle regeneration are regulated by the
LINC00961-encoded SPAR polypeptide. *Nature* **541**, 228-232 (2017).
53   Nelson, B. R. *et al.* A peptide encoded by a transcript annotated as long noncoding
RNA enhances SERCA activity in muscle. *Science* **351**, 271-275 (2016).
54   Bittel, D. C. *et al.* Annexin A2 mediates dysferlin accumulation and muscle cell
membrane repair. *Cells* **9** (2020).
55   Carmeille, R. *et al.* Membrane repair of human skeletal muscle cells requires
Annexin-A5. *Biochim Biophys Acta* **1863**, 2267-2279 (2016).
56   Jaiswal, J. K. *et al.* S100A11 is required for efficient plasma membrane repair and
survival of invasive cancer cells. *Nat Commun* **5**, 3795 (2014).
57   Cagliani, R. *et al.* Mutation finding in patients with dysferlin deficiency and role of the
dysferlin interacting proteins annexin A1 and A2 in muscular dystrophies. *Hum Mutat*
26, 283 (2005).
58   Leikina, E. *et al.* Extracellular annexins and dynamin are important for sequential
steps in myoblast fusion. *J Cell Biol* **200**, 109-123 (2012).
59   Zhu, B. *et al.* Monoubiquitination of human histone H2B: the factors involved and
their roles in HOX gene regulation. *Mol Cell* **20**, 601-611 (2005).
60   Needham, D. M. Red and white muscle. *Physiol Revs* **6**, 1-27 (1926).
61   Petrany, M. J. *et al.* Single-nucleus RNA-seq identifies transcriptional heterogeneity in
multinucleated skeletal myofibers. *Nat Commun* **11**, 6374 (2020).
62   Musalgaonkar, S., Black, J. J. & Johnson, A. W. The L1 stalk is required for efficient
export of nascent large ribosomal subunits in yeast. *Rna* **25**, 1549-1560 (2019).

63   Fernández, I. S., Bai, X. C., Murshudov, G., Scheres, S. H. & Ramakrishnan, V. Initiation
of translation by cricket paralysis virus IRES requires its translocation in the ribosome.
Cell **157**, 823-831 (2014).

64   Turner, D. C. *et al.* DNA methylation across the genome in aged human skeletal
muscle tissue and muscle-derived cells: the role of HOX genes and physical activity.
Sci Rep **10**, 15360 (2020).

65   Robinson, M. M. *et al.* Enhanced protein translation underlies improved metabolic
and physical adaptations to different exercise training modes in young and old
humans. *Cell Metab* **25**, 581-592 (2017).

66   D'Ercole, C. *et al.* Spatially resolved transcriptomics reveals innervation-responsive
functional clusters in skeletal muscle. *Cell Rep* **41**, 111861 (2022).

67   Fitzgerald, G. *et al.* MME+ fibro-adipogenic progenitors are the dominant adipogenic
population during fatty infiltration in human skeletal muscle. *Commun Biol* **6**, 111
(2023).

68   Babaeijandaghi, F. *et al.* Metabolic reprogramming of skeletal muscle by resident
macrophages points to CSF1R inhibitors as muscular dystrophy therapeutics. *Sci*
Trans Med **14**, eabg7504 (2022).

69   Chaillou, T., Zhang, X. & McCarthy, J. J. Expression of Muscle-Specific Ribosomal
Protein L3-Like Impairs Myotube Growth. *J Cell Physiol* **231**, 1894-1902 (2016).

70   Seaborne, R. A. E. & Ochala, J. The dawn of the functional genomics era in muscle
physiology. *J Physiol* **601**, 1343-1352 (2023).

71   Van der Stede, T. *et al.* Histamine H(1) and H(2) receptors are essential transducers of
the integrative exercise training response in humans. *Sci Adv* **7** (2021).

72   Bergström, J. Percutaneous needle biopsy of skeletal muscle in physiological and
clinical research. *Scand J Clin Lab Invest* **35**, 609-616 (1975).

73   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587.e3529 (2021).

74   Choudhary, S. & Satija, R. Comparison and evaluation of statistical error models for
scRNA-seq. *Genome Biol* **23**, 27 (2022).

75   Sahl, R. E. *et al.* Variation in mitochondrial respiratory capacity and myosin heavy
chain composition in repeated muscle biopsies. *Anal Biochem* **556**, 119-124 (2018).

76   Ranu, N. *et al.* NEB mutations disrupt the super-relaxed state of myosin and remodel
the muscle metabolic proteome in nemaline myopathy. *Acta Neuropathol Commun*
10, 185 (2022).

77   Demichev, V. *et al.* dia-PASEF data analysis using FragPipe and DIA-NN for deep
proteomics of low sample amounts. *Nat Commun* **13**, 3944 (2022).

78   Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural
networks and interference correction enable deep proteome coverage in high
throughput. *Nat Methods* **17**, 41-44 (2020).

79   Ritchie, M. E. *et al.* LIMMA powers differential expression analyses for RNA-
sequencing and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

80   Müller, C. *et al.* Removing batch effects from longitudinal gene expression - Quantile
normalization plus ComBat as best approach for microarray transcriptome data. *PLoS*
One **11**, e0156594 (2016).

81   Kim, H. J. *et al.* PhosR enables processing and functional analysis of
phosphoproteomic data. *Cell Rep* **34**, 108771 (2021).

82 Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics
data. *Innovation (Camb)* **2**, 100141 (2021).
83 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
84 Xiao, Y. *et al.* A novel significance score for gene selection and ranking. *Bioinformatics*
30, 801-807 (2014).
85 Panda, A. *et al.* Tissue- and development-stage–specific mRNA and heterogeneous
CNV signatures of human ribosomal proteins in normal and cancer samples. *Nucleic*
Acids Res **48**, 7079-7098 (2020).
86 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82 (2021).
1320