Organ-specific prioritization and annotation of non-coding regulatory variants in the human genome

Nanxiang Zhao[1,*], Rintsen N. Sherpa[1,*], Shengcheng Dong[2,*], Kobe B. Howcroft[1], Alan P. Boyle[1,3,†]

1. Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

2. Department of Genetics, Stanford University, Stanford, CA, USA

3. Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA

* These authors contributed equally

† Co-Correspondence: apboyle@umich.edu

Note: For the purpose of tracking manuscript status in the IGVF package, please include this e-mail in all manuscript correspondence: publications@igvf.org

Abstract

Identifying non-coding regulatory variants in the human genome remains a challenging task in genomics. Recently, we released the second version of our leading regulatory variant database, RegulomeDB. Building upon this comprehensive database, we developed a novel machine-learning architecture, TLand, which utilizes RegulomeDB-derived features to predict regulatory variants at the cell- or organ-specific level. In our holdout benchmarking, TLand consistently outperformed state-of-the-art models, demonstrating its ability to generalize to new cell lines or organs. We trained three types of organ-specific TLand models to overcome the common model bias toward high data availability cell lines or organs. These models accurately prioritize relevant organs for 2 million GWAS SNPs associated with GWAS traits. Moreover, our analysis of top-scoring variants in specific organ models showed a high enrichment of relevant GWAS traits. We expect that TLand and

1   RegulomeDB will further advance our ability to understand human regulatory variants

2   genome-wide.

Keywords

4   Genomics, non-coding variants, gene regulation, machine learning, disease genetics

Background

Understanding the biological impact of variants located in non-coding regions of the human

genome is a significant challenge. Nearly 90% of disease-risk-associated single-nucleotide

polymorphisms (SNPs) identified from genome-wide association studies (GWAS) are within

non-coding regions. Similarly, 75% of patients with Mendelian disease have mutations

outside of protein-coding regions (1). Delving into the functional consequences of these

disease-associated non-coding variants may uncover potential causal SNPs, expanding our

insights beyond those within the coding region.

Prioritizing non-coding variants requires integrating multiple layers of functional information,

including regulatory annotations identified from high-throughput sequencing datasets (e.g.,

DNase-seq (2), ChIP-seq (3), and ATAC-seq (4)). Such annotations provide additional

information for pinpointing causal variants, which are often not the lead variants identified in

GWAS. Despite the benefit of incorporating functional genomics assay-based evidence

when examining non-coding variants, the lack of available variant annotation tools limits the

use of such data. Most resources developed for clinical purposes have focused on coding

regions as an application of exome sequencing-based data (5,6), which captures less than

5% of human variation (7–9).

Previously, we developed RegulomeDB, a comprehensive database for prioritizing and

annotating variants in non-coding regions, which has been widely used in the research

community (10). RegulomeDB intersects query variants with regulatory regions predicted by functional genomics assays and, by utilizing ranking heuristics, informs users about putative functional consequences to prioritize variants. Recently, RegulomeDB has been upgraded to version 2 (11), improving its annotation power by incorporating thousands of new functional genomics datasets from the ENCODE project (12), Roadmap Epigenomics Consortium (13), and the Genomics of Gene Regulation Consortium. A suite of models, namely SURF and TURF, was developed and integrated in this version to provide accurate probabilistic scores for general and cell-type specific regulatory activities (14,15).

However, a drawback of this model suite is that it was trained using hg19-referenced ENCODE datasets for only six common cell lines. The resulting models are biased and lack sufficient statistical power to generalize their predictions to less-studied cell lines. As a result, RegulomeDB scores are less informative for cell lines, tissues, and organs that lack the abundance of data available for commonly-studied cell lines. This can make it challenging for users to identify variants of interest and formulate hypotheses regarding their regulatory functions. Nonetheless, we anticipate that RegulomeDB will continue to improve in the years to come as new datasets are released by the Encyclopedia of DNA Elements (ENCODE) (12) and Impact of Genomic Variation on Function (IGVF) (16) consortia, spanning additional cell lines, tissues, and assay types, such as Hi-C (17) and Enformer predictions (18). However, the current SURF and TURF models were not designed to incorporate data continuously, and are potentially prone to overfitting as the feature space expands due to the simple model architecture. Given the breadth and volume of functional genomics datasets being released each day, these are serious limitations.

Here, we present TLand, a flexible architecture based on stacked generalization (19) that learns RegulomeDB-derived features to predict regulatory variants at the cell- or organ-specific level. TLand's stacked generalization approach groups features into biologically meaningful subspaces, training individual estimators before assembly to reduce overfitting

and enable further integration of features. Cell-specific TLand consistently outperformed state-of-the-art models in benchmarking with a hold-out cell line. By developing a suite of models rather than a single, monolithic model, organ-specific TLand addresses the data availability bias and further improves on cell-specific TLand in identifying regulatory variants relevant to the target organ, as predicted by GWAS. Furthermore, analysis of the top-scoring variants in specific organ models revealed a high enrichment for correlated GWAS traits. Given its superior performance relative to its predecessors and competing methods, we expect TLand to address the ongoing challenge of reliably prioritizing variants, even in less-studied cell lines and organs, thus advancing our ability to identify regulatory variants genome-wide.

Methods

Allele-specific binding (ASB) variants

We define a variant as likely-regulatory if it shows evidence for allele-specific binding (ASB), the molecular signature of which is significantly different when ChIP-seq reads are phased between the two alleles in a heterozygous individual. We trained our models on ASB variants for all TFs with ChIP-seq data in ENCODE. We included a total of 7,530 ASB variants in 6 cell lines (GM12878, HepG2, A549, K562, MCF7, and H1hESC) called by *AlleleDB* (20). We utilized negative training sets previously produced in our lab (15), encompassing non-ASB variants and randomly-selected background variants. In total, we included 14,773 unique variants in our training set. The complementary ASB data of IMR-90 and H9 used for evaluation were downloaded from Adastra (21) at

https://adastra.autosome.org/bill-cipher/search/advanced?fdr=0.05&es=0&cl=IMR90%20(lung%20fibroblasts) and

https://adastra.autosome.org/bill-cipher/search/advanced?fdr=0.05&es=0&cl=H9.

Model architecture

TLand is a one-layer stacked architecture that consists of two parts: three base classifiers

and one meta-classifier (Fig 1b and Supplementary Fig 1). TLand takes input as 19 generic

features, 40 deep learning prediction-derived features, and 5 cell-specific features or 13

organ-specific features (Supplementary Table 1), which are directly derived from

RegulomeDB queries. Zeros would be assigned if no hits were found in RegulomeDB

queries. Features were bagged into 3 subspaces: experimental set (generic features and

cell/organ-specific features), deep learning set (deep learning features and cell/organ-

specific features), and cell/organ-specific set (only cell/organ-specific features). We selected

lightGBM (22), random forest (23), and neural network (24) as base classifiers due to their

distinct decision boundaries. Our base models were fine-tuned with Optuna (25). We used

300 estimators in random forest models, 250 boost rounds, and a 0.049 learning rate in

lightGBM models, 3 layers with 128 neurons per layer, batch size of 128, adaptive learning

rate, and with the max iteration of 30 in neuron network models. Probabilities were used as

the output of base classifiers and to train our meta-classifier. We calculated interaction terms

of probabilities up to the degree of 2 before feeding into our meta-classifier. We customized

a ridge classifier to output probabilities with hyperparameter alpha as 1.9 as our final meta-

classifier. The ridge classifier was trained with 4-fold group cross-validation. We grouped the

training data based on the genomic positions (i.e., variants at the same genomic positions

regardless of whether cell lines were in the same group). All models were specified with

balanced class weights. LightGBM was implemented with the Python package (22). Random

forest, neural network, and relevant pipelines were implemented with scikit-learn (26). The

stacked generalization algorithm was implemented with mlxtend (27).

Model training and evaluation

TLand was trained, validated, and tested on the generated ASB datasets. After

concatenating across six cell lines, 88,638 (i.e., 14,773 x 6) variants were used for training

and validation. The concatenation allowed us to have more data per prediction task. For the

task of predicting an unseen cell line, we held out data for one cell line as a test case and

then used the rest as training data for the TLand model. Simultaneously, during the train and

test split, we ensured that unique variants, irrespective of cell lines, were exclusively present

either in the training or test dataset. This precaution aimed to prevent information leakage,

as features across cell lines are often correlated. The test and train split ratio for negative

data is □. Similarly, for organ-specific models, we held out one organ and unique variants as

a test dataset and used organ-specific labels for splitting the data. We calculated both the

Area Under the Receiver Operating Characteristic Curve (AUROC) and the Area Under the

Precision Recall Curve (AUPR) to evaluate the classifiers. We also performed a separate

evaluation using leave-one-chromosome-out cross-validation to account for any sequence-

specific data leakage, as the Sei-derived features are conditional on 4096bp of sequence

context. We aggregated the variant sets across cell lines, calculated the AUPR for each held

out chromosome set, reported mean AUPR, and computed 95% confidence intervals using

1000 bootstrapped samples. After evaluations, the final models were trained with all 88,638

variants.

DNase signal quantile normalization

All 1372 DNase bigwig default files processed on GRCh38 assembly (up to May 2022) were

obtained from ENCODE (Supplementary Table 5). We designed an efficient pipeline to

quantile normalize all signals to achieve the balance between accuracy, runtime, and

storage (Supplementary Fig 2). BigWig files were first converted to BedGraph format using

bigWigToBedGraph (28). Average signals were extracted for 10bp, non-overlapping

windows extracted from each BedGraph file with bedmap (29) and stored in bed files. These

bed files were concatenated and converted to parquet format for processing with qnorm (30),

a Python package that applies a multithreaded incremental normalization strategy to

efficiently normalize extremely large datasets (30). We quantile normalized 10-bp average

signals for all 1372 DNase datasets with a batch size of 343 files per normalization iteration.

Details of storage, runtime, and memory are shown in Supplementary Fig 2.

Benchmarking other models and organ definition

We benchmarked TLand with state-of-the-art models, including GenoNet, DeepSEA, Sei,

and TURF. Pre-calculated GenoNet scores were downloaded from

https://zenodo.org/record/3336209/files/. We imputed the missing prediction scores of

GenoNet as the average of all available scores. DeepSEA and Sei models were downloaded

from the original paper (31,32). We ran the predictions locally. The previous best model,

TURF, was retrained to predict regulatory variants on GRCh38 by querying and inputting the

GRCh38 features (15). TURF predictions for unseen cell lines were derived by training

separate models on each of the seen (i.e., training) cell lines, then taking the average of all

trained models as an estimate for the unseen cell line. This resembles the ensemble method

of majority vote. For organ-level predictions, we averaged relevant cell line model predictions

to estimate the predictions, using the human organ definitions from ENCODE

(https://www.encodeproject.org/summary/?type=Experiment&control_type!=*&replicates.libra

ry.biosample.donor.organism.scientific_name=Homo+sapiens&status=released.). For

example, we averaged TURF model predictions of A549 and IMR-90 to predict lung

regulatory variants.

Evaluation with MPRA data

We download the new MPRA data of five cell lines from ENCODE, including GM12878

(ENCODE Accession ID: ENCSR560QIE), K562 (ENCSR971PLA), A549 (ENCSR793OLW),

HCT116 (ENCSR873QHK), and SK-N-SH (ENCSR579QGV). We trained the final set of

TLand models using all available data (i.e., from six training cell lines). Then we evaluated

the final models by calculating the AUROC and AUPR given the cell line's p-value threshold

(Skew -log10(FDR) at least > 5) to define positive and negative labels.

1 GWAS catalog variants and LD data

We first downloaded all known SNV positions significantly associated with traits described in

the GWAS Catalog (https://www.ebi.ac.uk/gwas/api/search/downloads/alternative) (33). We

then completed linkage disequilibrium (LD) expansion for each SNV, incorporating all SNVs

from the 1000 Genome Project in strong LD (R^2 threshold of 0.6). R^2 values were

downloaded from (gs://genomics-public-data/linkage-disequilibrium). In total, we calculated

their TLand model scores for 1,974,549 SNVs.

Summary statistics for 94 GWAS traits from the UK BioBank (34) were downloaded from

https://console.cloud.google.com/storage/browser/finucane-requester-pays/ukbb-

finemapping. Metadata for these traits (trait definitions and number of samples, cases, and

controls) were downloaded from https://www.finucanelab.org/data. Summary statistics were

prepared for use in LDSC using the munge_sumstats tool from

https://github.com/belowlab/ldsc/tree/2-to-3, and LD scores were calculated for each organ-

specific TLand score and trait combination using the ldsc tool from

https://github.com/bulik/ldsc.

17 Target organs annotations for GWAS traits

18 Gold-standard target-organ associations for 44 GWAS traits were annotated with Open

19 Targets (https://platform.opentargets.org/), EMBL-EBI Ontology Lookup Service

20 (https://www.ebi.ac.uk/ols/index), and by reviewing literature for selected GWAS traits

21 (Supplementary Table 3). These annotations were used to evaluate model performance for

22 each GWAS trait.

23 Prioritization of relevant organs for GWAS traits

We predicted approximately 2 million GWAS variants and strongly linked SNPs across 51

organs. For each manually annotated GWAS trait, we selected all associated SNPs. For

each organ, we calculated the p-value from a one-tailed Student's t-test (35) comparing the

sampled organ-specific SNP scores versus the population distribution, defined as the 2

million SNP score distribution for the organ. We ranked organs by the inverse of their p-

values, using a significance threshold of 0.05 to select only organs with statistical support for

a functional association. We evaluated performance by comparing the ranked (i.e.,

prioritized) organs with the manually annotated target organs for each trait. Accuracy was

defined as the fraction of correct overlaps between model-based and gold-standard organ

annotations among the top 4 organs for each model. We combined any two models' ranking

lists by taking the union of their lists then ranked by p-value, and selected the top 5 organs

for each combination.

GWAS trait enrichment

We began by gathering the top-scoring GWAS variants, specifically those with organ-specific

scores exceeding 0.5 for each organ model. We traced back from these variants to identify

their associated traits. For each trait, we defined and computed the trait score as the sum of

the organ-specific scores of the associated variants. To refine our results for each organ, we

excluded traits with low total counts, where total counts represent the sum of trait counts for

all associated GWAS and strongly-linked SNPs. Finally, we calculated organ-specific GWAS

trait enrichment scores by normalizing the trait score to the total counts.

Stratified LD Score Regression (S-LDSC)

S-LDSC (36) partitions complex trait heritability by genomic and functional annotations and

is used to help identify which annotations are most relevant to the genetic architecture of a

given trait. It estimates per-SNP heritability by regressing GWAS test statistics on LD scores

that are stratified by these annotations.

The expected GWAS χ^2 statistic of SNP j is modeled as

$$\mathbb{E}\left[\chi_j^2\right] = N \sum_C \tau_C \ell(j, C) + 1 \tag{1}$$

1 where N is the sample size, C is the set of annotations indexed by C (referred to as

2 categories in the S-LDSC paper), τ_C is the per-SNP contribution to heritability by annotation

3 C, and $\ell(j, C)$ is the LD score of SNP j with respect to annotation C.

4 Enrichment of an annotation for a given trait is calculated using

$$Enrichment_C = \frac{\tau_C}{\sum_C \tau_C / M} \tag{2}$$

where M is the total number of SNPs used in the computation. This is interpreted as the

proportion of SNP heritability contributed by an annotation divided by the proportion of SNPs

in that annotation.

TLand scores for the 51 organs were calculated for all HapMap 3 SNPs (37) excluding those

within the major histocompatibility complex (MHC) region of the genome (1,217,311 SNPs).

TLand scores were binarized at a threshold of 0.7, covering an average of 1% of SNPs

across organs. We performed S-LDSC (https://github.com/bulik/ldsc) for each of the 94

UKBB traits conditioned on 97 baseline-LD (v2.2) model annotations + one TLand organ-

specific annotation. We ran this for all trait-organ combinations, resulting in heritability

enrichment estimates and standard errors for each combination. An organ was considered

prioritized for a given trait if the lower bound of the heritability enrichment estimate's 95%

confidence interval was greater than 0. Specifically, if the contribution of the organ-specific

TLand annotation to the trait's narrow-sense heritability (h^2) is significantly greater than the

proportion of SNPs covered by the annotation.

Chromatin accessibility QTL evaluation and organ prioritization

We downloaded the cluster-specific chromatin accessibility QTL (caQTL) results from Wenz

et al. (38) (https://zenodo.org/records/12706263) and obtained coordinates for all tested

peaks from personal correspondence with the authors. Cluster 9 was chosen for this

evaluation due to its homogenous compositions of biosamples relative to other clusters and

a large number of caQTL calls to maximize power for statistical testing. To reduce the

number of false positive caQTL calls, we filtered for tested variants within 100bp of their

corresponding peak midpoint, hereby referred to as the evaluation set of variants. Positive

caQTL calls were defined as variants from this set passing a false discovery rate (FDR) of

0.05. We generated TLand scores and Sei scores for the evaluation sets and calculated the

corresponding AUPRC and AUROC metrics with these labels.

Organ prioritization of the cluster-specific caQTLs was done by performing one-sided

Welch's t-tests for each of the 51 organ-specific scores against the caQTL calls. Multiple test

correction was applied using the Benjamini–Yekutieli method to control for FDR. The

resulting q-values for each organ-specific score were log-transformed and ranked in

descending order. The top significant organs represent the most relevant organs for a set of

cluster-specific caQTLs.

Results

TLand incorporates comprehensive datasets to predict regulatory

variants

We developed a new model architecture, TLand, to predict regulatory variants from a

comprehensive set of features derived from RegulomeDB (Fig 1b and Supplementary Fig 1).

TLand takes input as genomic positions or dbSNP IDs, queries RegulomeDB for features,

and then outputs the probability that each variant is a cell-type specific (Supplementary Fig.

1) or organ-specific (Fig. 1b) regulatory variant. Importantly, by combining stacked

generalization, feature grouping, and interaction terms in meta-classifier training, the TLand

model can incorporate growing training datasets and novel data types into its feature space

while combating overfitting, whereas its predecessors cannot.

The continuously growing corpus of genomic data captures an increasingly complete view of

human regulatory variation. RegulomeDB recently upgraded to version 2, expanding to >650

million and >1.5 billion genomic intervals in hg19 and GRCh38, respectively (11). The large

discrepancy in the data availability between the two assemblies, for example, in ChIP-seq

and open chromatin data (RegulomeDB TF ChIP-seq availability in Fig 1a, open chromatin

in Supplementary Fig 3, histone ChIP-seq in Supplementary Fig 4), results from better

representation of complex variation and correction of sequencing artifacts in the GRCh38

assembly (12). Features used to predict regulatory variants included experimental and

computational features derived in GRCh38, the majority of which were derived from

Figure 1. TLand improves regulatory variant predictions. (a) TF ChIP-seq data availability across organs on RegulomeDB v2. Green bar plots represent counts on GRCh38. Orange bar plots represent counts on hg19. The total number of counts for each assembly is in the middle of the gray box. Notice that the summation is not simply adding all numbers together due to some cell lines having multiple corresponding organs. (b) Organ-specific TLand architecture. Organ-specific TLand was trained to predict human regulatory variants in an organ-specific manner by using RegulomeDB-derived features. (c) Benchmarking TLand performance by AUROC and AUPR. The X-axis is holdout cell lines or organs. The Y-axis is AUPR on the top panel and AUROC on the bottom panel.

RegulomeDB v2 (Supplementary Table 1). 1,372 DNase-seq BigWig files were quantile-

normalized and used to derive five quantile features for each genomic locus. The DeepSEA

disease score feature (31) was replaced by its successor Sei (32). The Sei model

simultaneously predicted 21,907 binary assay labels, which were dimension-reduced to 40

features representing sequence classes. Assembling deep learning model predictions and

training with comprehensive features reduced the variance of our final models and improved

generalization to new cell lines and less-studied organs (39,40).

Training data from six common cell lines were concatenated to train an agnostic model,

while the specificities of input features determined the cell- or organ-specificity of model

predictions. Features were bagged into three biologically meaningful subspaces before each

set was fed into an individual base classifier. Subspaces included a generic experimental

feature set, a computational feature set, and a cell/organ-specific experimental feature set

for regulatory variants.

We adapted the stacked generalization algorithm to stack the output of individual classifiers

and use a meta-classifier to compute the final prediction (19). Stacking enables the use of

each individual classifier's strengths by using their outputs as inputs to a final meta-classifier.

Cross-validation is required to prevent information leakage during meta-classifier training.

We made several modifications to the algorithm. We used group cross-validation to make

base classifiers to learn the regulatory function as the conditional probability between

generic features and cell/organ-specific features. Interaction terms were calculated prior to

their inclusion in the meta-classifier. We used probability rather than binary classification in

the first layer of base classifiers to train the meta-classifier, achieving higher accuracy (41).

TLand improves regulatory variant predictions

Cell-specific TLand models substantially outperformed state-of-the-art models in predicting

unseen cell-line regulatory variants (Fig. 1c and Supplementary Fig. 5). On average, across

holdout cell lines, cell-specific TLand models outperformed the previous best model (TURF)

with the area under the precision-recall curve (AUPR) and the area under the receiver

operating characteristic (AUROC) increasing from 0.389 to 0.471, and 0.729 to 0.776,

respectively. While GenoNet outperformed TURF in three held-out cell lines, the observed

gains were predominantly attributed to information leakage. This is because the GenoNet

models for held-out cell lines, which were initially trained and tested within that held-out cell

line, were directly used in the benchmarking. GenoNet performance was used to evaluate

the upper limit of the held-out prediction task. Cell-specific TLand models still outperformed

GenoNet, improving the average AUPR by 0.09 and the AUROC by 0.04. The high

performance of cell-specific TLand can be explained in two ways. First, we derived

comprehensive feature sets for predicting regulatory variants from RegulomeDB,

categorizing them into experimental feature sets and a complementary non-correlated

computational feature set (i.e., deep learning feature set) (Supplementary Fig. 7). The

inclusion of experimental features enabled a ~30% performance gain compared to DeepSEA

and Sei models, which use only deep learning features, with average TLand average

AUPR=0.471 compared to AUPR=0.336 for DeepSea and AUPR=0.364 for Sei. Second,

TLand's architecture models different biological domains separately and then calculates their

conditional probabilities. The meta-classifier finds the best combination of these probabilities

and subtracts redundant information through the inclusion of negative coefficients

(Supplementary Fig 8 and Supplementary Fig 9) to cancel out noise, thus improving

predictions, yielding improvements of 0.10 in average AUPR and 0.04 in average AUROC

when compared to TURF (Supplementary Fig 10).

Organ-specific TLand models address data availability bias

TLand's flexible architecture enables substituting cell-specific features with organ-specific

features, which can then be retrained with organ-specific labels to develop organ-specific

TLand models. Organ-specific TLand models can predict organ-specific regulatory variants,

even when the organ is represented by only less-studied cell lines or tissues. These models

performed better than their cell-specific counterparts in two out of four cell-type specific

tasks, specifically in HepG2 (AUPR 0.547 vs. 0.514, AUROC 0.781 vs. 0.692) and MCF-7

(AUPR 0.512 vs. 0.512, AUROC 0.879 vs. 0.840). However, because the GM12878 and

K562 cell lines have conflicting labels for the same organ (blood), the blood organ-specific

model was evaluated only in the organ-specific (i.e., heart) task, not in either of the cell-line-

specific tasks. The superior performance of organ-specific models, even in hold-out cell line

tasks, can be attributed to the fact that cell-type specific regulatory variants in HepG2 and

MCF-7 are well-represented by corresponding organ-specific regulatory variants within the

RegulomeDB database (see Supplementary Fig 11). In the A549 cell line, which is from the

least-represented organ (lung), the organ-specific TLand model underperformed the cell-

specific TLand model holdout on A549 (AUPR 0.329 vs. 0.347), as the organ model

predicted regulatory variants in the lung that were not observed in the A549 cell line. To

better evaluate model performance in the lung, the ASB dataset from the second-most

representative lung cell line, IMR-90, was added to the holdout dataset. The inclusion of this

dataset led to TLand outperforming the cell-specific model when held out on the lung with

the IMR-90 dataset (AUPR 0.483 vs. 0.432; AUROC 0.841 vs. 0.814), indicating that the

organ-specific TLand model can predict a comprehensive set of organ-specific regulatory

variants.

However, adding the second most representative cell line in the embryo, H9, did not improve

the TLand organ-specific model when compared to the cell-specific model when holding out

on the embryo organ (AUPR 0.536 vs. 0.597). Deep learning features, such as DeepSEA

disease impact score (31) and Sei sequence classes (32), were more representative of the

cell lines or organs with more data availability. This is because the scores were

dimensionally reduced from 919 and 21,907 prediction tasks of experimental assays for

DeepSEA and Sei, respectively (including TF and histone ChIP-seq and open chromatin

assays). Cell lines and organs with more available data dominated the dimension-reduced

results. Thus, we removed deep learning features and developed the organ-specific TLand

light model (Supplementary Fig 12) to predict regulatory variants in organs with low data

availability, such as the embryo. We defined organs with >=100 available ChIP-seq assays

as "high data availability" and those with <100 assays as "low data availability". We observed

that the TLand (full) model consistently outperformed the TLand light model in organs with

high data availability (Fig. 1a and c; Supplementary Fig. 5), whereas the light model

outperformed the full model in organs with low data availability. For example, the organ-

specific TLand light model performed best when holding out the embryo organ (AUPR 0.639,

AUROC 0.774). Those findings indicate that TLand and light models are suitable for

predicting regulatory variants in organs with low data availability, whereas TLand (full)

models are more suitable for organs with high data availability. We then proceeded with our

analysis by training the TLand and TLand light models on all available data. We trained an

additional TLand model, TLand lightest, in which the organ-specific ChIP-seq features were

removed to further reduce bias toward overrepresented organs (benchmarking results

shown in Supplementary Table 2).

We also evaluated the TLand models using leave-one-chromosome-out cross-validation

(LOCO CV) across aggregated cell line data to ensure that data leakage via Sei features

was not the source of TLand's improved performance (Supplementary Fig 6). The Sei

features depend on a 4096-bp DNA sequence as input, with the variant positioned at the

center. This allows for train and test set variants to potentially overlap the same input

sequence, resulting in data leakage between the train and test sets. This potential data

leakage is relevant only to the full TLand models, as the light and lightest models do not use

Sei features. We show that the TLand still outperforms other models using LOCO CV, and

that organ-specific models substantially outperform the cell-specific model. The complete

organ-specific model performs best, consistent with the held-out cell line results, as the

blood and liver variants are included in this cell line-aggregated dataset.

Finally, we evaluated our final models with independent MPRA datasets. We observed

similar trends, with TLand models outperforming alternative methods (Supplementary Fig.

13). However, MPRA evaluation can only test for a subset of our predicted variants since

MPRA variants affect gene expression. In contrast, TLand models predict variants that have

any regulatory effects (defined as altering TF binding), which may not affect gene

expression.

TLand prioritizes relevant organs for GWAS traits

To systematically evaluate organ-specific TLand models, we predicted approximately 2

million GWAS SNPs, including those within the same LD blocks, across 51 organs defined

by ENCODE (12). We found that TLand predictions were more correlated with TLand light

predictions than TLand lightest (e.g., in the heart organ in Supplementary Fig. 14). We

manually curated a list of GWAS traits with relevant organs. TLand, TLand light, and TLand

lightest models prioritized the relevant organs for 44 GWAS traits (Supplementary Table 3),

sourced from the GWAS catalog with an average accuracy of 0.311, 0.340, and 0.466,

respectively (Fig. 2b., see the definition of accuracy in Methods). By integrating prioritized

organs from two distinct models, TLand and TLand lightest, the average accuracy was

increased to 0.482. The overall low accuracy was partially due to uncertainty about whether

low scores reflect low data availability or indicate true negatives, with the target organ

lacking regulatory function.

To better evaluate TLand models in organs with limited data availability, we focused on the

top-scoring variants in those organs and asked whether the top-scoring variants identified by

those models were biologically meaningful. We found that top-scoring variants from organ-

specific models were enriched for GWAS traits associated with their target organs (Methods,

Fig. 2a, and results in Supplementary Table 4). For example, two of the three most-enriched

GWAS traits identified by the TLand lightest model were relevant to the heart organ: atrial

fibrillation and resting heart (Fig 2a). Originally, TLand lung models were unable to prioritize

the lung organ given the corresponding GWAS SNPs for relevant traits (Supplementary

Table 3). However, we found that traits such as physical activity measurement and peak

expiratory flow were enriched by top-scored variants in the TLand model. Traits that were

associated with multiple organs were properly identified by multiple-organ models. For

example, both the bone element and the immune TLand lightest models pinpointed

ankylosing spondylitis, a type of inflammatory arthritis affecting the spine and large joints

(42), as one of the most enriched GWAS traits for their top-scored variants (Fig 2a).

We also explore if organ-specific TLand scores are enriched for heritability in 94 GWAS

traits from the UK Biobank. This is a well-powered cohort, allowing us to utilize the stratified

LD score regression (S-LDSC) method to estimate the contribution of TLand scores to each

trait's narrow-sense heritability (Methods). We evaluated distributions of TLand scores at the

50, 90, 95, and 99 percentiles across all organs to identify an appropriate score threshold

(Fig. 2c). The top 50% of scores are largely between 0.1 and 0.2. The top 5% of scores span

a very wide range, from 0.2 to 0.7. The top 1% of scores contains a reasonable distribution

of high-scoring variants between 0.6 and 0.8, with the median around 0.7. We thus used a

threshold of 0.7 to binarize TLand scores for use in S-LDSC. We computed enrichment

scores for each trait-organ combination and identified significantly enriched organs for 71

traits (Methods, Supplementary Table 6 and Supplementary Table 7). Traits were enriched

for a median of 3 organs, with stage 4 balding having the highest number of enriched organs

at 17. We see that traits largely decompose into their relevant organs. Lymph node and

thyroid gland TLand scores are enriched in hypothyroidism, and body musculature and heart

TLand scores are enriched in atrial fibrillation (Figure 2d). Liver TLand scores are also

expectedly enriched in HbA1c levels and type 2 diabetes (T2D), but stomach TLand scores

are too. Gastroparesis - the delayed movement of food from the stomach - is a common

complication of T2D (43) and is hypothesized to be due to vagus nerve damage from T2D-

induced hyperglycemia. Still, the cause is ultimately unclear (44). There is also limited

research on the genetic factors of gastroparesis. The enrichment of stomach TLand scores

in T2D could suggest a pleiotropic model in which T2D-associated variants affect both T2D

and stomach traits, explaining the frequent co-occurrence of T2D and gastroparesis. Overall,

we show that organ-specific TLand models can accurately prioritize relevant organs for

GWAS traits and be used to propose hypotheses for complex genetic risk factors.

Figure 2. TLand prioritizes relevant organs for GWAS traits. (a) Prioritization of GWAS traits given

top-scored variants by TLand and TLand lightest models. TF ChIP-seq data availability plot is re-

plotted in the top left where green represents high data availability and TLand was the model used,

and blue represents low data availability, and TLand lightest was the model used. Each circle next to

the organs indicates the data availability of each organ. The colors of enrichment bars correspond to

the model used. (b) Performance of TLand models prioritizing organs. X-axis are TLand models. Y-

axis is accuracy (definition see Methods). (c) Distribution of TLand scores using 50, 75, 90, 95, and

99% quantile thresholds across organs. (d) Significant S-LDSC enrichment results using TLand

scores for atrial fibrillation (AFib), mean corpuscular volume (MCV), hemoglobin A1C (HbA1c),

hypothyroidism, uric acid (UA), and type 2 diabetes (T2D). Error bars denote 95% confidence

intervals.

TLand partitions caQTLs into relevant organs

We evaluated TLand on chromatin accessibility QTL (caQTL) prediction and ability to

recover sample tissue identities from cluster-specific ATAC-seq datasets. Wenz et al.

collected 10,293 publicly available ATAC-seq datasets, performed genotyping directly from

the ATAC-seq reads, and called caQTLs both globally and on clustered samples (38) (Figure

3a). Clustered samples generally reflected similar tissue types or treatment/disease

conditions. We calculated organ-specific TLand scores for these cluster-specific caQTLs and

limited the evaluations to tested variants within 100bp of their corresponding peak midpoints

to minimize false-positive caQTL calls and ensure that TLand was differentiating between

likely causal vs non-causal variants instead of simply predicting presence/absence of peaks

at variant positions. TLand blood scores clearly separate non-caQTLs and caQTLs in cluster

9 (Figure 3b), which is largely composed of lymphoblastoid cell lines, and outperform Sei in

classifying caQTLs in this cluster (Figures 3c and 3d). For each of the 51 organ-specific

TLand scores, we tested if the mean organ-specific score of caQTLs was greater than the

mean organ-specific score of non-caQTLs using a one-sided Welch's t-test. After performing

multiple test correction, we ranked the resulting q-values for each of the tests (where each

test corresponds to an organ) in order to prioritize organs associated with the set of caQTLs.

Because of shared ATAC-seq peaks and variant effects across organs, almost all tests yield

highly significant q-values; however, we expect the most relevant organs to be ranked

highly. Blood was the top-ranked organ, and the other highly ranked organs (bone marrow,

bone element, immune organ, lymph node, thymus) are also largely composed of immune

cell types (Figure 3e). We show that TLand can also be used to prioritize variant effects on

chromatin accessibility and partition caQTLs into relevant organs, allowing researchers to

impute potential cross-tissue effects.

Figure 3. TLand partitions caQTLs into relevant organs. (a) Diagram of organ prioritization of

caQTLs. Public ATAC-seq datasets were collected and uniformly processed through a genotyping

and caQTL calling pipeline by Wenz et al. They clustered the datasets based on accessibility profiles,

and cluster-specific caQTLs were called. We filtered these caQTLs and tested high-confidence

caQTLs against high-confidence negatives using organ-specific TLand scores. Multiple test correction

was performed, and organs were ranked by q-value. (b) Distribution of TLand blood scores for non-

caQTL variants and caQTL variants in cluster 9. (c) AUPRC and (d) AUROC of TLand blood and Sei

scores for classifying caQTL variants from (b). (e) Top 10 prioritized organs for cluster 9 caQTLs

ranked by q-value.

Discussion

The identification and characterization of non-coding regulatory variants in the human

genome remain a major challenge in genomics. We have developed a novel model

architecture, TLand, to accompany the recent advancement of RegulomeDB. TLand uses

RegulomeDB-derived features to infer regulatory variants at the cell- and organ-specific

levels. By applying stacked generalization to incorporate comprehensive datasets on the

GRCh38 reference genome, including experimental and computational features, TLand

models consistently outperformed state-of-the-art models in holdout benchmarking. This

demonstrates the TLand models' ability to generalize to new cell lines or

organs, a major advancement over the SURF and TURF models they extend. Much of this

improvement can be attributed to improved handling of differential data availability. In

particular, TLand light and lightest models are applicable to organs with limited data, e.g.,

the embryo. In contrast, current leading classification models, such as DeepSEA's disease

impact scores, perform poorly in such organs.

One of TLand's main strengths is that it covers nearly all human tissue types, enabling

hypothesizing pleiotropic effects via GWAS-trait heritability enrichment or caQTL organ

prioritization. Many existing variant-effect models either capture global, tissue-agnostic

pathogenicity or are limited to a few tissue types with maximal data modality coverage. Our

generalization approach, based on careful feature bagging and training separate models,

enables us to investigate variant effects while accounting for the vast majority of biological

contexts.

The TLand models presented here represent a promising foundation for further

improvements. These are facilitated mainly by TLand's ability to ingest new datasets as they

are released. We aim to continuously incorporate new datasets into RegulomeDB as they

are released, thus extending its performance and applicability as the number and diversity of

training datasets grow across different cells, tissues, and organs. Many of these, including

gkm-SVM model predictions (45) and Hi-C contact maps (17), are new features for TLand

models. TLand's flexible, modular design allows grouping features into biologically

meaningful sets that can be evaluated against one another before making the final decision

about whether to include novel features in the model.

One limitation of this study is the limited number of allele-specific binding (ASB) sites in our training dataset. Because of data availability constraints, ASB training datasets were derived from the personal genomes of only six common cell lines. In addition, there is an extensive disagreement among existing ASB prediction methods, further reducing the number of high-confidence training variants. Recently, Adastra, a new database that hosts 652,595 ASBs passing 5% FDR across 647 cell lines and 1,043 TFs, was published (21). However, Adastra predicts ASB using a statistical model, not by using personal genomes directly. Therefore, it is unclear whether incorporating these ASB predictions in the high-confidence training datasets will yield meaningful improvements to the TLand models. This remains an area of ongoing interest.

Conclusions

Overall, TLand advances the field of non-coding variant analysis by incorporating comprehensive datasets to predict functional regulatory variants at cell-specific and organ-specific levels. To foster downstream applications, we have provided pre-calculated TLand scores for 30 million genomic variants through the IGVF data portal (See Availability of data and materials section). These scores are being used by the IGVF consortium to prioritize disease and trait-associated variants for functional assays and benchmarking studies. TLand's flexible models and ability to integrate novel datasets in a principled way significantly improve our ability to identify human regulatory variants genome-wide. This ability will only improve as larger, more comprehensive training datasets spanning additional organs, tissues, cell types, and assays are released.

List of abbreviations

SNPs: Single-nucleotide polymorphisms

GWAS: Genome-wide association study

ASB: Allele-specific binding

AUROC: Area Under the Receiver Operating Characteristic Curve

AUPR/AUPRC: Area Under the Precision Recall Curve

MPRA: Massively parallel reporter assay

FDR: False discovery rate

LD: Linkage disequilibrium

MHC: Major histocompatibility complex

caQTL: Chromatin accessibility quantitative trait locus

TF: Transcription factor

LOCO CV: Leave-one-chromsome-out cross-validation

S-LDSC: Stratified linkage disequilibrium score

T2D: Type 2 diabetes

AFib: Atrial fibrillation

MCV: Mean corpuscular volume

HbA1c: Hemoglobin A1c

UA: Uric acid

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Scripts to reproduce the analyses can be found at

https://github.com/rnsherpa/tland_supplementary. A Snakemake workflow to run TLand

predictions on input variants is available at https://github.com/rnsherpa/TLand-predict.

The datasets supporting the conclusions of this article are available on Zenodo at

https://doi.org/10.5281/zenodo.18341395. TLand predictions for 30 million variants are

available on the IGVF data portal at https://data.igvf.org/prediction-sets/IGVFDS0038SURQ/.

caQTL data from Wenz et al. are available on Zenodo at

https://doi.org/10.5281/zenodo.12706262.

Competing interests

The authors declare that they have no competing interests.

Funding

This work is supported by NIH grants U01HG011952 and U24HG009293

Authors' contributions

NZ and SD developed the model. RNS developed the Snakemake workflow. NZ, RNS, and

KBH performed the model evaluations and downstream analyses. NZ, RNS, and APB

contributed to interpretation of the results. NZ, RNS, and APB wrote the manuscript with

input from all authors. APB supervised the work.

Acknowledgements

We thank Christopher Castro, Benjamin Hitz, Shengcheng Dong, and Zhewei Shen for their

assistance in data submission to the IGVF data portal.

References

1. 1. Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, et al. Clinical whole-exome sequencing for the diagnosis of mendelian disorders. N Engl J Med. 2013 Oct 17;369(16):1502–11.

2. 2. Boyle AP, Davis S, Shulha HP, Meltzer P, Margulies EH, Weng Z, et al. High-resolution mapping and characterization of open chromatin across the genome. Cell. 2008 Jan 25;132(2):311–22.

3. 3. Robertson G, Hirst M, Bainbridge M, Bilenky M, Zhao Y, Zeng T, et al. Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. Nat Methods. 2007 Aug;4(8):651–7.

4. 4. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat Methods. 2013 Dec;10(12):1213–8.

5. 5. Worthey EA, Mayer AN, Syverson GD, Helbling D, Bonacci BB, Decker B, et al. Making a definitive diagnosis: Successful clinical application of whole exome sequencing in a child with intractable inflammatory bowel disease. Genet Med. 2011 Mar 1;13(3):255–62.

6. 6. Ng SB, Buckingham KJ, Lee C, Bigham AW, Tabor HK, Dent KM, et al. Exome sequencing identifies the cause of a mendelian disorder. Nat Genet. 2010 Jan;42(1):30–5.

7. 7. International HapMap Consortium. A haplotype map of the human genome. Nature. 2005 Oct 27;437(7063):1299–320.

8. 8. International HapMap Consortium, Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, et al. A second generation human haplotype map of over 3.1 million SNPs. Nature. 2007 Oct 18;449(7164):851–61.

9. 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A global reference for human genetic variation. Nature. 2015 Oct 1;526(7571):68–74.

10. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, Kasowski M, et al. Annotation of functional variation in personal genomes using RegulomeDB. Genome Res. 2012 Sep;22(9):1790–7.

11. Dong S, Zhao N, Spragins E, Kagda MS, Li M, Assis P, et al. Annotating and prioritizing human non-coding variants with RegulomeDB v.2. Nat Genet. 2023 Apr 25;1–3.

12. ENCODE Project Consortium, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N, et al. Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature. 2020 Jul;583(7818):699–710.

13. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. Nature. 2015 Feb 19;518(7539):317–30.

14. Dong S, Boyle AP. Predicting functional variants in enhancer and promoter elements using RegulomeDB. Hum Mutat. 2019 Sep;40(9):1292–8.

15. Dong S, Boyle AP. Prioritization of regulatory variants with tissue-specific function in the non-coding regions of human genome. Nucleic Acids Res. 2022 Jan 11;50(1):e6.

16. Pazin M, Gilchrist DA, Morris SA. Genome.gov. [cited 2023 Apr 2]. Impact of Genomic Variation on function (IGVF) Consortium. Available from: https://www.genome.gov/Funded-Programs-Projects/Impact-of-Genomic-Variation-on-Function-Consortium

17. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, et al. Comprehensive mapping of long-range interactions reveals folding principles of the

human genome. Science. 2009 Oct 9;326(5950):289–93.

18. Avsec Ž, Agarwal V, Visentin D, Ledsam JR, Grabska-Barwinska A, Taylor KR, et al. Effective gene expression prediction from sequence by integrating long-range interactions. Nat Methods. 2021 Oct;18(10):1196–203.

19. Wolpert DH. Stacked generalization. Neural Netw. 1992 Jan 1;5(2):241–59.

20. Chen J, Rozowsky J, Galeev TR, Harmanci A, Kitchen R, Bedford J, et al. A uniform survey of allele-specific binding and expression over 1000-Genomes-Project individuals. Nat Commun. 2016 Apr 18;7:11101.

21. Abramov S, Boytsov A, Bykova D, Penzar DD, Yevshin I, Kolmykov SK, et al. Landscape of allele-specific transcription factor binding in the human genome. Nat Commun. 2021 May 12;12(1):2751.

22. Ke G, Meng Q, Finley T, Wang T, Chen W, Ma W, et al. LightGBM: A highly efficient gradient boosting decision tree. Adv Neural Inf Process Syst [Internet]. 2017 [cited 2023 Feb 21];30. Available from: https://proceedings.neurips.cc/paper/2017/hash/6449f44a102fde848669bdd9eb6b76fa-Abstract.html

23. Ho TK. Random decision forests. In: Proceedings of 3rd International Conference on Document Analysis and Recognition. 1995. p. 278–82 vol.1.

24. McCulloch WS, Pitts W. A logical calculus of the ideas immanent in nervous activity. Bull Math Biophys. 1943 Dec 1;5(4):115–33.

25. Akiba T, Sano S, Yanase T, Ohta T, Koyama M. Optuna: A Next-generation Hyperparameter Optimization Framework. In: Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining. New York, NY, USA: Association for Computing Machinery; 2019. p. 2623–31. (KDD '19).

26. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine Learning in Python [Internet]. arXiv [cs.LG]. 2012 [cited 2023 Feb 21]. p. 2825–30. Available from: https://www.jmlr.org/papers/volume12/pedregosa11a/pedregosa11a.pdf?ref=https:/

27. Raschka S. MLxtend: Providing machine learning and data science utilities and extensions to Python's scientific computing stack. J Open Source Softw. 2018 Apr 22;3(24):638.

28. Kent WJ, Zweig AS, Barber G, Hinrichs AS, Karolchik D. BigWig and BigBed: enabling browsing of large distributed datasets. Bioinformatics. 2010 Sep 1;26(17):2204–7.

29. Neph S, Scott Kuehn M, Reynolds AP, Haugen E, Thurman RE, Johnson AK, et al. BEDOPS: high-performance genomic feature operations [Internet]. Vol. 28, Bioinformatics. 2012. p. 1919–20. Available from: http://dx.doi.org/10.1093/bioinformatics/bts277

30. van der Sande M, van Heeringen S. qnorm [Internet]. 2021. Available from: https://github.com/Maarten-vd-Sande/qnorm

31. Zhou J, Troyanskaya OG. Predicting effects of noncoding variants with deep learning–based sequence model. Nat Methods. 2015 Aug 24;12(10):931–4.

32. Chen KM, Wong AK, Troyanskaya OG, Zhou J. A sequence-based global map of regulatory activity for deciphering human genetics. Nat Genet. 2022 Jul;54(7):940–9.

33. Cerezo M, Sollis E, Ji Y, Lewis E, Abid A, Bircan KO, et al. The NHGRI-EBI GWAS Catalog: standards for reusability, sustainability and diversity. Nucleic Acids Res. 2025 Jan 6;53(D1):D998–1005.

34. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of

middle and old age. PLoS Med. 2015 Mar 31;12(3):e1001779.

35. Student. The probable error of a mean. Biometrika. 1908;6(1):1–25.

36. Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh PR, et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat Genet. 2015 Nov 28;47(11):1228–35.

37. International HapMap 3 Consortium, Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA, et al. Integrating common and rare genetic variation in diverse human populations. Nature. 2010 Sep 2;467(7311):52–8.

38. Wenz BM, He Y, Chen NC, Pickrell JK, Li JH, Dudek MF, et al. Genotype inference from aggregated chromatin accessibility data reveals genetic regulatory mechanisms. Genome Biol. 2025 Mar 30;26(1):81.

39. Dey KK, van de Geijn B, Kim SS, Hormozdiari F, Kelley DR, Price AL. Evaluating the informativeness of deep learning annotations for human complex diseases. Nat Commun. 2020 Sep 17;11(1):4703.

40. Dey KK, Kim SS, Gazal S, Nasser J, Engreitz JM, Price AL. Integrative approaches to improve the informativeness of deep learning models for human complex diseases [Internet]. bioRxiv. 2021 [cited 2022 Dec 13]. p. 2020.09.08.288563. Available from: https://www.biorxiv.org/content/10.1101/2020.09.08.288563

41. Ting KM, Witten IH. Issues in Stacked Generalization. jair. 1999 May 1;10:271–89.

42. Zhu W, He X, Cheng K, Zhang L, Chen D, Wang X, et al. Ankylosing spondylitis: etiology, pathogenesis, and treatments. Bone Res. 2019 Aug 5;7:22.

43. Lee AA, Hasler WL. Diabetes and the stomach. Curr Treat Options Gastroenterol. 2017 Dec 6;15(4):441–59.

44. Bharucha AE, Kudva YC, Prichard DO. Diabetic gastroparesis. Endocr Rev. 2019 Oct 1;40(5):1318–52.

45. Lee D, Gorkin DU, Baker M, Strober BJ, Asoni AL, McCallion AS, et al. A method to predict the impact of regulatory variants from DNA sequence. Nat Genet. 2015 Aug;47(8):955–61.