Title

Children exhibit a developmental advantage in the offline processing of a learned motor sequence

Authors

Anke Van Roy[1], Geneviève Albouy[1], Ryan D. Burns[1], Bradley R. King[1, *]

Affiliation

[1] Department of Health and Kinesiology, College of Health, University of Utah, Salt Lake City,

UT 84112, USA

[*] **Corresponding author**

Bradley King, PhD

Department of Health & Kinesiology, College of Health, University of Utah

250 S 1850 E

Salt Lake City, Utah 84112

Bradley.ross.king@utah.edu

Abstract

Research that examines changes in a specific behavior across the lifespan frequently reports an

inverted-U trajectory. That is, young adults exhibit optimal performance, children are

conceptualized as developing systems progressing towards this ideal state, and older adulthood is

characterized by performance decrements. However, not all behaviors follow this trajectory, as

there are instances in which children outperform young adults. This study aimed to reveal instances

of childhood advantage in the domain of motor learning and memory consolidation. Using a

visually cued serial reaction time task, data were acquired from 7-35 and >55-year-old participants

(n=238 datasets) across two experiments that assessed initial encoding as well as memory

consolidation over 5-hour and 24-hour offline periods. Results revealed comparable initial learning

dynamics among age groups, but 7- to 12-year-old children exhibited smaller sequence-specific

learning relative to adolescents (13-17 year-olds), young adults (18-35 year-olds) and older adults

(> 55 years). Interestingly, children demonstrated the greatest performance gains in the 5-hour and

24-hour delayed retests, reflecting enhanced motor memory consolidation. These results suggest

that children exhibit an advantage in the offline processing of recently learned motor sequences.

1. Introduction

The ability to learn novel movements is vital to the everyday functioning of individuals across the human lifespan (e.g., from a child learning to ride a bike to an older individual learning to operate a touchscreen). One of the dominant motor learning paradigms in the field is motor sequence learning (MSL), whereby participants acquire a novel series of interrelated actions (i.e., a motor sequence). MSL has been extensively studied with multiple task variants, including the serial reaction time task (SRTT; i.e., participants respond to visual cues presented in a repeating order) and the finger tapping task (i.e., participants repeatedly reproduce a known/shown sequence of finger movements in a self-initiated manner). There has been extensive previous research that has employed these MSL task variants in healthy young adults (~18-35 years), facilitating the development of a framework that characterizes the time course of sequence learning in this age group (see [1–4] for reviews). Typically, a fast online (i.e., during repeated practice) acquisition phase with substantial performance improvements occurs during an initial training session. This is followed by a slow offline (i.e., in the absence of active task practice) consolidation phase that spans hours during which the acquired memory trace is stabilized, strengthened and reorganized into a more robust form.

Although previous research has examined MSL in school-aged children (i.e., between 5 and 12 years), the results are quite heterogenous. While some studies have reported comparable learning within the initial practice session in children and adults[5–8], others have shown that children exhibit worse[9–13] or better learning[10,14–17]. There is a general consensus, however, that children exhibit impaired consolidation of a recently acquired motor sequence over offline periods that include sleep [e.g., [13,15,18], but see [8,19] for specific examples of sleep-related benefits). Specifically, whereas

post-learning sleep is known to boost offline consolidation processes in young adults [see [3] for a

review], such a beneficial effect is absent in children. Interestingly, there is some evidence from

sequential finger tapping task variants suggesting that this degraded consolidation over sleep may

be the by-product of enhanced or accelerated consolidation over the wake epochs shortly following

initial learning[9,14,20]. Specifically, children showed a rapid performance stabilization and

enhancement 15 minutes[9] and 1 hour[14] after learning, which were not observed in young adults.

Interestingly, this enhanced offline processing does not appear to be limited to the timescale of

hours following initial learning but is also observed on shorter timescales. Specifically, Du et al.[5]

employed a stepping version of a SRTT and found that young children (i.e., 6-year-olds) showed

larger performance improvements over the short rest intervals in between blocks of practice within

a session as compared to older children (i.e., 10-year-olds) and young adults. These results

collectively suggest that young children may exhibit a developmental advantage with respect to

the micro- (i.e., timescale of seconds between practice blocks) and macro- (i.e., timescale of hours

between practice sessions) offline processing of recently practiced motor sequences during wake

intervals. This childhood advantage could then be added to the list of examples in which children

outperform adults in specific learning and memory behaviors (for review see [21]).

At the other end of the lifespan, there is considerable evidence that older adults (> ~60 years)

exhibit comparable initial learning of a novel motor sequence as young adults[22–26]. And, whereas

children appear to have a developmental advantage in the offline processing of recently acquired

sequences, older adults are known to exhibit deficits. Specifically, the magnitude of offline gains

over macro-offline intervals that include sleep and/or wake is significantly less than those observed

in younger adults[22,24–28].

Although the existing literature suggests that the offline processing of recently acquired movement

sequences appears to be superior in children and then declines throughout adulthood, systematic

examinations into changes across the human lifespan are relatively limited (see [10,11,16,29] for

examples in the domain of implicit sequence learning). This study thus aimed to provide an

exhaustive characterization of motor learning and memory consolidation behaviors across the

lifespan (i.e., in groups of children, adolescents, young adults and older adults). Consistent with

previous research[5–8], we hypothesized that the initial learning of a motor sequence (i.e., within a

single training session) would be comparable across the 4 age groups. We expected children to

exhibit enhanced micro- and 5-hour macro-offline consolidation, indicative of a developmental

advantage in the offline processing of a recently acquired motor sequence across periods of

wakefulness. This developmental advantage in macro-offline consolidation was expected to be

absent in the 24-hour retest, as previous research has indicated impaired sleep-facilitated

consolidation in children[8,13,15,18]. Last, and consistent with earlier studies, older adults were

expected to demonstrate intact initial learning but impaired macro-offline consolidation across

both 5- and 24-hour intervals.

2. Methods and materials

The project consisted of two experimental protocols. Specifically, Experiments 1 and 2 examined

the effect of age on the initial acquisition of a movement sequence and the time course of motor

memory consolidation, respectively. Data collection and analysis plans were pre-registered via

Open Science Framework and can be accessed at https://doi.org/10.17605/OSF.IO/WBK9H and

https://doi.org/10.17605/OSF.IO/ZMJ75. Any additional analyses that were not included in the

pre-registrations are labelled in this text as exploratory. Note that 4 (out of 130) and 12 (out of

108) datasets were acquired prior to the pre-registrations of Experiments 1 and 2, respectively.

This was done to verify that keypresses and their timing were logged adequately, and that saved

data were complete. The processed data used for the results presented in this text as well as the

raw data are publicly available on Zenodo (https://zenodo.org/doi/10.5281/zenodo.8274118).

2.1 Participants

Healthy volunteers between 7-35 or 55-75 years old of all genders were recruited by

advertisements on relevant websites and research databases. Note that as our experiments were

conducted through a web-based data collection platform, recruitment was not constrained to a

specific geographical region. Individuals interested in participation were sent an online screening

questionnaire to assess eligibility, which was subsequently reviewed by a member of the research

team. Exclusion criteria were: 1) reported history of medical, neurological, psychological or

psychiatric conditions, 2) use of psychoactive or sleep-influencing medications, 3) indications of

abnormal or irregular sleep, 4) mobility limitations of the fingers or hands, or 5) considered a

professional typist or prior extensive training on a musical instrument requiring dexterous finger

movements (e.g., piano, guitar). All experimental procedures were approved by the local Ethics

Committee (IRB_00136894). Adult participants and parents of child participants gave informed

consent and participants below 18 years of age provided informed assent. Participants received

compensation for their participation.

A convenience sample of 224 individuals met the inclusion criteria and thus initiated participation

in an experimental protocol. They were divided into the following four age groups: children

(operationally defined as 7-12 years old), adolescents (13-17 years old), young adults (18-35 years

old) and older adults ($\geq$ 55 years old). Pre-registered analyses compared differences among these

age *groups.* Yet, to provide more fine-grained analyses of age-related trajectories of motor learning

and memory consolidation behaviors, we also conducted exploratory analyses with age as a

continuous variable (see below for details).

Sample size computations were conducted *a priori* and with the software G*Power[30]. For the

assessment of age-group differences in initial motor sequence learning (i.e., Experiment 1), and to

detect an effect size of f = 0.3 (based on the comparison of initial learning between 9-year-olds

and young adults in Adi-Japha et al.[9]), with an alpha of 0.05 and power of 0.80, the desired sample

size was 128 subjects (32 per group). Of the participants that initiated the motor learning protocol,

data from 7 participants were excluded from analyses due to a failure to comply to experimental

instructions (e.g., repeatedly pressing the same key; n = 3 children), missing data because of

software issues (n = 1 young adult), or a failure to correctly perform the motor task (i.e., statistical

outliers (> 3SD from group mean) on sequence accuracy as defined in the preregistration; n = 1

child, 1 young adult, 1 older adult). Excluded participants were replaced and thus the final sample

size for analyses consisted of 130 participants (see Table 1 for participant demographics).

For the assessment of age-group differences in macro-offline consolidation processes (i.e.,

Experiment 2), the detection of an effect size of f = 0.20 (slightly more conservative than Adi-

Japha et al.[9]), with an alpha of 0.05, a power of 0.80 and a correlation among repeated

measurements of 0.25, this experiment required a sample size of 108 (27 participants per group).

Of those participants that initiated the motor learning protocol, 21 individuals were excluded from

data analysis. Specifically, 17 participants failed to comply to experimental instructions (i.e., 4

children repeatedly pressed the same key and 5 children, 2 young adults and 6 older adults either

did not complete the protocol or did not adhere to the specific schedule of the experimental

sessions). One older adult was excluded due to inaccurate data because of software issues. And,

based on our preregistration, additional exclusions were due to a lack of performance

improvements across training (n = 1 young adult, 1 older adult) and an inability to correctly

perform the motor task (statistical outliers on sequence accuracy; n = 1 young adult). Similar to

above, excluded participants were replaced and thus the final sample for analyses consisted of 108

participants (see Table 2 for participant details). Data from the third session of one young adult

and the post-learning random data of one adolescent were missing; thus, these individuals were

excluded from the contrasts involving these specific task runs only.

Note that the procedures of Experiment 1 (assessing initial motor sequence learning) were identical

to the first session of Experiment 2 (assessing macro-offline consolidation). Accordingly, data

from a subset of early participants from Experiment 2 (n = 42; 16 children, 19 adolescents, 4 young

adults, 3 older adults) were included in the analyses of Experiment 1. Thus, these participants were

included in analyses for both experiments.

169   *Table 1: Participant demographics, and sleep and vigilance scores for each age group in Experiment 1*

Variable	Children	Adolescents	Young adults	Older adults	p-value
n	33	33	32	32	/
Female *(n)*	17	12	23	24	0.004*
Age (years)	10.2 (1.6)	15.3 (1.3)	26.3 (5.0)	64.4 (5.2)	/
M/E Preference	2.9 (0.9)	3.3 (0.8)	2.9 (1.0)	2.3 (0.8)	<0.001*
Time of testing (SD in min)	12:47 (172)	12:29 (133)	14:52 (175)	12:05 (169)	<0.001*
Sleep quality	5.1 (0.8)	5.0 (0.7)	4.8 (0.6)	4.8 (0.8)	0.12
Sleep duration (hours)	9.8 (0.9)	9.1 (1.4)	8.5 (1.0)	7.7 (1.1)	<0.001*
SSS score	2.0 (0.9)	2.0 (0.8)	2.0 (0.7)	1.5 (0.7)	0.031*

Numbers represent the mean, with standard deviation in parentheses. Morningness/Eveningness (M/E)
Preference was added as an exploratory variable to assess age-related differences in circadian preferences
and was defined on a 5-point Likert scale (i.e., 1 = extreme morning person, 5 = extreme evening person).
The average time of testing is specified in 24-hour time notation, with the SD in minutes. Sleep quality is
defined on a 6-point Likert scale (i.e., 1 = very bad, 6 = very good). SSS = Stanford Sleepiness Scale[31],
with higher numbers indicative of increased sleepiness. The p-values resulted from one-way ANOVAs
assessing group differences. See Appendix 1 as part of the Supporting Information for full statistical
information.
178

179   ***Table* 2:** *Participant demographics, and sleep and vigilance scores for each age group in Experiment 2.*

Variable		Children	Adolescents	Young adults	Older adults	p-value
n		27	27	27	27	/
Female *(n)*		16	12	21	18	0.08
Age (years)		10.6 (1.6)	15.4 (1.4)	26.6 (4.4)	63.0 (4.4)	/
M/E Preference		3.0 (0.8)	3.2 (0.9)	3.2 (0.8)	2.4 (0.9)	0.002*
Session 1	*Time of testing*	11:34 (97)	11:34 (96)	11:39 (82)	11:30 (92)	0.99
	Sleep quality	4.9 (0.7)	5.1 (0.6)	5.0 (0.6)	4.7 (0.8)	0.19
	Sleep duration (hrs.)	9.7 (1.1)	9.4 (1.2)	8.3 (1.1)	7.7 (1.3)	<0.001*
	SSS score	2.2 (0.9)	2.1 (0.8)	2.0 (0.7)	1.5 (0.7)	0.012*
	PVT score (ms)	360.9 (81.5)	341.9 (102.3)	320.9 (62.0)	327.5 (75.7)	0.31
Session 2	*Time of testing*	16:42 (99)	16:56 (112)	16:43 (89)	16:29 (96)	0.81
	Offline period (hrs.)	5.1 (0.8)	5.4 (0.9)	5.1 (1.1)	5.0 (0.7)	0.43
	SSS score	2.0 (1.0)	1.8 (0.8)	2.0 (0.7)	1.7 (0.7)	0.36
	PVT score (ms)	386.0 (88.4)	318.5 (61.0)	320.3 (69.1)	323.6 (68.0)	0.002*
Session 3	*Time of testing*	11:57 (148)	12:25 (156)	11:30 (97)	11:24 (109)	0.31
	Offline period (hrs.)	24.4 (1.8)	24.8 (1.9)	23.8 (1.4)	23.9 (1.4)	0.095
	Sleep quality	5.0 (0.8)	4.7 (0.8)	4.7 (0.7)	4.6 (0.7)	0.38
	Sleep duration (hrs.)	10.3 (0.9)	9.3 (1.5)	8.0 (1.3)	7.6 (1.0)	<0.001*
	SSS score	1.9 (0.7)	2.3 (1.3)	2.3 (0.8)	1.5 (0.6)	0.003*
	PVT score (ms)	378.7 (97.3)	325.8 (74.6)	308.0 (54.4)	306.3 (33.6)	<0.001*

180
Numbers represent the mean, with standard deviation in parentheses. Morningness/Eveningness (M/E)
Preference was added as an exploratory variable to assess age-related differences in circadian preferences
and is defined on a 5-point Likert scale (i.e., 1 = extreme morning person, 5 = extreme evening person).
The average time of testing is specified in 24-hour time notation, with the SD in minutes. The offline periods
represent the time periods between session 1 and sessions 2 and 3, respectively. Sleep quality is defined on
a 6-point Likert scale (i.e., 1 = very bad, 6 = very good). SSS = Stanford Sleepiness Scale[31], with higher
numbers indicative of increased sleepiness. The PVT score indicates the average simple RT across the 30
trials. The p-values resulted from one-way ANOVAs assessing group differences. See Appendix 1 as part
of the Supporting Information for full statistical information.
190

2.2. General Experimental Procedures

After verification of eligibility, participants were contacted via email with the necessary

information to continue participation. This email contained a link to a video created by our study

team that provided detailed instructions on how to complete our motor learning task (details on

the task are provided in Section 2.3. below). Participants were also provided links to our series of

tasks created on the web-based data collection platform PsyToolkit[32,33]. As our experiments were

conducted online, participants were free to choose which day(s) to complete experiments but were

instructed to complete the protocols on days in which they were well-rested and could adhere to

the schedule of the sessions. The precise times-of-day that the experiments were completed were

constrained and followed experiment-specific schedules provided below (see Sections 2.4 and

2.5). Participants were instructed to receive a good night of sleep and to refrain from alcohol the

night before and throughout the experiment. Compliance to these instructions was verified through

a brief questionnaire prior to completion of the motor task. Lastly, parents of the child participants

were asked to be present when their child completed the experiments.

For both experiments and at the beginning of each experimental session, a brief questionnaire was

completed in which participants reported their sleep patterns in the previous 24 hours and their

subjective feelings of alertness (Stanford Sleepiness Scale[31]). Additionally, and for Experiment 2

only, participants completed the Psychomotor Vigilance Task (PVT[34]) to provide an objective

measure of vigilance at the time of testing. The PVT required participants to fixate on the middle

of the computer screen and press the space bar as fast as possible when a visual stimulus appeared

after a varying delay interval. Response time (i.e., time between appearance of stimulus and

keypress) was logged and used to assess vigilance (i.e., higher RTs indicative of lower vigilance).

We administered a shortened version of the PVT[35,36] that consisted of 35 trials, of which the first

5 were discarded as participants became familiar with the task.

217
Figure 1. a) *Serial reaction time task (SRTT). A stimulus appeared in 1 of 8 spatial locations and the*
participant responded with the corresponding key/finger as fast as possible. Stimuli appeared in either a
pseudorandom or sequential (i.e., 4-7-3-8-6-2-5-1) manner. ***b)*** *To increase motivation and attention, the*
task was built around a story in which participants were asked to catch butterflies. ***c)*** *Design of Experiments*
1 and 2. Experiment 1 consisted of the first session only (i.e., Day 1 in blue) and Experiment 2 of all three
sessions displayed. Session 1 included four blocks of pseudorandom SRTT, followed by 20 blocks of
sequential SRTT, which were divided into 16 blocks of training and 4 blocks of test after a 1-minute rest,
and four additional blocks of pseudorandom SRTT. Approximately 5 and 24 hours after session 1, session
2 and 3 were completed, respectively. Both sessions 2 and 3 consisted of 4 blocks of sequential SRTT and
4 blocks of pseudorandom SRTT. Note that the first session of Experiment 2 was constrained between 9am
and 2pm to give ample time for the 5hr delayed retest. Those participants in Experiment 1 were instructed
to complete the session anytime between 9am and 7pm.

2.3 Serial Reaction Time Task (SRTT)

All participants performed an adapted version of the explicit Serial Reaction Time Task (SRTT)

similar to our previous research[37,38], which was coded in the online data acquisition platform

PsyToolkit[32,33]. The SRTT was chosen for this research, in part, because of the online data

acquisition protocol. It was assumed that participants, and young children in particular, could more

easily comprehend and follow the instructions for an SRTT variant (i.e., press the key that spatially

corresponds to the visual stimulus). This in contrast to the sequential finger tapping task, where

fingers are assigned numerical values and participants are then instructed to perform the sequence

of finger movements that corresponds to the explicitly provided sequence of numbers.

The SRTT employed in this research consisted of an 8-choice reaction time task in which

participants were instructed to react to visual cues shown on a screen (see Figure 1a). Eight squares

that spatially corresponded to the eight fingers used to perform the task (i.e., no thumbs) and the

eight keys on the keyboard were presented on the screen. The color of the squares alternated

between red and green, indicating rest and practice blocks, respectively. During practice, a visual

stimulus (i.e., a butterfly) appeared consecutively in one of the squares and participants responded

to the visual cues by pressing the corresponding key with the corresponding finger as fast and

accurately as possible. The task was presented as a game, designed to increase motivation in child

participants, in which participants of all four age groups were instructed to move quickly to catch

as many butterflies as possible (see Figure 1b). The visual cues succeeded each other in either a

pseudorandom or a sequential order, depending on the experimental run (i.e., pseudorandom SRTT

vs. sequential SRTT, respectively). In the sequential SRTT, participants were aware that the

stimuli and keypresses followed a deterministic sequential pattern (i.e., 4-7-3-8-6-2-5-1, in which

1 through 8 are the left pinky to the right pinky fingers from left to right), but they were not given

any information about the structure or length of the sequence. All participants completed the same

8-element sequence. During the pseudorandom SRTT runs, the visual stimuli (and thus

corresponding key presses) appeared in an order that pseudo-randomly changed every 8 elements.

Specifically, the stimulus appeared in each location once every eight elements, but the stimulus

never appeared in the same location consecutively. The order of the stimuli thus changed every

eight elements within a block, across blocks of practice and differed across participants. For both

pseudorandom and sequential task variants, practice blocks contained 48 key presses (i.e., 6

repetitions of the 8-element sequence in the sequence condition) and were separated by 15-second

rest periods.

Measures reflecting performance speed (i.e., response time between visual cue and keypress) and

accuracy (i.e., correct or incorrect keypress) of the participants' motor responses were recorded

for each trial and used to assess performance on the task. Per our pre-registrations, individual trials

(i.e., responses) were excluded from analyses if the measured response time was greater than 3

standard deviations above or below the participant's mean response time for that block. For

Experiments 1 and 2, an average of 1.78% (children: 2.03%, adolescents: 1.76%, young adults:

1.68%, older adults: 1.65%) and 1.69% (children: 2.0%, adolescents: 1.83%, young adults: 1.58%,

older adults: 1.34%) of trials were excluded from analyses, respectively.

2.4 Experiment 1 Design

The first experiment consisted of a single experimental session in which participants performed

both the random and sequential variants of the SRTT (see Day 1 [blue] of Figure 1c). Participants

were instructed to complete this session between 9am and 7pm (see Table 1 in the main text and

Appendix 1 as part of the Supporting Information for details on time of testing). The experimental

session started with one block of familiarization (pseudorandom; data not analyzed) followed by

four blocks of the pseudorandom SRTT (pre-learning random), which afforded an assessment of

general motor execution on the task prior to any sequence learning. Subsequently, participants

completed 20 blocks of the sequential SRTT. These blocks were divided into two runs: a training

run that consisted of 16 blocks and then a post-learning test run of 4 blocks. The post-learning test

run was completed approximately one minute after the end of training, affording the assessment

of end-of-training performance following the further dissipation of mental and physical fatigue[39].

Lastly, four additional blocks of the pseudorandom SRTT (post-learning random) were completed,

allowing us to assess whether improvements in performance on the sequential SRTT could be

attributed to learning of the movement sequence *per se* (i.e., sequence-specific learning) or simply

a general improvement in movement speed due to task familiarization.

2.5 Experiment 2 Design

The second experiment consisted of three experimental sessions (see Figure 1c). Participants were

instructed to complete the first session between 9 am and 2 pm. This session was identical to what

was described for Experiment 1. The second session took place approximately five hours after the

first session, allowing the assessment of motor memory consolidation after an interval of post-

learning wakefulness. This second session consisted of four blocks of sequential SRTT during

which participants again performed the same specific sequence as in session 1, and four blocks of

pseudorandom SRTT (post-test random). Finally, the third session was completed 24 hours after

the first session, allowing the examination of motor memory consolidation after a night of sleep.

Similar to the second session, this third session included four blocks of sequential SRTT and four

blocks of pseudorandom SRTT (post-test random). See Table 2 in the main text and Appendix 1

as part of the Supporting Information for details on actual time of testing. For the intervals between

sessions, participants were instructed to refrain from napping, consuming alcohol or recreational

drugs and to receive a good night of sleep.

2.6 Data Processing and Statistical Analyses

We first describe general data processing and statistical analysis information prior to providing

details specific to each of the two experiments. All null hypothesis statistical testing described

below was performed using IBM SPSS Statistics for Windows, version 28 (IBM, Armonk , NY,

USA). All significance tests were two-sided, considered significant when $p < 0.05$ and performed

for both normalized outcome measures (i.e., reaction time and accuracy). For the sake of

completeness, we also highlight those effects when $0.05 < p < 0.1$ as non-significant trends. In the

event of a violation of the sphericity assumption, Greenhouse-Geisser correction was applied. All

significant main effects and interactions were followed up by pairwise comparisons using the

Tukey test and one-way ANOVAs, respectively. Depending on the statistical test, eta-squared,

partial eta-squared or Hedges' g are reported as effect size measures. The correspondence among

reported p-values, test statistics and degrees of freedom reported in this main text was verified via

statcheck[40].

In addition to the null hypothesis significance testing outlined above, the anovabf, ttestbf and

regressionbf functions in RStudio version 1.3.1093 (Posit, PBC, Boston, Massachusetts, USA)

were used to compute Bayes Factors (BF) for all effects presented in the main text. Note that this

Bayesian analysis was *not* included in the pre-registrations. BFs reflect the likelihood that the

observed data favor an alternative model (e.g., evidence for differences among groups, blocks,

sessions, etc.) relative to the null model (e.g., evidence for no differences). For t-tests, the null

model was the difference between means equal to zero. For mixed model ANOVAs, the null

models were specified as the random effect of subjects. For one-way ANOVAs and simple linear

regressions, the null models were only the intercept term. For multiple linear regressions that

assessed age group differences in the relationship between micro- and macro-offline performance

gains (see below for details), the reported BFs reflect the comparison of two models with and

without the age group x micro-offline gain interaction term. Default parameters in the R functions

were used, with the exception of "whichModels" that was set to "all" for mixed model ANOVAs

and multiple regressions in order to compute the appropriate BFs. In our results, we report BF_{10}

values and adhere to the interpretations offered in Wagenmakers et al.[41]; larger values are

indicative of greater likelihood that the observed data favor the alternative as compared to the null

hypotheses.

To assess potential differences among age groups and experimental sessions in certain

demographic/participant characteristics, one-way ANOVAs or chi-square tests (Experiment 1) and

two-way ANOVAs (Experiment 2) were performed. These measures included gender, vigilance

(both subjective and objective) at the time of testing, self-reported sleep quantity and quality of

the night prior to the experimental session(s), and the time of day in which the motor task or session

was completed. Any significant effects from these analyses were followed up by pairwise

comparisons to determine which age groups or sessions differed from each other. Group means

for these dependent measures are provided in Tables 1 and 2 and full output from the corresponding

statistical analyses can be found in Appendix 1 of the Supporting Information.

For the SRTT, the averaged response times for correct keypresses and the percentage of correct

keypresses were computed for each block and task variant. Additionally, as previous research has

indicated age differences in motor performance independent of sequence learning[6,7,22,23,42],

baseline differences in performance were accounted for by normalizing both movement speed and

accuracy of the sequential and post-learning random SRTT runs relative to the outcome variables

of the pre-learning random run. More specifically, for each outcome measure and each block of

the SRTT runs, performance was divided by the mean outcome measure across the four blocks of

the pre-learning random run. Statistical analyses of these normalized measures are presented in the

main text. For completeness, results based on non-normalized performance measures are reported

in Appendix 2 of the Supporting Information.

2.6.1 Experiment 1

Sequential learning dynamics (i.e., block-to-block performance changes) were compared among

age groups using a mixed ANOVA, with *group* (i.e., children, adolescents, young adults, older

adults) as between-subject and *block* (i.e., 16 training blocks or 4 test blocks) as within-subject

factors.

To assess whether observed improvements in performance across practice blocks reflected

sequence-specific learning (as opposed to general skill learning as a result of familiarization with

the motor task), a learning magnitude measure was computed for performance speed. Specifically,

the difference between averaged normalized performance across the four blocks in the post-

learning test run of the sequential SRTT and the averaged normalized performance in the post-

learning random run was divided by the average normalized performance in the post-learning

random run. This measure thus reflects the relative difference between the random and sequential

task variants at the end of training, affording the assessment of sequence-specific learning. Larger

values are indicative of greater sequence-specific learning. The effect of age *group* on learning

magnitude was assessed with a one-way ANOVA. Appendix 3 of the Supporting Information

includes corresponding results for when this learning magnitude measure was computed on non-

normalized RT data. It is worth noting here that the findings were consistent across the two

computations.

To provide a more fine-grained characterization of the age-related changes in sequence-specific

learning magnitude, we conducted exploratory analyses with age as a continuous variable.

Specifically, we tested five potential fit options (i.e., single exponential, double exponential, linear,

quadratic, and power functions) to characterize the relationship between age and learning

magnitude. As we did not acquire data from participants between 35 and 55 years of age, separate

models were built to characterize age-related changes from 7 to 35 and 55 to 75 years of age. The

model with the lowest Akaike Information Criterion value was deemed as the best fitting model

and the corresponding results are presented.

Similar to previous research[5,43,44], we distinguished between micro-online (i.e., within blocks of

task practice) and micro-offline (i.e., across rest epochs in between blocks of practice) performance

improvements. Micro-online changes were computed as the difference between the averaged

normalized RT for the correct keypresses of the first and last sequence repetitions within each

block (i.e., first sequence block n – last sequence block n). For micro-offline changes, the

difference between the averaged normalized RT of the last sequence repetition of one block and

the first sequence repetition of the subsequent block (i.e., last sequence block n – first sequence

block n+1) was computed. These micro-online and micro-offline measures were analyzed with

separate *group(4)* x *block(16)* or *group(4)* x *rest block (15)* mixed ANOVAs, respectively.

Appendix 3 in the Supporting Information includes corresponding results when micro-online and

-offline measures were computed with non-normalized RT data. Last, and following the same

exploratory procedure as described above for the learning magnitude measure, we assessed age-

related changes in these micro measures with age as a continuous variable.

2.6.2. Experiment 2

As in Experiment 1, learning dynamics (i.e., block-to-block performance changes) across training

and the post-learning test were analyzed using *group(4)* x *block(16 or 4)* mixed ANOVAs to assess

potential group differences in initial encoding of the motor sequence within the first session (i.e.,

before the assessment of consolidation).

The main focus of Experiment 2 was the effect of age on motor memory consolidation across the

5hr and 24hr macro-offline intervals. To this end, and for both the sequential and random SRTT,

a macro-offline change measure was computed for each outcome variable and offline interval.

Specifically, sequential offline changes were calculated as the difference between the average

normalized performance on the sequential SRTT run in the 5hr and 24hr retest sessions and the

average normalized performance in the post-learning test run of the first session. Similarly, offline

changes for the random task variant were computed as the difference between the averaged

normalized performance on the post-test random runs in the 5hr and 24hr retest sessions and the

averaged normalized performance in the post-learning random run of the first session. These

measures reflect the offline changes in sequential performance and general motor execution from

the end of the initial training session to the two retest sessions, respectively. Mixed ANOVAs with

group (4) as between-subject and *offline period* (i.e., 5-hour and 24-hour offline periods) as within-

subject factors were conducted. Appendix 4 in the Supporting Information includes corresponding

results when macro-offline measures were computed with non-normalized RT data. Consistent

with the exploratory analyses described above for Experiment 1, age-related changes in the macro-

offline performance gains were also assessed with age as a continuous variable.

Last, and similar to previous research[44,45], we assessed the relationship between offline changes in

performance on the micro- and macro-offline timescales. Specifically, we performed exploratory

multiple regressions with the micro-offline gains as independent and the 5h or 24h macro-offline

gains as dependent variables. To examine how these relationships differed among age groups, we

included age group (dummy coded) main effects as well as age group x micro-offline gains

interaction variables in the regression models. The young adult group was selected as the reference

group and thus the interaction variables compare the regression slopes of the other 3 age groups

with those of the young adults.

4. Results

4.1 Experiment 1: Initial motor sequence learning

4.1.1 Participant characteristics, sleep and vigilance

Group means from participant characteristics, the time of when the experimental session was

completed, self-reported sleep quality and duration of the night preceding the experimental

session, and a subjective measure of vigilance are provided in Table 1. Results from the

corresponding statistical analyses and a brief discussion can be found in Appendix 1 of the

Supporting Information.

4.1.2 Dynamics of Performance Improvements

Normalized performance measures on the SRTT are depicted in Figure 2a and 2b. Performance

accuracy remained stable across training, as evidenced by no main effect of block ($F_{(7.011,883.403)}$ =

0.501, p = 0.835, partial η^2 = 0.004, BF_{10} = 3.153e^{-06}), and did not significantly differ among age

groups ($F_{(3,126)}$ = 0.322, p = 0.810, partial η^2 = 0.008, BF_{10} = 0.091). Furthermore, no significant

group x block interaction was found ($F_{(21.033,883.403)}$ = 0.0.715, p = 0.821, partial η^2 = 0.017, BF_{10} =

1.821e^{-05}). Similarly, no block ($F_{(2.283,287.681)}$ = 1.523, p = 0.217, partial η^2 = 0.012, BF_{10} = 0.060),

group ($F_{(3,126)}$ = 1.796, p = 0.151, partial η^2 = 0.041, BF_{10} = 0.456), or interaction effects

($F_{(6.850,287.681)}$ = 0.814, p = 0.574, partial η^2 = 0.019, BF_{10} = 0.013) were found for the post-learning

test.

Figure 2. a*) Average normalized response time for both task variants. **b****) Average normalized accuracy for*
*both task variants. For panels a and b, shaded regions represent standard errors of the mean. **c****) Average*
learning magnitude for performance speed per age group. Shaded regions represent the kernel density
estimate of the data, colored circles depict individual data, open circles represent group medians, and the
*horizontal lines depict group means[46]. * p < 0.05 for pairwise group comparisons. **d****) Learning magnitude*
as a function of age. Quadratic fit from childhood into young adulthood: Learning magnitude = (-
0.0008572*age²) + (0.04021*age) + (-0.2226). Quadratic fit for older adults: Learning magnitude =
(0.00157*age²) + (-0.1996*age) + 6.4675.

For an assessment of performance speed, response times significantly decreased across sequential

practice blocks, as shown by a main effect of block ($F_{(7.900,995.358)}$ = 64.246, p < 0.001, partial η^2 =

0.338, BF_{10} = 5.881e^{152}). These performance improvements were similar among age groups, as

indicated by the lack of a significant group effect ($F_{(3,126)}$ = 1.578, p = 0.198, partial η^2 = 0.036,

BF_{10} = 0.291) and group x block interaction ($F_{(23.699,995.358)}$ = 1.337, p = 0.130, partial η^2 = 0.031,

BF_{10} = 1.21e^{-04}). During the post-learning test run, there was no significant block effect

($F_{(2.102,264.890)}$ = 0.404, p = 0.678, partial η^2 = 0.003, BF_{10} = 0.012), indicating a performance plateau

was reached. Moreover, consistent with the training run, there were no significant differences

among groups (Group main effect: $F_{(3,126)}$ = 1.099, p = 0.352, partial η^2 = 0.025, BF_{10} = 0.318;

Group x Block interaction: $F_{(6.307,264.890)}$ = 1.148, p = 0.335, partial η^2 = 0.027, BF_{10} = 0.032).

Collectively, the assessment of performance across blocks of practice indicates that the four groups

acquired the novel motor sequence, as evidenced by the reduction in RTs in the training run. And,

we found no credible evidence that the four groups differed in these performance improvements.

4.1.3 Sequence-specific learning

A learning magnitude measure was calculated for performance speed that reflects the relative

difference between the random and sequential task variants at the end of training and therefore

affords the assessment of the magnitude of sequence-specific learning (see Figure 2c). Results

revealed a significant group effect ($F_{(3,129)}$ = 5.440, p = 0.002, η^2 = 0.115, BF_{10} = 20.731). More

specifically, a significantly smaller sequence-specific learning magnitude was observed in children

as compared to adolescents ($p = 0.014$, $G = 0.714$, $BF_{10} = 8.645$) and young adults ($p = 0.001$, G

$= 0.970$, $BF_{10} = 124.212$). Follow-up analyses indicated that the difference in learning magnitude

could be attributed to significant group differences in RT for the post-learning random run ($F_{(3,126)}$

$= 5.795$, $p < 0.001$, partial $\eta^2 = 0.121$, $BF_{10} = 35.104$). Children demonstrated greater

improvements in performance speed from the pre-learning to post-learning *random* runs as

compared to young ($p < 0.001$, $G = 0.876$, $BF_{10} = 43.280$) and older adults ($p = 0.022$, $G = 0.591$,

$BF_{10} = 3.018$). Accordingly, a portion of the performance improvements across the sequential

training blocks in children can be attributed to general learning of the task that is *not* specific to

the sequence.

We also conducted exploratory regression analyses with age as a continuous variable to provide a

more fine-grained characterization of the age-related trajectories of sequence-specific learning

magnitude. Consistent with the differences among *groups*, age was a significant predictor of

learning magnitude from childhood into young adulthood (see Figure 2d; the best fit model was a

quadratic function: $R^2 = 0.156$, $F_{(1,95)} = 8.808$, $p = 0.004$). Specifically, learning magnitude

increased across childhood, reaching peak values at approximately 24 years of age, and then

decreased thereafter. Age was also a significant predictor of learning magnitude within the older

adult group, exhibiting an approximate U-shaped trajectory from 55 to 75 years (quadratic model:

$R^2 = 0.247$, $F_{(1,29)} = 4.746$, $p = 0.038$).

4.1.4 Micro-learning across sequential training

For micro-online learning, the block x group interaction was not significant ($F_{(34.573, 1429.030)}$ =

1.159, p = 0.243, partial η^2 = 0.027, BF_{10} = 0.001; see Figure 3a). There was, however, a significant

block effect ($F_{(11.524, 1429.030)}$ = 2.358, p = 0.006, partial η^2 = 0.019, BF_{10} = 0.140) as well as a non-

significant trend for a group effect ($F_{(3,124)}$ = 2.663, p = 0.051, partial η^2 = 0.061, BF_{10} = 0.259; see

Figure 3c). Although these findings suggest differences in micro-online learning across blocks of

practice and among age groups, the corresponding BFs indicate that the data are more consistent

with no differences as compared to the alternative (i.e., evidence of differences). Consistent with

these results, there was only a non-significant trend for age being a predictor of micro-online gains

from childhood into young adulthood (power model: R^2 = 0.032, $F_{(1,96)}$ = 3.141, p = 0.080; see

Figure 3e). No age-related changes were revealed within the older adult group (power model: R^2

= 0.045, $F_{(1,30)}$ = 1.424, p = 0.242).

The assessment of micro-offline learning showed no rest block ($F_{(11.662, 1446.066)}$ = 1.209, p = 0.272,

partial η^2 = 0.010, BF_{10} = 2.28e^{-04}; see Figure 3b), nor a rest block x group interaction

($F_{(34.985, 1446.066)}$ = 0.836, p = 0.739, partial η^2 = 0.020, BF_{10} = 7.098e^{-05}). There was a non-significant

trend for a group effect ($F_{(3,124)}$ = 2.490, p = 0.063, partial η^2 = 0.057, BF_{10} = 0.217), yet the

corresponding BF indicated that the data are more in favor with the null hypothesis as compared

to the alternative. Similarly, the characterizations of age-related trajectories revealed that there was

only a non-significant trend for age being a predictor of micro-offline gains from childhood into

young adulthood (power model: R^2 = 0.029, $F_{(1,96)}$ = 2.873, p = 0.093; see Figure 3f). No significant

age-related changes were revealed within older adulthood (power model: $R^2 = 0.043$, $F_{(1,30)} =$

1.362, p = 0.252).

528
Figure 3. *Micro-online (**left**) and -offline (**right**) gains. Panels a and b depict micro gains displayed as a*
function of practice blocks. Panels c and d contain violin plots of micro-online and -offline, respectively,
gains averaged across blocks. Note that pairwise comparisons in these panels included blocks or rest
periods in the statistical models. Shaded regions represent the kernel density estimate of the data, colored
circles depict individual data, open circles represent group medians, and the horizontal lines depict group
means[46]. Panels e and f show averaged micro-online and -offline gains as a function of age from childhood
into young adulthood and within older adulthood. Lines of fit childhood into young adulthood: micro-online
gains = (-0.318*age)^{-0.459}; micro-offline gains = 0.298*exp(-0.377*age). Lines of fit within older
adulthood: micro-online gains = (0.003*age) + (-0.238); micro-offline gains = (227.361*age) + (-1.897).
538

4.2 Experiment 2: Motor memory consolidation

4.2.1 Participant characteristics, sleep and vigilance

Group means for participant characteristics, the times of when the experimental sessions were

completed, the duration of the offline periods between sessions, self-reported sleep quality and

duration for the nights preceding each experimental session, and subjective (SSS) and objective

(PVT) measures of vigilance are provided in Table 2. Results from the corresponding statistical

analyses and a brief discussion can be found in Appendix 1 of the Supporting Information.

4.2.2 Initial learning dynamics

Similar to Experiment 1, we assessed the dynamics of learning during the initial training session

of Experiment 2. Normalized performance measures are shown in Figure 4. For brevity, the

corresponding statistical results can be found in Appendix 5 of the Supporting Information. In

brief, the results from the assessment of the performance improvements during the initial training

session of Experiment 2 largely mirrored those presented above for Experiment 1. We found little

credible evidence for differences among age groups in the performance changes across training

blocks.

555
Figure 4. *Average normalized response time (**a**) and accuracy (**b**) for the three experimental sessions and*
two task variants of Experiment 2. Shaded regions represent standard errors of the mean. Corresponding
statistical analyses are in Appendix 5 of the Supporting Information.

4.2.3 Macro-offline performance gains

Results on the sequential macro-offline changes in performance (Figure 5) revealed both offline

period ($F_{(1,104)}$ = 104.357, p < 0.001, partial η^2 = 0.501, BF_{10} = 1.037e^{14}) and group ($F_{(3,104)}$ =

14.541, p < 0.001, partial η^2 = 0.296, BF_{10} = 90620.35) main effects, as well as a significant offline

period x group interaction effect ($F_{(3,104)}$ = 3.382, p = 0.021, partial η^2 = 0.089, BF_{10} = 0.384).

Follow-up comparisons revealed significant group effects for both the 5-hour ($F_{(3,107)}$ = 5.756, p =

0.001, η^2 = 0.142, BF_{10} = 28.76) and the 24-hour ($F_{(3,107)}$ = 20.582, p < 0.001, η^2 = 0.373, BF_{10} =

26.800e^{06}) sequential macro-offline gains. Post-hoc pairwise comparisons indicated that the

sequential macro-offline gains after a 5-hour period of wakefulness were higher in children and

adolescents as compared to the young (p = 0.032, G = 0.590, BF_{10} = 1.943; p = 0.053, G = 0.548,

BF_{10} = 1.497, respectively) and older adults (p = 0.008, G = 1.382, BF_{10} = 3765.434; p = 0.015, G

= 1.290, BF_{10} = 1289.955, respectively). It is worth emphasizing that the corresponding BFs

suggest only anecdotal evidence that the data are in favor of the alternative hypothesis (i.e.,

evidence for group differences) as compared to the null hypothesis for the 5-hour offline gains.

The children showed higher 24-hour macro-offline gains in comparison to the young ($p = 0.001$,

$G = 1.285$, $BF_{10} = 574.756$) and older ($p < 0.001$, $G = 2.449$, $BF_{10} = 1.080e^{08}$) adults, and the older

adults displayed significantly lower 24-hour macro-offline gains than all the other age groups

(adolescents: $p < 0.001$, $G = 1.256$, $BF_{10} = 658.860$; young adults: $p = 0.002$, $G = 1.237$, $BF_{10} =$

458.448).

In line with the age group comparisons presented above, our analyses with age as a continuous

variable revealed that age significantly predicted the 5h macro-offline gains from childhood into

young adulthood (linear model: $R^2 = 0.068$, $F_{(1,78)} = 5.650$, $p = 0.020$; see Figure 5b). Age was not

a significant predictor of the 5hr offline gains within the older adult group (double exponential

model: $R^2 = 0.183$, $F_{(1,23)} = 1.72$, $p = 0.203$). Lastly, age significantly predicted the 24-hour macro-

offline gains from childhood to young adulthood (Power model: $R^2 = 0.157$, $F_{(1,78)} = 14.537$, $p <$

0.001; see Figure 5d) as well as within older adulthood (linear model: $R^2 = 0.136$, $F_{(1,25)} = 3.95$, p

$= 0.058$).

Figure 5. *Sequential offline gains across the 5-hour (**Panel a**) and 24-hour (**Panel c**) offline periods for the*
four age groups. Shaded regions represent the kernel density estimate of the data, colored circles depict
*individual data, open circles represent group medians, and the horizontal lines depict group means[46]. * p*
*< 0.05 and ~ p < 0.10 for pairwise group comparisons. Five-hour (**Panel b**) and 24-hour (**Panel d**)*
sequential offline gains are plotted as a function of age. Lines of fit from childhood into young adulthood:
5h gains = (-0.003 * age) + 0.075; 24h gains = (0.872*age)$^{-0.682}$. Lines of fit within older adulthood: 5h
gains = (1.814e^{-6} * exp(0.256 * age)) + (-1.858e^{-6} * exp(0.256 * age)); 24h gains = (-0.006 * age) + 0.381.
For better visualization of the age-related changes, the scale of the y-axes was set to range from -0.4 to 0.4
and thus the young adult with the outlier 5hr sequential gain (visible in Panel a) is not depicted in Panel b.
This individual, however, was included in the fitting procedure. Note that if this individual was excluded
from statistical analyses, the pattern of results observed in Panel a remains largely similar, with the
exception that the pairwise difference between young and older adults becomes statistically significant (i.e.,
older adults exhibit worse offline consolidation over a 5hr wake interval as compared to young adults).

4.2.4 Link between the offline performance changes on a micro- and macro-timescale

The regression model assessing the relationship between the micro- and 5h macro-offline gains –

independent of age group – revealed a significantly positive relationship between these two offline

measures (b = 0.581, p < 0.001, BF$_{10}$ = 831.113; see Appendix 6 in Supporting Information). This

suggests that greater performance changes during the short rest periods throughout training were

associated with greater performance gains across the 5-hour offline period of wakefulness

following training. We then assessed whether such a relationship differed between pairs of age

groups. Young adults exhibited a steeper slope as compared to children (b = -1.371, p = 0.003,

BF_{10} = 0.381), adolescents (b = -1.287, p = 0.002, BF_{10} = 0.476) and older adults (b = -1.509, p =

0.003, BF_{10} = 0.475). However, the reported BFs indicate that the data are more consistent with

no differences and these between-group differences were no longer statistically significant if the

young adult participant with the extremely negative 5hr offline gain (see Figure 5a) was removed

from the analyses.

Micro-offline gains were also significantly and positively related to the 24h macro-offline gains

(b = 0.609, p < 0.001, BF_{10} = 16736.4; see Appendix 6). Specifically, greater performance changes

during the short rest periods were also associated with greater performance gains across the 24-

hour period following training. This relationship was significantly different between the children

and young adults (b = -0.669, p = 0.047, BF_{10} = 6.268). Specifically, whereas young adults

exhibited a significant positive relationship between micro- and macro-offline performance gains

(b = 0.647, p = 0.007, BF_{10} = 6.380), children did not (b = -0.023, p = 0.90, BF_{10} = 0.360). The

relationship between micro-offline and 24-hr macro-offline gains did not differ between young

adults and the other two age groups (adolescents: b = 0.126, p = 0.679, BF_{10} = 0.594; older adults:

b = -0.217, p = 0.556, BF_{10} = 0.189).

5. Discussion

The current study examined age-related differences among children, adolescents, young adults and

older adults in initial motor sequence learning and motor memory consolidation. Our results

revealed that: (1) the dynamics of initial motor sequence learning, as demonstrated by performance

improvements during training, were comparable (i.e., not statistically different) across age groups;

(2) children exhibited less sequence-specific learning during the initial training session relative to

adolescents and young adults; (3) although micro-online and -offline performance changes were

the smallest and largest, respectively, in children, there was not sufficient evidence to support the

hypothesis of differences among age groups; (4) children exhibited greater changes in performance

from the end of initial training to both the 5-hour and 24-hour retests, suggesting a developmental

advantage in the macro-offline consolidation of a newly acquired movement sequence; and, (5)

there was a significant positive relationship between the magnitudes of micro- and macro-offline

performance changes.

Initial performance improvements are comparable across age groups, but children exhibit less
sequence-specific learning

On the one hand, our results provided no evidence that performance improvements across training

differed among age groups, which was consistent with a subset of the previous literature[5–8,22–26,47].

This finding thus agrees with the lifespan invariance model that has been discussed in previous

literature in the context of memory consolidation[29]. On the other hand, our results indicated that

children exhibited less sequence-specific learning and thus their performance improvements across

sequential training blocks are partially due to general motor improvements (i.e., improvements on

the motor task that are *independent* of sequence learning). This finding – which resembles the

prototypical developmental trajectory where young adults perform the best and children are

progressing towards this optimal performance level - is consistent with a subset of previous

literature in which children exhibited smaller sequence-specific motor learning[11,12,48]. The pattern

of results may explain some of the heterogeneity that is present in the literature with respect to the

development of motor sequence learning. That is, previous research that assessed performance

improvements across blocks of practice may report comparable learning between children and

young adults (e.g., [8,20]), whereas studies that afford the extraction of sequence-specific learning

may reveal differences between age groups (e.g., [11,48]). Presumably, the greater general learning

that is evident in children is due to this group simply learning how to perform the task itself (i.e.,

spatially mapping the stimuli on the screen to the appropriate fingers). Future research that aims

to isolate and examine sequence learning *per se* could consider incorporating an extended task

familiarization phase consisting of additional random runs (i.e., no sequence to be learned). This

would allow children to exhaust their general learning of the task itself prior to sequence

acquisition. Nevertheless - and as a point of emphasis - even though sequence-specific learning

was smaller in children, they still exhibited significant learning of the motor sequence. That is,

their sequence-specific learning magnitude was significantly greater than zero (see Appendix 7)

and thus a memory trace of the acquired sequence was presumably formed. Interestingly, whereas

children showed a smaller learning magnitude, the adolescents were not different from young

adults. This suggests that sequence-specific learning reaches adult-like levels by the teenage years.

Micro-offline consolidation during interspersed rest breaks

In line with previous literature[5] and our hypothesis, children exhibited the largest micro-offline

performance gains. However, the group main effect within the omnibus group (4 levels) x block

ANOVA was a non-significant trend ($p = 0.063$) and the corresponding Bayes Factors did not

provide substantial evidence in favor of age group differences. Specifically, although pairwise

comparisons indicated that micro-offline gains were larger in children as compared to both young

and older adults ($F_{(1,61)} = 4.829$, $p = 0.032$, partial $\eta^2 = 0.073$ and $F_{(1,61)} = 5.374$, $p = 0.024$, partial

$\eta^2 = 0.081$, respectively), the corresponding BFs were 1.109 and 1.411, respectively, and thus

indicative of only anecdotal evidence in support of age group differences[41]. It is worth noting here

that the assessment of micro-offline gains on non-normalized RT data did reveal significant age

group effects, with children exhibiting larger gains than the other 3 age groups (see Appendix 3).

This analysis, however, was not part of our pre-registration. Nonetheless, the result of our pre-

registered analysis of micro-offline gains presented in the main text can be viewed as weaker in

comparison to Du et al.[5]. One potential explanation could be the specific ages included in the child

groups. Specifically, Du et al.[5] found a micro-offline advantage in younger children (i.e., 6-year-

olds) as compared to older children (i.e., 10-year-olds) and adults. In the current research, the

online nature of our protocol made it difficult to acquire data in such young children (i.e., 6-year-

olds). Moreover, as data acquisition in Du et al.[5] was limited to these discrete age groups in (i.e.,

data from 7-9-year-old children were not acquired), an estimate of when this childhood advantage

in micro-offline processing disappears was not possible. We speculate that the differences between

children and young adults in our study would have been larger in magnitude if the sample included

a greater number of younger children (i.e., 6 to 7-year-olds). Another potential consideration is the

considerably longer rest breaks (i.e., 3 min) between practice blocks that Du et al.[5] implemented.

In comparison to our 15-second rest intervals, this allowed more time for the recently acquired

memory trace to be consolidated over these micro-offline epochs. Of note, recent research in young

adults has produced inconclusive results with respect to the impact of the rest period durations. In

a probabilistic sequence task, there was no effect of duration (15s vs. 30s vs. self-selected) on

micro-offline performance changes[49], whereas a temporal gradient (10s vs. 20s) was found in

resistance to an interfering sequence in a deterministic motor sequence task[50]. In summary, future

studies are needed to further examine micro-learning processes in children, with systematic

examinations into certain methodological choices such as the age of the participants, the length of

the rest epochs and the choice of sequence task (i.e., probabilistic vs. deterministic).

Although this fast consolidation process occurring over micro-offline intervals has received

considerable attention over the last 5 years[43–45,50,51], the current study is the first, to the best of our

knowledge, to examine micro-learning processes in older adults. Our results found no evidence

for differences in micro-online and -offline performance gains between young and older adults.

Accordingly, whereas older adults have deficits in the macro-offline consolidation processes

occurring between two practice sessions[24–26,28], they do not exhibit deficits in the rapid

consolidation process that occurs during the rest periods interspersed with blocks of task practice.

Lastly, similar to previous studies that employed the self-initiated, sequential finger tapping task

in young adults[5,43–45], our results with the SRTT also provide evidence for a significant

contribution of the micro-offline gains to overall learning in all four age groups. That is,

performance improvements across blocks of training were largely due to micro-offline – as

compared to micro-online – performance gains. However, and independent of the specific task

variant, one cannot rule out the effect of fatigue on these results. Gupta and Rickard[52,53] have

argued that the magnitude of the micro-offline gains in young adults can be attributed to the build-

up of fatigue or reactive inhibition as a function of task practice within blocks. Specifically, it was

suggested that performance deteriorates across practice within a block due to reactive inhibition

and thus micro-offline performance gains are inflated due to this performance variable that can be

considered independent of learning and memory processes. However, it is worth noting that this

view cannot account for recent neuroimaging data showing reactivation of sequence learning-

related patterns of hippocampal activity during the interspersed rest periods[43,44,51]. These

neuroimaging data then suggest that micro-offline epochs afford an opportunity for a rapid

memory consolidation process. Additional research is necessary to definitively conclude whether

these micro-offline performance gains are reflective of a memory consolidation process or are

simply attributed to reactive inhibition.

Children exhibited a developmental advantage in macro-offline consolidation

Our results revealed larger sequential performance gains across both the 5-hour and 24-hour

macro-offline periods in children. While the young adults showed a performance maintenance (i.e.,

offline gains that were not significantly different from zero; see Appendix 7 in Supporting

Information) across the 5-hour interval, children and adolescents significantly improved their

performance on the sequential task. Interestingly, children also demonstrated higher offline gains

across the 24-hour period as compared to adults. While the enhanced 5-hour consolidation in

children is in line with the findings of Ashtamker & Karni[14] that showed an accelerated

consolidation in children across a one-hour offline period, the greater 24-hour gains stand in

contrast to previous research that demonstrated impaired overnight consolidation in children [[13,15];

see below for expanded discussion on sleep-related consolidation].

It could be argued that the larger macro-offline processing in children could be explained by

performance levels reached at the end of the training session[54,55]. For example, perhaps children

show greater macro-offline gains due to a continuation of initial learning rather than the offline

processing of previously acquired information. This potential explanation, however, is not in line

with our data. Notably, during the post-training test phase, there was an absence of a block effect

in all groups, including in the children. This indicates that all groups reached a performance plateau

at the end of the first session and thus macro-offline gains do not appear to be the result of

continued learning. Along the same lines, one could speculate that the young adults reached a

performance ceiling effect, and thus their potential to further increase their performance over

macro-offline intervals was saturated (see plot of absolute reaction times in Appendix 2 of the

Supporting Information). Again, this is not consistent with our data. Notably, there was a

significant session effect in sequential offline gains, whereby gains were larger in the 24-hour

retest as compared to the 5-hour retest. Within-group comparisons revealed that this session effect

was present for all age groups, including young adults. This suggests that a performance ceiling

effect was not reached and thus this explanation cannot fully account for our results. Last, one

could raise the question whether perhaps the macro-offline performance gains are inflated in

children due to non-sequence-specific improvements (i.e., improvements due to motor task

familiarization as opposed to sequence learning). However, the children did not show greater

macro-offline gains on the random task variant (see Appendix 8 in Supporting Information).

Moreover, and to assess this possibility further, we examined the offline changes in the sequence-

specific learning magnitude (i.e., difference between the learning magnitude (i.e., sequence vs.

random) of the 5hr and 24hr retest sessions and the learning magnitude of the first session; see

Appendix 8 in Supporting Information). These results revealed significant main effects of offline

period and group, but no offline period x group interaction effect. Post-hoc pairwise comparisons

indicated that children exhibit enhanced sequence-specific offline gains – across both offline

periods – in comparison to young and older adults. This finding is consistent with the assessment

of the sequential macro-offline gains presented in the main text (Figure 5). Altogether, these data

suggest that the offline changes in children likely involved the strengthening of their sequential

memory and that this strengthening was greater as compared to young and older adults.

Offline performance gains on the 24-hour retest were largest in children, a result that was

unexpected based on previous literature[13,15]. Specifically, children have typically exhibited

impaired consolidation over post-learning intervals that include sleep as observed in young adults.

It is worth noting that most of the previous studies comparing sleep and wake conditions in children

employed an AM/PM design. In this design, participants acquire the motor sequence at a different

time of day based on the condition and then are retested after a delay of 12 hours. For example,

the wake (i.e., AM-PM) group initially learns the sequence in the morning and then is retested

following 12 hours of wakefulness. The sleep (i.e., PM-AM) group completes initial learning in

the evening and this is retested after a 12-hour interval that included a night of sleep. Although

informative, these designs are sensitive to time-of-day effects. And there is evidence that the time

of initial practice can affect performance during acquisition and offline consolidation in young

adults[56,57]. Specifically, Kvint et al.[57] found a greater sequential knowledge acquisition but similar

12-hour consolidation in an AM-trained as compared to a PM-trained group. Furthermore, Truong

et al.[56] showed an enhanced consolidation when the training and retest were completed at 3 pm (as

compared to 10 am). These findings indicate time-of-day effects for the acquisition and

consolidation outcomes in young adults. Although previous research that employed AM/PM

designs in children did not find differences in motor skill acquisition between morning and evening

groups[8], the impact on offline consolidation processes has yet to be systematically and

exhaustively examined. Although certainly speculative, it is possible that the impaired overnight

consolidation that previous studies have found in children could be – at least partially – attributed

to the choice of experimental design. An alternative, albeit related explanation, is that the

beneficial effect of sleep requires additional time to emerge at the behavioral level and thus that

these previous studies did not show overnight gains due to their retest immediately after the sleep

episode (i.e., 12 hours after initial learning). Consistent with this explanation, Desrochers et al.[19]

found that a post-learning nap facilitated the consolidation of a recently acquired motor sequence

in pre-school children, but only after an extended period of time that included a night of sleep.

Additional research is necessary to further disentangle these potential explanations.

Potential neural correlates of enhanced macro-offline processing in children

An open question is: What are the neural correlates of the greater macro-offline performance gains

observed in children? Based on (a) recent evidence in young adults demonstrating that the patterns

of activity observed in the hippocampus and putamen during motor learning are spontaneously

reactivated in macro-offline intervals[58]; and, (b) the known role of reactivation processes in

memory consolidation and longer-term retention[59,60], it is tempting to speculate that the childhood

advantages observed in this research can be attributed to enhanced reactivation of learning-related

patterns of brain activity. This possibility is pure speculation at this point but deserves attention in

future research.

Older adults exhibit impaired overnight consolidation

On the other end of the lifespan, our results showed similar 5-hour but smaller 24-hour offline

gains in the older as compared to the young adults. The comparable 5-hour offline gains are in line

with previous studies that showed a similar consolidation between young and older adults across

an offline period of wakefulness[24,25]. Similarly, the smaller 24-hour offline gains observed in the

older adults are in line with several studies that demonstrated impaired overnight consolidation

with aging[25,26,28]. These studies have largely attributed this deficit to age-related decreases in sleep

quality and quantity, as well as alterations in the sleep-related electrophysiological markers of

plasticity such as sleep spindles and slow waves[22,24] (but see [3,23,25] for discussion of alternative

explanations). Whereas our older adult participants did report shorter sleep duration as compared

to both children and adolescents, no differences in sleep quality were found. However, it is

important to note that these measures were self-reported and previous research has shown that

older adults tend to overestimate their sleep quality[61]. Nonetheless, our results replicate the often

demonstrated finding that aging negatively impacts overnight consolidation of recently acquired

motor sequences.

Considerations and future recommendations

There are some limitations and methodological considerations of the current research that warrant

further attention. *First*, these experiments were conducted online and thus were not monitored by

researchers in real-time. Although we cannot completely discard the possibility that the online

nature influenced our results, it is worth nothing that previous studies that compared online and

in-person experiments in children[62] and young adults[50,63] found comparable results between these

study modalities. For instance, Cubillos et al.[63] showed similar learning curves for the performance

of the finger-tapping task between a supervised-lab and unsupervised-online young adult group.

Additionally, Bönstrup et al.[50] obtained similar micro-offline learning in young adults through an

online crowdsourcing platform. It is also worth noting that our protocol was not implemented in

common crowdsourcing platforms (e.g., Mechanical Turk) that typically acquire full datasets

within hours of posting an experiment and have come under scrutiny for unreliable data[64].

Furthermore, although there are drawbacks to employing web-based data acquisition, a significant

advantage is the recruitment from a large and broad pool of participants, enabling us to include

individuals that otherwise may not have been able to complete our traditional in-person studies.

Second, and specific to experiment 2, each session ended with a pseudorandom run. The

implementation of this random run is often used in sequence learning studies and has the benefit

of assessing sequence specificity of performance improvements[5,6,11,48,65]. However, we cannot rule

out the possibility that the random runs at the end of each session interfered with the macro-offline

processing of the sequential motor memory.

Third, explicit awareness was not assessed at the end of any session or experiment. No explicit

recognition or recall test was implemented due to the online nature of the experiment and the

potential interference with the offline processes in Experiment 2. Nevertheless, the lack of such a

test impedes our ability to draw any conclusions regarding age groups differences in explicit

awareness and how such a difference may affect the learning and memory results.

Fourth, it is important to note that, due to concerns surrounding the acquisition of quality data

from young children, our minimum age for this research was 7 years. Moreover, this youngest age

was relatively under-sampled in our experiments (n = 3 and 1 7-year-olds included in analyses of

Experiments 1 and 2, respectively). Due to difficulties with completing the online motor task,

several 7-year-old children had to be excluded from data analyses. Future studies could consider

implementing certain measures (e.g., monitoring through videocall) to increase the number of

young children that successfully complete the experiments.

Fifth, and as outlined in our pre-registration, the age groups were operationally defined based on

age (i.e., children and adolescents were between 7 and 12, and 13 and 17 years, respectively),

which does not consider the participant's pubertal status. Our questionnaire battery did ask the

parents to indicate whether their child exhibited signs of puberty onset at the time of participation.

Note that all participants 13 years and older (i.e., classified as an adolescent based on age) were

reported to having started puberty. And only a minimal number of child subjects (6 in each of the

experiments) were considered to have initiated puberty. Given the relatively few subjects that

would be re-classified based on parental report of puberty onset, we do not anticipate our results

to substantially change if pubertal status – as opposed to age - was used to demarcate childhood

and adolescence.

6. Conclusion

The current study examined initial motor sequence learning and the time course of motor memory

consolidation across the human lifespan. Our results demonstrated comparable dynamics of

performance improvements during the initial learning session, yet 7- to 12-year-old children

exhibited smaller sequence-specific learning. Children demonstrated the largest performance gains

over macro-offline intervals, suggesting that children exhibit a developmental advantage in the

offline processing of recently practiced movement sequences. The neural underpinnings of this

childhood advantage are not yet known and thus future research is warranted to characterize them.

Data Availability

The source data used for the results presented in this text as well as the raw data are publicly

available without restriction on Zenodo (https://zenodo.org/doi/10.5281/zenodo.8274118).

Code Availability

Customized data processing codes in MATLAB are publicly available without restriction on

Zenodo (https://zenodo.org/doi/10.5281/zenodo.8274118).

Author Contributions

AVR: Methodology, Software, Formal Analysis, Investigation, Resources, Data curation, Writing

– original draft, Visualization, Project administration. **RDB**: Methodology, Writing – Reviewing

and Editing. **GA**: Methodology, Writing - Reviewing and Editing, Supervision. **BRK**:

Conceptualization, Methodology, Resources, Writing – Reviewing and Editing, Supervision,

Project administration.

REFERENCES

1. Albouy, G., King, B. R., Maquet, P. & Doyon, J. Hippocampus and striatum: Dynamics and interaction during acquisition and sleep-related motor sequence memory consolidation. *Hippocampus* **23**, 985–1004 (2013).

2. Doyon, J., Gabitov, E., Vahdat, S., Lungu, O. & Boutin, A. Current issues related to motor sequence learning in humans. *Current Opinion in Behavioral Sciences* **20**, 89–97 (2018).

3. King, B. R., Hoedlmoser, K., Hirschauer, F., Dolfen, N. & Albouy, G. Sleeping on the motor engram: The multifaceted nature of sleep-related motor memory consolidation. *Neuroscience & Biobehavioral Reviews* **80**, 1–22 (2017).

4. Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L. & Haith, A. M. Motor Learning. in *Comprehensive Physiology* (ed. Terjung, R.) 613–663 (Wiley, 2019). doi:10.1002/cphy.c170043.

5. Du, Y., Valentini, N. C., Kim, M. J., Whitall, J. & Clark, J. E. Children and Adults Both Learn Motor Sequences Quickly, But Do So Differently. *Front. Psychol.* **08**, (2017).

6. Salehi, S. K., Sheikh, M., Hemayattalab, R. & Humaneyan, D. The Effect of Different Ages levels and explicit-implicit Knowledge on Motor Sequence Learning. *Internation journal of environmental & science education* **11**, 13157–13165 (2016).

7. Meulemans, T., Van der Linden, M. & Perruchet, P. Implicit sequence learning in children. *J Exp Child Psychol* **69**, 199–221 (1998).

8. Wilhelm, I. *et al.* The sleeping child outplays the adult's capacity to convert implicit into explicit knowledge. *Nat Neurosci* **16**, 391–393 (2013).

9. Adi-Japha, E., Badir, R., Dorfberger, S. & Karni, A. A matter of time: rapid motor memory stabilization in childhood. *Developmental Science* **17**, 424–433 (2014).

10. Janacsek, K., Fiser, J. & Nemeth, D. The best time to acquire new skills: age-related

differences in implicit sequence learning across the human lifespan. *Developmental Science*

15, 496–505 (2012).

11. Lukács, Á. & Kemény, F. Development of Different Forms of Skill Learning Throughout the

Lifespan. *Cognitive Science* **39**, 383–404 (2015).

12. Thomas, K. M. *et al.* Evidence of developmental differences in implicit sequence learning:

an fMRI study of children and adults. *J Cogn Neurosci* **16**, 1339–1351 (2004).

13. Wilhelm, I., Diekelmann, S. & Born, J. Sleep in children improves memory performance on

declarative but not procedural tasks. *Learn. Mem.* **15**, 373–377 (2008).

14. Ashtamker, L. & Karni, A. Motor memory in childhood: Early expression of consolidation

phase gains. *Neurobiology of Learning and Memory* **106**, 26–30 (2013).

15. Fischer, S., Wilhelm, I. & Born, J. Developmental Differences in Sleep's Role for Implicit

Off-line Learning: Comparing Children with Adults. *Journal of Cognitive Neuroscience* **19**,

214–227 (2007).

16. Juhasz, D., Nemeth, D. & Janacsek, K. Is there more room to improve? The lifespan

trajectory of procedural learning and its relationship to the between- and within-group

differences in average response times. *PLoS ONE* **14**, e0215116 (2019).

17. Wilhelm, I., Prehn-Kristensen, A. & Born, J. Sleep-dependent memory consolidation – What

can be learnt from children? *Neuroscience & Biobehavioral Reviews* **36**, 1718–1728 (2012).

18. Zinke, K., Wilhelm, I., Bayramoglu, M., Klein, S. & Born, J. Children's initial sleep-

associated changes in motor skill are unrelated to long-term skill levels. *Developmental*

Science **20**, e12463 (2017).

19. Desrochers, P. C., Kurdziel, L. B. F. & Spencer, R. M. C. Delayed benefit of naps on motor

learning in preschool children. *Exp Brain Res* **234**, 763–772 (2016).

20. Dorfberger, S., Adi-Japha, E. & Karni, A. Reduced Susceptibility to Interference in the

Consolidation of Motor Memory before Adolescence. *PLoS ONE* **2**, e240 (2007).

21. Gualtieri, S. & Finn, A. S. The Sweet Spot: When Children's Developing Abilities, Brains,

and Knowledge Make Them Better Learners Than Adults. *Perspect Psychol Sci* **17**, 1322–

1338 (2022).

22. Bottary, R., Sonni, A., Wright, D. & Spencer, R. M. C. Insufficient chunk concatenation may

underlie changes in sleep-dependent consolidation of motor sequence learning in older

adults. *Learn. Mem.* **23**, 455–459 (2016).

23. Fitzroy, A. B., Kainec, K. A., Seo, J. & Spencer, R. M. C. Encoding and consolidation of

motor sequence learning in young and older adults. *Neurobiology of Learning and Memory*

185, 107508 (2021).

24. Fogel, S. M. *et al.* fMRI and sleep correlates of the age-related impairment in motor memory

consolidation. *Human Brain Mapping* **35**, 3625–3645 (2014).

25. Gudberg, C., Wulff, K. & Johansen-Berg, H. Sleep-dependent motor memory consolidation

in older adults depends on task demands. *Neurobiology of Aging* **36**, 1409–1416 (2015).

26. Spencer, R. M. C., Gouw, A. M. & Ivry, R. B. Age-related decline of sleep-dependent

consolidation. *Learn. Mem.* **14**, 480–484 (2007).

27. Brown, R. M., Robertson, E. M. & Press, D. Z. Sequence Skill Acquisition and Off-Line

Learning in Normal Aging. *PLoS ONE* **4**, e6683 (2009).

28. Wilson, J. K., Baran, B., Pace-Schott, E. F., Ivry, R. B. & Spencer, R. M. C. Sleep modulates

word-pair learning but not motor sequence learning in healthy older adults. *Neurobiology of*

Aging **33**, 991–1000 (2012).

29. Tóth-Fáber, E., Nemeth, D. & Janacsek, K. Lifespan developmental invariance in memory

consolidation: evidence from procedural memory. *PNAS Nexus* **2**, pgad037 (2023).

30. Faul, F., Erdfelder, E., Lang, A.-G. & Buchner, A. G*Power 3: A flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*

Methods **39**, 175–191 (2007).

31. Maclean, A. W., Fekken, G. C., Saskin, P. & Knowles, J. B. Psychometric evaluation of the

Stanford Sleepiness Scale. *Journal of Sleep Research* **1**, 35–39 (1992).

32. Stoet, G. PsyToolkit: A software package for programming psychological experiments using

Linux. *Behavior Research Methods* **42**, 1096–1104 (2010).

33. Stoet, G. PsyToolkit: A Novel Web-Based Method for Running Online Questionnaires and

Reaction-Time Experiments. *Teaching of Psychology* **44**, 24–31 (2017).

34. Dinges, D. F. & Powell, J. W. Microcomputer analyses of performance on a portable, simple

visual RT task during sustained operations. *Behavior Research Methods, Instruments, &*

Computers **17**, 652–655 (1985).

35. Basner, M., Mollicone, D. & Dinges, D. F. Validity and sensitivity of a brief psychomotor

vigilance test (PVT-B) to total and partial sleep deprivation. *Acta Astronautica* **69**, 949–959

(2011).

36. Loh, S., Lamond, N., Dorrian, J., Roach, G. & Dawson, D. The validity of psychomotor

vigilance tasks of less than 10-minute duration. *Behavior Research Methods, Instruments, &*

Computers **36**, 339–346 (2004).

37. King, B. R. *et al.* Schema and Motor-Memory Consolidation. *Psychol Sci* **30**, 963–978

(2019).

38. Reverberi, S., Dolfen, N., Van Roy, A., Albouy, G. & King, B. R. Sleep does not influence

schema-facilitated motor memory consolidation. *PLoS ONE* **18**, e0280591 (2023).

39. Pan, S. C. & Rickard, T. C. Sleep and motor learning: Is there room for consolidation?

Psychological Bulletin **141**, 812–834 (2015).

40. Epskamp, S. & Nuijten, M. B. statcheck: Extract statistics from articles and recompute p-

values. (2018).

41. Wagenmakers, E.-J., Wetzels, R., Borsboom, D. & Van Der Maas, H. L. J. Why

psychologists must change the way they analyze their data: The case of psi: Comment on

Bem (2011). *Journal of Personality and Social Psychology* **100**, 426–432 (2011).

42. Sugawara, S. K. *et al.* Sleep Is Associated with Offline Improvement of Motor Sequence

Skill in Children. *PLoS ONE* **9**, e111635 (2014).

43. Buch, E. R., Claudino, L., Quentin, R., Bönstrup, M. & Cohen, L. G. Consolidation of

human skill linked to waking hippocampo-neocortical replay. *Cell Reports* **35**, 109193

(2021).

44. Jacobacci, F. *et al.* Rapid hippocampal plasticity supports motor sequence learning. *Proc.*

Natl. Acad. Sci. U.S.A. **117**, 23898–23903 (2020).

45. Bönstrup, M. *et al.* A Rapid Form of Offline Consolidation in Skill Learning. *Current*

Biology **29**, 1346-1351.e4 (2019).

46. Bechtold, B., Fletcher, P., Seamusholden & Gorur-Shandilya, S. bastibe/Violinplot-Matlab:

A Good Starting Point. (2021) doi:10.5281/ZENODO.4559847.

47. Jongbloed-Pereboom, M., Nijhuis-van Der Sanden, M. W. G. & Steenbergen, B. Explicit and

implicit motor sequence learning in children and adults; the role of age and visual working

memory. *Human Movement Science* **64**, 1–11 (2019).

48. Thomas, K. M. & Nelson, C. A. Serial Reaction Time Learning in Preschool- and School-

Age Children. *Journal of Experimental Child Psychology* **79**, 364–387 (2001).

49. Szücs-Bencze, L. *et al.* Manipulating the Rapid Consolidation Periods in a Learning Task

Affects General Skills More than Statistical Learning and Changes the Dynamics of

Learning. *eNeuro* **10**, ENEURO.0228-22.2022 (2023).

50. Bönstrup, M., Iturrate, I., Hebart, M. N., Censor, N. & Cohen, L. G. Mechanisms of offline

motor learning at a microscale of seconds in large-scale crowdsourced data. *npj Sci. Learn.*

5, 7 (2020).

51. Gann, M. A., Dolfen, N., King, B. R., Robertson, E. M. & Albouy, G. Prefrontal stimulation

as a tool to disrupt hippocampal and striatal reactivations underlying fast motor memory

consolidation. *Brain Stimulation* **16**, 1336–1345 (2023).

52. Gupta, M. W. & Rickard, T. C. Dissipation of reactive inhibition is sufficient to explain post-

rest improvements in motor sequence learning. *npj Sci. Learn.* **7**, 25 (2022).

53. Gupta, M. W. & Rickard, T. C. *A Quantitative Model that Incorporates Reactive Inhibition

but No Facilitating Consolidation Explains Single-Session Motor Sequence Practice Effects*.

https://osf.io/d5pv9 (2023) doi:10.31234/osf.io/d5pv9.

54. Wilhelm, I., Metzkow-Mészàros, M., Knapp, S. & Born, J. Sleep-dependent consolidation of

procedural motor memories in children and adults: the pre-sleep level of performance

matters. *Developmental Science* **15**, 506–515 (2012).

55. Kuriyama, K., Stickgold, R. & Walker, M. P. Sleep-dependent learning and motor-skill

complexity. *Learn. Mem.* **11**, 705–713 (2004).

56. Truong, C. *et al.* Time-of-day effects on skill acquisition and consolidation after physical and

mental practices. *Sci Rep* **12**, 5933 (2022).

57. Kvint, S. Acquisition and retention of motor sequences: The effects of time of the day and

sleep. *Archives Italiennes de Biologie* 303–312 (2011) doi:10.4449/aib.v149i3.1244.

58. King, B. R., Gann, M. A., Mantini, D., Doyon, J. & Albouy, G. Persistence of hippocampal

and striatal multivoxel patterns during awake rest after motor sequence learning. *iScience* **25**,

105498 (2022).

59. Deuker, L. *et al.* Memory Consolidation by Replay of Stimulus-Specific Neural Activity. *J.*

Neurosci. **33**, 19373–19383 (2013).

60. Tambini, A. & Davachi, L. Persistence of hippocampal multivoxel patterns into

postencoding rest is related to memory. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 19591–19596

(2013).

61. Vitiello, M. V., Larsen, L. H. & Moe, K. E. Age-related sleep change. *Journal of*

Psychosomatic Research **56**, 503–510 (2004).

62. Chuey, A., Boyce, V., Cao, A. & Frank, M. C. *Conducting developmental research online*

vs. in-person: A meta-analysis. https://osf.io/qc6fw (2022) doi:10.31234/osf.io/qc6fw.

63. Cubillos, L. H., Augenstein, T. E., Ranganathan, R. & Krishnan, C. Breaking the barriers to

designing online experiments: A novel open-source platform for supporting procedural skill

learning experiments. *Computers in Biology and Medicine* **154**, 106627 (2023).

64. Webb, M. A. & Tangney, J. P. Too Good to Be True: Bots and Bad Data From Mechanical

Turk. *Perspect Psychol Sci* 174569162211200 (2022) doi:10.1177/17456916221120027.

65. Savion-Lemieux, T., Bailey, J. A. & Penhune, V. B. Developmental contributions to motor

sequence learning. *Exp Brain Res* **195**, 293–306 (2009).

1052
1053

1054   **FIGURE CAPTIONS**
1055
Figure 1. a) Serial reaction time task (SRTT). A stimulus appeared in 1 of 8 spatial locations and the
participant responded with the corresponding key/finger as fast as possible. Stimuli appeared in either a
pseudorandom or sequential (i.e., 4-7-3-8-6-2-5-1) manner. b) To increase motivation and attention, the
task was built around a story in which participants were asked to catch butterflies. c) Design of Experiments
1 and 2. Experiment 1 consisted of the first session only (i.e., Day 1 in blue) and Experiment 2 of all three
sessions displayed. Session 1 included four blocks of pseudorandom SRTT, followed by 20 blocks of
sequential SRTT, which were divided into 16 blocks of training and 4 blocks of test after a 1-minute rest,
and four additional blocks of pseudorandom SRTT. Approximately 5 and 24 hours after session 1, session
2 and 3 were completed, respectively. Both sessions 2 and 3 consisted of 4 blocks of sequential SRTT and
4 blocks of pseudorandom SRTT. Note that the first session of Experiment 2 was constrained between 9am
and 2pm to give ample time for the 5hr delayed retest. Those participants in Experiment 1 were instructed
to complete the session anytime between 9am and 7pm.
1068
1069
Figure 2. a) Average normalized response time for both task variants. b) Average normalized accuracy for
both task variants. For panels a and b, shaded regions represent standard errors of the mean. c) Average
learning magnitude for performance speed per age group. Shaded regions represent the kernel density
estimate of the data, colored circles depict individual data, open circles represent group medians, and the
*horizontal lines depict group means[46]. * p < 0.05 for pairwise group comparisons. d) Learning magnitude*
as a function of age. Quadratic fit from childhood into young adulthood: Learning magnitude = (-
*$0.0008572*age^2$) + (0.04021*age) + (-0.2226). Quadratic fit for older adults: Learning magnitude =*
*($0.00157*age^2$) + (-0.1996*age) + 6.4675.*
1078
1079
*Figure 3. Micro-online (**left**) and -offline (**right**) gains. Panels a and b depict micro gains displayed as a*
function of practice blocks. Panels c and d contain violin plots of micro-online and -offline, respectively,
gains averaged across blocks. Note that pairwise comparisons in these panels included blocks or rest
periods in the statistical models. Shaded regions represent the kernel density estimate of the data, colored
circles depict individual data, open circles represent group medians, and the horizontal lines depict group
means[46]. Panels e and f show averaged micro-online and -offline gains as a function of age from childhood
into young adulthood and within older adulthood. Lines of fit childhood into young adulthood: micro-online
*gains = ($-0.318*age)^{-0.459}$; micro-offline gains = 0.298*exp(-0.377*age). Lines of fit within older*
*adulthood: micro-online gains = (0.003*age) + (-0.238); micro-offline gains = (227.361*age) + (-1.897).*
1089
1090
1091   *Figure 4. Average normalized response time (**a**) and accuracy (**b**) for the three experimental sessions and*
1092   *two task variants of Experiment 2. Shaded regions represent standard errors of the mean. Corresponding*
1093   *statistical analyses are in Appendix 5 of the Supporting Information.*
1094
*Figure 5. Sequential offline gains across the 5-hour (**Panel a**) and 24-hour (**Panel c**) offline periods for the*
four age groups. Shaded regions represent the kernel density estimate of the data, colored circles depict
*individual data, open circles represent group medians, and the horizontal lines depict group means[46]. * p*
*< 0.05 and ~ p < 0.10 for pairwise group comparisons. Five-hour (**Panel b**) and 24-hour (**Panel d**)*
sequential offline gains are plotted as a function of age. Lines of fit from childhood into young adulthood:
*5h gains = (-0.003 * age) + 0.075; 24h gains = ($0.872*age)^{-0.682}$. Lines of fit within older adulthood: 5h*
*gains = ($1.814e^{-6}$ * exp(0.256 * age)) + ($-1.858e^{-6}$ * exp(0.256 * age)); 24h gains = (-0.006 * age) + 0.381.*
For better visualization of the age-related changes, the scale of the y-axes was set to range from -0.4 to 0.4

and thus the young adult with the outlier 5hr sequential gain (visible in Panel a) is not depicted in Panel b.
This individual, however, was included in the fitting procedure. Note that if this individual was excluded
from statistical analyses, the pattern of results observed in Panel a remains largely similar, with the
exception that the pairwise difference between young and older adults becomes statistically significant (i.e.,
older adults exhibit worse offline consolidation over a 5hr wake interval as compared to young adults).