Title

Children exhibit a developmental advantage in the offline processing of a learned motor sequence

Authors

Anke Van Roy[1], Geneviève Albouy[1], Ryan D. Burns[1], Bradley R. King[1, *]

Affiliation

[1] Department of Health and Kinesiology, College of Health, University of Utah, Salt Lake City,

UT 84112, USA

[*] **Corresponding author**

Bradley King, PhD

Department of Health & Kinesiology, College of Health, University of Utah

250 S 1850 E

Salt Lake City, Utah 84112

Bradley.ross.king@utah.edu

Abstract

Research that examines changes in a specific behavior across the human lifespan frequently reports

an inverted-U trajectory. That is, young adults exhibit optimal performance, children are

conceptualized as developing systems progressing towards this ideal state, and older adulthood is

characterized by performance decrements. However, not all behaviors follow this trajectory.

Notably, there are instances – and in the learning and memory domain in particular – in which

children outperform young adults and thus can be considered the model of optimal performance.

This study aimed to reveal instances of childhood advantage in the domain of motor learning and

memory consolidation. Motor sequence learning data were acquired from 7 to 35 and >55-year-

old participants across two experiments via an online data acquisition platform. In Experiment 1

(n=130), participants learned an 8-element, bimanual motor sequence in a single practice session.

In Experiment 2 (n=108), participants acquired the same sequence and were retested 5 and 24

hours after initial acquisition, affording the assessment of memory consolidation across offline

periods including both wakefulness and sleep. Results from Experiment 1 revealed comparable

initial learning dynamics among age groups, but 7- to 12-year-old children exhibited smaller

sequence-specific learning relative to adolescents (13-17 years), young adults (18-35 years) and

older adults (> 55 years). Interestingly, children tended to demonstrate the greatest performance

gains over the short rest epochs interspersed with practice (i.e., micro-offline intervals). Results

from Experiment 2 revealed that this developmental advantage was not limited to the micro-offline

timescale. Children and adolescents exhibited larger performance gains in the delayed retests,

reflecting enhanced "macro"-offline motor memory consolidation. Collectively, these results

suggest that children exhibit a developmental advantage in the offline processing of recently

practiced motor sequences.

1. Introduction

The ability to learn novel movements is vital to the everyday functioning of individuals across the human lifespan (e.g., from a child learning to ride a bike to an older individual learning to operate a touchscreen). One of the most dominant motor learning paradigms in the field is motor sequence learning (MSL), whereby participants acquire a novel series of interrelated actions (i.e., a motor sequence). MSL has been extensively studied in healthy young adults (~18-35 years), facilitating the development of a framework that characterizes the time course of learning in this age group. Typically, a fast online (i.e., during repeated practice) acquisition phase with substantial performance improvements occurs during an initial training session. This is followed by a slow offline (i.e., in the absence of active task practice) consolidation phase that spans several hours during which the acquired memory trace is stabilized, strengthened and reorganized into a more robust form. Interestingly, recent research has indicated that offline consolidation is also evident on a much shorter timescale. More specifically, young adults exhibit large performance gains over the short rest intervals in between blocks of practice within a session, providing evidence for a fast consolidation process during these micro-offline epochs [Bönstrup et al., 2019; Bönstrup et al., 2020; Buch et al., 2021; Gann et al., 2023; Jacobacci et al., 2020; but see Gupta & Rickard, 2022 for opposing view].

Although previous research has examined motor sequence learning in school-aged children (i.e., between 5 and 12 years), the results are heterogenous. While some studies have reported comparable learning within the initial practice session in children and adults (Du et al., 2017; Salehi et al., 2016; Meulemans , Van der Linden & Perruchet, 1998; Wilhelm et al., 2013), others have shown that children exhibit worse (Adi-Japha et al., 2014; Janacsek et al., 2012; Lukács &

Kemény, 2015; Thomas et al., 2004; Wilhelm et al., 2008) or better learning (Ashtamker & Karni,

2013; Fischer et al., 2007; Janacsek et al., 2012; Juhasz et al., 2019; Savion-Lemieux et al., 2009;

Wilhelm, Prehn-Kristensen & Born., 2012). There is a general consensus, however, that children

exhibit impaired consolidation of a recently acquired motor sequence over offline periods that

include sleep (e.g., Fischer et al., 2007; Wilhelm et al., 2008, 2013; Zinke et al., 2017).

Specifically, whereas post-learning sleep is known to boost offline consolidation processes in

young adults [see King et al., 2017 for a review], such a beneficial effect is absent in children.

Interestingly, there is some evidence suggesting that this degraded consolidation over sleep may

be the by-product of enhanced or accelerated consolidation over the wake epochs shortly following

initial learning (Adi-Japha et al., 2014; Ashtamker & Karni, 2013; Dorfberger et al., 2007).

Specifically, children showed a rapid performance stabilization and enhancement 15 minutes (Adi-

Japha et al., 2014) and 1 hour (Ashtamker & Karni, 2013) after learning, which were not observed

in young adults. Interestingly, this enhanced offline processing does not appear to be limited to the

timescale of hours following initial learning (herein referred to as *macro*-offline intervals) but is

also observed on shorter timescales. Specifically, Du et al. (2017) found that young children (i.e.,

6-year-olds) showed higher *micro*-offline gains as compared to older children (i.e., 10-year-olds)

and young adults. These results collectively suggest that young children may exhibit a

developmental advantage with respect to the offline processing of recently practiced motor

sequences during wake intervals. This childhood advantage could then be added to the list of

examples in which children outperform adults in specific learning and memory behaviors (for

review see Gualtieri & Finn, 2022).

At the other end of the lifespan, there is considerable evidence that older adults (> ~60 years)

exhibit comparable initial learning of a novel motor sequence as young adults (Bottary et al., 2016;

Fitzroy et al., 2021; Fogel et al., 2017; Fogel et al., 2014; Gudberg et al., 2015; Spencer et al.,

2007; Vien et al., 2016). And, whereas children appear to have a developmental advantage in the

offline processing of recently acquired sequences, older adults are known to exhibit deficits.

Specifically, the magnitude of offline gains over macro-offline intervals that include sleep and/or

wake is significantly less than those observed in younger adults (Bottary et al., 2016; Fogel et al.,

2017; Fogel et al., 2014; Gudberg et al., 2015; Spencer et al., 2007; Wilson et al., 2012).

Although the existing literature suggests that the offline processing of recently acquired movement

sequences appears to be superior in children and then declines throughout adulthood, systematic

examinations into changes across the human lifespan are relatively limited (see Janacsek et al.,

2012; Juhasz et al., 2019 for examples in the initial learning of probabilistic sequences). This study

thus aimed to provide an exhaustive characterization of motor learning and memory consolidation

behaviors across the lifespan (i.e., in groups of children, adolescents, young adults and older

adults). Consistent with Du et al., 2017; Salehi et al., 2016; Meulemans, Van der Linden &

Perruchet, 1998; and Wilhelm et al., 2013, we hypothesized that the initial learning of a motor

sequence (i.e., within a single training session) would be comparable across the 4 age groups. We

expected children to exhibit enhanced micro- and 5-hour macro-offline consolidation, indicative

of a developmental advantage in the offline processing of a recently acquired motor sequence

across periods of wakefulness. This developmental advantage in macro-offline consolidation was

expected to be absent at the 24-hour retest, as previous research has indicated impaired sleep-

facilitated consolidation in children (Fischer et al., 2007; Wilhelm et al., 2008, 2013; Zinke et al.,

2017). Last, and consistent with earlier studies, older adults were expected to demonstrate intact

initial learning but impaired macro-offline consolidation across both 5- and 24-hour intervals.

2. Methods and materials

The project consisted of two experimental protocols. Specifically, Experiments 1 and 2 examined

the effect of age on the initial acquisition of a movement sequence and the time course of motor

memory consolidation, respectively. Data collection and analysis plans were pre-registered via

Open Science Framework and can be accessed at https://doi.org/10.17605/OSF.IO/WBK9H and

https://doi.org/10.17605/OSF.IO/ZMJ75. Any additional analyses that were not included in the

pre-registrations are labelled in this text as exploratory. The source data used for the results

presented in this text as well as the raw data are publicly available on Zenodo

(https://doi.org/10.5281/zenodo.8274119).

2.1 Participants

Healthy volunteers between 7-35 or 55-75 years old of all genders were recruited by

advertisements on relevant websites and research databases. Note that as our experiments were

conducted through a web-based data collection platform, recruitment was not constrained to a

specific geographical region. Individuals interested in participation were sent an online screening

questionnaire to assess eligibility, which was subsequently reviewed by a member of the research

team. Exclusion criteria were: 1) reported history of medical, neurological, psychological or

psychiatric conditions, 2) use of psychoactive or sleep-influencing medications, 3) indications of

abnormal or irregular sleep, 4) mobility limitations of the fingers or hands, or 5) considered a

professional typist or prior extensive training on a musical instrument requiring dexterous finger

movements (e.g., piano, guitar). All experimental procedures were approved by the local Ethics

Committee (IRB_00136894). Adult participants and parents of child participants gave informed

consent and participants below 18 years of age provided informed assent. Participants received

compensation for their participation.

A convenience sample of 224 individuals met the inclusion criteria and thus initiated participation

in an experimental protocol. They were divided into the following four age groups: children

(operationally defined as 7-12 years old), adolescents (13-17 years old), young adults (18-35 years

old) and older adults (≥ 55 years old). Pre-registered analyses compared differences among these

age *groups*. Yet, to provide more fine-grained analyses of age-related trajectories of motor learning

and memory consolidation behaviors, we also conducted exploratory analyses with age as a

continuous variable (see below for details).

Sample size computations were conducted *a priori* and with the software G*Power (Erdfelder et

al., 2007). For the assessment of age-group differences in initial motor sequence learning (i.e.,

Experiment 1), and to detect an effect size of $f = 0.3$ (based on the comparison of initial learning

between 9-year-olds and young adults in Adi-Japha et al., 2014), with an alpha of 0.05 and power

of 0.80, the desired sample size was 128 subjects (32 per group). Data from 7 participants were

excluded from data analysis due to a failure to comply to experimental instructions (e.g., repeatedly

pressing the same key; $n = 3$ children), missing data because of software issues ($n = 1$ young adult),

or a failure to correctly perform the motor task (i.e., statistical outliers (> 3SD from group mean)

on sequence accuracy; $n = 1$ child, 1 young adult, 1 older adult). Excluded participants were

replaced and thus the final sample size for analyses consisted of 130 participants (see Table 1 for

participant demographics).

For the assessment of age-group differences in macro-offline consolidation processes (i.e.,

Experiment 2), the detection of an effect size of f = 0.20 (slightly more conservative than group

Adi-Japha et al., 2014), with an alpha of 0.05, a power of 0.80 and a correlation among repeated

measurements of 0.25, this experiment required a sample size of 108 (27 participants per group).

In total, 21 individuals were excluded from data analysis. Specifically, 17 participants failed to

comply to experimental instructions (i.e., 4 children repeatedly pressed the same key and 5

children, 2 young adults and 6 older adults did not adhere to the specific schedule of the

experimental sessions). Additional exclusions were due to inaccurate data because of software

issues (n = 1 older adult), a lack of performance improvements across training (n = 1 young adult,

1 older adult), and an inability to correctly perform the motor task (statistical outliers on sequence

accuracy; n = 1 young adult). Similar to above, excluded participants were replaced and thus the

final sample for analyses consisted of 108 participants (see Table 2 for participant details). Note

that data from the third session of one young adult and the post-learning random data of one

adolescent were missing; thus, these individuals were excluded from the contrasts involving these

specific task runs only.

Note that the procedures of Experiment 1 (assessing initial motor sequence learning) were identical

to the first session of Experiment 2 (assessing macro-offline consolidation). Accordingly, data

from a subset of early participants from Experiment 2 (n = 42; 16 children, 19 adolescents, 4 young

adults, 3 older adults) were included in the analyses of Experiment 1. Thus, these participants were

included in analyses for both experiments.

Table 1: Participant demographics, and sleep and vigilance scores for each age group in Experiment 1

Variable	Children	Adolescents	Young adults	Older adults
n	33	33	32	32
Female *(n)*	17	12	23	24
Age (years)	10.2 (1.6)	15.3 (1.3)	26.3 (5.0)	64.4 (5.2)
M/E Preference	2.9 (0.9)	3.3 (0.8)	2.9 (1.0)	2.3 (0.8)
Time of testing (SD in min)	12:47 (172)	12:29 (133)	14:52 (175)	12:05 (169)
Sleep quality	5.1 (0.8)	5.0 (0.7)	4.8 (0.6)	4.8 (0.8)
Sleep duration (hours)	9.8 (0.9)	9.1 (1.4)	8.5 (1.0)	7.7 (1.1)
SSS score	2.0 (0.9)	2.0 (0.8)	2.0 (0.7)	1.5 (0.7)

Numbers represent the mean, with standard deviation in parentheses. Morningness/Eveningness (M/E)
Preference was added as an exploratory variable to assess age-related differences in circadian preferences
and was defined on a 5-point Likert scale (i.e., 1 = extreme morning person, 5 = extreme evening person).
The average time of testing is specified in 24-hour time notation, with the SD in minutes. Sleep quality is
defined on a 6-point Likert scale (i.e., 1 = very bad, 6 = very good). SSS = Stanford Sleepiness Scale
(Maclean et al., 1992), with higher numbers indicative of increased sleepiness. See Appendix 1 as part of
the Supporting Information for corresponding statistical analyses.

***Table* 2:** *Participant demographics, and sleep and vigilance scores for each age group in Experiment 2.*

Variable		Children	Adolescents	Young adults	Older adults
n		27	27	27	27
Female *(n)*		16	12	21	18
Age (years)		10.6 (1.6)	15.4 (1.4)	26.6 (4.4)	63.0 (4.4)
M/E Preference		3.0 (0.8)	3.2 (0.9)	3.2 (0.8)	2.4 (0.9)
Session 1	*Time of testing*	11:34 (97)	11:34 (96)	11:39 (82)	11:30 (92)
	Sleep quality	4.9 (0.7)	5.1 (0.6)	5.0 (0.6)	4.7 (0.8)
	Sleep duration (hrs.)	9.7 (1.1)	9.4 (1.2)	8.3 (1.1)	7.7 (1.3)
	SSS score	2.2 (0.9)	2.1 (0.8)	2.0 (0.7)	1.5 (0.7)
	PVT score (ms)	360.9 (81.5)	341.9 (102.3)	320.9 (62.0)	327.5 (75.7)
Session 2	*Time of testing*	16:42 (99)	16:56 (112)	16:43 (89)	16:29 (96)
	Offline period (hrs.)	5.1 (0.8)	5.4 (0.9)	5.1 (1.1)	5.0 (0.7)
	SSS score	2.0 (1.0)	1.8 (0.8)	2.0 (0.7)	1.7 (0.7)
	PVT score (ms)	386.0 (88.4)	318.5 (61.0)	320.3 (69.1)	323.6 (68.0)
Session 3	*Time of testing*	11:57 (148)	12:25 (156)	11:30 (97)	11:24 (109)
	Offline period (hrs.)	24.4 (1.8)	24.8 (1.9)	23.8 (1.4)	23.9 (1.4)
	Sleep quality	5.0 (0.8)	4.7 (0.8)	4.7 (0.7)	4.6 (0.7)
	Sleep duration (hrs.)	10.3 (0.9)	9.3 (1.5)	8.0 (1.3)	7.6 (1.0)
	SSS score	1.9 (0.7)	2.3 (1.3)	2.3 (0.8)	1.5 (0.6)
	PVT score (ms)	378.7 (97.3)	325.8 (74.6)	308.0 (54.4)	306.3 (33.6)

188
Numbers represent the mean, with standard deviation in parentheses. Morningness/Eveningness (M/E)
Preference was added as an exploratory variable to assess age-related differences in circadian preferences
and is defined on a 5-point Likert scale (i.e., 1 = extreme morning person, 5 = extreme evening person).
The average time of testing is specified in 24-hour time notation, with the SD in minutes. The offline periods
represent the time periods between session 1 and sessions 2 and 3, respectively. Sleep quality is defined on
a 6-point Likert scale (i.e., 1 = very bad, 6 = very good). SSS = Stanford Sleepiness Scale (Maclean et al.,
1992), with higher numbers indicative of increased sleepiness. The PVT score indicates the average simple
RT across the 30 trials. See Appendix 1 as part of the Supporting Information for corresponding statistical
analyses.
198

2.2. General Experimental Procedures

After verification of eligibility, participants were contacted via email with the necessary

information to continue participation. This email contained a link to a video created by our study

team that provided detailed instructions on how to complete our motor learning task (details on

the task are provided in Section 2.3. below). Participants were also provided links to the web-based

data collection platform PsyToolkit (Stoet, 2010, 2017). As our experiments were conducted

online, participants were free to choose which day(s) to complete experiments but were instructed

to complete the protocols on days in which they were well-rested and could adhere to the schedule

of the sessions. The precise time-of-day that the experiments were completed was constrained and

followed experiment-specific schedules provided below (see Sections 2.4 and 2.5). Participants

were instructed to receive a good night of sleep and to refrain from alcohol the night before and

throughout the experiment. Compliance to these instructions was verified through a brief

questionnaire prior to completion of the motor task. Lastly, parents of the child participants were

asked to be present when their child completed the experiments.

For both experiments and at the beginning of each experimental session, a brief questionnaire was

completed in which participants reported their sleep patterns in the previous 24 hours and their

subjective feelings of alertness (Stanford Sleepiness Scale; Maclean et al., 1992). Additionally,

and for Experiment 2 only, participants completed the Psychomotor Vigilance Task (PVT; Dinges

& Powell, 1985) to provide an objective measure of vigilance at the time of testing. The PVT

required participants to fixate on the middle of the computer screen and press the space bar as fast

as possible when a visual stimulus appeared after a varying delay interval. Response time (i.e.,

time between appearance of stimulus and keypress) was logged and used to assess vigilance (i.e.,

higher RTs indicative of lower vigilance). We administered a shortened version of the PVT

(Basner et al., 2011; Loh et al., 2004) that consisted of 35 trials, the first 5 of which were discarded

as participants became familiar with the task.

Figure 1. A) Serial reaction time task (SRTT). A stimulus appeared in 1 of 8 spatial locations and the participant responded with the corresponding key/finger as fast as possible. Stimuli appeared in either a pseudorandom or sequential (i.e., 4-7-3-8-6-2-5-1) manner. B) To increase motivation and attention, the task was built around a story in which participants were asked to catch butterflies. C) Design of Experiments 1 and 2. Experiment 1 consisted of the first session only (i.e., Day 1 in blue) and Experiment 2 of all three sessions displayed. Session 1 included four blocks of pseudorandom SRTT, followed by 20 blocks of sequential SRTT, which were divided into 16 blocks of training and 4 blocks of test after a 1-minute rest. Approximately 5 and 24 hours after session 1, session 2 and 3 were completed, respectively. Both sessions 2 and 3 consisted of 4 blocks of sequential SRTT and 4 blocks of pseudorandom SRTT. Note that the first session of Experiment 2 was constrained between 9am and 2pm to give ample time for the 5hr delayed retest. Those participants in Experiment 1 were instructed to complete the session anytime between 9am and 7pm.

2.3 Serial Reaction Time Task (SRTT)

All participants performed an adapted version of the explicit Serial Reaction Time Task (SRTT) similar to our previous research (King et al., 2019; Reverberi et al., 2023) which was coded in the online data acquisition platform PsyToolkit (Stoet, 2010, 2017). The task consisted of an 8-choice reaction time task in which participants were instructed to react to cues shown on a screen (see Figure 1A). Eight squares that spatially corresponded to the eight fingers used to perform the task (i.e., no thumbs) and the eight keys on the keyboard were presented on the screen. The color of the

squares alternated between red and green, indicating rest and practice blocks, respectively. During

practice, a visual stimulus (i.e., a butterfly) appeared consecutively in one of the squares and

participants responded to the visual cues by pressing the corresponding key with the corresponding

finger as fast and accurately as possible. The task was presented as a game, designed to increase

motivation in child participants, in which participants of all four age groups were instructed to

move quickly to catch as many butterflies as possible (see Figure 1B). The visual cues succeeded

each other in either a pseudorandom or a sequential order, depending on the experimental run (i.e.,

pseudorandom SRTT vs. sequential SRTT, respectively). In the sequential SRTT, participants

were aware that the stimuli and keypresses followed a deterministic sequential pattern (i.e., 4-7-3-

8-6-2-5-1, in which 1 through 8 are the left pinky to the right pinky fingers from left to right), but

they were not given any information about the structure or length of the sequence. During the

pseudorandom SRTT runs, there was no repeating sequence, but each key was pressed once every

eight elements (i.e., no repeating elements). For both pseudorandom and sequential task variants,

practice blocks contained 48 key presses (i.e., 6 repetitions of the 8-element sequence in the

sequence condition) and were separated by 15-second rest periods.

Measures reflecting performance speed (i.e., response time between visual cue and keypress) and

accuracy (i.e., correct or incorrect keypress) of the participants' motor responses were recorded

for each trial and used to assess performance on the task. Per our pre-registrations, individual trials

(i.e., responses) were excluded from analyses if the measured response time was greater than 3

standard deviations above or below the participant's mean response time for that block. For

Experiments 1 and 2, an average of 1.78% (children: 2.03%, adolescents: 1.76%, young adults:

1.68%, older adults: 1.65%) and 1.69% (children: 2.0%, adolescents: 1.83%, young adults: 1.58%,

older adults: 1.34%) of trials were excluded from analyses, respectively.

2.4 Experiment 1 Design

The first experiment consisted of a single experimental session in which participants performed

both the random and sequential variants of the SRTT (see Day 1 [blue] of Figure 1C). Participants

were instructed to complete this session between 9am and 7pm (see Table 1 in the main text and

Appendix 1 as part of the Supporting Information (Table S1 and Figure S1) for details on time of

testing). The experimental session started with one block of familiarization (pseudorandon; data

not analyzed) followed by four blocks of the pseudorandom SRTT (pre-learning random), which

afforded an assessment of general motor execution on the task prior to any sequence learning.

Subsequently, participants completed 20 blocks of the sequential SRTT. These blocks were

divided into two runs: a training run that consisted of 16 blocks and then a post-learning test run

of 4 blocks. The post-learning test run was completed approximately one minute after the end of

training, affording the assessment of end-of-training performance following the further dissipation

of mental and physical fatigue (Pan & Rickard, 2015). Lastly, four additional blocks of the

pseudorandom SRTT (post-learning random) were completed, allowing us to assess whether

improvements in performance on the sequential SRTT could be attributed to learning of the

movement sequence *per se* (i.e., sequence-specific learning) or simply a general improvement in

movement speed due to task familiarization.

2.5 Experiment 2 Design

The second experiment consisted of three experimental sessions (see Figure 1C). Participants were

instructed to complete the first session between 9 am and 2 pm. This session was identical to what

was described for Experiment 1. The second session took place approximately five hours after the

first session, allowing the assessment of motor memory consolidation after an interval of post-

learning wakefulness. This second session consisted of four blocks of sequential SRTT during

which participants again performed the same specific sequence as in session 1, and four blocks of

pseudorandom SRTT (post-test random). Finally, the third session was completed 24 hours after

the first session, allowing the examination of motor memory consolidation after a night of sleep.

Similar to the second session, this third session included four blocks of sequential SRTT and four

blocks of pseudorandom SRTT (post-test random). See Table 2 in the main text and Appendix 1

as part of the Supporting Information (Table S2) for details on actual time of testing. For the

intervals between sessions, participants were instructed to refrain from napping, consuming

alcohol or recreational drugs and to receive a good night of sleep.

2.6 Data Processing and Statistical Analyses

We first describe general data processing and statistical analysis information prior to providing

details specific to each of the two experiments. All statistical analyses described below were

performed using IBM SPSS Statistics for Windows, version 28 (IBM, Armonk , NY, USA). All

statistical tests were performed for both normalized outcome measures (i.e., reaction time and

accuracy) and were considered significant when $p < 0.05$. For the sake of completeness, we also

highlight those effects when $0.05 < p < 0.1$ as non-significant trends. In the event of a violation of

the sphericity assumption, Greenhouse-Geisser correction was applied. All significant main effects

and interactions were followed up by pairwise comparisons and one-way ANOVAs, respectively.

Depending on the statistical test, eta-squared, partial eta-squared or Hedges' g are reported as effect

size measures. The correspondence among reported p-values, test statistics and degrees of freedom

reported in this main text were verified via statcheck (Epskamp & Nuijten, 2018).

To assess potential differences among age groups and experimental sessions in certain

demographic/participant characteristics, one-way ANOVAs or chi-square tests (Experiment 1) and

two-way ANOVAs (Experiment 2) were performed. These measures included gender, vigilance

(both subjective and objective) at the time of testing, self-reported sleep quantity and quality of

the night prior to the experimental session(s), and the time of day in which the motor task or session

was completed. Any significant effects from these analyses were followed up by pairwise

comparisons to determine which age groups or sessions differed from each other. Group means

for these dependent measures are provided in Tables 1 and 2 and results from the corresponding

statistical analyses can be found in Appendix 1 of the Supporting Information (Tables S1, S2;

Figures S1 and S2).

For the SRTT, the averaged response times for correct keypresses and the percentage of correct

keypresses were computed for each block and task variant. Additionally, as previous research has

indicated age differences in motor performance independent of sequence learning (Bottary et al.,

2016; Fitzroy et al., 2021; Kavos Salehi et al., 2016; Meulemans et al., 1998; Sugawara et al.,

2014), baseline differences in performance were accounted for by normalizing both movement

speed and accuracy of the sequential and post-learning random SRTT runs relative to the outcome

variables of the pre-learning random run. More specifically, for each outcome measure and each

block of the SRTT runs, performance was divided by the mean outcome measure across the four

blocks of the pre-learning random run. Statistical analyses of these normalized measures are

presented in the main text. For completeness, results based on non-normalized performance

measures are reported in Appendix 2 of the Supporting Information (Tables S3, S4; Figures S3,

S4).

2.6.1 Experiment 1

Sequential learning dynamics (i.e., block-to-block performance changes) were compared among

age groups using a mixed ANOVA, with *group* (i.e., children, adolescents, young adults, older

adults) as between-subject and *block* (i.e., 16 training blocks or 4 test blocks) as within-subject

factors.

To assess whether observed improvements in performance across practice blocks reflected

sequence-specific learning (as opposed to general skill learning as a result of familiarization with

the motor task), a learning magnitude measure was computed for performance speed. Specifically,

the difference between averaged normalized performance across the four blocks in the post-

learning test run of the sequential SRTT and the averaged normalized performance in the post-

learning random run was divided by the average normalized performance in the post-learning

random run. This measure thus reflects the relative difference between the random and sequential

task variants at the end of training, affording the assessment of sequence-specific learning. Larger

values are indicative of greater sequence-specific learning. The effect of age *group* on learning

magnitude was assessed with a one-way ANOVA. To provide more fine-grained analyses of the

age-related changes in sequence-specific learning magnitude, we conducted exploratory analyses

with age as a continuous variable. Specifically, we tested five potential fit options (i.e., single

exponential, double exponential, linear, quadratic, and power functions) to characterize the

relationship between age and learning magnitude. As we did not acquire data from participants

between 35 and 55 years of age, separate models were built to characterize age-related changes

from 7 to 35 and 55 to 75 years of age. The model with the lowest Akaike Information Criterion

value was deemed as the best fitting model and the corresponding results are presented.

Similar to previous research (Buch et al., 2021; Du et al., 2017; Jacobacci et al., 2020), we

distinguished between micro-online (i.e., within blocks of task practice) and micro-offline (i.e.,

across rest epochs in between blocks of practice) performance improvements. Micro-online

changes were computed as the difference between the averaged normalized RT for the correct

keypresses of the first and last sequence repetitions within each block (i.e., first sequence block n

– last sequence block n). For micro-offline changes, the difference between the averaged

normalized RT of the last sequence repetition of one block and the first sequence repetition of the

subsequent block (i.e., last sequence block n – first sequence block n+1) was computed. These

micro-online and micro-offline measures were analyzed with separate *group(4)* x *block(16)* or

group(4) x *rest block (15)* mixed model ANOVAs, respectively. Moreover, and following the same

exploratory procedure as described above for the learning magnitude measure, we assessed age-

related changes in these micro measures with age as a continuous variable.

2.6.2. Experiment 2

As in Experiment 1, learning dynamics (i.e., block-to-block performance changes) across training

and the post-learning test were analyzed using a *group(4)* x *block(16 or 4)* mixed ANOVAs to

assess potential group differences in initial encoding of the motor sequence within the first session

(i.e., before the assessment of consolidation).

The main focus of Experiment 2 was the effect of age on motor memory consolidation across the

5hr and 24hr macro-offline intervals. To this end, and for both the sequential and random SRTT,

a macro-offline change measure was computed for each outcome variable and offline interval.

Specifically, sequential offline changes were calculated as the difference between the average

normalized performance on the sequential SRTT run in the 5hr and 24hr retest sessions and the

average normalized performance in the post-learning test run of the first session. Similarly, offline

changes for the random task variant were computed as the difference between the averaged

normalized performance on the post-test random runs in the 5hr and 24hr retest sessions and the

averaged normalized performance in the post-learning random run of the first session. These

measures reflect the offline changes in sequential performance and general motor execution from

the end of the initial training session to the two retest sessions, respectively. Mixed ANOVAs with

group (4) as between-subject and *offline period* (i.e., 5-hour and 24-hour offline periods) as within-

subject factors were conducted. Consistent with the exploratory analyses described above for

Experiment 1, age-related changes in the macro-offline performance gains were also assessed with

age as a continuous variable.

Last, and similar to previous research (Bönstrup et al., 2019; Jacobacci, Armony, Yeffal, Lerner,

Amaro Jr., et al., 2020), we assessed the relationship between offline changes in performance on

the micro- (i.e., during the rest epochs in between practice blocks during training) and macro-

offline timescales (i.e., during the 5h and 24h offline periods). Specifically, we performed

exploratory multiple regressions with the micro-offline gains as independent and the 5h or 24h

macro-offline gains as dependent variables. To examine how these relationships differed among

age groups, we included age group (dummy coded) x micro-offline gains interaction variables in

the regression models. The young adult group was selected as the reference group and thus the

interaction variables compare the regression slopes of the other 3 age groups with those of the

young adults.

4. Results

4.1 Experiment 1: Initial motor sequence learning

4.1.1 Participant characteristics, sleep and vigilance

Group means from participant characteristics, the time of when the experimental session was

completed, self-reported sleep quality and duration of the night preceding the experimental

session, and a subjective measure of vigilance are provided in Table 1. Results from the

corresponding statistical analyses and a brief discussion can be found in Appendix 1 of the

Supporting Information (see Figure S1 and Table S1).

4.1.2 Dynamics of Performance Improvements

Normalized performance measures on the SRTT are depicted in Figure 2A and 2B. Performance

accuracy remained stable across training, as evidenced by no main effect of block ($F_{(7.011,883.403)}$ =

0.501, p = 0.835, partial η^2 = 0.004), and did not significantly differ among age groups ($F_{(3,126)}$ =

0.322, p = 0.810, partial η^2 = 0.008). Furthermore, no significant group x block interaction was

found ($F_{(21.033,883.403)}$ = 0.0.715, p = 0.821, partial η^2 = 0.017). Similarly, no block ($F_{(2.283,287.681)}$ =

1.523, p = 0.217, partial η^2 = 0.012), group ($F_{(3,126)}$ = 1.796, p = 0.151, partial η^2 = 0.041), or

interaction effects ($F_{(6.850, 287.681)} = 0.814$, p = 0.574, partial η^2 = 0.019) were found for the post-

learning test.

431
Figure 2. A*) Average normalized response time for both task variants. **B**) Average normalized accuracy*
for both task variants. For panels A and B, shaded regions represent standard errors of the mean. ***C****)*
*Average learning magnitude for performance speed per age group. * p < 0.05 for pairwise group*
comparisons. ***D****) Learning magnitude as a function of age. Quadratic fit from childhood into young*
adulthood: Learning magnitude = (-0.0008572*age^2) + (0.04021*age) + (-0.2226). Quadratic fit for older
adults: Learning magnitude = (0.00157*age^2) + (-0.1996*age) + 6.4675.
438

For an assessment of performance speed, response times significantly decreased across sequential

practice blocks, as shown by a main effect of block ($F_{(7.900, 995.358)} = 64.246$, p < 0.001, partial η^2 =

0.338). These performance improvements were similar among age groups, as indicated by the lack

of a significant group effect ($F_{(3,126)} = 1.578$, p = 0.198, partial η^2 = 0.036) and group x block

interaction ($F_{(23.699, 995.358)} = 1.337$, p = 0.130, partial η^2 = 0.031). During the post-learning test run,

there was no significant block effect ($F_{(2.102, 264.890)} = 0.404$, p = 0.678, partial η^2 = 0.003), indicating

a performance plateau was reached. Moreover, consistent with the training run, there were no

significant differences among groups (Group main effect: $F_{(3,126)} = 1.099$, p = 0.352, partial η^2 =

0.025; Group x Block interaction: $F_{(6.307, 264.890)} = 1.148$, $p = 0.335$, partial $\eta^2 = 0.027$). Collectively,

the assessment of performance across blocks of practice indicates that the four groups acquired the

novel motor sequence and exhibited similar performance improvements.

4.1.3 Sequence-specific learning

A learning magnitude measure was calculated for performance speed that reflects the relative

difference between the random and sequential task variants at the end of training and therefore

affords the assessment of the magnitude of sequence-specific learning (see Figure 2C). Results

revealed a significant group effect ($F_{(3,129)} = 5.440$, $p = 0.002$, $\eta^2 = 0.115$). More specifically, a

significantly smaller sequence-specific learning magnitude was observed in children as compared

to adolescents ($p = 0.014$, $G = 0.714$) and young adults ($p = 0.001$, $G = 0.970$). Follow-up analyses

indicated that the difference in learning magnitude could be attributed to significant group

differences in response time for the post-learning random run ($F_{(3,126)} = 5.795$, $p < 0.001$, partial

$\eta^2 = 0.121$). Children demonstrated greater improvements in performance speed from the pre-

learning to post-learning *random* runs as compared to young ($p < 0.001$, $G = 0.876$) and older

adults ($p = 0.022$, $G = 0.591$). Accordingly, a portion of the performance improvements across the

sequential training blocks in children can be attributed to general learning of the task that is *not*

specific to the sequence.

We also conducted exploratory regression analyses with age as a continuous variable to provide a

more fine-grained analyses of the age-related trajectories of sequence-specific learning magnitude.

Consistent with the differences among *groups*, age was a significant predictor of learning

magnitude from childhood into young adulthood (see Figure 2D; the best fit model was a quadratic

function: $R^2 = 0.156$, $F_{(1,95)} = 8.808$, $p = 0.004$). Specifically, learning magnitude increased across

childhood, reaching peak values at approximately 24 years of age, and then decreased thereafter.

Age was also a significant predictor of learning magnitude within the older adult group, exhibiting

an approximate U-shaped trajectory from 55 to 75 years (quadratic model: $R^2 = 0.247$, $F_{(1,29)} =$

4.746, $p = 0.038$).

4.1.4 Micro-learning across sequential training

For micro-online learning, there was a significant block effect ($F_{(11.524,1429.030)} = 2.358$, $p = 0.006$,

partial $\eta^2 = 0.019$) and a non-significant trend for a group effect ($F_{(3,124)} = 2.663$, $p = 0.051$, partial

$\eta^2 = 0.061$), but no block x group interaction ($F_{(34.573,1429.030)} = 1.159$, $p = 0.243$, partial $\eta^2 = 0.027$;

see Figure 3A). For the sake of completeness, we conducted follow-up pairwise comparisons on

the trend for an effect of age group. Results indicated significantly smaller micro-online gains in

children as compared to young ($F_{(1,61)} = 5.211$, $p = 0.026$, partial $\eta^2 = 0.079$; see Figure 3C) and

older ($F_{(1,61)} = 6.240$, $p = 0.015$, partial $\eta^2 = 0.093$) adults. Consistent with these group

comparisons, there was a non-significant trend for age being a predictor of micro-online gains

from childhood into young adulthood (power model: $R^2 = 0.032$, $F_{(1,96)} = 3.141$, $p = 0.080$; see

Figure 3E), but not within the older adult group (power model: $R^2 = 0.045$, $F_{(1,30\}} = 1.424$, $p =$

0.242).

Micro-offline learning also showed a non-significant trend for a group effect ($F_{(3,124)} = 2.490$, $p =$

0.063, partial $\eta^2 = 0.057$), but no rest block ($F_{(11.662,1446.066)} = 1.209$, $p = 0.272$, partial $\eta^2 = 0.010$;

see Figure 3B) nor a rest block x group interaction effect ($F_{(34.985,1446.066)} = 0.836$, $p = 0.739$, partial

$\eta^2 = 0.020$). Pairwise comparisons following-up on the trend for a group effect indicated that

children exhibited significantly larger micro-offline gains as compared to young ($F_{(1,61)}$ = 4.829, p

= 0.032, partial η^2 = 0.073; see Figure 3D) and older adults ($F_{(1,61)}$ = 5.374, p = 0.024, partial η^2 =

0.081). Similarly, there was a non-significant trend for age being a predictor of micro-offline gains

from childhood into young adulthood (power model: R^2 = 0.029, $F_{(1,96)}$ = 2.873, p = 0.093; see

Figure 3F), but not within older adulthood (power model: R^2 = 0.043, $F_{(1,30)}$ = 1.362, p = 0.252).

499
Figure 3. *Micro-online (**left**) and -offline (**right**) gains. Panels A and B depict micro gains displayed as a*
function of practice blocks. Panels C and D contain violin plots of micro-online and -offline, respectively,
gains averaged across blocks. Note that pairwise comparisons in these panels included blocks or rest
periods in the statistical models.* p < 0.05 for differences between pairs of groups. Panels E and F show
averaged micro-online and -offline gains as a function of age from childhood into young adulthood and
within older adulthood. Lines of fit childhood into young adulthood: micro-online gains = (-0.318*age)
$^{0.459}$; micro-offline gains = 0.298*exp(-0.377*age). Lines of fit within older adulthood: micro-online gains
= (0.003*age) + (-0.238); micro-offline gains = (227.361*age) + (-1.897).
508

4.2 Experiment 2: Motor memory consolidation

4.2.1 Participant characteristics, sleep and vigilance

Group means for participant characteristics, the times of when the experimental sessions were

completed, the duration of the offline periods between sessions, self-reported sleep quality and

duration for the nights preceding each experimental session, and subjective (SSS) and objective

(PVT) measures of vigilance are provided in Table 2. Results from the corresponding statistics

analyses and a brief discussion can be found in Appendix 1 of the Supporting Information (see

Figure S2 and Table S2).

4.2.2 Initial learning dynamics

Similar to Experiment 1, we assessed the dynamics of learning during the initial training session

of Experiment 2. Normalized performance measures are shown in Figure 4. For brevity, the

corresponding statistical results can be found in Appendix 3 of the Supporting Information (Table

S7). In brief, the results from the assessment of the initial training session of Experiment 2 largely

mirrored those presented above for Experiment 1 (i.e., no differences among age groups in the

performance improvements across training blocks).

525
Figure 4. *Average normalized response time (**A**) and accuracy (**B**) for the three experimental sessions and*
two task variants of Experiment 2. Shaded regions represent standard errors of the mean. Corresponding
statistical analyses are in Appendix 3 of the Supporting Information.

4.2.3 Macro-offline performance gains

Results on the sequential macro-offline changes in performance (Figure 5) revealed both offline

period ($F_{(1,104)}$ = 104.365, p < 0.001, partial η^2 = 0.501) and group ($F_{(3,104)}$ = 14.551, p < 0.001,

partial η^2 = 0.296) main effects, as well as a significant offline period x group interaction effect

($F_{(3,104)}$ = 3.367, p = 0.021, partial η^2 = 0.089). Follow-up comparisons revealed significant group

effects for both the 5-hour ($F_{(3,107)}$ = 5.764, p = 0.001, η^2 = 0.143) and the 24-hour ($F_{(3,107)}$ = 20.577,

p < 0.001, η^2 = 0.372) sequential macro-offline gains. Post-hoc pairwise comparisons indicated

that the sequential macro-offline gains after a 5-hour period of wakefulness were higher in children

and adolescents as compared to the young (p = 0.032, G = 0.590; p = 0.053, G = 0.548,

respectively) and older adults (p = 0.008, G = 1.382; p = 0.015, G = 1.290, respectively). The

children also showed higher 24-hour macro-offline gains in comparison to the young (p = 0.001,

G = 1.285) and older (p < 0.001, G = 2.448) adults, and the older adults displayed significantly

lower 24-hour macro-offline gains than all the other age groups (adolescents: $p < 0.001$, $G = 1.256$;

young adults: $p = 0.002$, $G = 1.237$).

In line with the age group comparisons presented above, our analyses with age as a continuous

variable revealed that age significantly predicted the 5h macro-offline gains from childhood into

young adulthood (linear model: $R^2 = 0.068$, $F_{(1,78)} = 5.650$, $p = 0.020$; see Figure 5B). Age was not

a significant predictor of the 5hr offline gains within the older adult group (quadratic model: $R^2 =$

0.117, $F_{(1,24)} = 1.587$, $p = 0.220$). Lastly, age significantly predicted the 24-hour macro-offline

gains from childhood to young adulthood (Power model: $R^2 = 0.157$, $F_{(1,78)} = 14.537$, $p < 0.001$;

see Figure 5D) as well as within older adulthood (linear model: $R^2 = 0.139$, $F_{(1,25)} = 4.047$, $p =$

0.055).

554
Figure 5. *Sequential offline gains across the 5-hour (**Panel A**) and 24-hour (**Panel C**) offline periods for*
*the four age groups. * p < 0.05 and ~ p < 0.10 for pairwise group comparisons. Five-hour (**Panel B**) and*
*24-hour (**Panel D**) sequential offline gains are plotted as a function of age. Lines of fit from childhood into*
*young adulthood: 5h gains = (-0.003 * age) + 0.075; 24h gains = (0.872*age)$^{-0.682}$. Lines of fit within older*
*adulthood: 5h gains = (0.001 * age^2) + (-0.110 * age) + 3.534; 24h gains = (-0.006 * age) + 0.377. For*

better visualization of the age-related changes, the scale of the y-axes was set to range from -0.4 to 0.4 and
thus the young adult with the outlier 5hr sequential gain (visible in Panel A) is not depicted in Panel B.
This individual, however, was included in the fitting procedure. Note that if this individual was excluded
from statistical analyses, the pattern of results observed in Panel A remains largely similar, with the
exception that the pairwise difference between young and older adults becomes statistically significant (i.e.,
older adults exhibit worse offline consolidation over a 5hr wake interval as compared to young adults).

4.2.4 Link between the offline performance changes on a micro- and macro-timescale

The regression model assessing the relationship between the micro- and 5h macro-offline gains –

independent of age group- revealed a significantly positive relationship between these two offline

measures (b = 0.569, p < 0.001; see Figure S5). This suggests that greater performance changes

during the short rest periods throughout training were associated with greater performance gains

across the 5-hour offline period of wakefulness following training. We then assessed whether such

a relationship differed between pairs of age groups. When age group was modeled, young adults

exhibited a steeper slope as compared to children (b = -1.136, p = 0.010), adolescents (b = -1.153,

p = 0.004) and older adults (b = -1.358, p = 0.005). This indicates that the relationship between

micro-offline and 5h macro-offline gains was strongest in young adults as compared to the other

3 age groups. However, note that these between-group differences were no longer statistically

significant if the young adult participant with the extremely negative 5hr offline gain (see Figure

5A) was removed from the analyses.

Micro-offline gains were also significantly and positively related to the 24h macro-offline gains

(b = 0.592, p < 0.001; see Supplemental Figure S6). Specifically, greater performance changes

during the short rest periods were also associated with greater performance gains across the 24-

hour period following training. This relationship was not significantly different between young

adults and any of the other age groups (children: b = -0.222, p =0.581; adolescents: b = 0.245, p =

0.496; older adults: b = -0.048, p = 0.913).

5. Discussion

The current study examined age-related differences among children, adolescents, young adults and

older adults in initial motor sequence learning and motor memory consolidation. Our results

revealed that: (1) the dynamics of initial motor sequence learning, as demonstrated by performance

improvements during training, were comparable (i.e., not statistically different) across age groups;

(2) children exhibited less sequence-specific learning during the initial training session relative to

adolescents and young adults; (3) micro-online and -offline performance changes tended to be the

smallest and largest, respectively, in children; (4) children exhibited greater changes in

performance from the end of initial training to both the 5-hour and 24-hour retests, suggesting a

developmental advantage in the macro-offline consolidation of a newly acquired movement

sequence; (5) there was a significant positive relationship between the magnitudes of micro- and

macro-offline performance changes.

Initial performance improvements are comparable across age groups, but children exhibit less
sequence-specific learning

On the one hand, our results showed comparable performance improvements across training

among age groups, which was consistent with a subset of the previous literature (Du et al., 2017;

Jongbloed-Pereboom et al., 2019; Meulemans et al., 1998; Salehi et al., 2016; Wilhelm et al., 2013;

Bottary et al., 2016; Fitzroy et al., 2021; Fogel et al., 2017; Fogel et al., 2014; Gudberg et al., 2015;

Spencer et al., 2007; Vien et al., 2016). This finding thus agrees with the lifespan invariance model

that has been discussed in previous literature (Tóth-Fáber et al., 2023). On the other hand, our

results indicated that children exhibited less sequence-specific learning, demonstrating that their

performance improvements across sequential training blocks are partially due to general motor

improvements (i.e., improvements on the motor task that are *independent* of sequence learning).

This finding – which resembles the prototypical developmental trajectory where young adults

perform the best and children are progressing towards this optimal performance level - is consistent

with a subset of previous literature in which children exhibited smaller sequence-specific motor

learning (Lukács & Kemény, 2015; Thomas et al., 2004; Thomas & Nelson, 2001). The pattern of

results may explain some of the heterogeneity that is present in the literature with respect to the

development of motor sequence learning. That is, previous research that assessed performance

improvements across blocks of practice may report comparable learning between children and

young adults (e.g., Dorfberger et al., 2007; Wilhelm et al., 2013), whereas studies that afford the

extraction of sequence-specific learning may reveal differences between age groups (e.g., Lukács

& Kemény, 2015; Thomas & Nelson, 2001). Presumably, the greater general learning that is

evident in children is due to this group simply learning how to perform the task itself (i.e., spatially

mapping the stimuli on the screen to the appropriate fingers). Future research that aims to isolate

and examine sequence learning *per se* could consider incorporating an extended task

familiarization phase consisting of additional random runs (i.e., not a sequence to be learned). This

would allow children to exhaust their general learning of the task itself prior to sequence

acquisition. Nevertheless - and as a point of emphasis - even though sequence-specific learning

was smaller in children, they still exhibited significant learning of the motor sequence. That is,

their learning magnitude was still significantly greater than zero (see Table S6) and thus a memory

trace of the acquired sequence was presumably formed. Interestingly, whereas children showed a

smaller learning magnitude, the adolescents were not different from young adults. This suggests

that sequence-specific learning reaches adult-like levels by the teenage years.

Micro-offline consolidation during interspersed rest breaks tended to be larger in children

In line with previous literature (Du et al., 2017) and our hypothesis, children exhibited greater

micro-offline performance gains in comparison to the young adults. However, it is important to

note that the group main effect within the omnibus group (4 levels) x block ANOVA was a non-

significant trend and thus our age effects on the micro-offline gains were weaker in comparison to

Du et al. (2017). A potential explanation for our weaker findings could be the specific ages

included in the child groups. Specifically, Du et al. (2017) found a micro-offline advantage in

younger children (i.e., 6-year-olds) as compared to older children (i.e., 10-year-olds) and adults.

In the current research, the online nature of our protocol made it difficult to acquire data in such

young children (i.e., 6-year-olds). Moreover, as data acquisition in Du et al. (2017) was limited to

these discrete age groups in (i.e., data from 7-9-year-old children were not acquired), an estimate

of when this childhood advantage in micro-offline processing disappears was not possible. We

speculate that the differences between children and young adults in this study would have been

larger in magnitude if the sample included a greater number of younger children (i.e., 6 to 7 year-

olds). Another potential consideration is the considerably longer rest breaks (i.e., 3 min) between

practice blocks that Du et al. (2017) implemented. In comparison to our 15-second rest intervals,

this allowed more time for the recently acquired memory trace to be consolidated over these micro-

offline epochs. Future studies are needed to further examine the micro-learning processes in

children, with systematic examinations into certain methodological choices such as the age of the

participants and the length of the rest epochs.

Although this fast consolidation process occurring over micro-offline intervals has received

considerable attention over the last 5 years (Bönstrup et al., 2019; Bönstrup et al., 2020; Buch et

al., 2021; Gann et al., 2023; Jacobacci et al., 2020), the current study is the first, to the best of our

knowledge, to examine micro-learning processes in older adults. Our results demonstrated

comparable micro-online and -offline performance gains in young and older adults. Accordingly,

whereas older adults have deficits in the macro-offline consolidation processes occurring between

two practice sessions (Fogel et al., 2017; Fogel et al., 2014; Gudberg et al., 2015; Spencer et al.,

2007; Vien et al., 2016; Wilson et al., 2012), they do not exhibit deficits in the rapid consolidation

process that occurs during the rest periods interspersed with blocks of task practice.

Lastly, similar to previous studies in young adults (Bönstrup et al., 2019; Buch et al., 2021; Du et

al., 2017; Jacobacci et al., 2020), our results provide evidence for a significant contribution of the

micro-offline gains to overall learning in all four age groups (see Table S6.C). That is, performance

improvements across blocks of training are largely due to micro-offline – as compared to micro-

online – performance gains. One could argue, however, that the magnitude of the micro-offline

gains could be partially attributed to the build-up of fatigue or reactive inhibition as a function of

task practice within blocks (Gupta & Rickard, 2022, 2023). Specifically, it has been suggested that

performance deteriorates across practice within a block due to reactive inhibition and thus micro-

offline performance gains are inflated due to this performance variable that is independent of

learning and memory processes. However, it is worth noting that this view would conflict with

recent data demonstrating that these micro-offline performance gains are linked to hippocampal

activity during the interspersed rest periods (Buch et al., 2021; Jacobacci et al., 2020). Additional

research is necessary to definitively conclude whether these micro-offline performance gains are

reflective of a memory consolidation process or are simply attributed to reactive inhibition.

Children exhibited a developmental advantage in macro-offline consolidation

Our results revealed larger sequential performance gains across both the 5-hour and 24-hour

macro-offline periods in children. While the young adults showed a performance maintenance (i.e.,

offline gains that were not significantly different from zero; see Appendix 6, Table S7A in

Supporting Information) across the 5-hour interval, children and adolescents significantly

improved their performance on the sequential task. Interestingly, children also demonstrated

higher offline gains across the 24-hour period as compared to adults. While the enhanced 5-hour

consolidation in children is in line with the findings of Ashtamker & Karni (2013) that showed an

accelerated consolidation in children across a one-hour offline period, the greater 24-hour gains

stand in contrast to previous research that demonstrated impaired overnight consolidation in

children [(Fischer et al., 2007; Wilhelm et al., 2008); see below for expanded discussion on sleep-

facilitated consolidation].

It could be argued that the larger macro-offline processing in children could be explained by

performance levels reached at the end of the training session (Kuriyama et al., 2004; Wilhelm,

Metzkow-Mészàros, et al., 2012). For example, perhaps children show greater macro-offline gains

due to a continuation of initial learning rather than the offline processing of previously acquired

information. This potential explanation, however, is not in line with our data. Notably, during the

post-training test phase, there was an absence of a block effect in all groups, including in the

children. This indicates that all groups reached a performance plateau at the end of the first session

and thus macro-offline gains are likely not the result of continued learning. Along the same lines,

one could speculate that the young adults reached a performance ceiling effect, and thus their

potential to further increase their performance over macro-offline intervals was saturated (see plot

of absolute reaction times in Figure S4 of the Supporting Information). Again, this is not consistent

with our data. Notably, there was a significant session effect in sequential offline gains, whereby

gains were larger in the 24-hour retest as compared to the 5-hour retest. Within-group comparisons

revealed that this session effect was present for all age groups, including young adults. This

suggests that a performance ceiling effect was not reached and thus this explanation cannot fully

account for our results. Last, one could raise the question whether perhaps the macro-offline

performance gains are inflated in children due to non-sequence-specific improvements (i.e.,

improvements due to motor task familiarization as opposed to sequence learning). However, the

children did not show greater macro-offline gains on the random task variant (see Appendix 5,

Figure S7 in Supporting Information). Moreover, and to assess this possibility further, we

examined the offline changes in the sequence-specific learning magnitude (i.e., difference between

the learning magnitude (i.e., sequence vs. random) of the 5hr and 24hr retest sessions and the

learning magnitude of the first session; see Appendix 5, Figure S7 in Supporting Information).

These results revealed significant main effects of offline period and group, but no offline period x

group interaction effect. Post-hoc pairwise comparisons indicated that children exhibit enhanced

sequence-specific offline gains – across both offline periods - in comparison to young and older

adults. This finding is consistent with the assessment of the sequential macro-offline gains

presented in the main text (Figure 5). Altogether, these data suggest that the offline changes in

children likely involved the strengthening of their sequential memory and that this strengthening

was greater as compared to young and older adults.

Offline performance gains on the 24-hour retest were largest in children, a result that was

unexpected based on previous literature (Fischer et al., 2007; Wilhelm et al., 2008). Specifically,

children have typically exhibited impaired sleep-facilitated consolidation as observed in young

adults. It is worth noting that most of the previous studies comparing sleep and wake conditions in

children employed an AM/PM design. In this design, participants acquire the motor sequence at a

different time of day based on the condition and then are retested after a delay of 12 hours. For

example, the wake (i.e., AM-PM) group initially learns the sequence in the morning and then is

retested following 12 hours of wakefulness. The sleep (i.e., PM-AM) group completes initial

learning in the evening and this is retested after a 12-hour interval that included a night of sleep.

Although informative, these designs are sensitive to time-of-day effects. And there is evidence that

the time of initial practice can affect performance during acquisition and offline consolidation in

young adults (Kvint et al., 2011; Truong et al., 2022). Specifically, Kvint et al. (2011) found a

greater sequential knowledge acquisition but similar 12-hour consolidation in an AM-trained as

compared to a PM-trained group. Furthermore, Truong et al. (2022) showed an enhanced

consolidation when the training and retest were completed at 3 pm (as compared to 10 am). These

findings indicate time-of-day effects for the acquisition and consolidation outcomes in young

adults. Although previous studies that employed AM/PM designs in children did not find

differences in motor skill acquisition between morning and evening groups (Wilhelm et al., 2013),

the impact on offline consolidation processes has yet to be systematically and exhaustively

examined. Although certainly speculative, it is possible that the impaired overnight consolidation

that previous studies have found in children could be – at least partially – attributed to the choice

of experimental design. An alternative, albeit related explanation, is that the beneficial effect of

sleep – as reported in the current research - requires additional time to emerge at the behavioral

level and thus that these previous studies did not show overnight gains due to their retest

immediately after sleep (i.e., 12 hours after initial learning). Consistent with this explanation,

Desrochers et al. (2016) found that a post-learning nap facilitated the consolidation of a recently

acquired motor sequence in pre-school children, but only after an extended period of time that

included a night of sleep. Additional research is necessary to further disentangle these potential

explanations.

Potential neural correlates of enhanced offline processing in children

An open question is: What are the neural correlates of the greater micro- and macro-offline

performance gains observed in children? Based on (a) recent evidence in young adults

demonstrating that the patterns of activity observed in the hippocampus and putamen during motor

learning are spontaneously reactivated during subsequent micro- and macro-offline intervals

(Buch et al., 2021; King et al., 2022); and, (b) the known role of reactivation processes in memory

consolidation and longer-term retention (Deuker et al., 2013; Tambini & Davachi, 2013), it is

tempting to speculate that the childhood advantages observed in this research can be attributed to

enhanced reactivation of learning-related patterns of brain activity. This possibility is pure

speculation at this point but deserves attention in future research.

Older adults exhibit impaired overnight consolidation

On the other end of the lifespan, our results showed similar 5-hour but smaller 24-hour offline

gains in the older as compared to the young adults. The comparable 5-hour offline gains are in line

with previous studies that showed a similar consolidation between young and older adults across

an offline period of wakefulness (Fogel et al., 2014; Gudberg et al., 2015; Vien et al., 2016).

Similarly, the smaller 24-hour offline gains observed in the older adults are in line with several

studies that demonstrated impaired overnight consolidation with aging (Fogel et al., 2017; Fogel

et al., 2014; Gudberg et al., 2015; Spencer et al., 2007; Vien et al., 2016; Wilson et al., 2012).

These studies have largely attributed this deficit to age-related decreases in sleep quality and

quantity, as well as alterations in the sleep-related electrophysiological markers of plasticity such

as sleep spindles and slow waves (Bottary et al., 2016; Fogel et al., 2017; Fogel et al., 2014; but

see Fitzroy et al., 2021; Gudberg et al., 2015; and King et al., 2017 for discussion of alternative

explanations). Whereas our older adult participants did report shorter sleep duration as compared

to both children and adolescents, no differences in sleep quality were found. However, it is

important to note that these measures were self-reported and previous research has shown that

older adults tend to overestimate their sleep quality (Vitiello et al., 2004). Nonetheless, our results

replicate the often demonstrated finding that aging negatively impacts overnight consolidation of

recently acquired motor sequences.

Considerations and future recommendations

There are some limitations and methodological considerations of the current research that warrant

further attention. *First*, these experiments were conducted online and thus were not monitored by

researchers in real-time. Although we cannot completely discard the possibility that the online

nature influenced our results, it is worth nothing that previous studies that compared online and

in-person experiments in children (Chuey et al., 2022) and young adults (Bönstrup et al., 2020;

Cubillos et al., 2023) found comparable results between these study modalities. For instance,

Cubillos et al. (2023) showed similar learning curves for the performance of the finger-tapping

task between a supervised-lab and unsupervised-online young adult group. Additionally, Bönstrup

et al. (2020) obtained similar micro-offline learning in young adults through an online

crowdsourcing platform. It is also worth noting that our protocol was not implemented in common

crowdsourcing platforms (e.g., Mechanical Turk) that typically acquire full datasets within hours

of posting an experiment and have come under scrutiny for unreliable data (Webb & Tangney,

2022). Furthermore, although there are drawbacks to employing web-based data acquisition, a

significant advantage is the recruitment from a large and broad pool of participants, enabling us to

include individuals that otherwise would not be able to complete our traditional in-person studies.

Second, and specific to experiment 2, each session ended with a pseudorandom run. The

implementation of this random run is often used in sequence learning studies and has the benefit

of assessing sequence specificity of performance improvements (Du et al., 2017; Lukács &

Kemény, 2015; Salehi et al., 2016; Savion-Lemieux et al., 2009; Thomas & Nelson, 2001).

However, we cannot rule out the possibility that the random runs at the end of each session

interfered with the macro-offline processing of the sequential motor memory.

Third, it is important to note that, due to concerns surrounding the acquisition of quality data from

young children, our minimum age for this research was 7 years. Moreover, this youngest age was

relatively under-sampled in our experiments (n = 3 and 1 7-year-olds included in analyses of

Experiments 1 and 2, respectively). Due to difficulties with completing the online motor task,

several 7-year-old children had to be excluded from data analyses. Future studies could consider

implementing certain measures (e.g., monitoring through videocall) to increase the number of

young children that successfully complete the experiments.

Fourth, and as outlined in our pre-registration, the age groups were operationally defined based on

age (i.e., children and adolescents were between 7 and 12, and 13 and 17 years, respectively),

which does not consider the participant's pubertal status. Our questionnaire battery did ask the

parents to indicate whether their child exhibited signs of puberty onset at the time of participation.

Note that all participants 13 years and older (i.e., classified as an adolescent based on age) were

reported to having started puberty. And only a minimal number of child subjects (6 in each of the

experiments) were considered to have initiated puberty. Given the relatively few subjects that

would be re-classified based on parental report of puberty onset, we do not anticipate our results

to substantially change if pubertal status – as opposed to age - was used to demarcate childhood

and adolescence.

6. Conclusion

The current study examined initial motor sequence learning and the time course of motor memory

consolidation across the human lifespan. Our results demonstrated comparable dynamics of

performance improvements during the initial learning session, yet 7- to 12-year-old children

exhibited smaller sequence-specific learning. Interestingly, children demonstrated the largest

performance gains over micro- and macro-offline intervals. This indicates that children exhibit a

developmental advantage in the offline processing of recently practiced movement sequences. The

neural underpinnings of this childhood advantage are not yet known and thus future research is

warranted to characterize them.

REFERENCES

Adi-Japha, E., Badir, R., Dorfberger, S., & Karni, A. (2014). A matter of time: Rapid motor memory stabilization in childhood. *Developmental Science*, *17*(3), 424–433. https://doi.org/10.1111/desc.12132

Ashtamker, L., & Karni, A. (2013). Motor memory in childhood: Early expression of consolidation phase gains. *Neurobiology of Learning and Memory*, *106*, 26–30. https://doi.org/10.1016/j.nlm.2013.07.003

Basner, M., Mollicone, D., & Dinges, D. F. (2011). Validity and sensitivity of a brief psychomotor vigilance test (PVT-B) to total and partial sleep deprivation. *Acta Astronautica*, *69*(11–12), 949–959. https://doi.org/10.1016/j.actaastro.2011.07.015

Bönstrup, M., Iturrate, I., Hebart, M. N., Censor, N., & Cohen, L. G. (2020). Mechanisms of offline motor learning at a microscale of seconds in large-scale crowdsourced data. *Npj Science of Learning*, *5*(1). https://doi.org/10.1038/s41539-020-0066-9

Bönstrup, M., Iturrate, I., Thompson, R., Cruciani, G., Censor, N., & Cohen, L. G. (2019). A Rapid Form of Offline Consolidation in Skill Learning. *Current Biology*, *29*(8), 1346-1351.e4. https://doi.org/10.1016/j.cub.2019.02.049

Bottary, R., Sonni, A., Wright, D., & Spencer, R. M. C. (2016). Insufficient chunk concatenation may underlie changes in sleep-dependent consolidation of motor sequence learning in older adults. *Learning and Memory*, *23*(9), 455–459. https://doi.org/10.1101/lm.043042.116

Buch, E. R., Claudino, L., Quentin, R., Bönstrup, M., & Cohen, L. G. (2021). Consolidation of human skill linked to waking hippocampo-neocortical replay. *Cell Reports*, *35*(10). https://doi.org/10.1016/j.celrep.2021.109193

Chuey, A., Boyce, V., Cao, A., & Frank, M. C. (2022). *Conducting developmental research online vs. in-person: A meta-analysis*. https://doi.org/10.31234/osf.io/qc6fw

Cubillos, L. H., Augenstein, T. E., Ranganathan, R., & Krishnan, C. (2023). Breaking the barriers to designing online experiments: A novel open-source platform for supporting procedural skill learning experiments. *Computers in Biology and Medicine*, *154*. https://doi.org/10.1016/j.compbiomed.2023.106627

Desrochers, P. C., Kurdziel, L. B. F., & Spencer, R. M. C. (2016). Delayed benefit of naps on motor learning in preschool children. *Experimental Brain Research*, *234*(3), 763–772. https://doi.org/10.1007/s00221-015-4506-3

Deuker, L., Olligs, J., Fell, J., Kranz, T. A., Mormann, F., Montag, C., Reuter, M., Elger, C. E., & Axmacher, N. (2013). Memory Consolidation by Replay of Stimulus-Specific Neural Activity. *The Journal of Neuroscience*, *33*(49), 19373–19383. https://doi.org/10.1523/JNEUROSCI.0414-13.2013

Dinges, D. F., & Powell, J. W. (1985). Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. In *Behavior Research Methods, Instruments, & Computers* (Vol. 17, Issue 6).

Dorfberger, S., Adi-Japha, E., & Karni, A. (2007). Reduced susceptibility to interference in the consolidation of motor memory before adolescence. *PLoS ONE*, *2*(2), 1–6. https://doi.org/10.1371/journal.pone.0000240

Du, Y., Valentini, N. C., Kim, M. J., Whitall, J., & Clark, J. E. (2017). Children and adults both learn motor sequences quickly, but do so differently. *Frontiers in Psychology*, *8*(FEB), 1–10. https://doi.org/10.3389/fpsyg.2017.00158

Epskamp, S., & Nuijten, M. B. (2018). statcheck: Extract statistics from articles and
recompute p-values. R package version 1.3.0. Web implementation at
https://statcheck.io.
Erdfelder, E., Faul, F., Lang, A.-G., & Buchner, A. (2007). G * Power 3 : A flexible statistical
power analysis program for the social , behavioral , and biomedical sciences. *Behavior
Research Methods*, *39*(2), 175–191.
Fischer, S., Wilhelm, I., & Born, J. (2007). Developmental differences in sleep's role for
implicit off-line learning: Comparing children with adults. *Journal of Cognitive
Neuroscience*, *19*(2), 214–227. https://doi.org/10.1162/jocn.2007.19.2.214
Fitzroy, A. B., Kainec, K. A., Seo, J., & Spencer, R. M. C. (2021). Encoding and consolidation
of motor sequence learning in young and older adults. *Neurobiology of Learning and
Memory*, *185*(July), 107508. https://doi.org/10.1016/j.nlm.2021.107508
Fogel, S. M., Albouy, G., Vien, C., Popovicci, R., King, B. R., Hoge, R., Jbabdi, S., Benali,
H., Karni, A., Maquet, P., Carrier, J., & Doyon, J. (2014). fMRI and sleep correlates of
the age-related impairment in motor memory consolidation. *Human Brain Mapping*,
35(8), 3625–3645. https://doi.org/10.1002/hbm.22426
Fogel, S., Vien, C., Karni, A., Benali, H., Carrier, J., & Doyon, J. (2017). Sleep spindles: a
physiological marker of age-related changes in gray matter in brain regions supporting
motor skill memory consolidation. *Neurobiology of Aging*, *49*, 154–164.
https://doi.org/10.1016/j.neurobiolaging.2016.10.009
Gann, M. A., Dolfen, N., King, B. R., Robertson, E. M., & Albouy, G. (2023). Prefrontal
stimulation as a tool to disrupt hippocampal and striatal reactivations underlying fast
motor memory consolidation. *Brain Stimulation*.
https://doi.org/10.1016/j.brs.2023.08.022
Gualtieri, S., & Finn, A. S. (2022). The Sweet Spot: When Children's Developing Abilities,
Brains, and Knowledge Make Them Better Learners Than Adults. *Perspectives on
Psychological Science*, *17*(5), 1322–1338. https://doi.org/10.1177/17456916211045971
Gudberg, C., Wulff, K., & Johansen-Berg, H. (2015). Sleep-dependent motor memory
consolidation in older adults depends on task demands. *Neurobiology of Aging*, *36*(3),
1409–1416. https://doi.org/10.1016/j.neurobiolaging.2014.12.014
Gupta, M. W., & Rickard, T. C. (2022). Dissipation of reactive inhibition is sufficient to
explain post-rest improvements in motor sequence learning. *Npj Science of Learning*,
7(1). https://doi.org/10.1038/s41539-022-00140-z
Gupta, M. W., & Rickard, T. C. (2023). A Quantitative Model that Incorporates Reactive
Inhibition but No Facilitating Consolidation Explains Single-Session Motor Sequence
Practice Effects. *PsyArXiv*. https://doi.org/10.31234/osf.io/d5pv9
Jacobacci, F., Armony, J. L., Yeffal, A., Lerner, G., Amaro, E., Jovicich, J., Doyone, J., &
Della-Maggiore, V. (2020). Rapid hippocampal plasticity supports motor sequence
learning. *Proceedings of the National Academy of Sciences of the United States of
America*, *117*(38), 23898–23903. https://doi.org/10.1073/pnas.2009576117
Janacsek, K., Fiser, J., & Nemeth, D. (2012). The best time to acquire new skills : age-related
differences in implicit sequence learning across the human lifespan. *Devel*, *4*, 496–505.
https://doi.org/10.1111/j.1467-7687.2012.01150.x
Jongbloed-Pereboom, M., Nijhuis-van der Sanden, M. W. G., & Steenbergen, B. (2019).
Explicit and implicit motor sequence learning in children and adults; the role of age and

visual   working   memory.   *Human   Movement   Science*,   *64*,   1–11.
https://doi.org/10.1016/j.humov.2018.12.007

Juhasz, D., Nemeth, D., & Janacsek, K. (2019). Is there more room to improve? The lifespan
trajectory of procedural learning and its relationship to the between? The within-group
differences   in   average   response   times.   *PLoS   ONE*,   *14*(7),   1–20.
https://doi.org/10.1371/journal.pone.0215116

Kavos Salehi, S., Sheikh, M., Hemayattalab, R., & Humaneyan, D. (2016). The Effect of
Different Ages levels and explicit-implicit Knowledge on Motor Sequence Learning.
International Journal of Environmental & Science Education, *11*(18), 13157–13165.

King, B. R., Dolfen, N., Gann, M. A., Renard, Z., Swinnen, S. P., & Albouy, G. (2019).
Schema and Motor-Memory Consolidation. https://doi.org/10.1177/0956797619847164

King, B. R., Fogel, S. M., Albouy, G., & Doyon, J. (2013). Neural correlates of the age-
related changes in motor sequence learning and motor adaptation in older adults.
Frontiers   in   Human   Neuroscience,   *APR   2013*.
https://doi.org/10.3389/fnhum.2013.00142

King, B. R., Gann, M. A., Mantini, D., Doyon, J., & Albouy, G. (2022). Persistence of
hippocampal and striatal multivoxel patterns during awake rest after motor sequence
learning. *IScience*, *25*(12). https://doi.org/10.1016/j.isci.2022.105498

King, B. R., Hoedlmoser, K., Hirschauer, F., Dolfen, N., & Albouy, G. (2017). Sleeping on
the motor engram: The multifaceted nature of sleep-related motor memory
consolidation. In *Neuroscience and Biobehavioral Reviews* (Vol. 80, pp. 1–22). Elsevier
Ltd. https://doi.org/10.1016/j.neubiorev.2017.04.026

Kuriyama, K., Stickgold, R., & Walker, M. P. (2004). Sleep-dependent learning and motor-
skill   complexity.   *Learning   and   Memory*,   *11*(6),   705–713.
https://doi.org/10.1101/lm.76304

Kvint, S., Bassiri-Tehrani, B., Pruski, A., Nia, J., Nemet, I., Lopresti, M., Moisello, C.,
Perfetti, B., Tononi, G., Ghilardi, M. F., & Ghilardi, M. F. (2011). Acquisition and
retention of motor sequences: the effects of time of the day and sleep. In *Arch Ital Biol*
(Vol. 149, Issue 3).

Loh, S., Lamond, N., Dorrian, J., Roach, G., & Dawson, D. (2004). The validity of
psychomotor vigilance tasks of less than 10-minute duration. *Behavior Research
Methods, Instruments, & Computers*, *36*(2), 339–346.

Lukács, Á., & Kemény, F. (2015). Development of different forms of skill learning
throughout   the   lifespan.   *Cognitive   Science*,   *39*(2),   383–404.
https://doi.org/10.1111/cogs.12143

Maclean, A. W., Fekken, G. C., Saskin, P., & Knowles, J. B. (1992). Psychometric evaluation
of the Stanford Sleepiness Scale. In *Journal of Sleep Research*.
https://doi.org/10.1111/j.1365-2869.1992.tb00006.x

Meulemans, T., Van Der Linden, M., & Perruchet, P. (1998). Implicit Sequence Learning in
Children. *Journal of Experimental Child Psychology*, *69*(3), 199–221.
https://doi.org/10.1006/jecp.1998.2442

Pan, S. C., & Rickard, T. C. (2015). *Sleep and Motor Learning: Is There Room for
Consolidation? 141*(4), 812–834.

Reverberi, S., Dolfen, N., Van Roy, A., Albouy, G., & King, B. R. (2023). Sleep does not
influence schema-facilitated motor memory consolidation. *PLOS ONE*, *18*(1),
e0280591. https://doi.org/10.1371/journal.pone.0280591

Salehi, S. K., Sheikh, M., Hemayattalab, R., & Humaneyan, D. (2016). The Effect of Different Ages levels and explicit-implicit Knowledge on Motor Sequence Learning. In *INTERNATIONAL JOURNAL OF ENVIRONMENTAL & SCIENCE EDUCATION* (Vol. 11, Issue 18).

Savion-Lemieux, T., Bailey, J. A., & Penhune, V. B. (2009). Developmental contributions to motor sequence learning. *Experimental Brain Research*, *195*(2), 293–306. https://doi.org/10.1007/s00221-009-1786-5

Spencer, R. M. C., Gouw, A. M., & Ivry, R. B. (2007). Age-related decline of sleep-dependent consolidation. *Learning and Memory*, *14*(7), 480–484. https://doi.org/10.1101/lm.569407

Stoet, G. (2010). *PsyToolkit : A software package for programming psychological experiments using Linux*. *42*(4), 1096–1104. https://doi.org/10.3758/BRM.42.4.1096

Stoet, G. (2017). *PsyToolkit : A Novel Web-Based Method for Running Online Questionnaires and Reaction-Time Experiments*. *44*(1), 24–31. https://doi.org/10.1177/0098628316677643

Sugawara, S. K., Tanaka, S., Tanaka, D., Seki, A., Uchiyama, H. T., Okazaki, S., Koeda, T., & Sadato, N. (2014). Sleep is associated with offline improvement of motor sequence skill in children. *PLoS ONE*, *9*(11), 7–12. https://doi.org/10.1371/journal.pone.0111635

Tambini, A., & Davachi, L. (2013). Persistence of hippocampal multivoxel patterns into postencoding rest is related to memory. *Proceedings of the National Academy of Sciences*, *110*(48), 19591–19596. https://doi.org/10.1073/pnas.1308499110

Thomas, K. M., Hunt, R. H., Vizueta, N., Sommer, T., Durston, S., Yang, Y., & Worden, M. S. (2004). Evidence of developmental differences in implicit sequence learning: An fMRI study of children and adults. *Journal of Cognitive Neuroscience*, *16*(8), 1339–1351. https://doi.org/10.1162/0898929042304688

Thomas, K. M., & Nelson, C. A. (2001). Serial Reaction Time Learning in Preschool- and School-Age Children. *Journal of Experimental Child Psychology*, *79*(4), 364–387. https://doi.org/10.1006/jecp.2000.2613

Tóth-Fáber, E., Nemeth, D., & Janacsek, K. (2023). Lifespan developmental invariance in memory consolidation: evidence from procedural memory. *PNAS Nexus*, *2*(3). https://doi.org/10.1093/pnasnexus/pgad037

Truong, C., Hilt, P. M., Bouguila, F., Bove, M., Lebon, F., Papaxanthis, C., & Ruffino, C. (2022). Time-of-day effects on skill acquisition and consolidation after physical and mental practices. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-09749-x

Vien, C., Boré, A., Lungu, O., Benali, H., Carrier, J., Fogel, S., & Doyon, J. (2016). Age-related white-matter correlates of motor sequence learning and consolidation. *Neurobiology of Aging*, *48*, 13–22. https://doi.org/10.1016/j.neurobiolaging.2016.08.006

Vitiello, M. V, Larsen, L. H., & Moe, K. E. (2004). Age-related sleep change. *Journal of Psychosomatic Research*, *56*(5), 503–510. https://doi.org/10.1016/S0022-3999(04)00023-6

Webb, M. A., & Tangney, J. P. (2022). Too good to be true: Bots and Bad data from Mechanical Turk. *Perspectives on Psychological Science*, 1–4. https://doi.org/https://doi.org/10.1177/17456916221120027

Wilhelm, I., Diekelmann, S., & Born, J. (2008). Sleep in children improves memory
performance on declarative but not procedural tasks. *Learning and Memory*, *15*(5), 373–
377. https://doi.org/10.1101/lm.803708
Wilhelm, I., Metzkow-Mészàros, M., Knapp, S., & Born, J. (2012). Sleep-dependent
consolidation of procedural motor memories in children and adults: The pre-sleep level
of   performance   matters.   *Developmental   Science*,   *15*(4),   506–515.
https://doi.org/10.1111/j.1467-7687.2012.01146.x
Wilhelm, I., Rose, M., Imhof, K. I., Rasch, B., Büchel, C., & Born, J. (2013). The sleeping
child outplays the adult's capacity to convert implicit into explicit knowledge. *Nature
Neuroscience*, *16*(4), 391–393. https://doi.org/10.1038/nn.3343
Wilson, J. K., Baran, B., Pace-Schott, E. F., Ivry, R. B., & Spencer, R. M. C. (2012a). Sleep
modulates word-pair learning but not motor sequence learning in healthy older adults.
Neurobiology   of   Aging,   *33*(5),   991–1000.
https://doi.org/10.1016/j.neurobiolaging.2011.06.029.Sleep
Zinke, K., Wilhelm, I., Bayramoglu, M., Klein, S., & Born, J. (2017). Children's initial sleep-
associated changes in motor skill are unrelated to long-term skill levels. *Developmental
Science*, *20*(6). https://doi.org/10.1111/desc.12463
1032