1

Clonal stochasticity in early NK cell response to mouse cytomegalovirus is generated by

mature subsets of varying proliferative ability

Darren Wethington[1,2,*], Saeed Ahmad[1], Marc Potempa[6], Giuseppe Giuliani[1,5], Oscar A. Aguilar[6],

Maheshwor Poudel[1], Simon Grassmann[7], William Stewart[8], Nicholas M. Adams[7,9], Joseph C.

Sun[7], Lewis L. Lanier[6,*], Jayajit Das[1,2,3,4,*]

[1]Steve and Cindy Rasmussen Institute for Genomic Medicine, Nationwide Children's Hospital,

Columbus OH

[2]Biomedical Sciences Graduate Program, [3]Department of Pediatrics, [4]Pelotonia Institute for

Immuno-Oncology, College of Medicine, The Ohio State University, Columbus OH

[5]Department of Physics, The Ohio State University, Columbus OH

[6]Department of Microbiology and Immunology and the Parker Institute for Cancer

Immunotherapy, University of California, San Francisco CA

[7]Immunology Program, Memorial Sloan Kettering Cancer Center, New York, NY

[8]GIG Statistical Consulting, LLC, Columbus OH

[9]Department of Pathology, New York University Grossman School of Medicine, New York, NY

*Corresponding Authors, wethington.8@osu.edu, Lewis.Lanier@ucsf.edu, jayajit@gmail.com

Abstract

Natural killer (NK) cells are classically defined as innate immune cells, but experiments show

that mouse cytomegalovirus (MCMV) infection in C57BL/6 mice can cause NK cells to undergo

antigen-specific proliferation and memory formation, similar to adaptive CD8+ T cells. One

shared behavior between CD8+ T cells and NK cells is clonal expansion, where a single

stimulated cell proliferates rapidly to form a diverse population of cells. For example, clones

derived from single cells are most abundant during expansion when they are primarily CD27- for

NK cells and CD62L- for T cells, phenotypes derived from precursor CD27+ and CD62L+ cells,

respectively. Here we determined the mechanistic rules involving proliferation, cell death, and

differentiation of endogenous and adoptively transferred NK cells in the expansion phase of the

response to MCMV infection. We found that the interplay between cell proliferation and cell

death of mature CD27- NK cells and a highly proliferative CD27-Ly6C- mature subtype and

intrinsic stochastic fluctuations in these processes play key roles in regulating the heterogeneity

and population of the NK cell subtypes. Furthermore, we estimate rates for maturation of

endogenous NK cells in homeostasis and in MCMV infection and found that only NK cell

growth rates, and not differentiation rates, are appreciably increased by MCMV. Taken together,

these results quantify the differences between the kinetics of NK cell antigen-specific expansion

from that of CD8+ T cells and unique mechanisms that give rise to the observed heterogeneity in

NK cell clones generated from single NK cells in the expansion phase.

Introduction

Natural Killer (NK) cells are lymphocytes of the innate immune system and provide important

protection against viral infections and tumors (Caligiuri, 2008; Vivier *et al*, 2008). NK cells

develop in the bone marrow and express a diverse set of germline-encoded activating and

inhibitory receptors (Abel *et al*, 2018; Biassoni, 2008; Huntington *et al*, 2007). Mouse

cytomegalovirus (MCMV) infection induces antigen-specific expansion of naïve Ly49H+ NK

cells in mice, followed by a contraction phase, and then the formation of long-lived adaptive NK

cells displaying recall responses to the same antigen (Cerwenka & Lanier, 2016; Robbins *et al*,

2004; Schlub *et al*, 2011; Sun *et al*, 2009). The activating Ly49H receptor, encoded by *Klra8*,

recognizes the MCMV-encoded m157 glycoprotein and leads to activation of Ly49H+ NK cells

(Orr *et al*, 2009; Voigt *et al*, 2003). In humans, a similar activation of NK cells by human

cytomegalovirus is observed with the preferential expansion of NKG2C+ NK cells (Guma *et al*,

2004; Hendricks *et al*, 2014; Ishiyama *et al*, 2022; Kim *et al*, 2019; Lopez-Verges *et al*, 2011).

The expansion, contraction, and memory formation of Ly49H+ NK cells during MCMV

infection bears similarity to antigen-specific responses of adaptive CD8+ T cells that generate

memory CD8+ T cells (Adams *et al*, 2020; Buchholz *et al*, 2013; Kaech *et al*, 2002; Schlub *et*

al., 2011).

A range of experiments investigated epigenetic changes, RNA transcription, and surface

protein expression in single and bulk Ly49H+ NK cells undergoing expansion and contraction in

mice responding to MCMV infection (Flommersfeld *et al*, 2021; Grassmann *et al*, 2019; Lau *et*

al, 2018; Martinet *et al*, 2015; Min-Oo *et al*, 2014; Nabekura *et al*, 2014; Potempa *et al*, 2022;

Riggan *et al*, 2022a). These observations show that the NK cell compartment undergoes

phenotypic changes, resulting in changes in the abundances of varying subpopulations of cells.

The development of these subpopulations depends on the three main signal inputs, antigen

simulation of a specific receptor (e.g., Ly49H), co-stimulation, and cytokine signaling (Ni *et al*,

2012; Peng & Tian, 2017; Romee *et al*, 2016). These subpopulations express distinct epigenetic,

transcriptomic, and protein markers imprinted by signaling and gene regulatory processes.

Observations of clonal expansion of barcoded single Ly49H+ NK cells adoptively transferred

into $Rag2^{-/-}Il2rg^{-/-}$ mice revealed another layer of heterogeneity where sizes and compositions

of NK cell clones originating from single naïve Ly49H+ NK cells vary widely (Flommersfeld *et*

al., 2021; Grassmann *et al.*, 2019). This heterogeneity in the NK cell population is regulated by

the stochastic kinetics of cell proliferation, cell death, and differentiation involving the

subpopulations that emerge as the NK cells respond to the infection. Analysis of similar

heterogeneities in clonal CD8+ T cell populations (Buchholz *et al.*, 2013; Gerlach *et al*, 2013)

have shown generation of a highly proliferative short-lived effector phenotype and intrinsic

stochastic fluctuations in the differentiation and proliferation processes, which account for

heterogeneity of the CD8+ T cell clones (Buchholz *et al.*, 2013). These effector CD8+ T cells are

CD62L- and differentiate during the expansion phase from upstream memory precursors that are

CD62L+(Schlub *et al*, 2010; Yang *et al*, 2011). More recently, tracking proliferation, death, and

differentiation of CD8+ T cells during clonal expansion has introduced the notion of division

destiny where antigen and cytokine stimulation in an individual T cell determine the division

destiny or the number of divisions the T cell undergoes before differentiating into a state with

negligible proliferation(Cheon *et al*, 2021; De Boer & Yates, 2023; Hawkins *et al*, 2007; Heinzel

et al, 2017). More generally, these studies demonstrated that the fates of individual T or B cells

during clonal expansion are linked to the number of divisions executed by those individual

cells(Bresser *et al*, 2022; De Boer & Yates, 2023). Modeling the expansion phase of CD8+ T cell

clones incorporating division destiny, heterogeneous antigen stimulation times, inheritance of

division and death times in daughter cells from mother T cell, and cell division dependent

kinetics of cell lineage marker (e.g., CD62L) expressions Pandit and DeBoer (Pandit & De Boer,

2019) showed that the model can qualitatively recapitulate the observed heterogeneity (Buchholz

et al., 2013)) in CD8+ T cell population in the expansion phase. The above models known as

Cyton models investigate stochastic kinetics of individual T/B cells during clonal expansion and

contraction. The existence of division destiny in antigen stimulated Ly49H+ NK cells still needs

to be elucidated (Rückert & Romagnani, 2024).

Despite the qualitative similarities between the antigen-specific responses of NK cells

and CD8+ T cells there are several marked differences. Single-cell NK clones typically undergo

a relatively smaller 1000× increase in number compared to the 40,000× increase in CD8+ T cell

populations (Adams *et al.*, 2020). Additionally, mature CD27- NK cells grow more slowly than

their immature CD27+ counterparts (Buchholz *et al.*, 2013; Hayakawa & Smyth, 2006; Paul *et*

al, 2016), whereas effector CD62L- CD8+ T cells proliferate faster than precursor CD62L+

subsets (Adams *et al.*, 2020; Kretschmer *et al*, 2020). These data suggest there may be

mechanistic differences in antigen-specific NK and CD8+ T cell responses.

Here, we addressed whether distinct mechanisms underlie the expansion of NK cells and

CD8+ T cells during antigen-specific responses by performing stochastic population dynamic

modeling on published bulk and single-cell NK cell data collected during the expansion phase

following MCMV infection in C57BL/6 mice. Our analysis and modeling show that the

mechanistic rules that describe the expansion of CD8+ T cells are inadequate for describing the

development of MCMV-specific NK cells. We found that mature CD27-Ly6C- NK cells

undergo a greater rate of cell proliferation compared to immature CD27+ NK cells and more

mature CD27-Ly6C+ NK cells, and mature CD27- NK cells undergo increased death relative to

the CD27+ immature cells. The variations in the timescales of proliferation and death processes

coupled with the intrinsic stochastic fluctuations within these processes among the NK cell

subsets regulate the heterogeneity in the sizes of the NK cell clones originated from individual

immature NK cells during MCMV infection. Quantification of the population kinetics of

Ly49H+ NK cells in MCMV infection in the expansion phase shows a 10-fold increase in the

cell growth rate compared to that in homeostatic conditions, whereas the effect of the infection

on the maturation rate appears to be negligible. We also find that growth rates of adoptive NK

cells in transfer experiments during MCMV infection in the expansion phase are higher than

those of endogenous NK cells.

Results

A minimal progressive two-state immature to mature differentiation model cannot capture

the heterogeneity of the NK cell clones in the expansion phase

We began our investigations by studying whether a minimal two-state progressive differentiation

model (Fig. 1a) could describe the size and composition of NK cell clones that originated from

single immature Ly49H+ CD27+ NK cells at day 8 post-infection with MCMV, a time typically

at the peak of expansion. The differentiation of immature CD27+ to mature CD27- NK cells

during MCMV infection and in homeostasis has been widely documented and established in

previous experiments (Chiossone *et al*, 2009; Grassmann *et al.*, 2019; Min-Oo *et al.*, 2014;

Mitrovic *et al*, 2012). We used data collected by Flommersfeld *et al.(Flommersfeld et al., 2021)*,

in which they adoptively transferred single genetic marker barcoded Ly49H+ NK cells to a host

that is subsequently infected with MCMV. Splenocytes are then harvested, and resulting NK cell

clones are distinguished with flow cytometry, and CD27+ and CD27- cells are determined with

flow cytometry gates. We characterized the NK cell clones by the means and variances of the

populations of CD27+ or CD27- NK cells and the co-variance between the two cell types in the

clones. The NK cell clones reported by Flommersfeld *et al.* contained mixtures of CD27+ and

CD27- NK cells in the spleen. We evaluated the percentage of CD27+ NK cells in each clone

and computed the correlation ($C_{size\text{-}CD27+}$) of the size of the clone with the percentage of CD27+

NK cells in the clones. The larger size NK cell clones contained a greater proportion of the

CD27- NK cells and thus produced a negative correlation between the clone size and the

percentage of CD27+ NK cells in individual clones (Fig. 1b). This finding is in line with that

found in a similar experiment by Grassmann *et al. (Grassmann et al., 2019).* These authors also

transferred congenically-marked CD27- and CD27+ Ly49H+ NK cells in 1:1 ratio at day 0 and

then assayed the expanded CD27+/- NK cells at day 8 post infection, showing a higher

proportion of mature CD27- NK cells compared to the immature CD27+ NK cells (Figure S2E in

Grassmann *et al. (Grassmann et al., 2019)*). This result is supported by other experiments

(Hayakawa & Smyth, 2006; Paul *et al.*, 2016), and we used this observation to constrain our

model parameter space. Heterogeneity of clones of CD8+ T cells generated by antigenic

SIINFEKL stimulation showed a similar pattern where the largest clones in the expansion phase

(day 7 post-stimulation) are composed of higher fractions of the mature CD62L- CD8+ T cells

(Buchholz *et al.*, 2013). Because mature CD62L- CD8+ T cells proliferate faster than the

immature CD62L+ CD8+ T cells (Buchholz *et al.*, 2013; Kretschmer *et al.*, 2020), it is intuitive

that a larger size of a clone can be attributed to the larger proportion of the mature CD8+ T cells

in the clone (Fig. 1c). In contrast, NK cells proliferate slower when they differentiate to the

mature CD27- phenotype. Thus, it is unclear how the similar relationship between the clone size

and the composition of the clones can be understood. In order to probe this question further we

used a stochastic model (Fig. 1a) that describes the proliferation and differentiation processes in

single Ly49H+ NK cells in response to MCMV infection. We do not model the MCMV infection

explicitly and assume that the parameters of the model capture the effects of antigen, co-receptor,

and cytokine stimulation and signaling and epigenetic modifications on the rates of

differentiation and the proliferation of the cell types. In this model, immature CD27+ NK cells

differentiate with a rate r to mature CD27- NK cells, and each cell type proliferates with rates, k_I

and k_M, respectively. The proliferation and differentiation processes are modeled as stochastic

events occurring with the above rates. We start our simulations with a single CD27+ NK cell and

evaluated the number of CD27+ and CD27- NK cells at 8 days, which represent the size and the

composition of a single NK cell clone. Repetitions of the stochastic simulations with the same

initial configuration and rates will produce clones of different sizes and compositions due to

intrinsic noise fluctuations in the stochastic simulations. We computed the correlation (C_{size-}

$_{CD27+}$) between the clone sizes and the percentages of the CD27+ NK cells in each clone from our

simulations. Our parameter scan of randomly sampled values of k_I, k_M, and r revealed that C_{size-}

$_{CD27+}$ is negative *only* when $k_I < k_M$ (Fig. 1d). However, the observations that CD27+ cells should

grow faster than CD27- cells imply $k_I > k_M$ for the NK cells. Therefore, the two-state model

cannot capture the heterogeneity of the NK cell clones.

Next, we investigated if inclusion of death of NK cells can make the model consistent

with the negative correlation between the clone size and the percentage of CD27+ NK cells in

each clone. We updated our proposed model to include both proliferation and death of each cell

population (Fig. 1e). In such a model, each cell has a probability of proliferating according to

birth rate b and probability of dying indicated by the death rate d. Cells can go through many

proliferation events before undergoing death, which would indicate a positive growth rate. The

net growth of the CD27+ and CD27- NK cell populations are determined by $k_I=b_I-d_I$ and $k_M=b_M-$

d_M, respectively, where b_I and d_I (or b_M and d_M) are the proliferation and death rates of the

immature CD27+ (or mature CD27-) NK cells. Repeating a parameter scan by randomly

sampling the 5 parameters (b_I, d_I, b_M, d_M, and r) from uniform distributions and evaluating the

correlation $C_{size\text{-}CD27+}$ shows that some parameter configurations can generate the negative values

at the parameter values, $k_I > k_M$ (Fig. 1f). Further inspection shows that parameter values in the

regime $k_I > k_M$ that produce the negative correlation share a common trait: mature CD27- cells

proliferate *and* die faster than immature CD27+ cells. To gain further insight into this behavior

we investigated the dependence of $C_{size\text{-}CD27+}$ on the parameter values and stochastic fluctuations

analytically and found that larger proliferation and death rates of the CD27- NK cells produce

large stochastic fluctuations in the sizes of the CD27- cell populations. Thus, mature CD27- NK

cells in a clone could undergo many proliferation events before encountering death, creating

larger clones with higher proportion of mature CD27- NK cells. Conversely, the CD27- cells

could go through many death events before a proliferation event causing the clone size to shrink

(details in Supplementary Text 1).

We investigated whether this model could accurately fit the first (means) and second

(variances and covariance) moments of the measured CD27+ and CD27- NK cell populations at

8 days post-infection. We fit the model by minimizing the sum of square errors (SSE) subject to

the constraints $C_{size\text{-}CD27+} < -0.2$ and $k_I > k_M$. This parameter configuration cannot accurately

capture the moments of the clonal burst data (Fig. 1g). Thus, although the model that

incorporates both NK cell proliferation and death is able produce the negative correlation

between the clone size and the percentage of CD27+ NK cells in each clone, it poorly describes

the distributions of the immature and mature NK cells in the clones.

To further investigate the role of NK cell death in explaining the clone size distribution

we compared the above two-state models without or with NK cell death based on the theory of

model selection using modified Akaike Information Criterion (AIC$_c$) (Burnham *et al*, 2011). We

computed the probability distribution function of the number of immature CD27+ (N$_I$) and

mature CD27- (N$_M$) cells at any time obtained from our stochastic model by solving the

corresponding Master equation numerically and computed the Likelihood function for the

available clone size distribution data (details in the Materials and Methods section), and then

evaluated modified-AIC (AIC$_c$) for the two models. To keep the computation in a feasible range

we used a part (38 out of 56 clone sizes) of the available data where the maximum clone size is ~

1000. The subset of clone size data used here also shows a negative value of $C_{size\text{-}CD27+}$. The

AICc value for the two-state model with death is over 50 points lower than that for the two state

model without death, which clearly show that the data favor the model with NK cell death.

Although this approach only models a subset of the data, it also finds that birth and death rates

for mature cells are greater than that of immature cells, and that the growth rate of immature cells

is greater than that of mature cells. Further details regarding the analysis are provided in the

Materials and Methods section and the Supplementary Material. Because of the computationally

intensive nature of this approach and its agreement with the simpler method of fitting moments

with a constraint, we continue with the later approach for subsequent models. In addition, we

investigated the roles of features considered in Cyton models for T cells(Pandit & De Boer,

2019) such as heterogeneity in the duration of antigen stimulation (Fig. S2), and the dependence

of cell differentiation rates on the number of cell division (Fig. S3). The model where the

durations are distributed in a log-normal distribution, as well the model where the rate of

differentiation increased linearly with the number of cell divisions both generated negative

values of $C_{size\text{-}CD27+}$ when $k_I > k_M$, without cell death (Fig. S2b-d, S3b). Since the distribution of

the duration of activation of NK cells in CMV infections as well as the increasing rate of

differentiation with increasing number of cell divisions are not established for NK cells, we did

not move forward with fitting these models to clonal burst moments. We also investigated an NK

cell proliferation model where the immature NK cell divides asymmetrically to generate a

mature and an immature NK cell (Fig. S4). T cells can undergo asymmetric division where the

daughter cells proximal and distal to the antigen presenting cell are predisposed toward the

effector and the memory fates, respectively(Chang *et al*, 2007); though asymmetric division has

not been observed in NK cells. This model generated negative values of $C_{size\text{-}CD27+}$ but did not

provide good agreement with the moments. The above alternative models contain more

parameters as well as additional mechanisms that not validated in experiments than the minimal

model we considered here. However, these explorations suggest that additional features needed

to be included in the minimal model to capture the heterogeneities observed in clonally expanded

Ly49H+ NK cells. In the next sections, we study some extensions of our two-state model that

can be rationalized against available experiments.

Figure 1: Two-stage linear models cannot capture desired properties of experimental clonal

burst data. (**a**) Schematic representation of the two-stage model. The model describes immature

(CD27+) cells and mature (CD27-) cells. Each cell type has a distinct growth rate (k_I for

immature, k_M for mature), and immature cells differentiate to mature cells at rate r. (**b**)

Experimental data from Flommersfeld *et al.* display negative value for the correlation $C_{size\text{-}CD27+}$

between clonal burst size and %CD27+ cells. (**c**) Schematic description of antigen specific clonal

proliferation and differentiation in CD8+ T cells. Because mature CD62L- cells proliferate more

rapidly compared to the immature CD62L+ cells, initial clones made primarily of CD62L- cells

can grow to generate the larger size clones. **(d)** A parameter scan of all possible combinations of

k_I, k_M, and r shows that negative correlations ($C_{size\text{-}CD27+} < 0$) are not realized when $k_I > k_M$. Each

point represents a value of $C_{size\text{-}CD27+}$ obtained from the model at a unique value of set of model

parameters. The parameter configurations that populate the bottom-right quadrant satisfy our two

constraints imposed by experimental observations regarding the negative values of $C_{size\text{-}CD27+}$ and

higher growth rate of immature NK cells compared to their mature counterparts. **(e)** Schematic

representation of the three-state model including cell death. The model has separate parameters

for birth of immature (CD27+) cells (b_I), death of immature cells (d_I), birth of mature (CD27-)

cells (b_M), death of mature cells (d_M), and differentiation of immature cells to mature (r). In this

model, the growth rates are defined as $k_I = b_I - d_I$ and $k_M = b_M - d_M$. **(f)** A parameter scan for this

model shows that some parameter configurations can meet the two constraints where $k_I > k_M$ and

the parameters result in negative values for $C_{size\text{-}CD27+}$. Each point represents a unique parameter

configuration. Red points denote parameter configurations where $b_M > b_I$ <u>and</u> $d_M > d_I$. **(g)** Shows

the best fit of the model in (e) to the moments with the constraints $k_I > k_M$ and $C_{size\text{-}CD27+} < -0.2$.

Circles represent the fitted values of the first and second moments for the distribution of the NK

cell clones obtained from the model. Blue and red circles represent mean populations of CD27+,

and CD27- NK cells, respectively. The variances of the CD27+ and CD27- NK cells are shown

in cyan and pink circles, and the covariance between the CD27+ and CD27- NK cells is shown in

purple. Horizontal lines around the circles show the standard deviations of bootstrapped

moments. The solid line is the line $y=x$, which demonstrates where values would lie if we had a

perfect fit.

A progressive three-stage differentiation model can quantitatively recreate clonal burst

dynamics

We modified our two-state model to include an intermediate state (Fig. 2a) in the model. This

reflects many investigations probing NK cell maturation, which show NK cell development goes

through several maturation states and some of those states can potentially represent the proposed

intermediate state in the model. A well-established NK cell maturation scheme is CD27+CD11b-

→ CD27+CD11b+ → CD27-CD11b+ (Chiossone *et al.*, 2009). However, the clonal burst data

show a negligible proportion of CD11b- NK cells at 8 days post-infection. Another potential

maturation scheme could be CD27+KLRG1- → CD27+KLRG1+ → CD27-KLRG1+

(Kamimura & Lanier, 2015). In addition, recent single cell RNA-seq experiments show the

presence of CD27-Ly6C+ and CD27-Ly6C- cells at 8 days post-MCMV infection (Riggan *et al.*,

2022a), thus the intermediate state could represent a CD27-Ly6C- state as well. Here we present

results for the model where the intermediate state represents a mature CD27- state, the other

variations of the model are shown in the Supplementary Material (Fig. S5-S7). In order to

determine parameter ranges that give rise to higher growth of mature NK cells compared to their

immature counterparts, we computed the average size of a clone that originates from either a

single CD27+ NK cell or single CD27- NK cell by solving ODEs describing the kinetics of the

average NK cell populations. We constructed an effective growth difference metric Δ_k, which

describes the difference between the average NK cell populations at day 8 that originate from a

single CD27+ or CD27- NK cell at t=0. If Δ_k is positive, then CD27+ cells grow faster than

CD27- cells (Materials and Methods). We performed a parameter scan on this three-stage model

and find that some parameter configurations with $\Delta_k > 0$ can capture the negative values of $C_{\text{size-}}$

$_{CD27+}$ (Fig. 2b). We then evaluated whether this three-stage model can accurately describe the

distributions of the immature and mature NK cells in clones. We fitted the means, variances, and

the covariance of the NK subpopulations in the clones by varying the model parameters with

constraints $\Delta_k > 0$ and $C_{\text{size-CD27+}} < -0.2$ and found that the best-fit three-stage model shows an

excellent fit to the data (Fig. 2c, Table 1). This best-fit parameter set generates a $C_{\text{size-CD27+}}$ of

-0.351, well within the confidence bounds of the correlation coefficient of the data (Fig. 2d).

Thus, adding an intermediate CD27- stage to the model can capture all the key features of the

heterogeneity in the NK cell clonal populations.

Table 1: Best fit parameters to 3-stage model describing NK clonal bursts given in Fig. 2a.

310   Confidence intervals are determined by bootstrapping.

Parameter	Estimate (confidence interval)
b_I	1.064 day^{-1} (1.051 - 1.215)
b_{INT}	1.613 day^{-1} (1.561 - 1.996)
b_M	0.700 day^{-1} (0.001 - 0.727)
d_I	0.054 day^{-1} (0.052 - 0.232)
d_{INT}	0.285 day^{-1} (0.261 - 0.433)
d_M	1.370 day^{-1} (1.330 – 2.000)
r_1	0.126 day^{-1} (0.088 - 0.172)
r_2	0.164 day^{-1} (0.151 - 0.330)

The three-stage model above makes no assumptions about the identities of the differing

phenotypes of the CD27- cell subsets. However, the best fit parameters indicate that the first

CD27- subset has a high proliferation rate, and the final CD27- subset has a high death rate.

Riggan *et al.* (Riggan *et al.*, 2022a) previously determined that, in the contraction phase, Ly6C+

NK cells differentiate from Ly6C- cells, and that Ly6C+ cells die more rapidly than Ly6C- cells.

Our model shows that similar activity may be observed during the expansion phase as well, as

we show higher death of Ly6C+ cells and a unidirectional maturation from Ly6C- $\rightarrow$ Ly6C+

cells. Because our analysis pertains to the expansion phase of the NK cell response to MCMV

we thus aimed to validate if the mature states proposed in the model can be identified with

CD27-Ly6C- and CD27-Ly6C+ NK cells. We found that our best fit parameters correctly

recreate the experimentally observed strong negative correlation between the %CD27+ and

%Ly6C+ NK cells across clones (Grassmann *et al.*, 2019), and correctly estimate that most

clones have <20% Ly6C+ cells at day 8 post-infection (Fig. 2e-f). This independent validation

gives evidence that our parameter estimates can be meaningfully interpreted for a model where

NK cells differentiate along a trajectory from CD27+Ly6C- $\rightarrow$ CD27-Ly6C- $\rightarrow$ CD27-Ly6C+.

We sought to understand the mechanism by which the three-stage model can generate

negative values for the correlation $C_{\text{size-CD27+}}$ as well as quantitatively describe the heterogeneity

in the NK cell populations in the clones. The two-stage model predictions for the first and the

second moments for the NK cell populations are substantially less than the actual values,

indicating that our best-fit parameters are not producing enough NK cells (Fig. 1g). The presence

of the intermediate state in the three-stage model alleviates this issue, which can rapidly grow

and increase the number of CD27-Ly6C+ NK cells while the rapidly dying CD27-Ly6C- NK

cells can help meet the constraint that a population of CD27- cells grows slower than one of

CD27+ cells. This variation increases stochastic fluctuations in the observed numbers of CD27-

cells, as some clones will have many CD27- cells from high growth of the intermediate Ly6C-

subset, while other clones will have few CD27- cells from low growth of the Ly6C+ subset.

Furthermore, the presence of the CD27-Ly6C+ NK cells expand the range of the model

parameters that can give rise to negative values of $C_{\text{size-CD27+}}$.

Figure 2: A three-stage linear model can capture experimental observations of NK cell

clones if intermediate double-positive NK cells are the fastest growing subset. (a) Schematic

representation of the three-stage model. In this model there are three stages of NK cell

maturation: an immature (CD27+), intermediate (CD27-), and terminally mature (CD27-) subset.

Each subset has a distinct proliferation and death rate, and cells progress from immature to

intermediate maturity according to rate r_1 and from intermediate to terminal maturity according

to rate r_2. **(b)** A parameter scan for the model in (a) shows that adding a third stage of maturation

can account for the negative values in $C_{size\text{-}CD27+}$. Each point represents a unique parameter

configuration. Positive values of Δ_k indicate that immature CD27+ cells grow faster than mature

CD27- cells and vice versa. See the Materials and Methods for further details. **(c)** Best fit of this

model to the moments with the constraints $\Delta_k > 0$ and $C_{size\text{-}CD27+} < -0.2$. Circles represent the

fitted values of the moments to the model. Blue and red circles represent mean populations of

CD27+, and CD27- NK cells, respectively. The variances of the CD27+ and CD27- NK cells are

shown in cyan and pink circles, and the covariance between the CD27+ and CD27- NK cells is

shown in purple. Horizontal lines around the circles show the standard deviations of

bootstrapped moments. The solid line is the line $y=x$, which demonstrates where values would lie

if we had a perfect fit. **(d)** Clones stochastically simulated with Gillespie's algorithm (Materials

and Methods) from the best fit parameters and the resulting clone sizes, CD27+ percentages, and

correlation. **(e)** Observed clonal compositions of Ly6C+ and CD27+ cells from stochastically

simulated clones. **(f)** Simulated clones from the three-stage model in (a) at the best fit parameter

values shown in (c) when the three stages are taken as immature CD27+Ly6C-, mature CD27-

LyC- and the terminally mature CD27-Ly6C+, respectively. Each point is a simulated clone

resulting from a single NK cell.

Comparison of kinetics between endogenous NK cells, adoptively transferred cells, and

homeostatic cells identifies potential identical maturation rates in all conditions

Here we compare the kinetics of NK cell maturation occurring in response to MCMV infection

and in homeostasis. In particular, we investigated the changes in the rates of cell growth and

differentiation induced by MCMV infection. First, we quantified the growth and the

differentiation rates using time-stamped flow cytometry data on homeostatic NK cells in

peripheral blood with a one-time treatment of tamoxifen in fluorescent NK cell reporter mice

reported by Adams *et al*.(Adams *et al*, 2021). Treating the mice with tamoxifen determines

whether observed NK cells are newly populated from bone marrow or existed in peripheral blood

at the onset of the infection. We fitted a two-state CD27+ → CD27- maturation model described

by ordinary differential equations (see Materials and Methods) to the data collected at days 3, 7,

14, 23, 29, and 35 to estimate the growth rates of the immature CD27+ (k_{CD27+}) and the mature

CD27- (k_{CD27-}) NK cells, the rate of differentiation (*r*) of the CD27+ to CD27- NK cells, and the

rate (λ) with which CD27+ NK cells are populated from bone marrow. We estimated these rates

both for Ly49H+ and Ly49H- NK cells. The estimated rates for Ly49H+ cells (Table 2, Fig. 3a)

show that a CD27+ cell population would double every 6.9 days if there were no differentiation,

whereas CD27- cells have a negative growth rate with a half-life of 5.6 days. This implies that

the death rate of a population of CD27- cells is greater than their proliferation rate. The

differentiation rate from CD27+ → CD27- phenotypes is similar to that of the death rate of

Ly49H+ CD27- cells. Approximately 1.37% of the Ly49H+ NK pool is replenished from bone

marrow every day. Comparing these rates to those of the Ly49H- NK pool reveals that Ly49H-

NK cells are repopulated from bone marrow at almost twice the rate (~2.4%/day). All other

estimated rates for Ly49H- NK cells are within the confidence bounds of the Ly49H+ estimated

rates (Table 2, Fig. 3b). Given the constraint governing λ which dictates

$$\lambda \propto N_{CD27-}$$

390   (Materials and Methods) the difference in estimated λ values can be attributed to a higher

391   percentage of CD27- cells. Next, we compared these rates with corresponding rates estimated in

our three-state model using the clonal burst data. The estimate for CD27+ → CD27-

differentiation rate is equivalent between Ly49H+ NK cells in homeostasis and in MCMV

infection. Our estimations for the clonal burst data (Table 1) show that varying subsets of CD27-

cells have varying growth rates ranging -0.7 day^{-1} to 1.3 day^{-1} in MCMV infection, a wide range

not in disagreement with the estimate for homeostatic data. Ly49H+ CD27+ cells grow ~10×

faster in MCMV infection than in homeostatic conditions.

Table 2: Best fit parameters to homeostatic Ly49H+ and Ly49H- NK cells. Confidence intervals

are determined by bootstrapping.

Parameter	Ly49H+ Estimate (confidence interval)	Ly49H- Estimate (confidence interval)
λ	1.366 percent/day (1.117– 1.594)	2.412 percent/day (2.028 – 2.915)
r	0.125 day^{-1} (0.095 – 0.159)	0.147 day^{-1} (0.118 – 0.173)
k_{CD27+}	0.097 day^{-1} (0.064 – 0.135)	0.088 day^{-1} (0.053 – 0.119)
k_{CD27-}	-0.120 day^{-1} (-0.152 – -0.091)	-0.107 day^{-1} (-0.126 – -0.086)

We investigated whether the kinetics of NK cell maturation in the expansion phase during

MCMV infection in single-cell adoptive transfer experiments is quantitatively similar to that of

an endogenous population of NK cells. To this end, we collected time-stamped mass cytometry

data on C57BL/6 mice splenocytes at days 0, 4, and 7 post-MCMV infection (Potempa *et al.*,

2022). We then gated Ly49H+ NK cells from each mouse and determined the mean relative

abundances across mice of CD27+Ly6C-, CD27-Ly6C-, and CD27-Ly6C+ subsets among the

NK cell population (Fig. S8). We multiplied these relative abundances by mean absolute

Figure 3: Kinetics of homeostatic NK cells and endogenous NK cells responding to MCMV

infection. (a-b) Best fit to **(a)** Ly49H+ and **(b)** Ly49H- homeostatic NK cells. Y axis refers to

percentage of the total Ly49H+ or Ly49H- population occupied by that NK subset. Circles and

Xs represent the mean values of tamoxifen-induced td-Tomato positive and negative NK cells

observed in the data, respectively. Solid and dashed error bars represent standard deviations

around the means of tamoxifen-induced positive and negative NK abundances respectively.

Smooth curves represent model fits for tamoxifen-induced positive (solid) and negative (dashed)

NK cells. Shaded regions refer to model 95% confidence bands, as determined by bootstrapping

model fits. **(c)** Best fit to profiled endogenous NK cells responding to MCMV infection. Points

represent mean cell abundances, and error bars represent standard deviations of bootstrapped

means. Smooth curves represent model fit.

numbers of Ly49H+ endogenous NK cells during MCMV infection from Robbins *et al.*(Robbins

et al., 2004) to obtain time-stamped population sizes of NK cell subsets. We then modeled these

data with the ordinary differential equation (ODE) solution in Equations S17-S25 using the

parameters from Table 1 and an initial condition of the mean uninfected *in vivo* abundances of

endogenous NK cells. The ODE result is multiple orders of magnitude above the measured

abundances (Fig. S9a). This indicates that the kinetics of endogenous NK cells differ

significantly from that of NK cells in adoptive transfer experiments. Further indication that the

kinetics differ is the expansion of cells in endogenous conditions over 7 days is less than a 10-

fold increase, while clonal expansion accounts for >1000-fold increase.

To further quantify the differences in the kinetic rates between the adoptively transferred

and endogenous NK cells, we then fit the 3-stage model with 2 mature stages to the endogenous

cell population. We used the gated Ly49H+ populations of CD27+/Ly6C-, CD27-/Ly6C-, and

CD27-/Ly6C+ NK cells and fit to a system of ODEs that describe the net growth rates of each

population and the differentiation rates between them (Materials and Methods). The best fit to

this data has unrealistic estimates for the death rate of CD27-/Ly6C+ NK cells and the

differentiation rate from CD27+ → CD27- cells (Table S5, Fig. S9b). We then attempted to

constrain the fitting procedure by setting the CD27+ → CD27- differentiation rate to those of the

previous estimations for the clonal burst kinetics and homeostatic conditions (Tables 1 and 2).

Under this constraint, the only rate estimated to be within the confidence interval for the

equivalent rate in adoptive transfer experiments is the differentiation rate from Ly6C- → Ly6C+

(Table 3, Fig. 3c). The growth rate of CD27+Ly6C- cells is ~4× faster for adoptively transferred

cells, the growth rate of CD27-Ly6C- cells is 2-3× faster, and the death rate of mature cells is

~1.5× faster. However, the resulting best fit to the endogenous NK cell population does show

that the intermediate CD27-/Ly6C- cell population has the fastest growth rate, and the mature

CD27-/Ly6C+ cells have a large death rate, confirming the qualitative results from the adoptive

transfer experiments. Thus, although the growth kinetics are different between endogenous and

adoptive transfer experiments, both systems have the intermediate population proliferating

rapidly and the terminally mature population dying rapidly.

Table 3: Best fit parameters to endogenous NK cells responding to MCMV infection with

constrained r_1. Confidence intervals are determined by bootstrapping.

Parameter	Estimate (confidence interval)
k_I	0.236 day^{-1} (0.153 - 0.350)
k_{INT}	0.537 day^{-1} (0.065 - 0.874)
k_M	-0.484 day^{-1} (-1.194 - 0.371)
r_1	0.126 day^{-1} (fixed)
r_2	0.447 day^{-1} (0.000 - 0.874)

Both the adoptively transferred cells and endogenous population are estimated to have a large

death rate for CD27- cells. To test this result, we analyzed NK cell viability in mice that were

infected with MCMV at different times post-infection (Fig. 4a). Surprisingly, mature (CD27-)

cells indeed had a greater fraction of dead cells than immature (CD27+) cells during expansion

(days 4 and 7 post-infection) (Fig. 4b). To determine if the higher rates of cell death in the

CD27- NK cell population in the three-state model give rise to a higher fraction of dead NK cells

we extended our three-state model to evaluate the number of dead mature and immature NK cells

(Fig. S10c) which showed that the best fit parameters for the adoptive transfer experiments are

consistent with the results reported above (Fig. 4c). Because we only have best fit growth rates of

the endogenous NK cells and not separate proliferation and death rates, we tested a variety of

birth and death rates consistent with the best fit parameters such that $b\text{-}d\text{=}k$ for each cell subset.

Many of the possible configurations agree with the higher percentage of live CD27+ cells. This

gives additional confirmation that CD27- cells may in fact be dying at faster rates than CD27+

cells.

Figure 4: Mature CD27- NK cells undergo rapid death during the expansion phase. (a)

Schematic of experimental setup. **(b)** Relative abundances of dead cells for CD27+ (blue) and

CD27- (red) *ex vivo* NK cell subsets as measured by flow cytometry experiment. Data are

representative of experiments with n = 4-5 mice per timepoint. Graph shows means ± SEM; *P

<0.033, and ***P <0.001 represent statistically significant difference between CD27$^+$ and CD27$^-$

NK cells as determined by 2-way ANOVA. **(c)** Parameter scans for determining percentage of

live cells for CD27+ and CD27- subsets using the model shown in Fig. S10c. Blue points

represent a parameter scan where birth and death rates were varied by random sampling such that

the growth rates $k_I = b_I-d_I$, $k_{INT} = b_{INT}-d_{INT}$, and $k_M = b_M-d_M$ are equivalent to the kinetic

estimates for endogenous NK cells shown in Table 3. Red points represent randomly sampled

dead cell clearance rates with other rates set to those describing adoptive transfer kinetics in

Table 1. Black dotted line is *y=x*, and points below this line show a higher percentage of live

cells for immature CD27+ cells than for mature CD27- cells.

Figure 5: Proposed mechanism of NK clonal expansion in response to MCMV infection.

Three distinct subsets of NK cells are necessary to generate full NK response to infection. The

first is CD27+, Ly6C-, and moderately proliferative, with a doubling time of ~0.3 days and

negligible cell death. The second is CD27-, Ly6C-, and highly proliferative, with a doubling time

of ~0.2 days. The last is CD27-, Ly6C-, and prone to rapid cell death with a half-life of ~0.45

days. This picture paints CD27-Ly6C- NK cells as having a highly proliferative and highly

effective phenotype, similar to effector CD8+ T cells, and CD27+Ly6C- cells as long-lived,

similar to memory precursor CD8+ T cells.

Discussion

We employed stochastic minimal mechanistic *in silico* models to investigate mechanisms that

underlie development of MCMV-specific NK cells at early times (0 – 8 days post-infection). We

found Ly49H+ NK cells undergoing antigen-specific expansion follow a maturation scheme

from (immature) CD27+Ly6C- → (mature I) CD27-Ly6C- → (mature II) CD27-Ly6C+,

characterized by a highly proliferative mature CD27-Ly6C- phenotype and a more mature CD27-

Ly6C+ phenotype with higher rates of cell death (Fig. 5). The differences in the rates of

proliferation and cell death are crucial to give rise to the heterogeneous clones that originate

from single immature CD27+ NK cells, in particular, the presence of larger sized NK cell clones

composed of mature CD27- NK cells at 8 days post-infection. Similar heterogeneity in the clones

of CD8+ T cells generated in the expansion phase (<8 days post-infection) from an acute

antigen-specific expansion of single immature CD62L+ cells in C57BL/6 mice has been

observed (Buchholz *et al.*, 2013); however, the larger CD8+ T cell clones composed of effector

CD62L- arise from the higher rate of proliferation of the CD62L- CD8+ T cells. This is in

contrast with the NK cell maturation program where the larger sized clones of mature NK cell

populations arise due to the interplay of uneven increase in the proliferation rates and increase in

the rates of cell death as the NK cells mature, and intrinsic stochastic fluctuations occurring in

these processes. Innate heterogeneity in adoptively transferred single NK cell phenotypes may

also contribute to observed stochasticity. In CD8+ T cells, precursor CD62L+ cells are deemed

memory precursors, matching their lowly proliferative phenotype (Kretschmer *et al.*, 2020). An

equivalent population of memory precursor NK cells has not been identified. CD27-Ly6C+ cells

match previously defined phenotypes for memory NK cells and could potentially be memory

precursors. However, our model shows that these cells experience rapid cell death. Therefore, for

these cells to live longer to become memory cells additional processes such as a change in

environmental conditions or an epigenetic change that promotes survival in these cells at later

times during the contraction phase could become relevant. This differentiation trajectory is also

in contrast with CD8+ T cells, as NK cell memory precursors would be formed downstream of

effector (Ly6C-) NK cells, whereas CD8+ effectors (CD62L-) are formed downstream of CD8+

T cell memory precursors (CD62L+) in acute infection. Alternately, another mature NK subset

such as Ly6C-KLRG1- that is lowly proliferative and long-lived could give rise to the adaptive

NK cell phenotype. Inclusion of these processes as well as the later time kinetics could be a

future extension of our approach. We note a similarity of our three state NK cell maturation

model with maturation of T cells in persistent (13-140 days) *T. gondi* infection in genetically

resistant mice reported by Robey and colleagues (Chu *et al*, 2016) where the infection gave rise

to a continuous production of CXCR3-KLRG1+ terminally differentiated effector T cells from a

quiescent memory CXCR3+KLRG1- T cells via an intermediate CXCR3+KLRG1+ state. The

intermediate T cell state showed higher proliferation and increased death compared to the

memory and effector T cells. However, these T cells were uniformly CD62L- in contrast to the

memory and effector T cells that are produced in response to acute antigenic SIINFEKL

infection.

Integration of signals in NK cells during MCMV infection leads to epigenetic remodeling

affecting NK cell differentiation, proliferation and death(Riggan *et al*, 2022b; Rückert &

Romagnani, 2024; Santosa *et al*, 2023). In T cells, the sum of signals received during clonal

expansion has been found to regulate the number of division and the time of death or the division

destiny in individual T cells which can arise due to a global remodeling of the transcriptional and

epigenetic program(Iwata *et al*, 2017; Rückert & Romagnani, 2024). We considered a variant of

the Cyton models(De Boer & Yates, 2023) that have been employed in modeling clonal

expansion in T cells where the rate of differentiation from the immature to the mature state

depends on the number of divisions in individual NK cells. This model can give rise to negative

values in the correlation between the clone size and the percentage of immature CD27+ NK cells

in the absence of NK cell death, suggesting that inheritance of epigenetic states regulating cell

fates in individual NK cells can play an important role in clonal expansion. Furthermore, we also

find the presence of different time delays in NK cells to encounter appropriate activation signal

in a pure birth model with two states can give rise to negative values in $C_{\text{size-CD27+}}$ implicating a

role of different activation times during the clonal expansion. The exploration of division destiny

and activation times in NK cells during clonal expansion could be an exciting future direction.

NK cell clones in MCMV infection go through a smaller (~1000×) fold increase in the

expansion phase compared that (~40,000×) of antigen-specific expansion of CD8+ T cells. This

could be because CD8+ T cells can generate autocrine cytokines such as IL-2 that help the cells

to proliferate (Kalia & Sarkar, 2018), whereas NK cells rely on other cells to make the necessary

cytokines for proliferation (such as IL-2 and IL-15) and survival (such IL-21) (Fauriat *et al*,

2010). Furthermore, NK cells are more dependent on cytokines such as IL-15 compared to the

CD8+ T cells (DeGottardi *et al*, 2016). Therefore, cytokine competition for IL-15 and IL-2 might

also explain increased death of mature NK cells in the expansion phase. A test of this hypothesis

could be investigation of MCMV-induced NK cell maturation in *Il15+/-* mice.

We quantified the population kinetics of antigen-specific Ly49H+ NK cells during the

expansion phase of MCMV infection (<8 days post-infection) using both endogenous and

adoptively transferred NK cells. This antigen-specific response was further compared with the

kinetics of NK cell maturation in the absence of MCMV infection. Adoptively transferred

Ly49H+ NK cells grow at an increased rate compared to the endogenous Ly49H+ NK cells (~3×

faster for Ly6C- cells). A possible explanation of this phenomenon is the stronger competition

for resources (e.g. antigens and/or cytokines) that would occur in endogenous NK cells could

prevent them from growing as quickly as the less dense population of adoptively transferred NK

cells. Interestingly, our estimate of the differentiation rate of the immature CD27+ to mature

CD27- NK cells showed negligible effect of MCMV infection on this process. This shows that

the signaling processes governing NK cell antigen-specific expansion and cytotoxicity are not

impacting transcription and translation of CD27, but rather preferential expansion of certain

subsets is creating altered phenotype abundances observed in MCMV infection.

Our stochastic modeling of heterogeneous NK cell clones showed that larger stochastic

variations in the mature NK cell populations arising due to higher intrinsic noise fluctuations

generated by the interplay between the higher proliferation and cell death processes are

important for producing large clones of mature NK cells at 8 days post-infection. This is in

contrast with stochastic kinetics of CD8+ T cells where the larger clones of effector CD62L-

CD8+ T cells can arise due to a higher rate of proliferation of these cells even in the absence of

any cell death (Buchholz *et al.*, 2013). Our effort to use different models to quantitatively

describe the heterogeneity of NK cell clones demonstrate the critical role of the correlation

between the clone size and the proportion of the mature NK cells in the clones, $C_{\text{size-CD27+}}$, in

screening out mechanisms that are unable to give rise to the observed heterogeneity.

Materials and Methods

Calculation of the first and second moments and $C_{size\text{-}CD27+}$ for the NK cell subpopulations

in the clones

The first and the second moments for the number of immature CD27+ ($N_I^{(c)}$) and mature CD27-

($N_M^{(c)}$) NK cells in n_{clone} number of clones ({c}) are defined as follows:

$$\overline{N_I} = 1/n_{clone} \sum_{\{c\}} N_I^{(c)}$$

$$\overline{N_M} = 1/n_{clone} \sum_{\{c\}} N_M^{(c)}$$

$$\overline{N_I^2} = 1/n_{clone} \sum_{\{c\}} \left(N_I^{(c)}\right)^2$$

$$\overline{N_M^2} = 1/n_{clone} \sum_{\{c\}} \left(N_M^{(c)}\right)^2$$

$$\overline{N_I N_M} = 1/n_{clone} \sum_{\{c\}} N_M^{(c)} N_I^{(c)}$$

590   The correlation $C_{size\text{-}CD27+}$ is defined as,

$$C_{size-CD27+} = \left(\sigma_T \sigma_{\frac{N_I}{T}}\right)^{-1} \left[1/n_{clone} \sum_{\{c\}} T^{(c)} \frac{N_I^{(c)}}{T^{(c)}} - \left[1/n_{clone} \sum_{\{c\}} T^{(c)} \right] \left[1/n_{clone} \sum_{\{c\}} \frac{N_I^{(c)}}{T^{(c)}} \right] \right]$$

where, $T^{(c)} = N_I^{(c)} + N_M^{(c)}$, $\sigma_T^2 = 1/n_{clone} \sum_{\{c\}} \left(T^{(c)}\right)^2 - \left(1/n_{clone} \sum_{\{c\}} T^{(c)}\right)^2$, and, $\sigma_{N_I/T}^2 =$

$1/n_{clone} \sum_{\{c\}} \left(N_I^{(c)}/T^{(c)}\right)^2 - \left(1/n_{clone} \sum_{\{c\}} N_I^{(c)}/T^{(c)}\right)^2$.

NK cell development model

We modeled the NK cell proliferation, death, and maturation as a continuous time Markov

processes where a proliferation or a death event in a NK cell population increases or decreases

the population number, respectively, by one cell. A differentiation event changing a NK cell

from immature (*I*) to mature (*M*) decreases the NK cell population in state *I* and increases the

population in state *M* simultaneously by a single NK cell. The kinetics of the system are given by

the Master equation describing the time evolution of the conditional probability for the system to

be in a particular state containing different numbers of immature and mature NK cells at any

time *t* given a specific state at *t=0*. The ODEs describing the kinetics of first and second

moments can be derived from the Master equation. The solutions are shown in the

Supplementary Materials. The first and second moments for the clone size distributions for all

our models were calculated by solving the corresponding ODEs.

Stochastic simulations

We consider the cell proliferation, differentiation, and death processes as continuous time

Markov processes which are modeled as simple birth (n # of cells $\rightarrow$ n+1 # of cells), death (n# of

cells $\rightarrow$ n-1 # of cells), and differentiation (N_I # of immature cells, N_M # of mature cells $\rightarrow$ N_I - 1

# of immature cells, N_M +1 # of mature cells) processes. The propensities for the above

transitions are computed for a range of rates described in Table 1, tables in the supplement, and

in the parameter scan section. We use the Gillespie algorithm(Gillespie, 1977) for all our

stochastic simulations. For two-stage and three-stage models, we initialized the simulation with a

single CD27+ NK cell. We evolved the system until 8 days and numbers of CD27+ and CD27-

cells in the NK cell clones were recorded. We calculated the correlation $C_{\text{size-CD27+}}$ for the clones

by averaging over 10^4 stochastic trajectories.

Two-state model with differentiation rate increasing with cell division number

This is variant of a Cyton model where the rates of proliferation, differentiation, and death

individual cells are inherited from mother to the daughter cells in dividing cells. In our model,

we considered a two-state model where immature CD27+ NK cells differentiate into mature

CD27- NK cells, and both immature and mature NK cells replicate with rates k_I and k_M (k_I>k_M).

In the model, we tracked the number of divisions experienced by an individual immature cell

where the rate of differentiation proportional to the number divisions (n_{division}) the cell has

undergone, i.e., the rate of differentiation in the cell after n_{division} is r_{diff}=$r_0 \times n_{\text{division}}$, where r_0 is

the differentiation rate in the founder cell after the first division (Fig. S3). We do not consider

any cell death in the model. We implement a continuous time Monte Carlo simulation following

Gillespie's approach to simulate cell proliferation and differentiation in the system. More

computational details of this multi-scale method that tracks single-cell divisions can be found in

Wethington et. al (Wethington *et al*, 2025). We adapt the framework in (Wethington *et al.*, 2025)

to include, 1) a propensity for division of immature cells, 2) a propensity for division of mature

cells, and 3) a propensity of differentiation that is linearly related to the number of divisions a

cell has undergone.

Two-state model with log-normally distributed activation times

638 For the model with log-normally distributed activation times, we assume an NK cell received
639 stimulation at time 8-*t* days and then expanded for a duration of *t* days until day 8. We assign
640 based on the distribution

$$X \sim Lognormal(1, 0.5)$$

$$t = \min(X, 8)$$

641 Each clone is evaluated for a randomly sampled *t* according to this distribution.

Fitting model to moments

645 The first and second moments $f(\vec{\theta})$ for the clone size distributions obtained for two- and three-

646 stage models were computed by solving ODEs (Eqs. S2-S6 in Supplementary Text 1 and the

647 ODEs in Supplementary Text 2). The parameter vector $\vec{\theta}$ represents the rates for proliferation,

648 differentiation, and death in the models. To compare the predicted moments $f(\vec{\theta})$ from the

649 model with the actual moments *y* computed from the data, we used the cost function below:

$$\chi^2 = \sum_i \left(\frac{f_i(\vec{\theta}) - y_i}{\sigma_i} \right)^2 + g(\vec{\theta})$$

where, σ_i is the standard deviation corresponding to the data used for computing y_i. A penalty

function, $g(\vec{\theta})$ is introduced to implement the constrained imposed by sign of the correlation

$C_{size\text{-}CD27+}$. We set $g(\vec{\theta})$ to a high value (= 10^4) when $C_{size\text{-}CD27+}$ > -0.2 at any value of $\varDelta_k$ or

$C_{size\text{-}CD27+}$ < 0 and $\varDelta_k$ < 0, otherwise $g(\vec{\theta})$=0. This constraint describes the experimental

observation that immature CD27- NK cells grow faster than mature CD27+ NK cells in the

expansion phase. σ_i is calculated by taking the standard deviation of each moment across 10,000

bootstrapped samples of the clones. Minimization of the cost function was performed with the

lmfit python package, using the 'least_squares' Trust Region Reflective algorithm. Parameter

search space was confined from 0.0 to 2.0 day^{-1} for all parameters. We also performed a

parameter fitting without any constraint on Δ_k , instead fitting the ratio of cells arising from a 1:1

mix of CD27+/CD27- NK cells following the experiments in Flommersfeld *et al.(Flommersfeld*

et al., 2021), adding one additional data point and residual to the model. The resulting best-fit

parameters represent Δ_k <0, thus we reasoned these estimated parameters do not capture the NK

cell expansion observed in experiments (Table S6). This is in line with the previous findings that

setting CD27- as the faster growing population can match observed moments data, but this must

be balanced with the finding that CD27+ cells grow faster (i.e. Δ_k >0).

Effective Growth Rate for 3-stage system

We developed a metric to quantify the difference in the growth rate of immature CD27- and

mature CD27+ NK cells for the three-stage model. We computed mean populations of the NK

cell subsets at day 8 by solving Eqs. S17-S25 in Supplementary Text 2 for two different initial

conditions. In one case we initialized with the immature CD27+Ly6C- cells, and in the other

case we initialized with equal proportions of mature CD27-Ly6C- and mature CD27-Ly6C+

cells. We then computed the "effective growth difference" ($\varDelta_k$) by evaluating the difference in

the mean populations of the immature CD27+ and the mature CD27- NK cells at day 8. We

performed an asinh(Δ_k) transformation to help visualization of the data in the graphs shown in

the manuscript.

Parameter scans

For parameter scans showing the relationship between growth rates and the correlation between

clone size and %CD27+ cells, each parameter is randomly selected from a uniform distribution

from 0 to 1.6 day^{-1}. For each parameter set, 10^4 Gillespie simulations were carried out to

compute the correlation $C_{\text{size-CD27+}}$ for the NK cell clones at day 8. We performed scans for 10^5

different parameter sets.

Ordinary differential equation fits to homeostatic data

The growth and differentiation kinetics of populations of CD27+Tom+ (n_1), CD27-Tom+ (n_2),

CD27+Tom- (n_3), and CD27-Tom- (n_4) cells in the periphery in homeostasis are given by the

ODEs below:

$$\frac{dn_1}{dt} = k_{CD27+}n_1 - rn_1$$

$$\frac{dn_2}{dt} = k_{CD27-}n_2 + rn_1$$

$$\frac{dn_3}{dt} = k_{CD27+}n_3 - rn_3 + \alpha$$

$$\frac{dn_4}{dt} = k_{CD27-}n_4 + rn_3$$

692   The CD27+ and CD27- cells grow with rates k_{CD27+} and k_{CD27-} , respectively, and the

693   differentiation CD27+ → CD27- occurs at a rate, r. The influx of immature CD27- cells from the

694   bone marrow to the periphery occurs at a rate α. At homeostasis it is reasonable to assume that

695   the sum of all Ly49H+ and the sum of all Ly49H- NK cells were assumed to be constant through

696   time, *i.e.*,

$$n_{TOT}(t = 0) = n_{TOT}(t) = \sum_i n_i(t)$$

697   The kinetics of the percentages ($N_i = 100 \times (n_i/n_{TOT})$) of each cell type follow the ODEs below:

$$\frac{dN_1}{dt} = k_{CD27+}N_1 - rN_1$$

$$\frac{dN_2}{dt} = k_{CD27-}N_2 + rN_1$$

$$\frac{dN_3}{dt} = k_{CD27+}N_3 - rN_3 + \frac{\alpha}{n_{TOT}}$$

$$\frac{dN_4}{dt} = k_{CD27-}N_4 + rN_3$$

699   We define $\lambda = \dfrac{\alpha}{n_{TOT}}$, which describes the influx rate of either the total Ly49H+ or total Ly49H-

700   population as a percentage of the given population. This makes the ODEs which we fit:

$$\frac{dN_1}{dt} = k_{CD27+}N_1 - rN_1$$

$$\frac{dN_2}{dt} = k_{CD27-}N_2 + rN_1$$

$$\frac{dN_3}{dt} = k_{CD27+}N_3 - rN_3 + \lambda$$

$$\frac{dN_4}{dt} = k_{CD27-}N_4 + rN_3$$

701   We assume the total number of NK cells and the relative abundances of varying subsets do not

702   change with time in homeostasis, therefore,

$$\frac{d(N_1 + N_2 + N_3 + N_4)}{dt} = k_{CD27+}(N_1 + N_3) + k_{CD27-}(N_2 + N_4) + \lambda = 0$$

$$\frac{d(N_1 + N_3)}{dt} = k_{CD27+}(N_1 + N_3) - r(N_1 + N_3) + \lambda = 0$$

$$\frac{d(N_2 + N_4)}{dt} = k_{CD27-}(N_2 + N_4) + r(N_1 + N_3) = 0$$

703   These constraints lead to equalities between parameters,

$$r = -\frac{k_{CD27-}N_{CD27-}}{N_{CD27+}}$$

$$\lambda = -(k_{CD27-}N_{CD27-} + k_{CD27+}N_{CD27+})$$

Thus, by estimating only k_{CD27+} and k_{CD27-}, and using the mean relative abundances of CD27+

and CD27- cells across mice and time, we can solve all four rate parameters in the homeostatic

state.

Ordinary differential equation fits to mass cytometry data

We gated Ly49H+ NK cells in the CyTOF studies into three subsets: CD27+Ly6C-, CD27-

Ly6C-, and CD27-Ly6C+. Details of the gating are described in the Supplementary Material. We

describe the growth and differentiation kinetics of the populations the immature CD27+Ly6C-

(N_I), mature CD27-Ly6C- (N_{INT}), and mature CD27-Ly6C+ (N_M) by the ODEs below,

$$\frac{dN_I}{dt} = k_I N_I - r_1 N_I$$

$$\frac{dN_{INT}}{dt} = k_{INT}N_{INT} + r_1 N_I - r_2 N_{INT}$$

$$\frac{dN_M}{dt} = k_M N_M + r_2 N_{INT}$$

Where, k_I, k_{INT}, and k_M denote the rates of growth for N_I, N_{INT}, and N_M, respectively. The rates of

differentiation for CD27+Ly6C- → CD27-Ly6C- and CD27-Ly6C- → CD27-Ly6C+ are given

by r_1 and r_2, respectively. Since the above system of ODEs is autonomous and linear, we can

obtain the time dependent solutions analytically and exactly. The analytical solution was

numerically evaluated for different values of the rate parameters represented by $\vec{\theta}$. The analytical

solutions were evaluated numerically for specific rate constants and initial conditions. We used

the estimated population sizes (details in the main text) for the above NK cell subsets in the

CyTOF data at t=0 as the initial conditions and computed the population sizes at day 4

($\{N_i(\vec{\theta}, \text{t=4d})\}$) and day 7 ($\{N_i(\vec{\theta}, \text{t=7d})\}$) by solving the above ODEs for a parameter set $\vec{\theta}$. We

then estimated the parameters by $\vec{\theta}$ by fitting the population sizes at day 4($\{y_i(\vec{\theta}, \text{t=4d})\}$) and day

7 ($\{y_i(\vec{\theta}, \text{t=7d})\}$) using the cost function below.

$$\chi^2 = \sum_i \left(\frac{N_i(\vec{\theta}, t = 4d) - y_i(t = 4d)}{\sigma_i} \right)^2 + \left(\frac{N_i(\vec{\theta}, t = 7d) - y_i(t = 7d)}{\sigma_i} \right)^2$$

Bootstrapping confidence intervals for parameter estimates

To assess the confidence intervals for parameter estimates, we randomly sampled the clones with

replacement with equal probability, calculated each moment and its standard deviation, and fit

the moments as described above. We repeated the process 1,000 times, yielding 1,000 estimates

for the parameters θ. We then sorted the parameter estimates by magnitude and took the 2.5[th] and

97.5[th] percentile for each. These estimates made up the 95% confidence intervals reported here.

For estimations of the homeostatic and endogenous NK populations, we sampled a

population of mice before sampling errors in measurements for each bootstrap estimate.

Bootstrapping confidence intervals for model fits

The model fits in Figures 3a and 3b have confidence intervals around them. These are calculated

by bootstrapping parameter estimates as described above, and simulating a trajectory for each

bootstrapped parameter estimate set. At each time t, we take the 2.5[th] and 97.5[th] percentile of the

model trajectory for each cell type. These make up the lower and upper limit for the 95%

confidence interval at time t.

Animals and MCMV infection

C57BL/6 (B6) B6 mice were purchased from the National Cancer Institute at 6 weeks of age and

housed in the specific pathogen-free animal facility at the University of California San Francisco

(UCSF) in accordance with the guidelines of the Institutional Animal Care and Use Committee.

Experiments used mice of both genders, aged 8-12 weeks.

B6 mice were infected with 1000 plaque-forming units (PFU) of Smith strain MCMV via

intraperitoneal injection. Spleens were acquired different time point following MCMV infection

and examined by flow cytometry.

Flow Cytometry

Splenocytes from uninfected or infected mice were processed into single cell suspensions, RBC lysed, and then stained for flow cytometry. Surface staining was performed using pre-conjugated antibodies in the presence of the viability marker ZombieRed (BioLegend) and were then analyzed on an LSR II cytometer (BD Biosciences). The antibodies used for flow cytometry were: CD3ε (145-2C11), CD19 (6D5), CD49b (DX5), NK1.1 (PK136), KLRG1 (2F1), Ly6C (HK1.4), CD27 (LG.3A10), and Ly49H (3D10). All antibodies were purchased from BioLegend, BD Biosciences, or ThermoFisher Scientific. An unconjugated anti-CD16/CD32 antibody (clone 2.4G2, UCSF Antibody Core) was used to block non-specific binding.

Mass cytometry

Samples were prepared for mass cytometry as previously described (Potempa *et al.*, 2022). Briefly, splenocytes from each mouse were obtained by mechanical dissociation through a 40-µm filter, treated with ACK lysis buffer, and resuspended in PBS + 5 mM EDTA (PBS/EDTA). Cells were stained for viability by mixing 1:1 with PBS/EDTA + 100 mM cisplatin (Enzo Life Sciences) for 1 minute before quenching with PBS/EDTA + 0.5% BSA (Staining Media). One million splenocytes from each mouse was stained with metal-conjugated antibodies against CD16 (Nd144) and CD32 (Eu153) for 15 minutes at room temperature in Staining Media before supplementation with an Fc blocking Ab (clone 2.4G2). After washing in Staining Media, cells

were stained with a master mix of 35 additional metal-conjugated antibodies (Table S7) for 30

minutes at room temperature. At the conclusion of the staining period, cells were washed in

PBS/EDTA before fixation with PBS/EDTA + 3.2% paraformaldehyde and stored at 4º C for 3-5

days. One day prior to data acquisition, samples were barcoded using a 20-plex Pd barcoding kit

(Fluidigm) per manufacturer protocol, pooled, and fixed again, overnight at 4º C, in a

PBS/EDTA + 3.2% paraformaldehyde solution supplemented with a 1:1000 dilution of a

191/193 DNA intercalator (Fluidigm). Just prior to acquisition, barcoded samples were pelleted,

washed in double-deionized water, and resuspended in double-deionized water with a 1:20

dilution of EQ four Element Calibration Beads (Fluidigm). Data were acquired using a CyTOF2-

Helios mass cytometer (Fluidigm), which were then normalized and debarcoded prior to

analysis.

Acknowledgements

This work was supported by NIH grant R01AI146581 to J.D. N.M.A. was supported by a

Medical Scientist Training Program grant from the National Institute of General Medical

Sciences (T32GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-

PhD Program) and by an F30 Predoctoral Fellowship from the National Institute of Allergy and

Infectious Diseases (F30 AI136239). We would like to thank the Ohio Supercomputer Center for

computational resources.

Abel AM, Yang C, Thakar MS, Malarkannan S (2018) Natural Killer Cells: Development,
Maturation, and Clinical Utilization. *Front Immunol* 9: 1869

Adams NM, Diaz-Salazar C, Dang C, Lanier LL, Sun JC (2021) Cutting Edge: Heterogeneity in Cell
Age Contributes to Functional Diversity of NK Cells. *J Immunol* 206: 465-470
Adams NM, Grassmann S, Sun JC (2020) Clonal expansion of innate and adaptive lymphocytes.
Nat Rev Immunol 20: 694-707
Biassoni R (2008) Natural killer cell receptors. *Adv Exp Med Biol* 640: 35-52
Bresser K, Kok L, Swain AC, King LA, Jacobs L, Weber TS, Perié L, Duffy KR, de Boer RJ, Scheeren
FA (2022) Replicative history marks transcriptional and functional disparity in the CD8+ T cell
memory pool. *Nature immunology* 23: 791-801
Buchholz VR, Flossdorf M, Hensel I, Kretschmer L, Weissbrich B, Graf P, Verschoor A, Schiemann
M, Hofer T, Busch DH (2013) Disparate individual fates compose robust CD8+ T cell immunity.
Science 340: 630-635
Burnham KP, Anderson DR, Huyvaert KP (2011) AIC model selection and multimodel inference
in behavioral ecology: some background, observations, and comparisons. *Behavioral ecology*
and sociobiology 65: 23-35
Caligiuri MA (2008) Human natural killer cells. *Blood* 112: 461-469
Cerwenka A, Lanier LL (2016) Natural killer cell memory in infection, inflammation and cancer.
Nat Rev Immunol 16: 112-123
Chang JT, Palanivel VR, Kinjyo I, Schambach F, Intlekofer AM, Banerjee A, Longworth SA, Vinup
KE, Mrass P, Oliaro J (2007) Asymmetric T lymphocyte division in the initiation of adaptive
immune responses. *science* 315: 1687-1691
Cheon H, Kan A, Prevedello G, Oostindie SC, Dovedi SJ, Hawkins ED, Marchingo JM, Heinzel S,
Duffy KR, Hodgkin PD (2021) Cyton2: A model of immune cell population dynamics that includes
familial instructional inheritance. *Frontiers in Bioinformatics* 1: 723337
Chiossone L, Chaix J, Fuseri N, Roth C, Vivier E, Walzer T (2009) Maturation of mouse NK cells is
a 4-stage developmental program. *Blood* 113: 5488-5496
Chu HH, Chan S-W, Gosling JP, Blanchard N, Tsitsiklis A, Lythe G, Shastri N, Molina-París C,
Robey EA (2016) Continuous effector CD8+ T cell production in a controlled persistent infection
is sustained by a proliferative intermediate population. *Immunity* 45: 159-171
De Boer RJ, Yates AJ (2023) Modeling T cell fate. *Annual review of immunology* 41: 513-532
DeGottardi MQ, Okoye AA, Vaidya M, Talla A, Konfe AL, Reyes MD, Clock JA, Duell DM, Legasse
AW, Sabnis A *et al* (2016) Effect of Anti-IL-15 Administration on T Cell and NK Cell Homeostasis
in Rhesus Macaques. *J Immunol* 197: 1183-1198
Fauriat C, Long EO, Ljunggren HG, Bryceson YT (2010) Regulation of human NK-cell cytokine and
chemokine production by target cell recognition. *Blood* 115: 2167-2176
Flommersfeld S, Bottcher JP, Ersching J, Flossdorf M, Meiser P, Pachmayr LO, Leube J, Hensel I,
Jarosch S, Zhang Q *et al* (2021) Fate mapping of single NK cells identifies a type 1 innate
lymphoid-like lineage that bridges innate and adaptive recognition of viral infection. *Immunity*
54: 2288-2304 e2287
Gerlach C, Rohr JC, Perié L, Van Rooij N, Van Heijst JW, Velds A, Urbanus J, Naik SH, Jacobs H,
Beltman JB (2013) Heterogeneous differentiation patterns of individual CD8+ T cells. *Science*
340: 635-639
Gillespie DT (1977) Exact Stochastic Simulation of Coupled Chemical Reactions. *The Journal of*
Physical Chemistry 81: 2340-2361

Grassmann S, Pachmayr LO, Leube J, Mihatsch L, Andrae I, Flommersfeld S, Oduro J, Cicin-Sain
L, Schiemann M, Flossdorf M *et al* (2019) Distinct Surface Expression of Activating Receptor
Ly49H Drives Differential Expansion of NK Cell Clones upon Murine Cytomegalovirus Infection.
Immunity 50: 1391-1400 e1394
Guma M, Angulo A, Vilches C, Gomez-Lozano N, Malats N, Lopez-Botet M (2004) Imprint of
human cytomegalovirus infection on the NK cell receptor repertoire. *Blood* 104: 3664-3671
Hawkins ED, Turner ML, Dowling MR, Van Gend C, Hodgkin PD (2007) A model of immune
regulation as a consequence of randomized lymphocyte division and death times. *Proceedings*
of the National Academy of Sciences 104: 5032-5037
Hayakawa Y, Smyth MJ (2006) CD27 dissects mature NK cells into two subsets with distinct
responsiveness and migratory capacity. *J Immunol* 176: 1517-1524
Heinzel S, Binh Giang T, Kan A, Marchingo JM, Lye BK, Corcoran LM, Hodgkin PD (2017) A Myc-
dependent division timer complements a cell-death timer to regulate T cell and B cell
responses. *Nature immunology* 18: 96-103
Hendricks DW, Balfour HH, Jr., Dunmire SK, Schmeling DO, Hogquist KA, Lanier LL (2014) Cutting
edge: NKG2C(hi)CD57+ NK cells respond specifically to acute infection with cytomegalovirus
and not Epstein-Barr virus. *J Immunol* 192: 4492-4496
Huntington ND, Vosshenrich CA, Di Santo JP (2007) Developmental pathways that generate
natural-killer-cell diversity in mice and humans. *Nat Rev Immunol* 7: 703-714
Ishiyama K, Arakawa-Hoyt J, Aguilar OA, Damm I, Towfighi P, Sigdel T, Tamaki S, Babdor J,
Spitzer MH, Reed EF *et al* (2022) Mass cytometry reveals single-cell kinetics of cytotoxic
lymphocyte evolution in CMV-infected renal transplant patients. *Proc Natl Acad Sci U S A* 119
Iwata A, Durai V, Tussiwand R, Briseño CG, Wu X, Grajales-Reyes GE, Egawa T, Murphy TL,
Murphy KM (2017) Quality of TCR signaling determined by differential affinities of enhancers
for the composite BATF–IRF4 transcription factor complex. *Nature immunology* 18: 563-572
Kaech SM, Wherry EJ, Ahmed R (2002) Effector and memory T-cell differentiation: implications
for vaccine development. *Nat Rev Immunol* 2: 251-262
Kalia V, Sarkar S (2018) Regulation of Effector and Memory CD8 T Cell Differentiation by IL-2-A
Balancing Act. *Front Immunol* 9: 2987
Kamimura Y, Lanier LL (2015) Homeostatic control of memory cell progenitors in the natural
killer cell lineage. *Cell Rep* 10: 280-291
Kim KH, Yu HT, Hwang I, Park S, Park SH, Kim S, Shin EC (2019) Phenotypic and Functional
Analysis of Human NK Cell Subpopulations According to the Expression of FcepsilonRIgamma
and NKG2C. *Front Immunol* 10: 2865
Kretschmer L, Flossdorf M, Mir J, Cho YL, Plambeck M, Treise I, Toska A, Heinzel S, Schiemann
M, Busch DH *et al* (2020) Differential expansion of T central memory precursor and effector
subsets is regulated by division speed. *Nat Commun* 11: 113
Lau CM, Adams NM, Geary CD, Weizman OE, Rapp M, Pritykin Y, Leslie CS, Sun JC (2018)
Epigenetic control of innate and adaptive immune memory. *Nat Immunol* 19: 963-972
Lopez-Verges S, Milush JM, Schwartz BS, Pando MJ, Jarjoura J, York VA, Houchins JP, Miller S,
Kang SM, Norris PJ *et al* (2011) Expansion of a unique CD57(+)NKG2Chi natural killer cell subset
during acute human cytomegalovirus infection. *Proc Natl Acad Sci U S A* 108: 14725-14732

Martinet L, Ferrari De Andrade L, Guillerey C, Lee JS, Liu J, Souza-Fonseca-Guimaraes F,
Hutchinson DS, Kolesnik TB, Nicholson SE, Huntington ND *et al* (2015) DNAM-1 expression
marks an alternative program of NK cell maturation. *Cell Rep* 11: 85-97
Min-Oo G, Bezman NA, Madera S, Sun JC, Lanier LL (2014) Proapoptotic Bim regulates antigen-
specific NK cell contraction and the generation of the memory NK cell pool after
cytomegalovirus infection. *J Exp Med* 211: 1289-1296
Mitrovic M, Arapovic J, Jordan S, Fodil-Cornu N, Ebert S, Vidal SM, Krmpotic A, Reddehase MJ,
Jonjic S (2012) The NK cell response to mouse cytomegalovirus infection affects the level and
kinetics of the early CD8(+) T-cell response. *J Virol* 86: 2165-2175
Nabekura T, Kanaya M, Shibuya A, Fu G, Gascoigne NR, Lanier LL (2014) Costimulatory molecule
DNAM-1 is essential for optimal differentiation of memory natural killer cells during mouse
cytomegalovirus infection. *Immunity* 40: 225-234
Ni J, Miller M, Stojanovic A, Garbi N, Cerwenka A (2012) Sustained effector function of IL-
12/15/18-preactivated NK cells against established tumors. *J Exp Med* 209: 2351-2365
Orr MT, Sun JC, Hesslein DG, Arase H, Phillips JH, Takai T, Lanier LL (2009) Ly49H signaling
through DAP10 is essential for optimal natural killer cell responses to mouse cytomegalovirus
infection. *The Journal of experimental medicine* 206: 807-817
Pandit A, De Boer RJ (2019) Stochastic inheritance of division and death times determines the
size and phenotype of CD8+ T cell families. *Frontiers in immunology* 10: 436
Paul S, Kulkarni N, Shilpi, Lal G (2016) Intratumoral natural killer cells show reduced effector
and cytolytic properties and control the differentiation of effector Th1 cells. *Oncoimmunology*
5: e1235106
Peng H, Tian Z (2017) Natural Killer Cell Memory: Progress and Implications. *Front Immunol* 8:
1143
Potempa M, Aguilar OA, Gonzalez-Hinojosa MDR, Tenvooren I, Marquez DM, Spitzer MH, Lanier
LL (2022) Influence of Self-MHC Class I Recognition on the Dynamics of NK Cell Responses to
Cytomegalovirus Infection. *J Immunol* 208: 1742-1754
Riggan L, Ma F, Li JH, Fernandez E, Nathanson DA, Pellegrini M, O'Sullivan TE (2022a) The
transcription factor Fli1 restricts the formation of memory precursor NK cells during viral
infection. *Nat Immunol* 23: 556-567
Riggan L, Ma F, Li JH, Fernandez E, Nathanson DA, Pellegrini M, O'Sullivan TE (2022b) The
transcription factor Fli1 restricts the formation of memory precursor NK cells during viral
infection. *Nature immunology* 23: 556-567
Robbins SH, Tessmer MS, Mikayama T, Brossay L (2004) Expansion and contraction of the NK
cell compartment in response to murine cytomegalovirus infection. *J Immunol* 173: 259-266
Romee R, Rosario M, Berrien-Elliott MM, Wagner JA, Jewell BA, Schappe T, Leong JW, Abdel-
Latif S, Schneider SE, Willey S *et al* (2016) Cytokine-induced memory-like natural killer cells
exhibit enhanced responses against myeloid leukemia. *Sci Transl Med* 8: 357ra123
Rückert T, Romagnani C (2024) Extrinsic and intrinsic drivers of natural killer cell clonality.
Immunological reviews 323: 80-106
Santosa EK, Kim H, Rückert T, Le Luduec J-B, Abbasi AJ, Wingert CK, Peters L, Frost JN, Hsu KC,
Romagnani C (2023) Control of nutrient uptake by IRF4 orchestrates innate immune memory.
Nature immunology 24: 1685-1697

Schlub TE, Badovinac VP, Sabel JT, Harty JT, Davenport MP (2010) Predicting CD62L expression
during the CD8+ T-cell response in vivo. *Immunol Cell Biol* 88: 157-164
Schlub TE, Sun JC, Walton SM, Robbins SH, Pinto AK, Munks MW, Hill AB, Brossay L, Oxenius A,
Davenport MP (2011) Comparing the kinetics of NK cells, CD4, and CD8 T cells in murine
cytomegalovirus infection. *J Immunol* 187: 1385-1392
Sun JC, Beilke JN, Lanier LL (2009) Adaptive immune features of natural killer cells. *Nature* 457:
557-561
Vivier E, Tomasello E, Baratin M, Walzer T, Ugolini S (2008) Functions of natural killer cells. *Nat
Immunol* 9: 503-510
Voigt V, Forbes CA, Tonkin JN, Degli-Esposti MA, Smith HR, Yokoyama WM, Scalzo AA (2003)
Murine cytomegalovirus m157 mutation and variation leads to immune evasion of natural killer
cells. *Proc Natl Acad Sci U S A* 100: 13483-13488
Yang S, Liu F, Wang QJ, Rosenberg SA, Morgan RA (2011) The shedding of CD62L (L-selectin)
regulates the acquisition of lytic activity in human tumor reactive T lymphocytes. *PLoS One* 6:
e22560
945