Platelet extracellular vesicles and their mitochondrial content significantly improve survival and cytokine levels when transfused into mice with sepsis or H1N1 infections

Tom Benson[1], Samir P. Patel[3], Benedict C. Albensi[456], Vinit B. Mahajan[78], Aida Adlimoghaddam[9,10], and Hiroshi Saito,[2,3]

[1] Mitrix Bio Inc., Pleasanton, California, United States

[2] Department of Surgery, University of Kentucky, Lexington, Kentucky, USA

[3] Department of Physiology University of Kentucky, Lexington, Kentucky, USA

[4] Division of Neurodegenerative Disorders, St. Boniface Hospital Albrechtsen Research Centre, Winnipeg, Canada

[5] Department of Pharmacology & Therapeutics, Max Rady College of Medicine, University of Manitoba, Winnipeg, Canada

[6] Department of Pharmaceutical Sciences, College of Pharmacy, Nova Southeastern University, Fort Lauderdale, FL, USA

[7] Molecular Surgery Laboratory, Stanford University, Palo Alto, CA, USA

[8] Department of Ophthalmology, Byers Eye Institute, Stanford University, Palo Alto, CA, USA

[9] Department of Neurology, Neuroscience Institute, Center for Alzheimer's Disease and Related Disorders, Southern Illinois University School of Medicine, Springfield, IL, USA

[10] Division of Neurodegenerative Disorders, St. Boniface Hospital Research, Winnipeg, MB, Canada.

Correspondence
Tom Benson, CEO
Mitrix Bio Inc.
3695 Chabot Dr. Suite 200
Pleasanton, CA 94566
Tom@mitrix.bio
 -or-
Hiroshi Saito, PhD
Department of Surgery, University of Kentucky
800 Rose Street, MS-476 Medical Science Building,
Lexington, KY 40536-0298.
E-mail: hiroshi.saito@uky.edu

Keywords: mitochondria, circulating mitochondria intercellular communication mitochondrial transplantation, immune senescence

First published online: September 7, 2023

ABSTRACT

BACKGROUND: Mitochondrial transplantation is an emerging technique focused until now on ischemia-reperfusion injuries and rare mitochondrial mutation diseases. Platelets are the most common immune cells in the blood and emit mitochondria inside extracellular vesicles ("mitlets") during activation or at their end-of-life. Mitlets are preferentially absorbed by immune cells in the blood and increase cellular energetics. In this study, we test the hypothesis that transplanting "young" mitlets can reverse age-related immune senescence in animals.

STUDY DESIGN AND METHODS: We transplanted mitlets from a young, healthy murine animal model into both young and old animals which were infected with H1N1 influenza virus or polymicrobial bacteria to induce sepsis.

RESULTS: Transplantation of mitlets significantly improved survival, reduced bacterial load, and reduced cytokine levels of the recipients. As expected, this effect was temporally limited, lasting a few days after cessation of treatment, and was more impactful for aged animals.

DISCUSSION: In a first-ever result, we show that transplants of young mitochondria in mitlets appear to reverse key age-related factors of the immune system. This suggests that immune senescence is primarily a function of reduced mitochondrial energy production and thus can be reversed. This approach has potential as a novel immunotherapy for infectious disease.

Note, this paper is a companion to: *Pelletier et al, Platelet extracellular vesicles and their mitochondrial content improve the mitochondrial bioenergetics of cellular immune recipients.*[66]

1 INTRODUCTION

Mitochondria produce the majority of cellular energy and, therefore, the energy for immune system function. Mitochondrial energy production declines with age, and this decline is more pronounced in primates and humans compared to mice. For example, the energy decline in retinal mitochondria of 14-year-old)\ primates is 70%, while the same decline measured in 12-month-old mice is 30%.[85] Recent work has found that mitochondria are frequently transferred between cells via stem cells and extracellular vesicles[21], and researchers have tested many techniques for effecting such transfers from an exogenous source – isolating and transplanting the mitochondria from cell to cell or from animal to animal.[72,87,88] Platelets are the most common form of immune system blood cell.[27] Mitochondria can be found in great numbers in the bloodstream, emitted inside extracellular vesicles by platelets during activation or at end-of-life.[14] Research has found that such platelet-derived mitochondrial vesicles ("mitlets") are quickly and preferentially absorbed by other immune cells in the bloodstream, including white blood cells and other platelets.[11]

Mitochondrial transplantation is an emerging technique in which exogenous mitochondria, isolated from either the patient or a donor or grown in external cell cultures, are transplanted into organs to restore cellular energetics and cure injuries. This technique until now has been tested primarily on ischemia-reperfusion injuries, organ transplants, and rare mitochondrial mutation diseases but has not been explored for adult age-related or chronic diseases.

Historically, the elderly are more prone to death and disability from infection. The age-related mechanisms that reduce the effectiveness of the immune system are not fully understood. We hypothesized that the decline of the human immune system, and thus the increased levels of mortality and morbidity in the elderly due to infectious disease, is primarily due to reduced energy production by mitochondria inside white blood cells and immune cells. The functions of white blood cells are energy-intensive, particularly during infection. If they suffer reduced energy production due to age, then it follows that they will be unable to perform their functions effectively during a serious infection. Aged white blood cells move slower, act slower, and display a reduced ability to digest or kill pathogens compared to younger white blood cells.[1]

We tested this hypothesis by transplanting mitlets containing mitochondria. We isolated the mitlets from the blood of young animals and then transplanted them into aged animals with induced viral or bacterial infections. We performed similar transplants for mitochondria that were isolated from mouse liver, not encased in vesicles. We then ascertained survival, blood pathogen load, and cytokine levels for these mitochondrial types and a variety of delivery mechanisms.

2 METHODS

2.1 Facilities involved. Experiments below were accomplished at several locations. Charles River Labs in Bristol, England performed experiments using mice with an H1N1 influenza virus. The University of Kentucky, USA performed experiments using mice with an induced sepsis infection. Additional mitlet preparation and testing was performed by AA at St. Boniface Hospital, MB, Canada.

2.2 Ethics. All experimental procedures at the University of Kentucky were approved by the Institutional Animal Care and Use Committee. Animal handling techniques in these studies were performed as described and approved in our Animal Use Protocol #2009-0541 and were in accordance with the National Institutes of Health guidelines for ethical treatment. All experimental procedures at Charles River Labs in Bristol, England, were approved by the Institutional Animal Care and Use Committee. Animal handling techniques in these studies were performed as described following commonly-accepted guidelines for ethical treatment.

2.3 Mice for sepsis experiments, University of Kentucky. For these experiments, middle-aged (13-month-old) female C57BL/6 mice with jugular vein catheterization were obtained from Charles River Laboratories, Watsonville, CA. A Vascular Access Button (VAB, Intech Laboratories, Inc) was attached to the subcutaneous end of the catheter, allowing exogenous fluids infusion repeatedly. Animals were maintained in the Division of Laboratory Animal Resources at the University of Kentucky in pressurized intra-ventilated cages housed five animals per cage under controlled temperature (21 – 23°C), humidity (30% – 70%), and lighting (14/10 light/dark cycle) with free access to drinking water and chow (Teklad Global No. 2918, 18% Protein Rodent Diet, Madison, Wis). All animals were acclimated for at least 7 days prior to experimentation to eliminate the influence of transportation stress.

2.4 Induction of sepsis in mice at University of Kentucky. Experimental polymicrobial abdominal sepsis was induced by bolus injection of cecal slurry (CS) as we previously described in Steel et al.[66.] Survival and the health of each mouse was monitored multiple times daily for 14 days. A small amount of blood samples was collected aseptically from the tail vein of each mouse 12 hours after sepsis induction. Each blood sample (10-µL) was quickly diluted with 9-folds volume (90-µL) of 1x citrate solution (0.32% sodium citrate in 0.9% sodium chloride) and plated onto an agar plate as we described (Steel et al [66]). Anaerobic bacteria colonies were counted after incubating these agar plates for 2 days using the GasPak anaerobic gas generating pouch system (BD Biosciences).

2.6 Mice for influenza virus experiments, Charles River Labs, Bristol, England. Adult female BALB/c mice from internal stock were randomly allocated and allowed to acclimatize for one week.

2.7 Induction of H1N1 virus in mice at Charles River Labs, Bristol, England. On Day 0, animals were challenged with A/Puerto/Rico/8 (H1N1)(PR8) and monitored daily for clinical signs of ill-health until termination on Day 7. In-life bleeds were taken on Day -1 (pre-treatment) and Day 3 (post-treatment) and processed to plasma for Luminex analysis of cytokine levels. At termination, bronchoalveolar lavage (BAL) and blood was taken. Blood was processed to plasma and serum. BAL fluid and plasma samples were analyzed for cytokine levels using Luminex and serum was analyzed in a HAI assay to indicate the levels of functional anti-viral antibodies present.

2.8 Platelet isolation and preparation of platelet extracellular vesicles (mitlets) at University of Kentucky for sepsis experiments. We used the same approach described by Marcoux et al. to isolate platelets and mitlets[32]. In brief, C57BL/6 2-month-old mouse blood preserved with ACD 20% on ice was purchased from BioIVT Inc., Westbury, NY. Platelets were isolated from blood, and then washed with Tyrode's Buffer pH 7.4 and resuspended at 1×10^8 cells/mL in Tyrode's Buffer pH 7.4 with 5 mM $CaCl_2$. Platelets were stimulated overnight (16h) at room temperature with thrombin. 10 mM EDTA was added to stop the stimulation, then stimulated platelets were centrifuged at 300 g for 5 min to remove remnant platelets or cells. The resulting mix is concentrated by ultracentrifugation at 18,000 g for 60 min at 18°C, which isolates approximately 98% of the mitlets according to quantifications of protein contents[3.2] The mitlet pellet was resuspended in filtered phosphate-buffered saline (PBS) pH 7.4. The resulting mitlets in PBS were chilled to 4 degrees C for up to 1 week prior to administration.

2.9 Platelet isolation and preparation of platelet extracellular vesicles (mitlets) performed at St. Boniface Hospital in support of Bristol H1N1 virus experiments. We used the same approach described by Marcoux et al. to isolate platelets and mitlets[32]. In brief, 2-month-old male or female BALB/C mouse blood preserved with ACD 20% at room temperature was purchased from BioIVT Inc., Westbury, NY. Platelets were isolated from blood, and then washed with Tyrode's Buffer pH 7.4 and resuspended at 1×10^8 cells/mL in Tyrode's Buffer pH 7.4 with 5 mM $CaCl_2$. Platelets were stimulated overnight (16h) at room temperature with thrombin. 10 mM EDTA was added to stop the stimulation, then stimulated platelets were centrifuged at 300 g for 5 min to remove remnant platelets or cells. The resulting mix is concentrated by ultracentrifugation at 18,000 g for 60 min at 18°C, which isolates approximately 98% of the mitlets according to quantifications of protein contents[3.2] The mitlet pellet was resuspended in filtered phosphate-buffered saline

(PBS) pH 7.4. 50% of the resulting mitlet/IPBS mixture was mixed with an additional cryopreservative and chilled to 4 degrees C. The other 50% of the mitlets were left in PBS and chilled to 4 degrees C. All chilled mitlets were shipped on ice by medical courier to the Charles River Labs, Bristol, England location.

2.10 Liver mitochondrial preparation for sepsis tests at University of Kentucky. Mitochondria were isolated from young (2-month-old) healthy C57BL/6 mouse liver from internal stock, as described before[32]. In summary, naive anesthetized mice under isoflurane were euthanized by cervical dislocation, and livers were removed and placed in ice cold homogenization buffer. Livers were transferred into a Dounce tissue homogenizer along with 5 ml ice cold homogenizing buffer (300 mM sucrose, 10 mM K-HEPES, and 1 mM K-EGTA (pH 7.2)). The tissue was gently homogenized for 60 seconds. 250µL of Subtilisin A (96.61µM) was added to the homogenate and mixed by inversion. This was then incubated on ice for 10 minutes. Using a pre-wet 40µm filter, filter the homogenate into a 50mL falcon on ice. This step was repeated with a 40µm filter and then a 10µm filter. The filtrate was then transferred to 1.5mL Eppendorfs and centrifuged at 9,000 xg for 10 minutes at 4°C. The supernatant was discarded, and the pellet was resuspended in 1mL ice-cold PBS. The concentrated filtrate was then filtered through a 1.2µm and 0.8µm filter. A protein assay was performed to confirm the concentration of the mitochondria. This was adjusted to 10mg/kg and handed to the in vivo team on ice for administration.

2.11 Liver mitochondrial preparation for H1N1 Viral Tests at CRL Bristol, England. Mitochondria were isolated from young (2-month-old) healthy C57BL/6 mouse liver from internal stock, as described before[32]. In summary, naive anesthetized mice under isoflurane were euthanized by cervical dislocation, and livers were removed and placed in ice cold homogenization buffer. Livers were transferred into a Dounce tissue homogenizer along with 5 ml ice cold homogenizing buffer (300 mM sucrose, 10 mM K-HEPES, and 1 mM K-EGTA (pH 7.2)). The tissue was gently homogenized for 60 seconds. 250µL of Subtilisin A (96.61µM) was added to the homogenate and mixed by inversion. This was then incubated on ice for 10 minutes. Using a pre-wet 40µm filter, filter the homogenate into a 50mL falcon on ice. This step was repeated with a 40µm filter and then a 10µm filter. The filtrate was then transferred to 1.5mL Eppendorfs and centrifuged at 9,000 xg for 10 minutes at 4°C. The supernatant was discarded, and the pellet was resuspended in 1mL ice-cold PBS. The concentrated filtrate was then filtered through a 1.2µm and 0.8µm filter. A protein assay was performed to confirm the concentration of the mitochondria. This was adjusted to 10mg/kg and handed to the in vivo team on ice for administration.

2.12 Statistical analysis and software at University of Kentucky. Survival curves were analyzed by Kaplan–Meier LogRank test. Data for two-group comparisons were analyzed by Student t test. When multiple comparisons were made, the Shapiro–Wilk normally test was run. If the data passed the normality test, one-way ANOVA and Holm–Sidak post-hoc test were used to analyze the data. Alternatively, when the data was not normally distributed, the Kruskal–Wallis test and Dunn post-hoc test were used. In instances where one group was assessed multiple times (i.e., bacteria load), repeated-measures one-way ANOVA was used, and when multiple groups were assessed multiple times (i.e., body temperature data), repeated- measures two-way ANOVA was used and the Holm–Sidak post-hoc test was run. All data are expressed as means and standard deviations and p <0.05 was considered statistically significant.

2.13 Statistical analysis and software at Charles River Labs, Bristol, England. Samples were analyzed individually at a 1 in 4 dilution. Each dot represents the group mean + SEM at each time point. The functional Lower Limit of Quantification (LLOQ) (3.34 pg/mL) is plotted as a dotted black line. Values <LLOQ are plotted as 0 pg/mL. Statistical significance analysis was attempted using a Mixed-effects model, however due to values <LLOQ a p value could not be calculated. Normality was determined by a Shapiro-Wilk test. Statistical significance between the Naïve (uninfected) and Vehicle group was determined by a Mann-Whitney T test. Statistical significance between the vehicle, oseltamivir, Mitlets and Liver mitochondria-treated groups was determined using a Kruskal-Wallis test followed by Dunn's multiple comparison test, using the vehicle group as the comparator. No differences between the groups in any tests were found to be statistically significant.

3 RESULTS

Platelet extracellular vesicles (mitlets) were isolated from 2-month-old mice and transplanted into same-age mice with H1N1 influenza infection, showing a trend by slightly improving survival and reducing cytokines, but not with statistical significance. (Charles River Labs, Bristol, England location). Liver isolated mitochondria had no noticeable impact. Adult female BALB/c mice were randomly allocated and allowed to acclimatize for one week. Each immune-assay readout was performed individually. Readouts were: Bodyweights, Clinical Scores, Survival, Cytokine levels measured using Luminex: 5-plex testing IL-6, IL-10, IL-17, IFN-ɣ and TNFα Haemagglutination inhibition assay (HAI). Compounds were administered by tail-vein injection, according to the table below:

Groups	N=	Treatments			Intervention
		Dose	Route	Regimen	
Naïve	5	N/A	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	N/A
Vehicle control	5	N/A	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	Day 0: IN admin of PR8 H1N1 influenza virus
Oseltamivir	5	20mg/kg	PO	SID from Day -1 to end	
Mitlets (in PBS)	5	N/A	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	
Mitlets (in Cryopreservative)	5	N/A	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	
Liver Mitochondria	10	10mg/kg	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	

None of the results from the Bristol experiments showed statistical significance and are scheduled to be repeated with a larger number of mice and higher doses.

Platelet extracellular vesicles (mitlets) isolated from 2-month-old mice transplanted into 13-month-old mice with sepsis significantly improved survival, reduced bacterial count, and reduced cytokines. (University of Kentucky location). Liver-isolated mitochondria had no noticeable impact. 13-month-old C57BL/6 mice were surgically modified by the vendor to add a VAB (injection port) in the jugular vein for easy injection. Mice were allowed to acclimate in cages for 1 week after receipt, then infected with Sepsis per long-established protocol[66]. Mice were injected according to the following table:

Groups	N=	Treatments			Intervention
		Dose	Route	Regimen	
Vehicle control with mitlets	8	N/A	IV, 120 μL	Day -1, Day , Day 2 & Day 3 (x4)	Day 0: induction of Sepsis via Cecal Slurry injection
Mitlets (in PBS)	8	N/A	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	
Vehicle control with liver mitochondria	3	10mg/kg	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	
Liver mitochondria	5	10mg/kg	IV, 120 μL	Day -1, Day 1 and Day 3 (x3)	

They were injected at -1, 1, 2, and 3 days. The bacterial count and cytokines were collected at -2, 1, and 3 days. The study was duplicated several months later with identical results. An additional 3 treatments on Days 3, 4, and 5 were performed or Exp 2. In the first experiment, 2 mice died during injection, probably because of poor handling, and were removed from the experiment.

Mitlet treatments significantly prevent early death of mice one week after sepsis induction (P<0.01). Mitlet treatments (by Day 2 - 5) delayed but did not prevent later death. Mitlet treatment significantly reduced blood bacteria several folds but didn't eliminate it completely, and reduced IL-6 cytokine level ~ 60%. The effect of mitlet treatment was temporary and declined a few days after the last injection. In contrast to mitlets, injections of mitochondria isolated from the liver had no protective effect – mice in these cases died at the same time as negative controls.

4 DISCUSSION

In our study, we transplanted mitochondria at two different locations using two separate teams, at Charles River Labs in Bristol, England (Bristol) and the lab of Hiroshi Saito at the University of Kentucky (Kentucky). At Bristol, we transplanted same-age mitochondria for BALB/C mice, 2-3 months old. Recipient mice were dosed with an H1N1 flu virus. We used same-age donor mitochondria of three types: mitochondria isolated directly from the livers of donor mice and mitlets isolated from the blood of BALB/C mice in another lab in Winnipeg, Canada, either preserved in a cryopreservative solution or PBS, chilled to 4° C, and shipped by courier to Bristol. All three of these were injected into recipient mice three times over seven days. We did not expect to see a beneficial effect in the Bristol experiments because the mitochondria were from animals of the same age; however, there was a small noticeable effect, albeit not enough to reach statistical significance. For the Kentucky tests, we injected mitochondrial test products from 2-month-old C57BL/6 mice into aged 13-month-old C57BL/6 mice, which had been given sepsis via cecal slurry induction. The first injections were mitochondria isolated from the liver of the same-aged mice and injected immediately, which showed no results. The second injections were mitlets isolated from the blood of donor animals, chilled in PBS to 4° C, and then injected three different times over the course of the experiment, which showed highly significant improvements ($p < 0.05$). We duplicated the successful mitlet experiments later with the same results.

These experiments, taken together, correspond accurately to our predictions and to the in-vitro work done in parallel at our partner lab at the University of Laval Quebec. [68,32] We expected to see that mitochondria transplanted from younger to older mice would create a much more profound beneficial effect. We expected that mitlets would provide a robust response while liver-isolated mitochondria would produce little or no response because mitlets are preferentially targeted to immune cells, while liver mitochondria, with no vesicle and no receptors, become scattered into different tissues.

The benefits of mitlets are more pronounced in aged mice. Aged mice have mitochondria that are more dysfunctional compared to young mice. This has been shown in prior studies that looked at the energy production level of mitochondria in aged animals versus young animals.[85,86] According to the "Mitochondrial Theory of Aging[83]," mitochondria start as highly functional at birth, but their functionality and performance decline steadily over the years, causing an inexorable decline in the ATP energy produced in each cell. As a result, the total energy available to a given organ or sub-system of the body, in this case the immune system, declines with age.[84] A similar decline can also be seen in children with mitochondrial mutation diseases such as Leigh syndrome or Kearns-Sayre syndrome. They are born with mutated and dysfunctional mitochondria from birth and are subject to a large variety of morbidity and mortality due to insufficient cellular energetics.[63,64] These children are most affected when the ratio of dysfunctional mitochondria exceeds a certain threshold, generally 60-70%.[65] Thus, we see that mitochondrial energy production is a measurable and predictable number, which can be seen to have predictable consequences for the human body. In a similar way, mitochondrial energy production declines due to age, in some cases 70% in the elderly primate retinas.[85] We and other researchers hypothesize that these age-related energy reductions lead to cellular failure, lower activity, less performance, accumulation of waste products, and other effects that result in a large part of biological aging. In this work, we tested the ability of white blood cells to combat infection, which is reduced in the 13-month-old mice used in Kentucky. We injected mitlets from young animals, which are absorbed into the white blood cells and used as supplemental energy sources to support the increased energy demand associated with their antimicrobial functions, such as phagocytosis, degranulation, and NETosis. Thus, we are, in essence, making these white blood cells temporarily younger, or we may instead say, we are making them temporarily *act* younger.

Lesser results for same-age animals. We weren't sure if we would see measurable results for same-age animals, because there is no difference in the activity level and the energy production of mitochondria in the donor mitochondria versus the wild-type mitochondria in the white blood cells of the recipient animals. It was interesting that the transfusion of mitlets produced a modest effect. We interpret this to mean that the recipient white blood cells, although they received mitochondria of approximately the same functionality as those they already had, nevertheless benefited by getting additional mitochondria. According to our *in-vivo* studies[68], white blood cells have a certain "starting complement" of existing mitochondria. In the middle of an infection, the white blood cells might have the capacity to accept additional mitochondria – again, corresponding to our earlier analogy of mitochondria as a type of spare battery pack. We helped the white blood cells not by giving them *higher-quality* mitochondria, but simply by giving them *more* mitochondria.

Immune system increases the volume of mitochondria in response to infection. The body increases and decreases the number of white blood cells in circulation in response to infection. Chiu et al. showed that platelets produced in the bone marrow are supplied with more mitochondria, averaging 7.26 mitochondria per platelet when suffering from cancer verses 4.85 when healthy[70]. This implies that the body is actively increasing or decreasing the number of mitochondria in the immune system in response to infection. More numerous mitochondria in platelets yield more mitlets in circulation, which then transfer more mitochondria to the white blood cells to fight off infection. Thus,

the body is systematically supplementing the number of mitochondria in white blood cells to match the demands of a given infection. Presumably, although this has not been described, one or more hormones or cytokines up-regulates the mitochondrial quantity per platelet.

Mitochondria in white blood cells "burned out" by the stress of infection. According to Picard[71], stressful conditions cause increased energy demands and mitochondrial DNA degeneration. Mitochondria serve many different functions, not only energy production but intracellular communication, inflammation, triggering apoptosis, and much more.[71] Our findings show that immune cells quickly and preferentially absorb mitlets for extra mitochondria needed to support the system's function, especially when fighting infection. This theory might be seen as analogous to a system of supply trucks providing soldiers in battle with more ammunition, supplies, and fuel, increasing these deliveries as needed to match the severity of the battle. Platelets serve many functions in the immune system; now, we show that they also deliver energy supplies to other cells.

Mitlets yield measurable changes, whereas mitochondria isolated from the liver do not. Results confirm that the mitlets are much more effective than mitochondria isolated from the liver. We propose that mitlets are pre-designed and engineered by evolution to perform these duties. They are encased in extracellular vesicles containing various receptor patterns, causing them to be preferentially targeted to and quickly accepted by the white blood cells floating nearby in the blood.[11] Mitochondria isolated from livers have no such vesicle and receptor coating, are not quickly absorbed by nearby immune cells, randomly flowing through the bloodstream to eventually become absorbed by other tissues such as lung, liver, or brain.[77,78]

Many types of mitlets in various tissues. Other studies have shown that the immune system, in fact, has many different varieties of mitlets with different receptor patterns on the surface.[73] Therefore, we might say that the immune system has a tool kit of different mitochondrial-bearing mitlets. These mitlets have also been shown to decline with age.[73, 76] The brain also has neural-specific mitlets that transfer mitochondria between astrocytes and neurons[16,17] and a similar system to serve cardiac cells.[75] We note that cardiac cells, neurons, and white blood cells are all high-priority, mission-critical cells for the body. In case of fight-or-flight, escape from predators, or infections from constant injuries, the energy production efficiency of the mitochondria in these cells are critical to the organism's survival. It makes logical sense that these cells have evolved to offload lower-priority mitochondrial production and maintenance activities to other support cells nearby, in order to conserve all available energies for survival. This would provide a significant competitive advantage. Thus, we may logically identify mitlets as a pervasive, evolutionarily-conserved subsystem of the body, which distribute

scarce energy resources between cells and tissues in response to environmental conditions. Similar mitochondrial transfer occurs via stem cells and directly from cell-to-cell within tissues.[21] Researchers are only now beginning to uncover this extraordinarily sophisticated web of energetic distribution, which is the fundamental basis of complex life.[79]

The benefit of transplantation is temporary. From these results we see the mitlet transplant's survival benefits decline steadily after the initial three-day infusion. This is consistent with our understanding of white blood cells. First, many white blood cells have short life spans. Thus, the beneficial effect of transplanted mitochondria will slowly wear off over a few days as the white blood cells undergo apoptosis and are replaced by wild-type white blood cells lacking the transplanted mitochondria. Additionally, according to Picard's theory of mitochondrial damage due to stressful conditions[71], we would expect that the freshly- transplanted younger mitochondria might quickly age and decay as their DNA is damaged by stressful usage. In essence, the young mitochondria quickly become old, in the heat of battle. We aim to explore this process in future experiments by performing additional injections on days 5, 7, 9 to determine if a continuous infusion of fresh replacement mitlets will allow the animals to survive longer.

Potential for permanent regeneration of the immune system. It is interesting to note that the concept of exhaustion in immunotherapy is already a well-known issue in other types of immunotherapies, such as CAR-T or CAR-NK therapy.[82] Transplantation of mitochondria via mitlets may also potentially apply to other types of immune senescence in the thymus or bone marrow, which also absorb mitlets, although in less proportion to white blood cells.[32] For that reason, we hypothesize that with changes to the administration or dosing of mitlets, it might be possible to transplant large numbers of young mitochondria into solid immune tissues so that the young mitochondria become templates for mitochondrial replication and distribution to future generations of platelets and white blood cells. Thus, age-reversal of the immune system might potentially be permanent instead of temporary. This would contribute further to the health of the aged individual, potentially reducing inflammation or reducing the occurrence of cancer and other immune-mediated diseases of aging.

Limitations of the study

Although our Kentucky sepsis experiments showed significant improvements in survival and reductions in bacterial count, results from the Bristol H1N1 influenza experiments did not show statistical significance, although there was sufficient improvement to infer the effect, especially since it matched closely to the results from the sepsis experiments. The lack of significance in the H1N1 results is likely due to a low dosage of administration, the low number of animals, and, most

importantly, the fact that donor and recipient animals were the same age. As we noted earlier, same-age donations were not expected to show strong results. A replicate study may be conducted for the Bristol experiment, using a larger number of animals and higher doses, to achieve statistical significance. The Kentucky sepsis experiments did not include positive controls, such as an anti-bacterial/anti-inflammatory treatment which is often used in sepsis tests[66], nor did they include tests with same-age mice to provide context for the survival results, nor did they include dosing studies. These omissions were deemed acceptable in the short term since the tests are not intended to address sepsis treatments specifically but use Sepsis only as a convenient biomarker of immune system age. Nonetheless, these issues are planned for future experiments.

Prospects for future experimentation

Future tests should address safety, dosing, and improved controls. Tests in larger animals or humans will encounter many potential challenges, including increases in inflammatory mediators, conflicts between mitochondrial haplotypes or vesicle antigens, organ-specific challenges, effects from storage of mitochondria, auto-immune responses, high concentrations of calcium ions, routes of administration, and overall safety concerns. Assuming these challenges can be addressed, there seems no fundamental reason that these treatments could not be used in humans, for example, elderly patients with sepsis, influenza, COVID-19, some types of cancers, other infectious diseases, and soldiers with battlefield injuries or sepsis. Another potential target population would be children with mitochondrial mutation diseases such as Leigh Syndrome, who have weakened immune systems and can suffer very negative outcomes from common infections. These children might be given mitlet transplants to temporarily make their immune systems "normal" to fight infection.[63,64,65]

Blood banks, which already have platelet transfusions regularly, might isolate mitlets, refrigerate them, and use them as needed in exceptional cases. However, it seems unlikely that the number of platelets currently available would create enough mitlets to treat many patients. Just as donor livers or kidneys are in short supply, we might expect donor mitochondria to be chronically short as well. Mitochondrial transplantation requires large volumes; small numbers confer little advantage. Therefore, developing a bioreactor manufacturing process that could grow mitlets (mitochondria encased in extracellular vesicles or an equivalent coating, along with artificial receptors to target specific tissue types) would be more practical for mass-market distribution and global use. Mitrix Bio, the sponsor of this study, has an active program to build this mitochondrial bioreactor.[74]

Potential value to society

If this process is found to be safe and efficacious in larger animals and humans, then it would represent a new type of immunotherapy with broad application to human medicine and pandemic response. We would term this HISET therapy (Human Immune System Energetics Transplant). HISET therapy could potentially serve as a "general purpose" immune system booster, which could be administered in combination with other antiviral or anti-bacterial treatments. Many elderly patients with infectious diseases end up in the hospital or the Intensive Care Unit (ICU), both of which are high-cost and a significantly drain on national healthcare budgets. By boosting the immune system of elderly patients with mass-manufactured transplantation-quality mitochondria, even if such a treatment is temporary, we could help large numbers of patients avoid dangerous and expensive ICU stays. These treatments might also apply to chronic diseases and pandemic management. For example, studies have shown that COVID-19 can cause significant mitochondrial damage and might contribute to Long Covid.[80,81] Therefore, mitochondrial transplantation therapy has potential for chronic post-infection diseases such as Long Covid, Long Sepsis, or post-chemotherapy.

FUNDING INFORMATION

The study was sponsored by Mitrix Bio, Inc

CONTRIBUTIONS

HS, SP, AA, and SS designed and conducted the experiments. TB wrote the manuscript. BA and VM edited and consulted on the manuscript.

CONFLICT OF INTEREST STATEMENT

The study was sponsored by Mitrix Bio, Inc. HS, BA, VM, SP, and AA are scientific advisors for Mitrix Bio. TB is the CEO for Mitrix Bio.

ETHICS STATEMENT

All experimental procedures at the University of Kentucky were approved by the Institutional Animal Care and Use Committee. Animal handling techniques in these studies were performed as described and approved in our Animal Use Protocol #2009-0541 and were in accordance with the National Institutes of Health guidelines for ethical treatment. All experimental procedures at Charles River Labs in Bristol, England, were approved by the Institutional Animal Care and Use Committee.

Figure 1. Platelet extracellular vesicles (mitlets) isolated from young mice and transplanted into same-age recipient mice with H1N1 influenza infection, improved survival. Kaplan-Meier Survival Graphs. Some data points have been nudged for clarity. "Naïve control" received no H1N1 dosing. "Vehicle control" received the H1N1 dosing but was given no treatment. Others received the H1N1 dosing and were given mitlet or mitochondria as shown in legend. Improvement did not reach statistical significance.

Figure 2. Platelet extracellular vesicles (mitlets) isolated from young mice and transplanted into same-age recipient mice with H1N1 influenza infection, reduced cytokine levels (plasma IL-6), day -1 to day 7. Blood was collected from each animal and processed to plasma for cytokine analysis by Luminex (n=5-10 per group). Samples were analyzed in singlicate at a 1 in 4 dilution. Each dot represents the group mean + SEM at each time point. The functional Lower Limit of Quantification (LLOQ) (3.34 pg/mL) is plotted as a dotted black line. Values <LLOQ are plotted as 0 pg/mL. Statistical significance analysis was attempted using a Mixed-effects model, however due to values <LLOQ a p value could not be calculated. "Naïve control" received no H1N1 dosing. "Vehicle control" received the H1N1 dosing but was given no treatment. Others received the H1N1 dosing and were given mitlet or mitochondria as shown in legend. No results reached statistical significance.

Figure 3. Platelet extracellular vesicles (mitlets) isolated from young mice and transplanted into aged recipient mice with sepsis infection significantly improved survival. (A) Mitlet injections, first experiment. (B) Mitlet injections, second experiment. (C) Mitlet injections, first and second experiments combined. (D) Liver isolated mitochondria alone. Statistical significance estimated manually as p < 0.05.

Figure 4. (A/B/C/D) Platelet extracellular vesicles (mitlets) isolated from young mice and transplanted into aged recipient mice with sepsis infection, significantly reduced cytokine IL-6 and bacterial levels. (A) Bacterial levels after mitlet injections, first experiment. (B) Bacterial levels after mitlet injections, second experiment. (C) Bacterial levels after mitlet injections, first and second experiments combined. (D) Plasma IL-6 levels for first experiment only. Liver isolated mitochondria not shown. Statistical significance estimated manually as shown.

REFERENCES

1. Björn Ingelsson, Daniel Söderberg, Tobias Strid, Anita Söderberg, Ann-Charlotte Bergh, Vesa Loitto, Kourosh Lotfi, Mårten Segelmark, Giannis Spyrou, Anders Rosén. Lymphocytes eject interferogenic mitochondrial DNA webs in response to CpG and non-CpG oligodeoxynucleotides of class C. Proceedings of the National Academy of Sciences, 2018; 201711950 DOI: 10.1073/pnas.1711950115

2. Wiedemann, N. & Pfanner, N. Mitochondrial Machineries for Protein Import and Assembly. *Annu. Rev. Biochem.* **86**, 685-714 (2017).

3. Beignon, F., Gueguen, N., Tricoire-Leignel, H., Mattei, C. & Lenaers, G. The multiple facets of mitochondrial regulations controlling cellular thermogenesis. *Cell. Mol. Life Sci.* **79**, 525 (2022).

4. Schlame, M. Cardiolipin synthesis for the assembly of bacterial and mitochondrial membranes. *J. Lipid Res.* **49**, 1607-1620 (2008).

5. Anderson, S.*, et al.* Sequence and organization of the human mitochondrial genome. *Nature* **290**, 457-465 (1981).

6. Mercer, T.R.*, et al.* The human mitochondrial transcriptome. *Cell* **146**, 645-658 (2011).

7. Andersson, S.G.*, et al.* The genome sequence of Rickettsia prowazekii and the origin of mitochondria. *Nature* **396**, 133-140 (1998).

8. Buzas, E.I. The roles of extracellular vesicles in the immune system. *Nat. Rev. Immunol.*, 1-15 (2022).

9. Abels, E.R. & Breakefield, X.O. Introduction to Extracellular Vesicles: Biogenesis, RNA Cargo Selection, Content, Release, and Uptake. *Cell. Mol. Neurobiol.* **36**, 301-312 (2016).

10. Puhm, F., Boilard, E. & Machlus, K.R. Platelet Extracellular Vesicles: Beyond the Blood. *Arterioscler. Thromb. Vasc. Biol.* **41**, 87-96 (2021).

11. Marcoux, G.*, et al.* Platelet EVs contain an active proteasome involved in protein processing for antigen presentation via MHC-I molecules. *Blood* (2021).

12. Al Amir Dache, Z.*, et al.* Blood contains circulating cell-free respiratory competent mitochondria. *FASEB J.* **34**, 3616-3630 (2020).

13. Melki, I.*, et al.* Platelets release mitochondrial antigens in systemic lupus erythematosus. *Sci. Transl. Med.* **13**(2021).

14. Boudreau, L.H.*, et al.* Platelets release mitochondria serving as substrate for bactericidal group IIA-secreted phospholipase A2 to promote inflammation. *Blood* **124**, 2173-2183 (2014).

15. Becker, Y.L.C., Duvvuri, B., Fortin, P.R., Lood, C. & Boilard, E. The role of mitochondria in rheumatic diseases. *Nat Rev Rheumatol* **18**, 621-640 (2022).

16. Peruzzotti-Jametti, L.*, et al.* Neural stem cells traffic functional mitochondria via extracellular vesicles. *PLoS Biol.* **19**, e3001166 (2021).

17. Hayakawa, K.*, et al.* Transfer of mitochondria from astrocytes to neurons after stroke. *Nature* **535**, 551-555 (2016).

18. Puhm, F.*, et al.* Mitochondria Are a Subset of Extracellular Vesicles Released by Activated Monocytes and Induce Type I IFN and TNF Responses in Endothelial Cells. *Circ. Res.* **125**, 43-52 (2019).

19. Islam, M.N.*, et al.* Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nat. Med.* **18**, 759-765 (2012).

20. Lood, C.*, et al.* Neutrophil extracellular traps enriched in oxidized mitochondrial DNA are interferogenic and contribute to lupus-like disease. *Nat. Med.* **22**, 146-153 (2016).

21. Torralba, D., Baixauli, F. & Sanchez-Madrid, F. Mitochondria Know No Boundaries: Mechanisms and Functions of Intercellular Mitochondrial Transfer. *Front Cell Dev Biol* **4**, 107 (2016).

22. Nakajima, A., Kurihara, H., Yagita, H., Okumura, K. & Nakano, H. Mitochondrial Extrusion through the cytoplasmic vacuoles during cell death. *J. Biol. Chem.* **283**, 24128-24135 (2008).

23. Todkar, K.*, et al.* Selective packaging of mitochondrial proteins into extracellular vesicles prevents the release of mitochondrial DAMPs. *Nature communications* **12**, 1971 (2021).

24. D'Acunzo, P.*, et al.* Mitovesicles are a novel population of extracellular vesicles of mitochondrial origin altered in Down syndrome. *Sci Adv* **7**(2021).

25. Kapur, R., Zufferey, A., Boilard, E. & Semple, J.W. Nouvelle cuisine: platelets served with inflammation. *J. Immunol.* **194**, 5579-5587 (2015).

26. Semple, J.W., Italiano, J.E. & Freedman, J. Platelets and the immune continuum. *Nat. Rev. Immunol.* **11**, 264-274 (2011).

27. Machlus, K.R. & Italiano, J.E., Jr. The incredible journey: From megakaryocyte development to platelet formation. *J. Cell Biol.* **201**, 785-796 (2013).

28. Boilard, E., Duchez, A.C. & Brisson, A. The diversity of platelet microparticles. *Curr. Opin. Hematol.* **22**, 437-444 (2015).

29. Brisson, A.R., Tan, S., Linares, R., Gounou, C. & Arraud, N. Extracellular vesicles from activated platelets: a semiquantitative cryo-electron microscopy and immuno-gold labeling study. *Platelets* **28**, 263-271 (2017).

30. Marchi, S., Guilbaud, E., Tait, S.W.G., Yamazaki, T. & Galluzzi, L. Mitochondrial control of inflammation. *Nat. Rev. Immunol.* **23**, 159-173 (2023).

31. Hasuwa, H.*, et al.* Transgenic mouse sperm that have green acrosome and red mitochondria allow visualization of sperm and their acrosome reaction in vivo. *Exp. Anim.* **59**, 105-107 (2010).

32. Marcoux, G.*, et al.* Platelet EVs contain an active proteasome involved in protein processing for antigen presentation via MHC-I molecules. *Blood* **138**, 2607-2620 (2021).

33. Marcoux, G.*, et al.* Revealing the diversity of extracellular vesicles using high-dimensional flow cytometry analyses. *Sci. Rep.* **6**, 35928 (2016).

34. Becker, Y.L.*, et al.* Identification of mitofusin 1 and complement component 1q subcomponent-binding protein as mitochondrial targets in systemic lupus erythematosus. *Arthritis & rheumatology* (2022).

35. Pelletier, M., Billingham, L.K., Ramaswamy, M. & Siegel, R.M. Extracellular flux analysis to monitor glycolytic rates and mitochondrial oxygen consumption. *Methods Enzymol.* **542**, 125-149 (2014).

36. Chargaff, E. & West, R. The biological significance of the thromboplastic protein of blood. *J. Biol. Chem.* **166**, 189-197 (1946).

37. Wolf, P. The nature and significance of platelet products in human plasma. *Br. J. Haematol.* **13**, 269-288 (1967).

38. Berckmans, R.J., Lacroix, R., Hau, C.M., Sturk, A. & Nieuwland, R. Extracellular vesicles and coagulation in blood from healthy humans revisited. *J. Extracell. Vesicles* **8**, 1688936 (2019).

39. Burnouf, T., *et al.* An overview of the role of microparticles/microvesicles in blood components: Are they clinically beneficial or harmful? *Transfus. Apher. Sci.* **53**, 137-145 (2015).

40. Burnouf, T., Goubran, H.A., Chou, M.L., Devos, D. & Radosevic, M. Platelet microparticles: detection and assessment of their paradoxical functional roles in disease and regenerative medicine. *Blood Rev.* **28**, 155-166 (2014).

41. Jonavice, U., Tunaitis, V., Kriauciunaite, K., Jarmalaviciute, A. & Pivoriunas, A. Extracellular vesicles can act as a potent immunomodulators of human microglial cells. *J. Tissue Eng. Regen. Med.* **13**, 309-318 (2019).

42. Lam, S.M., *et al.* A multi-omics investigation of the composition and function of extracellular vesicles along the temporal trajectory of COVID-19. *Nat Metab* **3**, 909-922 .

43. Russell, A.E., *et al.* Extracellular Vesicles Secreted in Response to Cytokine Exposure Increase Mitochondrial Oxygen Consumption in Recipient Cells. *Front. Cell. Neurosci.* **13**, 51 (2019).

44. Chimen, M., *et al.* Appropriation of GPIbalpha from platelet-derived extracellular vesicles supports monocyte recruitment in systemic inflammation. *Haematologica* **105**, 1248-1261 (2020).

45. Risitano, A., Beaulieu, L.M., Vitseva, O. & Freedman, J.E. Platelets and platelet-like particles mediate intercellular RNA transfer. *Blood* **119**, 6288-6295 (2012).

46. Duchez, A.C., *et al.* Platelet microparticles are internalized in neutrophils via the concerted activity of 12-lipoxygenase and secreted phospholipase A2-IIA. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E3564-3573 (2015).

47. Laffont, B., *et al.* Activated platelets can deliver mRNA regulatory Ago2*microRNA complexes to endothelial cells via microparticles. *Blood* **122**, 253-261 (2013).

48. Melki, I., *et al.* FcgammaRIIA expression accelerates nephritis and increases platelet activation in systemic lupus erythematosus. *Blood* **136**, 2933-2945 (2020).

49. Arraud, N., *et al.* Extracellular vesicles from blood plasma: determination of their morphology, size, phenotype and concentration. *J. Thromb. Haemost.* **12**, 614-627 (2014).

50. Boilard, E. & Bellio, M. Platelet extracellular vesicles and the secretory interactome join forces in health and disease. *Immunol. Rev.* **312**, 38-51 (2022).

51. Bao, Y., *et al.* Mitochondria regulate neutrophil activation by generating ATP for autocrine purinergic signaling. *J. Biol. Chem.* **289**, 26794-26803 (2014).

52. Wang, X. & Chen, D. Purinergic Regulation of Neutrophil Function. *Front. Immunol.* **9**, 399 (2018).

53. Caicedo, A., *et al.* The diversity and coexistence of extracellular mitochondria in circulation: A friend or foe of the immune system. *Mitochondrion* **58**, 270-284 (2021).

54. Miliotis, S., Nicolalde, B., Ortega, M., Yepez, J. & Caicedo, A. Forms of extracellular mitochondria and their impact in health. *Mitochondrion* **48**, 16-30 (2019).

55. Becker, Y., *et al.* Anti-mitochondrial autoantibodies in systemic lupus erythematosus and their association with disease manifestations. *Sci. Rep.* **9**, 4530 (2019).

56. Becker, Y., *et al.* Autoantibodies in Systemic Lupus Erythematosus Target Mitochondrial RNA. *Front. Immunol.* **10**, 1026 (2019).

57. Berg, P.A. & Klein, R. Mitochondrial antigens and autoantibodies: from anti-M1 to anti-M9. *Klin. Wochenschr.* **64**, 897-909 (1986).

58. Zeeuw van der Laan, E.A.N., *et al.* Update on the pathophysiology of transfusion-related acute lung injury. *Curr. Opin. Hematol.* **27**, 386-391 (2020).

59. Toy, P., *et al.* Recipient clinical risk factors predominate in possible transfusion-related acute lung injury. *Transfusion (Paris)* **55**, 947-952 (2015).

60. Spees, J.L., Olson, S.D., Whitney, M.J. & Prockop, D.J. Mitochondrial transfer between cells can rescue aerobic respiration. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 1283-1288

61. King, M.P. & Attardi, G. Injection of mitochondria into human cells leads to a rapid replacement of the endogenous mitochondrial DNA. *Cell* **52**, 811-819 (1988).

62. Lin, H.Y., *et al.* Mitochondrial transfer from Wharton's jelly-derived mesenchymal stem cells to mitochondria-defective cells recaptures impaired mitochondrial function. *Mitochondrion* **22**, 31-44 (2015).

63. Lightowlers, R.N., Chrzanowska-Lightowlers, Z.M. & Russell, O.M. Mitochondrial transplantation-a possible therapeutic for mitochondrial dysfunction?: Mitochondrial transfer is a potential cure for many diseases but proof of efficacy and safety is still lacking. *EMBO Rep* **21**, e50964

64. Sofou, K., *et al.* A multicenter study on Leigh syndrome: disease course and predictors of survival. *Orphanet J. Rare Dis.* **9**, 52 (2014).

65. Jacoby, E., *et al.* Mitochondrial augmentation of hematopoietic stem cells in children with single large-scale mitochondrial DNA deletion syndromes. *Sci. Transl. Med.* **14**, eabo3724 (2022)

66. Steele AM, Starr ME, Saito H. Late therapeutic intervention with antibiotics and fluid resuscitation allows for a prolonged disease course with high survival in a severe murine model of Sepsis. Shock 47:726–734. (2017) DOI: https://doi.org/10.1097/SHK.0000000000000799, PMID: 27879561

67. Flurkey K M, Currer J, Harrison DE. 2007. Chapter 20 - Mouse Models in Aging Research A2 - Fox. In: James G, Davisson MT, Quimby FW, Barthold SW, Newcomer CE, Smith AL (Eds). The Mouse in Biomedical Research. Second Edition. Burlington: Academic Press. p. 637–672.

68. Pelletier M, Breton Y, Allaeys I, Becker Y, Benson T, Boilard E. Platelet extracellular vesicles and their mitochondrial content improve the mitochondrial bioenergetics of cellular immune recipients. Transfusion. (2023) https://doi.org/10.1111/trf.17524

69. Allison M Owen, Samir P Patel, Jeffrey D Smith, Beverly K Balasuriya, Stephanie F Mori, Gregory S Hawk, Arnold J Stromberg, Naohide Kuriyama, Masao Kaneki, Alexander G Rabchevsky, Timothy A Butterfield, Karyn A Esser, Charlotte A Peterson, Marlene E Starr, Hiroshi Saito. Chronic muscle weakness and mitochondrial dysfunction in the absence of sustained atrophy in a preclinical sepsis model eLife 8:e49920. (2019) https://doi.org/10.7554/eLife.49920

70. Wang R, Stone RL, Kaelber JT, Rochat RH, Nick AM, Vijayan KV, Afshar-Kharghan V, Schmid MF, Dong JF, Sood AK, Chiu W. Electron cryotomography reveals ultrastructure alterations in

platelets from patients with ovarian cancer. Proc Natl Acad Sci U S A. 2015 Nov 17;112(46):14266-71. doi: 10.1073/pnas.1518628112. Epub 2015 Nov 2. PMID: 26578771; PMCID: PMC4655568.

71. Natalia Bobba-Alves, Gabriel Sturm, Jue Lin, Sarah A. Ware, Kalpita R. Karan, Anna S. Monzel, Céline Bris, Vincent Procaccio, Guy Lenaers, Albert Higgins-Chen, Morgan Levine, Steve Horvath, Balaji S. Santhanam, Brett A. Kaufman, Michio Hirano, Elissa Epel, Martin Picard, Cellular allostatic load is linked to increased energy expenditure and accelerated biological aging, Psychoneuroendocrinology, Volume 155, 2023, 106322, https://doi.org/10.1016/j.psyneuen.2023.106322

72. McCully, J.D., D.B. Cowan, C.A. Pacak, and S. Levitsky, Mitochondrial transplantation for cardioprotection. Circulation, 2007. 116(16): p. 496-496.

73. Zhang, X., Hubal, M.J. & Kraus, V.B. Immune cell extracellular vesicles and their mitochondrial content decline with ageing. Immun Ageing 17, 1 (2020). https://doi.org/10.1186/s12979-019-0172-9

74. Mitrix Bio Inc. Website. https://www.mitrix.bio.

75. Ikeda G, Santoso MR, Tada Y, Li AM, Vaskova E, Jung JH, O'Brien C, Egan E, Ye J, Yang PC. Mitochondria-Rich Extracellular Vesicles From Autologous Stem Cell-Derived Cardiomyocytes Restore Energetics of Ischemic Myocardium. J Am Coll Cardiol. 2021 Mar 2;77(8):1073-1088. doi: 10.1016/j.jacc.2020.12.060.

76. Lazo S, Noren Hooten N, Green J, Eitan E, Mode NA, Liu QR, Zonderman AB, Ezike N, Mattson MP, Ghosh P, Evans MK. Mitochondrial DNA in extracellular vesicles declines with age. Aging Cell. 2021 Jan;20(1):e13283. doi: 10.1111/acel.13283.

77. Adlimoghaddam, A., T. Benson, and B.C. Albensi, Mitochondrial Transfusion Improves Mitochondrial Function Through Up-regulation of Mitochondrial Complex II Protein Subunit SDHB in the Hippocampus of Aged Mice. Mol Neurobiol, 2022. 59(10): p. 6009-6017.

78. Liu X, Khouri-Farah N, Wu CH, Wu GY. Targeted delivery of mitochondria to the liver in rats. J Gastroenterol Hepatol. 2020 Dec;35(12):2241-2247. doi: 10.1111/jgh.15091.

79. Lane, Nick. The Vital Question: Why is life the way it is? Profile, 2015 ISBN 1847658806, 9781847658807

80. Joseph W. Guarnieri et al. Core mitochondrial genes are down-regulated during SARS-CoV-2 infection of rodent and human hosts. Sci. Transl. Med.15,eabq1533(2023). DOI:10.1126/scitranslmed.abq1533

81. Singh KK, Chaubey G, Chen JY, Suravajhala P. Decoding SARS-CoV-2 hijacking of host mitochondria in COVID-19 pathogenesis. Am J Physiol Cell Physiol. 2020 Aug 1;319(2):C258-C267. doi: 10.1152/ajpcell.00224.2020.

82. Zhu X, Li Q, Zhu X. Mechanisms of CAR T cell exhaustion and current counteraction strategies. Front Cell Dev Biol. (2022) doi: 10.3389/fcell.2022.1034257.

83. Loeb LA, Wallace DC, Martin GM. The mitochondrial theory of aging and its relationship to reactive oxygen species damage and somatic mtDNA mutations. Proc Natl Acad Sci U S A. 2005 Dec 27;102(52):18769-70. doi: 10.1073/pnas.0509776102.

84. Melov, S., Mitochondrial dysfunction and aging. Gerontologist, 2001. 41: p. 65-65. '

85. Kam, J.H., Weinrich, T.W., Shinhmar, H. et al. Fundamental differences in patterns of retinal ageing between primates and

mice. Sci Rep 9, 12574 (2019). https://doi.org/10.1038/s41598-019-49121-0

86. Rosa FLL, de Souza IIA, Monnerat G, Campos de Carvalho AC, Maciel L. Aging Triggers Mitochondrial Dysfunction in Mice. International Journal of Molecular Sciences. 2023; 24(13):10591. https://doi.org/10.3390/ijms241310591

87. Hosseinian, S., P. Ali Pour, and A. Kheradvar, Prospects of mitochondrial transplantation in clinical medicine: Aspirations and challenges. Mitochondrion, 2022. 65: p. 33-44.

88. McCully, J., P.J. del Nido, and S.M. Emani, Therapeutic mitochondrial transplantation. Current Opinion in Physiology, 2022. 27.