Methylotrophic methanogenesis in the Archaeoglobi revealed by cultivation of *Ca.* Methanoglobus hypatiae from a Yellowstone hot spring

Mackenzie M. Lynes[1], Zackary J. Jay[1], Anthony J. Kohtz[1], Roland Hatzenpichler[1,2,*]

[1] Department of Chemistry and Biochemistry, Center for Biofilm Engineering, and Thermal Biology Institute, Montana State University, Bozeman, MT 59717, USA

[2] Department of Microbiology and Cell Biology, Montana State University, Bozeman, MT 59717, USA

[*] Corresponding author: Roland Hatzenpichler, email: roland.hatzenpichler@montana.edu, 111 Chemistry and Biochemistry Building, Montana State University, MT-59718, Bozeman, USA

ORCID Numbers of Authors: M.M.L., 0000-0002-9410-0285; Z.J.J., 0000-0003-3062-4933; A.J.K., 0000-0002-0561-8710; R.H, 0000-0002-5489-3444

Running title: *Ca.* Methanoglobus hypatiae

Abstract

Over the past decade, environmental metagenomics and PCR-based marker gene surveys have revealed that several lineages beyond just a few well-established groups within the Euryarchaeota superphylum harbor the genetic potential for methanogenesis. One of these groups are the Archaeoglobi, a class of thermophilic euryarchaeotes that have long been considered to live non-methanogenic lifestyles. Here, we enriched *Candidatus* Methanoglobus hypatiae, a methanogen affiliated with the family Archaeoglobaceae, from a hot spring in Yellowstone National Park. The enrichment is sediment-free, grows at 64-70 ºC and a pH of 7.8, and produces methane from mono-, di-, and tri-methylamine. *Ca.* M. hypatiae is represented by a 1.62 Mb metagenome-assembled genome with an estimated completeness of 100% and accounts for up to 67% of cells in the culture according to fluorescence *in situ* hybridization. Via genome-resolved metatranscriptomics and stable isotope tracing, we demonstrate that *Ca.* M. hypatiae expresses methylotrophic methanogenesis and energy-conserving pathways for reducing monomethylamine to methane. The detection of Archaeoglobi populations related to *Ca.* M. hypatiae in 36 geochemically diverse geothermal sites within Yellowstone National Park, as revealed through the examination of previously published gene amplicon datasets, implies a previously underestimated contribution to anaerobic carbon cycling in extreme ecosystems.

Keywords: archaea, MCR, methane, stable isotope tracing, thermophile, transcriptomics

Introduction

Methanogenesis is one of the most ancient metabolic pathways and plays a major role in the biogeochemical carbon cycle. Phylogenomic reconstructions and geological evidence suggest that methanogenesis was among the earliest metabolisms to evolve and that the last common ancestor of all extant archaea likely was a methanogen (1-9). Therefore, the study of methanogens is essential for understanding the co-evolution of life and the biosphere. Methanogenic archaea are the primary producers of biogenic methane (CH_4) and contribute approximately 60% to the estimated 576 Tg of annual global methane emissions to the atmosphere (10, 11). Methanogenic pathways are classified by their carbon and electron sources (12-14). All methanogenic pathways converge at the terminal methane-forming step catalyzed by the methyl-coenzyme M reductase (MCR) complex. MCR and its homologs also catalyze the reverse reaction in the anaerobic oxidation of alkanes in alkanotrophic archaea (15, 16). MCR is uniquely present in all methanogens and is commonly used to identify potential methane and/or alkane cycling archaea in sequencing surveys (12, 17).

The physiology and biochemistry of methanogens has near-exclusively been investigated in axenic cultures of microorganisms belonging to the Euryarchaeota superphylum (12, 17-19). These predominantly grow by acetoclastic or CO_2-reducing hydrogenotrophic methanogenesis, with only rare observations of Euryarchaeotal methyl-reducing methanogens (12, 20, 21). As a result, despite the dominance of methyl-based methanogenesis in anoxic environments with high salt and/or high sulfate concentration (*e.g.*, anoxic marine sediments, coastal wetlands, hypersaline lakes), methylotrophic methanogenesis has in the past often been considered to be of comparatively limited environmental distribution. The extensive use of environmental metagenomics has led to the discovery of metagenome-assembled genomes (MAGs) encoding MCR from new lineages that are prevalent in anoxic environments, both within and outside the Euryarchaeota (2, 12, 22-26).

The majority of MAGs affiliated with archaeal phyla outside the Euryarchaeota are predicted to be methyl-reducing methanogens, with the exception of *Candidatus* (*Ca.*) Nezhaarchaeota (25, 27) and *Ca.* Methanomixophus affiliated with the order Archaeoglobales, which have been hypothesized to be CO_2-reducing hydrogenotrophic methanogens (12, 25, 28). This result is consistent with the observation that methylated methanogenic substrates, including methylamines and methanol, are prevalent in the environment, although their concentrations in hot springs is currently unknown. Further, methyl-reducing methanogenesis is considered the predominant mode of methanogenesis in anoxic marine, freshwater, and hypersaline sediments (reviewed in (20)).

Members of the class Archaeoglobi have long been considered non-methanogenic with isolates characterized as dissimilatory sulfate reducers brought into culture as early as 1987 (29). To date, only nine species of the class Archaeoglobi have been obtained in axenic culture, and all were sourced from marine

hydrothermal systems or off-shore oil reservoirs (30). The discovery of both MCR (25, 31, 32) and methyl-

H₄M(S)PT:coenzyme M methyltransferase (MTR) complexes in genomes of the Archaeoglobaceae have

suggested that members of this family may live by methanogenesis (28).

Very recently, important progress towards experimental verification of methanogenesis by members

of this family has been made. Liu *et al.* reported the *in situ* expression of genes related to hydrogen-

dependent methylotrophic methanogenesis and heterotrophic fermentation within populations of

Archaeoglobi in a high-temperature oil reservoir (28). Lynes, Krukenberg *et al.* reported that Archaeoglobi

can be enriched in hot spring mesocosms under methanogenic conditions (33). Wang *et al.* reported that

mcrABG and other methanogenesis marker genes encoded by two Archaeoglobales MAGs were highly

expressed in hot spring microcosms incubated at 65 ºC and 75 ºC (34). Importantly, one of these

Archaeoglobales MAGs represented the only Mcr-encoding archaeon that expressed *mcrABG* genes in

methanogenic microcosms performed without substrate addition or with the addition of 10 mM methanol

at 75 ºC. This indirectly demonstrated the methanogenic nature of this archaeon (34). Last, Buessecker *et*

al. reported the establishment of a methanogenic enrichment culture of *Ca.* Methanoglobus nevadensis from

Great Boiling Spring (NV, USA) (35). The culture yields up to 158 µM methane after two weeks of

incubation at its optimal growth temperature of 75 ºC. *Ca.* M. nevadensis is represented by a 63% complete

MAG obtained from the culture and a 98% complete MAG obtained a decade earlier (35).

Here, we report on the enrichment cultivation of *Ca.* Methanoglobus hypatiae LCB24, a methanogen

affiliated with the family Archaeoglobaceae, from a hot spring in Yellowstone National Park (YNP). Using

a combination of targeted cultivation, growth experiments, microscopy, stable isotope tracing,

metagenomics, and metatranscriptomics, we demonstrate that *Ca.* M. hypatiae lives by methylotrophic

methanogenesis and converts different methylamines to methane. By examining previously published

datasets for the presence of Mcr-encoding Archaeoglobi, we demonstrate that these archaea are distributed

in geothermal features of YNP, where they likely contribute to anaerobic carbon cycling. Our study presents

direct evidence of methanogenesis within the Archaeoglobaceae and adds to the growing body of evidence

demonstrating that methanogenesis is widely spread within the Euryarchaeota superphylum.

Materials and Methods

All chemicals used in this study were sourced from Sigma Aldrich unless otherwise specified.

Sample Collection, Enrichment, and Cultivation

In November 2021, a slurry of sediment and water (1:9) was collected from an unnamed hot spring

in the Lower Culex Basin of Yellowstone National Park (YNP), WY, USA. In our previous survey of Mcr-

encoding archaea in YNP (33), this hot spring was given the identifier LCB024 (44.573294, -110.795388;

pH 7.8, 67 ˚C). A mixture of surface sediment (~1-2 cm deep) and hot spring water was collected into a

glass bottle and sealed headspace-free with a thick butyl rubber stopper. Collected material was transported

back to the lab and stored at room temperature. Using this material as inoculum, 30 mL enrichments were

established in February 2022 in 60 mL serum bottles. Material was homogenized by mixing and was then

diluted 1:10 (v/v) with anoxic medium in an anoxic glove box ($N_2/CO_2/H_2$; 90/5/5%).

Medium was prepared anoxically as described previously (36). Basal mineral medium contained a

base of (per liter): KH_2PO_4, 0.5 g; $MgSO_4 \cdot 7H_2O$, 0.4 g; NaCl, 0.5 g; NH_4Cl, 0.4 g; $CaCl_2 \cdot 2H_2O$, 0.05 g;

HEPES, 2.38 g; yeast extract, 0.1 g; and 0.002% (w/v) $(NH_4)_2Fe(SO_4)_2 \cdot 6H_2O$. Medium was transferred to

a Duran flask with a side opening and autoclaved for 20 m at 121 ˚C. Medium was then further

supplemented with 5 mM $NaHCO_3$, 1 mL trace element solution SL-10, 1 mL Selenite-Tungstate solution,

1 mL CCM vitamins (37), 0.0005% (w/v) resazurin, 10 mg of coenzyme-M, 2 mg sodium dithionite, 1 mM

dithiothreitol, 1 mM $Na_2S \cdot 9H_2O$, with pH adjusted to 7.8 using sodium hydroxide (NaOH, 12 N). Serum

bottles were sealed with butyl rubber stoppers and aluminum crimps before the headspace was exchanged

with N_2 (99.999%) for 5 minutes and set to a 200 kPa N_2 atmosphere. Monomethylamine (MMA) was added

from a filter-sterilized methylamine-hydrochloride anoxic stock solution to a final concentration of 10 mM.

The bacterial antibiotics streptomycin (50 mg/L; inhibitor of protein synthesis) and vancomycin (50 mg/L;

inhibitor of peptidoglycan synthesis) were added from filter-sterilized anoxic stock solutions. The

enrichments were incubated at 70 ˚C in the dark without shaking. Cultures were maintained by regular

transfer of 10% v/v into fresh media, which contained MMA and antibiotics. A sediment-free culture was

obtained after the third transfer after which it was transferred at 10% v/v to 50 mL in 125 mL serum bottles.

Stable Isotope Tracing

The conversion of ^{13}C- or D_3-MMA ($^{13}CH_3$-NH_2, CD_3-NH_2) to $^{13}CH_4$ or CD_3H was tracked by

incubating active enrichment cultures in the presence of 20% labeled substrate (98%; Cambridge Isotope

Laboratories). Incubations were carried out in 30 mL culture volumes in 60 mL serum bottles with 8% v/v

inoculum, 50 mg/L streptomycin, 50 mg/L vancomycin, 10 mM MMA, and N_2 gas (99.999%) incubated in

anoxic media (pH 7.8, 70 ˚C) in six replicates (SI Appendix, Fig. S3). Duplicate control incubations

included (i) ^{12}C-MMA and (ii) inoculum without MMA. Triplicate control incubations were performed with

(iii) ^{12}C-MMA plus 10 mM bromoethanesulfonate (BES) added in mid-exponential phase (day 33) to inhibit

methanogenesis and (iv) 10 mM BES added at time of inoculation (day 0) without substrate. Headspace

samples were collected throughout the experiment as described above and analyzed using a Shimadzu

QP2020 NX GCMS equipped with a GS-CarbonPLOT column (30 m × 0.35 mm; 1.5 μm film thickness;

Agilent) and operated in Selected Ion Monitoring mode. The instrument was operated using the method

described in Ai et al., 2013 (38) with helium as a carrier gas. All injections were performed by a Shimadzu

AOC-6000 autosampler robot. Peak areas corresponding to m/z ratios of 16 for $^{12}CH_4$, 17 for $^{13}CH_4$, and 19

for CD_3H were used for quantification.

Metagenomic Sequencing, Assembly, and Annotation

Two metagenomes were obtained over the course of this study. A 42 mL aliquot of the fourth transfer

of the enrichment (Fig. 1 T4-MG) was filtered onto a 0.22 µm filter. The filter was transferred to a lysing

matrix E tube and DNA extracted immediately following filtration. Genomic DNA was extracted using the

FastDNA Spin Kit for Soil (MP Biomedicals, Irvine, CA) following the manufacturer's guidelines.

A second metagenome was recovered from one of the six culture replicates grown in the presence of

CD_3-NH_2 and used for recruiting transcriptomic reads from the other replicates (Fig. 1 SIT-MG). A 60 mL

syringe flushed with N_2 gas was used to transfer 30 mL of culture to a sterilized oak ridge tube. Cells were

harvested through centrifugation for 30 minutes at 10,000 rpm at 4 ˚C. The supernatant was removed, and

DNA extracted from the pellet using the FastDNA Spin Kit for Soil (MP Biomedicals, Irvine, CA)

following the manufacturer's guidelines. Genomic DNA for both metagenomes was shipped to SeqCenter

(Pittsburgh, PA) and sample libraries were prepared using the Illumina DNA Prep kit and 10 bp unique

dual indices (UDI). The first metagenome (T4-MG) was sequenced on a NextSeq 2000 System (Illumina)

and the second (SIT-MG) sequenced on a NovaSeq 6000 System (Illumina), each producing 2x151 bp

reads. Demultiplexing, quality control, and adapter trimming was performed with bcl-convert v3.9.3.

Quality of the reads were evaluated with FastQC before quality, linker and adapter trimming, artifact and

common contaminant removal, and error correction were performed with the rqcfilter2 pipeline (maxn=3,

maq=10, trimq=20) and bbcms (mincount=2, hcf=0.6). Resulting reads were assembled with SPAdes

v3.15.13 (Nurk, 2017) (-k 33,55,77,99,127 --meta –only-assembler) and coverage was determined with

bbmap v38.94 (ambiguous=random) (https://sourceforge.net/projects/bbmap) (39). In addition to the initial

assembly, co-assemblies using both T4-MG and SIT-MG metagenomes were also performed (1) with reads

directly fed into SPAdes with the –only-assembler option excluded; and (2) with the trimmed and error

corrected reads and the same SPAdes parameters as above. The statistics of MAGs generated through

various assembly and quality control methods were evaluated, and the approach that produced the highest

quality MAG was chosen for subsequent analysis (Dataset S1). Quality was determined by considering

factors such as the number of resulting sequences, total length, completeness, and the minimum, maximum,

and average sequence lengths. Annotation of the assembled sequences was performed with Prokka v1.14.16

(40). Assembled scaffolds ≥2000 bp were binned using Maxbin v2.2.7 (41), Metabat v2.12.1 (with and

without coverage) (42), Concoct v1.0.0 (43), Autometa v1 (bacterial and archaeal modes with the machine

learning step) (44), followed by bin refinement with DAS_Tool v1.1.6 (45), as previously described (46).

Bin completeness and redundancy were assessed with CheckM v1.2.2 (47).

RNA Extraction, Sequencing, and Transcriptomic Processing

Total RNA was extracted for transcriptomics from four of the six replicates of Archaeoglobus cultivated in the presence of labeled substrate ($^{13}CH_3-NH_2$ or CD_3-NH_2) for a total of eight replicates. Each replicate culture in the exponential growth phase (day 32) was moved from the 70 ˚C incubator to an ice bath placed at -20 ˚C for 40 minutes to stop cellular activity. A 60 mL syringe flushed with N_2 gas was used to transfer 30 mL of culture to a sterilized oak ridge tube and kept on ice. Cells were harvested through centrifugation for 30 minutes at 10,000 rpm at 4 ˚C. The supernatant was removed, and the pellet transferred to a lysing matrix E tube (MP Biomedicals, Irvine, CA) to which 600 µL of RNA lysis buffer was added. Samples were homogenized in a MP Bioscience FastPrep instrument for 40 seconds at a speed setting of 6.0 m/s followed by centrifugation for 15 minutes at 14,000 rpm. RNA was extracted using the Quick-RNA miniprep kit (Zymo Research, Irvine, CA) including a DNAse treatment step and eluted in 50 µL of RNAse free water. Centrifugation steps were performed at 15,000 rpm and the final spin for elution at 10,000 rpm. Of the eight replicates extracted, six measured >50 ng/µL (3x $^{13}CH_3-NH_2$ and 3x CD_3-NH_2) and were sent for transcriptomic sequencing at SeqCenter (Pittsburgh, PA). Samples were DNAse treated with Invitrogen DNAse (RNAse free). Library preparation was performed using Illumina's Stranded Total RNA Prep Ligation with Ribo-Zero Plus kit and 10bp UDI. Sequencing was done on a NovaSeq 6000, producing paired end 151bp reads. Demultiplexing, quality control, and adapter trimming was performed with bcl-convert (v4.1.5). Read quality was further evaluated with FastQC v0.11.9 (48) before quality trimming and artifact, rRNA, and common contaminant removal with the rqcfilter2 pipeline (trimq=28, maxns=3, maq=20), and error correction with bbcms (mincount=2, hcf=0.6) from the BBTools suite v38.94 (39). Additional rRNA gene reads were detected and removed with Ribodetector v0.2.7 (49) and any remaining rRNA gene reads were finally removed with bbmap, using rRNA genes recovered from the metagenomes (see below) as references. The resulting mRNA reads were mapped against annotated genes from the paired metagenomes with bbmap to calculate RPKM (ambig=random).

Data Availability

All metagenomic, metatranscriptomic, and amplicon data discussed in this manuscript are available under NCBI BioProject ID PRJNA1014417. McrA gene amplicon data from YNP hot springs discussed in this manuscript has been previously published (Lynes et al., 2023) and is available under NCBI under BioProject PRJNA859922.

Results and Discussion

Cultivation

In our recent survey on the diversity of Mcr-encoding archaea in the geothermal features of YNP,

mesocosms seeded with biomass from a hot spring located within the Lower Culex Basin (LCB024; pH 7-

8, 56-74 ˚C), had shown potential to enrich for methanogenic Archaeoglobi (33). Using a sediment slurry

collected from LCB024, we initiated incubations supplied with monomethylamine (MMA) and antibiotics

incubated in anoxic media (pH 7.8, 70 ˚C) under a N_2 headspace. The relative abundance, as determined by

16S rRNA gene amplicon sequencing, of Archaeoglobi-affiliated organisms in LCB024 was 0.32% in the

initial slurry and had fallen to 0.02% by the time incubations were initiated a few months after samples had

been collected (Fig. 1A).

Methane was detected after 36 days in the initial enrichment and the culture transferred to fresh media

after reaching the late exponential phase of methane production following 70 days of incubation (447 µM;

Fig. 1B). Five Archaeoglobi related 16S rRNA gene amplicon ASVs were identified in the initial

enrichment, however one ASV grew to dominate the microbial community after the first transfer and

reached 74.8% relative abundance after 62 days. In the transfers that followed, Archaeoglobi-related

sequences became the only archaeal reads detected by 16S rRNA gene amplicon sequencing with the

second most abundant organism a bacterium affiliated with the *Pseudothermotoga* at 6.8%. Although the

CO_2-reducing methanogen *Methanothermobacter* sp. was detected at 0.45% relative abundance in the

slurry material used for inoculation, it was not detected in any subsequent transfers, nor were any known

methanogens. Over subsequent transfers (238 days, T2-T5), the relative abundance of Archaeoglobi ASVs

ranged from 46 to 69% and the final methane yield steadily increased from 1,844 to 2,459 µM. A sediment-

free enrichment was obtained by the third transfer. Starting with the fourth transfer, the culture volume was

scaled from 30 mL to 50 mL. By the sixth transfer, the culture produced 3,943 µM methane within 34 days.

Metagenomic sequencing at two timepoints (day 335 of the enrichment and day 33 of the isotope tracing

experiment described below) and 16S rRNA gene amplicon sequencing over recurring transfers (Fig. 1A)

demonstrated that ASVs and MAGs affiliated with Archaeoglobi represented the only archaeon in culture

LCB24. A single MCR complex (*mcrAGCDB*) belonging to the Archaeoglobi MAG was present, indicating

this MAG represents the only methanogenic population.

Metagenomics and Phylogenetics

The reconstructed Mcr-encoding Archaeoglobi MAG from culture LCB24 was 1.62 Mbp in length

with an estimated completeness of 100% according to checkM (SI Appendix, Table S3). This MAG was

the result of a combined assembly of the T4-MG and SIT-MG metagenomes as this method yielded an

improved assembly. Therefore, it was used for phylogenomic analysis against Archaeoglobi reference

MAGs and genomes using 33 conserved single copy marker proteins and 16S rRNA genes (Fig. 2AB, SI

Appendix, Table S4). The phylogenomic analysis showed that MAGs encoding MCR complexes clustered

separately from those lacking *mcr* gene sequences. Consistently, 16S rRNA gene phylogeny supported this

clustering with a pronounced separation of hot spring reference genomes and MAGs from current known

isolates of Archaeoglobi, resulting in three main clusters: (i) those retrieved from North American hot

springs (YNP and Great Boiling Spring, GBS), (ii) those originating from hot springs in China, and (iii)

isolates, all of which were obtained from deep-sea marine hydrothermal systems (Fig. 2B).

LCB24 and closely related reference MAGs and isolate genomes exhibited a range of amino acid

identities (AAI, 52.6-98.6%; Fig. 2C). Altogether, the LCB24 MAG was found to be highly related to

previously obtained Archaeoglobi MAGs encoding the MCR complex and only distantly related to other

Archaeoglobales sp. ~~confirmed sulfate-reducing Archaeoglobales~~ (AAI, 58.9-65%; ANI, 70.3-70.6%; 16S

rRNA ANI, 91.6-93.8%; SI Appendix Fig. S1, S2). Based on AAI, MAG LCB24 was most closely related

to Archaeoglobi LCB024-003 MAG (AAI, 98.6%), which we had obtained from the same hot spring in a

previous study (33). The ANI and AAI values to the closest cultured methanogen, *Ca.* Methanoglobus

nevadensis GBS, are 80.2 and 83.3%, respectively. Based on these results, we designate this archaeon *Ca.*

Methanoglobus hypatiae strain LCB24, named after the philosopher Hypatia of Alexandria (for a

protologue, see the SI Appendix, Results and Discussion). The estimated relative abundance of *Ca.* M.

hypatiae based on the SIT-MG was 92.8%. Other community members in the LCB24 culture with >1%

relative sequence abundance included members of the *Pseudothermotoga* (3.2%), *Desulfovirgula* (1.7%),

and the family Moorellaceae (1.3%) (Fig. 1A, Dataset S1).

The only *mcrAGCDB* genes recovered from both metagenomes belong to the genome of *Ca.* M.

hypatiae. Phylogenetic analysis of the single copy of McrA indicated its close relationship to McrA

sequences found in members of the TACK superphylum (Fig. 2D). This contrasts with the placement of

Ca. M. hypatiae within the Euryarchaeota based on phylogenomics (Fig. 2A), suggesting that Archaeoglobi

could have obtained the MCR complex as a result of a horizontal gene transfer event from an archaeon in

the TACK superphylum (7, 8). Also, it could indicate that non-methanogenic Archaeoglobi lost the capacity

for anaerobic methane cycling after they had diverged from a shared methanogenic ancestor.

Methanogenic Activity of *Ca.* M. hypatiae

To gain insight into the activity of *Ca.* M. hypatiae under methanogenic and non-methanogenic

conditions, a stable isotope tracing (SIT) experiment was conducted. Cultures were incubated in the

presence of 10 mM of MMA; 8 mM of substrate were isotopically light, whereas the remaining 2 mM

consisted of either ^{13}C-monomethylamine ($^{13}CH_3$-NH_2) or D$_3$-monomethylamine (CD_3-NH_2). Addition of

the methanogenesis inhibitor bromoethanesulfonate (BES) was used as a non-methanogenic control (Fig.

1B, 3, SI Appendix, Fig. S3). On average across six replicates, the cultured converted $^{13}CH_3$-NH_2 to 356

µM $^{13}CH_4$ (17.8%) and 138.71 µM $^{13}CO_2$ (6.9%) by day 32 (Fig. 3AC, Dataset S2). The conversion of CD_3-

NH$_2$ was nearly identical yielding 355 μM CD$_3$H (Fig. 3B). In the exponential phase of methane production, five of the six replicates were harvested for metagenomic and metatranscriptomic sequencing while the sixth replicate was allowed to grow to stationary phase. The replicate allowed to grow in each respective experiment converted the provided ^{13}CH$_3$-NH$_2$ to 717.7 μM ^{13}CH$_4$ (35.9%) and 212.95 μM ^{13}CO$_2$ (10.65%) or CD$_3$-NH$_2$ to 394.76 μM CD$_3$H (19.7%) by day 38 (Fig. 3ABC). These results confirmed monomethylamine was converted to methane by the LCB24 culture. The production of ^{13}CO$_2$ may represent the dismutation of ^{13}CH$_3$-NH$_2$ to generate reducing power for methanogenesis via the methyl-branch of the Wood-Ljungdahl pathway or may be explained by other organisms in the culture catabolizing MMA. Yet, no transcriptomic evidence for this activity was present in this experiment. No methane production was observed for cultures treated with BES or in cultures incubated without MMA (Fig. 3D). When BES was added to cultures in the exponential phase, methane production ceased indicating the generation of methane is reliant on the Archaeoglobi MCR (Fig. 3E).

Visualization and Cell Enumeration

The growth of *Ca.* M. hypatiae was tracked in four replicates during the SIT experiment with catalyzed reporter deposition fluorescence *in situ* hybridization (CARD-FISH) using a general archaea-targeted probe Arch915 (50) and DNA-staining (DAPI) (Fig. 1C). As the production of methane increased throughout the experiment, there was a concurrent rise in the relative cell abundance of *Ca.* M. hypatiae (Fig. 3F, SI Appendix, Table S5). The initial assessment on day 22 across four replicates revealed the total cell density to be $3.45 \times 10^7 \pm 1.14 \times 10^7$ before substantial concentrations of methane had been detected in the headspace (<132 μM). By day 32, methane concentrations reached 1,777±739 μM and the total cell density increased to $6.97 \times 10^7 \pm 3.73 \times 10^7$ cells mL^{-1} with 54% (±9.6%) of cells labeled as *Ca.* M. hypatiae (Fig. 3F). All but one of these replicates were then sacrificed for further analysis. Finally on day 45, the remaining replicate reached a headspace methane concentration of 4,109 μM and a total cell density of 1.22 $\times 10^8$ with 53% of cells labeled as *Ca.* M. hypatiae.

Visualization of the enrichment culture via scanning electron microscopy (SEM) revealed that most cells exhibited a regular to irregular coccoid morphology, with a width ranging from 0.5-1 μm (Fig. 1D). This morphology has previously been described for other Archaeoglobi species (30, 51-53).

Alternative Substrates and Temperature Optimum

We determined the substrate and temperature range of *Ca.* M. hypatiae by growing the culture in the presence of several substrates at 70 ºC or with 10 mM MMA at 60-85 ˚C (Fig. 4AB). Conditions that lead to the production of methane included 10 mM trimethylamine (TMA), 10 mM dimethylamine (DMA), 10 mM MMA in media without yeast extract, and the control with 10 mM MMA and 0.01% yeast extract.

Methane production of cultures grown with MMA in the presence or absence of yeast extract were

indistinguishable (5,202±606 and 5,703±410 µM CH_4, respectively) indicating that yeast extract is not

essential for methanogenic growth. Observed methane concentrations were higher in incubations amended

with DMA (10,115±836 µM CH_4) and TMA (9,524±3,626 µM CH_4, with a wide range of 5,361-11,993

µM) on average more than the MMA controls, consistent with what has been observed for other

methylotrophic methanogens (54). Incubations amended with 10 mM methanol (MeOH) did not produce

methane after 47 days of incubation at 70 ˚C. Due to its use by sulfate reducing organisms as an electron

donor (55), 10 mM lactate (LAC) was tested, as well as 10 mM MMA with 10 mM LAC, but none of these

incubations produced methane. Production of methane has not been observed in any attempted transfers

where hydrogen (99.9999% purity) was present in the headspace, or hydrogen with MMA was added.

The enrichment grew optimally at both 64 and 70 ˚C with relative amounts of methane produced at

5,304±451 µM and 5,202±606 µM, respectively. This deviates from the predicted optimal growth

temperature of 74.4˚C, which was derived from the translation of proteins in the Ca. M. hypatiae MAG

using Tome (56). This is lower than the observed range of growth and optimum temperatures for type

strains of non-methanogenic Archaeoglobus which have been demonstrated to grow between 50 and 95 ˚C

with optimal temperatures between 75-83 ˚C in organisms sourced predominantly from deep sea vent

environments (30). No methane production was detected at temperatures 77 ˚C or above or lower than 64

˚C after 47 days of incubation (Dataset S4).

Genomic and Transcriptomic Basis for Methanogenesis

The assembled metagenome obtained at the end of the SIT experiment was used to align a total of

23,376,154 metatranscriptome mRNA reads obtained from six replicates harvested in the exponential

growth phase and to create a detailed reconstruction of the metabolism of *Ca.* M. hypatiae (Fig. 3AB, 5,

Dataset S5). A total of 22,891,651 reads, *i.e.,* 97.8% of all recovered reads, were recruited to the *Ca.* M.

hypatiae MAG. Only 2.1% of the total mRNA reads (484,503) were aligned with other co-enriched

organisms. Among these, only 13 genes across four MAGs were expressed above 200 reads per kilobase

of transcript per million mapped reads (RPKM) and just five genes exceeded >1,000 RPKM. Genes required

for the conversion of methylamine to methane were among the top 2% of highest expressed genes

transcribed by *Ca.* M. hypatiae, including genes encoding the MCR complex (*mcrAGCDB*; 13,046-18,098

RPKM), one of three monomethylamine methyltransferase copies (*mtmB*; 9,884 RPKM), dimethylamine

corrinoid (*mtbC*; 3,677 RPKM), and methanol:coenzyme M methyltransferase (*mtaA*; 12,577 RPKM) (Fig.

5). Seven copies of substrate-specific methyltransferases for MMA (*mtmB*; 3 copies), DMA (*mtbB*; 2

copies), and TMA (*mttB*; 2 copies) were present in the genome, but methanol methyltransferase (*mtaB*) was

not identified. These genes were differentially expressed with one copy for each type of methylamine

expressed above 3,200 RPKM. In addition to *mtbC*, two gene copies of the trimethylamine corrinoid protein

(*mttC*) were found in the genome but their expression was relatively low (<460 RPKM average).

Monomethylamine corrinoid (*mtmC*) or methanol corrinoid (*mtaC*) proteins were not identified in *Ca.* M.

hypatiae. Additionally, genes were expressed for pyrrolysine synthesis (*pylBCD*; 819, 343, 37 RPKM) and

the methyltransferase corrinoid activation protein (*ramA*; 1,076 RPKM), both of which support

methylamine methyltransferases in methylotrophic methanogenesis (57, 58). The absence of *mtmC* and the

high expression levels of *mtbC* (3,677 RPKM) and *mtaA* (12,577 RPKM) suggest that they are responsible

for the transfer of a methyl group from monomethylamine to coenzyme M (CoM) after it has been

transferred by a substrate-specific methyltransferase (*mtmB*). Consistent with the observed methane

production from DMA and TMA, *Ca.* M. hypatiae can use these methylamines and expressed the

corresponding genes (*mtbB*, *mttB*) at comparatively high levels (JOOIALLP_01813 *mtbB* 3,249 RPKM;

JOOIALLP_01787 *mttB* 5,324 RPKM; Fig. 4B, 5). It is worth noting that the expression of mtbB/mttB was

detected despite the culture not having been previously exposed to DMA or TMA at the time of the

transcriptomics experiment. We hypothesize that *Ca.* M. hypatiae could employ one of two strategies: it

either (i) constitutively expresses all substrate-specific methyltransferases and corrinoid proteins as a

precautionary measure to accommodate substrates potentially encountered *in situ*, or (ii) *Ca.* M. hypatiae

transcriptionally co-regulates the genes responsible for these functions.

Ca. M. hypatiae expressed the methyl-branch of the Wood-Ljungdahl pathway (WLP) and the acetyl-

CoA decarbonylase/synthase complex (Cdh, *cdhABCDE*), which is consistent with genes observed and

shown to be expressed in sulfate-reducing Archaeoglobi genomes (55). This includes two paralogous copies

of 5,10-methylenetetrahydromethanopterin reductase (*mer*) which might function as a traditional Mer,

considering that these genes are also members of the large luciferase-like monooxygenase family

(pfam00296)(35). The expression of genes in the WLP varied. Methylenetetrahydromethanopterin

dehydrogenase (*mtd*), methenyltetrahydromethanopterin cyclohydrolase (*mch*), formylmethanofuran-

tetrahydromethanopterin N-formyltransferase (*ftr*), formylmethanofuran dehydrogenase (*fwdABC*), and

one copy of the *mer* homologs were expressed at comparatively high levels (456-2,763 RPKM), whereas

FwdDEFG and the other *mer* copy were only minimally expressed (<180 RPKM) (Dataset S5). The high

expression of the Cdh complex (*cdhACDE*; 3,063±362, *cdhB* 677 RPKM average across subunits) suggests

that *Ca.* M. hypatiae is capable of autotrophically fixing CO_2 to acetyl-CoA as has been shown for other

Archaeoglobus species (59). Acetyl-CoA could also be derived from the degradation of fatty acids present

in yeast extract through the process of beta-oxidation. Enzymes involved in this pathway were expressed at

moderate to high levels during growth (Dataset S6). Pyruvate synthase (Por) was highly expressed

providing a way for acetyl-CoA to be converted to pyruvate and subsequently be fed into major biosynthetic

pathways. Specifically, *Ca.* M. hypatiae encodes pyruvate carboxylase (PycAB), an incomplete reductive

tricarboxylic acid cycle (rTCA), phosphoenolpyruvate synthase (Pps), most enzymes needed for
gluconeogenesis, and several enzymes associated with the pentose phosphate pathway in archaea, which
were all expressed at varying levels (Dataset S5). Together, these pathways provide *Ca*. M. hypatiae the
capacity to synthesize amino acids, carbohydrates, integral components of the cell wall, and vital sugars for
nucleic acids.

Several complexes related to energy conservation and electron transport were moderately to highly
expressed. *Ca*. M. hypatiae encodes a fused heterodisulfide reductase (*hdrDE*) that was highly expressed
($1,106\pm120$ RPKM) in addition to a fused *hdrD/mvhD* and four copies of *hdrD* that were all expressed at
much lower levels (<500 RPKM). The differing levels of transcription suggest the membrane-bound HdrDE
is responsible for the regeneration of coenzymes M and B through the reduction of heterodisulfide (CoM-
S-S-CoB). Additionally, the absence of HdrB, which contains the active site for disulfide reduction,
eliminates the possibility that disulfide reduction could occur via a HdrABC complex (60). As reported for
Ca. M. nevadensis (35), a unique gene cluster was identified containing F_{420}-non-reducing hydrogenase
(MvhAGD), two HdrA copies and a QmoC fused to a HdrC. One HdrA copy (JOOIALLP_01710) was
predicted by DiSCo analysis as a quinone-modifying oxidoreductase (QmoB), a protein related to the HdrA
of methanogens (61, 62). This cluster was expressed at high levels (995-2,431 RPKM average), suggesting
its importance for electron transfer in *Ca*. M. hypatiae. We hypothesize these subunits are associating
together *in vivo* to bifurcate electrons from hydrogen (H_2) to reduce both menaquinone (MQ) and ferredoxin
(Fd_{ox}), as proposed recently (35, 63). Lastly, *Ca*. M. hypatiae moderately expressed a membrane-bound
$F_{420}H_2$:quinone oxidoreductase (Fqo) complex (88-280 RPKM across subunits) and a V-type ATP synthase
(24-442 RPKM across subunits).

The electrons required for reducing the CoM-S-S-CoB heterodisulfide could originate from two
possible routes. The first possibility would rely on sourcing electrons from hydrogen, which could be
oxidized by the Mvh-Qmo-Hdr complex coupled to menaquinone reduction. H_2 may be produced through
the activity of a group 3b [NiFe]-sulfhydrogenase (HydABDG), which was the highest expressed
hydrogenase complex with an average RPKM of 4,421 across subunits (64, 65). To evolve hydrogen via
HydABDG, reducing power, via NADPH, could be supplied by sulfide dehydrogenase (SudAB; SudA,
1,088 RPKM; SudB, 495 RPKM). Alternatively, NADPH could instead be provided to biosynthesis
pathways and therefore be decoupled from methanogenic metabolism. H_2 could also potentially be sourced
from fermentative bacteria in the enrichment culture, however, the low number of hydrogenases encoded
by co-enriched organisms were only very lowly expressed at the time of sampling for metatranscriptomics
(<51 RPKM). At this point, the source of H_2 *Ca*. M. hypatiae uses remains uncertain, as no H_2 was added
to the headspace. Second, in a hydrogen-independent electron transport system, reduced F_{420} and ferredoxin
could be produced through the dismutation of methylated substrate to CO_2 through the WLP. Reduced F_{420}

could be oxidized by the Fqo complex and contribute to a reduced menaquinone pool that could be used by

the fused HdrDE complex to reduce CoM-S-S-CoB. Reduced ferredoxin could be oxidized at a soluble

FqoF to reduce F_{420} or at an Fqo complex lacking FqoF to reduce menaquinone (66, 67). Based on the low

expression levels of the Fqo complex (171±67 RPKM) and the absence of F_{420}-reducing hydrogenase (*frh*)

from the genome, it is not likely the WLP runs in the reductive direction as a source of reduced F_{420} would

be required. Resolving the exact configuration of the electron transport system encoded by *Ca.* M. hypatiae

will require biochemical confirmation in future investigations.

Importantly, genes necessary for dissimilatory sulfate reduction typically observed in sulfate-

reducing members of the Archaeoglobi, including dissimilatory sulfite reductase (*dsrAB*), sulfate

adenylyltransferase (*sat*), and adenylylsulfate reductase (*aprAB*), were neither identified in the genome of

Ca. M. hypatiae nor in the unbinned fraction of the metagenome. They were also absent from the

comparatively incomplete MAG of *Ca.* M. nevadensis GBS (35). However, *Ca.* M. hypatiae encodes

subunits *dsrMK* and *dsrOP* of the Dsr complex in addition to *dsrC*. This complex is strictly conserved in

sulfate-reducing organisms (68) where it mediates electron transfer from the periplasm to the cytoplasm

reducing the disulfide bond found in DsrC cysteines (69). The expression of the Dsr complex and *dsrC* was

low (450±63 RPKM) during growth on monomethylamine suggesting it is not vital to the metabolism of

Ca. M. hypatiae. The presence of the Dsr complex, DsrC, and subunits QmoC and QmoB in the genome

may be explained as evolutionary remnants from ancestral Archaeoglobi, growing initially as sulfate-

reducing organisms but later transitioning to a methanogenic lifestyle (7, 8). This raises the question

whether intermediate of this process, Archaeoglobi capable of both methanogenesis and sulfate-reduction

(and possible anaerobic oxidation of methane), still exist today (25, 28).

Collectively, the metagenomic and transcriptomic data confirmed that *Ca.* M. hypatiae is not only

the sole archaeon but the sole methanogen in our culture. The metabolic reconstruction and

metatranscriptomic results are consistent with methylotrophic methanogenesis from methylamines. The

absence of genes required for sulfate reduction eliminates the possibility for this metabolism in *Ca.* M.

hypatiae. A unique gene cluster (Mvh-Qmo-Hdr) potentially involved in energy conservation was

expressed, however future studies will be required to test how *Ca.* M. hypatiae internally cycles electrons

for methanogenesis and if it sources H_2, or other reductants, from the medium or co-enriched bacteria.

Distribution of *Ca.* Methanoglobus Across Geothermal Features in YNP

16S rRNA and *mcrA* gene amplicon sequence data generated in a recent microbial diversity survey

of 100 geothermal features in YNP (33) were used to analyze the distribution of Archaeoglobi related to

Ca. M. hypatiae (SI Appendix, Fig. S5). 16S rRNA gene amplicons closely related to *Ca.* M. hypatiae

(96.7-100% sequence identity) were found in seven DNA samples from six hot springs (pH 5.1-9.35, 31-

78 ºC) in addition to hot spring LCB024 (the source of this culture) at relative abundances ranging from

0.02-0.22%. In addition, *mcrA* gene ASVs affiliated with Archaeoglobi were PCR-amplified from 53 DNA

samples, out of 201 total samples that had been screened by PCR. These 53 samples had been collected

from microbial mats or sediments originating from 36 geothermal features distributed across various

thermal regions within YNP by Lynes, Krukenberg *et al.* (33). Archaeoglobi-related *mcrA* genes were found

in geothermal features with a pH range of 2.61 to 9.32 and a temperature range of 18.4 ºC to 93.8 ºC.

Collectively, our results and the studies by Wang *et al*. and Buessecker *et al*., who reported that Mcr-

encoding Archaeoglobi are present (35) and transcriptionally active in hot spring mesocosms (34),

demonstrate the previously overlooked role that Archaeoglobi might play in the anaerobic carbon cycle of

geothermal environments.

Conclusion

In summary, the cultivation of *Ca.* Methanoglobus hypatiae LCB24 provides direct experimental

evidence that members of the Archaeoglobi are methanogens. *Ca.* M. hypatiae can use MMA, DMA, and

TMA as methanogenic substrates and grows optimally at 64-70 ºC, as evidenced by metagenomics,

metatranscriptomics, and isotope tracing experiments. Metagenomic sequencing and phylogenomic

analysis confirmed the close relationship of *Ca.* M. hypatiae to other Mcr-encoding Archaeoglobi and the

relatedness of its *mcrA* to MAGs of the TACK superphylum, some of which have recently been shown to

also be methanogens (70, 71). Together, this supports the idea that the capacity for methanogenesis is deeply

rooted in the archaea and possibly dates to the last common ancestor of archaea (1, 3, 7, 8, 72). The wide

distribution of Archaeoglobi-affiliated *mcrA* gene sequences and *Ca.* M. hypatiae-related 16S rRNA gene

sequences in geothermal features across YNP suggests that members of this lineage play a hitherto

unaccounted-for role in anaerobic carbon cycling in these extreme ecosystems. Future studies of *Ca.* M.

hypatiae and other methanogens will provide valuable insights into the evolution of methane metabolism

and the significance of these archaea in biogeochemical cycles across geothermal and other environments.

Acknowledgements

This study was funded through a NASA Exobiology program award (80NSSC19K1633) to R.H. We

thank the US National Park Service for permitting work in YNP under permit number YELL-SCI-8010.

We thank George Schaible (MSU) for help with SEM imaging, Dr. Viola Krukenberg (MSU) for initial

FISH methodology development, Sylvia Nupp, Dr. Andrew Montgomery, and Paige Schlegel (all MSU)

for assistance with field sampling, Dr. Christopher Lemon (MSU) for allowing use of his cooling centrifuge,

and Dr. Marike Palmer (UN Las Vegas) for discussing naming of this archaeon.

Author Contributions

M.M.L. and R.H. developed the research project. M.M.L., Z.J.J., A.J.K, and R.H. designed experiments. M.M.L. and A.J.K. conducted field sampling. M.M.L. performed cultivation, extracted DNA for amplicon and metagenomic sequencing, extracted RNA for transcriptomic sequencing, and conducted physiology and stable isotope experiments. A.J.K. developed GC/GCMS protocols and processed GCMS samples. Z.J.J. processed and annotated metagenomic and transcriptomic data, assembled MAGs, mapped transcripts, assigned taxonomy, constructed 16S rRNA gene phylogeny, and performed phylogenetic analysis of MAGs. M.M.L. conducted phylogenetic analysis of amplicon data, refined gene annotations, reconstructed, and interpreted the metabolic potential of *Ca.* M. hypatiae with insight from Z.J.J and A.J.K. R.H. was responsible for funding and supervision of the project. M.M.L. and R.H. wrote the manuscript, which was then edited by all authors.

Conflict of interest: none declared.

References

1. Wolfe JM, Fournier GP. Horizontal gene transfer constrains the timing of methanogen evolution. Nature Ecology & Evolution. 2018;2(5):897-903.

2. Sorokin DY, Makarova KS, Abbas B, Ferrer M, Golyshin PN, Galinski EA, et al. Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into the evolutionary origin of methanogenesis. Nature Microbiology. 2017;2(8):1-11.

3. Martin WF, Sousa FL. Early microbial evolution: the age of anaerobes. Cold Spring Harbor Perspectives in Biology. 2016;8(2):a018127.

4. Sauterey B, Charnay B, Affholder A, Mazevet S, Ferrière R. Co-evolution of primitive methane-cycling ecosystems and early Earth's atmosphere and climate. Nature Communications. 2020;11(1):2705.

5. Ueno Y, Yamada K, Yoshida N, Maruyama S, Isozaki Y. Evidence from fluid inclusions for microbial methanogenesis in the early Archaean era. Nature. 2006;440(7083):516-9.

6. Spang A, Ettema TJG. Archaeal evolution: the methanogenic roots of Archaea. Nature Microbiology. 2017;2(8):1-2.

7. Adam PS, Kolyfetis GE, Bornemann TLV, Vorgias CE, Probst AJ. Genomic remnants of ancestral methanogenesis and hydrogenotrophy in Archaea drive anaerobic carbon cycling. Science Advances. 2022;8(44):eabm9651.

8. Wang Y, Wegener G, Williams TA, Xie R, Hou J, Tian C, et al. A methylotrophic origin of methanogenesis and early divergence of anaerobic multicarbon alkane metabolism. Science Advances. 2021;7(27):eabj1453.

9. Hinrichs K-U. Microbial fixation of methane carbon at 2.7 Ga: Was an anaerobic mechanism possible? Geochemistry, Geophysics, Geosystems. 2002;3(7):1-10.

10. Saunois M, Stavert AR, Poulter B, Bousquet P, Canadell JG, Jackson RB, et al. The Global Methane Budget 2000–2017. Earth System Science Data. 2020;12(3):1561-623.

11. Rosentreter JA, Borges AV, Deemer BR, Holgerson MA, Liu S, Song C, et al. Half of global methane emissions come from highly variable aquatic ecosystem sources. Nature Geoscience. 2021;14(4):225-30.

12. Garcia PS, Gribaldo S, Borrel G. Diversity and evolution of methane-related pathways in archaea. Annual Review of Microbiology. 2022;76:727-55.

13. Ferry JG, Kastead KA. Methanogenesis. Archaea: Molecular and Cellular Biology. 2007. p. 288-314.

14. Conrad R. The global methane cycle: recent advances in understanding the microbial processes involved. Environmental Microbiology Reports. 2009;1(5):285-92.

15. Scheller S, Goenrich M, Boecher R, Thauer RK, Jaun B. The key nickel enzyme of methanogenesis
catalyses the anaerobic oxidation of methane. Nature. 2010;465(7298):606-8.

16. Thauer RK. Methyl (alkyl)-coenzyme M reductases: nickel F-430-containing enzymes involved in
anaerobic methane formation and in anaerobic oxidation of methane or of short chain alkanes.
Biochemistry. 2019;58(52):5198-220.

17. Evans PN, Boyd JA, Leu AO, Woodcroft BJ, Parks DH, Hugenholtz P, et al. An evolving view of
methane metabolism in the Archaea. Nature Reviews Microbiology. 2019;17(4):219-32.

18. Thauer RK, Kaster AK, Seedorf H, Buckel W, Hedderich R. Methanogenic archaea: ecologically
relevant differences in energy conservation. Nature Reviews Microbiology. 2008;6(8):579-91.

19. Baker BJ, De Anda V, Seitz KW, Dombrowski N, Santoro AE, Lloyd KG. Diversity, ecology and
evolution of Archaea. Nature Microbiology. 2020;5(7):887-900.

20. Bueno de Mesquita CP, Wu D, Tringe SG. Methyl-based methanogenesis: an ecological and genomic
review. Microbiology and Molecular Biology Reviews. 2023;87(1):e00024-22.

21. Söllinger A, Urich T. Methylotrophic methanogens everywhere — physiology and ecology of novel
players in global methane cycling. Biochemical Society Transactions. 2019;47(6):1895-907.

22. Nobu MK, Narihiro T, Kuroda K, Mei R, Liu WT. Chasing the elusive Euryarchaeota class WSA2:
genomes reveal a uniquely fastidious methyl-reducing methanogen. The ISME Journal.
2016;10(10):2478-87.

23. Vanwonterghem I, Evans PN, Parks DH, Jensen PD, Woodcroft BJ, Hugenholtz P, et al.
Methylotrophic methanogenesis discovered in the archaeal phylum Verstraetearchaeota. Nature
Microbiology. 2016;1(12):1-9.

24. McKay LJ, Dlakic M, Fields MW, Delmont TO, Eren AM, Jay ZJ, et al. Co-occurring genomic
capacity for anaerobic methane and dissimilatory sulfur metabolisms discovered in the Korarchaeota.
Nature Microbiology. 2019;4(4):614-22.

25. Wang Y, Wegener G, Hou J, Wang F, Xiao X. Expanding anaerobic alkane metabolism in the domain
of Archaea. Nature Microbiology. 2019;4(4):595-602.

26. Evans PN, Parks DH, Chadwick GL, Robbins SJ, Orphan VJ, Golding SD, et al. Methane metabolism
in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. Science.
2015;350(6259):434-8.

27. Berghuis BA, Yu FB, Schulz F, Blainey PC, Woyke T, Quake SR. Hydrogenotrophic methanogenesis
in archaeal phylum Verstraetearchaeota reveals the shared ancestry of all methanogens. Proceedings
of the National Academy of Sciences. 2019;116(11):5037-44.

28. Liu YF, Chen J, Zaramela LS, Wang LY, Mbadinga SM, Hou ZW, et al. Genomic and transcriptomic
evidence supports methane metabolism in Archaeoglobi. mSystems. 2020;5(2):e00651-19.

29. Stetter KO, Lauerer G, Thomm M, Neuner A. Isolation of extremely thermophilic sulfate reducers:
evidence for a novel branch of archaebacteria. Science. 1987;236(4803):822-4.

30. Slobodkina G, Allioux M, Merkel A, Cambon-Bonavita MA, Alain K, Jebbar M, et al. Physiological
and genomic characterization of a hyperthermophilic archaeon *Archaeoglobus neptunius* sp. nov.
isolated from a deep-sea hydrothermal vent warrants the reclassification of the genus *Archaeoglobus*.
Frontiers in Microbiology. 2021;12:679245.

31. Boyd JA, Jungbluth SP, Leu AO, Evans PN, Woodcroft BJ, Chadwick GL, et al. Divergent methyl-
coenzyme M reductase genes in a deep-subseafloor Archaeoglobi. The ISME Journal.
2019;13(5):1269-79.

32. Hua ZS, Wang YL, Evans PN, Qu YN, Goh KM, Rao YZ, et al. Insights into the ecological roles and
evolution of methyl-coenzyme M reductase-containing hot spring Archaea. Nature Communications.
2019;10(1):4574.

33. Lynes MM, Krukenberg V, Jay ZJ, Kohtz AJ, Gobrogge CA, Spietz RL, et al. Diversity and function
of methyl-coenzyme M reductase-encoding archaea in Yellowstone hot springs revealed by
metagenomics and mesocosm experiments. ISME Communications. 2023;3(1):22.

34. Wang J, Qu Y-N, Evans PN, Guo Q, Zhou F, Nie M, et al. Evidence for nontraditional *mcr*-containing
archaea contributing to biological methanogenesis in geothermal springs. Science Advances.
2023;9(26):eadg6004.

35. Buessecker S, Chadwick GL, Quan ME, Hedlund BP, Dodsworth JA, Dekas AE. Mcr-dependent
methanogenesis in Archaeoglobaceae enriched from a terrestrial hot spring. The ISME Journal.
2023;17(10):1649-59.

36. Laso-Perez R, Krukenberg V, Musat F, Wegener G. Establishing anaerobic hydrocarbon-degrading
enrichment cultures of microorganisms under strictly anoxic conditions. Nature Protocols.
2018;13(6):1310-30.

37. Brandis A, Thauer RK. Growth of *Desulfovibrio* species on hydrogen and sulphate as sole energy
source. Microbiology. 1981;126(1):249-52.

38. Ai G, Zhu J, Dong X, Sun T. Simultaneous characterization of methane and carbon dioxide produced
by cultured methanogens using gas chromatography/isotope ratio mass spectrometry and gas
chromatography/mass spectrometry. Rapid Communications in Mass Spectrometry.
2013;27(17):1935-44.

39. Bushnell B. BBMap: a fast, accurate, splice-aware aligner. Lawrence Berkeley National Lab (LBNL),
Berkeley, CA (United States); 2014 Mar 17.

40. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30(14):2068-9.

41. Wu Y-W, Tang Y-H, Tringe SG, Simmons BA, Singer SW. MaxBin: an automated binning method
to recover individual genomes from metagenomes using an expectation-maximization algorithm.
Microbiome. 2014;2:26.

42. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately reconstructing single
genomes from complex microbial communities. PeerJ. 2015;3:e1165.

43. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning metagenomic
contigs by coverage and composition. Nature Methods. 2014;11(11):1144-6.

44. Miller IJ, Rees ER, Ross J, Miller I, Baxa J, Lopera J, et al. Autometa: automated extraction of
microbial genomes from individual shotgun metagenomes. Nucleic Acids Research. 2019;47(10):e57-
e.

45. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of genomes from
metagenomes via a dereplication, aggregation and scoring strategy. Nature Microbiology.
2018;3(7):836-43.

46. Kohtz AJ, Jay ZJ, Lynes MM, Krukenberg V, Hatzenpichler R. Culexarchaeia, a novel archaeal class
of anaerobic generalists inhabiting geothermal environments. ISME Communications. 2022;2(1):86.

47. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of
microbial genomes recovered from isolates, single cells, and metagenomes. Genome Research.
2015;25(7):1043-55.

48. Andrews S, Krueger F, Seconds-Pichon A, Biggins F, Wingett S. FastQC: A quality control tool for
high throughput sequence data. Babraham Bioinformatics: Babraham Institute; 2015.

49. Deng Z-L, Münch PC, Mreches R, McHardy AC. Rapid and accurate identification of ribosomal RNA
sequences via deep learning. Nucleic Acids Research. 2022;50(10):e60-e.

50. Stahl DA. Development and application of nucleic acid probes in bacterial systematics. Sequencing
and hybridization techniques in bacterial systematics. 1991:205-48.

51. Stetter KO. *Archaeoglobus fulgidus* gen. nov., sp. nov.: a new taxon of extremely thermophilic
archaebacteria. Systematic and Applied Microbiology. 1988;10:172-3.

52. Huber H, Jannasch H, Rachel R, Fuchs T, Stetter KO. *Archaeoglobus veneficus* sp. nov., a novel
facultative chemolithoautotrophic hyperthermophilic sulfite reducer, isolated from abyssal black
smokers. Systematic and Applied Microbiology. 1997;20(3):374-80.

53. Mori K, Maruyama A, Urabe T, Suzuki K-i, Hanada S. *Archaeoglobus infectus* sp. nov., a novel
thermophilic, chemolithoheterotrophic archaeon isolated from a deep-sea rock collected at Suiyo
Seamount, Izu-Bonin Arc, western Pacific Ocean. International Journal of Systematic and
Evolutionary Microbiology. 2008;58(4):810-6.

54. Watkins AJ, Roussel EG, Webster G, Parkes RJ, Sass H. Choline and N,N-dimethylethanolamine as
direct substrates for methanogens. Applied and Environmental Microbiology. 2012;78(23):8298-303.
55. Hocking WP, Stokke R, Roalkvam I, Steen IH. Identification of key components in the energy
metabolism of the hyperthermophilic sulfate-reducing archaeon *Archaeoglobus fulgidus* by
transcriptome analyses. Frontiers in Microbiology. 2014;5:95.
56. Li G, Rabe KS, Nielsen J, Engqvist MK. Machine learning applied to predicting microorganism
growth temperatures and enzyme catalytic optima. ACS Synthetic Biology. 2019;8(6):1411-20.
57. Mahapatra A, Patel A, Soares JA, Larue RC, Zhang JK, Metcalf WW, et al. Characterization of a
Methanosarcina acetivorans mutant unable to translate UAG as pyrrolysine. Molecular Microbiology.
2006;59(1):56-66.
58. Ferguson T, Soares JA, Lienard T, Gottschalk G, Krzycki JA. RamA, a protein required for reductive
activation of corrinoid-dependent methylamine methyltransferase reactions in methanogenic archaea.
Journal of Biological Chemistry. 2009;284(4):2285-95.
59. Estelmann S, Ramos-Vera WH, Gad'on N, Huber H, Berg IA, Fuchs G. Carbon dioxide fixation in
'*Archaeoglobus lithotrophicus*': are there multiple autotrophic pathways? FEMS Microbiology
Letters. 2011;319(1):65-72.
60. Ferry JG. How to make a living by exhaling methane. Annual Review of Microbiology. 2010;64:453-
73.
61. Chernyh NA, Neukirchen S, Frolov EN, Sousa FL, Miroshnichenko ML, Merkel AY, et al.
Dissimilatory sulfate reduction in the archaeon '*Candidatus* Vulcanisaeta moutnovskia' sheds light on
the evolution of sulfur metabolism. Nature Microbiology. 2020;5(11):1428-38.
62. Neukirchen S, Sousa FL. DiSCo: a sequence-based type-specific predictor of Dsr-dependent
dissimilatory sulphur metabolism in microbial data. Microbial Genomics. 2021;7(7).
63. Ramos AR, Keller KL, Wall JD, Pereira IAC. The membrane QmoABC complex interacts directly
with the dissimilatory adenosine 5′-phosphosulfate reductase in sulfate reducing bacteria. Frontiers in
Microbiology. 2012;3:137.
64. Adams MWW. The metabolism of hydrogen by extremely thermophilic, sulfur-dependent bacteria.
FEMS Microbiology Reviews. 1990;6(2-3):219-37.
65. Ma K, Schicho RN, Kelly RM, Adams M. Hydrogenase of the hyperthermophile *Pyrococcus furiosus*
is an elemental sulfur reductase or sulfhydrogenase: evidence for a sulfur-reducing hydrogenase
ancestor. Proceedings of the National Academy of Sciences. 1993;90(11):5341-4.
66. Hocking WP, Roalkvam I, Magnussen C, Stokke R, Steen IH. Assessment of the carbon monoxide
metabolism of the hyperthermophilic sulfate-reducing archaeon *Archaeoglobus fulgidus* VC-16 by
comparative transcriptome analyses. Archaea. 2015;2015.

67. Welte C, Deppenmeier U. Membrane-bound electron transport in *Methanosaeta thermophila*. Journal
of Bacteriology. 2011;193(11):2868-70.

68. Pereira IA, Haveman SA, Voordouw G. Biochemical, genetic and genomic characterization of
anaerobic electron transport pathways in sulphate-reducing delta-proteobacteria. Sulphate-Reducing
Bacteria: Environmental and Engineered Systems (LL, B and WA, H, eds), Cambridge University
Press, Cambridge, UK. 2007.

69. Grein F, Pereira IA, Dahl C. Biochemical characterization of individual components of the
Allochromatium vinosum DsrMKJOP transmembrane complex aids understanding of complex
function in vivo. Journal of Bacteriology. 2010;192(24):6369-77.

70. Wu K, Zhou L, Tahon G, Liu L, Li J, Zhang J, et al. Isolation of a methyl-reducing methanogen outside
the Euryarchaeota. In revision (https://doi.org/10.21203/rs.3.rs-2501667/v1).

71. Kohtz A, Krukenberg V, Petrosian N, Jay Z, Pilhofer M, Hatzenpichler R. Cultivation and
visualization of a methanogen of the phylum Thermoproteota. In revision
(https://doi.org/10.21203/rs.3.rs-2500102/v1)

72. Spang A, Caceres EF, Ettema TJG. Genomic exploration of the diversity, ecology, and evolution of
the archaeal domain of life. Science. 2017;357(6351):eaaf3883.

 Figure legends

 Fig. 1. Community composition and methane production of the methanogenic enrichment culture

 containing *Ca.* Methanoglobus hypatiae LCB24. *(A)* Relative abundance of 16S rRNA gene amplicons in

 the initial sediment from hot spring LCB024, the slurry collected in November 2021, slurry material used

 to initiate enrichments in February 2022, the initial enrichment, and five subsequent transfers (T1-T5) are

 shown. For comparison, the estimated relative abundance of two metagenomic samples (T4-MG and SIT-

 MG) is included. The metagenome recovered from a replicate from the stable isotope tracing experiment

 incubated in the presence of deuterated methylamine (SIT-MG) revealed *Ca.* M. hypatiae grew to 92.8%

 relative abundance during the experiment. The two most abundant ASVs across enrichment transfers are

 shown with other taxa collapsed. Other methanogenic archaea were not identified in the initial enrichment

 or in any subsequent transfers. Relative sequence abundance for all ASVs is reported in SI Appendix, Table

 S1. *(B)* Headspace methane produced over long-term cultivation. The time between transfers decreased

 while the average maximum concentration of methane increased over time. Culture 1A represents the initial

 enrichment. A history of methane measurements can be found in SI Appendix, Table S2. *(C)* Visualization

 of *Ca.* M. hypatiae cells at T6 labeled via CARD-FISH by the general archaea probe Arch915 (red). DAPI

 staining of cells is in blue. *(D)* Cell morphologies in enrichment culture LCB24 at T7 as observed by

 scanning electron microscopy.

 Fig. 2. Phylogenetic affiliation of *Ca.* M. hypatiae LCB24. *(A)* Maximum-likelihood tree, inferred with

 fasttree and WAG model (midpoint root), using a concatenated alignment of 33 conserved single copy

 proteins (list provided in SI Appendix, Table S4). References are colored by the habitat or type from which

 sequences had been recovered: hot springs in Yellowstone National Park (YNP), yellow; Great Boiling

 Spring (GBS), green; hot springs in China, blue; isolates from marine hydrothermal vent systems, dark

 blue. *(B)* Maximum-likelihood tree inferred with fasttree using 16S rRNA genes with length in base pairs

 (bp). *(C)* ANI and AAI analysis of reference Archaeoglobales MAGs and genomes. Asterisks (*) indicate

 MAGs containing *mcrA*, apart from the MAG of *Ca.* M. hydrogenotrophicum bin74 which encodes a *mcrA*

 that is interrupted by a stop codon. AAI and ANI values are provided in SI Appendix, Fig. S1. *(D)*

 Maximum-likelihood tree, inferred with IQtree2 and the LG+C60+F+G model, from the amino acid

 alignment of McrA. Dashed lines indicate McrA/AcrA groups: (I) McrA from methanogens and ANME

 (MCR-type), (II) McrA from TACK lineages (MCR-type), (III) McrA-like from proposed and

 experimentally confirmed alkane oxidizing archaea (ACR-type). Insert shows MAGs closely related to *Ca.*

 M. hypatiae LCB24.

Fig. 3. Conversion of stable isotope labeled monomethylamine to methane by culture LCB24. *(A)*

Production of $^{13}CH_4$ in cultures amended with $^{13}CH_3$-NH_2 vs. $^{12}CH_3$-NH_2 (6 replicates). *(B)* Production of

CD_3H in cultures amended with CD_3-NH_2 (6 replicates). *(C)* Production of $^{13}CO_2$ in cultures amended with

$^{13}CH_3$-NH_2 vs. $^{12}CH_3$-NH_2. For plots A and B, ten total replicates across treatments were sacrificed during

mid-exponential phase for metagenomic or metatranscriptomic sequencing indicated by red arrows. $^{13}CH_4$,

CD_3H, or $^{13}CO_2$ production for the replicate allowed to reach stationary phase is shown as a dashed line

through open diamond symbols. *(D)* Production of $^{12}CH_4$ in cultures amended with ^{12}C-monomethylamine

($^{12}CH_3$-NH_2; 2 replicates). Cultures incubated without substrates (2 replicates) and those to which the

inhibitor BES was added on day 0 (3 replicates) did not produce $^{12}CH_4$ over the course of the experiment.

(E) Production of $^{12}CH_4$ in cultures amended with $^{12}CH_3$-NH_2 to which BES was added on day 33 of

incubation (black arrow; 3 replicates). The average production of $^{12}CH_4$ leveled off and ceased after the

introduction of BES, indicating methane generation by *Ca.* M. hypatiae is MCR-dependent. Error bars

indicate standard deviation of biological replicates when applicable. Measurements of $^{12}CH_4$, $^{13}CH_4$, CD_3H,

and $^{13}CO_2$ for all replicates and controls are reported in Dataset S2. $^{12}CH_4$ measurements for all controls

and replicates are shown in SI Appendix, Fig. S4 and Dataset S3. *(F)* $^{12}CH_4$ production and fraction of *Ca.*

M. hypatiae cells in biological replicates incubated with $^{13}CH_3$-NH_2. Relative abundance of cells was

determined at three time points (day 22, 32, 45) based on the fraction of *Ca.* M. hypatiae specific CARD-

FISH counts (orange) versus total counts of DAPI stained cells (blue). Error bars indicate the standard

deviation for four biological replicates on days 22 and 32.

Fig. 4. Temperature and substrate range of culture LCB24. *(A)* Methane production from MMA was

observed between 64-70 ˚C. *(B)* Substrate range. Methane production was observed for MMA, DMA,

TMA, and in media prepared without yeast extract. LAC, lactate; MeOH, methanol. Both experiments

performed in triplicate. All measurements can be found in Dataset S4.

Fig. 5. Transcriptional activity in *Ca.* M. hypatiae grown under methanogenic conditions (N_2 headspace,

10 mM MMA, and 0.01% yeast extract). Transcriptionally active proteins are shown in bold black font.

Proteins not encoded in the MAG are colored in white and denoted in bold red font. Average reads per

kilobase of transcript per million mapped reads (RPKM) values of six biological replicates are depicted.

RPKM values are represented by boxes or colored subunits close to each protein and are colored according

to their expression level with the RPKM value of the lowest expressed gene depicted, 37 RPKM. For

enzymes comprising multiple subunits, the beta-oxidation pathway, and the TCA cycle, an average RPKM

value representing the transcribed enzymes is used. *Ca.* M. hypatiae is transcriptionally active under

methanogenic conditions and encodes the ability to convert methyl-groups from mono-, di-, and

trimethylamine to methane. This ability is enabled by several copies of substrate-specific

methyltransferases and corrinoid proteins highlighted in the box to the bottom right. A complete list of

genes described in this figure, their transcription levels, and their abbreviations is provided in Dataset S5.

Supporting Information

SI Results and Discussion

Protologue

Methanoglobus hypatiae sp. nov.

Me.tha.no.glo.bus. Gr. pref. *methano-*, pertaining to methane; L. masc. n. -*globus*, sphere; Gr.L.
masc. n. *Methanoglobus*, methane producing organism spherical in shape. This genus was named by
Buessecker *et al.* (1). Hy.pa.ti.ae. Gr. fem. hypatiae, to honor Hypatia of Alexandria, a respected and
renowned philosopher of ancient Alexandria, Egypt, who made significant contributions to the
understanding of mathematics and astronomy. A symbol of intellectual courage and scholarly achievement.
This archaeon was cultured from an unnamed hot spring in the Lower Culex Basin of Yellowstone National
Park identified as feature LCB024 (2). This archaeon is an obligately anaerobic thermophile that performs
methylotrophic methanogenesis using methylamines and grows as regular to irregular coccoid cells
approximately 0.5 to 1 μm in width. The type genome of this archaeon is deposited at NCBI under
BioProject PRJNA1014417, accession number will be added upon publication.

SI Materials and Methods

Amplicon Sequencing and Analysis

DNA was extracted from environmental slurry samples and enrichment cultures sampled on the day
of transfer using the FastDNA Spin Kit for Soil (MP Biomedicals, Irvine, CA) following the manufacturer's
guidelines. Archaeal and bacterial 16S rRNA genes were amplified with the updated Earth Microbiome
Project primer set 515F and 806R (3). Amplicon libraries were prepared as previously described (2) and
sequenced by the Molecular Research Core Facility at Idaho State University (Pocatello, ID) using an
Illumina MiSeq platform with 2 x 250 bp paired end read chemistry. Gene reads were processed using
QIIME 2 version 2022.8 (4). Primer sequences were removed from demultiplexed reads using cutadapt (5)
with error rate 0.12 and reads truncated (130 bp forward, 150 bp reverse), filtered, denoised and merged in
DADA2 with default settings (6). Processed 16S rRNA gene amplicon sequence variants (ASVs) were
taxonomically classified with the sklearn method and the SILVA 138 database (7). The R package
decontam (version 1.18.0) (8) was used to remove contaminants using the "Prevalence" model with a
threshold of 0.5.

Annotation and Reconstruction of Metabolic Potential

Genes associated with methanogenesis pathways, dissimilatory sulfur metabolism pathways,
coenzyme and cofactor biosynthesis, energy conservation, and beta-oxidation, were inventoried.
Annotations assigned by Prokka were refined through manual evaluation using KofamKOALA, NCBI
BLASTP, NCBI's Conserved Domain Database, InterPro, the hydrogenase classifier HydDB, and DiSCo
(9-14).

Phylogenetic and Phylogenomic Analyses

Average nucleotide identities (ANI) of 16S rRNA genes were calculated with blastn, with ANI and
average amino acid identities (AAI) calculated by pyani v02.2.12 (ANIb) and CompareM v0.0.23 (--
fragLen 2000) (https://github.com/dparks1134/CompareM), respectively for selected Archaeoglobales
genomes and MAGs (Table 1). Phylogenetic analysis of 16S rRNA genes was performed with fasttree (15)
using masked alignments generated by ssu-align.

Archaeoglobales MAGs and reference genomes were screened for 54 phylogenetically informative
single copy proteins (16, 17) of which a subset of 33 proteins were identified in them all (SI Appendix,
Table S4). In order to maximize the number of proteins compared across references, MAGs LMO1 and
LMO3 were excluded from this analysis, as they lacked 2 and 7 proteins out of the total 33, respectively.
These were then aligned with muscle (18), concatenated, and phylogenomically analyzed with maximum
likelihood analysis with fasttree (WAG model). McrA alignments were performed with MAFFT-LINSi
v7.522 (19), trimmed with trimAL v1.4.rev22 (20) using a 0.5 gap threshold, and maximum likelihood trees
were built with IQTree2 v2.0.6 (21) using LG+C60+F+G model and 1,000 ultrafast bootstraps.
53
Temperature and Substrate Optimum Experiments
Methane production and growth of Archaeoglobus was evaluated at different temperatures and in the
presence of methylated substrates (i.e., methanol and mono-, di-, and trimethylamine), lactate, and media
prepared without yeast extract. The sixth transfer of the enrichment was used to inoculate triplicate 30 mL
serum bottles containing 15 mL of medium with 8% v/v inoculum, streptomycin (50 mg/L), vancomycin
(50 mg/L), and 10 mM of each substrate tested. Cultures were evaluated at 60 ˚C, 64 ˚C, 70 ˚C, 77 ˚C, 80
˚C, and 85 ˚C with 10 mM MMA. Separately, we tested whether the culture would grow on the following
substrate (combinations): 10 mM dimethylamine (DMA); 10 mM trimethylamine (TMA); 10 mM methanol
(MeOH); 10 mM lactate (LAC); 10 mM MMA and 10 mM LAC); 10 mM MMA with media without yeast
extract; and a control in media without yeast or any methanogenic substrate. The 70 ˚C cultures amended
with 10 mM MMA served as the control. All incubations were performed in biological triplicate.
65
Methane Measurements
During cultivation, 250 µL subsamples of the headspace were taken using a gas tight syringe
(Hamilton) and injected into a 10 mL autosampler vial that had been sealed with grey chlorobutyl septa.
Samples were taken from the autosampler vials and injected into a Shimadzu 2020-GC gas chromatograph
equipped with a GS-CarbonPLOT column (30 m x 0.32 mm; 1.5 μm film thickness; Agilent) and a Rt-Q-
BOND column (30 m x 0.32 mm; 1.5 μm film thickness; Restek) using helium as a carrier gas. All injections
were performed by a Shimadzu AOC-6000 autosampler robot. The injector, column, and flame ionization
detector (FID) were maintained at 200 ˚C, 50 ˚C, and 240 ˚C, respectively. Methane concentrations were
calculated based on injection of a standard curve.
75
Fluorescence *in situ* hybridization and cell counts
Aliquots of enrichment cultures incubated with ^{13}C-MMA during the SIT experiment were treated
with 2% paraformaldehyde (PFA) and fixed for 1 hr at room temperature. Following fixation, cells were
washed twice with 1x PBS, followed by centrifugation at $16,000 \times g$ to remove the supernatant, resuspended
in 1x PBS, and stored at 4 ˚C. For direct cell counts, aliquots of fixed cell suspensions were filtered onto
polycarbonate filters (0.2 μm pore size, 25 mm diameter, GTTP Millipore, Germany) and air dried before
filter pieces were cut and embedded in 0.2% low melting agarose. We attempted to use the
Archaeoglobales-specific probe Arglo32 (22), however fluorescent signal was insufficient. Given *Ca.* M.
hypatiae was the sole archaeon in the enrichment culture, the relative abundance of *Ca.* M. hypatiae cells
was determined via catalyzed reporter deposition fluorescence *in situ* hybridization (CARD-FISH) using
the general archaea-targeted 16S rRNA oligonucleotide probe Arch915 (23). Total cell counts were based
on DNA-stained cells using DAPI (4,6-diamidino-2-phenylindole). CARD-FISH was performed as
previously described (24). Cell wall permeabilization was achieved with a brief treatment of 0.1 M HCl (1

min, RT) followed by treatment with 0.01 M HCl (15 min, RT). Endogenous peroxidases were inactivated
with 0.15% H_2O_2 in methanol (30 min, RT). A formamide concentration of 35% was used for all
hybridization reactions (2.5 hrs, 46˚C). CARD was performed using Alexa Fluor 594 labeled tyramides for
30 min at 46˚C. Following signal amplification, an additional washing step in 1x PBS was included to
reduce background fluorescence (15 min, RT, dark). Samples were stained with DAPI, embedded in
Citifluor-Vectashield, and enumerated using an epifluorescence microscope (Leica DM4B).

Scanning electron microscopy (SEM)
An aliquot of the enrichment culture at transfer 7 (T7) was treated with 2% paraformaldehyde (PFA) and
fixed for 1 hr at room temperature. Following fixation, cells were washed twice through centrifugation at
16,000 × g to remove the supernatant, resuspended in 1x phosphate buffered saline (PBS), and stored at 4
˚C. Samples for imaging were prepared according to Schaible et al., 2022 (25). Briefly, a square coupon
of mirror-finished 304 stainless steel (25 mm diameter, 0.6 mm thickness) was purchased from Stainless
Supply (Monroe, NC). The coupon was cleaned by washing with a 1% solution of Tergazyme (Alconox,
New York, NY) and rinsed with Milli-Q water. The coupon was dried under compressed air and stored at
room temperature. 5 µL of fixed sample was spotted on the coupon and air-dried at 46 °C for 3 min. The
coupon was then dried for 1 m each step in a successive ethanol series starting with 10% ethanol and
increasing by increments of 10% with the last step 90% ethanol. SEM images were captured using a Zeiss
(Jena, Germany) SUPRA 55VP field emission scanning electron microscope (FE-SEM). The microscope
was operated at 1 keV under a vacuum of 0.2–0.3 mPa, with a working distance of 5.4-6.2 mm at the
Imagining and Chemical Analysis Laboratory (ICAL) of Montana State University (Bozeman, MT). No
conductivity coating was applied before SEM analysis as the microscope was operated at 1 keV.

SI Tables

Table S1. Extended community composition history of methanogenic enrichment cultures via estimated
relative abundance (%) from 16S rRNA gene amplicon sequencing.

	LCB024 sediment	Slurry 11/2021	Slurry 02/2022	Initial Enrichment	T1	T2	T3	T4	T4 - MG	T5	SIT-MG
Archaeoglobaceae ASV_78ad2	0.46	0.32	0.02	6.42	74.8	68.9	48.5	46.0	84.7	62.2	92.8
Pseudothermotoga ASV_17231	2.14	1.06	1.08	18.93	6.8	17.1	28.5	37.4	14.5	15.7	3.2
Other Archaeoglobaceae uncultured	0.00	0.00	0.00	1.54	0.0	0.01	0.0	0.0	0.0	0.0	0.0
Confirmed methanogenic archaea	0.25	0.38	0.45	0.00	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other Archaea	13.51	11.87	10.49	2.67	0.0	0.0	0.0	0.04	0.0	0.0	0.0
Other Bacteria	83.63	86.37	87.95	70.44	18.4	14.1	23.1	16.5	0.8	22.1	4.0

117   **Table S2.** Methane production of the enrichment culture over time. T, transfer.

Culture ID	Days continuous	Days	CH$_4$ (µM)	CH$_4$ (mM)	CH$_4$ (ppm)	CH$_4$ (%)
1A – initial enrichment	6	6	4.91	0.00	118.67	0.01
	13	13	16.73	0.02	404.73	0.04
	20	20	12.98	0.01	314.13	0.03
	27	27	12.86	0.01	311.20	0.03
	36	36	32.92	0.03	796.48	0.08
	48	48	146.28	0.15	3538.80	0.35
	53	53	279.15	0.28	6753.17	0.68
	63	63	419.11	0.42	10139.33	1.01
	70	70	447.12	0.45	10816.36	1.08
T1	82	12	14.40	0.01	348.40	0.03
	91	21	50.85	0.05	1230.06	0.12
	98	27	198.41	0.20	4800.09	0.48
	106	35	390.47	0.39	9445.97	0.94
	113	42	565.32	0.57	13676.04	1.37
	117	46	688.44	0.69	16654.06	1.67
	126	55	1442.03	1.44	34885.74	3.49
	133	62	1844.13	1.84	44611.75	4.46
T2	146	7	15.22	0.02	368.32	0.04
	158	18	16.11	0.02	389.83	0.04
	175	34	49.40	0.05	1195.06	0.12
	197	57	680.54	0.68	16463.41	1.65
	221	81	2200.24	2.20	53227.90	5.32
	229	89	2654.34	2.65	64213.66	6.42
	238	98	2230.72	2.23	53964.66	5.40
T3	243	5	18.17	0.02	439.53	0.04
	259	21	10.70	0.01	258.82	0.03
	266	28	536.13	0.54	12969.71	1.30
	274	36	2413.23	2.41	58379.47	5.84
	280	42	2283.95	2.28	55253.34	5.53
	284	46	2193.96	2.19	53077.04	5.31
T4	294	10	17.36	0.02	420.07	0.04
	301	17	18.12	0.02	438.37	0.04
	309	25	12.51	0.01	302.54	0.03
	319	35	1944.01	1.94	47028.08	4.70
	327	43	3054.00	3.05	73881.94	7.39
	335	51	2619.45	2.62	63370.12	6.34
T5	347	10	13.41	0.01	324.42	0.03
	355	18	13.14	0.01	317.78	0.03
	365	28	137.59	0.14	3328.60	0.33
	371	34	1023.35	1.02	24756.27	2.48
	375	38	2218.75	2.22	53674.81	5.37
	384	46	2459.29	2.46	59493.85	5.95
T6	386	2	16.29	0.02	394.19	0.04
	393	9	21.33	0.02	516.14	0.05
	400	16	18.68	0.02	451.92	0.05
	407	23	245.86	0.25	5947.76	0.59
	412	28	2555.27	2.56	61815.61	6.18
	414	30	3432.93	3.43	83051.24	8.31
	416	32	3621.90	3.62	87623.79	8.76
	418	34	3942.97	3.94	95385.88	9.54
T7	428	9	34.33	0.03	830.38	0.08
	435	16	296.91	0.30	7182.55	0.72
	440	21	2931.05	2.93	70907.06	7.09
	444	25	3164.45	3.16	76553.52	7.66
	446	27	3338.04	3.34	80751.55	8.08

Table S3. Extended Archaeoglobales metagenome assembled genome and isolate genome statistics. A combined assembly of metagenomes from T4-MG and SIT-MG (Fig. 1A) for the *Ca.* M. hypatiae LCB24 MAG was used as it yielded an improved assembly. GTDB classified the YNP, GBS, and China MAGs as: d_Archaea;p_Halobacteriota;c_Archaeoglobi;o_Archaeoglobales; f_Archaeoglobaceae;g_WYZ-LMO2;s_WYZ-LMO2. Len., length; Compl., completeness; Redun, redundancy; Strain Hetero., strain heterogeneity; pOGT, predicted optimal growth temperature. [*] stop codon interrupts *mcrA* sequence; [a] Both sequences 5ʹ start; not identical; [b] Consists of 1 chromosome and 1 plasmid.

	Seqs	Len (Mb)	GC (%)	Compl. (%)	Redun. (%)	16S	tRNA	CDS	*mcrA*	CRISPRs	pOGT (°C)	Citation
Ca. **M. hypatiae LCB24**	19	1.624	46.12	100	1.31	1	44	1,760	1	2	74.44	This study
Archaeoglobales LCB024-003	179	1.157	45.72	88.48	1.31	0	31	1,226	1	1	73.90	Lynes *et al.* 2023
Archaeoglobales WYZ-LMO2	220	1.514	45.92	97.60	0	0	40	1,614	1	2	72.40	Wang *et al.* 2019
Ca. M. hydrogenotrophicum bin74	127	1.547	45.63	92.81	1.31	1	34	1,679	0[*]	9	75.32	Liu *et al.* 2020
Archaeoglobales SJ3.Bin34	73	1.468	46.00	98.69	0.65	0	39	1,581	1	2	73.13	Colman *et al.* 2019
Ca. Methanoglobus nevadensis GBS	71	1.602	47.16	98.04	3.66	1	47	1,785	1	2	74.75	Peacock *et al.* 2013
Archaeoglobales GMQP bin32	20	1.727	41.26	100	0.98	1	39	1,931	1	2	71.88	Hua *et al.* 2019
Archaeoglobales GMQP_D bin 18	22	1.565	42.01	97.39	0.98	1	45	1,747	1	3	69.41	Wang *et al.* 2023
Archaeoglobales JZ-3 D bin 138	32	1.550	42.08	99.35	1.31	1	43	1,701	1	2	69.11	Wang *et al.* 2023
Archaeoglobales WYZ-LMO1	140	1.557	43.85	88.89	1.31	1	41	1,701	1	1	72.72	Wang *et al.* 2019
Archaeoglobales WYZ-LMO3	135	1.568	43.93	88.03	1.96	2[a]	35	1,670	1	2	73.09	Wang *et al.* 2019
Ca. M. hydrogenotrophicum bin11	252	1.220	47.59	91.83	6.17	1	27	1,387	1	0	72.64	Liu *et al.* 2020
Ca. M. hydrogenotrophicum bin16	46	1.668	45.49	96.51	1.31	0	41	1,864	1	2	72.69	Liu *et al.* 2020
Archaeoglobus JZ bin24	35	1.492	44.66	99.35	1.31	0	44	1,642	0	2	68.52	unpublished
Archaeoglobales LCB024-002	66	1.334	42.28	97.39	0.03	0	41	1,459	0	0	73.46	Lynes *et al.* 2023
Archaeoglobales LCB003-04	128	1.335	42.27	96.73	1.96	0	38	1,447	0	0	73.71	Lynes *et al.* 2023
Archaeoglobales SJ3.bin61	45	1.270	41.68	89.54	0	1	40	1,392	0	0	73.34	Colman *et al.* 2019
Archaeoglobales JZ_75 SW bin109	16	1.562	39.48	99.35	0.65	1	45	1,762	1	5	74.34	Wang *et al.* 2023
Archaeoglobus fulgidus DSM 4304	1	2.178	48.58	100	0	1	46	2,440	0	3	79.88	Klenk *et al.* 1997
Archaeoglobus neptunius SE56	32	2.116	46.05	100	0	1	47	2,336	0	2	78.26	Slobodkina *et al.* 2021
Archaeoglobus profundus DSM 5631	2[b]	1.563	46.05	100	0	1	48	1,784	0	0	85.86	von Jan *et al.* 2010
Ferroglobus placidus DSM 10642	1	2.196	44.14	100	0	1	49	2,467	0	1	83.83	Anderson *et al.* 2011
Geoglobus acetivorans SBH6	1	1.861	46.84	99.84	0	1	48	2,168	0	6	78.22	Mardanov *et al.* 2015
Geoglobus ahangari 234	1	1.770	53.11	100	0	1	46	1,985	0	7	84.11	Manzella *et al.* 2015
Archaeoglobus sulfaticallidus PM70-1	1	2.077	43.24	100	0	1	51	2,237	0	1	76.12	Stokke *et al.* 2013
Archaeoglobus veneficus SNP6	1	1.902	47.05	99.35	0	1	46	2,055	0	2	76.18	Mukherjee *et al.* 2017
Ca. Polytropus marinifundus rG16	21	2.129	40.22	99.84	1.96	1	44	2,286	2	0	66.39	Boyd *et al.* 2019

122   **Table S4.** Conserved single copy proteins used in phylogenomic analysis of MAGs and isolates.

arCOG	Gene	Product
arCOG00405	GRS1	Glycyl-tRNA synthetase (class II)
arCOG00779	RplO	Ribosomal protein L15
arCOG00785	RpmC	Ribosomal protein L29
arCOG01001	Map	Methionine aminopeptidase
arCOG01183	Kae1p/TsaD	Subunit of KEOPS complex, contains a domain with ASKHA fold and RIO-type kinase (AP-endonuclease activity)
arCOG01228	Ffh	Signal recognition particle GTPase
arCOG01722	RpsM/rps13p	Ribosomal protein S13
arCOG01758	RpsJ/rps10p	Ribosomal protein S10
arCOG04070	RplC	Ribosomal protein L3
arCOG04071	RplD	Ribosomal protein L4
arCOG04072	RplW	Ribosomal protein L23
arCOG04086	RpmD	Ribosomal protein L30
arCOG04087	RpsE	Ribosomal protein S5
arCOG04088	RplR	Ribosomal protein L18
arCOG04090	RplF/rpl6p	Ribosomal protein L6P
arCOG04091	RpsH/rps8p	Ribosomal protein S8
arCOG04094	RplX/rpl24p	Ribosomal protein L24
arCOG04095	RplN/rps14p	Ribosomal protein L14
arCOG04096	RpsQ/rps17p	Ribosomal protein S17
arCOG04097	RpsC/rps3p	Ribosomal protein S3
arCOG04098	RplV/rpl22p	Ribosomal protein L22
arCOG04113	RplP	Ribosomal protein L10AE/L16
arCOG04121	RnhB	Ribonuclease HII
arCOG04169	SecY	Preprotein translocase subunit SecY
arCOG04185	RpsO	Ribosomal protein S15P
arCOG04239	RpsD/rps4p	Ribosomal protein S4 or related protein
arCOG04242	RplM/rpl13p	Ribosomal protein L13
arCOG04243	RpsI/rps9p	Ribosomal protein S9
arCOG04245	RpsB/rps2p	Ribosomal protein S2
arCOG04255	RpsL/rps12p	Ribosomal protein S12
arCOG04256	RpoC/Rpo11	DNA-directed RNA polymerase subunit A"
arCOG04257	RpoC/Rpo3/rpoA1	DNA-directed RNA polymerase subunit A'
arCOG04277	Efp	Translation elongation factor P (EF-P)/translation initiation factor 5A (eIF-5A)

123   **Table S5.** Calculated cell density of replicates in the SIT experiment. FID measurements of replicates used
124   to determine cell density. Density is calculated based on cell counts of DAPI and CARD-FISH labeled
125   samples. Stdev., standard deviation. Letters A-F identify each replicate.
126

Summary	$^{12}CH_4$		Cell Density, cells mL^{-1}		
Replicate ID	μM	ppm	DAPI	FISH	% labeled with Arch915
5A Day 22	25	610	3.83×10^7	1.34×10^6	3.5
5B Day 22	57	1,374	1.88×10^7	5.07×10^6	12.7
5E Day 22	50	1,204	3.49×10^7	0	0
5F Day 22	132	3,196	4.59×10^7	2.21×10^6	4.8
Day 22 Average	66	1,596	3.45×10^7	2.16×10^6	5.3
Day 22 Stdev.	46	1,116	1.14×10^7	2.15×10^6	5.4
5A Day 32	741	17,917	4.26×10^7	2.85×10^7	66.9
5B Day 32	2,176	52,635	1.11×10^8	6.00×10^7	54.2
5E Day 32	1,779	43,035	3.40×10^7	1.74×10^7	51.2
5F Day 32	2,412	58,355	9.14×10^7	4.03×10^7	44.1
Day 32 Average	1,777	42,985	6.97×10^7	3.65×10^7	54.1
Day 32 Stdev.	739	17,868	3.73×10^7	1.82×10^7	9.6
5A Day 45	4,109	99,398	12.2×10^7	6.41×10^7	52.7

127
128

SI Figures

Fig. S1. Detailed ANI (lower half of matrix) and AAI (upper half of matrix) analysis of related Archaeoglobales MAGs and reference genomes.

159
160
161 **Fig. S2.** 16S rRNA nucleotide identity analysis of closely related Archaeoglobales MAGs and reference
162 genomes.

164
Fig. S3. Experimental setup of the stable isotope tracing (SIT) experiment. Incubations were carried out in
30 mL culture volumes in 60 mL serum bottles with 8% v/v inoculum, 50 mg/L streptomycin, 50 mg/L
vancomycin, 10 mM MMA, and N_2 gas (99.999%) incubated in anoxic media (pH 7.8, 70 ˚C). Replicates
sacrificed for analysis during mid-log phase are indicated. Of the eight samples harvested for
metatranscriptomics, six were sequenced and used for analysis as two replicates did not yield sufficient
RNA for sequencing. AB, antibiotics streptomycin and vancomycin, MG, metagenome sample; MT,
metatranscriptome sample; BES, bromoethanesulfonate/methanogenesis inhibitor.
172

174
175 **Fig. S4.** $^{12}CH_4$ measurements by GC-FID during the stable isotope tracing experiment. Measurements can
176 be found in Dataset S3.
177
178

180
181

Fig. S5. Geographical distribution of geothermal features in Yellowstone National Park in which
Archaeoglobi-related *mcrA* genes (n = 36) and *Ca.* M. hypatiae-related 16S rRNA genes (n = 6) were
detected. Features are located in the (*A*) Map of Yellowstone National Park Wyoming, USA modified from
Vaughan *et al.* 2014 (26) *(B)* Lower Culex Basin (n = 36), (*C*) Mud Volcano Region (n = 2), and (*D*) the
White Creek Area (n = 1). These features spanned a wide pH (2.61-9.35) and temperature (18.4-93.8 ºC)
range. Features in which *mcrA* were detected are marked in red and features with related 16S rRNA genes
are shown in blue. Features in which both amplicons were detected are colored in purple. For details on
these sites, their *mcrA* data, water geochemistry, and exact location, see Lynes & Krukenberg et al., 2023.
Image source: Google Earth.
191

Description of Available Supplementary Datasets

SI Dataset S1. Extended metagenome assembled genome (SIT-MG) and isolate genome statistics. Seqs, sequences; avg_cov, average coverage; avg_gc, average G+C content; % rel. abund., percent relative abundance.

SI Dataset S2. GCMS measurements of masses 16 (CH_4), 17 ($^{13}CH_4$), and 19 ($^{12}CD_3H$) during the isotope tracing experiment. Percent of labeled methane is calculated as a fraction of provided labeled substrate. Stdev, standard deviation.

SI Dataset S3. Gas chromatograph FID measurements of $^{12}CH_4$ during isotope tracing experiment. NA, not available/measured.

SI Dataset S4. Gas chromatograph FID measurements of CH_4 during temperature optimum experiment. NA, not available/measured.

SI Dataset S5. Inventory of genes expressed by *Ca.* M. hypatiae LCB24 under methanogenic conditions and as depicted in Fig. 5. Expression levels averaged across six replicates are reported in reads per kilobase of transcript per million mapped reads (RPKM).

SI Dataset S6. Inventory of genes expressed by Ca. M. hypatiae LCB24 under methanogenic conditions belonging to the beta-oxidation pathway. Expression levels averaged across six replicates are reported in reads per kilobase of transcript per million mapped reads (RPKM).

References

1. Buessecker S, Chadwick GL, Quan ME, Hedlund BP, Dodsworth JA, Dekas AE. Mcr-dependent methanogenesis in Archaeoglobaceae enriched from a terrestrial hot spring. The ISME Journal. 2023;17(10):1649-59.

2. Lynes MM, Krukenberg V, Jay ZJ, Kohtz AJ, Gobrogge CA, Spietz RL, et al. Diversity and function of methyl-coenzyme M reductase-encoding archaea in Yellowstone hot springs revealed by metagenomics and mesocosm experiments. ISME Communications. 2023;3(1):22.

3. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial Ecology. 2015;75(2):129-37.

4. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology. 2019;37(8):852-7.

5. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal. 2011;17(1):10-2.

6. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13(7):581-3.

7. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Research. 2012;41(D1):D590-D6.

8. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. Microbiome. 2018;6(1):226.

9. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics. 2020;36(7):2251-2.

10. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: the conserved domain database in 2020. Nucleic Acids Research. 2020;48(D1):D265-D8.

11. Sondergaard D, Pedersen CN, Greening C. HydDB: A web tool for hydrogenase classification and analysis. Scientific Reports. 2016;6(1):34212.

12. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al. The InterPro protein families and domains database: 20 years on. Nucleic Acids Research. 2021;49(D1):D344-D54.

13. Neukirchen S, Sousa FL. DiSCo: a sequence-based type-specific predictor of Dsr-dependent dissimilatory sulphur metabolism in microbial data. Microb Genom. 2021;7(7).

14. Neukirchen S, Sousa FL. DiSCo: a sequence-based type-specific predictor of Dsr-dependent dissimilatory sulphur metabolism in microbial data. Microbial Genomics. 2021;7(7).

15. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for large alignments. PLoS One. 2010;5(3):e9490.

16. Jay ZJ, Beam JP, Dlakic M, Rusch DB, Kozubal MA, Inskeep WP. Marsarchaeota are an aerobic archaeal lineage abundant in geothermal iron oxide microbial mats. Nature Microbiology. 2018;3(6):732-40.

17. Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Backstrom D, Juzokaite L, Vancaester E, et al. Asgard archaea illuminate the origin of eukaryotic cellular complexity. Nature. 2017;541(7637):353-8.

18. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research. 2004;32(5):1792-7.

19. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular Biology and Evolution. 2013;30(4):772-80.

20. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25(15):1972-3.

21. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Molecular Biology and Evolution. 2020;37(5):1530-4.

22. Rusch A, Amend JP. Order-specific 16S rRNA-targeted oligonucleotide probes for (hyper) thermophilic archaea and bacteria. Extremophiles. 2004;8:357-66.

23. Stahl DA. Development and application of nucleic acid probes in bacterial systematics. Sequencing and
hybridization techniques in bacterial systematics. 1991:205-48.

24. Pernthaler A, Pernthaler J, Amann R. Fluorescence in situ hybridization and catalyzed reporter deposition for
the identification of marine bacteria. Applied and Environmental Microbiology. 2002;68(6):3094-101.

25. Schaible GA, Kohtz AJ, Cliff J, Hatzenpichler R. Correlative SIP-FISH-Raman-SEM-NanoSIMS links identity,
morphology, biochemistry, and physiology of environmental microbes. ISME Communications. 2022;2(1):52.

26. Vaughan RG, Heasler HP, Jaworowski C, Lowenstern JB, Keszthelyi LP. Provisional maps of thermal areas in
Yellowstone National Park based on satellite thermal infrared imaging and field observations: US Department
of the Interior, US Geological Survey; 2014.

27. Wang Y, Wegener G, Hou J, Wang F, Xiao X. Expanding anaerobic alkane metabolism in the domain of
Archaea. Nature Microbiology. 2019;4(4):595-602.

28. Liu YF, Chen J, Zaramela LS, Wang LY, Mbadinga SM, Hou ZW, et al. Genomic and transcriptomic evidence
supports methane metabolism in Archaeoglobi. mSystems. 2020;5(2):e00651-19.

29. Colman DR, Lindsay MR, Amenabar MJ, Boyd ES. The intersection of geology, geochemistry, and
microbiology in continental hydrothermal systems. Astrobiology. 2019;19(12):1505-22.

30. Peacock JP, Cole JK, Murugapiran SK, Dodsworth JA, Fisher JC, Moser DP, et al. Pyrosequencing reveals
high-temperature cellulolytic microbial consortia in Great Boiling Spring after in situ lignocellulose enrichment.
PLoS One. 2013;8(3):e59927.

31. Hua ZS, Wang YL, Evans PN, Qu YN, Goh KM, Rao YZ, et al. Insights into the ecological roles and evolution
of methyl-coenzyme M reductase-containing hot spring Archaea. Nature Communications. 2019;10(1):4574.

32. Wang J, Qu Y-N, Evans PN, Guo Q, Zhou F, Nie M, et al. Evidence for nontraditional *mcr*-containing archaea
contributing to biological methanogenesis in geothermal springs. Science Advances. 2023;9(26):eadg6004.

33. Klenk H-P, Clayton RA, Tomb J-F, White O, Nelson KE, Ketchum KA, et al. The complete genome sequence
of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. Nature. 1997;390(6658):364-70.

34. Slobodkina G, Allioux M, Merkel A, Cambon-Bonavita MA, Alain K, Jebbar M, et al. Physiological and
genomic characterization of a hyperthermophilic archaeon *Archaeoglobus neptunius* sp. nov. isolated from a
deep-sea hydrothermal vent warrants the reclassification of the genus *Archaeoglobus*. Frontiers in
Microbiology. 2021;12:679245.

35. von Jan M, Lapidus A, Glavina Del Rio T, Copeland A, Tice H, Cheng J-F, et al. Complete genome sequence
of *Archaeoglobus profundus* type strain (AV18T). Standards in Genomic Sciences. 2010;2(3):327-46.

36. Anderson I, Risso C, Holmes D, Lucas S, Copeland A, Lapidus A, et al. Complete genome sequence of
Ferroglobus placidus AEDII12DO. Standards in Genomic Sciences. 2011;5(1):50-60.

37. Mardanov AV, Slododkina GB, Slobodkin AI, Beletsky AV, Gavrilov SN, Kublanov IV, et al. The *Geoglobus
acetivorans* genome: Fe (III) reduction, acetate utilization, autotrophic growth, and degradation of aromatic
compounds in a hyperthermophilic archaeon. Applied and Environmental Microbiology. 2015;81(3):1003-12.

38. Manzella MP, Holmes DE, Rocheleau JM, Chung A, Reguera G, Kashefi K. The complete genome sequence
and emendation of the hyperthermophilic, obligate iron-reducing archaeon "*Geoglobus ahangari*" strain 234T.
Standards in Genomic Sciences. 2015;10(1):1-19.

39. Stokke R, Hocking WP, Steinsbu BO, Steen IH. Complete genome sequence of the thermophilic and
facultatively chemolithoautotrophic sulfate reducer *Archaeoglobus sulfaticallidus* strain PM70-1T. Genome
Announcements. 2013;1(4):e00406-13.

40. Boyd JA, Jungbluth SP, Leu AO, Evans PN, Woodcroft BJ, Chadwick GL, et al. Divergent methyl-coenzyme
M reductase genes in a deep-subseafloor Archaeoglobi. The ISME Journal. 2019;13(5):1269-79.

41. Mukherjee S, Seshadri R, Varghese NJ, Eloe-Fadrosh EA, Meier-Kolthoff JP, Göker M, et al. 1,003 reference
genomes of bacterial and archaeal isolates expand coverage of the tree of life. Nature Biotechnology.
2017;35(7):676-83.