Contrasting Effects of SARS-CoV-2 Vaccination vs. Infection on Antibody and TCR Repertoires

Jasper Braun[1], Elliot D. Hill[1], Elisa Contreras[1], Michie Yasuda[1], Alexandra Morgan[1], Sarah

Ditelberg[1], Ethan Winter[1], Cody Callahan[1], Gabrielle Mazzoni[1], Andrea Kirmaier[1], Ghee Rye

Lee[2*], Hamid Mirebrahim[3], Hosseinali Asgharian[3], Dilduz Telman[3], Ai-Ris Y. Collier[4,5,6], Dan H.

Barouch[4,6,7,8], Stefan Riedel[1,4], Sanjucta Dutta[1], Florian Rubelt[3], and Ramy Arnaout[1,4,9,†]

[1]Division of Clinical Pathology, Department of Pathology, Beth Israel Deaconess Medical Center,

Boston, MA 02215, USA; [2]Department of Radiology, Beth Israel Deaconess Medical Center,

Boston, MA 02215, USA; [3]Roche Sequencing Solutions, Pleasanton, CA 94588, USA; [4]Harvard

Medical School, Boston, MA 02215, USA; [5]Department of Obstetrics and Gynecology, Beth

Israel Deaconess Medical Center, Boston, Massachusetts; [6]Center for Virology and Vaccine

Research, Beth Israel Deaconess Medical Center, Boston, Massachusetts; [7]Ragon Institute of

MGH, MIT, and Harvard, Cambridge, Massachusetts; [8]Department of Medicine, Beth Israel

Deaconess Medical Center, Boston, Massachusetts; [9]Division of Clinical Informatics,

Department of Medicine, Beth Israel Deaconess Medical Center, Boston, MA 02215, USA

*Current address: Ohio State University College of Medicine, Columbus, OH 43210

Contact Information

18   †Corresponding author: Ramy Arnaout, Beth Israel Deaconess Medical Center,330 Brookline

19   Avenue, DA709, Boston, MA 02215, USA · rarnaout@bidmc.harvard.edu ·617-538-5681

Keywords

21   SARS-CoV-2, COVID-19, BCR repertoire, TCR repertoire, AIRRseq, ELISA, immunoPETE

Abstract

Antibodies and helper T cells play important roles in SARS-CoV-2 infection and vaccination. We sequenced B- and T-cell receptor repertoires (BCR/TCR) from the blood of 251 infectees, vaccinees, and controls to investigate whether features of these repertoires could predict subjects' SARS-CoV-2 neutralizing antibody titer (NAbs), as measured by enzyme-linked immunosorbent assay (ELISA). We sequenced recombined immunoglobulin heavy-chain (IGH), TCRβ (TRB), and TCRδ (TRD) genes in parallel from all subjects, including select B- and T-cell subsets in most cases, with a focus on their hypervariable CDR3 regions, and correlated this AIRRseq data with demographics and clinical findings from subjects' electronic health records. We found that age affected NAb levels in vaccinees but not infectees. Intriguingly, we found that vaccination and infection have an effect on non-productively recombined IGHs, suggesting an effect that precedes clonal selection. We found that repertoires' binding capacity to known SARS-CoV-2-specific CD4+ TRBs performs as well as the best hand-tuned approximate or "fuzzy" matching at predicting a protective level of NAbs, while also being more robust to repertoire sample size and not requiring hand-tuning. The overall conclusion from this large, unbiased, clinically well annotated dataset is that B- and T-cell adaptive responses to SARS-CoV-2 infection and vaccination are surprising, subtle, and diffuse. We discuss methodological and statistical challenges faced in attempting to define and quantify such strong-but-diffuse repertoire signatures and present tools and strategies for addressing these challenges.

Introduction

Since COVID-19's emergence, there has been great interest in identifying antibody (BCR) and T-cell receptor (TCR) gene sequences specific to SARS-CoV-2. The pandemic presented a high-profile opportunity to test the extent to which TCRs and BCR sequences against respiratory viruses would be public, i.e. appearing across many different individuals, or private, present in only one or a few individuals. New SARS-CoV-2-specific BCRs could form the basis for new treatments. Understanding how to identify and characterize commonalities in such a setting could help evaluate the viability of new diagnostics based on adaptive immune-receptor repertoire sequencing (AIRRseq).[1] number of studies have succeeded in identifying public immunoglobulin heavy-chain (IGH) and TCRβ (TRB) sequences. However, in part due to exigencies and constraints imposed by the pandemic, and in part because it was impossible to know a priori what study size would be adequate to identify public sequences comprehensively in COVID-19, many of these studies involved relatively few individuals. Small sample sizes have known limitations for AIRRseq studies. Because small samples may not be representative of larger populations, results on small samples may not generalize. Small studies may be insufficiently powered to detect subtle patterns. And the smaller the sample size, the more likely it is that random fluctuations in the data—literally, the luck of the draw—will produce results that appear to be statistically significant but do not reflect underlying relationships. Moreover, most studies investigated only BCRs or only TCRs, but not both in the same cohort, despite the importance of both antibody and T-cell responses in SARS-CoV-2 infection.[2,3] To our knowledge only two previous COVID-19 studies[4,5] have sequenced TCR from Ɣδ T cells, a little-studied subset that may be important in mucosal antimicrobial immunity.[6,7] How SARS-CoV-2 virus or vaccine exposure affects different B- and T-cell subsets (IgM-positive vs. IgM-negative B-cells, CD4-positive vs. CD8-positive T cells) has also been insufficiently explored.[2,3]

To our knowledge previous studies have identified 20 IGH V genes to be enriched in sequences produced during various immune responses to SARS-CoV-2.[8–16] Given that human genomes encode 54 IGH V genes,[17] collectively these studies implicate 37% of all V genes in the response

to this single viral exposure, indicating that the SARS-CoV-2 response is either quite broad

within individuals, quite heterogeneous among individuals, or both. There is no obvious reason

to think each V gene would contribute to only a single SARS-CoV-2-specific recombined IGH

sequence, much less a single IGH:immunoglobulin light chain (IGL) pair; therefore, collectively

these studies also suggest that the IGH response to SARS-CoV-2 might account for a significant

fraction of a given repertoire, a possibility that requires more comprehensive AIRRseq

investigation. Note that studies that compare only a single non-control cohort to a control

cohort cannot distinguish between features (clones, motifs, genes, CDR3 lengths, etc.) that

signify a disease-specific vs. a general immune response.

Regarding TCRs, one study[18] searched repertoires of 140 COVID-19 patients and another 140

pre-pandemic (and therefore unexposed) controls for the presence of each of 1,267 TRB

sequences that had independently been shown to recognize epitopes of the SARS-CoV-2 spike

protein. These authors showed that while the presence of some of the TRB sequences in almost

all of the repertoires suggested a public response to SARS-CoV-2 infection, the fraction of the

repertoire that matched the query sequences was similar between infectees and controls.

These authors also looked for SARS-CoV-2 specific TRBs in the brain tissue of COVID-19 patients,

since T-cell infiltration of the brain, an organ otherwise seldom infiltrated by T cells, is known to

occur during COVID-19 infection.[19–21] The 68 TRBs they identified were found in 40% of COVID-

19 repertoires vs. 17% of pre-pandemic controls. This suggests significant enrichment even as

the majority of individuals with COVID-19 lacked these sequences and a significant minority of

controls (1 in 6) had them despite these control samples having been collected before the

pandemic (which has been observed in other contexts,[22] perhaps indicating cross-reactivity

with previously circulating coronaviruses, which are very common in human populations[3]).

In addition, one study[23] identified a large database of SARS-CoV-2-specific TRB sequences as

being shared among infectees and unenriched among healthy controls. Subsequently, another

study[24] sequenced repertoires of individuals 0 and 4 weeks after vaccination with the Oxford-

AstraZeneca COVID-19 vaccine AZD1222 or the meningococcus vaccine MenACWY and matched

the database sequences to these repertoires. An increase of database sequences among

AZD1222-vaccinee repertoires but not MenACWY-vaccinee controls was seen between the two

timepoints.

Heterogeneity in clinical settings across studies complicates the interpretation of private vs.

public responses, for several reasons. First, there are important antigenic differences between

exposure and vaccination. This is especially true for the mRNA vaccines, which immunize

subjects with only spike protein, in contrast to the full complement of SARS-CoV-2 proteins to

which infectees are exposed. Second, demographics may play a role.

For example, it has long been recognized that individuals respond differently to vaccines by age,

with older individuals generally mounting less-robust and shorter-lasting responses as

measured by ELISA.[25–27] Other clinical features are also known to affect the adaptive immune

response to infection and vaccination, including immunosuppressive conditions such as organ

transplant or cancer therapy as well as metabolic disease.[28] Third, studies from different

periods of the pandemic likely measured responses to different strains. Fourth, the signal or

signature detected may differ depending on whether the controls were healthy, which might

result in detecting generalized responsiveness (e.g. bystander activation), or were instead

presenting with a non-COVID illness, making any signal/signature more likely to be specific for

COVID-19. Fifth, exposure, whether to replicating virus or to an inactivated or subcomponent

vaccine, may not be as clinically relevant as whether a substantial NAb response was mounted.

This is because NAbs are a marker of protection in SARS-CoV-2.[29–31] (T-cell-mediated immunity

may also play an important role[32]). And sixth, accessing clinical annotations from electronic

medical records can be challenging.[33] As a result, the effect of clinical heterogeneity in AIRRseq

studies in the setting of COVID-19 has been under-explored to date.

In all, the work above supports the view that there are commonalities in IGH and TRB at the

gene and sequence level in response to SARS-CoV-2 infection; however, the nature of the

signature is not well understood. One open question is to what extent infection affects

antibody and TCR repertoires as a whole vs. enriching specific clones within it. One can refer to

these ends of the continuum of possible effects as "diffuse" vs. "precise." From previous work

on repertoires, "diffuse" features include CDR3 length, the frequency of usage of specific V or J

genes, and repertoire diversity as measured any of several ways (richness, Shannon entropy,

Simpson's index, or their Hill-number equivalents).[34] At the other extreme, the most "precise"

features are the frequency of clones   with specific sequences. Between these extremes is a

set of features that includes approximate or "fuzzy" matching of sequences[35] and other

clustering methods.[36,37] This middle ground has been less explored. Recently the concept of

binding capacity has been developed to measure the fraction of a repertoire that is "like" a

given query sequence in terms of target specificity (weighting the repertoire by the predicted

dissociation constants of its constituent antibodies or TCRs and by their sequence frequencies).

Whether or how binding capacity might be affected by SARS-CoV-2 infection and/or vaccination

is unknown.

Given this background, we sought to investigate the effects of SARS-CoV-2 infection and

vaccination on both antibody and TCR repertoires in a large clinical cohort, with attention to

major B- and T-cell subsets where possible, using NAbs via ELISA as a functional readout, with a

special focus on diffuse repertoire features and how they compare to both more traditional

features and to clinical correlates.

Results

NAbs vary with exposure, age, and immune status

Using immunoPETE (Roche; research use only),[5] we deep-sequenced IGH, TRB, and TRD from

the blood of 251 individuals: 36 vaccinees, all of whom had received an mRNA vaccine, 145

infectees, 53 healthy controls, and 20 with unknown SARS-CoV-2 exposure status. Three

individuals belonged to both the vaccinee and infectee groups. Forty-seven subjects were

considered immunosuppressed and the remaining 204 immunocompetent. Blood samples for

129/145 (89%) infectees were within 6 months of the most recent positive PCR test on record

and 121/145 (83%) were ≥7 days from the presumed most recent infection date (assuming a

mean of 4 days from exposure to testing). Fig. S1 presents a summary of the timeline and

sequencing yield. Tables S1 and S2 present demographics and relevant comorbidities for the

different cohorts. We measured plasma NAbs against SARS-CoV-2 spike for 237/251 subjects.

Fig. 1 presents a summary of the measured NAbs concentrations by cohort,

immunosuppression status, and age.

NAbs were undetectable in some infectees and one vaccinee (Fig. 1). The odds of producing

NAbs were significantly higher in immunocompetent subjects compared to immunosuppressed

subjects (OR=3.9, p_c=0.008—note, all p-values in this work have been corrected for multiple-

hypothesis testing; we write p_c to indicate this). Odds were also significantly higher in the

infectees (OR=5.8, p_c<0.001) and vaccinees (OR=4.7, p_c=0.034) compared to controls. Age was

not significantly associated with NAbs titer (p_c=0.801) and therefore age was excluded from

most of the subsequent models (below).

Among the subjects who did produce detectable NAbs, NAb concentration was notably higher

in the vaccinated and infected groups compared to the control group (Fig. 1a). The relationship

between concentration and age was more complex and depended on the cohort (significant

age × cohort interaction). Age affected titer only in vaccinees, with NAbs being lower in older

individuals: above age 65, individuals had higher NAbs with infection than vaccination (Fig. 1a).

Meanwhile, age did not affect NAbs in the control or infected groups. We found no significant

effect of immunocompetence on NAbs in subjects who had non-zero NAbs.

Vaccination is associated with shorter IGH CDR3s in productive joins

The characteristic (e.g. mean or median) length of CDR3s is known to vary during development

and in response to various exposures, at least in productively recombined IGH genes, a.k.a.

"productive joins."[38] Because only productively recombined IGH genes can be expressed as

(BCR) proteins, such differences are generally considered evidence that the B cells that express

them are selected for having e.g. longer CDR3s. We found that vaccinees had shorter IGH

CDR3s than controls (p_c=0.024; Fig. 2a) or infectees (p_c=0.0046; Fig. 2b), indicating a repertoire-

wide difference in the B-cell response to vaccination vs. infection (Table S3, Figs. S2-S4).

The length of IGH CDR3s depends on the lengths of the constituent IGHV, IGHD, and IGHJ

genes, as well as the number of non-templated (N) and palindromic (P) nucleotides inserted at

the junctions between them.[39] Annotating IGHD and distinguishing mutated/truncated IGHD

sequence from N and P sequences is challenging due to insertions, chewbacks, and somatic

hypermutation. However, IGHV and IGHJ can be annotated reliably, and so we tested whether

the overall differences in CDR3 length were attributable to differences in the use of longer vs.

shorter IGHV and IGHJ genes.

We grouped IGHV genes by the number of amino acids that their germline contributes to

CDR3s, and similarly for IGHJ genes. The 54 IGHV genes hard-coded as part of the human

germline contribute either 3 or 4 amino acids to the CDR3, depending on the gene. We found

that vaccinees generally used more of the IGHV genes that contribute 3 residues (p_c=0.021 vs.

controls and p_c=0.0028 vs. infectees) and fewer of the IGHV genes that contribute 4 residues

(again p_c=0.021 vs. controls and p_c=0.0028 vs. infectees; Fig. 2c and Table S4). The same pattern

held for IgM-positive and IgM-negative (mainly IgG; hereafter "IgG") in controls and infectees.

Meanwhile, the six IGHJ germline genes contribute 5 (IGH J4), 6 (IGH J3 and J5), 7 (IGH J1 and

J2), or 10 (IGH J6) amino acids to the CDR3 (Fig. 2d). We found that vaccinees used more J4

(p_c=9.4e-5 vs. controls and p_c=1.9e-6 vs. infectees) and fewer J3 & J5 (p_c=0.0002 vs. controls and

p_c=0.0021 vs. infectees; Table S4). Thus, the preference of shorter IGH CDR3s after vaccination

can at least partially be explained by selection for V and J genes that contribute fewer residues

to the CDR3.

No such differences were observed in TCR CDR3s, which have a far narrower length

distribution.

Vaccination is associated with longer IGH CDR3s in non-productive joins

Next we sought to estimate the strength of selection for IGH CDR3s of different lengths in

vaccinees, infectees, and controls. This can be done by comparing the length distribution of

productive joins to the distribution in non-productive joins, i.e. those in which VDJ

recombination occurs out of frame or produces stop codons. Because non-productive joins do

not produce functional antibodies, the B cells that contain them cannot be selected for or

against based on them. Nevertheless, the lengths of the CDR3 regions in non-productive joins

can be measured. Thus, any differences in length between non-productive joins and productive

joins reflect selection on (some aspect of) the productive joins, for example by exposure to

SARS-CoV-2 (in infectees) or vaccine contents (vaccinees).

Our null hypothesis was that the lengths of non-productive joins would be similar for vaccinees,

infectees, and controls. Surprisingly, we found that CDR3s in non-productive joins differed

across these three cohorts. In fact, we observed reverse relationships from the ones we saw in

productive joins: CDR3s in non-productive joins were longer in vaccinees and infectees than in

controls (p_c=0.039 and 0.0021, respectively). Vaccinees' non-productive CDR3s used the

shortest J gene, J4, less often and the longest J, J6, more often than controls' (p_c=0.00011 and

0.022, respectively; Fig. S5). Thus, selection for shorter CDR3s in vaccinees is even stronger than

indicated from the comparison of productive joins in the previous section, because in

vaccinees, recombination, which precedes selection, is biased toward longer CDR3s. Again, no

such differences were observed in TCR CDR3s (Figs. S6 and S7).

Vaccination affects at least one-sixth of the pre-selection IGH repertoire

We next sought to better characterize this apparent effect of vaccine exposure on IGH

recombination. The results in the previous section were regarding differences in subjects' entire

IGH CDR3 repertoires. However, vaccine exposure is generally thought to affect only a portion

of the repertoire. The rest of the repertoire, the unaffected portion, should be the same as a

control's. Therefore, conceptually, each vaccinee's repertoire can be thought of as a weighted

sum of two parts: a vaccine-responsive part and a control part. We asked what the minimum

size of the vaccine-responsive part would have to be, in order to explain the difference in the

length distribution of non-productive joins between vaccinees and controls.

To do this, we analyzed the differences between the mean IGH CDR3 length-distribution curves

of vaccinees and controls. By calculating differences at each length, we generated the length

distribution that the putative vaccine-responsive part would have to have, in order for the

vaccinee curve to be a weighted sum of the control curve and the vaccine-responsive part, for a

given size of the vaccine-responsive part (Fig. 2e). Inevitably, there will an inverse relationship

between how different the length distribution of the vaccine-responsive part is, and its size.

This fact sets a floor on the size of the vaccine-responsive part: any smaller, and the vaccine-

responsive part would have to be so different that at least one of its lengths would have a

negative frequency.

For example, 20-amino-acid-long CDR3s constituted an average of 9% of non-productive joins in

controls but 8% in vaccinees. Considering just this length for the moment, if length-20 CDR3s

constituted 7% in the vaccine-responsive part, then the vaccine-responsive part would have to

be 50% of the repertoire, since 50%×9% + 50%×7% = 8%. If instead length-20 CDR3s constituted

3.5%, the vaccine-responsive part would only have to be 18%, since (100-18)%×9% + 18%×3.5%

= 8%. In this example, the vaccine-responsive part could never be as small as 1%, since in that

case length-20 CDR3s would have to have a negative frequency. By this approach, we found

that the size of the vaccine-responsive part could be no smaller than 16%, or one-sixth, of the

vaccinees' non-productive joins.

Vaccinees and infectees with more SARS-CoV-2-specific TRBs have higher NAbs

Next, we tested whether TRB and IGH CDR3s that had been previously found to be associated

with SARS-CoV-2 exposure, including by structural studies, were enriched among our vaccinee

and infectee cohorts (see Methods). We obtained SARS-CoV-2-specific TCRs from CD4 and CD8

T cells from Nolan et al.[23] and obtained non-CD-restricted SARS-CoV-2- and non-SARS-CoV-2-

specific TRBs and IGHs from CoV-AbDab, PDB, and VDJDb.[40–42] These comprised totals of

184,100 unique SARS-CoV-2-specific TRBs and 1,630 unique SARS-CoV-2-specific IGHs (Table

S5).

We found a much higher proportion of SARS-CoV-2-specific TRB sequences than IGH sequences

had exact matches in our samples: ≥ 12% vs. 0.1%, respectively, with the 0.1% representing just

a single sequence (Table S5). The fraction of each repertoire that matched SARS-CoV-2-specific

TRBs correlated positively with NAbs, as measured by ELISA titer, in infectees and vaccinees

(Fig. 3a-b, Table S6, and Fig. S9). In infectees, for whom we had separate CD4-positive and CD4-

negative (mainly CD8-positive; hereafter "CD8") TRB repertoires, the positive correlation was

confined to CD4 repertoires. In contrast, no correlation was seen for controls. Likewise, no

correlation was seen for TRBs that were not specific for SARS-CoV-2 in infectees, supporting the

interpretation that this correlation is causal. Nevertheless, this correlation alone performed

poorly as a classifier of who had high enough NAbs to be considered positive (per the ELISA test

manufacturer), with an area under the receiver-operator characteristic curve (AUROC) of 0.55

(95%CI, 0.46-0.63). Notably, there was also a positive relationship between non-specific TRBs

and NAbs in vaccinees, although the 95% CI on the regression slope only narrowly missed

including zero (Table S6).

Binding capacity outperforms fuzzy matching for measuring similarity

That subjects had almost no exact matches to SARS-CoV-2-specific IGH sequences did not

exclude the possibility that they have sequences that are functionally similar to these reference

sequences. The same possibility exists for TRBs. A standard method for finding similar

sequences is using the Levenshtein distance (i.e. edit distance). Sequences with a distance of

less than or equal to a tolerance t are considered similar (for example, sequences that differ by

no more than t=1 amino acid). This is known as "fuzzy matching" with tolerance t. (Note that

exact matches are just fuzzy matches with tolerance 0.) Unfortunately, there is no consensus on

what t should be chosen. Also, the fraction of a repertoire that fuzzy-matches a set of

references could depend on repertoire size because of the nature of sampling, potentially

complicating the use of fuzzy matching.

To test this possibility, we subsampled 30 subjects' repertoires (10 controls, 10 infectees, and

10 vaccinees) and measured the fraction of the repertoire that fuzzy-matched SARS-CoV-2-

specific CD4 TRBs at tolerances of 0, 2, 4, 6, 8, and 10 amino acids. We fit a linear mixed model

grouped by subject for all repertoires with at least 1,000 sequences. We found the fraction of

fuzzy matches depended strongly on repertoire size for all repertoire sizes measured (up to 1

million sequences), falling steeply and continuously throughout (Fig. 3c). Thus, fuzzy matching

was shown to not be a robust measure of repertoire content in this study.

We therefore tested a recently proposed alternative method for finding similar sequences:

measuring repertoires' binding capacity for the targets of reference sequences.[43] Binding

capacity is the average similarity of a repertoire to one or more reference sequences, with

similarity estimated according to a general model of the likelihood of a given sequence in the

repertoire to bind the same antigen as a reference sequence. In contrast to fuzzy matching, we

found the binding capacity remained robust for sample sizes above 1,000 sequences, with only

minimal dependence on repertoire size (Fig. 3d). Binding capacity was more robust to

repertoire size than fuzzy matching at all tolerances tested (Fig. 3e, Table S7; note that binding

capacity does not require a choice of tolerance; it is independent of and therefore robust to

tolerance; technically it is a nonlinear weighted mean across all tolerances). Thus, binding

capacity provides a robust way to measure the fraction of these TRB repertoires that is similar

to reference SARS-CoV-2-specific TRB sequences.

Repertoire features predict levels of NAbs consistent with exposure comparably to clinical

data

Finally, we compared how well above feature sets predicted exposure-level NAbs titers. To do

so, we trained machine-learning models that used each of these feature sets. Because there

were many reference SARS-CoV-2-specific TCR sequences to consider, each of which produces

one exact-matching fraction, several fuzzy-matching fractions (one for each chosen tolerance),

and one binding capacity measurement, there was a risk of overfitting (true whenever the

number of features exceeds the number of datapoints). Therefore we first filtered out

uninformative features.

To do this, we calculated exact/fuzzy matches and binding capacities for SARS-CoV-2 specific

and non-specific sequences (from VDJDB) and measured their correlations to NAb titer. (Based

on the results above, we only used repertoires with ≥ 1,000 sequences.) We used non-specific

sequences as a null model and kept only SARS-CoV-2-specific sequences with correlations

outside the middle 95% of the null model: specific sequences on the high end correlated more

with NAb titer than was expected by chance, while specific sequences on the low end were

correlated inversely with NAb titer to a larger degree than expected by chance (Fig. 4a). Of

7,804 SARS-CoV-specific features with non-zero fractions or binding capacities, this process

filtered out all but 323. To reduce redundancy and further reduce the number of features, we

performed Principal Component Analysis on the results (the number of Principal Components

to keep was tunable and fit by each model). We did the same to reduce the number of V-gene

features. To avoid data leakage, we performed this dimensionality reduction procedure on

training data only.

We performed 700 replicate logistic-regression fits on each of the above feature sets and

measured performance by AUROC (Fig. 4b). As a comparator, we also fit 700 replicates on

subjects' infection and vaccination status, which we reasoned would approximate the

maximum possible performance that should be achievable on this dataset. As expected, this

comparator resulted in the highest median AUROC of all the feature sets tested, at 0.72 (inter-

quartile range across the replicates, 0.66-0.79; Fig. 4b). Strong performance was also seen

when training on fuzzy matches with tolerance 2 on all TRB sequences (AUROC 0.71; IQR, 0.64-

77) and on TRJ frequencies for CDR4 TRB sequences (0.70; 0.62-0.77). Binding capacities on all

TRB sequences showed similar performance to these two (0.68; 0.61-0.74), while exact matches

on the same sequences showed poor performance (0.59; 051-0.66).

In sum, being infected and/or vaccinated—the gold-standard clinical model—lacked high

predictive power for Nab titer, although binding capacities, fuzzy matches with a tolerance of 2,

and TRJ frequencies on CD4 TRB sequences performed nearly as well and much better than

exact matches.

To better understand the characteristics of different feature sets, we also calculated sensitivity,

specificity, and precision for all replicates (see Fig. 4c). The clinical feature set's performance

metrics are relatively well balanced. In contrast, for most binding-capacity and fuzzy-matching

feature sets, sensitivity and precision were low, making them less desirable for screening.

Interestingly, features based on IgG diversity had the highest sensitivities while IgM diversities

had the lowest sensitivities. The reverse was true for specificities, with IgM diversities having

among the highest specificities and IgG diversities among the lowest. While it should be noted

that repertoire diversity is not disease specific, these observations suggest that trends in

diversity measurements taken for different repertoire subsets might give insights about

exposure status in very different ways.

Discussion

Detecting and defining signatures in repertoire sequence is challenging in part due to the large

number of features that can contribute to a signature. These include high-level features such as

CDR3 length and repertoire diversity, mid-level features such as frequencies of V and J gene

usage and VJ combinations (D genes are harder to assign), and low-level features such as the

frequencies of specific reference sequences. Repertoire diversity itself is actually a set of

features, some of which incorporate sequence similarity, which furthermore can be defined in

multiple ways[38,43]. Ideally the features above should be measured in both antibody and TCR

repertoires, since they act cooperatively[44], and in cell subsets defined by isotype (for B cells) or

CD4 vs. CD8 expression (for T cells). Thus, overall, the total number of features that can be used

to detect and define signatures reaches into the hundreds of thousands.

As a result, statistical confidence requires large study sizes, which are challenging to obtain;

methods that can avoid spurious associations, which are common in high-dimensional systems;

appropriate controls, so that signatures are specific and not related to e.g. general immune

activation; and detailed clinical annotation, which we obtained from our electronic medical

record (as detailed in Materials and Methods). Even with these design safeguards in place, the

signature of exposure to a specific immunogen, such as SARS-CoV-2, may be broad or diffuse,

with different individuals' repertoires reacting in different ways. And factors and features

outside of repertoires may be important for determining exposure.

Given these considerations, our study was fairly large, with over 250 subjects, and involved

sequencing IGH and TRD as well as TRB, to a median depth of over 10^5 cells/subject, made

possible by ImmunoPETE's integrated library preparation.[5] To focus analysis on SARS-CoV-2-

specific signatures and patterns, controls in our study were not typical "healthy controls" but

rather patients presenting for care who had sufficient concern for SARS-CoV-2 infection, and

who were therefore tested, and were negative. At the time, hospital policy involved widespread

testing with very sensitive tests (limit of detection, 100 copies of viral mRNA/mL), so we

consider the probability of false negatives to be low. In addition, for infectees and controls, we

separately analyzed IGM- and non-IGM (predominantly IGG+)-isotype antibodies and CD4+ and

CD4- (i.e. mainly CD8+) T cells. Scheduling issues related to vaccine rollout prevented separate

subset analysis for vaccinees, a limitation of the study. We also limited dimensionality, thereby

increasing statistical confidence, by filtering for features that correlate with the outcome

measure of NAb titer. And instead of simply combining all features into a single model, we

compared models with different feature sets to tease apart where signals might lie. Finally, we

compared these to the simplest model we could think of, made up of readily available clinical

information: whether or not a person was infected and/or vaccinated, to test how repertoire

data compares (and what, if anything, it could add). To our knowledge this is the largest such

study, and possibly the first. It led to the discovery of several previously unreported patterns

across multiple feature sets, for both IGH and TRB, as well as in multiple subtypes of B and T

cells, that merit discussion.

First, the pattern in IGH CDR3 lengths in vaccinees was curious for several reasons. First, it

involved a change in non-productive joins (which in our reading of the literature are usually

treated as a baseline and not compared between cohorts, as we did). This was unexpected

because B cells are selected for survival based on expressed B-cell receptors, and non-

productive joins are not expressed. Our finding seems to indicate selection independent of

expression (non-productive joins are not expressed). Second, this is a much larger effect than

would be expected from an antigen-specific adaptive immune response. Immunogen-specific B

cells rarely exceed low-single-digit percentages of the repertoire. Yet the effect we found

appears to involve at least one-sixth (~17%) of the repertoire. Third, the direction of the length

change in non-productive joins is opposite that of productive joins: Non-productive CDR3s are

longer in vaccinees than in controls and infectees, but productive CDR3s are shorter. And

fourth, while other patterns we found were fairly similar between vaccinees and infectees, this

CDR3 length effect appears confined to vaccinees.

We conclude that vaccination may have some undescribed effect on the V-D-J recombination

machinery, biasing recombination toward use of IGHJ genes (and secondarily IGHV genes) that

result in longer CDR3s. This effect would have to be due to some difference between the

vaccine and natural infection, or else it would have been seen in infectees. If our interpretation

is correct, it would mean the effect of selection for shorter CDR3s in productive joins is quite

strong, because there are fewer short joins from which to select. In any event, both vaccination

(in nonproductive and productive joins) and infection (in productive joins) affect a larger

proportion of IGH repertoires than is typically considered "specific."

Second, binding capacity was shown to have essentially the same predictive power as the best-

performing version of fuzzy matching. Recall that both fuzzy matching and binding capacity

measure the size of groups of similar antibodies or TCRs. Here they were applied by taking a

reference sequence, for example a sequence previously reported in the literature to be

associated with SARS-CoV-2 (a "SARS-CoV-2-specific sequence") and ask what fraction of a

given subject's repertoire was similar to that index sequence. The methods differ in how they

view similarity. Fuzzy matching requires choice of tolerance: above a set number of amino-acid

mismatches, a query sequence is considered different to the index sequence. If the tolerance is

2, a query with 3 mismatches is considered just as different from the index sequence as a query

with 20 mismatches.

Binding capacity has neither problem. It is based on the measured relationship between

number of mismatches and change in dissociation constant (K_d), i.e. binding similarity (cite

Arora Arnaout 2023). This empirical data essentially substitutes for having to choose a

tolerance. In addition, binding capacity is continuous: a query with 3 mismatches is more similar

to the index than a query with 20 mismatches. Consequently, binding capacity can detect the

potential presence of a large group of sequences with low similarity, which collectively might

play as important a role as a small group of high-similarity sequences (or in the limit, the

presence of the index sequence as a high-frequency clone). The magnitude of the CDR3 length

effect supports the importance of being able to detect such diffuse/weak signals. We showed

that different tolerances had different ability to predict NAb titer. To us there is no obvious

reason that tolerance of 2 should outperform, e.g., a tolerance of 10. Possibly which tolerance

is best may differ by exposure. That binding capacity performs comparably to the best-

performing tolerance supports its utility for immune-repertoire analysis.

We note that in principle, diffuse effects could reflect superantigen-type activity.[45] However, to

our knowledge, clinical evidence for a superantigen in SARS-CoV-2 is scant and has focused on T

cells. Although proposed to explain the rare sequela of multisystem inflammatory syndrome in

children,[46] wet-lab investigation has not demonstrated superantigen activity in SARS-CoV-2.[47]

This study has several limitations. First, we were unable to sort vaccinee samples to obtain

separate IgM vs. IgG B-cell repertoires and CD4 vs. CD8 T-cell repertoires due to exigencies at

the height of the pandemic. Different subtypes may follow different (even opposite) trends, as

did the sensitivities and specificities of classifiers trained on IGG and IGM diversities. Any such

patterns in vaccinees were beyond our ability to measure. Second, we used concentrations of

SARS-CoV-2 anti-spike NAbs as our proxy of protection. Signals may be present that do not

correlate with antibodies binding this particular immunogen. For example, a signal might be

seen in T cells or antibodies that bind other SARS-CoV-2 proteins, which we are unable to

evaluate given NAbs as a readout. Third, although the sequence data in this study was

quantitative, it contained only single-chain, not paired-chain data. Fourth, the ability to define

signatures is limited by uncertainty about the specificity of reference sequences. Much effort is

being put into methods that predict receptor-antigen binding, but a unified, accepted, and

feasible approach to identifying all sequences that bind a given immunogen has yet to be

established. Fifth, the quality of binding capacity measurements is limited by the current

measure of binding similarity being based on mean behavior[43]; this is expected to improve with

additional data and advances in protein structure prediction. And sixth, we did not have the

benefit of HLA typing for these samples, which could be valuable because of its influence

especially on the TCR repertoire.

It will be valuable to see the methodology presented here, with its many steps taken to

maximize robustness and avoid statistical artifacts, applied to additional datasets. This will give

additional evidence of how well these results and this approach generalize for SARS-CoV-2 in

general, for immune responses to variants of the virus, and for other pathogens and

immunogens. A careful statistical approach applied to multiple, functional features, measured

on unbiased repertoire sequence from TCR and BCR subsets from large cohorts, is, in our

opinion, the best way to decipher the rich information that the adaptive immunome encodes.

Acknowledgements

This work was supported by the National Institutes of Health (NIH) National Institute for Allergy and Infectious Diseases (NIAID) R01AI148747; the Massachusetts Life Sciences Center Bits to Bytes program; the Extreme Science and Engineering Discovery Environment, which is supported by NSF Grant ACI-1548562; and the Comet supercomputer at the San Diego Supercomputer Center, which is supported by NSF Grant ACI-1341698.

Conflict of Interest Disclosures

HM, HA, DT, and FR are employees of Roche Molecular Systems Inc. JB, EDH, EC, MY, AM, SD, EW, CC, GM, AK, A-RYC, DHB, SR, SD, and RA have no conflicts of interest to declare.

Materials and Methods

Study subjects

The subjects in this study were patients seeking clinical care at the Beth Israel Deaconess Medical Center (BIDMC), a 743-bed tertiary care medical center in Boston, MA, USA. BIDMC serves a large and diverse population in and around eastern Massachusetts, USA, centered on the Boston metropolitan area.

Institutional review board approval

All work was carried out in accordance with BIDMC's Institutional Review Board protocols 2020P000634, 2021P000109 and 2020P000361.

Cohort assignment

All subjects from whom samples were obtained received RT-qPCR tests performed on two Abbott Molecular platforms: m2000 and Alinity m (Abbott Molecular, Des Plaines, IL, U.S.A.). These detect identical SARS-CoV-2 N and RdRp gene targets and are extremely sensitive for SARS-CoV-2 infection, with limit of detection of 100 copies/mL.[48–50] Infectees had a positive

result at the time of sample acquisition. Controls were tested, but negative. COVID-19 test and

vaccination information were obtained using SQL queries from BIDMC's clinical data repository

and via a dedicated REDCap database set up to facilitate research involving vaccinees.[51]

Using these records, subjects were considered infectees if there was a record of a positive

COVID-19 test result dated before or on the sample collection date and non-infected otherwise.

If no medical record number was available for a subject, their infection status was considered

unknown. Subjects were considered vaccinees if vaccination prior to or on the day of sample

collection was indicated as the appropriate procedure code in the clinical data repository,

recorded in REDCap, or identified from Massachusetts' state Immunization Information System.

The entire vaccinated cohort had received mRNA-based vaccines, with 80% receiving Pfizer

Comirnaty and the rest receiving Moderna Spikevax. Samples were collected from the vaccinee

cohort 4 to 84 days (mean = 44.3 days, standard deviation = 15.3 days) after administration of

the initial vaccine dose.

Subjects were considered non-vaccinated if the sample collection date preceded 12/15/2020

(the date of the first administered COVID-19 vaccine); if there was record of vaccination after

sample collection that was annotated as the first dose; if there were two vaccinations after

sample collection where the second was annotated as the second dose; or if there were two

vaccinations after sample collection within 42 days of each other (consistent with being the

primary series). Subjects that did not satisfy vaccinee or non-vaccinated criteria were

considered to have unknown vaccination status. Subjects were annotated as unexposed

controls if they were non-infected and non-vaccinated. Subjects whose vaccination status was

unknown or whose infection status was unknown and were neither vaccinees nor infectees

were considered to have an "unknown" SARS-CoV-2 exposure status.

Clinical annotations

Immunosuppression

Subjects were labelled either "immunosuppressed" or "immunocompetent." Subjects were

designated immunosuppressed if at least one of the following criteria was met:

● the most recent CD4+ cell count was less than 100 cells/µl;

● there was a diagnosis of lymphoma or leukemia associated with a healthcare encounter

(visit, admission, or phone call) either before or within 60 days after sample collection;

or

● the subject was prescribed any of the following medications on an ongoing basis prior to

sample collection and with enough refills to include up to 30 days **after**: abatacept,

adalimumab, anakinra, azathioprine, basiliximab, budesonide, certolizumab,

cyclosporine, daclizumab, dexamethasone, everolimus, etanercept, golimumab,

infliximab, ixekizumab, leflunomide, lenalidomide, methotrexate, mycophenolate,

natalizumab, pomalidomide, prednisone, rituximab, secukinumab, sirolimus, tacrolimus,

tocilizumab, tofacitinib, ustekinumab, and vedolizumab.

If none of these criteria were met, subjects were considered immunocompetent.

Demographics

If a subject had a COVID test, the sex and date of birth were read from the corresponding

record. Otherwise, we read sex and date of birth from other records of lab specimens, the

electronic health record (EHR), or the project's REDCap database (always in structured fields,

not using natural-language processing). Self-reported race was read from the EHR.

Risk factors

A semi-automated review of EHRs for ICD-10 diagnosis codes and related entries was used to

identify subjects having any of the medical conditions highlighted by the CDC as increasing risk

of severe illness from COVID-19.[52] Where feasible, the list of ICD-10 codes indicative of each

comorbidity was taken from the Elixhauser Comorbidity Software Refined for ICD-10-CM,[53]

version v2022.1, developed for the Healthcare Cost and Utilization Project (HCUP), which is

based on the work of Elixhauser et al.[54] In addition to these, another widely used set of

comorbidity measures is the Charlson Comorbidity Index.[55] For comorbidities not defined in the

HCUP software, the lists of ICD-10 codes defined by this study[56] were used where possible.

Comorbidities that were not codified in either resource were identified, where possible, using

ICD-10 codes or other automated chart queries, detailed as follows:

● Cancer: identified using ICD-10 codes in the HCUP software for "Leukemia,"

"Lymphoma," "Metastatic cancer," or "Solid tumor without metastasis, malignant."

● Chronic Kidney Disease: identified using ICD-10 codes in the HCUP software for "Renal

failure, moderate," and "Renal failure, severe."

● Chronic Liver Disease: identified using ICD-10 codes in the HCUP software for "Liver

disease, mild," and "Liver disease, moderate to severe."

● Chronic Lung Disease: The CDC website stipulates that asthma is of concern "if it's

moderate to severe," implying mild asthma is not of concern. The HCUP software

includes codes for all degrees of severity of asthma in the definition of "Chronic

pulmonary disease." Thus, chronic lunch disease was identified using ICD-10 codes in

the HCUP software for "Chronic pulmonary disease," excluding any ICD-10 codes

beginning with J452 or J453 (mild intermittent or mild persistent asthma, respectively).

● Cystic Fibrosis: Identified by any ICD-10 code beginning with E84.

● Dementia or other neurological condition: identified using ICD-10 codes in the HCUP

software for "Dementia," "Neurological disorders affecting movement," "Seizures and

epilepsy," and "Other neurological disorders."

● Diabetes: identified using ICD-10 codes in the HCUP software for "Diabetes with chronic

complications" and "Diabetes without chronic complications."

● Disabilities: identified using ICD-10 codes in the HCUP software for "Paralysis" plus any

ICD-10 code beginning with Q (birth defects and chromosomal abnormalities). Note that

this omits many, possibly most, forms of disabilities, including non-congenital blindness

and deafness, cognitive impairments not due to chromosomal abnormalities, autism

spectrum disorders of unknown etiology, etc., but these are of dubious connection to

COVID-19.

- Heart conditions: identified using ICD-10 codes in the HCUP software for "Heart failure," the ICD-10 codes listed in the referenced study[50] for "Myocardial Infarction," and/or any ICD-10 code starting with any of these prefixes: A1884, A3282, A3681, A381, A395, A5203, B2682, B332, B376, B5881, C452, D8685, G130, G712, G713, G720, G721, G722, G7249, G7281, G7289, G729, G737, I01, I02, I05, I06, I07, I08, I09, I11, I13, I20, I23, I24, I25, I3, I4, I5, I70, I9713, J1082, J1182, O101, OO2912, O903, Q2, R570, S26, T82, and Z95.

- HIV: identified using ICD-10 codes in the HCUP software for "Acquired immune deficiency syndrome."

- Mental health conditions: identified using ICD-10 codes in the HCUP software for "Depression" and "Psychoses." Note that this may omit many other forms of mental illness, such as obsessive-compulsive disorder, post-traumatic stress syndrome, borderline personality disorder, etc. Note that there is overlap between conditions considered mental health conditions and those considered disabilities (such as autism spectrum disorders) as well as between mental health conditions and other medical conditions (such as substance abuse disorders).

- Overweight or obese: Subjects were considered to be overweight or obese if their BMI was ≥25. If multiple BMI or height-and-weight values were recorded in the database over time for a given subject, the value(s) used were those closest in time to the date of sample collection.

- Pregnancy or recent pregnancy: Electronic medical records of all female subjects under the age of 69 were searched for: ICD-10 codes starting with Z3A and records of hospital admissions which include a baby delivery time. The timespans of the pregnancy and puerperium periods were estimated from either type of record. In the case of ICD-10 codes starting with Z3A, the final digits of the ICD-10 code encode weeks of gestation at the time of the encounter, from which a start and end date of the pregnancy can be estimated. If only a delivery date is known, the pregnancy is estimated to have begun 40 weeks earlier, unless "PRETERM" is found in the free-text diagnosis. Subjects were marked as "pregnancy or recent pregnancy" only if their COVID-19 test date fell

between the estimated start date of the pregnancy and 42 days after the estimated end

date (to allow for post-term pregnancy). Where there was no COVID test date, the date

of the blood sample collection was used.

● Sickle cell or Thalassemia: Identified by any ICD-10 code beginning with D56 or D57.

● Smoking, current or former: Electronic medical records were searched for any non-zero

"Tobacco pack years," and for a free-text description of their tobacco usage including

the text "current smoker," "former," "every day," "some days," "light," "heavy," "less

than 10," "10+," "yes," or "counseling provided."

● Solid organ or blood stem cell transplant: Identified by any ICD-10 code beginning with

Z94.

● Stroke or cerebrovascular disease: identified using ICD-10 codes in the HCUP software

for "Cerebrovascular disease," which includes ICD-10 codes for both CBVD POA and

CBVD SQLA.

● Substance abuse: identified using ICD-10 codes in the HCUP software for "Drug abuse"

and for "Alcohol abuse."

● Tuberculosis: Identified by any ICD-10 code beginning with A15.

Sample collection, cell separation, and DNA extraction (Fig. S13-S14)

2mL aliquots were taken from EDTA-anticoagulated venous blood collected in the course of

standard clinical care (via "purple-top" tubes; Becton, Dickinson, Franklin Park, NJ). Tubes were

stored at 4°C between collection and processing, never more than 12 hours. Each aliquot was

mixed 1:1 dilution in phosphate-buffered saline (PBS) and centrifuged over Ficoll-Paque-plus

(Cytiva, Marlborough) to obtain peripheral blood mononuclear cells (PBMCs). Plasma was

collected and stored at 80°C. PBMCs were washed with PBS and resuspended in a sorting buffer

of PBS, 1% bovine serum albumin (BSA), and 0.01% sodium azide.

Magnetically-labeled anti-CD4 and anti-IgM microbeads (Miltenyi, Bergisch Gladbach) were

used to label and column-separate for infectee and control samples; vaccinee samples cells

were not separated. This process divided the samples into CD4-positive T cells and IgM-positive

B cells in one fraction, and CD4-negative T cells and IgM-negative B cells (principally CD8-

positive T cells and IgG-positive B cells) in the other fraction. DNA was isolated for each fraction

using EZ1&2 DNA Blood 350μL kits (Qiagen, Hilden) and the EZ1 Advanced XL automated

system (Qiagen, Hilden). DNA concentration was assessed via Nanodrop (Thermo Fisher,

Waltham).

Sequencing library preparation

AIRRseq libraries were generated using the immunoPETE method as described.[5] ImmunoPETE is

a two-step primer extension based targeted gene enrichment assay designed to specifically

target and quantitatively amplify recombined human TRB, TRD, and IGH from genomic DNA

simultaneously. Briefly, V gene-based primers containing unique molecular identifiers (UMI) as

well as universal PCR amplification handles were annealed to the chromosomal VDJ rearranged

loci. The first primer extension products, spanning the VDJ rearrangement, were purified from

any remaining oligos by a combination of beads (KAPA HyperPure, Roche) and enzymatic

treatment with Thermolabile Exonuclease I (New England Biolabs). A second primer extension

and amplification master mix containing a pool of J-gene oligos and an Illumina i7 primer

generated VDJ amplicons after 10 cycles of target amplification. Illumina sequencing library

amplification was performed using the i7/i5 primer pairs with dual sample indexes. All primer

extensions and amplifications were performed using the KAPA Long Range HotStart Ready Mix

(Roche). The resulting libraries underwent purification using KAPA HyperPure beads (Roche),

followed by quantification with the Qubit dsDNA HS Assay kit (Thermo Fisher) and fragment

analysis (Agilent TapeStation). Individual sample libraries were pooled in equal mass. A final

round of quantification and fragment analysis was then performed. Finally, libraries were

sequenced using the Illumina NextSeq 500/550 High Output Kit v2.5 (300 cycles).

Sequencing and bioinformatics

ImmunoPETE sequencing libraries were analyzed using the Roche in-house bioinformatics

pipeline, Daedalus (https://github.com/bioinform/Daedalus). After quality filtering of reads and

trimming off primers, the pipeline identified V and J genes using a Smith-Waterman alignment

approach (https://github.com/pgngp/swift) against an in-house curated V and J gene database.

Original V and J gene data and sequences were sourced from HGNC

(https://www.genenames.org/) and ENSEMBL (https://ensemblgenomes.org/). CDR3

sequences were identified for all V-J pairs, capturing both functional (functional V/J gene AND

coding CDR3) and non-functional (annotated non-functional or pseudogene V/J gene in the

database OR stop codon/frameshift in CDR3) rearrangements. Sequences are deduplicated by

clustering UMI and CDR3 sequences to identify UMI families. Consensus sequences were

derived for the CDR3 and UMI segments of each UMI family, suppressing sequencing and PCR

errors, and identifying CDR3 rearrangements at single molecule resolution. High quality CDR3

rearrangements were further analyzed for cell counting, clonal diversity, and other calculations.

Terms used are listed alphabetically and defined as follows:

●   Cell count: the total number of functional IGH, TRD, and TRB rearrangements in a

651   sample

652   ●   Cell type percentages: the total number of functional rearrangements from each heavy

653   chain divided by the total cell count × 100

654   ●   CDR3 clone: BCR or TCR sequences from the same individual with matching V gene,

655   CDR3 amino acid sequence (CDR3-AA), and J gene assignment arising from two or more

656   UMI families

657   ●   CDR3 clonal type: BCR or TCR sequences from multiple UMI families from multiple

658   individuals with matching V gene, CDR3-AA, and J gene assignment

659   ●   Clone count: total number of UMI families from the same individual with the same V

660   gene, CDR3-AA, and J gene

661   ●   UMI family: a set of reads that have been clustered together based on the similarities of

662   the 9-nt UMI sequence and the CDR3-nt region

Both UMI and CDR3 sequences are clustered based on a Levenshtein edit distance of 1,

capturing likely PCR and sequencing errors. A UMI family represents a single captured DNA

molecular fragment from the immunoPETE reaction.

NAbs ELISA titers

The SARS-CoV-2 Surrogate Virus Neutralization Test Kit (GenScript, L00847-A) is designed to

quantify the circulating antibodies capacity of blocking the interaction between the receptor

binding domain of SARS-CoV-2 spike glycoprotein and the ACE2 human cell surface receptor.

The kit was used according to the manufacturer's instructions as follows. A standard curve was

generated using a serial dilution of the standard (GenScript, A02087-20) with a dilution factor of

1:2. Each subject's serum sample was mixed with sample dilution buffer (1:10) and horseradish

peroxidase-conjugated recombinant SARS-CoV-2 receptor-binding domain (HRP-RBD). The

mixture was incubated at 37°C for 30 min to allow the circulating NAbs to bind to HRP-RBD. The

mixture was then added to an ACE2 protein-coated plate and incubated for an additional 15

min at 37°C. Unbound HRP-RBD and HRP-RBD bound to non-neutralizing antibodies were

bound to the plate while circulating neutralization antibody HRP-RBD complexes remained in

the supernatant for subsequent wash steps. After washing, tetramethylbenzidine solution was

added, followed by a stop solution to quench the reaction, turning wells yellow. The plate was

read immediately at 450 nm in a microtiter plate reader. Statistical analysis was performed with

GraphPad Prism using a 4PL model for linear regression. Results were reported by interpolating

the OD450 values to the standard curve values.

Results are reported in the WHO standard units (BAU/mL), which can be converted to the

manufacturer's U/mL by dividing numbers by a factor of 1.626 (per the manufacturer).

pymmunomics

Code used for the analyses was written up as a python package and made publicly available on

github (https://www.github.com/ArnaoutLab/pymmunomics). Reference is made in the

following sections wherever that is the case.

Validation of pooled sequences from sorted subsets vs. unsorted sequences

Additional test samples were split into two replicates. One replicate was sorted into IgM/CD4

and IgG/CD8 fractions as above. The other replicate was unsorted. Each of the IgM/CD4,

IgG/CD8, and unsorted fractions had libraries prepared as above and were sequenced as above.

V- and J-gene frequencies from the IgM/CD4 and IgG/CD8 fractions and from the unsorted

fraction were measured, and the frequencies from the IgM/CD4 and IgG/CD8 fractions were

averaged together. These averages were compared to the frequencies in the unsorted fraction,

and were shown to be indistinguishable by t-test (p-values ranging from 0.19 to 0.95 for

comparisons of the frequencies of each gene), demonstrating that IgM/CD4 + IgG/CD8 is

statistically indistinguishable from unsorted, supporting comparisons between infectees

(sorted) and vaccinees (unsorted) and between controls (sorted) and vaccinees. Dependence of

antibody concentrations on age, immunocompetence, and SARS-CoV-2 exposure

Univariate and bivariate exploratory plots suggested zero antibody concentration to be a

special category. Therefore, we first modeled the ability to produce zero vs. non-zero amounts

of antibody using logistic regression. We then performed linear regression to model the $\log_{10}$-

transformed concentration of the nonzero values on our set of covariates. In both cases, we

started with a full model incorporating age, immunocompetence status, cohort, and all of their

two-way and three-way interactions. Starting with the interaction terms and then proceeding

to the main effects, we sequentially eliminated covariates that were not significant at $\alpha=0.05$.

This did not change the regression coefficients of any of the significant terms by >20% (i.e. were

not confounders). Finally, we confirmed that the best model had lower AIC (logistic regression)

or higher adjusted R^2 (linear regression) compared with the alternative models.

CDR3 length analysis

For calculating lengths, the CDR3 region was defined to include the preceding Cysteine anchor

residue and the terminating Tryptophan (for IGH) or Phenylalanine (for TCR) anchor residue.

CDR3 length frequencies for each available functional and non-functional pooled IGH, TRB, TRD,

and subtyped IGG, IGM, CD4 TRB/D, CD8 TCB/D repertoire of immunocompetent subjects were

calculated using the pymmunomics python package (above). Since vaccinee samples were not

sorted into subtypes, pooled repertoire CDR3 length frequency distributions were used to

compare vaccinees to controls and infectees. CD4$^+$/IGM$^+$ and CD8/IGG repertoire CDR3 length

frequency distributions were compared independently between controls to infectees.

To compare CDR3 length distributions between cohorts without simplifying them down to their

mean or median distribution, which ignores variance within groups, we chose a threshold CDR3

length ℓ and compared the cumulative frequencies of sequences on each side of that length

using a two-tailed Mann-Whitney-U test. The threshold length was determined by estimating

the difference of length frequencies between cohorts for each CDR3 length. These estimates

were calculated by taking the median difference in frequency between members of one cohort

and members of the other. The dividing line is then placed between the lengths ℓ and $\ell+1$,

where ℓ is the CDR3 length that maximizes the magnitudes of the total areas under the curve of

estimated frequency differences to the left and right of the line, i.e. the best dividing line

between patterns:

730
$$\left| \sum_{\ell' < \ell} d_{\ell'} \right| + \left| \sum_{\ell' > \ell} d_{\ell'} \right|$$

Here $d_{\ell'}$ denotes the estimated difference of frequencies of CDR3s of length ℓ' between the two

cohorts. Note that the absolute values are taken after summing group differences on one side

of the dividing line (making positive and negative differences cancel each other out before

taking the absolute value), favoring a dividing line that splits the median differences into large

same-signed runs. P-values were corrected for multiple hypotheses via the Holm-Bonferroni

method (Table S3).

To identify trends among lengths of V and J genes, V and J genes (from IMGT) of the relevant

cell types (IGH for the functional pooled IGH comparisons and pooled IGH, IGG, and IGM for the

non-functional comparisons) which had a corrected p-value below 0.05 were grouped into the

number of residues that fall into the CDR3 region. Usage frequencies of V- and J-gene groups

were compared between cohorts using two-tailed Mann-Whitney-U and a second correction

round was conducted to correct all original p-values of the CDR3 length comparisons at the

same time as the p-values obtained from the follow-up tests.

Sets of known SARS-CoV-2 binders and binders to other pathogens

MIRA-identified SARS-CoV-2 specific T-cell receptor sequences[23] were downloaded from

https://clients.adaptivebiotech.com/pub/covid-2020 on April 19, 2021.

Query B-cell and T-cell receptor sequences (CDR3) of cells known to bind to SARS-CoV-2 were

downloaded from CoVAbDab, PDB, and VDJDB. The CoVAbDab sequences were downloaded on

April 20, 2022 and consists of all SARS-CoV-2-WT-neutralizing human antibodies with CDRH3

sequence listed in the database at the time and added since May 04, 2020. PDB sequences

were download on May 03, 2022 searching for all structures of source organism Homo sapiens,

containing in the title one of "antibody" or "Fab," and one of "CMV," "cytomegalovirus,"

"DENV" (i.e. dengue virus), "dengue," "EBV," "Epstein-Barr," "hepatitis," "HIV," "human

immunodeficiency virus," "influenza," "SARS-CoV-2," or "tetanus." The resulting entries were

filtered for sequences in which a CDRH3 sequence of length at least 6 and at most 40 could be

detected using in-house Python code. For each sequence, the name of the binding target was

extracted from the structure title. VDJDB sequences were also downloaded on April 20, 2022 to

obtain human TRB sequences with CDR3 and J-gene specified that bind to their listed target

with a non-zero score.

To conform with the gene database used for V- and J-gene assignment of repertoire sequences

(see Sequencing and bioinformatics), the same gene sequences were aligned (blastp and blastp-

short for V genes and J genes, respectively; BLAST+ v2.12.0) to the sequences from PDB and

CoVAbDab, setting the max target seqs parameter to 10,000—a number much larger than the

total number of genes in the query to avoid missing the best matching genes.[57] V-gene matches

with query coverage less than 30% or percent identity less than 40% and J-gene matches with

query coverage less than 50% or percent identity less than 40% were filtered out. From the

remainder, the best V- and J-gene matches according to percent identity and gene sequence

coverage (lexicographically) were assigned to each query sequence. Data downloaded from

VDJDB contained sequence only for the CDR3 region, so the V, and J-gene annotation provided

by the database was used (as opposed to using e.g. BLAST).

To calculate the fractions of query sequences sets matching subject repertoire sequences and

the fractions of subject TRB repertoires matching query TRB sequences sets, a pair of sequences

is considered to match if their V gene, J gene, and CDR3 sequence are identical.

Binding-capacity measurements

Binding capacities to the MIRA-identified HLA class II T-cell sequences were measured for all

subject pooled (CD4+CD8), and CD4 TRB repertoires, wherever possible. The binding capacity of

a repertoire R to a clone c is defined as:

$$\tau(c; R) = \sum_{c' \in R} p(c') \cdot s(c, c')$$

779   where $p(c')$ denotes the frequency of clone c' in repertoire R and s is the binding similarity

780   between sequences. Here, s as previously described,[43] which accounted only for the

781   relationship between Levenshtein distance of CDR3s and the predicted difference in strength of

782   their binding to the same target(s) (in terms of relative K_d), was constrained as follows to

783   require matching V and/or J genes:

$$s(c, c') = \begin{cases} 0.3^{Lev(c,c')} & \text{if V and J genes match} \\ 0 & \text{otherwise} \end{cases}$$

784   Here, $Lev(c, c')$ is the Levenshtein distance between the CDR3 amino acid sequences of

785   sequences c and c'. The pymmunomics Python package was used to calculate similarity

786   matrices and binding capacities.

Fuzzy query sequence matching

Fuzzy sequence matching measurements for each pooled CD4+CD8 and each CD4-only TRB

subject repertoire to the MIRA-identified HLA class II query sequences were tabulated from the

similarity matrices that are calculated as part of determining binding capacities. For each query

sequence and each subject repertoire, we measured the fraction of repertoire sequences for

with the same V and J genes as the query sequence, and whose CDR3 sequence was within

Levenshtein distances 0-10 of the query's CDR3. Note that exact matching is equivalent to fuzzy

matching with a Levenshtein distance of 0.

Binding-capacity and fuzzy-matching robustness experiments

To compare robustness to variations in repertoire size of binding capacity and fuzzy matching

features, we conducted subsampling experiments. We randomly chose 10 subjects from each

of the vaccinee, infectee, and control cohorts that had a pooled TRB repertoire size of at least

80,000 cells, i.e. 80,000 distinct corrected UMIs. (This size was chosen in order to guarantee at

least 10 subjects from the control cohort to choose from.) Each of these repertoires was

sampled down to 20 different subsample sizes chosen to be equidistantly spaced between 10

and 80,000 at log-scale. For each subsample, we calculated binding capacities as well as fraction

of fuzzy matches for fuzzy-match tolerances 0, 2, 4, 6, 8, and 10 amino acids to CD4 TRB

reference sequences from MIRA. The slopes and their surrounding 95% confidence intervals

were obtained by fitting a linear mixed model that groups the data by subject.

Feature selection

Preferring the use of domain knowledge over generic feature selection mechanisms for

selecting from the high-dimensional query sequence matching features (binding capacity and

fuzzy matching), a custom feature selection method is developed and implemented in the

python package pymmunomics. For this mechanism we use binding capacity and fuzzy

matching measurements to sequence specific to pathogens other than SARS-CoV-2 ("SARS-CoV-

2 non-specific sequences") as a null distribution to which to compare the measurements for

MIRA-identified SARS-CoV-2-specific sequences. We calculated the (Stuart-)Kendall Tau-c

correlation coefficient between each feature's measurement and NAb titer. For each feature

group (binding capacity, fuzzy matching with tolerances 0, 1, …, 10, etc.), the correlation

coefficients of measurements for non-SARS-CoV-2 specific sequences form the null distribution

and correlation coefficients of SARS-CoV-2 specific features below the 2.5^{th} and above the

97.5^{th} percentile are selected (cumulatively, the most correlated and anti-correlated 5%).

Following the same idea, V-gene frequencies were also selected from among the 54 total

possibilities (one for each V gene). Here, V-gene frequencies in non-functional repertoires were

taken as the null distribution against which to compare functional repertoires' V-gene

frequencies, since non-functional sequences do not undergo SARS-CoV-2 specific clonal

expansion. Since the functional and non-functional frequencies can be viewed as paired

measurements, the distribution of differences between their correlation coefficients was

calculated, and the most correlated and anti-correlated 5% (as defined above) were selected as

features.

Machine learning to classify subjects with a protective NAb titer

Machine learning classifiers of high or low neutralizing antibody concentration were fit to

various feature groups and for various cell types. For the CD4 and pooled TRB receptor

repertoires, binding capacities as well as fuzzy matching features with tolerances 0, 1, …, 10 to

the MIRA-identified CD4 clones from Nolan et al.[23] were used. Another set of models was

derived from these by adding a mechanism at the end of feature selection that aggregates the

selected features into their sums. For the pooled IGH, TRB, and TRD as well as the IGM, non-

IGM (predominately IGG), CD4 TRB and CD8 TRB repertoires models are fit on the following

feature sets:

● CDR3 length frequencies, summarized by 3 features: mean, variance and skewness;

● diversity, with Recon[58] (https://github.com/ArnaoutLab/Recon) being used to correct

Hill D_q numbers for $q=0, 1, …, \infty$ to correct for missing species;

●   J-gene frequencies (with only 6 J genes, no further feature selection was required);

●   V-gene frequencies for select V genes as described above;

●   Baseline/clinical features: age, sex, days since infection (runs of positive COVID-19 PCR

tests successively within 28 days of each other and not interrupted by negative tests are

considered infected periods; to account for incubation of the virus prior to taking the

test, the start date of an infection is predicted as 4 days before the first positive test in

the corresponding run of tests; when a negative test was performed within those 4

days, that test's date is considered the infection start date; for the model, the predicted

start date of the most recent infection before sample collection was used, or 0 if the

subject was not infected), and days since vaccination (the number of days between

sample collection and most recent vaccination on record).

The machine learning framework was set up as follows. For each feature group, 700 replicate

performances were measured via repetition of 7-fold cross-validation 100 times, each time

choosing a different split of the data into 7 folds at random. For each replicate, 10-fold cross-

validation was used to tune hyperparameters via Bayesian optimization. For each model fit, the

training data was standardized, then underwent principal component analysis, and finally was

used to train an L2-regularized regression. There were two tuned hyperparameters:

regularization strength (with a log-uniform search space distribution between 10^{-8} and 10^{-2})

and the amount of variance to be explained by chosen principal components (with uniform

search space distribution between 0.50 and 0.99; e.g. if the value was 0.75 and the first four

PCs account for 75% of variance, these four PCs would be chosen). For feature sets relating to

similarity—binding capacities, fuzzy-matching features at various tolerances, and their

aggregated versions—and for V-gene features, feature selection was performed on the training

data before standardization for each model fit. To facilitate avoidance of train-test leakage, the

mechanisms are implemented in the pymmunomics python package to fit into the popular

scikit-lean API framework.

Data availability

All raw data for this study can be found at: https://data.entrop.ai/cdr3/data/index.html

Figure 1: Anti-SARS-CoV-2 ELISA trends and distributions by age for immunocompetent and immunosuppressed vaccinees, infectees, and controls

NAbs target SARS-CoV-2 spike protein. (a) ELISA titers for each subject. Solid lines indicate regression fits; shaded areas indicate 95% confidence intervals. Dotted black line at ~10^3 indicates manufacturer's cutoff for positive vs. negative. Note strong negative trend with age in vaccinees (blue) but not infectees (salmon). Note mild positive trend with age in controls (olive), even as titers in this cohort remain below the cutoff for almost all individuals. (b)

876   Distribution of titers in the three cohorts, split by immune status. (c) Distribution of ages in

877   these cohorts, again split by immune status, with numbers of subjects in each sub-cohort.

Figure 2: IGH CDR3 length distributions

(a) CDR3 length comparison plots for productive IGH repertoires of vaccinees vs. controls. Left

inset: the median differences of frequencies at each length, showing that CDR3s of length 16 or

shorter are more frequent in vaccinees, whereas CDR3s of length 17 or longer are less frequent.

The pattern reverses at the dividing line between 16 and 17 amino acids (vertical dotted line).

Right inset: total fraction of the repertoire up to the dividing line. The p-value is obtained by

applying Mann-Whitney U to the cumulatives followed by Holm-Bonferroni multiple-hypothesis

correction. **(b)** The same for vaccinees vs. infectees, showing the same pattern but with a

dividing line between 18 and 19 amino acids. **(c)**-**(d)** Frequencies of V and J genes grouped by

the number of residues each gene contributes to the CDR3 according to germline (Fig. S8). Note

the only J gene that contributes 5 residues is IGHJ4. **(e)** Assuming the nonproductive IGH

vaccinee repertoire (blue) is made up of a part that is unaffected by vaccination and therefore

looks like the control repertoire (green) and a part that is affected by vaccination (salmon), this

plot shows what the distribution of the affected part would have to look like so the two parts

add up correctly, for different fractions affected (dark to light salmon lines). Estimated means

for vaccinee and control distributions are shown. The smaller the affected portion, the more

extreme the effect must be. The minimum possible effect size is that for which a CDR3 length

for the affected portion is zero; any smaller, and a negative frequency at that CDR3 length

would be required (negative frequencies are not possible). Boxes show the 25th and 75th

percentiles. Horizontal lines show medians. Whiskers show the min and max unless there are

outliers (diamonds).

Figure 3: SARS-CoV-2-specific TRBs vs. NAb titers

(a)-(b) Fraction of TRB repertoires matching the SARS-CoV-2-specific CD4 TRB sequences

obtained from Nolan et al.[23] against SARS-CoV-2 NAb titer. Panel (a) shows repertoires from

CD4+ T cells, which were available for infectees and controls but not vaccinees, while panel (b)

shows repertoires from all T cells, which were available for all three cohorts. Theil-Sen

regression fits (solid lines) show positive relationships for infectees and vaccinees but not

controls. **(c)** The fraction of a repertoire that matches reference TRBs within a chosen tolerance

(here, 2 amino-acid differences) depends strongly on the number of cells in the repertoire (i.e.,

repertoire size; fig. S10). **(d)** In contrast, binding capacity is much more robust. The slope of the

dependency on size for repertoires above 1,000 cells are shown as black lines. **(e)** Slope as a

function of fuzzy-binding tolerance, demonstrating binding capacity is more robust regardless

of tolerance.

Figure 4: Predicting positive NAbs

(a) Feature selection mechanism used for binding capacity and fuzzy matching features on the

binding capacity measurements of all TCR repertoires of size at least 1,000 using the SARS-CoV-

2-specific CD4 TCR sequences and non-SARS-CoV-2-specific TCR sequences obtained from

VDJDB. Of the 7,804 SARS-CoV-2-specific features' correlations, 323 fall outside the selection

boundaries set by the 95% boundaries of the correlations of non-SARS-CoV-2-specific features

with NAb titer. **(b)-(c)** Machine learning performance results for a selected group of feature sets

and cell types across all 700 replicates (100 repeats of 7-fold cross-validation). (a) shows areas

under receiver operating curves and (b) breaks down the performances into sensitivity,

precision, and specificity. The same plots for all feature sets and cell types can be found in Fig.

S11 and S12. Median values and interquartile ranges for all metrics are reported in Table S8.

Boxes and whiskers as in Fig. 2.

References

References

1. Arnaout, R.A., Prak, E.T.L., Schwab, N., Rubelt, F., and the Adaptive Immune Receptor
Repertoire Community (2021). The Future of Blood Testing Is the Immunome. Front.
Immunol. *12*, 626793. 10.3389/fimmu.2021.626793.

2. Primorac, D., Vrdoljak, K., Brlek, P., Pavelić, E., Molnar, V., Matišić, V., Erceg Ivkošić, I., and
Parčina, M. (2022). Adaptive Immune Responses and Immunity to SARS-CoV-2. Frontiers in
Immunology *13*.

3. Shen, J., Fan, J., Zhao, Y., Jiang, D., Niu, Z., Zhang, Z., and Cao, G. (2023). Innate and adaptive
immunity to SARS-CoV-2 and predisposing factors. Front Immunol *14*, 1159326.
10.3389/fimmu.2023.1159326.

4. Joseph, M., Wu, Y., Dannebaum, R., Rubelt, F., Zlatareva, I., Lorenc, A., Du, Z.G., Davies, D.,
Kyle-Cezar, F., Das, A., et al. (2022). Global patterns of antigen receptor repertoire
disruption across adaptive immune compartments in COVID-19. Proc Natl Acad Sci U S A
119, e2201541119. 10.1073/pnas.2201541119.

5. Dannebaum, R., Suwalski, P., Asgharian, H., Du Zhipei, G., Lin, H., Weiner, J., Holtgrewe, M.,
Thibeault, C., Müller, M., Wang, X., et al. (2022). Highly multiplexed immune repertoire
sequencing links multiple lymphocyte classes with severity of response to COVID-19.
EClinicalMedicine *48*, 101438. 10.1016/j.eclinm.2022.101438.

6. Fujihashi, K., McGhee, J., Yamamoto, M., Hiroi, T., and Kiyono, H. (1996). Role of gamma
delta T cells in the regulation of mucosal IgA response and oral tolerance. Ann N Y Acad Sci
778, 55–63. 10.1111/j.1749-6632.1996.tb21114.x.

7. Rezende, R.M., Cox, L.M., Moreira, T.G., Liu, S., Boulenouar, S., Dhang, F., LeServe, D.S.,
Nakagaki, B.N., Lopes, J.R., Tatematsu, B.K., et al. (2023). Gamma-delta T cells modulate the
microbiota and fecal micro-RNAs to maintain mucosal tolerance. Microbiome *11*, 32.
10.1186/s40168-023-01478-1.

8. Brouwer, P.J.M., Caniels, T.G., van der Straten, K., Snitselaar, J.L., Aldon, Y., Bangaru, S.,

Torres, J.L., Okba, N.M.A., Claireaux, M., Kerster, G., et al. (2020). Potent neutralizing

antibodies from COVID-19 patients define multiple targets of vulnerability. Science *369*,

643–650. 10.1126/science.abc5902.

9. Cao, Y., Su, B., Guo, X., Sun, W., Deng, Y., Bao, L., Zhu, Q., Zhang, X., Zheng, Y., Geng, C., et

al. (2020). Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-

Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. Cell *182*, 73-84.e16.

10.1016/j.cell.2020.05.025.

10. Chen, E.C., Gilchuk, P., Zost, S.J., Suryadevara, N., Winkler, E.S., Cabel, C.R., Binshtein, E.,

Chen, R.E., Sutton, R.E., Rodriguez, J., et al. (2021). Convergent antibody responses to the

SARS-CoV-2 spike protein in convalescent and vaccinated individuals. Cell Reports *36*,

109604. 10.1016/j.celrep.2021.109604.

11. Cerutti, G., Guo, Y., Zhou, T., Gorman, J., Lee, M., Rapp, M., Reddem, E.R., Yu, J., Bahna, F.,

Bimela, J., et al. (2021). Potent SARS-CoV-2 neutralizing antibodies directed against spike N-

terminal domain target a single supersite. Cell Host & Microbe *29*, 819-833.e7.

10.1016/j.chom.2021.03.005.

12. Galson, J.D., Schaetzle, S., Bashford-Rogers, R.J.M., Raybould, M.I.J., Kovaltsuk, A.,

Kilpatrick, G.J., Minter, R., Finch, D.K., Dias, J., James, L., et al. (2020). Deep sequencing of B

cell receptor repertoires from COVID-19 patients reveals strong convergent immune

signatures. bioRxiv, 2020.05.20.106294. 10.1101/2020.05.20.106294.

13. Lima, N.S., Mukhamedova, M., Johnston, T.S., Wagner, D.A., Henry, A.R., Wang, L., Yang,

E.S., Zhang, Y., Birungi, K., Black, W.P., et al. (2022). Convergent epitope specificities, V gene

usage and public clones elicited by primary exposure to SARS-CoV-2 variants. bioRxiv,

2022.03.28.486152. 10.1101/2022.03.28.486152.

14. McCallum, M., De Marco, A., Lempp, F.A., Tortorici, M.A., Pinto, D., Walls, A.C., Beltramello,

M., Chen, A., Liu, Z., Zatta, F., et al. (2021). N-terminal domain antigenic mapping reveals a

site of vulnerability for SARS-CoV-2. Cell *184*, 2332-2347.e16. 10.1016/j.cell.2021.03.028.

15. Rapp, M., Guo, Y., Reddem, E.R., Yu, J., Liu, L., Wang, P., Cerutti, G., Katsamba, P., Bimela,

J.S., Bahna, F.A., et al. (2021). Modular basis for potent SARS-CoV-2 neutralization by a

prevalent VH1-2-derived antibody class. Cell Reports *35*, 108950.

10.1016/j.celrep.2021.108950.

16. Wec, A.Z., Wrapp, D., Herbert, A.S., Maurer, D.P., Haslwanter, D., Sakharkar, M., Jangra,

R.K., Dieterle, M.E., Lilov, A., Huang, D., et al. (2020). Broad neutralization of SARS-related

viruses by human monoclonal antibodies. Science *369*, 731–736. 10.1126/science.abc7424.

17. Manso, T., Folch, G., Giudicelli, V., Jabado-Michaloud, J., Kushwaha, A., Nguefack Ngoune,

V., Georga, M., Papadaki, A., Debbagh, C., Pégorier, P., et al. (2022). IMGT® databases,

related tools and web resources through three main axes of research and development.

Nucleic Acids Research *50*, D1262–D1272. 10.1093/nar/gkab1136.

18. Simnica, D., Schultheiß, C., Mohme, M., Paschold, L., Willscher, E., Fitzek, A., Püschel, K.,

Matschke, J., Ciesek, S., Sedding, D.G., et al. (2021). Landscape of T-cell repertoires with

public COVID-19-associated T-cell receptors in pre-pandemic risk cohorts. Clinical &

Translational Immunology *10*, e1340. 10.1002/cti2.1340.

19. Ellwardt, E., Walsh, J.T., Kipnis, J., and Zipp, F. (2016). Understanding the Role of T Cells in

CNS Homeostasis. Trends in Immunology *37*, 154–165. 10.1016/j.it.2015.12.008.

20. Matschke, J., Lütgehetmann, M., Hagel, C., Sperhake, J.P., Schröder, A.S., Edler, C.,

Mushumba, H., Fitzek, A., Allweiss, L., Dandri, M., et al. (2020). Neuropathology of patients

with COVID-19 in Germany: a post-mortem case series. The Lancet Neurology *19*, 919–929.

10.1016/S1474-4422(20)30308-2.

21. Schwabenland, M., Salié, H., Tanevski, J., Killmer, S., Lago, M.S., Schlaak, A.E., Mayer, L.,

Matschke, J., Püschel, K., Fitzek, A., et al. (2021). Deep spatial profiling of human COVID-19

brains reveals neuroinflammation with distinct microanatomical microglia-T-cell

interactions. Immunity *54*, 1594-1610.e11. 10.1016/j.immuni.2021.06.002.

22. Arora, R., and Arnaout, R. (2020). Private Antibody Repertoires Are Public.

10.1101/2020.06.18.159699.

23. Nolan, S., Vignali, M., Klinger, M., Dines, J., Kaplan, I., Svejnoha, E., Craft, T., Boland, K.,

Pesesky, M., Gittelman, R., et al. (2020). A large-scale database of T-cell receptor beta

(TCRβ) sequences and binding associations from natural and synthetic exposure to SARS-

CoV-2. 10.21203/rs.3.rs-51964/v1.

24. Swanson, P.A., Padilla, M., Hoyland, W., McGlinchey, K., Fields, P.A., Bibi, S., Faust, S.N.,

McDermott, A.B., Lambe, T., Pollard, A.J., et al. (2021). AZD1222/ChAdOx1 nCoV-19

vaccination induces a polyfunctional spike protein–specific T $_H$ 1 response with a diverse

TCR repertoire. Sci. Transl. Med. *13*, eabj7211. 10.1126/scitranslmed.abj7211.

25. Bates, T.A., Leier, H.C., Lyski, Z.L., Goodman, J.R., Curlin, M.E., Messer, W.B., and Tafesse,

F.G. (2021). Age-Dependent Neutralization of SARS-CoV-2 and P.1 Variant by Vaccine

Immune Serum Samples. JAMA *326*, 868–869. 10.1001/jama.2021.11656.

26. Bonfante, F., Costenaro, P., Cantarutti, A., Di Chiara, C., Bortolami, A., Petrara, M.R.,

Carmona, F., Pagliari, M., Cosma, C., Cozzani, S., et al. (2021). Mild SARS-CoV-2 Infections

and Neutralizing Antibody Titers. Pediatrics *148*, e2021052173. 10.1542/peds.2021-052173.

27. Dyer, A.H., Noonan, C., McElheron, M., Batten, I., Reddy, C., Connolly, E., Pierpoint, R.,

Murray, C., Leonard, A., Higgins, C., et al. (2022). Previous SARS-CoV-2 Infection, Age, and

Frailty Are Associated With 6-Month Vaccine-Induced Anti-Spike Antibody Titer in Nursing

Home Residents. Journal of the American Medical Directors Association *23*, 434–439.

10.1016/j.jamda.2021.12.001.

28. Sadighi Akha, A.A. (2018). Aging and the immune system: An overview. Journal of

Immunological Methods *463*, 21–26. 10.1016/j.jim.2018.08.005.

29. Haveri, A., Ekström, N., Solastie, A., Virta, C., Österlund, P., Isosaari, E., Nohynek, H., Palmu,
A.A., and Melin, M. (2021). Persistence of neutralizing antibodies a year after SARS-CoV-2
infection in humans. European Journal of Immunology *51*, 3202–3213.
10.1002/eji.202149535.

30. Korobova, Z.R., Zueva, E.V., Arsentieva, N.A., Batsunov, O.K., Liubimova, N.E., Khamitova,
I.V., Kuznetsova, R.N., Rubinstein, A.A., Savin, T.V., Stanevich, O.V., et al. (2022). Changes in
Anti-SARS-CoV-2 IgG Subclasses over Time and in Association with Disease Severity. Viruses
14, 941. 10.3390/v14050941.

31. Luo, Y.R., Chakraborty, I., Yun, C., Wu, A.H.B., and Lynch, K.L. (2021). Kinetics of Severe
Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Antibody Avidity Maturation and
Association with Disease Severity. Clinical Infectious Diseases *73*, e3095–e3097.
10.1093/cid/ciaa1389.

32. Moss, P. (2022). The T cell immune response against SARS-CoV-2. Nat Immunol *23*, 186–
193. 10.1038/s41590-021-01122-w.

33. Morgan, A., Contreras, E., Yasuda, M., Dutta, S., Hamel, D., Shankar, T., Balallo, D., Riedel,
S., Kirby, J.E., Kanki, P.J., et al. (2023). The Coviral Portal: Multi-Cohort Viral Loads and
Antigen-Test Virtual Trials for COVID-19. 10.1101/2023.05.05.23289582.

34. Brown, A.J., Snapkov, I., Akbar, R., Pavlović, M., Miho, E., Sandve, G.K., and Greiff, V. (2019).
Augmenting adaptive immunity: progress and challenges in the quantitative engineering
and analysis of adaptive immune receptor repertoires. Mol. Syst. Des. Eng. *4*, 701–736.
10.1039/C9ME00071B.

35. Brown, S.D., Raeburn, L.A., and Holt, R.A. (2015). Profiling tissue-resident T cell repertoires
by RNA sequencing. Genome Med *7*, 125. 10.1186/s13073-015-0248-x.

36. Huang, H., Wang, C., Rubelt, F., Scriba, T.J., and Davis, M.M. (2020). Analyzing the
Mycobacterium tuberculosis immune response by T-cell receptor clustering with GLIPH2

and genome-wide antigen screening. Nat Biotechnol *38*, 1194–1202. 10.1038/s41587-020-

0505-4.

37. Mayer, A. and Curtis G. Callan (2023). Measures of epitope binding degeneracy from T cell

receptor repertoires. Proceedings of the National Academy of Sciences *120*, e2213264120.

10.1073/pnas.2213264120.

38. Kaplinsky, J., Li, A., Sun, A., Coffre, M., Koralov, S.B., and Arnaout, R. (2014). Antibody

repertoire deep sequencing reveals antigen-independent selection in maturing B cells. Proc

Natl Acad Sci U S A *111*, E2622-9. 10.1073/pnas.1403278111.

39. Murphy, K., and Weaver, C. (2016). Janeway's Immunobiology (Garland Science).

40. Raybould, M.I.J., Kovaltsuk, A., Marks, C., and Deane, C.M. (2020). CoV-AbDab: the

Coronavirus Antibody Database. bioRxiv, 2020.05.15.077313. 10.1101/2020.05.15.077313.

41. Berman, H.M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T.N., Weissig, H., Shindyalov, I.N.,

and Bourne, P.E. (2000). The Protein Data Bank. Nucleic Acids Res. *28*, 235–242.

42. Goncharov, M., Bagaev, D., Shcherbinin, D., Zvyagin, I., Bolotin, D., Thomas, P.G., Minervina,

A.A., Pogorelyy, M.V., Ladell, K., McLaren, J.E., et al. (2022). VDJdb in the pandemic era: a

compendium of T cell receptors specific for SARS-CoV-2. Nature Methods *19*, 1017–1019.

10.1038/s41592-022-01578-0.

43. Arora, R., and Arnaout, R. (2022). Repertoire-scale measures of antigen binding. Proc. Natl.

Acad. Sci. U.S.A. *119*, e2203505119. 10.1073/pnas.2203505119.

44. Arnaout, R.A., and Nowak, M.A. (2000). Competitive coexistence in antiviral immunity. J

Theor Biol *204*, 431–441. 10.1006/jtbi.2000.2027.

45. Arnaout, R., Lee, R.A., Lee, G.R., Callahan, C., Cheng, A., Yen, C.F., Smith, K.P., Arora, R., and

Kirby, J.E. (2021). The Limit of Detection Matters: The Case for Benchmarking Severe Acute

Respiratory Syndrome Coronavirus 2 Testing. Clinical Infectious Diseases *73*, e3042–e3046.

10.1093/cid/ciaa1382.

46. Callahan, C., Lee, R.A., Lee, G.R., Zulauf, K., Kirby, J.E., and Arnaout, R. (In press). Nasal Swab
Performance by Collection Timing, Procedure, and Method of Transport for Patients with
SARS-CoV-2. J Clin Microbiol.

47. Callahan, C., Ditelberg, S., Dutta, S., Littlehale, N., Cheng, A., Kupczewski, K., McVay, D.,
Riedel, S., Kirby, J.E., and Arnaout, R. (2021). Saliva is Comparable to Nasopharyngeal Swabs
for Molecular Detection of SARS-CoV-2. Microbiol Spectr *9*, e0016221.
10.1128/Spectrum.00162-21.

48. Liu, J., Chandrashekar, A., Sellers, D., Barrett, J., Jacob-Dolan, C., Lifton, M., McMahan, K.,
Sciacca, M., VanWyk, H., Wu, C., et al. (2022). Vaccines elicit highly conserved cellular
immunity to SARS-CoV-2 Omicron. Nature *603*, 493–496. 10.1038/s41586-022-04465-y.

49. People with Certain Medical Conditions (2023). Centers for Disease Control and Prevention.
https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-
medical-conditions.html.

50. ELIXHAUSER COMORBIDITY SOFTWARE REFINED FOR ICD-10-CM Agency for Healthcare
Research and Quality. https://hcup-
us.ahrq.gov/toolssoftware/comorbidityicd10/comorbidity_icd10.jsp.

51. Elixhauser, A., Steiner, C., Harris, D.R., and Coffey, R.M. (1998). Comorbidity measures for
use with administrative data. Med Care *36*, 8–27. 10.1097/00005650-199801000-00004.

52. Charlson, M.E., Pompei, P., Ales, K.L., and MacKenzie, C.R. (1987). A new method of
classifying prognostic comorbidity in longitudinal studies: development and validation. J
Chronic Dis *40*, 373–383. 10.1016/0021-9681(87)90171-8.

53. Garry, E.M., Weckstein, A.R., Quinto, K., Bradley, M.C., Lasky, T., Chakravarty, A., Leonard,
S., Vititoe, S.E., Easthausen, I.J., Rassen, J.A., et al. (2022). Categorization of COVID-19

severity to determine mortality risk. Pharmacoepidemiol Drug Saf *31*, 721–728.

10.1002/pds.5436.

54. Shah, N., Nute, M.G., Warnow, T., and Pop, M. (2019). Misunderstood parameter of NCBI

BLAST impacts the correctness of bioinformatics workflows. Bioinformatics *35*, 1613–1614.

10.1093/bioinformatics/bty833.

55. Kaplinsky, J., and Arnaout, R. (2016). Robust estimates of overall immune-repertoire

diversity from high-throughput measurements on samples. Nat Commun *7*, 11881.

10.1038/ncomms11881.

1110