Post-ejaculatory Inhibition of Female Sexual Drive via Heterogeneous

Neuronal Ensembles in the Medial Preoptic Area

Kentaro K Ishii[1,2]**,** Koichi Hashikawa[1,2], Jane Chea[1,2], Shihan Yin[1,2], Rebecca Erin Fox[1,2], Suyang Kan[1,2], Meha Shah[1,2], Charles Zhou[1,2],

Jovana Navarrete[1,4], Alexandria D. Murry[1,4], Eric R Szelenyi[1,4], Sam A. Golden[1,4], Garret D. Stuber[1,2,3]

1. Center for the Neurobiology of Addiction, Pain, and Emotion, University of Washington, Seattle, WA, 98195, USA
2. Department of Anesthesiology and Pain Medicine, University of Washington, Seattle, WA, 98195, USA
3. Department of Pharmacology, University of Washington, Seattle, WA, 98195, USA
4. Department of Biological Structure, University of Washington, Seattle, WA, 98195, USA

Running title: Neural substrate for negative control of female sexual motivation

Highlights:

• **Female mice show decreased sexual motivation in the post-ejaculatory period.**
• **A subset of MPOA neurons in female respond specifically to male ejaculation.**
• **Male-ejaculation evokes persistent activity in MPOA inhibitory neurons in females.**
• **Activation of a subset of MPOA neurons is sufficient to suppress female sexual motivation.**

SUMMARY

Male ejaculation acutely suppresses sexual motivation in male mice. In contrast, relatively little is known about how male ejaculation
affects sexual motivation and sexual behavior in female mice. How the brain responds to completion of mating is also unclear. Here,
by using self-paced mating assay, we first demonstrate that female mice show decreased sexual motivation acutely after experiencing
male ejaculation. By using brain-wide analysis of activity-dependent labeling, we next pinpointed the medial preoptic area as a brain
region strongly activated during the post-ejaculatory period. Furthermore, using freely moving *in vivo* calcium imaging to compare
neural activity of inhibitory and excitatory neurons in the medial preoptic area, we revealed that a subset of the neurons in this region
respond significantly and specifically to male ejaculation but not to female-to-male sniffing or to male mounting. While there were
excitatory and inhibitory neurons that showed increased response to male ejaculation, the response magnitude as well as the proportion
of neurons responding to the event was significantly larger in the inhibitory neuron population. Next, by unbiased classification of their
responses, we also found a subpopulation of neurons that increase their activity late after the onset of male ejaculation. These neurons
were all inhibitory indicating that male ejaculation induces a prolonged inhibitory activity in the medial preoptic area. Lastly, we found
that chemogenetic activation of medial preoptic area neurons that were active during post-ejaculatory period, but not during appetitive

or consummatory periods, were sufficient to suppress female sexual motivation. Together, our data illuminate the importance of medial
preoptic area as a brain node which encodes a negative signal that sustains low sexual motivation state after the female mice
experience ejaculation.

11/29/2024
Kentaro Ishii

Introduction:

Sexual behavior is a fundamental and universal component of the observable behavioral spectrum in all mammalian species. This
behavior results in reproduction, the vital process for species propagation. However, reproduction carries significant burdens,
particularly for females, spanning pregnancy, birth, nursing, offspring rearing, and heightened vulnerability to environmental threats.
Consequently, the female ability to strictly control sexual drive is adaptive and confers individual survivability. This can involve both
increasing and decreasing motivation to engage in sexual behavior, dependent on anticipated beneficial outcome. Rodents provide a
powerful resource for studying female sexual behavior and the underlying neural basis (Agrati, 2022; Micevych and Meisel, 2017).
Typically, rodent sexual behavior is divided into three periods: an appetitive, consummatory, and post-ejaculatory phase (Hashikawa
et al., 2016; Ishii and Touhara, 2018; Lenschow and Lima, 2020). Importantly, in males, sexual motivation acutely reduces after
ejaculation (Hull et al., 2005). However, less is known about how the experience of male ejaculation in female mice impacts their sexual
behavior and motivation. Furthermore, how the transient experience of male ejaculation is represented in the brain to change the
behavioral outcome is not well understood. The medial preoptic area (MPOA), a molecularly and functionally heterogenous brain region,
has classically been known to regulate wide variety of social behaviors (Fang et al., 2018; Hashikawa et al., 2021; Karigo et al., 2021;
Kohl et al., 2018; Moffitt et al., 2018; Tsuneoka and Funato, 2021; Wei et al., 2018). Indeed, the MPOA has also been described to be
an important brain area for the regulation of female sexual behavior. Early studies using histological approaches to map neural activity
in rats has also showed MPOA, together with other hypothalamic regions, to be activated during sexual behavior, after male ejaculation
and after genital stimulations (Parada et al., 2010; Pfaus et al., 1996). Studies in female rats using electrophysiological recording or
histological approaches have reported that the MPOA contains neurons with various response specificity to sexual appetitive behaviors
such as sniffing, consummatory behaviors such as male mounting, intromission, and ejaculation (Kato and Sakuma, 2000; Pfaus et al.,
1993). Similarly, MPOA lesions increased sexual receptivity, suggesting that the MPOA negatively regulates female sexual behavior
when intact (Powers and Valenstein, 1972). Interestingly, the MPOA has also been found to positively regulate female sexual appetitive
behavior through their projection to the mesolimbic reward system (McHenry et al., 2017). However, how the MPOA both positively
and negatively regulate female sexual behavior, furthermore, sexual motivation, has been a long-lasting question in the field. Here,
we first assess sexual motivation in female mice by quantifying measurable approach or avoidance of a male conspecific. Next, by
combining cellular-resolution whole-brain mapping of activity-dependent labeling, *in vivo* calcium imaging, and chemogenetic neural
manipulation, we find that a subset of neurons in the MPOA, which are largely inhibitory, respond to male ejaculation but not to sexual
appetitive behavior. This population is sufficient to suppress female sexual motivation. Our findings illuminate the importance of the
MPOA as a brain region that encodes a negative signal that sustains low sexual motivation following male ejaculation in female mice.

Results:

Female sexual motivation is suppressed after male ejaculation.

Experimentally, sexual motivation can be quantified by the frequency of a subject displaying appetitive behavior, such as approach
and sniffing behavior, toward an opposite sex conspecific. However, in a traditional rodent mating assay, where the female and male
both freely interact with each other, it is difficult to isolate female-initiated appetitive behavior, and thus their sexual motivation. To
directly address aspects of female sexual motivation independent from the male's behavior, we used a female self-paced mating assay
(Erskine, 1989, 1985; Peirce and Nuttall, 1961). In this procedure, a female subject and a sexually experienced male are placed in a

behavior apparatus divided by a wall with a hole small enough for a female to move through, but not the male. The female subject may
move into the area where the male animal was placed ("Interaction zone") or move out to the other area ("Isolation zone") (**Figure 1A**).
Thus, in this assay the female has direct control of when interaction happens and provides a more ethologically valid measure of female
appetitive sexual behavior. Specifically, the amount of time the female subject spends in the interaction zone and the number of
transitions they make serve as metrics to quantify female appetitive behavior. On the experiment day, the female subject, which was
primed with estradiol to be behaviorally estrus, was placed in the apparatus with the partner ("Sexual interaction" trial) or alone ("Control"
trial). Each animal went through both trials on different experiment days. The position of the animal was tracked to quantify the time
spent in each zone. First, we found that the virgin female mice stayed in the isolation zone significantly longer than sexually experienced
females, suggesting that previous sexual experience increased sexual motivation in the female mice (**Figure S1**). Next, in the sexually
experienced animals, we observed robust behavioral changes after male ejaculation. First, we observed that female subjects tended
to spend more time in the isolation zone after male ejaculation compared to post- sniffing, mounting and intromission (**Figure 1B–F**).
To quantify the changes in animal behavior, we divided the sexual interaction trial into pre- and post- male ejaculation epochs (**Figure**
1G–M). As result, we found that there was a significant decrease in the number of zone transitions as well as a significant increase in
the time spent in the isolation zone after male ejaculation. Importantly, this change was not dependent on the state of the male animal
(**Figure S2**). When the female animal was introduced to a novel sexually motivated male animal immediately after they experienced
male ejaculation from another mouse, they showed similar reduction in zone transition and increase in the time spent in the isolation
zone. To further evaluate the female animal's motivation to engage in sexual behavior, we quantified the latency to return to the
interaction zone after a behavioral episode (Bermant, 1961; Bermant and Westbrook, 1966). This was the largest after the female
experienced male ejaculation when compared to after sniffing or after mounting (**Figure 1I**). In addition, while there was no change in
the amount of female-to-male sniffing, there was a significant decrease in the number of mounting and intromission episodes (**Figure**
1J–L). There was also a significant increase in the number of self-grooming behavioral episodes (**Figure 1M**). Animals that were in
their natural estrus cycle showed similar behavior changes after experiencing male ejaculation (**Figure 1N–T).** Interestingly, animals
that were in their natural estrus cycle showed strong reduction of sniffing after experiencing male ejaculation but no changes in self-
grooming behaviors (**Figure 1Q, 1T**). Collectively, these results suggest that male ejaculation acutely decreases sexual motivation.
92

Brain-wide mapping of neural response to male ejaculation using activity-dependent labeling in the female brain.
We next questioned how male ejaculation suppresses female sexual motivation. To investigate this question, we first determined which
brain regions were associated with neural activity during the post-ejaculatory period in an unbiased manner. We conducted a cellular-
resolution brain wide activity mapping analysis using targeted recombination in active populations method (TRAP), which allows genetic
labeling and access to neurons that are active during a specific time window determined by tamoxifen administration (DeNardo et al.,
2019). To capture the activity pattern, Fos-2A-iCreER^{T2}::RCL-tdTomato (TRAP2::Ai14) female mice were administered 4-
hydroxytamoxifen (4-OHT) immediately after male ejaculation ("post-ejaculatory" group) (**Figure 2A**). In this method, cells that show
behaviorally induced Fos activity were labeled with tdTomato. In addition to these animals, we also prepared a group of animals that
were administrated with 4-OHT while the animals were in the appetitive phase or in the consummatory phase. For the appetitive group,
female subjects interacted with male mice and displayed female-to-male sniffing but did not experience any mount or intromission
behavior, nor male ejaculation. For the consummatory group, female subjects interacted with male mice and displayed female-to-male

sniffing, experienced mount or intromission behavior, but not male ejaculation. After 2 weeks, the tissue was collected, cleared, imaged,
and processed for registration on a template brain atlas (Park et al., 2019; Renier et al., 2016) (**Figure 2A**). The tdTomato+ cells were
segmented using a trained classifier and normalized by the area of each brain region (**Figure 2B**). As result, we found the level of
tdTomato expression increased in a brain wide manner during the post-ejaculatory period when compared to the appetitive phase
(**Figure 2C–D**, **Table 2**). We further noticed that regions well known to be related to social behavior had significantly higher tdTomato
density in the post-ejaculatory group compared to appetitive group (**Figure 2E–I, Figure S3, S4, Table 3**). This includes paraventricular
hypothalamic areas (PVHd, PVH, PaV) which are important for endocrinological regulations, medial amygdala (MeA) which is critical
for olfactory sensory information processing, and the BNST (Flanigan and Kash, 2022; Marsh et al., 2021; Raam and Hong, 2021).
The medial preoptic area nucleus (MPN), a subregion of the MPOA, had significantly higher tdTomato density in the post-ejaculatory
group compared to the appetitive group. The MPN also had the 4th largest difference of density between the post-ejaculatory group
and appetitive group among all the brain regions (**Figure 2F**). While it was not statistically significant, the consummatory group also
tended to have a larger tdTomato density than the appetitive group in the MPN as well (**Figure 2G and 2H**). From this result, we
decided to further investigate the role of MPOA and how it is related to the regulation of female sexual motivation after experiencing
male ejaculation.
118
Male ejaculation is represented by both inhibitory and excitatory cell types in the MPOA.
MPOA is a molecularly heterogenous brain region. The area is a mixture of 80% inhibitory neurons and 20% of excitatory neurons
which can be further classified into neuronal subtypes by marker gene expressions (Hashikawa et al., 2021; Moffitt et al., 2018). These
subtypes are often associated with a biological function. One example are the *Nts* expressing cells, which also tend to co-express *Esr1*
and *Vgat* (McHenry et al., 2017). In female mice, these neurons were shown to respond to male odor when the subject was sexually
receptive, and the activation of these neurons were sufficient to increase sexual appetitive behaviors. Therefore, identifying the specific
cell types which respond to male ejaculation but not to appetitive behavior is critical to understand how the MPOA complex regulates
female sexual motivation. Here, we utilized multiplexed *in situ* RNA hybridization chain reaction (HCR) to identify previously
characterized cell type markers of MPOA and an immediate early gene *Fos* (**Figure S5A**) (Choi et al., 2014). Brain samples were
collected after the subject experienced male ejaculation ("post-ejaculatory" group) or during the appetitive phase, while the subject was
interacting with the male mice without any consummatory behavior ("Appetitive" group) (**Figure S5B**). First, to understand which
subregion of the preoptic area was activated, we quantified the number of *Fos*+ cells. As result, we found that the number of *Fos*+ cells
in the post-ejaculatory group were significantly higher than appetitive group in the MPN but not in the nearby regions (**Figure S5C–D**).
In the MPN, the amount of *Vglut2*+ and *Vgat*+ cells expressing *Fos* were significantly higher in the post-ejaculatory group than in the
appetitive group, suggesting that male ejaculation activates both excitatory and inhibitory neurons (**Figure S5E–I**). The proportion of
Fos+ cells was higher in *Vgat*+ population. To further investigate the cell types in the MPN that were activated by male ejaculation, we
analyzed the co-expression of *Fos* and the MPOA cell type markers (**Figure S5J**). As result, in the *Vglut2*+ expressing cells, we found
that *Fos* induction was only observed in the cells that co-expressed *Calcr*. In contrast, in the *Vgat*+ expressing cells, *Fos* induction was
observed in *Calcr, Gal* and *Prlr* co-expressing cells. Next, we conducted clustering on the gene expression matrix to classify cells and
analyzed the distribution of *Fos* among the clusters (**Figure S5K–M**). As result, we were able to identify 6 potential cell types which we
labeled as type 1: *Vglut2*+, type 2: *Vgat*+, type 3: *Gal*+, type 4: *Nts*+, type 5: *Calcr*+ and type 6: NA, or a cell group that could not be

categorized by a specific gene expression pattern. Type 1: *Vglut2+,* type 2: *Vgat+* and type 3: *Gal+* had significantly more *Fos+* cells
from the post-ejaculatory group compared to the appetitive group (**Figure S5M**). In contrast, *Fos+* cells were not induced in type 4:
Nts+ indicating that MPOA neurons which regulate appetitive sexual behavior was not activated by male ejaculation. Collectively, these
results indicate that male ejaculation induces more neural activity in the MPN when compared to appetitive behavior, both in excitatory
and inhibitory cell types.
145
A subpopulation of MPOA Vgat cells respond to male ejaculation in the female brain.
While mapping neural activity based on *Fos* activity provided critical insight in how the brain responds to male ejaculation, the low
temporal resolution of the method makes it difficult to identify cells that specifically respond to the onset to each behavior. For this
reason, we next conducted single-cell resolution *in vivo* imaging in a freely moving animal (**Figure 3A–B**). We virally labeled MPOA
inhibitory and excitatory cells with a calcium indicator GCaMP6s by using a Cre-dependent virus and female Vgat-Cre or Vglut2-Cre
respectively **(Figure 3C, Figure S6)** (Vgat-Cre: n = 6, 87 ± 27 cells. Vglut2-Cre: n = 4, 68 ± 42 cells). We recorded the calcium activity
of MPOA Vgat and Vglut2 cells in female mice during mating in a home cage mating assay (**Figure 3E**). First, we analyzed the
magnitude of calcium activity for 15-seconds after behavior onset. From peri-event analysis, we found that as a population, Vgat cells
showed significant changes in calcium activity after the onset of all behaviors observed, with the highest increase after male ejaculation
(**Figure 3F–G**). Vglut2 cells showed increased activity after sniffing and mounting, intromission, and male ejaculation. Furthermore,
when we compared the response of Vgat and Vglut2 cells, Vgat cells showed significantly higher average response magnitude after
sniffing, self-grooming, anogenital sniffing and male ejaculation than the Vglut2 cells (**Figure 3H**). Vgat cells also contained a
significantly higher proportion of cells that positively respond (> 2δ) to sniff and male ejaculation onset (**Figure 3I**). These results
suggest that male ejaculation signal is more dominantly represented in the MPOA inhibitory cells than in the excitatory cells. On the
other hand, the response magnitude and the proportion of cells that respond after mounting and intromission had no significant
difference between the two cell types. This indicates that the excitatory/inhibitory balance in the MPOA fluctuates during mating. The
MPOA has high inhibitory activity during appetitive phase, with more Vgat cells responding to sniffing, similar activity during the
consummatory phase, and higher inhibitory activity after male ejaculation.
To further investigate the heterogeneity of MPOA during mating, we subset the cells into those which positively responded to any of
sniff, mount or male ejaculation, and then classified the cells based on their specificity (**Figure 4A,B** and **Figure S7**). Consistent with
Figure 3I, the number of cells that specifically responded was largest to male ejaculation with 177 cells, next to mount with 46 cells
and to sniff with 12 cells. When compared to other cell populations, sniff cells and completion cells only showed significantly stronger
responses than the other cells to sniff and male ejaculation respectively (**Figure 4C**). Mount responsive cells showed significantly
stronger response than the other cells to mount, intromission and anogenital sniffing, but not to sniffing and male ejaculation (**Figure
4C**). We also found that the cells that respond to these 3 behaviors partially overlap but were mostly distinct (**Figure 4D**). The proportion
of cells that respond specifically to a behavior was highest for male ejaculation (Vgat: 80%, Vglut2: 82%), and lower in mounting (Vgat:
55%, Vglut2: 68%) and sniffing (Vgat: 36%, Vglut2: 0%). There was no notable bias in the anatomical distribution of these cells within

the field of view (**Figure 4E**). Overall, these results suggest that the MPOA contains highly heterogenous functional population which
contains a large proportion of cells specifically responding to male ejaculation in female mice.
175
MPOA Vgat cells display prolonged activity after the onset of male ejaculation.
Female mice avoid sexual interaction with a male animal after male ejaculation for hours and even days, suggesting a persistent
changes in the animal's behavior (McGill and Coughlin, 1970; Zhou et al., 2023). How the MPOA regulate this persistent change in
female sexual motivation is unclear. To analyze the response of MPOA cells on a longer time scale, we focused on the cells that
positively responded to male ejaculation and characterized the calcium response pattern for 120-seconds after the onset (**Figure 5**).
First, we observed that Vgat cells showed more diverse response patterns during this time-window (**Figure 5A,B**). To quantify this
difference, we calculated the decay length of calcium signal, the time for the increased calcium signal to return to baseline (**Figure 5C**).
As result, we found that Vgat cells had significantly longer decay length compared to the Vglut2 cells, suggesting increased activity
late after the onset of male ejaculation. To further classify the temporal dynamics of responses to male ejaculation, we first
concatenated calcium activity that occurred within this time-window from both Vgat and Vglut2 cells and conducted spectral clustering
in an unbiased fashion. As a result, we were able to classify the responses into 5 types that could be characterized as: 1) fast response,
2) slow response, 3) late response, 4) weak response and 5) negative response (**Figure 5E, Figure S8**). Type 3 contained cells that
show increased calcium dynamics late after the onset of male ejaculation and had significantly larger decay length than the other cell
types (**Figure 5F**). Then we separated the Vgat and Vglut2 cells to compare their proportions within each cluster. Interestingly, all the
cells in type 3 were Vgat cells, suggesting that the increased activity in the MPOA late after the onset of male ejaculation is mediated
by inhibitory neurons (**Figure 5H–K**). In contrast, Vglut2 cells were more likely to show fast responses than Vgat cells (**Figure 5K**).
Collectively, these results suggest that the MPOA contains a subpopulation of inhibitory neurons that increase their activity late after
the onset of male ejaculation, which may contribute to persistent suppression of the female animals' sexual motivation.
194
An activity defined subset of neurons in the MPOA is sufficient to suppress female sexual behavior.
Next, we examined whether the increased activity of neurons in the MPOA is sufficient to suppress female sexual motivation after male
ejaculation. We injected an AAV to express the chemogenetic neural activator hM3Dq or eYFP as control into the MPOA of TRAP2
female mice (**Figure 6A**). To capture the neural activity, each mouse was administrated 4-OHT immediately after the after male
ejaculation ("post-ejaculatory-hM3Dq" or "post-ejaculatory-eYFP" group). In addition to these animals, we also prepared a group of
animals in which we injected an AAV expressing hM3Dq and then administrated 4-OHT during the appetitive phase ("appetitive-hM3Dq"
group) or the consummatory phase ("consummatory-hM3Dq" group) (**Figure 6C**). We found that while the overall number of TRAP-
labeled cells had no difference between the post-ejaculatory groups, consummatory group and appetitive group, the proportion of cells
located in the MPN was significantly higher in the post-ejaculatory group (**Figure 6D** and **Figure S9A–C**). From *in situ* RNA
hybridization, we found that most TRAP-labeled cells in the MPN were *Vgat*+ (*Vgat*+: 0.76 ± 0.16. *Vglut2*+: 0.15 ± 0.13.). The proportion
of *Vgat*+ cells was significantly higher than the *Vglut2*+ cells (**Figure S7D–G**). Together these results suggest that the post-ejaculatory
group labels more inhibitory neurons in the MPN than in appetitive group. Furthermore, we analyzed the accuracy and efficiency of
TRAP labeling by allowing the female subject to experience male ejaculation before perfusion (**Figure S9H–L**). By immunolabeling c-
Fos protein, we analyzed how well the TRAP ensemble overlaps with the c-Fos positive-ensemble in the MPN. As a result, we first

found that the proportion of c-Fos+ and TRAP-labeled cells in the post-ejaculatory group tended to be higher than the appetitive group
and the consummatory group (**Figure S9I**). The proportion of c-Fos+ cells were consistent across each group (**Figure S9J**). Next, we
found that the accuracy of TRAP labeling (the proportion of TRAP-labeled cells that were also c-Fos positive) was similar between the
groups (**Figure S9K**). The efficiency of TRAP labeling (the proportion of c-Fos+ cells that were also TRAP labeled) tended to be higher
in the post-ejaculatory group (**Figure S9L**) than in the appetitive group and the consummatory group. This also suggests that the
appetitive group labels a different set of neurons than the post-ejaculatory group. These results indicate that by using the TRAP2
method, we were able to specifically label a proportion of MPOA neurons, which are primarily inhibitory, that contains neurons that
respond to male ejaculation.
To test the causality of these neurons, we first investigated how the activation affects female sexual behavior in a home cage mating
assay (**Figure S10A–F**). The subjects were intraperitoneal injection (i.p.) injected with CNO 30-minutes before the assay. As result,
first we found that post-ejaculatory-hM3Dq group but not the appetitive-hM3Dq and consummatory-hM3Dq group showed significantly
fewer sniffing events compared to post-ejaculatory-eYFP group (**Figure S10C**). For consummatory behaviors, while post-ejaculatory-
hM3Dq group showed no difference in mounting episodes compared to post-ejaculatory-eYFP group, there were significantly less
intromission episodes, resulting in lower receptive scores (**Figure S10C–F**). Thus, from these results, activation of MPOA neurons that
respond to male ejaculation suppresses both appetitive and consummatory sexual behavior in female mice. As a next step, we
investigated how the activation of these MPOA neurons affects female sexual motivation in the self-paced mating assay (**Figure 6**).
The subjects were i.p. injected with CNO or saline 30-minutes before the assay (**Figure 6B**). As result, only post-ejaculatory-hM3Dq
group showed significantly fewer number of transitions and spent more time in the isolation zone on the CNO trial than on the saline
trial (**Figure 6E–G**). We also found that post-ejaculatory-hM3Dq group displayed fewer numbers of self-grooming, mounting,
intromission and lower receptivity score on the CNO trial than on the saline trial (**Figure 6H–L**). Interestingly, activation of these neurons
did not induce conditional place preference or aversion (**Figure S10G–J**), implying that the suppression of sexual behavior is not
associated with negative valence. Next, we investigated how inhibition of these neurons affects female sexual behavior (**Figure S11**).
We prepared a new group of mice in which we injected an AAV expressing hM4Di or eYFP as control in TRAP2 female mice. Two-
weeks after injection, we administrated 4-OHT immediately after the subject experienced male ejaculation ("post-ejaculatory-hM4Di"
or "post-ejaculatory-eYFP" group). First, we tested the animals in the self-paced mating assay (**Figure S11A–L**). The female subjects
were i.p. injected with CNO or saline 30-minutes before the assay (**Figure S11B**). As result, the subjects did not show any significant
change in sexual behavior during this assay (**Figure S11D–L**). Next, we tested the animals in the home cage mating assay (**Figure
S11M–S**). The subjects were intraperitoneal injections (i.p.) injected with CNO 30-minutes before the assay (**Figure S11M**). As result,
the post-ejaculatory-hM4Di group experienced did not show significant changes in sniffing, self-grooming or mounting, but showed
more intromission and higher sexual receptivity than in the post-ejaculatory-eYFP group (**Figure S11O–S**). Collectively, these results
suggests that the MPOA neurons are sufficient but not necessary for the modulation for sexual motivation but is sufficient and necessary
for consummatory behaviors in female mice (**Figure 6M**).

Discussion:
How sexual motivation is represented in the brain has been a long-lasting question in the field. Here we found that MPOA plays a
critical role in regulating female sexual motivation. We found that MPOA neurons, especially the inhibitory neurons, respond significantly

and specifically to male ejaculation. These neurons show persistent activity after the onset of male ejaculation. Reactivation of these
neurons result in suppression of female sexual motivation. We propose that MPOA together with other neural substrates such as the
BNST (Zhou et al., 2023), encode a negative feedback signal that sustains low sexual motivation state after male ejaculation in female
mice. Here, we will further discuss how sensory inputs contribute to the male ejaculation response in the MPOA, how MPOA responds
to different sexual behaviors and how these neurons work in concert with other sexual behavior related hypothalamic regions.
249

Using self-paced mating paradigm to evaluate sexual motivation in female rodents
The self-paced mating paradigm is a classical yet powerful method to directly assess female sexual behaviors (Erskine, 1989, 1985;
Peirce and Nuttall, 1961). This paradigm is still in current use, mainly in rats (Ventura-Aquino and Paredes, 2023). Our paper
demonstrate that this simple behavioral paradigm is still a powerful method to study neural circuitry of female sexual behavior and
motivation in mice. In our results we first demonstrate that the latency to return to the interaction zone was significantly larger after the
animal experienced male ejaculation consistent with the results shown in rats. Direct measurements of both increase of withdrawal
and decrease of approach to sexual behavior are the strength of this paradigm. Furthermore, we have also demonstrated that female
animals show more time spent in the isolation zone regardless of the male partners sexual motivation state (Figure S2). From these
results, we believe that this paradigm reports important metrics that correlate with female sexual motivation. However, it is also clear
that other factors affect the females behavior in the self-paced mating paradigm. Recent studies developed an operant conditioning
apparatus which delivers an intruder to a male subject in response to a nose poke (Minakuchi et al., 2024). This apparatus is a powerful
approach to dissociate motivation and action which is difficult in commonly used self-paced paradigms. The self-paced mating paradigm
used in our study has limitations to studying sexual motivation in detail. An advancement in the design of the apparatus may be
essential to further study the neural circuitry that regulates female sexual motivation in fine detail.
264

Sensory representation of male ejaculation in the female brain and genitalia.
Suppression of female sexual behavior and motivation after male ejaculation is not a unique phenomenon found in mice and can be
observed in various insect species (Gillott, 2003). Studies in *Drosophila* has demonstrated that the primary trigger of this behavioral
switch is the sex peptide (SP), a peptide released in the male seminal fluid (Chapman et al., 2003; Chen et al., 1988; Liu and Kubli,
2003). Studies in flies have further identified the receptor for SP as well as the neural circuitry which conveys the sensory information
to the mating regulating neurons in the brain (Feng et al., 2014; Rezával et al., 2012; Yapici et al., 2008). In contrast to the well-studied
system in *Drosophila*, little is known about how the mammalian brain senses male ejaculation. Histological studies in rats have shown
that male ejaculation induces c-Fos expression widely in the brain, including the MPOA (Pfaus et al., 1993). Indeed, anesthesia to the
female genitalia has been shown to disrupt pacing of mating, indicating that sensory input from the genitalia can impact female sexual
behavior (Parada et al., 2014). Furthermore, artificial stimulation of the vagina or the clitoris of the female rat was further shown to
induce c-Fos expression in the MPOA in a similar fashion as male ejaculation, indicating that at least tactile sensory input can be
conveyed to this brain area (Parada et al., 2010; Pfaus et al., 1996). Ejaculation from vasectomized male animals were sufficient to
suppress sexual behavior suggesting that at least chemical components from the testis is not necessary for this effect (Zhou et al.,
2023). A recent study has shown that mechanical stretch induces serotonergic activity in the dorsal raphe, which resembles changes
after male ejaculation, suggesting that stretch sensation is mediating male ejaculation signal to the brain (Troconis et al., 2023). Mouse

genetics has also enabled identification of nerves innervating the genitalia (Qi et al., 2024). However, there are still to uncover on how
stretch sensation in the genitalia is conveyed to the brain, and equally unclear how this information is ultimately transmitted to the
MPOA. Our findings on how the male ejaculation signal is represented in the MPOA will be an ideal entry point to study the neural
basis of genital-brain communication.
284
Appetitive behavior and male ejaculation ensembles in MPOA.
The MPOA is a molecularly and functionally heterogenous brain region. One function MPOA has heavily been related to is parenting
behavior. MPOA neurons which express *Gal* and/or *Calcr* has been demonstrated to regulate parenting behavior (Kohl et al., 2018;
Wu et al., 2014; Yoshihara et al., 2021). From our data, neurons that respond to male ejaculation tended to co-express *Gal* and *Calcr*
more than neurons that responded to appetitive behavior (**Figure S3**). However, we also found that chemogenetic activation of male
ejaculation responsive MPOA neurons did not induce parenting behavior (**Figure S10K**). While it is necessary to conduct extensive
analysis of how MPOA neurons respond to other behaviors, these results suggest that male ejaculation responding MPOA neurons
may specifically regulate sexual behavior and motivation but not parenting behavior. A previous study also found that MPOA contains
a subset of neurons which regulate female appetitive sexual behavior. MPOA neurons which express *Vgat, Esr1* and *Nts*, and projects
to the ventral tegmental area (VTA) has been shown to respond to olfactory signals from the male mouse and to enhance female-to-
male sniffing (McHenry et al., 2017). There is also a significant release of dopamine after the first female to male sniffing interaction as
well as after male ejaculation (Dai et al., 2022). How the brain, especially the MPOA, increases sexual motivation before and during
the appetitive phase, and how it suppresses motivation after male ejaculation, has been a long-lasting question in the field that is
addressed throughout this work. The whole brain activity-dependent labeling experiment identified the MPOA as one of the brain
regions which strongly increases Fos activity (**Figure 2**).
However, there were many other brain regions showing increased activity associated with male ejaculation, including ones that are not
canonically involved in the regulation of social behavior. While it is possible that all these brain regions work in concert to suppress
sexual motivation after male ejaculation, our results demonstrate that a specific brain region, MPOA, is sufficient to evoke behavioral
response. This suggests that all brain regions showing higher Fos activity are not necessarily causal to behavioral changes. It is also
possible that these widespread brain regions may be a consequence of noise due to the long time-window in this method, which may
be resolved by further development of genetic methods for neural ensemble labeling. In the MPOA, our *in vivo* calcium imaging data
shows nearly half of the inhibitory neurons that positively respond to appetitive behavior also respond to male ejaculation (**Figure 4**).
Our data also identified that there was no difference between the number of *Nts+* cells activated by appetitive behavior and male
ejaculation (**Figure S5**). These results suggest that there is a small but significant number of neurons which respond both to appetitive
behavior and male ejaculation in the MPOA, and potentially regulate release of dopamine.
Although a large number of neurons that responded to female-to-male sniffing also respond to male ejaculation, the opposite
is not true (**Figure 4**). Importantly, the chemogenetic activation of neurons that were active after male ejaculation suppressed the
amount of sexual interaction in the self-paced mating assay, opposite to what was described when activating *Nts+* MPOA neurons

(McHenry et al., 2017). This suggests that there are at least two-distinct populations in the MPOA which regulate appetitive sexual
behavior or sexual motivation in opposing directions.
315

Neural circuitry for female sexual behavior regulation:
The neural circuitry that regulates female sexual behavior has been extensively studied in rodent models. From early lesion and
electrical stimulation studies, it has been proposed that the rodent brain has a positive and negative regulatory system for female
sexual behavior. The positive system has been proposed in rats, where Pfaff and Sakuma demonstrated that the lesion of the
ventromedial hypothalamus (VMH) abolished sexual receptivity while electrical stimulation of the same region induces receptivity (Pfaff
and Sakuma, 1979a, 1979b). Mouse genetics further showed that a subset of neurons in the ventrolateral part of the VMH is crucial
for the regulation of female sexual behavior (Inoue et al., 2019; Yang et al., 2013). More specifically, *Esr1+ Cckar+* (*Nts*-) cells in the
VMHvl were found to be necessary and sufficient for female sexual receptivity (Liu et al., 2022; Yin et al., 2022). These results suggest
that the VMHvl is the "command center" which positively regulate female sexual behavior.
The MPOA together with other brain regions has been proposed to form the negative regulating system. Lesion of the MPOA was
shown to increase the sexual receptivity of the female rat and electrical stimulation was shown to reduce receptivity, suggesting that
MPOA negatively regulates female sexual behavior (Pfaff and Sakuma, 1979b; Powers and Valenstein, 1972). In the current study, we
demonstrate that chemogenetic activation of neurons in the MPOA which respond to male ejaculation is sufficient to suppress female
sexual behavior and sexual motivation. This result suggests that these neurons in the MPOA constitute the negative regulating system.
Importantly, both our single-cell calcium imaging results and histological analysis of the chemogenetic activation cohort suggest that
the majority of these neurons are inhibitory (**Figure 3I** and **Figure S9G**). We further demonstrate that the inhibitory population contains
neurons that show prolonged responses to male ejaculation which resemble the persistent change of sexual motivation. Other studies
have also supported the idea that the inhibitory cell population is more related to the regulation of social behavior than the excitatory
population (Fang et al., 2018; Hashikawa et al., 2021; Moffitt et al., 2018; Tsuneoka and Funato, 2021; G.-W. Zhang et al., 2021). Our
results are also in line with this idea and suggest that the inhibitory neuronal population is the main regulator of the change in sexual
motivation. The role of excitatory neurons that respond to male ejaculation will be a future topic to examine.
Recently, BNST was also found to play a role in negatively controlling female sexual behavior (Zhou et al., 2023). There, they use a
behavioral assay which introduces 2 different males to the female subject before and after male ejaculation. They demonstrate that
after male ejaculation, the female will be less receptive toward the second male. Importantly, this assay clearly demonstrate that the
reduction of female sexual behavior is due to the female herself, not due to the change in their male conspecific. This result together
with the result from our self-paced mating assay (**Figure 1**) strongly indicates that the behavioral change after male ejaculation reflects
change in internal motivational states. The study further used single-cell resolution *in vivo* calcium imaging techniques to demonstrate
that BNST^{Esr2} neurons show specific and robust responses after male ejaculation, like what we observed in the MPOA. They then use
opto- and chemogenetic tools to study the causality of these neurons. They demonstrate that the inhibition of BNST^{Esr2} neurons rescued
female receptivity after male ejaculation, activation of these neurons did not affect sexual behavior in female mice, suggesting that
BNST^{Esr2} neurons are necessary but not sufficient to suppress female sexual behavior and sexual motivation. While there were some
differences in how modulation of BNST and MPOA neurons affect sexual behavior and motivation, overall, the role of BNST and MPOA
seems to be similar suggesting that the two regions work in concert to suppress female sexual behavior. This is contrary to the finding

from Mei et al. showing that the BNST^{Esr1} neurons and MPOA^{Esr1} neurons forms a bidirectional inhibitory circuit to antagonistically
regulate infanticidal behavior and maternal behavior respectively (Mei et al., 2023). A mechanism which is yet to be revealed can
override the negative feedback loop between MPOA and BNST neurons, potentially a disinhibitory circuit that connects the BNST^{Esr2}
and MPOA^{GABA}, is necessary for these neurons to regulate the same behavior change.
Interestingly, a subpopulation in the VMHvl expressing Cckar show strong inhibition of spontaneous bulk calcium events after male
ejaculation (Yin et al., 2022). This is opposite to the persistent increase of activity in the MPOA and BNST, suggesting that the activity
in these regions is reflected onto the VMHvl. Indeed, MPOA and BNST both send dense projection to the VMHvl, and VMHvl^{Esr1} neurons
project to the MPOA and to the BNST, making the MPOA-BNST-VMHvl network highly interconnected (Dimén et al., 2021; Lo et al.,
2019; Osakada et al., 2018). A recent study focusing on male aggression motivation further demonstrated that the long-range input
from inhibitory MPOA neurons to VMHvl promote and prolongs low motivational state (Minakuchi et al., 2024). From this finding and
the fact that the acute activation of MPOA neurons more drastically affected female sexual behavior than the BNST^{Esr2} neurons, we
speculate that the MPOA can directly inhibit the VMHvl, while the BNST utilizes a different circuit mechanism which either involves
multiple synaptic connections or different molecular mechanisms to act on the VMHvl.
Long-lasting change in neural activity within this network, potentially through monoamine or steroid molecules such as dopamine or
opioids may be the neural basis of suppressed sexual motivation after male ejaculation (Micevych and Meisel, 2017; S. X. Zhang et
al., 2021). Another candidate is serotonin, which is known to be an important mediator for sexual behavior (Uphouse, 2014). Recent
study has revealed release of serotonin in response to male ejaculation in female mice (Troconis et al., 2023). Our whole brain activity
mapping data also indicates increased Fos activity in the dorsal raphe (**Figure S3 and S4**). Whether serotonin mediates the changes
in persistent activity across the entire network and whether there is a difference in how it impacts MPOA and BNST and its output to
the VMHvl will be an important topic of future studies.
Taken together, our study identifies a subset of neurons in the MPOA which is dedicated to negatively sustain female sexual motivation.
This finding fills the missing hole in how the brain neurons regulate female sexual motivation and proposes that the basal forebrain
complex of MPOA and BNST work in concert as the negative regulating system of female sexual behavior.
372
373

Materials and methods:

Animals.

All experiments were approved by the Institutional Animals Care and Use Committee at the University of Washington (Protocol #4450-
01). Animals were group-housed with littermates on a 12-hr light cycle at ~22°C with food and water available *ad libitum*. Wild-type
Swiss-Webster mice were purchased from Taconic Biosciences (NY, U.S.). Wild-type C57BL/6 mice were purchased from the Jackson
laboratory and bred in-house. Vgat-Cre (also known as *Slc32a1*-ires-Cre, Jax#016962), Vglut2-Cre (also known as *Slc17a6*-ires-Cre,
Jax#028863) and Ai14 (Rosa-CAG-LSL-tdTomato-WPRE, Jax#007914) were purchased from Jackson laboratory. TRAP2 mice (also
known as Fos-2A-iCreER^{T2}, (DeNardo et al., 2019)) were provided by Dr. Palmiter (University of Washington) and crossed with Ai14
to generate TRAP2::Ai14 mice. All male mice which were used as intruders were singly housed to avoid male-male aggression. Female
mice with GRIN-lens implant were singly housed. All other mice were group housed.

Surgery.

For targeting AAV into a certain brain region, stereotactic coordinates were defined for each brain region based on the brain atlas
(Franklin and Paxinos, 2007). Mice were anesthetized isoflurane and head-fixed to stereotactic equipment (Kopf). For *in vivo* calcium
imaging experiments, 500 nL of AAVDj EF1a-DIO-GCaMP6s (2.8*e12, UNC viral core, lot#av78310) was injected into two coordinates
in the target brain region at a speed of 60 nl/min using Nanoject 3 pump (Drummond Scientific). The following coordinates were used
[Distance in millimeters from the Bregma for the anterior (A)-posterior (P) and lateral (L) positions, and from the brain surface for ventral
(V) direction]: MPOA, A 0.2, L 0.15, V 5.2. and A 0.4, L 0.35, V 5.2. Immediately following viral injection, 5 screws were placed into the
skull and a GRIN lens (Inscopix, 0.6 mm x 7.3 mm) was slowly lowered to position above the injection site: MPOA, A 0.3, L 0.3, V 4.95.
GRIN lens was glued to the skull with C&B Metabond Kit (Parkell, # S380) and dental cement (Lang Dental, #1304CL). After placement
of GRIN lens, a custom-made head ring was glued to the skull and screws. For bare GRIN lenses, more than 3 weeks after the virus
injection and GRIN lens placement surgery, a baseplate (Inscopix) was positioned to while visualizing GCaMP6s expression with a
microscope. Once a field of view with GCaMP6s expression was found, the baseplate was glued to the head ring and skull. For
pharmacogenetic experiments, 300-400 nL of AAV5 hSyn-DIO-hM3D(Gq)-mCherry (2.0*e13, Addgene, #44361-AAV5, lot#v141469),
AAV5 hSyn-DIO-hM4D(Gi)-mCherry (2.5*e13, Addgene, #44362-AAV5, lot#v178820) or AAV5 EF1a-DIO-eYFP (4.4*e12, UNC viral
core, lot#av4802B) was bilaterally injected into the target brain region at a speed of 60 nl/min using Nanoject 3 pump. The following
coordinates were used: MPOA, A 0.3, L 0.25, V 5.2. To avoid pregnancy during the first experience, all female mice that experienced
a mating assay except ones used in **Figure 1 N–T** was ovariectomized. Ovariectomized female mice that were primed with hormones
has been successfully used in other sexual behavior studies (Ring, 1944; Yang et al., 2013).Animals that were used *in vivo* calcium
imaging experiments or pharmacogenetic experiments were ovariectomized during the virus injection or GRIN lens placement surgery.
After surgery, the incision was sutured, and the animal was warmed using a heat pad to facilitate recovery from anesthesia.

Behavior experiments and analysis.

For all mating assays, female mice older than 6 weeks were ovariectomized (OVX) under isoflurane anesthesia. If necessary, mice
underwent surgeries for AAV injections, GRIN lens implantation on the same day. After more than 2 weeks of recovery, the OVX female
subjects were subcutaneously (s.c.) injected with 0.1 mL and 0.05 mL Estradiol (E8515, Sigma-Aldrich, 0.1 mg/ml) dissolved in sesame

oil (Sigma-Aldrich) at 24 and 48h before assay, and 0.05 mL Progesterone (P0130, Sigma-Aldrich, 10 mg/ml in sesame oil) at 4h before

assay. If necessary, mice underwent I.P. injection of CNO/saline. C57BL/6 or Swiss-Webster mice were used as a stud male. A 4 – 6

cm to head bar was placed onto the male mice under isoflurane anesthesia, which was used for the female self-paced mating assay

described below. The male animals were kept isolated until the assay and had been trained prior to the assay to show mounting and

ejaculation.

The home cage mating assay was conducted as previously described with modifications (Ishii et al., 2017). Each assay was conducted

5-10 hrs after the onset of dark period, under an IR light and recorded (30 Hz) for subsequent 30 to 120 min for analysis of sexual

behavior from a dorsal view. For neural activation experiments using hM3Dq (**Figure S9**), 0.1 mL of 1 mg/kg clozapine-N-oxide

dissolved in saline (CNO, RTI International, NIMH Code C-929) or 0.1 mL of saline was I.P. injected into the animal 30 minutes prior

to the male entry. For neural inhibition experiments using hM4Di (**Figure S11**), 0.1 mL of 5 mg/kg CNO dissolved in saline or 0.1 mL

of saline was I.P. injected into the animal 30 minutes prior to the male entry.

The female self-paced mating assay was conducted using a behavior apparatus modified from an apparatus used in rat (Nyuyki et al.,

2011). A 1-cm thick black wall with a 20 mm x 20 mm hole 3D-printed with PFA filament was placed in a 25 cm x 35 cm cage to divide

the room in a 1:2 ratio along the longer edge. The smaller area was defined as the isolation zone and the larger area was defined as

the interaction zone. The female subjects that were not able to pass through the hole during the habituation was eliminated from the

cohort. On the experiment day, the female subjects were placed into the behavior apparatus for 30 minutes for analysis of movement

tracking from a dorsal view, which was used as control (Control trial). After the habituation, the partner (Swiss-Webster) was placed in

the interaction zone. Each assay was conducted 5-10 hrs after the onset of dark period, under an IR light and recorded (15 Hz) for

subsequent 30 to 120 min for analysis of movement tracking and sexual behavior from a dorsal view (Sexual interaction trial).

For **Figure 6**, females underwent two mating assays at > 1 week interval, with different treatment (e.g., saline or CNO). Different males

were used for each assay. For neural activation experiments using hM3Dq (**Figure 6, Figure S9, Figure S10**), 0.1 mL of 1 mg/kg CNO

dissolved in saline or 0.1 mL of saline was I.P. injected into the animal 30 minutes prior to the male entry. For neural inhibition

experiments using hM4Di (**Figure S11**), 0.1 mL of 5 mg/kg CNO dissolved in saline or 0.1 mL of saline was I.P. injected into the animal

30 minutes prior to the male entry.

Sexual behavior was annotated from the recorded video using Behavioral Observation Research Interactive Software (BORIS) in a

frame-by-frame fashion (Friard and Gamba, 2016). Male mounts, intromission, male to female anogenital sniffing, female to male

sniffing, female self-grooming and male ejaculation were annotated based on characteristic postures described previously (Ishii et al.,

2017). Briefly, mount was defined as a stud male using both forepaws to climb onto a female from behind for copulation. Intromission

was defined by stable thrusting from the male animal during mount. Male ejaculation was identified by increase and then termination

of thrusting and more than 10 seconds of immobility from the male, which are postures frequently associated with male ejaculation.

Male to female anogenital sniffing was identified by the male animal following the female subjects anogenital area. Female to male

sniffing was identified by female subject nose approaching male subject. When analyzing sniff-evoked GCaMP activity, to restrict the

analysis to appetitive sniffing behaviors, sniffing behaviors which happened after the first mounting were excluded from the analysis.

Female self-grooming was identified by the female subject moving their head close to their genital region. The receptive ratio was

calculated by dividing the total number of intromissions with the total number of mountings. Behavior annotation was conducted by

volunteers who were blind for the stimulation or AAV condition. For the female self-paced mating assay, the position of the female

subject was tracked from the recorded video using DeepLabCut (Nath et al., 2019) or Ethovision. The amount of time spent in the
interaction zone was quantified using SimBA region of interest (ROI) interface (Nilson et al., 2020). Behavior data from *in vivo* calcium
imaging experiments were down-sampled to 10-Hz to match the imaging data.
For conditional place preference (**Figure S10G–J**), a two chambered apparatus was used with visual (horizontal and vertical stripes)
cues. The floor was backlit and an overhead video camera recorded position (Basler, 3.75 Hz frame rate with Ethovision software,
Noldus). The apparatus was in a sound isolation chamber. *Ad libitum* fed Fos-CreER mice with hM3Dq and eYFP controls were
acclimatized (60 min). The following day, the mouse's position was recorded (1,800 s) and tracked offline using Ethovision (pre-test).
Daily conditioning consisted of two 1,800 s sessions: (1) one side of the chamber paired with saline injection; (2) the other side paired
with CNO (1 mg/kg) injection. Saline or CNO was administrated 10 min prior to the entry to the chamber. The side on which saline or
CNO was administered was counterbalanced. After three conditioning days, mice were given free access to the entire apparatus and
their position was tracked (post-test). The difference of time spent in the CNO zone, the number of entries into the CNO zone and the
velocity between the post- and pre- test.
For pup retrieval test (**Figure S10K–O**), a home cage of the female subject was used. Female subject was i.p. injected with saline as
control or CNO in their home cage. 30-minutes after administration, three 7-day-old pups were placed in the corners of the home cage.
Animal behavior was recorded from the dorsal view for 10 minutes. Parenting behavior was annotated from the recorded video using
Behavioral Observation Research Interactive Software (BORIS) in a frame-by-frame fashion. Pup retrievals, nest building, pup
grooming and pup sniffing were annotated based on characteristic postures. Briefly, pup retrieval was defined as a female subject
carrying pup. Nest building was defined by female subject digging their bedding or carrying nesting material. Pup grooming was
identified by grooming pups without carrying. Female to pup sniffing was identified by female subject nose approaching pup. For neural
activation experiments using hM3Dq (**Figure S10 K–O**), 0.1 mL of 1 mg/kg
CNO dissolved in saline or 0.1 mL of saline was I.P. injected into the animal 30 minutes prior to the pup introduction.
467
***In vivo* calcium imaging experiments and analysis.**
Miniature microscope nVoke 2.0 or nVue (Inscopix) was used to obtain GCaMP6s signal (20 Hz). The microscope was connected
above the behavior apparatus using a commutator (Inscopix) to support the weight of the microscope. On the experiment day, the
subject was habituated to the microscope for 10 minutes. After the habituation, the LED power and gain was adjusted to 0.2-0.5 and
6.0-8.0 respectively depending on the intensity of GCaMP6s expression. Data acquisition started after the adjustment of imaging
parameters. The animal was kept habituated to the microscope for additional > 5 minutes to obtain background signal, and then the
intruder male animal was introduced to the subject's home cage. Calcium activity as well as animal behavior was recorded until 2-10
minutes after male ejaculation. Imaging data were spatially down-sampled by factor of 4 and temporally down-sampled to 10 Hz using
the Inscopix Data Processing software (1.8.0). The data was further filtered by a spatial band-pass filter and motion corrected using
the same software. Masks were manually drawn to extract data from single cell. The data frame containing GCaMP6s signal intensity
traces were exported from the Inscopix Data Processing software and further processed in Python. Signal intensity array was z-scored
using the background signal. For peri event activity analysis in **Figure 3** and **Figure 4**, z-scored signal was collected from -5 to + 15
seconds around the behavior onset. Peri event activity from multiple behavior events were concatenated and averaged. Finally, the
mean value of pre-behavior background (-5 to 0 seconds from onset) was subtracted. The magnitude of z-scored signal was quantified

by averaging 0 to +5 seconds from onset. Cells that had larger than 2 d z-scored signal were classified as "responding" cells. For
persistent activity analysis in **Figure 5**, z-scored signal was collected from -5 to + 120 seconds around the onset of male ejaculation.
The mean value of pre-behavior background (-5 to 0 seconds from onset) was subtracted. The decay length was quantified by
calculating the time the z-scored signal was smaller than 0. The magnitude of z-scored signal was quantified by averaging 0 to +5
seconds from onset. Spectral clustering was conducted using scikit-learn package and previously used custom script (Namboodiri et
al., 2019). Briefly, first dimensionality reduction using principal component analysis was conducted for z-scored signal from -5 to +120
seconds around the onset of male ejaculation. Next, n = 4 principal components were used to project the data. This was further used
as the input into the clustering algorithm while systematically testing nearest neighbor and number of cluster variables. The best variable
that resulted in maximizing the silhouette score was determined (number of clusters = 5, number of nearest neighbors = 55, average
silhouette score = 0.405). All clusters contained cells from multiple animal subjects, indicating the variability of the dataset can be
explained by heterogenous cell population, rather than subject variability.
493
TRAP experiments.
TRAP method was conducted as previously described with modifications (DeNardo et al., 2019). TRAP2::Ai14 or TRAP2 mice were
OVX during isoflurane anesthesia and AAV injected if necessary. After 2 – 3 weeks from OVX surgery, animals were prepared for the
activity labeling assay. Female subjects were subcutaneously (s.c.) injected with 0.1 mL and 0.05 mL Estradiol (E8515, Sigma-Aldrich,
0.1 mg/ml) dissolved in sesame oil (Sigma-Aldrich) at 24 and 48h before assay. On the day for the assay, female subjects were isolated
in their home cage and introduced with a male mating partner to either experience appetitive behaviors, consummatory behaviors or
male ejaculation. For the appetitive group, the male partner was removed after 10–20 minutes. To ensure that the female subject did
not experience any consummatory behavior from the male, the male partner was removed whenever an attempt to mount was observed.
For the consummatory group, the male partner was removed after the first male mounting and intromission was observed. For the
post-ejaculatory group, the male partner was removed just after male ejaculation. Immediately after the removal of the male animal,
the female subject was I.P. injected with 0.125 mL of 50 mg/kg of 4-Hydroxytamoxifen (Sigma-Aldrich, cat#H6278) dissolved in sesame
oil. After the injection, the female subject was isolated in their home cage and kept in the behavior room for overnight to avoid
background stimulation. On the following day, the mice returned to the regular housing.
507
HCR experiments and analysis.
Hybridization chain reaction (HCR) was performed based on the protocol provided by Molecular Instruments with some modifications
(**Figure S5** and **Figure S9D–G**). For *Fos* activity analysis (**Figure S5**), the female subject was first singly housed in the home cage for
1 hour. The male partner was introduced into the home cage and animal behavior was recorded. For the appetitive group, the male
partner was removed after 10–20 minutes until they first attempt mounting to the female subject. For the completion group, the male
partner was removed just after male ejaculation. Twenty-minutes after the removal of the male partner, the female subject was
immediately sacrificed to collect brain tissue. The brain tissue was then snap-frozen using dry ice. Every fifth 20-micron coronal sections
containing the MPOA (3 sections from Bregma +0.10 mm to – 0.10 mm) were collected on a slide glass and fixed with 4% PFA in PBS
for 30 min and rinsed with PBS, followed by dehydration with a series of 50%, 70%, 100%, 100% of ethanol. After final dehydration the
slides were dried. The tissue was treated with Protease4 (ACDBiosystem) for 5–7 minutes. The slides were then rinsed with PBS and

were applied with hybridization buffer (Molecular Instruments) for 10 minutes at 37°C for pre-hybridization. RNA probes for *Fos, Vgat,*
Vglut2, Calcr, Esr1, Gal, Nts, Prlr (1 mM, Molecular Instruments) were diluted in a ratio of 1:100–1:250 into the hybridization buffer.
After pre-hybridization, hybridization buffer (70 µl) containing the probes was applied to each slide on which the parafilm cover was
placed. After 12h–18h of incubation at 37°C in the moisture chamber, the sections were washed, with a series of probe wash buffer
and SSC-T buffer (5x SSC, 0.1% Tween 20) mixture. After the final wash, the sections were incubated in amplification buffer (Molecular
Instruments) for 30 minutes at room temperature for pre-amplification. Amplification hairpin probes (3 mM, Molecular Instruments) were
diluted in a ratio of 1:50 into the hybridization and then snap cooled for 1.5 minutes in 95°C. After pre-amplification, amplification buffer
containing hairpin probes (70 µl) was applied to each slide on which the parafilm cover was placed. After 12h–18h of incubation at
room temperature in the moisture chamber, the sections were washed in 5xSSC-T buffer for 30 minutes x2. The sections were
quenched for autofluorescence using Vector TrueView reagent (Vector, cat# SP-8400). After washed in 2xSSC, the sections were
stained with DAPI in PBS (5 µg/ml, Thermo Fisher Scientific, cat#D1306) for 8 minutes. The slide was then mounted using Vibrance
Antifade Mounting Medium (VECTASHIELD, cat#H-1800). Images were obtained using a conventional microscope (Zeiss, Apotome
Imager.M2) with a 20x objective. After image acquisition, the coverslip was removed and washed in 2x SSC for 10 minutes, then
incubated with DNase1 (0.25 U/µl, Roche, cat#4716728001) for 1.5 h at room temperature. The section was then washed in 2x SSC
for 5 minutes x6. The sections then proceeded to the pre-hybridization step for the next round of HCR. A total of 3 rounds of HCR were
conducted. Images from multiple rounds were aligned using HiPlex image registration software (ACDBiosystem) or manually aligned
using ImageJ package TrackEM2 (Cardona et al., 2012).
For anatomical characterization for TRAP2 animals used in **Figure S9D–G**, subjects were perfused with PBS and 4% PFA-PBS, then
brain was harvested and post-fixed with PFA overnight. Every fourth 40-µm coronal sections containing the MPOA were collected on
a slide glass. The sections went through 1 round of HCR and were stained for *Vgat* and *Vglut2* following the procedures described
above.
The images with each probe signal were aligned to the DAPI channel to identify MPOA subregions and adjacent regions. The image
was processed in HALO v3.2 software (Indica Labs) to segment each cell based on the DAPI signal and to quantify RNA transcript
signal in each cell. Parameters for detecting puncta and intensity of each gene were manually adjusted for each tissue section of each
animal by a volunteer blind to the experimental condition. The cutoff value to determine a gene "expressing cell" was defined as the
mean fluorescent intensity of 3 to 5 transcripts depending on the background fluorescent signal. For supervised clustering of the MPN
cell gene expression matrix (**Figure S5K–M**) was conducted using Seurat3.1.1 in R. Cells that had no mRNA signal were excluded
from the dataset. This expression matrix contained single cells detected across all 10 animals and the spot counts for every gene; we
also included signal intensity measurements as metadata. Once loaded, the HCR digital expression matrix was scaled but not
normalized. During scaling, we regressed out gene intensity measurements on the basis that signal intensity may indicate staining
quality. Next, cells were clustered using the same pipeline. The cells were UMAP plotted using nearest_neighbors = 100,

minimum_distance = 0.05 (**Figure S5K**). The cluster ID for each cell was exported and used for statistical analysis using custom Python
scripts.
551

Immunohistochemistry experiments.
To evaluate the accuracy and efficiency of TRAP labeling, we utilized subjects used in pharmacogenetic experiments (**Figure 6, Figure**
S9) and labeled neurons that respond to male ejaculation using immunohistochemistry against immediate early gene c-Fos. The
TRAP2 female subjects were labeled with eYFP or hM3Dq-mCherry in the appetitive or male ejaculation responding neurons in the
MPOA (appetitive-hM3Dq, consummatory-hM3Dq, post-ejaculatory-hM3Dq or post-ejaculatory-eYFP group). Ninety-minutes before
perfusion, the female subjects experienced male ejaculation during a home cage mating assay (described in "**Behavior experiments**
and analysis.") to activate the male ejaculation responding neurons in the MPOA. The subjects were then perfused with PBS and 4%
PFA-PBS, and the brain was harvested and post-fixed with PFA overnight. Every fourth 40-µm coronal sections containing the MPOA
were collected. were washed three times with PBS-T (1x PBS, 0.3% Triton-X, Sigma-Aldrich, cat# 11332481001) for 10 min x3, and
treated with 5% normal donkey serum (NDS, Sigma-Aldrich, cat# 566460) in PBST for 1 hr at room temperature for blocking. Sections
were then incubated with rabbit anti-cFos antibody (Abcam, cat#190289-Rb, 1:500), diluted into 5% BSA in PBST for overnight at 4°C.
Antibody signals were detected by goat anti-rabbit Alexa 488 or Alexa 594 (Invitrogen, cat#A21206 or cat#A21209, 1:500) for 2h at
room temperature. Sections were washed once with PBST for 10 min, treated with PBS containing DAPI for 20 min, rinsed with PBS,
and mounted with cover glass using Vibrance Antifade Mounting Medium (VECTASHIELD, cat#H-1800).
Sections were imaged by FV-3000 confocal microscope (Olympus) with a 20x objective. Images were processed in cellpose to
segment DAPI positive cells in the MPOA and their subregions (Stringer et al., 2021). DAPI positive cell masks were used to quantify
intensity of c-Fos protein or YFP/mCherry signal. The intensity value was used to determine the negative and positive cell.
569

Brain-wide analysis of activity-dependent labeling experiments.
TRAP2::Ai14 females were used to label neurons responding to appetitive behavior or male ejaculation. Two-weeks after activity
labeling (described in "**TRAP experiments.**"), the animals were perfused with PBS and 4% PFA-PBS, the brain was harvested and
post-fixed with PFA overnight. After post-fixation, the brains were processed for tissue clearing. For tissue clearing, we utilized
stabilization under harsh conditions via intramolecular epoxide linkages to prevent degradation method (SHIELD) (Park et al., 2019).
The procedures followed the SHIELD tissue clearing protocol (Full Active Pipeline Protocol v5.05, LifeCanvas). Briefly, the brain was
incubated in SHIELD OFF solution [2.5 mL DI water, 2.5 mL SHIELD-Buffer Solution, 5 mL SHIELD-Epoxy Solution] at 4oC for 4 days.
Then transferred to SHIELD ON buffer and incubated at 37°C for 24 hrs. Following, SHIELD ON buffer incubation, the brain was pre-
incubated in Delipidization buffer for overnight at room temperature. The tissue was then placed into SmartClear II Pro (LifeCanvas) to
conduct active tissue clearing for 24 hours. To match the refraction index, the tissue was incubated in 50% of EASYINDEX solution
(RI = 1.52, LifeCanvas) at 37°C shaking for 24 hrs and then 100% of EASYINDEX solution at 37°C shaking for 24 hrs. The tissue was
then embedded in EASYINDEX with 2% agar and mounted onto SmartSPIM light sheet microscope (LifeCanvas). The tissue was
mounted on the ventral side facing toward the objective, imaged from the horizontal view using a 3.6x objective with 4-µm zstep. 594
nm channel was collected as the signal channel, 488 or 647 nm channel was collected as the background channel. Background channel

was used for the alignment to the brain atlas. The images were first destriped and then stitched using a custom written script
(LifeCanvas). The images were pre-processed and down sampled by 2 times (final resolution (x,y,z) = (3.6 μm, 3.6 μm, 4 μm)). Images
were registered to a standard brain atlas Unified Anatomical Atlas (Yongsoo Kim Lab) (resolution (x,y,z) = (20 μm, 20 μm, 50 μm))
using ClearMap pipeline on a cloud-based cluster computer (Chon et al., 2019; Madangopal et al., 2022; Renier et al., 2016). tdTomato
expressing cells were segmented through the same pipeline using a classifier created using Ilastik (Berg et al., 2019). To train the
classifier, 10 chunks of images from the 4 of the subjects were randomly sampled and used as the training dataset (Image size, (x,y,z)
= 1.8 mm x 1.8 mm x 100 μm). One chunk image of the same size from each subject was used as the test dataset. The quality of the
classifier was visually examined. The total number of tdTomato positive cells per brain region was quantified and normalized by the
size of each brain region (density, mm^3). In the standard brain atlas, brain regions were hierarchically sorted from broader regions to
finer regions (level 1 to level 8). Level 4 was selected for broader analysis of brain regions (**Figure 2C**). Level 6 was selected for finer
analysis of brain regions (**Figure 2H, Figure S4**). To compare the means of density of tdTomato cell in each brain area between the
appetitive, consummatory and post-ejaculatory group, we conducted an ANOVA test and further corrected for multiple comparisons
with Benjamini/Hochberg correction. False discovery rate (FDR) = 0.05 was used for the broader analysis while a loose discovery rate
(FDR = 0.10) was used for finer analysis due to the large number of comparisons. A post-hoc Tukey's HSD post-hoc test (FWER =
0.05) was conducted for regions that tested significant in the ANOVA test.
599
Quantification, data processing and Statistical Analysis.
Data were presented as mean with error bar of 95% confidence interval unless otherwise mentioned. Data processing and plot
generation was done in Python and R. Plots were further processed in Adobe Illustrator to create Figures. The sample sizes were
chosen based on previous studies using similar behavioral experiments. Codes used for analysis and statistics are provided on Github
(https://github.com/stuberlab/Ishii-Mating-completion.git). The statistical details of each experiment including the statistical tests used,
exact value of n and what n represents, are shown in each figure legend. Each statistical test result is shown in **Table 1**. Significance
was shown as *p < 0.05, **p < 0.01, ***p < 0.001 and not significant values were not noted or shown as n.s.
The sample size for behavior experiments in this study were n = 5-9. This was predetermined based on previous studies examining
female sexual behavior (Ishii et al. 2017, Liu et al. 2022, Yin et al. 2022). To further examine the number of animals required for our
behavioral experiments, we pooled data used in this study and conducted a power analysis (n = 111 pooled data, control n = 94, stim
n = 17). We conducted a power analysis using the variance calculated from pooled average time in isolation zone. These data were
pooled from control animals in each experiment (eg. animals with GFP control virus injected, saline injected, etc.). The average time
in isolation zone in the after ejaculation or after reactivating the completion cells was 420 ± 210 seconds, and 49 ± 91 seconds in the

control group (mean ± s.d.). Within this population, we found that 5 animals were sufficient to detect the difference (p < 0.05, power =
0.8) in Students t-test.

Data distribution
617 Whole brain FosTRAP activity mapping data is available on DANDI (https://dandiarchive.org/dandiset/001249/0.241109.2357). Other
618 data used in this project is shared on Figshare (https://doi.org/10.6084/m9.figshare.27642768.v1). The code for analysis and data
619 generation will be shared on GitHub (https://github.com/stuberlab/Ishii-MPOA-Ejaculation-2023).

Author Contributions:
621 Experiments were conceived and designed by K.K.I. and G.D.S. G.D.S. supervised the project, provided resources, and funding. K.H.
622 provided transcriptional analysis data of the MPOA and assisted in gene selection of HCR experiments. J.C. conducted HCR
623 experiments and histological analysis. S.Y., R.E.F., S.K. and M.S. conducted behavioral experiments as well as behavioral annotations.
624 C.Z. provided *in vivo* imaging data analysis pipelines. J.N., A.D.M, E.R.S., and S.A.G. assisted and supervised tissue clearing
625 experiments and whole-brain data analysis.

Acknowledgments:
627 This work was supported by a JSPS oversea fellowship (K.K.I.), Uehara postdoctoral fellowship (K.K.I.), NIH grants R37 DA032750,
628 R01 DA038168, R01 DA054317, P30 DA048736 and Washington Research Foundation Postdoctoral Fellowship (E.R.S.)

References:
Agrati D. 2022. Adolescence and Postpartum: Two Life Periods to Deepen Our Understanding of the Complexity of Female Rat Sexual Behavior.
Sexes **3**:282–297. doi:10.3390/sexes3020022

Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J, Beier T, Rudy M, Eren K, Cervantes JI, Xu B, Beuttenmueller F,
Wolny A, Zhang C, Koethe U, Hamprecht FA, Kreshuk A. 2019. ilastik: interactive machine learning for (bio)image analysis. *Nature Methods*
16:1226–1232. doi:10.1038/s41592-019-0582-9

Bermant G. 1961. Response Latencies of Female Rats during Sexual Intercourse. *Science* **133**:1771–1773. doi:10.1126/science.133.3466.1771

Bermant G, Westbrook WH. 1966. Peripheral factors in the regulation of sexual contact by female rats. *Journal of Comparative and Physiological*
Psychology **61**:244–250. doi:10.1037/h0023152

Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, Tomancak P, Hartenstein V, Douglas RJ. 2012. TrakEM2 Software
for Neural Circuit Reconstruction. *PLoS ONE* **7**:e38011. doi:10.1371/journal.pone.0038011

Chapman T, Bangham J, Vinti G, Seifried B, Lung O, Wolfner MF, Smith HK, Partridge L. 2003. The sex peptide of Drosophila melanogaster: Female
post-mating responses analyzed by using RNA interference. *Proceedings of the National Academy of Sciences* **100**:9923–9928.
doi:10.1073/pnas.1631635100

Chen PS, Stumm-Zollinger E, Aigaki T, Balmer J, Bienz M, Böhlen P. 1988. A male accessory gland peptide that regulates reproductive behavior of
female D. melanogaster. *Cell* **54**:291–298. doi:10.1016/0092-8674(88)90192-4

Choi HMT, Beck VA, Pierce NA. 2014. Next-Generation in Situ Hybridization Chain Reaction: Higher Gain, Lower Cost, Greater Durability. *ACS Nano*
8:4284–4294. doi:10.1021/nn405717p

Chon U, Vanselow DJ, Cheng KC, Kim Y. 2019. Enhanced and unified anatomical labeling for a common mouse brain atlas. *Nature Communications*
10. doi:10.1038/s41467-019-13057-w

Dai B, Sun F, Tong X, Ding Y, Kuang A, Osakada T, Li Y, Lin D. 2022. Responses and functions of dopamine in nucleus accumbens core during social
behaviors. *Cell Reports* **40**:111246. doi:10.1016/j.celrep.2022.111246

DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, Guenthner CJ, Tessier-Lavigne M, Luo L. 2019. Temporal evolution of cortical
ensembles promoting remote memory retrieval. *Nat Neurosci* **22**:460–469. doi:10.1038/s41593-018-0318-7

Dimén D, Puska G, Szendi V, Sipos E, Zelena D, Dobolyi Á. 2021. Sex-specific parenting and depression evoked by preoptic inhibitory neurons.
iScience **24**:103090. doi:10.1016/j.isci.2021.103090

Erskine MS. 1989. Solicitation behavior in the estrous female rat: A review. *Hormones and Behavior* **23**:473–502. doi:10.1016/0018-
506X(89)90037-8

Erskine MS. 1985. Effects of paced coital stimulation on estrus duration in intact cycling rats and ovariectomized and ovariectomized-
adrenalectomized hormone-primed rats. *Behavioral Neuroscience* **99**:151–161. doi:10.1037/0735-7044.99.1.151

Fang Y-Y, Yamaguchi T, Song SC, Tritsch NX, Lin D. 2018. A Hypothalamic Midbrain Pathway Essential for Driving Maternal Behaviors. *Neuron*
98:192-207.e10. doi:10.1016/j.neuron.2018.02.019

Feng K, Palfreyman MT, Häsemeyer M, Talsma A, Dickson BJ. 2014. Ascending SAG Neurons Control Sexual Receptivity of Drosophila Females.
Neuron **83**:135–148. doi:10.1016/j.neuron.2014.05.017

Flanigan ME, Kash TL. 2022. Coordination of social behaviors by the bed nucleus of the stria terminalis. *Eur J of Neuroscience* **55**:2404–2420.
doi:10.1111/ejn.14991

Franklin KBJ, Paxinos G. 2007. The mouse brain in stereotaxic coordinates. Academic Press.

Friard O, Gamba M. 2016. BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. *Methods in*
Ecology and Evolution **7**:1325–1330. doi:10.1111/2041-210X.12584

Gillott C. 2003. Male Accessory Gland Secretions: Modulators of Female Reproductive Physiology and Behavior. *Annual Review of Entomology*
48:163–184. doi:10.1146/annurev.ento.48.091801.112657

Hashikawa K, Hashikawa Y, Falkner A, Lin D. 2016. The neural circuits of mating and fighting in male mice. *Current Opinion in Neurobiology* **38**:27–
37. doi:10.1016/j.conb.2016.01.006

Hashikawa K, Hashikawa Y, Liu Y, Rossi MA, Basiri ML, Chen JY, Ahmad OR, Mukundan RV, Johnston NL, McHenry JA, Palmiter RD, Rubinow DR,
Zweifel LS, Stuber GD. 2021. Pubertal sex hormones control transcriptional trajectories in the medial preoptic area. *bioRxiv* **1**:2021.09.02.458782.

Hull EM, Wood RI, McKenna KE. 2005. Neurobiology of Male Sexual BehaviorKnobil and Neill's Physiology of Reproduction. Elsevier. pp. 1729–
1824. doi:10.1016/B978-012515400-0/50038-5

Inoue S, Yang R, Tantry A, Davis C ha, Yang T, Knoedler JR, Wei Y, Adams EL, Thombare S, Golf SR, Neve RL, Tessier-Lavigne M, Ding JB, Shah NM.
2019. Periodic Remodeling in a Neural Circuit Governs Timing of Female Sexual Behavior. *Cell* **179**:1393-1408.e16. doi:10.1016/j.cell.2019.10.025

Ishii KK, Osakada T, Mori H, Miyasaka N, Yoshihara Y, Miyamichi K, Touhara K. 2017. A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual
Behaviors in Mice. *Neuron* **95**:123-137.e8. doi:10.1016/j.neuron.2017.05.038

Ishii KK, Touhara K. 2018. Neural circuits regulating sexual behaviors via the olfactory system in mice. *Neuroscience Research*.
doi:10.1016/j.neures.2018.10.009

Karigo T, Kennedy A, Yang B, Liu M, Tai D, Wahle IA, Anderson DJ. 2021. Distinct hypothalamic control of same- and opposite-sex mounting
behaviour in mice. *Nature* **589**:258–263. doi:10.1038/s41586-020-2995-0

Kato A, Sakuma Y. 2000. Neuronal activity in female rat preoptic area associated with sexually motivated behavior, Brain Research.

Kohl J, Babayan BM, Rubinstein ND, Autry AE, Marin-Rodriguez B, Kapoor V, Miyamishi K, Zweifel LS, Luo L, Uchida N, Dulac C. 2018. Functional
circuit architecture underlying parental behaviour. *Nature* **556**:326–331. doi:10.1038/s41586-018-0027-0

Lenschow C, Lima SQ. 2020. In the mood for sex: neural circuits for reproduction. *Current Opinion in Neurobiology* **60**:155–168.
doi:10.1016/j.conb.2019.12.001

Liu H, Kubli E. 2003. Sex-peptide is the molecular basis of the sperm effect in Drosophila melanogaster. *Proceedings of the National Academy of*
Sciences **100**:9929–9933. doi:10.1073/pnas.1631700100

Liu M, Kim DW, Zeng H, Anderson DJ. 2022. Make war not love: The neural substrate underlying a state-dependent switch in female social behavior.
Neuron **110**:841-856.e6. doi:10.1016/j.neuron.2021.12.002

Lo L, Yao S, Kim D-W, Cetin A, Harris J, Zeng H, Anderson DJ, Weissbourd B. 2019. Connectional architecture of a mouse hypothalamic circuit node
controlling social behavior. *Proceedings of the National Academy of Sciences* **116**:7503–7512. doi:10.1073/pnas.1817503116

Madangopal R, Szelenyi ER, Nguyen J, Brenner MB, Drake OR, Pham DQ, Shekara A, Jin M, Choong JJ, Heins C, Komer LE, Weber SJ, Hope BT,
Shaham Y, Golden SA. 2022. Incubation of palatable food craving is associated with brain-wide neuronal activation in mice. *Proc Natl Acad Sci USA*
119:e2209382119. doi:10.1073/pnas.2209382119

Marsh N, Marsh AA, Lee MR, Hurlemann R. 2021. Oxytocin and the Neurobiology of Prosocial Behavior. *Neuroscientist* **27**:604–619.
doi:10.1177/1073858420960111

McGill TE, Coughlin RC. 1970. Ejaculatory reflex and luteal activity induction in Mus musculus. *Journal of reproduction and fertility* **21**:215–220.
doi:10.1530/jrf.0.0210215

McHenry JA, Otis JM, Rossi MA, Robinson JE, Kosyk O, Miller NW, McElligott ZA, Budygin EA, Rubinow DR, Stuber GD. 2017. Hormonal gain control
of a medial preoptic area social reward circuit. *Nature neuroscience* **20**:449–458. doi:10.1038/nn.4487

Mei L, Yan R, Yin L, Sullivan RM, Lin D. 2023. Antagonistic circuits mediating infanticide and maternal care in female mice. *Nature*.
doi:10.1038/s41586-023-06147-9

Micevych PE, Meisel RL. 2017. Integrating Neural Circuits Controlling Female Sexual Behavior. *Frontiers in Systems Neuroscience* **11**:42.
doi:10.3389/fnsys.2017.00042

Minakuchi T, Guthman EM, Acharya P, Hinson J, Fleming W, Witten IB, Oline SN, Falkner AL. 2024. Independent inhibitory control mechanisms for
aggressive motivation and action. *Nat Neurosci* **27**:702–715. doi:10.1038/s41593-023-01563-6

Moffitt JR, Bambah-Mukku D, Eichhorn SW, Vaughn E, Shekhar K, Perez JD, Rubinstein ND, Hao J, Regev A, Dulac C, Zhuang X. 2018. Molecular,
spatial, and functional single-cell profiling of the hypothalamic preoptic region. *Science* **362**. doi:10.1126/science.aau5324

Namboodiri VMK, Otis JM, van Heeswijk K, Voets ES, Alghorazi RA, Rodriguez-Romaguera J, Mihalas S, Stuber GD. 2019. Single-cell activity tracking
reveals that orbitofrontal neurons acquire and maintain a long-term memory to guide behavioral adaptation. *Nature Neuroscience* **22**:1110–1121.
doi:10.1038/s41593-019-0408-1

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. 2019. Using DeepLabCut for 3D markerless pose estimation across species and
behaviors. *Nature Protocols* **14**:2152–2176. doi:10.1038/s41596-019-0176-0

Nilson SR, Goodwin NL, Choong JJ, Hwang S, Wright HR, Norville ZC, Tong X, Lin D, Bentzley BS, Eshel N, Mclaughlin RJ, Golden SA. 2020. Simple
Behavioral Analysis (SimBA) – an open source toolkit for computer classification of complex social behaviors in experimental animals.
doi:10.1101/2020.04.19.049452

Nyuyki KD, Waldherr M, Baeuml S, Neumann ID. 2011. Yes, I Am Ready Now: Differential Effects of Paced versus Unpaced Mating on Anxiety and
Central Oxytocin Release in Female Rats. *PLoS ONE* **6**:e23599. doi:10.1371/journal.pone.0023599

Osakada T, Ishii KK, Mori H, Eguchi R, Ferrero DM, Yoshihara Y, Liberles SD, Miyamichi K, Touhara K. 2018. Sexual rejection via a vomeronasal
receptor-triggered limbic circuit. *Nature communications* **9**:4463. doi:10.1038/s41467-018-07003-5

Parada M, Chamas L, Censi S, Coria-Avila G, Pfaus JG. 2010. Clitoral stimulation induces conditioned place preference and Fos activation in the rat.
Hormones and Behavior **57**:112–118. doi:10.1016/j.yhbeh.2009.05.008

Parada M, Sparks LM, Censi S, Pfaus JG. 2014. Clitoral anesthesia disrupts paced copulation in the female rat. *Physiology & Behavior* **123**:180–186.
doi:10.1016/j.physbeh.2013.10.018

Paredes R. 2003. Medial preoptic area / anterior hypothalamus and sexual motivation. *Scandinavian Journal of Psychology* **44**:203–212.

Park Y-G, Sohn CH, Chen R, McCue M, Yun DH, Drummond GT, Ku T, Evans NB, Oak HC, Trieu W, Choi H, Jin X, Lilascharoen V, Wang J, Truttmann
MC, Qi HW, Ploegh HL, Golub TR, Chen S-C, Frosch MP, Kulik HJ, Lim BK, Chung K. 2019. Protection of tissue physicochemical properties using
polyfunctional crosslinkers. *Nature Biotechnology* **37**:73–83. doi:10.1038/nbt.4281

Peirce JT, Nuttall DL. 1961. Self-paced sexual behavior in the female rat. *Journal of Comparative and Physiological Psychology* **54**:310–313.
doi:10.1037/h0040740

Pfaff DW, Sakuma Y. 1979a. Deficit in the lordosis reflex of female rats caused by lesions in the ventromedial nucleus of the hypothalamus. *The
Journal of physiology* **288**:203–210. doi:10.1113/jphysiol.1979.sp012690

Pfaff DW, Sakuma Y. 1979b. Facilitation of the lordosis reflex of female rats from the ventromedial nucleus of the hypothalamus. *The Journal of
physiology* **288**:189–202. doi:10.1113/jphysiol.1979.sp012690

Pfaus JG, Kleopoulos SP, Mobbs CV, Gibbs RB, Pfaff DW. 1993. Sexual stimulation activates c-fos within estrogen-concentrating regions of the
female rat forebrain. *Brain Research* **624**:253–267. doi:10.1016/0006-8993(93)90085-2

Pfaus JG, Marcangione C, Smith WJ, Manitt C, Abillamaa H. 1996. Differential induction of Fos in the female rat brain following different amounts
of vaginocervical stimulation: modulation by steroid hormones. *Brain Research* **741**:314–330. doi:10.1016/S0006-8993(96)00985-7

Powers B, Valenstein ES. 1972. Sexual receptivity: facilitation by medial preoptic lesions in female rats. *Science (New York, NY)* **175**:1003–5.

Qi L, Iskols M, Greenberg RS, Xiao JY, Handler A, Liberles SD, Ginty DD. 2024. Krause corpuscles are genital vibrotactile sensors for sexual behaviours.
Nature **630**:926–934. doi:10.1038/s41586-024-07528-4

Raam T, Hong W. 2021. Organization of neural circuits underlying social behavior: A consideration of the medial amygdala. *Current Opinion in*
Neurobiology **68**:124–136. doi:10.1016/j.conb.2021.02.008

Renier N, Adams EL, Kirst C, Wu Z, Azevedo R, Kohl J, Autry AE, Kadiri L, Umadevi Venkataraju K, Zhou Y, Wang VX, Tang CY, Olsen O, Dulac C, Osten
P, Tessier-Lavigne M. 2016. Mapping of Brain Activity by Automated Volume Analysis of Immediate Early Genes. *Cell* **165**:1789–1802.
doi:10.1016/j.cell.2016.05.007

Rezával C, Pavlou HJ, Dornan AJ, Chan Y-B, Kravitz EA, Goodwin SF. 2012. Neural Circuitry Underlying Drosophila Female Postmating Behavioral
Responses. *Current Biology* **22**:1155–1165. doi:10.1016/j.cub.2012.04.062

Ring JR. 1944. TTHE ESTROGEN-PROGESTERONE INDUCTION OF SEXUAL RECEPTIVITY IN THE SPAYED FEMALE MOUSE[1]. *Endocrinology* **34**:269–275.
doi:10.1210/endo-34-4-269

Stringer C, Wang T, Michaelos M, Pachitariu M. 2021. Cellpose: a generalist algorithm for cellular segmentation. *Nature Methods* **18**:100–106.
doi:10.1038/s41592-020-01018-x

Troconis EL, Seo C, Guru A, Warden MR. 2023. Serotonin neurons in mating female mice are activated by male ejaculation. *Current Biology* **33**:4926-
4936.e4. doi:10.1016/j.cub.2023.09.071

Tsuneoka Y, Funato H. 2021. Cellular Composition of the Preoptic Area Regulating Sleep, Parental, and Sexual Behavior. *Frontiers in Neuroscience*
15. doi:10.3389/fnins.2021.649159

Uphouse L. 2014. Pharmacology of serotonin and female sexual behavior. *Pharmacology Biochemistry and Behavior* **121**:31–42.
doi:10.1016/j.pbb.2013.11.008

Ventura-Aquino E, Paredes RG. 2023. Being friendly: paced mating for the study of physiological, behavioral, and neuroplastic changes induced by
sexual behavior in females. *Front Behav Neurosci* **17**:1184897. doi:10.3389/fnbeh.2023.1184897

11/29/2024
Kentaro Ishii

Wei Y, Wang S, Jiao Z, Zhang W, Lin J, Li X-Y, Li S, Zhang X, Xu X. 2018. Medial preoptic area in mice is capable of mediating sexually dimorphic
behaviors regardless of gender. *Nature Communications* **9**:279. doi:10.1038/s41467-017-02648-0

Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. 2014. Galanin neurons in the medial preoptic area govern parental behaviour. *Nature*
509:325–30. doi:10.1038/nature13307

Yang CF, Chiang MC, Gray DC, Prabhakaran M, Alvarado M, Juntti SA, Unger EK, Wells JA, Shah NM. 2013. Sexually dimorphic neurons in the
ventromedial hypothalamus govern mating in both sexes and aggression in males. *Cell* **153**:896–909. doi:10.1016/j.cell.2013.04.017

Yapici N, Kim YJ, Ribeiro C, Dickson BJ. 2008. A receptor that mediates the post-mating switch in Drosophila reproductive behaviour. *Nature*
451:33–37. doi:10.1038/nature06483

Yin L, Hashikawa K, Hashikawa Y, Osakada T, Lischinsky JE, Diaz V, Lin D. 2022. VMHvllCckar cells dynamically control female sexual behaviors over
the reproductive cycle. *Neuron* **110**:3000-3017.e8. doi:10.1016/j.neuron.2022.06.026

Yoshihara C, Tokita K, Maruyama T, Kaneko M, Tsuneoka Y, Fukumitsu K, Miyazawa E, Shinozuka K, Huang AJ, Nishimori K, McHugh TJ, Tanaka M,
Itohara S, Touhara K, Miyamichi K, Kuroda KO. 2021. Calcitonin receptor signaling in the medial preoptic area enables risk-taking maternal care.
Cell Reports **35**:109204. doi:10.1016/j.celrep.2021.109204

Zhang G-W, Shen L, Tao C, Jung A-H, Peng B, Li Z, Zhang LI, Tao HW. 2021. Medial preoptic area antagonistically mediates stress-induced anxiety
and parental behavior. *Nat Neurosci* **24**:516–528. doi:10.1038/s41593-020-00784-3

Zhang SX, Lutas A, Yang S, Diaz A, Fluhr H, Nagel G, Gao S, Andermann ML. 2021. Hypothalamic dopamine neurons motivate mating through
persistent cAMP signalling. *Nature* **597**:245–249. doi:10.1038/s41586-021-03845-0

Zhou X, Li A, Mi X, Li Yixuan, Ding Z, An M, Chen Y, Li W, Tao X, Chen X, Li Ying. 2023. Hyperexcited limbic neurons represent sexual satiety and
reduce mating motivation. *Science (New York, NY)* **379**:820–825. doi:10.1126/science.abl4038

Figures:

786
Figure 1: Female mice show decreased sexual motivation after male ejaculation.
(**A**) Schematic of female self-paced mating assay. The behavior apparatus was divided by a wall with a hole small enough for only the
female subject to go through. The female subject was allowed to freely choose between "interaction" with the male partner in the larger

zone or "isolation" in the smaller zone. Results using ovariectomized and hormone primed female mice are shown in **B – M**. Results
using naturally estrus female mice are shown in **N – T**.

(**B**) Raster plots of time spent in the interaction zone (gray). The onset of male ejaculation, start and the endpoint of the experiment is
shown in yellow, blue and red respectively.

(**C – E**) Raster plots of time spent in the interaction zone (gray) around the onset of the first female to male sniffing, male mounting,
intromission and male ejaculation.

(**F**) Representative heatmap showing the time spent in the behavior apparatus during a control trial or a sexual interaction trial. The
sexual interaction trial was further divided into pre- and post- male ejaculation.

(**G–H**) Behavior analysis for hormone primed animals. (**G**) Number of transitions between the zones per 10 minutes. RM ANOVA;
$F_{(2,16)}=10.89$, **$p=0.001035889$. Tukey's HSD; Control vs Post, **$p=0.0015$. Control vs Pre, $p=0.6284$. Post vs Pre, *$p=0.0142$. (**H**)
Average time spent in the isolation zone during the control trial, and pre- and post- male ejaculation. RM ANOVA; $F_{(2,16)}=14.34$,
$p=0.00027$. Control vs Post, ***$p=0.0004$. Control vs Pre, $p=0.5361$. Post vs Pre, **$p=0.0057$. (I**) Latency to return to the interaction
zone after sniff, male mount or male ejaculation. RM ANOVA; $F_{(2,16)}=11.87$, **$p=0.00069052$. Ejaculation vs Sniff, **$p=0.0016$.
Ejaculation vs Mount, **$p=0.0077$. Sniff vs Mount, $p=0.7951$. (**J**) Number of female-to-male sniffing per 10 minutes during pre- and
post- male ejaculation. $p=0.65234375$. (**K**) Number of male mounting per 10 minutes during pre- and post- male ejaculation.
$p=0.00390625$. (L**) Number of intromission per 10 minutes during pre- and post- male ejaculation. **$p=0.00390625$. (**M**) Number of
female self-grooming per 10 minutes during pre- and post- male ejaculation. *$p=0.025061844$.

(**N–T**) Behavior analysis for naturally estrus animals. (**N**) Number of transitions between the zones per 10 minutes. RM ANOVA; $F_{(2,10)}$
$= 7.54$, *$p=0.01007$. Tukey's HSD; Control vs Post, $p=0.0871$. Control vs Pre, $p=0.2727$. Post vs Pre, **$p=0.0038$. (**O**) Average time
spent in the isolation zone during the control trial, and pre- and post- male ejaculation. RM ANOVA; $F_{(2,10)} = 15.71$, ***$p=0.00082$.
Control vs Post, ***$p=0.0002$. Control vs Pre, $p=0.9862$. Post vs Pre, ***$p = 0.0002$. (**P**) Latency to return to the interaction zone after
sniff, male mount, or male ejaculation. RM ANOVA; $F_{(2,10)} = 38.36$, ***$p=0.00002$. Ejaculation vs Sniff, ***$p=0$. Ejaculation vs Mount,
***$p=0$. Sniff vs Mount, $p=0.6201$. (**Q**) Number of female-to-male sniffing per 10 minutes during pre- and post- male ejaculation. *p
$=0.03125$. (**R**) Number of male mounting per 10 minutes during pre- and post- male ejaculation. *$p=0.03125$. (**S**) Number of intromission
per 10 minutes during pre- and post- male ejaculation. *$p=0.03125$. (**T**) Number of female self-grooming per 10 minutes during pre-
and post- male ejaculation. $p=0.3452$.

All data are shown as mean ± 95% confidence interval and were analyzed by RM ANOVA with Tukey HSD post-hoc test (FWER =
0.05) (**G–I** and **N–P**) or Wilcoxon signed-rank test (**J–M** and **Q–T**). n = 9 (**D–K**), n = 6 (**N–T**). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; ns, not
significant.

11/29/2024
Kentaro Ishii

Figure 2: Brain wide analysis of cells responding to male ejaculation in the female brain.

(**A**) Overview of the experiment pipeline. After the activity labeling using TRAP, brain tissue was collected and cleared using SHIELD.

The cleared tissue was imaged using a light-sheet microscope. The dataset was registered to a reference brain atlas and the number

of tdTomato+ cells were quantified using ClearMap and ilastik.

(**B**) Representative image showing cell segmentation.

(**C**) Density of tdTomato+ cells per mm^3 in larger brain regions. CTXpl: Cortical plate, CTXsp: Cortical subplate, STR: Striatum, PAL:

Pallidum, VERM: Vermal regions, HEM: Hemispheric regions, TH: Thalamus, HY: Hypothalamus, MBsen: Midbrain, sensory related,

MBmot: Midbrain, motor related, MBsta: Midbrain, behavioral state related, P: Pons.

(**D**) Heatmap showing density of tdTomato+ cells per mm^3 in subregions. Each column indicates data from one subject. Each row

indicates data from one subregion.

(**E**) Heatmap of average tdTomato density for each group from representative coronal planes.

(**F**) Scatterplot showing difference of average tdTomato density and p-value after Tukey's HSD post-hoc test between the post-

ejaculatory group and the appetitive group.

(**G**) Scatterplot showing difference of average tdTomato density and p-value after Tukey's HSD post-hoc test between

theconsummatory group and the appetitive group.

(**H**) Density of tdTomato+ cells per mm^3 in a subset of subregions. The full list of brain regions is shown in **Figure S2.**

MPN: Medial preoptic nucleus, MPA: Medial preoptic area, BNST: Bed nuclei of the stria terminalis, LPO: Lateral preoptic area, VLPO:

Ventrolateral preoptic nucleus, Me: Medial amygdaloid nucleus, CO: cortical amygdaloid nucleus, BMP: Basomedial amygdaloid

nucleus, posterior part, BLA: Basolateral amygdaloid nucleus, anterior part, BLP: Basolateral amygdaloid nucleus, posterior part, VMH:

Ventromedial hypothalamic nucleus, Arc: Arcuate hypothalamic nucleus, DM: Dorsomedial hypothalamic nucleus, LH: Lateral

hypothalamic area, PMV: Premamillary nucleus, ventral part, PrG: Pregeniculate nucleus of the prethalamus, PHA: Posterior

hypothalamic area.

All data are shown as mean ± 95% confidence interval and were analyzed by ANOVA test with Tukey's HSD post-hoc test (FWER =

0.05) (**C**: FDR = 0.05, **I**: FDR = 0.10). Detailed statistical values are shown in **Table 2** and **Table 3.** Appetitive (green), n = 4.

Consummatory (blue), n = 5. Post-ejaculatory (orange), n = 5. *p < 0.05, ns, not significant.

Figure 3 Male ejaculation signal is strongly represented in the MPOA of the female mice.

(**A**) Schematic of *in vivo* calcium imaging from MPOA Vgat and Vglut2 neurons. AAVDj EF1a-DIO-GCaMP6s was injected into the

MPOA of Vgat-Cre or Vglut2-Cre female mice. A GRIN lens was placed above the MPOA following virus injection.

(**B**) Calcium activity was imaged from a female subject during a free moving mating assay with a sexually experienced male partner.

(**C**) Representative maximum projected image of the field of view during calcium imaging. Vgat: 87 ± 28 cells, n = 6. Vglut2: 68 ± 42

cells, n = 4.

(**D**) Representative coronal section showing GCaMP6s expression and GRIN lens placement.

(**E**) Representative raster plot showing behavior events during the mating assay and a heatmap showing calcium activity from cells

imaged during the assay. Behaviors quantified: female to male sniffing (Sniff), female self-grooming (Self-groom), male to female

anogenital sniffing (Ano. sniff), male mounting (Mount), male intromission (Intromission) and male ejaculation (Ejaculation).

(**F**) Heatmap showing peri-event calcium activity around behavior onset. Cells are ordered by the average response after male

ejaculation.

(**G**) Line plot showing average calcium activity around behavior onset. Vgat neurons, blue. Vglut cells, orange.

(**H**) Average calcium response magnitude following behavior onset per cell. Vgat-Cre: n = 522 cells, Vglut2-Cre: n = 271 cells. Sniff:

$t(791)=6.46$, ***p=1.08E-09. Grooming: $t(791)=-7.25$, ***p=5.89E-12. Male anogenital sniffing: $t(791)=6.64$, ***p=3.41E-10. Mount:

$t(791)=0.34$, p=1. Intromission: $t(791)=-0.32$, p=1. Ejaculation: $t(791)=4.00$, ***p=0.00042.

(**I**) Proportion of cells which responded positively or negatively to behavior onset (>2δ). Positive: Sniff; Vgat 29/522 cells vs. Vglut2

1/271 cells, **p=0.00169. Grooming; Vgat 2/522 cells vs. Vglut2 0/271 cells, p=1. Male Anogenital Sniffing; Vgat 9/522 cells vs. Vglut2

0/271 cells, p=0.178. Mount; Vgat 58/522 cells vs. Vglut2 22/271 cells, p=1. Intromission; Vgat 57/522 cells vs. Vglut2 26/271 cells,

p=1. Ejaculation; Vgat 184/522 cells vs. Vglut2 35/271 cells, ***p=1.52E-10. Negative: Sniff; Vgat 0/522 cells vs. Vglut2 0/271 cells, no

statistic. Grooming; Vgat 5/522 cells vs. Vglut2 0/271 cells, p=0.636. Male Anogenital Sniffing; Vgat 0/522 cells vs. Vglut2 0/271 cells,

no statistic. Mount; Vgat 7/522 cells vs. Vglut2 3/271 cells, p=1. Intromission; Vgat 17/522 cells vs. Vglut2 4/271 cells, p=0.831.

Ejaculation; Vgat 33/522 cells vs. Vglut2 17/271 cells, p=1.

All data are shown as mean ± 95% confidence interval and were analyzed by Student's t-test with Bonferroni correction (**H**) and

chisquare test with Bonferroni correction (**I**). *p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.

Figure 4: Heterogenous response properties to sexual behaviors in the MPOA.

(**A**) Heatmap showing peri-event calcium activity around behavior onset of cells that were categorized based on response magnitude. S: Sniff, M: Mount, E: Ejaculation.

(**B**) Line plot showing average calcium activity around behavior onset. Sniff cells, blue. Mount cells, green. Ejaculation cells, magenta.

(**C**) Average calcium response magnitude following behavior onset per cell in each cell types. Sniff: One-way ANOVA, $F_{(2,235)}=73.35$, ***$p=1.77E-25$. Tukey's HSD test, Ejaculation cells vs Mount cells, $p=0.9996$. Ejaculation cells vs Sniff cells, ***$p=0$. Mount cells vs Sniff cells, ***$p=0$. Grooming: One-way ANOVA, $F_{(2,235)}=0.09$, $p=0.915$. Tukey's HSD test, Ejaculation cells vs Mount cells, $p=0.9723$. Ejaculation cells vs Sniff cells, $p=0.9236$. Mount cells vs Sniff cells, $p=0.9695$. Male Anogenital Sniffing: One-way ANOVA,

F(2,235)=12.34, ***p=8.01E-06. Tukey's HSD test, Ejaculation cells vs Mount cells, ***p=0. Ejaculation cells vs Sniff cells, p=0.6794.
Mount cells vs Sniff cells, **p=0.0048. Mount: One-way ANOVA, F(2,235)=68.32, ***p=4.1E-24. Tukey's HSD test, Ejaculation cells vs
Mount cells, ***p=0. Ejaculation cells vs Sniff cells, p=0.8726. Mount cells vs Sniff cells, ***p=0. Intromission: One-way ANOVA,
F(2,235)=15.72, ***p=3.9E-07. Tukey's HSD test, Ejaculation cells vs Mount cells, ***p=0. Ejaculation cells vs Sniff cells, p=0.5451.
Mount cells vs Sniff cells, ***p=0.0009. Ejaculation: One-way ANOVA, F(2,235)=49.59, ***p=1.08E-18. Tukey's HSD test, Ejaculation
cells vs Mount cells, ***p=0. Ejaculation cells vs Sniff cells, ***p=0.0001. Mount cells vs Sniff cells, p=0.7295.
(**D**) Venn diagram showing the number of cells with positive and negative response in each cell type. (>2δ).
(**E**) Representative field of view showing spatial location of cells with different response properties to sexual behavior in Vgat (left) and
Vglut2 (right) cells.
All data are shown as mean ± 95% confidence interval and were analyzed by Student's t-test with Bonferroni correction (**C**). Sniff cells:
n = 12 cells. Mount cells: n = 48 cells. Ejaculation cells: n = 178 cells. *p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.
895
896

Figure 5 MPOA inhibitory neurons display prolonged activity late after the onset of male ejaculation in female mice.

(**A**) Heatmap showing peri-event calcium activity around male ejaculation onset for male ejaculation-responding cells in a longer time scale.

(**B**) Line plot showing average calcium activity of Vgat and Vglut2 population around behavior onset.

(**C**) The decay length for male ejaculation-responding cells from Vgat and Vglut2 population. Vgat-Cre: n = 184 cells. Vglut2-Cre: n = 35 cells. t(217)=3.039653717, **p=0.002659548.

(**D–G**) Cluster based analysis of the activity pattern of the male ejaculation-cells.

(**D**) Heatmap showing peri-event calcium activity around male ejaculation onset sorted by cluster.

(**E**) Representative calcium activity traces of cells from each cluster.

(**F**) The decay length for male ejaculation-responding cells from each cluster. Further details of statistical tests are shown in **Table 1**.

(**G**) Average z-scored response for male ejaculation-responding cells from each cluster. Further details of statistical tests are shown in
Table 1.
(**H–K**) Cluster based analysis of the activity pattern of the male ejaculation-cells in Vgat and Vglut2 population.
(**H–I**) Heatmap showing peri-event calcium activity of Vgat (H) and Vglut2 (I) cells around male ejaculation onset sorted by cluster.
(**J**) Line plot showing average calcium activity of Vgat and Vglut2 cells in each cluster around behavior onset.
(**K**) Proportion of Vgat and Vglut2 cells in each cluster over Vgat and Vglut2 cells in all clusters. Statistical results are shown next to
the cluster name. Type1; Vgat 26/184 cells vs Vglut2 16/35 cells, ***p=6.80E-05. Type2; Vgat 33/184 cells vs Vglut2 1/35 cells,
p=0.119810823. Type3; Vgat 37/184 cells vs Vglut2 0/35 cells, *p=0.018. Type4; Vgat 47/184 cells vs Vglut2 13/35 cells,
p=0.792188347. Type5; Vgat 41/184 cells vs Vglut2 5/35 cells, p=1.
(**L**) Representative field of view showing spatial location of cells with different response properties after male ejaculation in Vgat (left)
and Vglut2 (right) cells
All data are shown as mean ± 95% confidence interval and were analyzed by Student's t-test (**C**), ANOVA with post-hoc Tukey HSD
test (**F** and **G**), and chisquare test with Bonferroni correction (**K**). Type 1: n = 42 cells, Vgat-Cre: n = 26 cells, Vglut2: n = 16 cells. Type
2: n = 34 cells, Vgat-Cre: n = 33 cells, Vglut2: n = 1 cells. Type 3: n = 37 cells, Vgat-Cre: n = 37 cells, Vglut2: n = 0 cells. Type 4: n =
60 cells, Vgat-Cre: n = 47 cells, Vglut2: n = 13 cells. Type 5: n = 46 cells, Vgat-Cre: n = 41 cells, Vglut2: n = 5 cells.
(**C** and **K**) *p < 0.05, **p < 0.01, ∗∗∗p < 0.001; ns, not significant. (**F** and **G**) a vs. b: p < 0.001. b vs. c: p < 0.05. a vs. c: p < 0.001.
924

Figure 6: Activation of a subpopulation of MPOA labeled following mating is sufficient to suppress female sexual behavior.

(**A**) Schematic of selective labeling of MPOA neurons using TRAP2. AAV5 EF1a-DIO-hM3Dq-mCherry or AAV5 EF1a-DIO-eYFP as control was injected into the MPOA of the female TRAP2 subjects. After incubation of the virus, the female subjects went through procedures to TRAP cells in MPOA (further described in **C**).

(**B**) Schematic of pharmaco-genetic activation of MPOA neurons during a female self-paced mating assay. Female subjects were primed with hormone before the experiment day. On the experiment day, subjects were injected i.p. with CNO to activate TRAP cells,

or saline as control. 30 minutes after the injection, the male partner was placed into the apparatus and the animal behavior was
recorded for 2 hrs.
(**C**) After incubation of the virus, the female subjects were administered with hormones, then received an ejaculation from the male
partner or only interacted without mating. Immediately after, the subjects were administrated with 4-hydroxytamoxifen (4-OHT).
(**D**) Representative image showing TRAP cells MPN from post-ejaculatory-eYFP, appetitive-hM3Dq and post-ejaculatory-hM3Dq group.
Further histological analysis is shown in **Figure S9**. Scale bar, 100 μm.
(**E**) Raster plots of time spent in the interaction zone the female self-paced mating assay (post-ejaculatory-eYFP: gray. appetitive-
hM3Dq: green, consummatory-hM3Dq: blue, post-ejaculatory-hM3Dq: orange).
(**F**) Number of zone transitions per 10 minutes. Post-ejaculatory-hM3Dq; *p=0.03125. Post-ejaculatory-eYFP; p=1. Appetitive-hM3Dq;
p=0.875. Consummatory-hM3Dq; p=0.0625.
942
(**G**) Average time in isolation zone. Post-ejaculatory-hM3Dq; *p=0.03125. Post-ejaculatory-eYFP; p=1. Appetitive-hM3Dq; p=0.875.
Consummatory-hM3Dq; p=0.125.
(**H**) Number of female-to-male sniff. Post-ejaculatory-hM3Dq; p=0.11197. Post-ejaculatory-eYFP; p=0.375. Appetitive-hM3Dq; p=1.
Consummatory-hM3Dq; p=0.4594.
(**I**) Number of female self-grooming. Post-ejaculatory-hM3Dq; p=0.0625. Post-ejaculatory-eYFP; p=1. Appetitive-hM3Dq; p=1.
Consummatory-hM3Dq; p=0.552.
(**J**) Number of male mount. Post-ejaculatory-hM3Dq; p=0.0625. Post-ejaculatory-eYFP; p=0.875. Appetitive-hM3Dq; p=1.
Consummatory-hM3Dq; p=1.
(**K**) Number of male intromission. Post-ejaculatory-hM3Dq; *p=0.03125. Post-ejaculatory-eYFP; p=0.625. Appetitive-hM3Dq; p=1.
Consummatory-hM3Dq; p=0.2716.
(**L**) Receptive score (number of mount episodes with intromissions/number of all mount episodes). Post-ejaculatory-hM3Dq; p=0.03125.
Post-ejaculatory-eYFP; p=1. Appetitive-hM3Dq; p=0.625. Consummatory-hM3Dq; p=0.2716. (**M**) Schematic illustration summarizing
how MPOA neurons suppress female sexual motivation after male ejaculation.
All data are shown by thin lines for individual subjects and thick lines for mean. The data were analyzed by Wilcoxon rank sum test
with Bonferroni correction (**F–L**). post-ejaculatory-eYFP: n = 6, appetitive-hM3Dq: n = 7, consummatory-hM3Dq: n = 8. post-ejaculatory-
hM3Dq: n = 8. *p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.
959

Supplementary Figures:

Figure S1: Sexual experience increase sexual motivation in female mice, related to Figure 1.

(**A**) Diagram showing the experiment design. Virgin animals were tested in a self-paced mating assay without any sexual experience.

SE (sexually experienced) animals experienced male ejaculation in their home cage >7 days before the self-paced mating assay.

(**B**) The number of entries in the isolation zone per 10 minutes between virgin animals and sexually experienced animals.

*p=0.041125541.

(**C**) The average time spent in the isolation zone between virgin animals and sexually experienced animals. *p=0.025974026.

All data are shown as mean ± 95% confidence interval and were analyzed by Mann-Whitney U test (**B** and **C**). Virgin: n = 6, SE: n = 6.

*p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.

Figure S2: Reduction of sexual motivation after male ejaculation in female mice is not dependent on the state of the male
mouse, related to Figure 1.

(**A**) Diagram showing the experiment design. First animals were put in a self-paced mating assay for 30 minutes (pre-ejaculation assay).
After >7 days, the animals were put in a home cage mating assay until they experienced male ejaculation. Immediately after, they were
put in a self-paced mating assay for 30 minutes (post-ejaculation assay). During each assay, the female mouse encountered a novel
sexually active male mouse (male 1, 2 and 3).

(**B**) The number of entries in the isolation zone per 10 minutes between pre and post ejaculation assays. *p=0.027281171.

(**C**) The average time spent in the isolation zone between pre and post ejaculation assays. *p=0.027707849.

All data are shown as mean ± 95% confidence interval and were analyzed by Wilcoxon signed-rank test (**B** and **C**). n = 7.
*p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.

Figure S3: Brain-wide analysis of male ejaculation-responding cells in the female brain related to Figure 2 and Table 3.
Average density heatmap of tdTomato+ cells in the appetitive, consummatory and post-ejaculatory group. Appetitive, n = 4. Consummatory, n = 5. Post-ejaculator, n = 5.

Appetitive Consummatory Post-ejac.

Figure S4: Brain-wide analysis of neurons active during post-ejaculatory period in the female brain related to Figure 2 and Table 3.

Density of tdTomato+ cells in each brain region from post-ejaculatory, consummatory and appetitive group. Statistical significances are shown above each column. The regions are sorted from the top left to bottom right by the qvalue and then the difference in the density between post-ejaculatory group and appetitive group. Details of statistical values and full name of acronyms are shown in **Table 3**. Appetitive, n = 4. Completion, n = 5. Post-ejaculatory, n = 5. ANOVA with Benjamini/Hochberg correction (FDR = 0.10), *q < 0.10; ns, not significant.

1001

Figure S5: Male ejaculation responding cells are composed of a subset of excitatory and inhibitory cells in the MPOA.

(**A**) Brain sections went through multiplexed *in situ* RNA hybridization chain reaction (HCR) to identify cell type markers of MPOA and an immediate early gene *Fos. Vgat, Vglut2, Esr1, Gal, Calcr, Nts, Prlr* were used as cell type markers.

(**B**) Schematic of tissue sampling for *Fos* expression analysis in the MPOA. Wild type female subjects were administered with hormones before the experiment day. On the experiment day, male partners entered the female home cage. Male partners were removed

immediately after ejaculation ("post-ejaculatory" group) or during the appetitive phase before they showed mounting ("appetitive" group).

Brain tissue was harvested 20 minutes after the male exit.

(**C**) Representative coronal section showing *Fos* mRNA expression in the MPOA of Appetitive and Post-ejaculatory group.

(**D**) Proportion of *Fos+* cells in each brain subregion. MPN: medial preoptic nucleus, MPA: medial preoptic area, BNST: bed nucleus

of stria terminalis, LPO: lateral preoptic area, vLPO: ventral lateral preoptic area. BNST; $t(8)=-0.369$, $p=1$. LPO; $t(8)=-0.3094$, $p=1$.

MPA; $t(8)=-1.841$, $p=0.514$. MPN; $t(8)=-4.2709$, *$p=0.0136$. VLPO; $t(8)=-0.1751$, $p=1$.

(**E**) Representative coronal section showing expression of 7 genes: *Fos, Vgat, Vglut2, Esr1, Gal, Calcr, Nts*.

(**F–H**) Expanded image of area highlighted in yellow in (E). (F) Expression of *Fos, Vgat* and *Slc17a6*. (**G**) Expression of *Fos, Vgat* and

Gal. (H) Expression of *Fos, Calcr* and *Nts*. Colored arrow indicates cells expressing given gene. Scale bar, 50 μm.

(**I**) Proportion of *Vgat+* or *Vglut2+* and *Fos+* cells in the MPN of Appetitive and post-ejaculatory group. *Vglut2*; $t(8)=-3.3885$, *$p=0.019$.

Vgat; $t(8)=-3.5361$, *$p=0.0153$.

(**J**) Proportion of *Vgat+ Fos+* or *Vglut2+ Fos+* and cell type marker expressing cells in the MPN of Appetitive and post-ejaculatory

group. *Vgat+* Calcr+; $t(8)=-3.6646$, *$p=0.0318$. Vgat+ Esr1+; $t(8)=-3.1442$, $p=0.0686$. Vgat+ Gal+; $t(8)=-4.0896$, *$p=0.0174$. *Vgat+ Nts+*;

$t(8)=-2.6382$, $p=0.149$. *Vgat+ Prlr+*; $t(8)=-3.9487$, *$p=0.0212$. *Vglut2+ Calcr+*; $t(8)=-3.9487$, *$p=0.0146$. *Vglut2+ Esr1+*; $t(8)=-3.9487$,

$p=0.8223$. *Vglut2+ Gal+*; $t(8)=-3.9487$, $p=0.1299$. *Vglut2+ Nts+*; $t(8)=-3.9487$, $p=0.1951$. *Vglut2+ Prlr+*; $t(8)=-3.9487$, $p=0.1038$.

(**K**) UMAP plot of MPN cells.

(**L**) Disc plot showing proportion of gene expressing cell and the number of copies per cell for each gene and cluster. The proportion

of cells from each group are shown on the right. Cluster *Vglut2+*; $t(8)=-4.1631$, *$p=0.0189$. Cluster *Vgat+*; $t(8)=-5.9018$, **$p=0.0022$.

Cluster *Gal+*; $t(8)=-3.9208$, *$p=0.0265$. Cluster *Calcr+*; $t(8)=-1.5438$, $p=0.9993$. Cluster *Nts+*; $t(8)=-2.1595$, $p=0.377$. Cluster N.A.; $t(8)=-$

1.8254, $p=0.6322$.

(**M**) Proportion of *Fos+* cells from Appetitive and post-ejaculatory group in each cluster.

All data are shown as mean ± 95% confidence interval and were analyzed by Student's t-test with Bonferroni correction (**D, E, K** and

N). appetitive group, n = 5. post-ejaculatory group, n = 5. *p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.

1030

1031

1032

Figure S6: Lens placement and field of view of *in vivo* imaging experiment related to Figure 3, Figure 4, Figure 5.

(**A**) Schematic illustration showing position of GRIN lens placement from each animal. Positions shown on the left hemisphere are from Vglut2 animals, the ones on the right hemisphere are from Vgat animals.

(**B**) Field of view and region of interest (ROI) for each animal. The subject ID and the number of ROI in each FOV are shown above. Vgat-Cre: n = 6, Vglut2-Cre: n = 4.

1038

1039

1040

Figure S7: Distribution of subjects per cluster related to Figure 4.

Cells that had significant positive or negative response (>2δ) to each behavior onset were categorized as positive- or negative-responding cells respectively. Outer ring indicates the mouse line Vgat or Vglut2. The inner ring indicates each subject. Vgat-Cre: n = 6, Vglut2-Cre: n = 4.

1045

Figure S8: Distribution of subjects per cluster related to Figure 5.

Cells that had significant positive or negative response (>2δ) to each behavior onset were categorized as positive- or negative-responding cells respectively. Outer ring indicates the mouse line Vgat or Vglut2. The inner ring indicates each subject. Vgat-Cre: n = 6, Vglut2-Cre: n = 4.

Figure S9: Histological analysis of TRAP labeling related to Figure 6.

(**A**) Histological analysis of the distribution of TRAP labeled cells. Representative coronal section showing eYFP or hM3Dq-mCherry expression in the MPOA. Scale bar, 100 μm.

(**B–C**) Data from post-ejaculatory-eYFP and post-ejaculatory-hM3Dq were grouped into post-ejaculatory group. Appetitive group, n = 7. Consummatory group, n = 5. post-ejaculatory group, n = 14.

(**B**) Number of TRAP labeled cells. One-way ANOVA, $F_{(2,24)}=0.3072$, p=0.7383.

(**C**) The proportion of TRAP labeled cells in the MPN. One-way ANOVA, $F_{(2,24)}=4.3335$, *p=0.0247. Tukey HSD; Appetitive vs Consummatory, p=0.9975. Appetitive vs Post-ejaculatory, *p=0.0471. Consummatory vs Post-ejaculatory, p=0.0984.

(**D**) *In situ* RNA hybridization analysis of the excitatory and inhibitory marker gene expression in the MPN. appetitive-hM3Dq and post-ejaculatory-hM3Dq groups were used for analysis. Representative coronal section showing mCherry, *Vgat* and *Vglut2* expression in the MPOA. Enlarged image indicated by the yellow box, is shown on the right.

(**E–G**) Number of *Vgat*+ TRAP labeled cells in the MPN (**E,** t(11)=-1.3299, p=0.2105), number of *Vglut2*+ TRAP labeled cells in the MPN (**F**, t(11)=-0.7917, p=0.4453.) and the proportion of TRAP labeled cells that express *Vgat* or *Vglut2* (**G,** Appetitive, t(10)=6.6154, **p=0.0024. Post-ejaculatory; t(12)=13.9477, ***p=1.69E-05). Appetitive group, n = 6. Post-ejaculatory group, n = 7.

(**H**) Antibody analysis of c-Fos expression after reactivation of male ejaculation ensemble. Representative coronal section showing mCherry and c-Fos expression in the MPOA. Enlarged image indicated by the yellow box, is shown on the right.

(**I–L**) proportion of c-Fos+ and TRAP labeled cells in the MPN (**I,** One-way ANOVA, $F_{(2,17)} = 3.2569$, p=0.0635. Tukey HSD; Appetitive vs Consummatory, p=0.9991. Appetitive vs Post-ejaculatory, p=0.0942. Consummatory vs Post-ejaculatory, p=0.1268.), proportion of c-Fos+ cells in the MPN (**J,** One-way ANOVA, $F_{(2,17)}=3.1285$, p=0.0697. Tukey HSD; Appetitive vs Consummatory, p=0.1724. Appetitive vs Post-ejaculatory, p=0.8445. Consummatory vs Post-ejaculatory, p=0.0631), the accuracy score (proportion of c-Fos+ TRAP labeled cells among all TRAP labeled cells) (**K,** One-way ANOVA, $F_{(2,17)}=0.4666$, p=0.6349) and the efficiency score (proportion of c-Fos+ TRAP labeled cells among all c-Fos+ cells) (**L,** One-way ANOVA, $F_{(2,17)}=2.8313$, p=0.0868. Tukey HSD; Appetitive vs Consummatory, p=0.9028. Appetitive vs Post-ejaculatory, p=0.0882. Consummatory vs Post-ejaculatory, p=0.2666). Appetitive group, n = 6. Consummatory group, n = 5. Post-ejaculatory group, n = 8.

All data are shown by thin lines for individual subjects and thick lines for mean. The data were analyzed by ANOVA test with Tukey's HSD post-hoc test (FWER = 0.05) (**B, C, I, J, K, L**), Student's t-test (**E**, **F**) or Student's relative t-test (**H**). *p < 0.05, **p < 0.01, ***p < 0.001; ns, not significant.

1079

Figure S10: MPOA^{post-ejaculatory} cells supress female sexual behavior in home cage mating assay but not other behaviors, related to Figure 6.

(**A**) Schematic of pharmaco-genetic activation of MPOA neurons during a home cage mating assay. Female subjects were primed with hormone before the experiment day. On the experiment day, subjects were injected i.p. with CNO to activate TRAP cells 30 minutes before the experiment. Animal behavior was recorded for 90 minutes.

(**B**) The male partner was placed into the home cage of the female subject.

(**C–F**) Number of female-to-male sniff (**C,** Post-ejaculatory-eYFP vs Appetitive-hM3Dq, p=1. vs Consummatory-hM3Dq, p=0.2581. vs Post-ejaculatory-hM3Dq, **p=0.0072.), male mounting (**D,** Post-ejaculatory-eYFP vs Appetitive-hM3Dq, p=1. Post-ejaculatory-eYFP vs Consummatory-hM3Dq, p=1. Post-ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, p=0.4116.), male intromission (**E,** Post-ejaculatory-eYFP vs Appetitive-hM3Dq, p=1. Post-ejaculatory-eYFP vs Consummatory-hM3Dq, p=0.3042. Post-ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, *p=0.0165) and the receptive score (number of mounting episodes with intromissions over all mounting

episodes) (**F**, Post-ejaculatory-eYFP vs Appetitive-hM3Dq, p=1. Post-ejaculatory-eYFP vs Consummatory-hM3Dq, p=0.1888. Post-
ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, p=0.0626).
(**G**) Schematic of conditional place preference (CPP) assay to evaluate the valence of activation of MPOA neurons. A two-chamber
behavior apparatus, one room with vertical stripes and one room with horizontal stripes. On the first day animals were placed in the
behavior apparatus to test their initial preference to each room for 30 minutes (Pre-test). On the conditioning days, the animals were
placed in one side of the room where they were i.p. injected with either saline as control or CNO. CNO was injected 4 hours after the
saline injection. On the last day, the animals were placed in the behavior apparatus to test their preference 30 minutes (Post-test).
(**H–J**) The difference in the time spent in the CNO zone (Time spent in the CNO zone in Post-test – Pre-test) (**H,** Post-ejaculatory-
eYFP vs Appetitive-hM3Dq, p=0.2203. Post-ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, p=1. Appetitive-hM3Dq vs Post-ejaculatory-
hM3Dq, p=1.), the difference in the number of CNO zone entries (Entries in Post-test – Pre-test) (**I,** Post-ejaculatory-eYFP vs Appetitive-
hM3Dq, p=1. Post-ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, p=1. Appetitive-hM3Dq vs Post-ejaculatory-hM3Dq, p=1.) and the
difference in velocity of the animal (Velocity in Post-test – Pre-test) (**J,** Post-ejaculatory-eYFP vs Appetitive-hM3Dq, p=1. Post-
ejaculatory-eYFP vs Post-ejaculatory-hM3Dq, p=0.8472. Appetitive-hM3Dq vs Post-ejaculatory-hM3Dq, p=0.8472.).
(**K**) Representative image of pup retrieval test. Female subject was i.p. injected with saline as control or CNO in their home cage. 30-
minutes after injection, 3x 7w-old pup was placed in the corner of the home cage. Animal behavior was examined for 10 minutes.
(**L–O**) The number of pup retrievals (**L,** eYFP, p=1**,** hM3Dq, p=1), the number of nest building (**M,** eYFP, p=1, hM3Dq, p=0.635), the
number of pup grooming (**N,** eYFP, p=1, hM3Dq, p=1) and the number of pup sniffing (**O,** eYFP, p=1, hM3Dq, p=0.391).
All data are shown by thin lines for individual subjects and thick lines for mean. Appetitive-hM3Dq, consummatory-hM3Dq and post-
ejaculatory-hM3Dq group were tested against post-ejaculatory-eYFP group by Mann–Whitney U test with Bonferroni correction (**C–F**,
H–J) and Wilcoxon rank sum test (**L–O**) with Bonferroni correction. post-ejaculatory-eYFP: n = 6, appetitive-hM3Dq: n = 7,
consummatory-hM3Dq: n = 7. post-ejaculatory-hM3Dq: n = 8. $*$p < 0.05, $**$p < 0.01, $***$p < 0.001; ns, not significant.
1112
1113

Figure S11: MPOA$^{\text{post-ejaculatory}}$ cells is not necessary for sexual motivation but for sexual receptivitiy.

(**A**) Schematic of selective labeling of MPOA neurons using TRAP2. AAV5 EF1a-DIO-hM4Di-mCherry or AAV5 EF1a-DIO-eYFP as
control was injected into the MPOA of the female TRAP2 subjects. After incubation of the virus, the female subjects went through
procedures to TRAP cells in MPOA (further described in **C**).

(**B**) Schematic of pharmaco-genetic inhibition of MPOA neurons during a female self-paced mating assay. Female subjects were primed
with hormone before the experiment day. On the experiment day, subjects were injected i.p. with CNO to inhibit TRAP cells, or saline
as control. 30 minutes after the injection, the male partner was placed into the apparatus and the animal behavior was recorded for 2
hrs.

(**C**) After incubation of the virus, the female subjects were administered with hormones, then received an ejaculation from the male
partner. Immediately after, the subjects were administrated with 4-hydroxytamoxifen (4-OHT).

(**D**) Raster plots of time spent in the interaction zone the female self-paced mating assay (Post-ejaculatory-eYFP: gray. Post-
ejaculatory-hM3Dq: orange).

(**E–L**) Number of zone transitions per 10 minutes (**E**, eYFP, p=0.4375. hM4Di, p=1.), average time in isolation zone (**F**, eYFP, p=1.
hM4Di, p=1.), average time in interaction zone (**G**, eYFP, p=0.3125. hM4Di, p=0.15625.), number of female-to-male sniff (**H**, eYFP,
p=0.1592. hM4Di, p=0.9917.), female self-grooming (**I**, eYFP, p=0.2883. hM4Di, p=1.), male mount (**J**, eYFP, p=0.4456. hM4Di, p=1.),
male intromission (**K**, eYFP, p=0.4456. hM4Di, p=1.) and the receptive score (number of mount episodes with intromissions/number
of all mount episodes) (**L**, eYFP, p=1. hM4Di, p=0.1094).

(**M**) Schematic of pharmaco-genetic inhibition of MPOA neurons during a home cage mating assay. Female subjects were primed with
hormone before the experiment day. On the experiment day, subjects were injected i.p. with CNO to activate TRAP cells 30 minutes
before the experiment. Animal behavior was recorded for 90 minutes.

(**N**) The male mating partner was placed into the home cage of the female subject.

(**O–R**) Number of female-to-male sniff (**O**, p=0.6969.), male mounting (**P**, p=0.4003.), male intromission (**Q, **p=0.0045.) and the
receptive score (number of mounting episodes with intromissions over all mounting episodes) (**R**, *p=0.01998.).

All data are shown by thin lines for individual subjects and thick lines for mean. The data were analyzed by Wilcoxon rank sum test (**E–**
L) with Bonferroni correction. Post-ejaculatory-eYFP: n = 6, post-ejaculatory -hM4Di: n = 8, or by Mann-Whitney U test (**O–R**). Post-
ejaculatory -eYFP: n = 6, post-ejaculatory -hM4Di: n = 8∗p < 0.05, ∗∗p < 0.01, ∗∗∗p < 0.001; ns, not significant.

1141